Is there a trade-off between host generalism and aggressiveness across pathogen populations? A

synthesis of the global potato and tomato blight lesion growth data

Justin SH Wan₁*

*Corresponding author

Email: wanjustinsh@gmail.com

₁Institute of Environmental Health and Ecological Security, School of Environment and Safety

Engineering, Jiangsu University, Zhenjiang, P.R. China

ORCID: 0000-0002-3184-8683

Statements and Declarations: The sole author declares there is no conflict of interest.

Data availability statement: All data generated or analysed during this study are included in this

published article (and its supplementary information files).

Abstract

Phytophthora infestans populations and lineages vary widely in host specificity from specialisation to

generalism among potato and tomato. However, generalists that displace others and became dominant

on both hosts are relatively uncommon. Generalists may have lower fitness compared to specialists on

the host of the latter, which could explain their coexistence at many locations. Lesion size is an

aggressiveness metric closely related to fitness in *P. infestans*. A trade-off between generalism and

lesion growth rate on the original host can explain the variation among blight populations. I collated

the data from cross-inoculation trials on potato and tomato isolates to test for the trade-off. In addition,

other metrics related to disease symptoms were included to test whether the degree of specificity is

different between populations from potato and tomato, and to explore whether specificity had changed

over time. The results indicate a trade-off between generalism and lesion growth rate where higher

specificity was associated with significantly faster lesion growth on the original host at the

population-level, but not at the lineage-level. Potato and tomato isolates were overall not significantly

different in specificity, but tomato isolates tended towards generalism with time. These findings

indicate that specialists may avoid displacement on their host by generalists through faster lesion

growth, and help explain the common co-occurrence of generalists and specialists. However, a few

invasive generalists can rapidly displace competitors and became dominant on both hosts across a

broader region. It is likely that important exceptions exist to this trade-off.

Keywords: Fitness, local adaptation, pathogenicity, *Solanum tuberosum*, *Solanum lycopersicum*,

specialisation

Introduction

Local adaptation to hosts can determine the spatial and temporal population dynamics of plant pathogens (Kröner et al. 2017). Host specificity is indicated by a greater pathogenicity on the original host than on other possible hosts. In the extreme case, the pathogen population is host-specific where the pathogen is only pathogenic on the original host. Adaptive processes occurring within pathogen populations have strong effects on the severity of outbreaks (Sicard et al. 2007). Studying trade-offs and constraints in the population biology of pathogens are of fundamental interest because of their effects on population structure and epidemiology (Burdon, 1993). For example, the emergence of sister species *P. mirabilis*, *P. ipomaea*, and *P. phaseoli* were associated with specialization onto lesser hosts of *P. infestans* (Raffaele et al. 2010). In addition, severe outbreaks often involve the invasion of new aggressive strains with different properties such as superior performance on hosts or environmental tolerances. Adaptation to hosts is an important aspect of disease emergence (Engering et al. 2013), so it is imperative to study the evolutionary and population biology of emergent pathogens on their hosts.

Phytophthora infestans de Bary is a globally distributed heterothallic oomycete pathogen on solanaceous plants. It was the causal agent of the Great Irish Potato Famine that was responsible for millions of deaths. The pathogen was formally described in the 1800s (as *Botrytis infestans* by M.J. Berkeley 1846) and primarily causes disease on potato and tomato on which it is among the most destructive of pathogens (termed late/potato and tomato blight respectively). Populations of newly emergent lineages have occurred continuously, mainly with migration (Shattock 2002; Kildea et al. 2012). Strains vary widely in host specificity among the two host species. The co-occurrence, displacement, and turnover of genotypes through time have been as intriguing to investigators as it was impeding to managers. The global blight populations have undergone several remarkable changes throughout their history, from pan-globally dominant lineages (FAM-1 followed by US-1) to the expansive array of genotypes today (reviewed in Yoshida et al. 2013; Saville et al. 2016). Fitness is related to the effects of pathogens on their hosts and affects the persistence or invasiveness of strains (Pariaud et al. 2009). While no strain persists indefinitely, in invasive strains their persistence is prolonged (e.g., over several seasons or more) and are often dominant (i.e., with high local frequency) across a meta-population level or continental region. Therefore, a better understanding of fitness trade-offs within *P. infestans* on potato and tomato is needed to identify the factors that shape population structure (Andrivon et al. 2013; Saville & Ristaino, 2019).

Host specificity is readily quantified in cross-inoculation trials involving isolates from different host species or varieties. In the literature, aggressiveness is broadly defined as the negative effect that a pathogen has on the host, and can be considered to be any quantitative component of pathogenicity that is non-specific to host genotypes (reviewed in Pariaud *et al.* 2009). In this study, lesion growth

rate on host tissue was used as a measure of aggressiveness, which is a life-history trait that is a key

predictor of epidemic severity and is widely evaluated across studies on *P. infestans* (Birhman &

Singh, 1995; Pariaud et al. 2009; Montarry *et al*. 2010). Previous reports often found that potato

isolates (derived from potato as the original host) have a higher preference for potato, but tomato

isolates (derived from tomato as the original host) are generalist on both hosts (Legard et al. 1995;

Michalska et al. 2016; Kröner et al. 2017). Infectivity on tomato likely evolved relatively recently

during initial invasion from the centre of origin in the 1800s, because isolates from the native range

were non-pathogenic to tomato (Legard et al. 1995). Blight was long believed to spread from potato to

adjacent tomato crops, but not vice versa (Berg, 1926; Small, 1938; Legard et al. 1995; Cohen, 2002).

However, more recently it has been revealed that transmission regularly occurs from tomato to potato

in relatively new lineages such as 13_A2, US-22, and US-23 (Hu et al. 2012; Frost et al. 2016).

Although pathogenicity on tomato likely evolved more recently, the pathogen is typically considered

to be as impactful on tomato as it is on potato (Kröner et al. 2017).

Historical observations on host specificity in populations of the pathogen are particularly thought-

provoking. *Phytophthora infestans* likely originated from the central highlands of Mexico, where

there is a high diversity of tuber-bearing *Solanum* species (Garry et al. 2005; Shakya *et al.* 2018).

Prior to the early 1900s blight on tomato was uncommon (Berg, 1926). Reports from the earliest

cross-inoculation trials were mostly conducted after 1900, when tomato blight was becoming more

prevalent globally. Those reports (between 1900 and 1915) did not mention finding host specificity,

or they found that isolates from both hosts were destructive on both hosts under lab and field

conditions (Berg, 1926; Oyarzun et al. 1998). However, this changed in the after 1915 when

remarkable specificity was frequently noted either on both potato and tomato isolates or in potato

isolates only (from Holland, Australia, USA and the UK; Berg, 1926). This was followed by the

emergence of US-1 lineage which became dominant from the 1930s. During this period, studies on

populations from different regions showed the majority had no remarkable specificity, but potato

isolates were still more likely host-specific. For example, Wilson and Gallegly (1955) examined

isolates from Scotland, the Netherlands, Canada and the USA, and reported that 8 out of 29 potato

isolates were host-specific versus only 2 out of 16 tomato isolates. Isolates from Israel at the time

showed similar proportions (9 of 25 potato isolates vs. only 2 of 25 for tomato isolates; Kedar et al.

1959). Other cross-inoculation studies conducted during US-1 dominance in Japan (Kishi, 1962), the

USA (before 1970, Flier et al. 2003), and Germany (Günther et al. 1970) also reported little or no

specificity. This period was followed by the emergence of an array of successive lineages from the

mid- to late-1980s up to the present where US-1 became either extinct or rare relative to other

lineages. Studies on surviving (or remaining) US-1 populations after the early 90s overwhelmingly

found host specificity in populations from either host species, but mostly on tomato and not potato

(Erselius et al. 1997; Oyazun et al. 1998; Vega-Sanchez et al. 2000; Suassuna et al. 2004; Chen et al.

2009). This pattern is reversed In North America where the remaining US-1 potato isolates tended to
be host-specific, but not the tomato isolates (Legard et al. 1995; Platt, 1999). Subsequently emerged
lineages vary widely in the specificity spectrum. It must be noted that specificity mainly pertains to
the population-level rather than the lineage-level (or genotypic-level; Pariaud et al. 2009). For
example, in the 90s isolates in Uganda and Kenya from potato and tomato were both US-1, but there
were differences in host specificities and glucose-6-phosphate isomerase (*Gpi*) phenotypes among
populations on the two hosts (Erselius et al. 1997). In another example, US-8 is a well-known potato
specialist but in the late-90s populations were reported on tomato on Prince Edward Island, Canada.
Pathogenicity tests on those isolates did not find strong specificity to either host (Daayf and Platt
2003).
121
The overall evidence suggests *P. infestans* genotypes that are more aggressive and could effectively
overcome host defences should be more successful, especially those with a greater lesion size and/or
higher sporulation (Gisi et al. 2011). As such those two life-history metrics are closely related to
fitness across pathogen genotypes (modelled in Montarry et al. 2010). For instance, the relatively high
aggressiveness of US-8 (lesion expansion curve) is likely an important factor in the replacement of
US-1 in south-western Canada (Miller et al. 1998). Similarly the more aggressive 2_A1 displaced the
less aggressive US-1 in east Africa (Njoroge et al. 2018). In Brazil, the more aggressive BR-1 with
much faster lesion growth on potato (but less aggressive on tomato) displaced US-1 on potato only
(Suassuna et al. 2004). The 13_A2 lineage was dominant on several continents including Europe and
Asia, and had a relatively long presence over many seasons. It is more aggressive than other lineages,
and was demonstrated to competitively exclude other lineages in the field, which likely explains its
quick expansion across Europe within 3 years (Cooke et al. 2012). Meanwhile, strains that can
effectively infect a variety of hosts, including resistant hosts should also be favoured over host-
specific strains (Seidl Johnson & Gevens, 2014). Many of the invasive strains that spread quickly to
high frequency were aggressive on both hosts, including 13_A2 (Cooke et al. 2012), US-11 (Chen et
al. 2009; Daayf and Platt 2003), and US-23 (Saville & Ristaino, 2019). The long-standing question
remains: Why aren't pathogen populations all explicitly both highly aggressive and generalist across
hosts?
140
The relationship between pathogenicity components and fitness is complex (Montarry et al. 2010).
Although there is abundant evidence that a faster lesion growth is advantageous for fitness, conversely
there have been few suggestions high pathogenicity is costly to transmissibility because hosts are
weakened or killed too quickly (Pasco et al. 2016; Mariette et al. 2016). This is in line with the
'virulence-transmission trade-off' hypothesis which posits that high host mortality limits transmission
(note that the "virulence" term was used to mean quantitative pathogenicity on the host; Acevedo et al.
2019). Models and studies on trade-offs on *P. infestans* often did not consider the effects of alternate

host habitats, such as tomato plants or wild *Solanum* species (Montarry et al. 2010; Frost et al. 2016).

This is an important aspect since populations on tomato and potato often differ in genotypic

composition (Lebreton & Andrivon 1998). A strain that is extremely pathogenic on potato can have

slower lesion growth on tomato to compensate an assumed lowered transmission rate on potato if the

other host is locally available. Even if high pathogenicity is indeed linked with lower transmissibility,

a trade-off between host generalism versus aggressiveness could explain the variation in host

specificity among *P. infestans* potato and tomato strains (Thrall & Burdon, 2003; Pariaud et al. 2009).

Such a trade-off would be in line with generalists being 'the jack of all trades but a master of none'

and specialists being 'the master of some' (Remold, 2012). It is also evident that few lineages highly

aggressive on both hosts exist (i.e., 'the master of all' strategy). Perhaps a trade-off between the

ability to effectively infect both hosts and aggressiveness on the original host may help explain the

scarcity of highly aggressive and generalist genotypes.

A range of cross-inoculation trials on *P. infestans* from potato and tomato have been conducted over

the past few decades. Meta-analytic approaches use a standardized within-study effect size that can

account for the variation among different metrics used across studies (Chandrasekaran et al. 2016;

Acevedo et al. 2019), and have been used effectively in uncovering patterns or processes across

studies in plant pathology (Madden & Paul, 2011). It is also a good tool for assessing or comparing

the efficacy of different disease control measures across trials (e.g. in pepper blight, Wan & Liew

2020). Here, I collated the global lesion growth rate data to analyse the relationship between host

specificity and aggressiveness. In addition I tested for differences in specificity and lesion growth rate

among potato and tomato isolates. The following predictions were tested: (1) There will be a positive

relationship between host specificity and lesion growth rate on the original host; (2) overall, potato

isolates will have higher specificity for potato than tomato isolates for tomato; (3) host specificity will

tend to reduce over time (with selection for generalists); and (4) isolates of the old US-1 lineage will

have slower lesion growth rate than those of successive, more aggressive lineages.

Methods

Data collation

To identify studies performing controlled cross-inoculations using potato and tomato isolates of *P.*

infestans, I searched ISI Web of Science (Clarivate) and Google Scholar (https://scholar.google.com)

in January 2022 using the following terms: 'Phytophthora infestans' AND 'tomato' OR 'potato'. Only

trials conducted on potato and tomato hosts were selected. Studies that tested pathogenicity on only

one out of these two hosts were excluded, as well as studies that did not report the original host of

isolates (or was indeterminable based on available information). As such, 'potato isolates' or 'tomato

isolates' refer to the respective original host. To address predictions (2), (3) and (4), studies reporting

pathogenicity metrics other than lesion growth rate were included to assess the degree of specificity

across populations. Studies must describe some quantitative or semi-quantitative measure of
pathogenicity (e.g. lesion size, AUDPC, degree of infection). Results not in journal articles such as
research thesis and reports were accepted to reduce publication bias (Madden & Paul 2011). Thus,
those reporting only qualitative pathogenicity (i.e. pathogenic or not pathogenic) are excluded. Studies
that tested multiple isolates were pooled to one data point (separately for each original host). This was
done to avoid pseudoreplication leading to bias towards studies reporting more detailed data.

From each study, I recorded: (1) the original host (potato or tomato); (2) lesion growth rate (hereafter
'LGR' in millimetres per day) on both hosts. Alternatively another measurement of pathogenicity was
recorded to address the other predictions (2-4). Semi-quantitative metrics such as percentages, disease
severity scores or scaled grading were also accepted as host specificity was analysed using non-
parametric analyses. For studies reporting lesion areas, the length should ideally be assumed elliptical
and calculated using width (length = area / [$0.25\pi \cdot$ width]; Vleeshouwers et al. 2000). However, the
width of lesions was not reported in the studies, so lesions were assumed to be square-shaped and
converted using: length = $\sqrt{mm^2}$, divided by the number of days post-inoculation to obtain mm/day; (3)
location of the pathogen population; and (4) time of isolation from hosts (calendar year). The mean
calendar year was used where multiple isolates were pooled.

The within-study differences in pathogenicity of isolates on the two hosts were standardized using
$\log_{10}$ response ratio (LRR) as the effect size. Very negative LRR values (less than minus one) are
associated with extreme specificity (i.e., non-pathogenic on the other host). Conversely, a very
positive LRR (greater than one) indicates the pathogen is only pathogenic on the other host and
therefore the original host is likely a sink habitat. This standardized LGR metric was used to gauge
the aggressiveness of pathogen populations on the two hosts. Henceforth, LGR_{origin} and LGR_{other} refer
to the LGR on the original host and the other host respectively. LRR_{LGR} refers to the LRR calculated
from LGR, used in the analyses on aggressiveness.

To test the relationship between specificity and aggressiveness, the analysis on LGR_{origin} vs. LRR_{LGR} is
divided into two parts: 1) Population-level patterns. By extracting only one data point per host from
each study (i.e., pooling isolates and lineages from each host), each data point is representative of the
whole population on the hosts at that location. Thus, in this part I test whether host specificity is
associated with greater aggressiveness across pathogen populations strictly in general. 2) Lineage-
level patterns (sub-population). Studies reporting the LGR of multilocus genotypes (MLG) or lineages
were recorded as separate entries in another dataset. This allows for analysis on whether MLGs with
higher specificity also tend to be more aggressive. Where only a single lineage is found on a host the
pathogen population is represented by that lineage on that host.

In addition, I compared the host specificity and LGR of US-1 isolates and other lineages. From all the

studies collected from the literature search, studies that tested US-1 isolates were identified. Part of

these studies also tested isolates that are not US-1 along with other isolates (which were pooled with

US-1 in the other analyses). Those were un-pooled and entered as separate entries comprising of 'US-

1' and 'Other' lineage categories. Limiting the analysis to studies that tested both US-1 and

successive lineages within the same trial allows for more direct comparison.

Statistical analyses

Normality in distributions was checked using quantile-quantile plots (Q-Q plots) prior to analysis,

which indicated the LRR data did not follow a normal distribution. To assess whether host specificity

is associated with greater LGR, the relationship between LGR_{origin} and LRR_{LGR} was analysed in two

ways. For the population-level analysis, a median-based linear model with the Thiel-Sen single

median method was used (Komsta & Komsta, 2013). Theil-Sen estimates the slope from the median

among all possible combinations of slopes between points. This method is robust and assumes all

LGR values are different, so all duplicate values was adjusted by a token amount (0.001). For the

lineage-level analysis, the median model could not account for non-independence from data points

extracted from the same study (there were only 10 studies reporting the lineage and data points from

the same study ranged from 2-7), so linear mixed model was used with 'Study' as a random effect.

The LRR_{LGR} data was exponential transformed using $\exp(LRR_{LGR})$. Three outliers with $LRR_{LGR} < -1$

were removed prior to analysis.

For the analysis on whether potato and tomato isolates had statistically significant host specificities, a

non-parametric bootstrap approach was used to estimate the mean effect size (LRR) and confidence

intervals (bootstrapped 95% CI). Bootstrapping is based on the method used in van den Noortgate and

Onghena (2005) and involves random subsampling with replacement over 1000 iterations. This

approach is more suitable because pathogenicity trials often did not report the variance (as compared

to a classical meta-analysis approach). Non-parametric bootstrapping does not emphasize thresholds

for statistical significance, but emphasizes effect size and confidence intervals (Rillig et al. 2019).

Median-based linear models were used to test the relationships between LGR_{origin} vs. LRR_{LGR}, and

LRR vs. time (calendar year). The LRR and LGR differences between potato versus tomato isolates,

and US-1 versus other lineages were analysed using Mann-Whitney U tests.

All analyses were conducted using R (v.3.6.1, R Core Development Team 2019). Bootstrapping was

conducted using the *boot* package (v. 1.3-22) and the *metaphor* package (v. 2.1-0) was used to

calculate effect sizes. The *mblm* package (v.0.12.1; Komsta & Komsta, 2013) was used to run the

median-based models, and the *lmer* package was used to run the linear mixed model on the lineage

data. Host specificities (LRR) were considered statistically significant where the bootstrapped CI do

not overlap with zero (e.g., Chandrasekaran et al. 2016). GetData Graph Digitzier software (v. 2.26)

was used to extract data from figures (available at getdata-graph-digitizer.com).

Results

This search included 78 host specificity comparisons of *P. infestans* from 44 studies in total ($k = 44$

and 34 for potato and tomato respectively, Fig. 1). Of these, 43 were of lesion size data. Most studies

(32 of the 44 studies) conducted cross-inoculation trials on isolates from both hosts within the same

trial. The data was drawn from all over the globe and all inoculation trials were conducted under

controlled trial conditions (Fig. 2). Ten of the studies tested US-1 isolates along with isolates of

successively emergent lineages. An outlying data point with LRR value of greater than 1 (i.e., non-

pathogenic on the original host but highly pathogenic on the other host) was removed from all

analyses (summary statistics in Fig. S1). For aggressiveness (LGR), all data were derived from

detached leaf assays on potato and tomato. For the expanded host specificity data used in comparing

specificity (LRR) within and among potato and tomato isolates, the vast majority of pathogenicity

metrics were directly related to lesions (63 out of 78 data points, e.g., lesion size, growth, percentage,

or number), and 5 data points were based on the evaluation of lesions (proportion of plants or leaves

with lesions or disease ratings /grading based on the type of lesions – sporulating, expanding or non-

expanding). The remainder (10 data points) include sporulation (density or proportion of lesions with

sporulation – 5 data points) and overall disease severity (5 data points). The inoculation approach

used was mostly detached leaf assays (64 out of 78 data points, with one combining data from

detached stem and leaf inoculations), 4 data points were of leaf inoculation in planta, 10 used spray

inoculation and a single one was of tuber and stem inoculation. The isolation time of pathogen isolates

from plants in the field ranged from 1970 to 2017. The full dataset and references are provided in the

Supplementary Material (Table S1).

At the population-level, there was a significant positive relationship between host specificity (LRR_{LGR})

and aggressiveness on the original host, LGR_{origin} ($P < 0.05$, Table 1a, Fig. 3), but not at the lineage-

level (Table 1b). These together indicate that host specificity is generally associated with faster lesion

growth across pathogen populations across locations and regions, but not among genotypes. Extreme

specificity to potato or tomato was rare, and pertains to only few cases (such as certain US-8

populations; Table S1). Overall, potato and tomato isolates had statistically significant specificities for

their original host (LRR estimates with 95% bootstrap CI: -0.26 [-0.09, -0.52] and -0.18 [-0.01, -0.41]

respectively, $P < 0.05$; Fig. 4). This means that overall the pathogenicity when infecting the other host

is reduced to ~55% and 66% of when on the original host for potato and tomato isolates, respectively.

The specificity difference between potato versus tomato isolates (LRR -0.26 vs. -0.18 respectively)

was not significant (W = 673, $P = 0.45$; Fig. 4a), and neither were the differences when only the LRR

based on lesion growth was considered, LRR_{LGR} (W = 173, $P = 0.18$). There was also no difference in

aggressiveness (LGR) between potato and tomato isolates on their respective original hosts, LGR_{origin}

(W = 239, P = 0.99; Fig. 4b, c). There was no significant relationship between LRR and isolation time

for potato isolates (P = 0.54), but for tomato isolates there was a significant positive relationship,

where the median LRR increased from -0.18 (i.e. reduced to ~84% pathogenicity on the other host

compared to when on original host) to 0.06 (i.e. to ~106% pathogenicity on other host compared to

when on original host) over around 30 years (P < 0.01, Table 2; Fig. 5). This suggests that the most

recent tomato populations have little if any specificity – from year 2010 onwards the tomato isolates

data is mainly represented by generalists such as US-22, US-23, and 13_A2.

Focussing on US-1 versus successively emerged lineages from the studies, the sample size available

for analysis was relatively low (k = 11 and k = 10 for US-1 and other isolates respectively) and fewer

reported lesion growth rates (k = 7 and k = 6 respectively). Two US-1 tomato isolates had extreme

specificity for potato and was non-pathogenic on its original host which were removed from the

analyses (LRR 2.31 and 1.58). There were no significant differences in LRR among isolates of US-1

and other lineages (LRR -0.22 vs. -0.30 for US-1 and other lineages respectively; W = 51, P = 0.66).

Furthermore, there was no significant difference in LGR_{origin} among US-1 and other successively

emergent lineages (LGR 4.12 vs. 3.47 mm per day respectively; W = 5, P = 0.57). The low

availability of data between US-1 and new isolates means that the results for this comparison should

be taken with caution. Nonetheless as the analyses are on a subset that assessed US-1 and other

lineages within the same controlled study, it is indicated that those US-1 isolates do not necessarily

have lower LGR than those of newer lineages.

Discussion

The current synthesis of the global data suggests generalism tended to be costly to lesion growth rate

in *P. infestans* across pathogen populations on each host. *Phytophthora infestans* is a hemibiotrophic

pathogen with a clear necrotrophic phase, so aggressiveness should be under selection in the absence

of trade-offs (Jarosz & Davelos, 1995; Montarry et al. 2007). Aggressive lineages often succeed less

aggressive ones (e.g. Miller et al. 1998; Cooke et al. 2012), so a slower lesion growth rate is likely a

fitness cost for the ability to effectively infect both hosts for many generalist populations. From the

data, the most aggressive populations (with LGR on original host > 5 mm/day) were mostly host-

specific (5 out of 7 data points had LRR_{LGR} < -0.30). Moreover the top two (with LGR on original host

~6 mm/day were extreme in host specificity (LRR_{LGR} < -1, see Fig. 3). However, this trade-off was not

found at the lineage-level. Populations typically consist of multiple competing lineages. Within each

population, specialists may only require a small fitness advantage on its host (e.g. a slightly higher

LGR) relative to co-occurring generalists to avoid displacement by them (Kröner et al. 2017). This

difference may be too small to be detected in the lineage data, or that there is a lot of variance in the

data. Unfortunately, there were only 10 studies that reported lineage specific data, and other studies

combined data from different lineages or did not identify the MLG of isolates, which limits the

certainty in the conclusion. This main finding indicates that populations consisting of host-specific

genotype(s) tend to be more aggressive on that host versus populations that comprise of more

generalists. This trade-off is in line with the classic trade-off between virulence and aggressiveness in

the GFG model within a given host species (Thrall & Burdon, 2003), and may help explain the

existence of specialists when both hosts are often abundantly available in the environment. The

finding is consistent with the first prediction that there will be a positive relationship between host

specificity and LGR on the original host.

There are some genotypes where this trade-off may not be applicable. Invasive generalists can quickly

displace competitors and became dominant on both hosts across a greater region, such as 13_A2

across Asia (Dey et al. 2018). In those cases the whole populations on both hosts is represented by the

single lineage. Such dominant lineages are relatively few because the vast majority of lineages are not

and do not become dominant. There are a number of possibilities for this. Migration restrictions and

control measures (e.g. biosecurity on potato trade) can prevent introduction to new areas and inhibit

their spread. Other explanations include limitations in genetic mechanisms leading to the emergence

of aggressive generalists (e.g. epistatic pleiotropy, Remold, 2012) or compensatory selection that may

recover the costs of generalism (Pariaud et al. 2009). The genetic mechanisms and ecological context

for how highly aggressive generalists arise remain to be explored (Burdon, 1993). However whether

those invasive genotypes are more aggressive than most specialists could not be interpreted from the

data. While the current study found that populations with faster lesion growth tended to have higher

specificity through the standardised LGR, it is not suitable to directly compare specific data points

sourced from different studies for inferences on their relative pathogenicity or fitness. This is because

of variation among conditions connected to each individual trial (e.g., host cultivar, culturing

conditions, environmental parameters etc.) that replication may produce different results (Andrivon et

al. 2013). For instance, the widespread 13_A2 from India that was dominant on both hosts only had a

moderate lesion expansion rate in trials (~3.5 mm/day in Chowdappa et al. 2015; ~1.6 to 2.6 mm/day

in Dey et al. 2018). It is possible that a specialist more aggressive (on its host) than an invasive

generalist could persist on that host. Testing isolates together under controlled conditions (e.g. with

competitive interactions) is still necessary to measure the relative aggressiveness of genotypes.

Several factors could potentially explain why generalist populations tended to have slower lesion

growth than specialists on the same host. First, specialisation may be associated with increased fitness

in that environment (e.g., more effective suppression of host defences), regardless of any fitness costs

associated with generalism (i.e. 'the jack of all trades' strategy; Remold, 2012). Second, genetic

change involved in adaptation to alternative hosts may involve fitness costs due to antagonistic

pleiotropy. This includes costs associated with virulence genes to overcome host resistance factors

(Pariaud et al. 2009; Montarry et al. 2007, 2010). Third, specialists should have a lower accumulation

of deleterious alleles (Whitlock, 1996), which may hinder the competitiveness of a generalist in the

presence of specialists (Kawecki, 1994). Yang et al. (2016) found no evidence that mutation

accumulation was involved in a trade-off between thermal tolerance and pathogen growth rate in *P.*

infestans, but may be driven by selection under antagonistic pleiotropy. Virulence was traditionally

defined as the ability of a pathogen to overcome host defences, especially in the gene-for-gene (GFG)

model so that high virulence is associated with the ability to infect more hosts (Laine & Barrès, 2013).

Even though lesion size is regularly used to gauge aggressiveness and the severity of outbreaks (e.g.

Miller et al. 1998; Pariaud et al. 2009), both LGR and sporulation capacity are closely linked to

fitness (Montarry et al. 2010). Thus considering LGR alone may not accurately reflect aggressiveness

in cases where sporulation capacity (i.e. spore density within lesion area) is the key factor. There was

insufficient data for analyses based on sporulation. In the compiled literature, there were only 29 data

points in total for sporulation, and measurements differed in scaling (12 were scaled to area or length

and 17 were scaled to volume, Table S1). Other metrics (e.g. host mortality and infection efficiency)

were even less commonly reported. Some lineages had fast lesion growth on both hosts but low

sporulation density on one host (i.e., the host it is less often isolated from). An example where

sporulation is more important is in 23_A1 from Algeria that is mainly found on tomato (Beninal et al.

2022). A trial found that it has relatively fast lesion growth on both hosts (4.9 – 5.0 mm/day) but

lower sporulation capacity on potato (Belkhiter et al. 2019). This is especially compared to its

specialist competitor (13_A2 on potato), which had a slightly lower LGR (4.7 mm/day) but superior

sporulation on potato (see Figure 2a of Belkhiter et al. 2019). Conversely in North America (around

2010), US-23 is genetically similar to 23_A1 but caused major epidemics on both hosts (Danies et al.

2013). It has both higher sporulation capacity and LGR than competing lineages US-22 and potato

specific US-24, except US-8 on potato (Danies et al. 2013). A few years later it completely displaced

US-22 and US-24 (Saville & Ristaino, 2019). Whether sporulation capacity is involved in trade-offs

could be addressed in future research. Nonetheless genotypes with high LGR on both hosts should be

considered potentially invasive.

Interestingly, genotypes with weaker pathogenicity may sometimes persist. There was a suggestion

that potato specialist US-24 may persist in more northern locations despite being less aggressive than

generalists US-22 and US-23 due to its superior performance under cold conditions (Danies et al.

2013). The genetically similar potato-specific US-8 was almost entirely displaced but some remaining

isolated populations (mainly in the West Coast of the USA) may have been temporarily sustained by

inoculum sources at certain localities (Saville & Ristaino, 2019). Long-term studies taking account of

the complete life cycle of the pathogen conducted over multiple seasons are likely needed to more

reliably assess fitness (Andrivon et al. 2013). Other considerations that may influence the current

results include the effects of fungicide presence and other environmental variation (e.g. temperature)

that were not replicated in lab trials. There are alternative metrics such as infection efficiency and

competitive ability (Lebreton et al. 1999) that could not be addressed by this study that may also be

important in gauging aggressiveness.

Contrary to the second hypothesis that potato isolates will be more host-specific, potato and tomato

populations did not differ significantly in specificity, but there may be around twice as many highly

host-specific populations for potato than for tomato by proportion (12 out of 44 data points or 27% for

potato, versus 4 out of 34 data points or 12% for tomato) as defined by an LRR of less than -0.5 (i.e.

pathogenicity reduced to < 32% compared to infecting original host; Fig. 5). The reduction in host

specificity of pathogen populations on tomato over the last few decades is interesting, and only

partially supports the third hypothesis (that overall host specificity will reduce with time) since the

specificity of potato isolates did not change over time. Together, these results generally support the

long-held belief that tomato isolates are more generalist than potato isolates. This may be particularly

relevant for cold regions where potato isolates could survive on tubers but tomato isolates may need

other hosts, so specialisation to tomato may not be under selection.

In the current study, lower lesion growth rate on the original host or higher specificity in US-1

populations was not found. Past studies often observed the displacement of US-1 by more aggressive

lineages (Legard et al. 1995; Reis et al. 2003; Suassuna et al. 2004; Chen et al. 2009). The non-

significance difference in lesion growth could be due to the low number of data points available in

this case. Alternatively there is a possibility that the persistent US-1 reflects selection on a subset

equally aggressive as newer invading populations. This survivorship effect may have also been

observed in *P. cinnamomi*, where the A2 mating type is predominant and displaced the less virulent

endemic A1 throughout Asia (Arentz, 2017). The remaining extant *P. cinnamomi* A1 has similar

aggressiveness (i.e. lesion size) to the A2 populations (Dudzinski et al. 1993; Robin & Desprez-

Loustau, 1998). Nonetheless, future routinized controlled trials testing the pathogenicity (especially of

LGR and sporulation capacity) of pathogen populations on the two hosts would be greatly beneficial

for the management of extant and emergent lineages of potato and tomato blight (such as 36_A2 and

41_A2 in Europe; EuroBlight network: agro.au.dk/forskning/internationale-platforme/euroblight,

accessed Nov 2022), particularly under the effects of climate change (Pangga et al. 2011)..

Conclusions

The current synthesis revealed a trade-off between generalism and aggressiveness across

Phytophthora infestans populations on potato and tomato. Overall, populations consisting of more

specialists have faster lesion growth rate on their original host than populations consisting of

generalists on the same host, which may help explain the patterns in population composition of

generalists and specialists among the two hosts. However, this trade-off may not apply for few

aggressive and invasive generalists that became dominant on both hosts across a broader region.

Although the level of specificity among potato and tomato isolates was not significantly different,

there was an indication tomato isolates tended to become more generalist over the last few decades.

Due to the variability in how pathogenicity traits relate to fitness, long-term studies (e.g. over multiple

seasons) are likely needed to better understand the trade-offs involving pathogenicity. The trade-off

between generalism and aggressiveness may help explain the regular coexistence of specialist and

generalist lineages at many locations.

Acknowledgements

The author expresses gratitude to the researchers whose data contributed to this synthesis. Special

thanks go to Susan Rutherford for her support and help proofreading the manuscript and Edward Liew

for the earlier discussions on Phytophthora and for his advice. Justin SH Wan was supported by the

Jiangsu University Science Foundation Fund (20JDG056).

References

Andrivon, D., Montarry, J., Corbiere, R., Pasco, C., Glais, I., Marquer, B., (2013). The hard life of

461    *Phytophthora infestans*: when trade-offs shape evolution in a biotrophic plant pathogen.

462    *Plant Pathology, 62*, 28-35. 10.1111/ppa.12164

463 Arentz, F. (2017). *Phytophthora cinnamomi* A1: An ancient resident of New Guinea and Australia of

464    Gondwanan origin? *Forest Pathology, 47*, e12342. 10.1111/efp.12342

465 Acevedo, M.A., Dillemuth, F.P., Flick, A.J., Faldyn, M.J., & Elderd, B.D. (2019). Virulence-driven

466    trade-offs in disease transmission: a meta-analysis. *Evolution, 73*, 636-647.

467    10.1111/evo.13692

468 Belkhiter, S., Beninal, L., Khedidji, H., Krimi, A., & Bouznad, Z. (2019). Aggressiveness and host

469    adaptation of some Algerian *Phytophthora infestans* clonal lineage on potato and tomato.

470    *Pakistan Journal of Phytopathology, 31*, 147-154.

471 Beninal, L., Bouznad, Z., Corbiere, R., Belkhiter, S., Mabon, R., Taoutaou, A., Keddad, A., Runno-

472    Paurson, E., & Andrivon, D. (2022) Distribution of major clonal lineages EU_13_A2,

473    EU_2_A1, and EU_23_A1 of *Phytophthora infestans* associated with potato late blight

474    across crop seasons and regions in Algeria. *Plant Pathology, 71*, 458-469.

475    10.1111/ppa.13471

476 Berg, A. (1926) *Tomato late blight and its relation to late blight of potatoes*. Bulletin 2055,

477    Agricultural Experiment Station, West Virginia University. Available online 11-2022:

478    https://core.ac.uk/download/pdf/230443028.pdf

Birhman, R.K., & Singh, B.P. (1995) Path-coefficient analyses and genetic parameters of the
components of field resistance of potatoes to late blight. *Annals of Applied Biology* 127:
353-362. 10.1111/j.1744-7348.1995.tb06679.x

Blandón-Díaz, J.U., Widmark, A.K., Hannukkala, A., Andersson, B., Högberg, N., & Yuen, J.E.
(2012). Phenotypic variation within a clonal lineage of *Phytophthora infestans* infecting
both tomato and potato in Nicaragua. *Phytopathology, 102*, 323-330. 10.1094/PHYTO-02-
11-0033

Burdon, J.J. (1993). The structure of pathogen populations in natural plant communities. *Annual*
Review of Phytopathology, 31, 305–323. 10.1146/annurev.py.31.090193.001513

Chandrasekaran, M., Subramanian, D., Yoon, E., Kwon, T., & Chun, S.C. (2016). Meta-analysis
reveals that the genus *Pseudomonas* can be a better choice of biological control agent
against bacterial wilt disease caused by *Ralstonia solanacearum*. *The Plant Pathology*
Journal, 32, 216-217. 10.5423/PPJ.OA.11.2015.0235

Chen CH, Wang TC, Black L, Sheu ZM, Perez F, Deahl K (2009) Phenotypic and genotypic changes
in the *Phytophthora infestans* population in Taiwan - 1991 to 2006. *Journal of*
Phytopathology, 157, 248-255. 10.1111/j.1439-0434.2008.01483.x

Chowdappa, P., Nirmal Kumar, B.J., Madhura, S., Mohan, S.P., Myers, K.L., Fry, W.E., & Cooke,
D.E.L. (2015). Severe outbreaks of late blight on potato and tomato in South India caused
by recent changes in the *Phytophthora infestans* population. *Plant Pathology, 64*, 191-199.
10.1111/ppa.12228

Cohen, Y. (2002). Populations of *Phytophthora infestans* in Israel underwent three major genetic
changes during 1983 to 2000. *Phytopathology*, *92*, 300-307. 10.1094/PHYTO.2002.92.3.300

Cooke, D.E., Cano, L.M., Raffaele, S., Bain, R.A., Cooke, L.R., Etherington, G.J., Deahl, K.L., Farrer
R.A., Gilroy, E.M., Goss, E.M., & Grünwald, N.J. (2012). Genome analyses of an
aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathogens, 8*,
e1002940. 10.1371/journal.ppat.1002940

Daayf, F. & Platt, H.W. (2003). Differential pathogenicity on potato and tomato of *Phytophthora*
infestans US-8 and US-11 strains isolated from potato and tomato. *Canadian Journal of*
Plant Pathology. *25*, 150-154. 10.1080/07060660309507063

Danies, G., Small, I.M., Myers, K., Childers, R., & Fry, W.E. (2013) Phenotypic characterization of
recent clonal lineages of *Phytophthora infestans* in the United States. *Plant Disease, 97*,
873-881. 10.1094/PDIS-07-12-0682-RE

Dey, T., Saville, A., Myers, K., Tewari, S., Cooke, D.E., Tripathy, S., Fry, W.E., Ristaino, J.B., &
Guha Roy, S. (2018). Large sub-clonal variation in *Phytophthora infestans* from recent
severe late blight epidemics in India. *Scientific Reports, 8*, 4429. s41598-018-22192-1

Dudzinski, M.J., Old, K.M., & Gibbs, R.J. (1993). Pathogenic variability in Australian isolates of
Phytophthora cinnamomi. *Australian Journal of Botany, 41*, 721-732. 10.1071/BT9930721

Engering A, Hogerwerf L, Slingenbergh J (2013). Pathogen-host-environment interplay and disease
emergence. *Emerging Microbes and Infections, 2*, e5. 10.1038/emi.2013.5

Erselius, L.J., Vega-Sánchez, M.E., Rodríguez, A.M., Bastidas, O., Hohl, H.R., Ojiambo, P.S.,
Mukalazi, J., Vermeulen, T., Fry, W.E., & Forbes, G.A. (1997). Host specificity of
Phytophthora infestans on tomato and potato in Uganda and Kenya. *CIP Program Report*
1998, 49-55.

Flier, W.G., Van den Bosch, G.B., & Turkensteen, L.J. (2003). Epidemiological importance of
Solanum sisymbriifolium, *S. nigrum* and *S. dulcamara* as alternative hosts for *Phytophthora*
infestans. *Plant Pathology, 52*, 595-603. 10.1046/j.1365-3059.2003.00922.x

Frost, K.E., Seidl Johnson, A.C., & Gevens, A.J. (2016). Survival of isolates of the US-22, US-23,
and US-24 clonal lineages of *Phytophthora infestans* by asexual means in tomato seed at
cold temperatures. *Plant Disease, 100*, 180-187. 10.1094/PDIS-04-15-0462-RE

Garry, G., Forbes, G.A., Salas, A., Santa Cruz, M., Perez, W.G., & Nelson, R.J. (2005). Genetic
diversity and host differentiation among isolates of *Phytophthora infestans* from cultivated
potato and wild Solanaceous hosts in Peru. *Plant Pathology, 54*, 740-748. 10.1111/j.1365-
3059.2005.01250.x

Gisi, U., Walder, F., Resheat-Eini, Z., Edel, D., & Sierotzki, H. (2011). Changes of genotype,
sensitivity and aggressiveness in *Phytophthora infestans* isolates collected in European
countries in 1997, 2006 and 2007. *Journal of Phytopathology, 159*, 223-232. j.1439-
0434.2010.01753.x

Günther, E., Eggert, D., & Grümmer, G. (1970). Resistance of tomatoes to late blight (*Phytophthora*
infestans). *Annals of Applied Biology, 65*, 255-262. 10.1111/j.1744-7348.1970.tb04585.x

Hu, C.H., Perez, F.G., Donahoo, R., McLeod, A., Myers, K., Ivors, K., Secor, G., Roberts, P.D.,
Deahl, K.L., Fry, W.E., & Ristaino, J.B. (2012). Recent genotypes of *Phytophthora*
infestans in the eastern United States reveal clonal populations and reappearance of
mefenoxam sensitivity. *Plant Disease, 96*, 1323-1330. 10.1094/PDIS-03-11-0156-RE

Jarosz, A.M., & Davelos, A.L. (1995). Effects of disease in wild plant populations and the evolution
of pathogen aggressiveness. *New Phytologist, 129*, 371-387. 10.1111/j.1469-
8137.1995.tb04308.x

Kawecki, T.J. (1994). Accumulation of deleterious mutations and the evolutionary cost of being a
generalist. *The American Naturalist, 144*, 833-838. 10.1086/285709

Kedar, N., Rotem, J., & Wahl, I. (1959) Physiologic specialization of *Phytophthora infestans* in Israel.
Phytopathology, 49, 675-679.

Kildea, S., Quinn, L., Mehenni-Ciz, J., Cooke, D.E.L., Perez, F.M., Deahl, K.L., Griffin, D., & Cooke,
L.R. (2013). Re-emergence of the Ib mitochondrial haplotype within the British and Irish
Phytophthora infestans populations. *European Journal of Plant Pathology*, *135*, 237-242.
10.1007/s10658-012-0098-y

Kishi, K. (1962) Studies on the physiological specialization of *Phytophthora infestans* on tomatoes.
Japanese Journal of Phytopathology 27: 180-188.

Komsta, L., & Komsta, M.L. (2013). Package '*mblm*'. Available online: https://cran.r-
project.org/web/packages/mblm/index.html (accessed April 2022).

Kröner, A., Mabon, R., Corbière, R., Montarry, J., & Andrivon, D. (2017). The coexistence of
generalist and specialist clonal lineages in natural populations of the Irish Famine pathogen
Phytophthora infestans explains local adaptation to potato and tomato. *Molecular Ecology,*
26, 1891-1901. 10.1111/mec.14004

Laine, A.L., & Barrès, B. (2013) Epidemiological and evolutionary consequences of life-history
trade-offs in pathogens. *Plant Pathology* 62: 96-105. 10.1111/ppa.12129

Lebreton, L., & Andrivon, D. (1998) French isolates of *Phytophthora infestans* from potato and
tomato differ in phenotype and genotype. *European Journal of Plant Pathology, 104*, 583-
594. 10.1023/A:1008662518345

Lebreton, L., Lucas, J.M., Andrivon, D. (1999). Aggressiveness and competitive fitness of
Phytophthora infestans isolates collected from potato and tomato in France. *Phytopathology,*
89, 679-686. 10.1094/PHYTO.1999.89.8.679

Legard, D.E., Lee, T.Y., & Fry, W.E. (1995). Pathogenic specialisation in *Phytophthora infestans*:
aggressiveness on tomato. *Phytopathology, 85*, 1356-1361. 10.1094/Phyto-85-1356

Madden, L.V., & Paul, P.A. (2011). Meta-analysis for evidence synthesis in plant pathology: An
overview. *Phytopathology, 101*, 16-30. 10.1094/ PHYTO-03-10-0069

Mariette, N., Mabon, R., Corbiere, R., Boulard, F., Glais, I., Marquer, B., Pasco, C., Montarry, J., &
Andrivon, D. (2016). Phenotypic and genotypic changes in French populations of
Phytophthora infestans: are invasive clones the most aggressive? *Plant Pathology, 65*, 577-
586. 10.1111/ppa.12441

Michalska, A.M., Sobkowiak, S., Flis, B., & Zimnoch-Guzowska, E. (2016). Virulence and
aggressiveness of *Phytophthora infestans* isolates collected in Poland from potato and
tomato plants identified no strong specificity. *European Journal of Plant Pathology, 144*,
325-336. 10.1007/s10658-015-0769-6

Miller, J.S., Johnson, D.A., & Hamm, P.B. (1998). Aggressiveness of isolates of *Phytophthora*
infestans from the Columbia Basin of Washington and Oregon. *Phytopathology, 88*, 190-
197. 10.1094/PHYTO.1998.88.3.190

Montarry, J., Corbiere, R., & Andrivon, D. (2007). Is there a trade-off between aggressiveness and
overwinter survival in *Phytophthora infestans*? *Functional Ecology, 21*, 603-610.
10.1111/j.1365-2435.2007.01252.x

Montarry, J., Hamelin, F.M., Glais, I., Corbière, R., & Andrivon, D. (2010). Fitness costs associated
with unnecessary virulence factors and life history traits: evolutionary insights from the

potato late blight pathogen *Phytophthora infestans*. *BMC Evolutionary Biology, 10*, 1-9.

10.1186/1471-2148-10-283

Njoroge, A.W., Tusiime, G., Forbes, G.A., & Yuen, J.E. (2016). Displacement of US-1 clonal lineage

by a new lineage of *Phytophthora infestans* on potato in Kenya and Uganda. *Plant*

Pathology, 65, 587-592. 10.1111/ppa.12451

Oliva, R.F., Kroon, L.P., Chacón, G., Flier, W.G., & Ristaino, J.B., & Forbes, G.A. (2010).

Phytophthora andina sp. nov., a newly identified heterothallic pathogen of solanaceous

hosts in the Andean highlands. *Plant Pathology, 59*, 613-625. 10.1111/j.1365-

3059.2010.02287.x

Oyarzun, P.J., Pozo, A., Ordoñez, M.E., Doucett, K., & Forbes, G.A. (1998). Host specificity of

Phytophthora infestans on tomato and potato in Ecuador. *Phytopathology, 88*, 265-271.

10.1094/PHYTO.1998.88.3.265

Pangga, I.B., Hanan, J., & Chakraborty, S. (2011). Pathogen dynamics in a crop canopy and their

evolution under changing climate. *Plant Pathology, 60*, 70-81. 10.1111/j.1365-

3059.2010.02408.x

Pariaud, B., Ravigné, V., Halkett, F., Goyeau, H., Carlier, J., & Lannou, C. (2009). Aggressiveness

and its role in the adaptation of plant pathogens. *Plant Pathology, 58*, 409-424.

10.1111/j.1365-3059.2009.02039.x

Pasco, C., Montarry, J., Marquer, B., & Andrivon, D. (2016). And the nasty ones lose in the end:

foliar pathogenicity trades off with asexual transmission in the Irish famine pathogen

Phytophthora infestans. *New Phytologist, 209*, 334-342. 10.1111/nph.13581

Platt, H.W. (1999). Response of Solanaceous cultivated plants and weed species to inoculation with

A1 or A2 mating type strains of *Phytophthora infestans*. *Canadian Journal of Plant*

Pathology, *21*, 301-307. 10.1080/07060669909501195

Raffaele, S., Farrer, R.A., Cano, L.M., Studholme, D.J., MacLean, D., Thines, M., Jiang, R.H., Zody,

M.C., Kunjeti, S.G., Donofrio, N.M. & Meyers, B.C. (2010). Genome evolution following

host jumps in the Irish potato famine pathogen lineage. *Science, 330*, 1540-1543.

10.1126/science.1193070

Reis, A., Smart, C.D., Fry, W.E., Maffia, L.A., & Mizubuti, E.S. (2003). Characterization of isolates

of *Phytophthora infestans* from southern and southeastern Brazil from 1998 to 2000. *Plant*

Disease, 87, 896-900. 10.1094/PDIS.2003.87.8.896

Remold, S. (2012). Understanding specialism when the jack of all trades can be the master of all.

Proceedings of the Royal Society B: Biological Sciences, 279, 4861–4869.

10.1098/rspb.2012.1990

Rillig, M.C., Ryo, M., Lehmann, A., Aguilar-Trigueros, C.A., Buchert, S., Wulf, A., Iwasaki, A., Roy,

J., & Yang, G. (2019). The role of multiple global change factors in driving soil functions

and microbial biodiversity. *Science, 366*, 886-890. 10.1126/science.aay2832

Robin, C., & Desprez-Loustau, M. (1998). Testing variability in pathogenicity of *Phytophthora*
cinnamomi. *European Journal of Plant Pathology, 104*, 465–475.
10.1023/A:1008649806620

Saville, A.C., Martin, M.D., & Ristaino, J.B. (2016). Historic late blight outbreaks caused by a
widespread dominant lineage of *Phytophthora infestans* (Mont.) de Bary. *PLoS ONE,*
e0168381. 10.1371/journal.pone.0168381

Saville, A., & Ristaino, J.B. (2019). Genetic structure and subclonal variation of extant and recent US
lineages of *Phytophthora infestans*. *Phytopathology, 109*, 1614-1627. 10.1094/PHYTO-09-
18-0357-R

Seidl Johnson, A.C., & Gevens, A.J. (2014) Investigating the host range of the US-22, US-23, and
US-24 clonal lineages of *Phytophthora infestans* on solanaceous cultivated plants and weeds.
Plant Disease, 98, 754-760. 10.1094/PDIS-09-13-0924-RE

Shakya, S.K., Larsen, M.M., Cuenca-Condoy, M.M., Lozoya-Saldaña, H., & Grünwald, N.J. (2018).
Variation in genetic diversity of *Phytophthora infestans* populations in Mexico from the
center of origin outwards. *Plant Disease, 102*, 1534-1540. PDIS-11-17-1801-RE

Shattock, R.C. (2002). *Phytophthora infestans*: populations, pathogenicity and phenylamides. *Pest*
Management Science, 58, 944-950. 10.1002/ps.527

Sicard, D., Pennings, P.S., Grandclément, C., Acosta, J., Kaltz, O., & Shykoff, J.A. (2007).
Specialization and local adaptation of a fungal parasite on two host plant species as revealed
by two fitness traits. *Evolution, 61*, 27-41.

Small, T. (1938). The relation between potato blight and tomato blight. *Annals of Applied Biology, 25*,
271-276.

Suassuna, N.D., Maffia, L.A., & Mizubuti, E.S. (2004). Aggressiveness and host specificity of
Brazilian isolates of *Phytophthora infestans*. *Plant Pathology, 53*, 405-413. j.1365-
3059.2004.01043.x

Thrall, P.H., & Burdon, J.J. (2003). Evolution of virulence in a plant host-pathogen metapopulation.
Science, 299, 1735-1737. 10.1126/science.1080070

Van Den Noortgate, W., & Onghena, P. (2005). Parametric and nonparametric bootstrap methods for
meta-analysis. *Behavior Research Methods, 37*, 11-22. 10.3758/bf03206394

Vega-Sánchez, M.E., Erselius, L.J., Rodriguez, A.M., Bastidas, O., Hohl, H.R., Ojiambo, P.S.,
Mukalazi, J., Vermeulen, T., Fry, W.E., & Forbes, G.A. (2000). Host adaptation to potato
and tomato within the US- 1 clonal lineage of *Phytophthora infestans* in Uganda and Kenya.
Plant Pathology, 49, 531-539. 10.1046/j.1365-3059.2000.00487.x

Vleeshouwers, V.G.a.A., W. Van Dooijeweert, F. Govers, S. Kamoun & L.T. Colon. (2000). The
hypersensitive response is associated with host and nonhost resistance to *Phytophthora*
*infestan*s. *Planta 210*, 853-864. 10.1007/s004250050690

Wan, J.S.H., & Liew, E.C. (2020). Efficacy of chemical and biological agents against pepper blight
663   (*Phytophthora capsici* Leonion) in East Asia: a meta-analysis of laboratory and field trial
664   data. *Journal of Plant Pathology*, 102, 835-842. 10.1007/s42161-020-00519-0
665   Whitlock, M.C. (1996). The red queen beats the jack-of-all-trades: the limitations on the evolution of
666   phenotypic plasticity and niche breadth. *The American Naturalist, 148*, S65-S77.
667   http://www.jstor.org/stable/2463048.
668   Wilson, J.B. & Gallegly, M.E. (1955). The interrelationships of potato and tomato races of
669   *Phytophthora infestans*. *Phytopathology, 45*, 473-476.
670   Yang, L.N., Zhu, W., Wu, E.J., Yang, C., Thrall, P.H., Burdon, J.J., Jin, L.P., Shang, L.P., & Zhan, J.
671   (2016). Trade-offs and evolution of thermal adaptation in the Irish potato famine pathogen
672   *Phytophthora infestans*. *Molecular Ecology, 25*, 4047-4058. 10.1111/mec.13727
673   Yoshida, K., Schuenemann, V.J., Cano, L.M., Pais, M., Mishra, B., Sharma, R., Lanz, C., Martin,
674   F.N., Kamoun, S., Krause, J., & Thines, M. (2013) The rise and fall of the *Phytophthora*
675   *infestans* lineage that triggered the Irish potato famine. *eLife, 2*, e00731.
676   10.7554/eLife.00731
677   Young, G.K., Cooke, L.R., Kirk, W.W., Tumbalam, P., Perez, F.M., & Deahl, K.L. (2009). Influence
678   of competition and host plant resistance on selection in *Phytophthora infestans* populations
679   in Michigan, USA and in Northern Ireland. *Plant Pathology, 58*, 703-714. 10.1111/j.1365-
680   3059.2009.02043.x
681   Yuen, J.E., & Andersson, B. (2013). What is the evidence for sexual reproduction of *Phytophthora*
682   *infestans* in Europe? *Plant Pathology, 62*, 485-491. 10.1111/j.1365-3059.2012.02685.x

683 **Table 1**

684 Results on aggressiveness on the original host (LGR_{origin} in millimetres per day) versus host specificity (LRR_{LGR}) at the: a) Population-level from median-based

685 model ($k = 43$), and b) lineage-level from mixed effects linear model with 'Study' as a random effect level. LRR_{LGR} was exponential transformed for normality

686 in the residuals ($k = 28$). Three outliers $LRR_{LGR} < -1$ were excluded). Bold values indicate statistically significant effects ($P < 0.05$).

688 a)

Source	Estimate	Median absolute deviation	V	P
Intercept	-0.00	0.21	470	0.78
LRR_{LGR}	-0.03	0.10	316	**0.037**

695 b)

Source	Estimate	df	t	P
Intercept	0.93	26	12.41	**<0.001**
exp(LRR_{LGR})	-0.03	26	-0.26	0.797

697 **Table 2**

698 Results from median-based linear models on host specificity (LRR) versus isolation time (calendar year) in *Phytophthora infestans* potato and tomato isolates.

699 Bold values indicate statistically significant effects ($P < 0.05$).

Source	Estimate	Median absolute deviation	V	*P*
Potato isolates ($k = 40$)				
Intercept	-0.41	10.29	448	0.62
Calendar year	0.00	0.01	328	0.54
Tomato isolates ($k = 32$)				
Intercept	-17.00	26.98	126	**0.009**
Calendar year	0.01	0.01	402	**0.009**

Figure Captions

Fig. 1

Flow diagram of the literature search and the screening process, detailing the number of studies excluded

during screening up to the final number of studies included.

Fig. 2

World map detailing the locations of *Phytophthora infestans* potato and tomato populations tested in cross-

inoculation trials. Symbols with lighter shades represent imprecise locations specified to the general region

only. Geographically distant populations tested within the same study are joined by dotted lines.

Fig. 3

Relationships between host specificity based on lesion growth rate (LRR_{LGR}) and lesion growth rate on the

original host (LGR_{origin}) for potato and tomato isolates of *Phytophthora infestans* at the: a) Population-level,

where each data point is representative of the pathogen population present on each host at a location.

Populations with no specificity are indicated by $LRR_{LGR} = 0$; b) Lineage-level, with each data point of the

exponential transformed LRR_{LGR} representing a lineage tested within a given study. Lineages with no

specificity are indicated by $\exp(LRR_{LGR}) = 1$. More negative LRR values are associated with greater specificity

on the original host in all cases.

Fig. 4

Boxplots comparing *Phytophthora infestans* populations from potato and tomato, showing the 0.25, median

(0.5) and 0.75 quartiles: a) The overall degree of host specificity (LRR) as indicated by different pathogenicity

metrics; b) host specificity in the subset as calculated from lesion growth rate (LRR_{LGR}), and c) lesion growth

rate on the original host (LGR_{origin}).

Fig. 5

Relationship between host preference (LRR), as indicated by different pathogenicity metrics, versus isolation

time (calendar year) in *Phytophthora infestans* a) potato and b) tomato isolates, where each data point

represents a pathogen population present during the time period.

Supplementary materials

Table S1

The datasets used in the synthesis of host specificity and lesion growth rate in *Phytophthora infestans*

pathogen populations and lineages on potato and tomato.

Fig. S1

Summary median and quantiles for host specificity (LRR) among potato and tomato populations of

Phytophthora infestans in global regions (k = sample size). More negative LRR values indicate higher

specificity. Very negative values (such as less than -1) are host-specific to the original host. One population

had a very positive LRR and was host-specific to the other host which was excluded from all analyses

(indicated by the grey arrow). A value of zero indicates identical pathogenicity on both hosts indicating no

specificity.