Is there a trade-off between aggressiveness and host generalism across *Phytophthora infestans*

(potato and tomato blight) populations? A synthesis of global cross-inoculation data

Justin SH Wan[1]*

*Corresponding author

[1]Institute of Environmental Health and Ecological Security, School of Environment and Safety

Engineering, Jiangsu University, Zhenjiang, China

ORCID: 0000-0002-3184-8683

Conflict of Interest: The sole author declares there is no conflict of interest.

Abstract

Phytophthora infestans is among the most destructive of plant diseases. Pathogen lineages have

varying degrees of host preference among potato and tomato, from no specificity to extreme

specialisation. How host specificity impacts fitness among pathogen populations is unclear.

Comparing the pathogenicity of both population types can shed light on the determinants of host

specificity in *P. infestans*. A trade-off between generalism and quantitative aggressiveness can explain

the variation among pathogen populations. I assessed the existence of such a trade-off by analysing

the global cross-inoculation data on potato and tomato isolates. In addition, host specificity was

assessed for change over time. The synthesis included 44 and 34 data points for potato and tomato

isolates respectively. Potato isolates overall did not prefer their original host significantly more than

tomato isolates. Isolates from potato can generally infect tomato with similar aggressiveness (in terms

of lesion growth rate) and specificity as in reverse. There was a modest trade-off between generalism

and aggressiveness, where high specificity was associated with significantly greater aggressiveness.

However, this trade-off did not explain the presence of generalist strains that are equally aggressive as

specialist strains. A portion of generalists has high aggressiveness equal to the most host specific

strains. For tomato isolates, there was a significant change towards generalism over the last few

decades. This synthesis reveals several novel observations on the evolutionary ecology of the blight

pathogen. Many unresolved questions on host specificity and aggressiveness remain which are

discussed.

Keywords: Local adaptation, specialisation, fitness, *Solanum tuberosum*, *Solanum lycopersicum*

Introduction

Local adaptation and specialisation to hosts can determine the spatial and temporal evolutionary

dynamics of plant pathogens (Kröner *et al.,* 2017). It is a process where populations under divergent

selection achieve better performance in their habitat of origin than all other possible habitats. In plant

pathogens, local adaptation manifests as specificity for the original host over other potential hosts; and

in the extreme case, host specialisation where the pathogen is only pathogenic to the original host.

Host specificity can be readily quantified in cross-inoculation trials of isolates from different host

species or varieties. Adaptation to hosts, environmental conditions, and fungicide application can

change the genetic composition of pathogen populations and thus their epidemiological properties

(Cohen, 2002; Pangga *et al.,* 2011; Engering *et al.,* 2013). Understanding the effects of these drivers

is important in predicting their impacts under climate change (Anderson *et al.,* 2004; Pangga *et al.,*

2011).

Phytophthora infestans de Bary is a globally distributed heterothallic oomycete pathogen on

solanaceous plants. It was the causal agent of the Great Irish Potato Famine that was responsible for

millions of deaths. It is among the most destructive of plant pathogens (Chowdappa *et al.,* 2015; Fry

et al., 2015). The pathogen was formally described in the 1800s (as *Botrytis infestans* by M.J.

Berkeley 1846) and primarily causes disease on potato and tomato (termed late / potato and tomato

blight respectively). Populations of *P. infestans* have been continuously re-emerging through time and

space (Fry *et al.,* 2015). Strains vary dramatically in host specificity among the two host species, from

non-specific generalists (i.e., infects both hosts with equal pathogenicity) to extreme obligate

specialists (i.e., only infects on one host and zero or near-zero pathogenicity on the other host). The

dominance and turnover of genotypes through time have been as intriguing to investigators as it was

impeding to managers (Fry *et al.,* 2015). The global blight populations have undergone several

remarkable changes throughout their history, from pan-globally dominant lineages (FAM-1 followed

by US-1) to the diversity of genotypes seen today (reviewed in Yoshida *et al.,* 2013; Saville *et al.,*

2016). Adaptive processes within the pathogen and environmental conditions underlie the severity of

the impacts on each host (Sicard *et al.,* 2007; Fry *et al.,* 2015). The fluxes in the global blight

populations are likely explained by variation in the effects of pathogens on their hosts (Pariaud *et al.,*

2009). Therefore, a better understanding of how *P. infestans* adapt and specialise to potato and tomato,

and the determinants of pathogenicity is needed to effectively predict the emergence of disease.

Phytophthora infestans on potato and tomato hosts also presents a great opportunity to test whether

generalism tends to carry fitness costs or 'no-cost' generalism prevails (Remold, 2012). In the

literature, aggressiveness is broadly defined as the negative effect that a pathogen has on the host, and

can be considered to be any quantitative component of pathogenicity that is non-specific to host

genotypes (reviewed in Pariaud *et al.* 2009). In this study, aggressiveness is defined explicitly by

lesion growth rate on host tissue. In *P. infestans* it is a life-history trait that is a key predictor of

epidemic severity, and is widely evaluated across studies (Birhman & Singh, 1995; Pariaud et al.,
2009; Montarry *et al*. 2010).

Potato and tomato are traditionally recognised as primary and secondary hosts for *P. infestans*
respectively. Previous reports often found that potato isolates have a higher preference for potato, but
tomato isolates are generalist on both hosts (Legard *et al.,* 1995; Michalska *et al.,* 2016; Kröner *et al.,*
2017). Infectivity on tomato likely evolved relatively recently during initial invasion from the centre
of origin in the 1800s, because isolates from the native range were non-pathogenic to tomato (Legard
et al., 1995). Blight was long believed to spread from potato to adjacent tomato crops, but not vice
versa (Berg, 1926; Small, 1938; Legard *et al.,* 1995; Cohen, 2002). However, more recently it has
been revealed that transmission regularly occurs from tomato to potato in relatively new strains such
as 13_A2, US-22, and US-23 (Hu *et al.,* 2012; Frost *et al.,* 2016). The disease is now generally
considered to be as impactful on tomato as it is on potato (Fry *et al.,* 2015; Kröner *et al.,* 2017).

The history of the disease is particularly thought-provoking owing in part to a long timeline of
observations on host specificity. Prior to the early 1900s blight on tomato was uncommon (Berg,
1926). Reports from the earliest cross-inoculation trials were mostly published after 1900, when
tomato blight was becoming more prevalent globally. Those reports did not mention finding host
specificity, or they found that isolates from both hosts were destructive on both hosts under lab or
field conditions (Berg, 1926; Oyarzun *et al.,* 1998). However, this changed in the following period
after 1915 when remarkable specificity was frequently noted either on both potato and tomato isolates
or in potato isolates only (including isolates from Holland, Australia, USA and the UK; Berg, 1926).
This was followed by the emergent US-1 lineage which became dominant from the 1930s. During this
period, studies on isolates from different regions showed the majority had no remarkable specificity,
but potato isolates were still more likely host specific. For example, Wilson and Gallegly (1955)
examined isolates from Scotland, the Netherlands, Canada and the USA, and reported that 8 out of 29
potato isolates were host specific versus 2 out of 16 tomato isolates. Isolates from Israel at the time
showed similar proportions (9 out of 25 potato isolates and 2 out of 25 for tomato isolates; Kedar *et*
al., 1959). Other cross-inoculation studies conducted during US-1 dominance in Japan (Kishi, 1962),
the USA (before 1970, Flier *et al.,* 2003), and Germany (Günther *et al.,* 1970) also reported little or
no specificity. This period was followed by the emergence of an array of successive lineages from the
mid- to late-1980s up to the present where US-1 becomes either extinct or rare relative to other strains.
Studies on surviving (or remaining) US-1 isolates overwhelmingly found host specificity in isolates
from either host species within a given site, and mostly in tomato isolates, but not potato isolates
(Erselius *et al.,* 1997; Oyazun *et al.,* 1998; Vega-Sanchez *et al.,* 2000; Suassuna *et al.,* 2004; Chen *et*
al., 2009). In North America this pattern is reversed where the remaining US-1 potato isolates tend to
be host specific, but not the tomato isolates (Goodwin *et al.,* 1995; Legard *et al.,* 1995; Platt, 1999).

Other more recently emerged strains may or may not be host specific, such as US-8 and BR-1

primarily pathogenic on potato (Legard *et al.,* 1995; Suassuna *et al.,* 2004), US-21 was host specific

on tomato (Hu *et al.,* 2012), and EC-1 has low specificity (Oliva *et al.,* 2010). Importantly, host

specificity is a population-level trait so there may be variation within a lineage. Different populations

of the same lineage can also inhabit the two hosts at the same sites where their host specificity may

vary (e.g. Erselius *et al.,* 1997).

On the one hand, *P. infestans* strains that are more aggressive with a greater lesion size and higher

sporulation on their host should become dominant (Gisi *et al.,* 2011). For instance, the greater

aggressiveness of US-8 (in terms of area under lesion expansion curve) is likely an important factor in

the replacement of US-1 in south-western Canada (Miller *et al.,* 1998), and similarly the more

aggressive 2_A1 displaced the less aggressive US-1 in east Africa (Njoroge *et al.,* 2018). In Brazil,

the more aggressive BR-1 with much faster lesion growth on potato (but less aggressive on tomato)

displaced US-1 from potato only, (Suassuna *et al.,* 2004). The 13_A2 lineage is dominant in several

continents and is more aggressive than other lineages, and was demonstrated to competitively exclude

other lineages in the field, which likely explains its quick expansion across Europe within 3 years

(Cooke *et al.,* 2012). On the other hand, strains that can infect a wider range of hosts, including

resistant hosts should also be favoured (Seidl Johnson & Gevens, 2014). The 13_A2 lineage was also

able to infect previously resistant potato cultivars (Cooke *et al.,* 2012). In Taiwan, the US-11 is

aggressive on both potato and tomato and rapidly displaced a potato specific US-1 (Chen *et al.,* 2009).

The generalist US-6 and US-7 lineages similarly replaced the potato specific US-1 in North America

(Hwang *et al.,* 2014). The question then remains: why aren't pathogen populations all explicitly both

highly aggressive and generalist across hosts?

The relationship between pathogen pathogenicity components and fitness is complex. Although there

is abundant evidence that greater aggressiveness is advantageous, conversely there have been a few

suggestions that the most pathogenic strains are not most transmissible because hosts are weakened or

killed too quickly (Pasco *et al.,* 2016; Mariette *et al.,* 2016). This is in line with the 'virulence-

transmission trade-off' hypothesis which posits that high host mortality limits transmission rate (note

that the "virulence" term was used to mean quantitative pathogenicity on the host) (Acevedo *et al.,*

2019). However, the effects of relative population size (of both pathogens and hosts) are often not

considered. In the case of *P. infestans* this hypothesis also does not consider the effects of alternate

host habitats, such as tomato plants or wild *Solanum* species (Frost *et al.,* 2016). For instance, a strain

that is extremely pathogenic on potato can have lower aggressiveness on tomato to compensate the

assumed lowered transmission rate on potato if the other host is locally available. Even if high

pathogenicity is indeed linked with lower transmissibility, a trade-off between a generalist strategy

(among potato and tomato hosts) versus aggressiveness could explain the variation in host specificity

among *P. infestans* potato and tomato strains (Thrall & Burdon, 2003; Pariaud *et al.,* 2009). Such a

trade-off would be in line with generalists being 'the jack of all trades but a master of none' and

specialists being 'the master of some' (Remold, 2012). Conversely, it is also evident that strains

highly aggressive on both hosts exist (i.e., 'the master of all' strategy), including some invasive

isolates of US-11 (Chen *et al.,* 2009) and 13_A2 (Chowdappa *et al.,* 2015). Thus our current

understanding of the relationship between aggressiveness and host specificity remains unclear.

A range of cross-inoculation trials on *P. infestans* from potato and tomato have been conducted over

the past few decades. Meta-analytic approaches can be used to synthesise the evidence to uncover

patterns or processes across studies in plant pathology (Madden & Paul, 2011). A quantitative

synthesis of this global data is likely to enable us to draw general conclusions on the relationship

between host specificity and aggressiveness. Also, given the historical population changes associated

with potential major host specificity shifts, a data synthesis can reveal how host specificity is

changing over time. Selection for strains virulent on both hosts can decrease overall host specificity

over time. Hence, the following hypotheses were tested: (1) there will be a positive association

between host specificity and aggressiveness on the original host (as defined by lesion growth rate), (2)

potato isolates will overall have greater host specificity than tomato isolates, (3) host specificity will

tend to reduce over time, and (4) isolates of the older US-1 lineage will have lower aggressiveness but

higher specificity than those of successive lineages.

Methods

Data collation

To identify studies performing cross-inoculations using potato and tomato isolates of *P. infestans*, I

searched ISI Web of Science (Clarivate) and Google Scholar (https://scholar.google.com) in January

2022 using the following terms: 'Phytophthora infestans' AND 'tomato' OR 'potato'. Only trials

conducted on potato and tomato hosts were included. Studies that tested pathogenicity on only one out

of these two hosts were excluded, as well as studies that did not report the host of origin for isolates

(or the host is not determinable based on the isolate information). As such, isolates were classified

either as potato isolates or tomato isolates based on the host of origin. Studies must describe some

quantitative measure of pathogenicity (e.g., lesion size, AUDPC, degree of infection). Thus, those

reporting only qualitative pathogenicity are excluded. Studies that tested multiple isolates were pooled

to one data point (for each host of origin) to avoid pseudoreplication.

From each study, the following were recorded: (1) the host of origin of the isolates (potato or tomato).

Data of isolates from the same host species were pooled; (2) a quantitative measurement of pathogen

pathogenicity on each of potato and tomato under controlled conditions and variance. Most commonly

these are from the inoculation of plant parts or whole plants (e.g., mortality, sporulation rate, lesion

area). Semi-quantitative metrics such as disease severity scores were also accepted as host specificity

was analysed using non-parametric tests. Where more than one measure of aggressiveness was

reported, lesion size on host tissue (converted to lesion growth rate per day) was preferred over other

measures such as AUDPC and plant mortality; (3) original geographical location of the isolate; and (4)

time of isolation from hosts (calendar year). The average isolation year was recorded if data from

multiple isolates were pooled.

The differences in pathogenicity of isolates on the two hosts were standardized using effect size as

$\log_{10}$ response ratio (LRR). Highly negative LRR values (< -1) are associated with host specificity

(i.e., non-pathogenic on the other host). Conversely, a highly positive LRR (> 1) indicates the

pathogen is only aggressive on the other host and therefore the host of origin is likely a sink habitat

(such as the US-8 isolate from tomato in Legard *et al.,* 1995). Lesion growth rate (hereafter 'LGR' in

millimetres expansion per day) was estimated from the lesion size data in the time following

inoculation. The standardized LGR is used to compare the pathogen aggressiveness on the two hosts

across studies.

In addition, I compared the host specificity and aggressiveness of US-1 isolates and other lineages.

From all the studies collected from the literature search, studies that tested US-1 isolates were

identified. Part of these studies also tested isolates that are not US-1 along with other isolates (which

were pooled with US-1 in the main analyses). Those were un-pooled and entered as separate entries

comprising of 'US-1' and 'Other' lineage categories. By limiting the analysis on studies that tested

both US-1 and successive lineages within the same experiment, it allows for a more direct comparison.

Successive lineages are predicted to be more aggressive than US-1 (e.g. Legard *et al.,* 1995) and more

generalist across both hosts (e.g. Chen *et al.,* 2009; Njoroge *et al.,* 2016).

Statistical analyses

For the LRR data, a non-parametric bootstrap approach was used to estimate the mean effect sizes and

confidence intervals (95% CI). Bootstrapping is based on the method used in Van den Noortgate and

Onghena (2005) and involves randomly subsampling with replacement over 1000 iterations. This

approach is more suited to cases where many pathogenicity trials did not report the variance (as

compared to a classical meta-analysis approach). Non-parametric bootstrapping does not emphasise

thresholds for statistical significance, but emphasises effect size and confidence intervals (Rillig *et al.,*

2019). Normality in the lesion growth data was checked using quantile-quantile plots (Q-Q plots),

which showed that the distribution of the data was normal (Fig. S1). To assess whether high

specificity is associated with greater aggressiveness, the relationship between LGR on the original

host and the difference with LGR on other host was analysed using linear regressions. The LGR of

US-1 versus other lineages in subset data was analysed using t-tests. The time and LRR data were not

normally distributed, so median-based linear models (mblm) were used to test the relationships

between LRR and time (year of isolation), and between LRR and aggressiveness on the original host.

All analyses were conducted using R (v.3.6.1, R Core Development Team 2019). Bootstrapping was

conducted using the *boot* package (v. 1.3-22) and the *metaphor* package (v. 2.1-0) was used to

calculate effect sizes. The *mblm* package (v.0.12.1) was used to run the median-based models.

Results

This search included 78 host specificity comparisons from 44 studies (Fig. 1). Most studies (32

studies) conducted cross-inoculation trials on both potato and tomato isolates of *P. infestans* within

the same trial. There were a total of 44 and 34 data points for potato and tomato isolates respectively.

The isolates are from all over the globe (Fig. 2), and all inoculation trials were conducted in controlled

conditions. Of the 78 data points, 43 reported lesion size (or growth) that was usable for the

aggressiveness data. Ten of the studies tested US-1 isolates along with isolates of more successively

emergent lineages (including BR-1, EC-1, US-6, US-7, US-8, US-11, and other unnamed isolates of

A2 mating type). An outlying data point with LRR value of greater than 1 (i.e., non-pathogenic on the

original host but highly pathogenic on the other host) was removed from all analyses (shown in Fig.

3).

For the host specificity data (LRR), the vast majority of pathogenicity traits were directly related to

lesions (63 out of 78 cases, e.g., lesion size, growth, percentage, or number), and 5 cases were based

on the evaluation of lesions (proportion of plants or leaves with lesions or disease ratings based on the

type of lesions – sporulating, expanding or non-expanding). The remainder (10 cases) include

sporulation (density or proportion of lesions with sporulation) and overall disease severity. The

aggressiveness (lesion growth rate) data were entirely from detached leaf assays. The inoculation

approach used was mostly detached leaf assays (64 out of 78 cases, with one combining data from

detached stem and leaf inoculations), 4 cases were of leaf inoculation *in planta*, 10 cases used spray

inoculation and a single case was of tuber and stem inoculation. Data sources and references are

provided in the Supplementary Material (Table S1). The isolation time of pathogen isolates from

plants in the field ranged from 1970 to 2017.

Potato and tomato isolates generally had a modest but statistically significant specificities for their

host of origin (LRR estimate -0.26 [-0.09, -0.52] and -0.18 [-0.01, -0.41] 95% bootstrap CI for potato

and tomato isolates respectively, $P < 0.05$); that is, pathogenicity on the other host is overall reduced

to ~55% and ~66% of that on the original host for potato and tomato isolates, respectively. The

specificity difference between potato (LRR -0.26) versus tomato isolates (LRR -0.18) was not

significant ($P = 0.81$). The host specificity data from each region is summarised in Fig. 3. From the

LGR data, there was a significant positive relationship between aggressiveness on the original host

and the difference in aggressiveness among hosts for potato ($P < 0.001$), but not for tomato, although

this relationship is also strongly positive ($P = 0.07$, Table 1; Fig. 4). This overall result is also

supported in the omnibus analysis between host specificity (LRR) versus aggressiveness (LGR) on the

original host ($P < 0.05$, Table 2, Fig. 5). These together indicate that host specificity is associated with

greater aggressiveness, and overall the most host specific strains may be up to 3-5 times more

aggressive than generalist strains (Fig. 4). There was no significant relationship between LRR and

isolation time for potato isolates ($P = 0.54$), but for tomato isolates there was a significant positive

relationship ($P < 0.01$, Table 3; Fig. 6).

Focussing on 'surviving' isolates of the old US-1 lineage versus other successively emerged lineages,

the sample size available for analysis was relatively low ($k = 11$ and $k = 10$ for US-1 and other isolates

respectively) and fewer reporting lesion growth rates ($k = 7$ and $k = 6$ respectively). Two US-1 tomato

isolates had extreme specificity for potato and was non-pathogenic on its original host which were

removed from the analyses (LRR 2.31 and 1.58). There were no significant differences in LRR among

isolates of US-1 and other lineages (LRR -0.22 vs. -0.30 for US-1 and other lineages respectively; t =

-0.30, df = 15.94, $P = 0.77$). Furthermore, there was no significant difference in aggressiveness on the

host of origin among US-1 and other successively emergent lineages (LGR 4.12 vs. 3.47 mm per day

respectively; t = 0.54, df = 2.84, $P = 0.63$). The low availability of data between US-1 and new

isolates means that the results for this comparison should be taken with caution. Nonetheless as the

analyses pertains to a subset of studies that assessed US-1 and other isolates in tandem, it is indicated

that the remaining US-1 isolates are not necessarily less aggressive than those of other lineages.

Discussion

The current synthesis of the global data suggests a generalist strategy tends to be costly to

aggressiveness, which can help explain the persistence of host specific strains in the presence of

generalist strains even when both hosts are often abundantly available. *Phytophthora infestans* is a

hemibiotrophic pathogen with a clear necrotrophic phase, so aggressiveness towards hosts should

generally be under selection in the absence of trade-offs (Jarosz & Davelos, 1995; Montarry *et al.,*

2007). Aggressive strains often succeed less aggressive ones (e.g. Miller *et al.,* 1998; Cooke *et al.,*

2012), therefore a lower aggressiveness is likely a fitness cost for the ability to effectively infect both

hosts. This finding is consistent with the first hypothesis that host specific *P. infestans* strains will be

more aggressive than generalist strains. Several factors could potentially explain why generalist

strains tend to be lower in aggressiveness. First, evolutionary theory predicts that specialisation may

be associated with increased fitness in that environment, so naturally specialists would be fitter

regardless of any fitness costs associated with generalism (i.e. 'the jack of all trades' strategy; Remold,

2012). Second, genetic change involved in adaptation to the other host environment may involve

fitness costs due to antagonistic pleiotropy. This includes costs associated with virulence genes to

overcome host resistance factors (Pariaud *et al.,* 2009; Montarry *et al.,* 2007, 2010). Third, specialists
should have a lower accumulation of deleterious alleles (Whitlock, 1996), which in theory can hinder
the competitiveness of a generalist in the presence of specialists (Kawecki, 1994). To my knowledge
this has not yet been investigated in blight. Yang et al. (2016) found no evidence that mutation
accumulation was involved in a trade-off between thermal tolerance and pathogen growth rate in *P.*
infestans, but may be explained by antagonistic pleiotropy. Virulence is traditionally defined as the
ability of a pathogen to overcome host defences, especially in the gene-for-gene (GFG) model so that
high virulence is associated with the ability to infect more hosts (Laine & Barrès, 2013). This finding
is in line with the classic trade-off between virulence and aggressiveness in the GFG model within a
given host species (Thrall & Burdon, 2003).
302
This trade-off between generalism and aggressiveness however does not account for the presence of
generalist strains that are equally aggressive as specialist strains, or the co-occurrence of specialists
and generalist strains (even where strains are of the same lineage, Dey *et al.,* 2018). The synthesis
results showed that a portion of generalists (both potato and tomato isolates) has high aggressiveness
equal to the most host specific strains (Fig. 4). The underrepresentation of aggressive generalists is
unexpected since populations that cause severe disease (such as 13_A2) often quickly becomes
prevalent over a very wide area, so more trials should have been conducted on such populations as
those become more common. There are a number of possibilities for this. Migration limitations and
stochasticity can limit aggressive generalists, where they are restricted to some continents due to the
lack of an introduction event (e.g. quarantine measures). Another possible explanation is the epistatic
pleiotropy model where alleles can be conditionally beneficial (Remold, 2012). Under this model,
generalists with high relative fitness can arise in a population consisting of normal (less fit) generalists
or specialists through sexual recombination or selection on cryptic variation. This would require some
level of mutation or sexual reproduction. Subsequent compensatory selection against genetic costs of
generalism may also recover the lost aggressiveness, as demonstrated in other systems such as insects
and fungi (Pariaud *et al.,* 2009). However, the genetic mechanisms for how highly aggressive
generalists arise remain largely unknown.
320
Even though lesion size is regularly measured to gauge the pathogenicity of isolates and the severity
of outbreaks (e.g. Miller *et al.* 1998, Pariaud *et al.,* 2009), results from single controlled laboratory
trials may be misleading on the actual fitness of pathogen populations across seasons (Andrivon *et al.*
2013). Fitness is related to a suite of factors other than aggressiveness, including environmental and
intrinsic factors such as host availability, population size, infection efficiency, competition, fungicide
use and seasonal variation. Long-term studies taking account of the complete life cycle of the
pathogen conducted over multiple seasons are likely needed to more reliably assess fitness (Andrivon

et al. 2013). Furthermore, studies on the competitive interactions between co-occurring genotypes (e.g.

Lebreton et al. 1999) should also shed more light on this problem.

There are indications that aggressive generalists can co-occur with other strains, including specialists

under specific conditions. For instance, 13_A2 and 23_A1 (genetically similar to US-23) co-occur in

northern Africa (Beninal *et al.,* 2022) and Pakistan (Belkhiter *et al.,* 2019) where both crops are

grown in proximity almost year round. In this case, controlled trials showed that 13_A2 was

marginally less aggressive than 23_A1 with a moderate specificity on potato, while the latter was

overall more aggressive on both hosts (Belkhiter *et al.,* 2019). Yet, the more aggressive 'generalist'

23_A1 is found predominantly on tomato. In Algeria, 23_A1 is similarly only dominant on tomato

during late-season (Belkhiter *et al.,* 2019; Beninal *et al.,* 2021). Pathogenicity trials showed that US-

23 in North America has specificity similar to that of 23_A1. Conversely, there US-23 has caused

major epidemics on both hosts (Danies *et al.,* 2013). It has also completely displaced the co-occurring

generalist US-22 and a very potato-specific US-24 (Saville & Ristaino, 2019). These together indicate

that the more restricted 23_A1 could potentially be successful, but undetected factors may be limiting

its fitness. It has been suggested that the potato specialist US-24 may persist in more northern

locations despite being less aggressive than either generalists (US-22 and US-23) due to its superior

performance under cold conditions (Danies *et al.,* 2013). The genetically similar potato-specific US-8

is almost entirely displaced, but isolated populations (mainly in the West Coast of the USA) may have

been sustained by inoculum sources at certain localities (Saville & Ristaino, 2019).

Contrary to the second hypothesis that potato isolates will be more host specific, potato and tomato

isolates do not differ significantly in specificity, but there were around twice as many highly host-

specific populations for potato than for tomato by proportion (12 out of 44 cases or 27% for potato,

versus 4 out of 34 cases or 12% with LRR less than -0.5 for tomato; Fig. 5). The reduction in host

specificity of tomato strains over the last few decades is interesting, and only partially supports the

third hypothesis since the specificity of potato isolates did not change over time. Together, these

results generally support the long-held belief that tomato isolates are more generalist than potato

isolates. Possible explanations for the reduction in specificity in tomato isolates include a much

broader resistance complex for potato than tomato (Lebreton & Andrivon, 1998; Oyarzun *et al.,* 1998;

Brouwer *et al.,* 2004) or more successful sexual reproduction in tomato isolates. For tomato isolates, a

simpler virulence spectra required to overcome resistance may allow generalism to arise more easily

than potato isolates. Rare sexual admixture can produce prospective invasive generalist genotypes

such as US-11 (Gavino *et al.,* 2000). Perhaps sexual recombination might help reduce costs of

generalism through selective compensatory mechanisms, but sexual populations in *P. infestans* are

exceedingly rare and diverse populations are only known to occur in Mexico and northern Europe

(Yuen and Andersson 2013). Moreover, in cold regions potato isolates could survive on tubers but

tomato isolates may need other hosts, so specialisation to tomato may not be under selection; although

tomato isolates could also be reintroduced from other sources (e.g., seed sources). Differences in

regions could not explain the specificity reduction however as tomato isolates from colder regions

(i.e., North America and Europe) did not seem different in specificity than warmer regions (i.e., Asia,

Africa and South America) (LRR -0.21 and -0.28 respectively). It is clear that the evolutionary forces

contributing to host specificity in *P. infestans* are poorly understood, and will require further study

before the increasing generalism in tomato isolates (but not potato) could be explained. Nevertheless,

this finding provides critical information about how the global pathogen populations are interacting

and changing with time. Pathogen transmission from tomato to potato appears likely to increase in the

future.

The question of how specialization to hosts impacts on the evolution of pathogens is an important one

because of the potential for new and unforeseen outbreaks (Dittmar *et al.,* 2016). For example, the

evolution of the sister species *P. mirabilis*, *P. ipomaea*, and *P. phaseoli* closely-related to *P. infestans*

is associated with adaptation and specialization on new host species (Raffaele *et al.,* 2010). In *P.*

infestans, complete host specialization to potato or tomato is rare, and pertains to only very few

isolates (such as certain US-8 populations). Most of the time some level of pathogenicity on the other

host is retained (Fig. 3). This may perhaps be due to the relative closeness in physiology among the

two hosts and their global metropolitan nature, ensuring genetic exchange among populations.

In the current study, lower aggressiveness or higher specificity in US-1 strains was not found. Past

studies often observed the displacement of US-1 by more aggressive strains (Legard *et al.,* 1995; Reis

et al., 2003; Suassuna *et al.,* 2004; Chen *et al.,* 2009). The non-significance difference in

aggressiveness could be due to the low number of data points available in this case. Alternatively

there is a possibility that these studies were testing extant US-1 populations that represent a subset of

survivors, which may be equally aggressive as the newer invading populations. This survivorship

effect may have also been observed in *P. cinnamomi*, where the A2 mating type is the predominant

strain that displaced the less virulent endemic A1 throughout Asia (Arentz, 2017). The remaining

extant *P. cinnamomi* A1 has similar aggressiveness (i.e. lesion size) to the A2 strains (Dudzinski *et al.,*

1993; Robin & Desprez-Loustau, 1998). Comparing tomato and potato isolates of US-1 and those of

other lineages under controlled conditions (e.g., competitiveness or pathogenicity under different

environmental conditions; Young *et al.,* 2009) should shed some light on the selection pressures

within each host species.

An overview of the historical literature on host specificity following the early 1900s (when tomato

blight was spreading worldwide) clearly shows that transitions from generalism to specialism can

occur over time. Specialists are favoured in various models, including selection for their higher

aggressiveness, mutation accumulation, and genetic fixation (i.e., under low gene flow and strong

selection, Yeaman & Whitlock, 2011). As these displacements continue and populations shift in

properties, it is important to track these changes. Controlled pathogenicity trials (such as detached leaf

assays) can be employed to monitor populations. Lesion growth data from pathogenicity trials (i.e.

inoculations on potato and tomato) could be plotted onto a Cartesian plane to compare pathogen

populations (as in Fig. 4). This approach has limitations; however, as trade-offs involving other life-

history traits that are more seldom measured (such as infection efficiency, inter-seasonal transmission

and ploidy) could also be important (Andrivon et al. 2013). Identifying common principles among

different invasions should aid in disentangling the complexity of the problem. While a trade-off

between generalism and aggressiveness may help explain the variation among pathogen populations,

host specificity likely interplays with a suite of other dependencies such as competitiveness, infection

efficiency, relative host abundance, and other environmental factors.

Conclusions

The current synthesis revealed a slight trade-off between generalism and aggressiveness across

Phytophthora infestans populations on potato and tomato. In general, host specificity was associated

with greater aggressiveness in isolates from both hosts, but a proportion of cases had high

aggressiveness on both hosts which this trade-off could not explain. Although the level of specificity

among potato and tomato isolates were not significantly different, tomato isolates tended to become

more generalist over the last few decades. Due to the variability in how pathogenicity traits relate to

fitness, long-term studies (e.g. over multiple seasons) are likely needed to better understand the trade-

offs involving pathogenicity. The mechanisms driving the evolution of host specificity are complex,

and are likely shaped by demography and other evolutionary factors.

Acknowledgements

The author expresses sincere gratitude to the researchers whose data contributed to this synthesis, and

to reviewers for their helpful comments that improved the manuscript. Special thanks go to Susan

Rutherford for her support and help proofreading the manuscript and Edward Liew for the discussions

on Phytophthora. Justin SH Wan was supported by the Jiangsu University Science Foundation Fund

(20JDG056).

References

Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P (2004) Emerging

infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers.

Trends in Ecology and Evolution 19: 535-544.

Andrivon D, Montarry J, Corbiere R, Pasco C, Glais I, Marquer B, Clément JA, Castel M, Hamelin FM (2013) The hard life of *Phytophthora infestans*: when trade-offs shape evolution in a biotrophic plant pathogen. *Plant Pathology* 62: 28-35.

Arentz F (2017) *Phytophthora cinnamomi* A1: An ancient resident of New Guinea and Australia of Gondwanan origin? *Forest Pathology* 47: e12342.

Acevedo MA, Dillemuth FP, Flick AJ, Faldyn MJ, Elderd BD (2019) Virulence-driven trade-offs in disease transmission: a meta-analysis. *Evolution* 73: 636-647.

Belkhiter S, Beninal L, Khedidji H, Krimi A, Bouznad Z (2019) Aggressiveness and host adaptation of some algerian phytophthora infestans clonal lineage on potato and tomato. *Pakistan Journal of Phytopathology* 31:147-154.

Beninal L, Bouznad Z, Corbiere R, Belkhiter S, Mabon R, Taoutaou A, Keddad A, Runno-Paurson E, Andrivon D (2022) Distribution of major clonal lineages EU_13_A2, EU_2_A1, and EU_23_A1 of *Phytophthora infestans* associated with potato late blight across crop seasons and regions in Algeria. *Plant Pathology* 71: 458-469.

Berg A (1926) *Tomato late blight and its relation to late blight of potatoes*. Bulletin 2055, Agricultural Experiment Station, West Virginia University.

Birhman RK, Singh BP (1995) Path-coefficient analyses and genetic parameters of the components of field resistance of potatoes to late blight. *Annals of Applied Biology* 127: 353-362.

Blandón-Díaz JU, Widmark AK, Hannukkala A, Andersson B, Högberg N, Yuen JE (2012) Phenotypic variation within a clonal lineage of *Phytophthora infestans* infecting both tomato and potato in Nicaragua. *Phytopathology* 102: 323-330.

Brouwer DJ, Jones ES, Clair DA (2004) QTL analysis of quantitative resistance to *Phytophthora infestans* (late blight) in tomato and comparisons with potato. *Genome* 47: 475-492.

Chen CH, Wang TC, Black L, Sheu ZM, Perez F, Deahl K (2009) Phenotypic and genotypic changes in the *Phytophthora infestans* population in Taiwan -1991 to 2006. *Journal of Phytopathology* 157: 248-255.

Chowdappa P, Nirmal Kumar BJ, Madhura S, Mohan SP, Myers KL, Fry WE, Cooke DEL (2015) Severe outbreaks of late blight on potato and tomato in South India caused by recent changes in the *Phytophthora infestans* population. *Plant Pathology* 64: 191-199.

Cohen Y (2002) Populations of *Phytophthora infestans* in Israel underwent three major genetic changes during 1983 to 2000. *Phytopathology* 92: 300-307.

Cooke DE, Cano LM, Raffaele S, Bain RA, Cooke LR, Etherington GJ, Deahl KL, Farrer RA, Gilroy EM, Goss EM, Grünwald NJ (2012) Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathogens* 8: e1002940.

Daayf F, Platt HW. 2003. Differential pathogenicity on potato and tomato of *Phytophthora infestans*
US-8 and US-11 strains isolated from potato and tomato. *Canadian Journal of Plant*
Pathology 25: 150-154.

Danies G, Small I M, Myers K, Childers R, Fry WE (2013) Phenotypic characterization of recent
clonal lineages of *Phytophthora infestans* in the United States. *Plant Disease* 97: 873-881.

Dey T, Saville A, Myers K, Tewari S, Cooke DE, Tripathy S, Fry WE, Ristaino JB, Guha Roy S.
(2018) Large sub-clonal variation in *Phytophthora infestans* from recent severe late blight
epidemics in India. *Scientific Reports* 8: 4429.

Dittmar EL, Oakley CG, Conner JK, Gould BA, Schemske DW (2016) Factors influencing the effect
size distribution of adaptive substitutions. *Proceedings of the Royal Society B: Biological*
Sciences 283: 20153065.

Drenth A, Janssen EM, Govers F (1995) Formation and survival of oospores of *Phytophthora*
infestans under natural conditions. *Plant Pathology* 44: 86-94.

Dudzinski MJ, Old KM, Gibbs RJ (1993) Pathogenic variability in Australian isolates of
Phytophthora cinnamomi. *Australian Journal of Botany* 41: 721-732.

Engering A, Hogerwerf L, Slingenbergh J (2013) Pathogen-host-environment interplay and disease
emergence. *Emerging Microbes and Infections* 2: e5.

Erselius LJ, Vega-Sánchez ME, Rodríguez AM, Bastidas O, Hohl HR, Ojiambo PS, Mukalazi J,
Vermeulen T, Fry WE, Forbes GA (1997) Host specificity of *Phytophthora infestans* on
tomato and potato in Uganda and Kenya. *CIP Program Report* 1998, 49-55.

Flier WG, Van den Bosch GB, Turkensteen LJ (2003) Epidemiological importance of *Solanum*
sisymbriifolium, *S. nigrum* and *S. dulcamara* as alternative hosts for *Phytophthora infestans*.
Plant Pathology 52: 595-603.

Frenkel O, Brewer, MT, Milgroom MG (2010) Variation in pathogenicity and aggressiveness of
Erysiphe necator from different *Vitis* spp. and geographic origins in the eastern United
States. *Phytopathology* 100: 1185-1193.

Frost KE, Seidl Johnson AC, Gevens AJ (2016) Survival of isolates of the US-22, US-23, and US-24
clonal lineages of *Phytophthora infestans* by asexual means in tomato seed at cold
temperatures. *Plant Disease* 100: 180-187.

Fry WE, Goodwin SB (1997) Re-emergence of potato and tomato late blight in the United States.
Plant Disease 81: 1349-1357.

Fry WE, Birch PR, Judelson HS, Grünwald NJ, Danies G, Everts KL, Gevens AJ, Gugino BK,
Johnson DA, Johnson SB, McGrath MT (2015) Five reasons to consider *Phytophthora*
infestans a reemerging pathogen. *Phytopathology* 105: 966-981.

Garry G, Forbes GA, Salas A, Santa Cruz M, Perez WG, Nelson RJ (2005) Genetic diversity and host
differentiation among isolates of *Phytophthora infestans* from cultivated potato and wild
Solanaceous hosts in Peru. *Plant Pathology* 54: 740-748.

Gavino PD, Smart CD, Sandrock RW, Miller JS, Hamm PB, Lee TY, Davis RM, Fry WE (2000)
Implications of sexual reproduction for *Phytophthora infestans* in the United States:
Generation of an aggressive lineage. *Plant Disease* 84:731-735.

Ghimire SR, Hyde KD, Hodgkiss IJ, Shaw DS, Liew ECY (2003) Variations in the *Phytophthora*
infestans population in Nepal as revealed by nuclear and mitochondrial DNA
polymorphisms. *Phytopathology* 93: 236–243.

Gisi U, Walder F, Resheat-Eini Z, Edel D, Sierotzki H (2011) Changes of genotype, sensitivity and
aggressiveness in *Phytophthora infestans* isolates collected in European countries in 1997,
2006 and 2007. *Journal of Phytopathology* 159: 223-232.

Goodwin SB, Sujkowski LS, Fry WE (1995) Rapid evolution of pathogenicity within clonal lineages
of the potato late blight disease fungus. *Phytopathology* 855: 669-676.

Goodwin SB, Smart CD, Sandrock RW, Deahl KL, Punja ZK, Fry WE (1998) Genetic change within
populations of *Phytophthora infestans* in the United States and Canada during 1994 to 1996:
Role of migration and recombination. *Phytopathology* 88: 939-949.

Günther E, Eggert D, Grümmer G (1970) Resistance of tomatoes to late blight (*Phytophthora*
infestans). *Annals of Applied Biology* 65: 255-262.

Hu CH, Perez FG, Donahoo R, McLeod A, Myers K, Ivors K, Secor G, Roberts PD, Deahl KL, Fry
WE, Ristaino JB (2012) Recent genotypes of *Phytophthora infestans* in the eastern United
States reveal clonal populations and reappearance of mefenoxam sensitivity. *Plant Disease*
96: 1323-1330.

Hwang YT, Wijekoon C, Kalischuk M, Johnson D, Howard R, Prüfer D, Kawchuk L (2014)
Evolution and management of the Irish potato famine pathogen Phytophthora infestans in
Canada and the United States. *American Journal of Potato Research* 91: 579-593.

Jarosz AM, Davelos AL (1995) Effects of disease in wild plant populations and the evolution of
pathogen aggressiveness. *New Phytologist* 129: 371-387.

Kato M, Mizubuti ES, Goodwin SB, Fry WE (1997) Sensitivity to protectant fungicides and
pathogenic fitness of clonal lineages of *Phytophthora infestans* in the United States.
Phytopathology 87: 973-978.

Kawecki TJ (1994) Accumulation of deleterious mutations and the evolutionary cost of being a
generalist. *The American Naturalist* 144: 833-838.

Kedar N, Rotem J, Wahl I (1959) Physiologic specialization of *Phytophthora infestans* in Israel.
Phytopathology 49: 675-679.

Kishi K (1962) Studies on the physiological specialization of *Phytophthora infestans* on tomatoes.
Japanese Journal of Phytopathology 27: 180-188.

Komsta L, Komsta ML. 2013. Package '*mblm*'. Available online: https://cran.r-
project.org/web/packages/mblm/index.html (accessed April 2022).

Kröner A, Mabon R, Corbière R, Montarry J, Andrivon D (2017) The coexistence of generalist and
specialist clonal lineages in natural populations of the Irish Famine pathogen *Phytophthora*
infestans explains local adaptation to potato and tomato. *Molecular Ecology* 26: 1891-1901.

Laine AL, Barrès B (2013) Epidemiological and evolutionary consequences of life-history trade-offs
in pathogens. *Plant Pathology* 62: 96-105.

Lebreton L, Andrivon D (1998) French isolates of *Phytophthora infestans* from potato and tomato
differ in phenotype and genotype. *European Journal of Plant Pathology* 104: 583-594.

Lebreton L, Lucas JM, Andrivon D (1999) Aggressiveness and competitive fitness of *Phytophthora*
infestans isolates collected from potato and tomato in France. *Phytopathology* 89: 679-686.

Legard DE, Lee TY, Fry WE (1995) Pathogenic specialisation in *Phytophthora infestans*:
aggressiveness on tomato. *Phytopathology* 85: 1356-1361.

Madden LV, Paul PA (2011) Meta-analysis for evidence synthesis in plant pathology: An overview.
Phytopathology 101: 16-30.

Mariette N, Mabon R, Corbiere R, Boulard F, Glais I, Marquer B, Pasco C, Montarry J, Andrivon D
(2016) Phenotypic and genotypic changes in French populations of *Phytophthora infestans*:
are invasive clones the most aggressive? *Plant Pathology* 65: 577-586.

Michalska AM, Sobkowiak S, Flis B, Zimnoch-Guzowska E (2016) Virulence and aggressiveness of
Phytophthora infestans isolates collected in Poland from potato and tomato plants identified
no strong specificity. *European Journal of Plant Pathology* 144: 325-336.

Miller JS, Johnson DA, Hamm PB (1998) Aggressiveness of isolates of *Phytophthora infestans* from
the Columbia Basin of Washington and Oregon. *Phytopathology* 88: 190-197.

Mills WR (1940) *Phytophthora infestans* on tomato. *Phytopathology* 30: 830-839.

Montarry J, Corbiere R, Lesueur S, Glais I, Andrivon D (2006) Does selection by resistant hosts
trigger local adaptation in plant-pathogen systems? *Journal of Evolutionary Biology* 19:
522-531.

Montarry J, Corbiere R, Andrivon D (2007) Is there a trade-off between aggressiveness and
overwinter survival in *Phytophthora infestans*? *Functional Ecology* 21: 603-610.

Montarry J, Hamelin FM, Glais I, Corbière R, Andrivon D (2010) Fitness costs associated with
unnecessary virulence factors and life history traits: evolutionary insights from the potato
late blight pathogen *Phytophthora infestans*. *BMC Evolutionary Biology* 10: 1-9.

Nishikawa K, Kinjo AR (2018) Mechanism of evolution by genetic assimilation: Equivalence and
independence of genetic mutation and epigenetic modulation in phenotypic expression.
Biophysical Reviews 10: 667-676.

Njoroge AW, Tusiime G, Forbes GA, Yuen JE (2016) Displacement of US-1 clonal lineage by a new
lineage of *Phytophthora infestans* on potato in Kenya and Uganda. *Plant Pathology* 65: 587-
592.

Oliva RF, Kroon LP, Chacón G, Flier WG, Ristaino JB, Forbes GA (2010) *Phytophthora andina* sp.
nov., a newly identified heterothallic pathogen of solanaceous hosts in the Andean highlands.
Plant Pathology 59: 613-625.

Oyarzun PJ, Pozo A, Ordoñez ME, Doucett K, Forbes GA (1998) Host specificity of *Phytophthora*
infestans on tomato and potato in Ecuador. *Phytopathology* 88: 265-271.

Palti J, Netzer D (1963) Development and control of *Phytophthora infestans* (Mont.) De By. under
semi-arid conditions in Israel. *Phytopathologia Mediterranea* 1: 265-274.

Pangga IB, Hanan J, Chakraborty S (2011) Pathogen dynamics in a crop canopy and their evolution
under changing climate. *Plant Pathology* 60: 70-81.

Pariaud B, Ravigné V, Halkett F, Goyeau H, Carlier J, Lannou C (2009) Aggressiveness and its role
in the adaptation of plant pathogens. *Plant Pathology* 58: 409-424.

Pasco C, Montarry J, Marquer B, Andrivon D (2016) And the nasty ones lose in the end: foliar
pathogenicity trades off with asexual transmission in the Irish famine pathogen
Phytophthora infestans. *New Phytologist* 209: 334-342.

Platt HW (1999) Response of Solanaceous cultivated plants and weed species to inoculation with A1
or A2 mating type strains of *Phytophthora infestans*. *Canadian Journal of Plant Pathology*
21: 301-307.

Raffaele S, Farrer RA, Cano LM, Studholme DJ, MacLean D, Thines M, Jiang RH, Zody MC,
Kunjeti SG, Donofrio NM *et al*. (2010) Genome evolution following host jumps in the Irish
potato famine pathogen lineage. *Science* 330: 1540-1543.

Reddick D, Crosier W (1933) Biological specialization in *Phytophthora infestans*. *American Potato*
Journal 10: 129-134.

Reis A, Smart CD, Fry WE, Maffia LA, Mizubuti ES (2003) Characterization of isolates of
Phytophthora infestans from southern and southeastern Brazil from 1998 to 2000. *Plant*
Disease 87: 896-900.

Remold S. 2012. Understanding specialism when the jack of all trades can be the master of all.
Proceedings of the Royal Society B: Biological Sciences 279: 4861–4869.

Rillig MC, Ryo M, Lehmann A, Aguilar-Trigueros CA, Buchert S, Wulf A, Iwasaki A, Roy J, Yang
G (2019) The role of multiple global change factors in driving soil functions and microbial
biodiversity. *Science* 366: 886-890.

Robin C, Desprez-Loustau M (1998) Testing variability in pathogenicity of *Phytophthora cinnamomi*.
European Journal of Plant Pathology 104: 465–475.

Rubin E, Baider A, Cohen Y (2001) *Phytophthora infestans* produces oospores in fruits and seeds of
tomato. *Phytopathology* 91: 1074-1080.

Saville AC, Martin MD, Ristaino JB (2016) Historic late blight outbreaks caused by a widespread
dominant lineage of *Phytophthora infestans* (Mont.) de Bary. *PLoS ONE* e0168381.

Saville A, Ristaino JB (2019) Genetic structure and subclonal variation of extant and recent US
617   lineages of *Phytophthora infestans*. *Phytopathology* 109: 1614-1627.
618   Seidl Johnson AC, Gevens AJ (2014) Investigating the host range of the US-22, US-23, and US-24
619   clonal lineages of *Phytophthora infestans* on solanaceous cultivated plants and weeds. *Plant
620   Disease* 98: 754-760.
621   Sicard D, Pennings PS, Grandclément C, Acosta J, Kaltz O, Shykoff JA (2007) Specialization and
622   local adaptation of a fungal parasite on two host plant species as revealed by two fitness
623   traits. *Evolution* 61: 27-41.
624   Small T (1938) The relation between potato blight and tomato blight. *Annals of Applied Biology* 25:
625   271-276.
626   Suassuna ND, Maffia LA, Mizubuti ES (2004) Aggressiveness and host specificity of Brazilian
627   isolates of *Phytophthora infestans*. *Plant Pathology* 53: 405-413.
628   Thrall PH, Burdon JJ (2003) Evolution of virulence in a plant host-pathogen metapopulation. *Science*
629   299: 1735-1737.
630   Van Den Noortgate W, Onghena P (2005) Parametric and nonparametric bootstrap methods for meta-
631   analysis. *Behavior Research Methods* 37: 11-22.
632   Vartanian VG, Endo RM (1985) Overwintering hosts, compatibility types, and races of *Phytophthora
633   infestans* on tomato in southern California. *Plant Disease* 69: 516-519.
634   Vega-Sánchez ME, Erselius LJ, Rodriguez AM, Bastidas O, Hohl HR, Ojiambo PS, Mukalazi J,
635   Vermeulen T, Fry WE, Forbes GA (2000) Host adaptation to potato and tomato within the
636   US- 1 clonal lineage of *Phytophthora infestans* in Uganda and Kenya. *Plant Pathology* 49:
637   531-539.
638   Wang X, El Hadrami A, Adam L, Daayf F (2004) US-1 and US-8 genotypes of *Phytophthora
639   infestans* differentially affect local, proximal and distal gene expression of phenylalanine
640   ammonia-lyase and 3-hydroxy, 3-methylglutaryl CoA reductase in potato leaves.
641   Physiological and Molecular *Plant Pathology* 65: 157-167.
642   Whitlock MC (1996) The red queen beats the jack-of-all-trades: the limitations on the evolution of
643   phenotypic plasticity and niche breadth. *The American Naturalist* 148: S65-S77.
644   Wilson JB, Gallegly ME (1955) The interrelationships of potato and tomato races of *Phytophthora
645   infestans*. *Phytopathology* 45: 473-476.
646   Yang LN, Zhu W, Wu EJ, Yang C, Thrall PH, Burdon JJ, Jin LP, Shang LP, Zhan J (2016) Trade-offs
647   and evolution of thermal adaptation in the Irish potato famine pathogen *Phytophthora
648   infestans*. *Molecular Ecology* 25: 4047-4058.
649   Yeaman S, Whitlock MC (2011) The genetic architecture of adaptation under migration-selection
650   balance. *Evolution* 65: 1897–1911.

Yoshida K, Schuenemann VJ, Cano LM, Pais M, Mishra B, Sharma R, Lanz C, Martin FN, Kamoun
S, Krause J, *et al.* (2013) The rise and fall of the *Phytophthora infestans* lineage that
triggered the Irish potato famine. *eLife* 2: e00731.
Young GK, Cooke LR, Kirk WW, Tumbalam P, Perez FM, Deahl KL (2009) Influence of
competition and host plant resistance on selection in *Phytophthora infestans* populations in
Michigan, USA and in Northern Ireland. *Plant Pathology* 58: 703-714.
Yuen JE, Andersson B (2013) What is the evidence for sexual reproduction of *Phytophthora infestans*
in Europe? *Plant Pathology* 62: 485-491.

Table 1

Aggressiveness as defined by lesion growth rate (LGR in millimetres per day) on host of origin versus difference in aggressiveness on the other host (defined

as LGR on host of origin minus LGR on other host). The data was collated from cross-inoculation trials conducted on potato and tomato isolates of

Phytophthora infestans ($k = 24$ and $k = 19$ respectively). Bold values indicate statistically significant effects (at $P < 0.05$).

Source	Estimate	SE	d.f.	t	*P*
Potato isolates					
Intercept	-0.72	0.55	22	-1.32	0.20
Aggressiveness host of origin	0.55	0.14		3.80	**<0.001**
Tomato isolates					
Intercept	-0.33	0.68	17	-0.49	0.63
Aggressiveness host of origin	0.36	0.18		1.94	0.07

Table 2

Aggressiveness as defined by lesion growth rate (LGR in millimetres per day) on host of origin versus effect size (LRR) for all *Phytophthora infestans* isolates ($k = 68$). Bold values indicate statistically significant effects ($P < 0.05$).

Source	Estimate	Median absolute deviation	V	*P*
Intercept	-0.00	0.21	470	0.77
LRR	-0.03	0.10	316	**0.04**

668 **Table 3**

669 Year of isolation (Calendar year) versus effect size (LRR) for *Phytophthora infestans* potato ($k = 40$) and tomato ($k = 32$) isolates. Bold values indicate

670 statistically significant effects (at $P < 0.05$).

Source	Estimate	Median absolute deviation	V	*P*
Potato isolates				
Intercept	-0.41	10.29	448	0.62
Year	0.00	0.01	328	0.54
Tomato isolates				
Intercept	-17.00	26.98	126	**0.009**
Year	0.01	0.01	402	**0.009**

Figure Captions

Fig. 1

Flow diagram of the literature search and the screening process, detailing the number of studies excluded

during screening up to the final number of studies included.

Fig. 2

World map detailing the locations of *Phytophthora infestans* potato and tomato populations tested in cross-

inoculation trials. Symbols with lighter shades represent imprecise locations specified to the general region

only. Geographically distant populations tested within the same study are joined by dotted lines.

Fig. 3

Summary median and quantiles for host specificity (effect size LRR) among potato and tomato isolates of

Phytophthora infestans across continental regions (where k is the sample size). More negative LRR values

indicate higher specificity. Very negative values (such as < -1) are associated with specialisation to the

original host. Populations with very positive values are specialised on the other host and are considered

outliers (indicated by the grey arrow). A value of zero indicates equal aggressiveness on both hosts where

there is no specificity.

Fig. 4

Relationship between aggressiveness versus the difference in aggressiveness among hosts in *Phytophthora*

infestans potato and tomato isolates. Aggressiveness is defined as lesion growth rate on host tissue (in

millimetres per day). A greater difference in aggressiveness among hosts indicates greater specificity. Potato

isolates are indicated in brown (solid line and circles) and tomato isolates are indicated in red (dashed line and

triangles).

Fig. 5

Relationship between host specificity (LRR) and aggressiveness on the original host (LGR) for potato and

tomato isolates.

Fig. 6

Relationship between host preference (LRR) versus year of isolation in *Phytophthora infestans* a) potato and b)

tomato isolates. More negative LRR values are associated with greater specificity for the host of origin than

the other host.

Supplementary material captions

Figure S1

Quantile-quantile plots for the lesion growth rate data of *Phytophthora infestans* from a) potato and b) tomato

(LGR on the original host).

Table S1

List of studies and data used in the synthesis of *Phytophthora infestans* host specificity and aggressiveness of

potato and tomato isolates.

Fig. 2

Tomato isolates

Potato isolates

Fig. 3

Fig. 5

a)

b)

Fig. 1