Genetic basis of *Arabidopsis thaliana* responses to infection by naïve and adapted isolates of turnip mosaic virus

Anamarija Butković[1,†,§], Thomas James Ellis[2,†], Rubén González[1,†,#], Benjamin Jaegle[2], Magnus Nordborg[2,*], Santiago F. Elena[1,3,*]

[1] Instituto de Biología Integrativa de Sistemas (I²SysBio), CSIC-Universitat de València, Parc Científic UV, Catedrático Agustín Escardino 9, Paterna, 46980 València, Spain,

[2] Gregor Mendel Institute (GMI), Austrian Academy of Sciences, Vienna BioCenter, Doktor-Bohr-Gasse 3, 1030 Vienna, Austria,

[3] The Santa Fe Institute, Santa Fe, 1399 Hyde Park Road, NM 87501, USA.

[†] These authors contributed equally

[§] Current address: Archaeal Virology Unit, Institut Pasteur, 75015 Paris, France.

[#] Current address: Institut de Biologie de l'Ecole Normale Supérieure, CNRS, INSERM, 75005 Paris, France.

[*] **Corresponding authors' e-mail:** magnus.nordborg@gmi.oeaw.ac.at and santiago.elena@csic.es

ORCIDs:
A.B.: 0000-0002-1435-0912
T.J.E.: 0000-0002-8511-0254
R.G.: 0000-0001-9588-1127
B.J.: 0000-0002-2679-1328
M.N.: 0000-0001-7178-9748
S.F.E.: 0000-0001-8249-5593

Summary

Plant viruses account for enormous agricultural losses worldwide, and the most effective way to combat them is to identify genetic material conferring plant resistance to these pathogens. Aiming to identify genetic associations with responses to infection, we screened a large panel of *Arabidopsis thaliana* natural inbred lines for four disease-related traits caused by infection by *A. thaliana*-naïve and -adapted isolates of the natural pathogen turnip mosaic virus (TuMV). We detected a strong, replicable association in a 1.5 Mb region on chromosome 2 with a 10-fold increase in relative risk of systemic necrosis. The region contains several plausible causal genes as well as abundant structural variation, including an insertion of a *Copia* transposon into a Toll/interleukin receptor (TIR-NBS-LRR) coding for a gene involved in defense, that could be either a driver or a consequence of the disease-resistance locus. When inoculated with TuMV, loss-of-function mutant plants of this gene exhibited different symptoms than wild-type plants, being the magnitude and sign of the difference dependent on the degree of adaptation of the viral isolate to *A. thaliana*. This increase in symptoms severity was specific for infections with the adapted isolate. Necrosis-associated alleles are found worldwide, and their distribution is consistent with a trade-off between resistance during viral outbreaks and a cost of resistance otherwise, leading to negative frequency-dependent selection.

Keywords: emerging viruses; experimental evolution; frequency-dependent selection; GWAS; *Potyvirus*; resistance genes; virus evolution

Introduction

Plant viruses represent an enormous threat to crop yields and food security (Tomlinson, 1987; Oerke, 2006; Jones, 2021). Infected plants are difficult to treat and cure, so it is of vital importance to identify genetic material that is resistant to infection (Monnot et al., 2021). In spite of this, the genetic architecture of plant responses to viral infections has received much less attention than the response to bacterial and fungal pathogens (Bartoli & Roux, 2017; Monnot et al., 2021). In agricultural settings, plants are predominantly grown as monocultures, which results in more virulent and specialized viruses that cause more detrimental effects on the host (McDonald & Stukenbrock, 2016; González et al., 2019). Through their specialization in one host species, viruses also evolve better counter defenses than their naïve counterparts (Brosseau et al., 2020).

In this study, we investigated the response of the model plant *Arabidopsis thaliana* (L.) HEYNH to its pathogen turnip mosaic virus (TuMV; species *Turnip mosaic virus*, genus *Potyvirus*, family *Potyviridae*). Potyviruses affect a wide variety of crops, especially from the families *Brassicaceae* and *Solanaceae*, and are among the most widespread crop pathogens worldwide (Revers & García, 2015). TuMV is among the most damaging of the potyviruses (Tomlinson, 1987), and also has a high incidence in wild populations of *A. thaliana* (Pagán et al., 2010). The extensive genetic resources available in *A. thaliana* make it a useful system for investigating mechanisms of viral resistance in plants. In addition, *A. thaliana* belongs to the same family (*Brassicaceae*) as agriculturally important crops such as cauliflower, cabbage, turnip, or rapeseed; thus, a large array of relevant crop viruses can infect and be studied in *A. thaliana* (Pagán et al., 2010; Ouibrahim & Caranta, 2013). Viral outbreaks are frequent in natural populations of *A. thaliana*, and there is substantial genetic variation in resistance, indicating that viral coevolution represents a meaningful selection pressure in this species (Pagán et al., 2010).

Several previous genome-wide association (GWA) studies of the response of *A. thaliana* to viral infection have been carried out. Rubio et al. (2019) used TuMV and 317 lines grown under field conditions, while Butković et al. (2021) also used TuMV but 450 lines kept in laboratory conditions. Montes et al. (2021) and Liu et al. (2022) used 156 and 496 inbred lines, respectively, to measure the response of *A. thaliana* to infection by cucumber mosaic virus under controlled conditions. These studies have

demonstrated that genetic variation for virus response exists, and that individual loci

with large effects on virus response segregate in *A. thaliana* populations.

Here, we report the results of GWA studies using two isolates of TuMV and one

of the largest sets of *A. thaliana* (1,050) inbred lines studied so far. We compare an

"ancestral" isolate of TuMV that was obtained from a calla lily (*Zantedeschia* sp.) and

was naïve to *A. thaliana* (Chen et al., 2003), to its "evolved" descendant that had been

submitted to 12 serial passages of experimental evolution on *A. thaliana* Col-0 line

(González et al., 2019). The two virus isolates differ in two non-synonymous mutations

fixed during their adaptation to Col-0 and in their infection phenotypes. The evolved

virus has mutations in amino acids T1293I (cylindrical inclusion protein; CI) and

N2039H (viral genome-linked protein; VPg) and induces stronger symptoms and faster

disease progression (Deng et al., 2015; Wu et al., 2018; González et al., 2019; Navarro

et al., 2022). We aim to identify plant genes that play a role in TuMV infection and

whether some of these genes may respond differentially to each viral isolate.

100　　**Results**

101　　**Stronger disease phenotypes in response to the evolved TuMV isolate**

We inoculated 1,050 lines of *A. thaliana* (Alonso-Blanco et al., 2016) with the ancestral

(Chen et al., 2003) or evolved (González et al., 2019) isolates of TuMV and

characterized the response to each one based on four phenotypes: (*i*) infectivity

(proportion of plants showing symptoms), (*ii*) AUDPS (area under the disease progress

stairs), a measure of disease progress (Simko & Piepho, 2012), (*iii*) severity of

symptoms on a semi-quantitative scale from 0 - 5 (Fig. 1A; Butković et al., 2020), and

(*iv*) necrosis (a binary trait reflecting presence/absence of systemic necrosis, or stage 5

of the severity-of-symptoms scale). Note that throughout the manuscript we refer to

necrosis as the most severe stage of infection: systemic necrosis that results in the death

of the plant (Fig. 1A). We do not refer to the hypersensitive response that induces local

necrotic spots and limits the spread of a pathogen. As a negative control, we inoculated

2,100 plants in the same trays as the treated plants and did not observe disease

symptoms in any case, indicating that there was no cross contamination between plants.

Patterns of correlations between traits were very similar between the two viral

isolates (off-diagonal elements of the correlation matrix between phenotypes were

largely symmetric), with strong correlations between AUDPS and infectivity, and weak
correlations between necrosis and other traits (Fig. 1B). Traits showed low to moderate
SNP heritability, with a stronger correlation between genotype and severity of
symptoms and necrosis in response to the evolved virus (Fig. 1C). This shows that
phenotyping was repeatable, and the disease traits measured show genetic variation.

Lines infected with the evolved isolate showed more severe disease symptoms
than those infected with the ancestral one (Fig. 1D-G). We assessed phenotype
differences using linear models that account for cohort effects, and calculate P-values
by permutation. On average, we found a 37% increase in AUDPS ($P < 0.001$), a 27%
increase in infectivity ($P < 0.001$), and a 23% increase in severity of symptoms ($P <$
0.001). We also found a 31% increase in necrosis, although this is not statistically
significant when cohort effects are taken into account ($P = 0.08$). On average, the
history of adaptation to Col-0 is associated with increased virulence across the panel of
A. thaliana inbred lines studied here.

Despite this overall trend, there was substantial variation in the direction of effects
between lines. Although necrosis was highly repeatable between viral isolates, we found
only moderate correlations for AUDPS, infectivity and severity of symptoms in
response to the two isolates (Fig. 1B, diagonal elements). When comparing the
responses of *A. thaliana* lines to the evolved isolate relative to the ancestral one, 53.21%
of lines exhibited increased infectivity, 23.08% showed the same level of infection, and
23.70% were less infected by the evolved virus. In terms of disease progression
(AUDPS values), the evolved viral strain performed better in 64.49% of the lines, the
same in 9.25%, and worse than the ancestral strain in 26.25%. Two lines exhibited no
necrosis in response to the evolved isolate, despite previously showing necrosis when
exposed to the ancestral virus. Conversely, 14 lines that did not display necrosis for the
ancestral virus exhibited necrosis when infected by the evolved isolate (Fig. 1I-L).
Adaptation of TuMV to Col-0 thus tends to enhance the virus performance in other lines
but does not guarantee increased infectivity across the range of *A. thaliana* lines,
pointing to a complex interaction between viral isolates and plant genotypes.
146

Figure 1. Disease phenotypes. (A) Illustration of the scale used to evaluate the severity of symptoms: (0) no symptoms or healthy plant (1) mild symptoms without chlorosis (2) visible chlorosis, (3) advanced chlorosis, (4) strong chlorosis and incipient necrosis, (5) necrosis and death of the plant. (B) Correlation matrix between disease phenotypes in response to the ancestral (upper left) and evolved (lower right) isolates. The diagonal shows correlations between the same phenotype in response to each viral isolate. (C) SNP heritability for each trait in response to each virus. (D-G) Disease phenotypes across lines in response to the ancestral (red) and evolved (blue) TuMV isolates. (I-K) Differences in disease phenotypes in response to each viral isolate (evolved minus ancestral). (L) Number of lines showing necrosis in response to the ancestral or/and evolved isolates; 1 indicates the presence of necrosis. For clarity, the 997 lines showing no necrosis for either isolate (0/0) are not shown.

149   **A major locus associated with necrosis and severity of symptoms**

We used the multi-trait GWA analysis implemented in the software package LIMIX to identify individual genetic loci that correlate with the response to each TuMV isolate (Lippert et al., 2014). LIMIX assesses the response to each viral isolate jointly, and identifies loci associated with (*i*) a shared response to both isolates and (*ii*) specific responses to individual viral isolates.

We found a strong peak of association with the common response to both isolates via both severity of symptoms and necrosis on chromosome 2 (Fig. 2A-B). 56% and 49% of the lines showing necrosis in response to the ancestral and evolved lineages, respectively, harbor the minor (susceptible) allele at the most strongly associated SNP at position 5,927,469. Lines with the minor allele showed a 1.83-fold increase in severity of symptoms in response to the ancestral virus and a 1.75-fold increase in response to the evolved virus (Fig. 2E). Moreover, the region contains five strong candidate genes (Table S1). In particular, the most strongly associated SNP for both phenotypes is found inside the annotated TIR-LRR-NBS disease-resistance gene *AT2G14080*. This region shows a marked association with disease phenotypes, and contains multiple plausible candidate genes [*e.g.*, *AGD2-LIKE DEFENSE RESPONSE PROTEIN 1* (*ALD1*), involved in systemic acquired resistance, and *DYNAMIN RELATED PROTEIN 3B* (*DRP3B*) involved in membranes organization], making it a strong candidate for variation in disease symptoms.

We used two approaches to validate this association. First, we repeated the experimental procedure using 51 lines that showed systemic necrosis in the initial experiment and 65 that did not, and recovered the association in the same region on chromosome 2, with a peak at position 5,927,469 (Fig. S1). Second, we compared disease symptoms in the Col-0 genotype to those in T-DNA mutants for two candidate loci: *AT2G14080* and *DRP3B* (Table S1; Fig. 2F). Consistent with the evolved isolate being adapted to Col-0, there was a significant increase in the severity of symptoms in Col-0 in response to the evolved isolate ($U = 15.5$, $P = 0.007$). Mutant *at2g14080* showed significantly increased severity of symptoms relative to Col-0 in response to the ancestral isolate ($U = 9$, $P = 0.001$), but no significant difference from Col-0 in response to the evolved one ($U = 49$, $P = 0.971$), suggesting and antiviral role for this gene that has been overcome by the evolved isolate. In contrast, *drp3b* plants showed a significant difference from Col-0 in response to the evolved isolate ($U = 20.5$, $P = 0.023$), but not the ancestral isolate ($U = 47.5$, $P = 0.853$), suggesting a potential proviral role for this gene gained by the evolved isolate. The major association with disease phenotypes is

184   readily repeatable, and two candidate genes in the region show a significant difference

185   in severity of symptoms from Col-0.

Figure 2. Genetic associations with severity of symptoms and systemic necrosis. (A, B) associations with a common response to both viral isolates. (C, D) associations with isolate-specific responses. SNPs in red indicate markers with $-\log_{10}P$-values above the Bonferroni-corrected significance threshold. (E) Severity of symptoms in response to each viral isolate for accessions with the major

and minor alleles at the most strongly associated SNP. (F) Severity of symptoms in response to each virus for plants of T-DNA knockout mutants for two major candidate genes and Col-0 wild-type controls. (G) Severity of symptoms in response to each virus for lines with and without a TE insertion in *AT2G14080*.

Additional associations with necrosis

In addition to the major association with severity of symptoms and necrosis we
identified 13 loci associated with necrosis (Fig. 2; Table S1). Six of these associations
were common responses to both viral isolates and seven with isolate-specific responses.
These loci include the annotated disease resistance genes *RECOGNITION OF*
PERONOSPORA PARASITICA 7 (*RPP7*; encoding an LRR and NB-ARC domains-
containing protein), *BAK1-INTERACTING RECEPTOR-LIKE KINASE 2* (*BIR2*) and
AT1G61100, which are strong candidates for an involvement in the response to TuMV.
However, minor-allele frequencies at all but one of these loci are below 5%, and the
presence of the strong association on chromosome 2 may cause false-positive
associations at other loci. We therefore repeated the GWA analysis, including the
genotype at the most strongly associated SNP as a covariate (Segura et al., 2012). This
revealed two loci with a significant association with the common response to both
viruses, and eight loci with a virus-specific response (Fig. S2 and Table S2). All but one
(chromosome 4 position 273,465) of the isolate-specific associations overlapped
between the analyses with and without the major association on chromosome 2.
Moreover, linkage disequilibrium between these loci is weak (Fig. S3), suggesting that
associations are not due to chance long-range correlations. Notably, *AT1G61100*
encodes a TIR class disease resistance gene and is detected both with and without the
major association as a cofactor.

We did not identify any significant associations with AUDPS or infectivity (Fig.
S4), consistent with the low heritability for these traits (< 30%; Fig. 1C).

A *Copia* element insertion in the primary candidate gene

We explored haplotypes in the region associated with necrosis in 157 genomes
assembled *de novo* from long-read PacBio sequencing data as part of a different study
(https://1001genomes.org/). We aligned the sequences in the region around the strong
association for necrosis and severity of symptoms. To assess patterns of synteny, we
looked for homology with the two most strongly associated genes (*AT2G14080* and

DRP3B) and the eight annotated genes or transposable elements from the TAIR10

genome annotation on either side of these genes using BLAST. This analysis revealed

many structural polymorphisms (Fig. 3), including abundant intergenic indel variation,

duplications and a large presence/absence polymorphism downstream of *AT2G14080*,

containing gene *CYTOCHROME P450 FAMILY 705 SUBFAMILY A POLYPEPTIDE*

13 (*CYP705A13*) and a block of transposable elements. At least one line (9470) shows a

large-scale inversion for the entire region. The region around the strong association for

necrosis and severity of symptoms appears to be a hotspot for structural variation.

Figure 3. Multiple sequence alignment around the major association with severity of symptoms and necrosis on chromosome 2. (A) An enlarged view of the peak of association with necrosis in Fig. 2B. (B) A summary of structural variation in assembled genomes. Transposable elements are shown in orange and coding genes in grey, with candidate genes *AT2G14080* and *DRP3B* highlighted in purple and pink respectively. Boxes on the right indicate necrosis (red) in response to either virus or

no necrosis (grey). For clarity, only half of the non-necrotic lines are plotted.

We next looked for structural polymorphism that might be causal for the disease

phenotypes. Thirteen inbred lines harbored a *Copia*-element inside the first intron of

AT2G14080. We tried to genotype the presence/absence of this element in the full

sample using short-read sequencing data, but this turned out to be unachievable due to

the repetitiveness of the sequence (see Materials and methods). Among those lines for

which PacBio genomes are available, there was a marked increase in severity of

symptoms for lines with the TE insert (Fig. 2G). Nine and ten lines showed necrosis in

response to the ancestral and evolved viral isolates respectively, corresponding to a 23-

and 21- folder increase in risk of necrosis for lines with this insertion. This element

remains a promising candidate for a causal polymorphism.

The minor allele at the major association is geographically

overdispersed

The minor allele at the major association with necrosis on chromosome 2, associated

with increased disease symptoms, is spread throughout the global range of the GWA

panel (Fig. 4A). This observation is curious, because we would expect selection to limit

the spread of deleterious mutations. To investigate whether this broad spatial

distribution could be due to chance, we quantified the mean geographic distance

between lines with the minor allele at the most strongly associated SNP at major

association on chromosome 2, and at each of the top SNPs identified by GWA that were

significant after accounting for the major association as a cofactor (Table S2, Fig. 2, and

Fig. S2). We then compared these distances to the mean distances between lines with

the minor allele at 10,000 randomly chosen loci with similar minor allele counts. For

most associated loci, distances between lines with susceptible alleles are well within the

distribution of randomly chosen loci, consistent with genetic drift allowing minor alleles

to reach modest frequency on a local spatial scale. In contrast, lines harboring the minor

alleles at the major association on chromosome 2 are further apart on average than

98.5% of randomly chosen alleles at similar frequencies (Fig. 4B). This allele is found

at a similar frequency in all but three admixture groups identified by reference (Alonso-

Blanco et al., 2016), so this overdispersion cannot be accounted for by an association

with a particular ancestry with a high dispersal rate (Fig. 4C). We found similar spatial

overdispersion for the associated loci on chromosome 4 at positions 273,465 and

7,917,355. Alleles associated with systemic necrosis at the major association seem to be

spatially overdispersed, in a way that is independent of genetic background, and that is

unlikely to be due to genetic drift alone.

Figure 4. Global distribution of necrotic alleles. (A) Global distribution of genotypes (minor/major alleles at the SNP most strongly associated with systemic necrosis), and phenotypes (whether lines showed systemic necrosis or not). The inset table shows the numbers of each genotype/phenotype; colors correspond to those on the map. (B) Frequencies of minor alleles at chromosome 2 position 5,927,469 within each admixture group. The horizontal line indicates global frequency. (C) Mean distances between pairs of loci harboring the minor allele at the 13 associated loci (red) and between minor alleles at 10,000 randomly chosen loci.

Discussion

We report results from the largest GWA study done so far aiming to identify *A. thaliana*

genes involved in the infection response to one of its natural pathogens, TuMV, using

an ancestral isolate naïve to *A. thaliana*, and a second isolate that had been

experimentally evolved on *A. thaliana* line Col-0. The comparison of the results for the
two isolates has pinpointed candidate host targets for virus adaptation.

Natural variation of disease-related traits in response to different
TuMV isolates

The majority of inbred lines infected with the evolved viral lineage showed more severe
disease symptoms compared to plants infected with the ancestral isolate (Fig. 1D - G).
Our study confirms that the adaptation to one susceptible *A. thaliana* line, in this case
Col-0, enables potyviruses to perform better in otherwise poorly susceptible lines. This
observation was first made by Lalić et al. (2010) after evolving a tobacco etch virus
isolate from tobacco in the susceptible line L*er*-0. Upon inoculation of 18 other lines,
they observed that the evolved isolate exhibited greater infectivity compared to the
ancestral one. However, using a much larger panel of lines, we also found substantial
variation in both the magnitude and direction of the response to the two viral lineages
between the inbred lines, showing that adaptation to one genotype results in a complex
response across host genotypes. Subsequent plant-viral co-evolution is thus likely to
depend on the range of host genotypes the virus encounters and, hence, the extent to
which it is able to specialize.

The increase in virulence of the evolved isolate used in this study is associated
with two non-synonymous mutations acquired during adaptation to *A. thaliana* (Navarro
et al., 2022). First, mutation T1293I was found in the CI protein that is involved in viral
genome replication and cell-to-cell movement (Deng et al., 2015), and has been shown
to interact with the photosystem I PSI-K protein in *A. thaliana* (Jiménez et al., 2006).
Second, mutation N2039H was found in the VPg protein that is involved in replication
and intracellular movement (Wu et al., 2018), and is a hub for interactions with other
viral proteins (Bosque et al., 2014; Martínez et al. 2023). Mutations in VPg have been
pervasively observed in evolution experiments of potyviruses in *A. thaliana*, in all cases
resulting in increased infectivity, virulence and viral load (Agudelo-Romero et al.,
2018; Hillung et al., 2014; González et al., 2021; Navarro et al., 2022; Melero et al.,
2023). Together, these observations indicate that genome replication and viral
movement underlie the increased virulence of the evolved virus.

The genetic architecture of TuMV resistance

We identified a 1.5 Mb region on chromosome 2 that was strongly associated with the

severity of symptoms, and accounted for approximately half the cases of systemic

necrosis. This region encompasses several genes (Table 1), two of which are especially

strong candidates for causing these phenotypes. First, *AT2G14080* encodes a known

TIR-NBS-LRR family protein. These proteins recognize and block translation of viral

transcripts in the cytoplasm, induce a hypersensitive response leading to cell death, act

as a second line of defense if the initial antiviral response fails to restrict virus

accumulation, and are involved in signaling pathways (Li et al., 2001; Meyers et al.,

2003; Bhattacharjee et al., 2009; Kobayashi et al., 2010; Marone et al., 2013; van de

Weyer et al., 2019). TIR-NBS-LRR genes are the most numerous disease-resistance

genes in the *A. thaliana* genome and are under diversifying selection to respond to as

broad a spectrum of pathogens as possible (Ellis et al., 2003). This gene appears to have

a significant role in development of systemic necrosis during infection with both TuMV

strains. Second, *DRP3B* encodes a self-assembling GTPase involved in fission and

fusion of membranes in mitochondria and peroxisomes (Fujimoto et al., 2009; Wu et al.,

2018). Its analogue, *DRP2B,* has been shown to be co-opted by TuMV to enhance

virulence, and treatment with dynamin-specific inhibitors suppresses growth by

interfering with VPg (Wu et al., 2018). As previously noted, one of the two amino-acid

differences between the ancestral and evolved TuMV isolates is in VPg (Navarro et al.,

2022). These observations make *AT2G14080* and *DRP3B* strong candidate genes for a

role in viral resistance, either independently or in tandem.

To test this possibility, the ancestral and evolved isolates were inoculated into KO

mutants *at2g14080* and *drp3b*. We observed that loss of function of *AT2G14080* led to

increased disease symptoms in response to the ancestral virus, but not the evolved virus

(Fig. 2F). This suggests that *AT2G14080* is efficient for the antiviral defense against the

ancestral but not for the evolved isolate. The evolved viral isolate may be able to

overcome the response imposed by *AT2G14080* due to the mutations acquired during its

adaptation to Col-0, most probably because changes in the VPg viral protein allow it to

evade detection by the TIR-NBS-LRR protein encoded by *AT2G14080*. In contrast, loss

of function of DRP3B decreased symptoms relative to those in Col-0 in response to the

ancestral, but not the evolved virus. This points to a possible proviral effect of the

dynamin *DRP3B*. Potyviruses are able to recruit DRP2B, a homolog of DRP3B, for the

viral replication complex (Wu et al., 2018). One explanation for the reduction in

symptoms in *drp3b* plants is that the evolved isolate acquired the ability to co-opt

DRP3B in a similar way to DRP2B, in a way that the ancestral isolate is not. Whether

these two loci act independently or in an epistatic manner should be considered as a

possibility, considering the complex interplay between viral genetic factors and

environmental cues in determining the phenotypic outcomes of viral infections

(Bomblies et al., 2007; Lalić and Elena, 2013).

Thirteen additional loci (besides the major association on chromosome 2) were

significantly associated with the common or specific responses to the two viral isolates

(Fig. S2, Table S2). The low linkage disequilibrium between these loci indicates that

they segregate independently of the major association on chromosome 2, and of each

other (Fig. S3). However, these alleles are also at very low global frequencies (Table

S3). On the one hand, this is consistent with the expectation that selection against such

deleterious alleles keeps susceptible alleles rare. On the other hand, there is a substantial

risk that alleles at such low frequency should be associated with a binary phenotype at

similar frequency just by chance. As such, caution is warranted in interpreting a

causative role for these associations without further evidence.

We found very little overlap with associations found in two previous GWA

studies of TuMV resistance in *A. thaliana*. First, Butković et al. (2021) identified

genome-wide associations for AUDPS, severity of symptoms and infectivity under very

similar experimental conditions to those used here, but using two viral isolates that

differed in their host range (one evolved as a specialist in more permissive *A. thaliana*

lines and the other evolved as a generalist able to successfully infect more resistant

lines) and fewer host lines. That analysis recovered the association with *AT2G14080* for

both isolates, and identified eleven additional loci associated with the response to one or

both viruses. These eleven loci identified were not recovered in our larger dataset. This

lack of consistency can either indicate that these associations were spurious artifacts of

a limited sample size or, alternatively, a true biological effect due to the different TuMV

isolates used by Butković et al. (2021) and in this study. Second, we did not detect any

of the genes associated with viral load or infectivity in 317 lines grown under field

conditions identified by Rubio et al. (2019), and nor did that study detect the association

on chromosome 2 reported here. In this case, the lack of overlap may be explained not

only by the differences in sample sizes, but also by the differences in traits measured,

viral isolate and environmental conditions.

The *Copia*-element insertion is a candidate causative variant

We identified a *Copia*-element polymorphism in the first intron of *AT2G14080* which is
strongly associated with severity of symptoms, and especially necrosis. Given this
strong association, and the fact that first introns often harbor regulatory elements
(Gallegos and Rose, 2017), this insertion is a strong candidate for a causative mutation.
We note also that it need not be the only causal mutation. Whether it turns out to be
casual or not, the abundance of structural variation in this region highlights the need for
restraint when trying to identify any causal polymorphisms from SNP data alone, and
more generally the limitation of reference-biased polymorphism data. Moreover, recent
studies in *Drosophila melanogaster* showed that the insertion of a transposable element
into a protein-coding sequence resulted in the acquisition of antiviral function (Brosh et
al., 2022). This highlights the potential of transposable elements to contribute to
adaptation (Lisch, 2013), in this case to response to viral challenges.

It is also interesting to consider whether structural variation is driven by selection
for diversification of resistance genes, or whether particular structural variants
themselves cause variation in disease resistance. It has been previously observed that
plant disease-resistance genes often form clusters of duplicated copies within the
genome, which would contribute to structural variation between genomes (Leister,
2004). In particular, TIR-NBS-LRR genes, of which *AT2G14080* is an example, are
known to have undergone expansion by numerous large- and small-scale duplication
and translocation events (Leister, 2004). However, we did not observe widespread
duplications of *AT2G14080* or its domains, nor evidence of synteny with other regions
of the genome that would indicate translocations, so in this case a simple transposon
insertion remains the most likely explanation.

The distribution of susceptible alleles is consistent with frequency-dependent selection

Given that potyvirus outbreaks are common in nature (Pagán et al., 2010) and
susceptibility is unquestionably deleterious, it is curious that (minor) susceptible alleles
should be at sufficiently high frequency to be detectable. There are three explanations
for this. First, selection may be sufficiently weak or the virus geographically restricted,
such that alleles can arise in one location and increase in frequency by genetic drift. If
this were true, we would expect to see a clustered geographic distribution of susceptible
alleles, reflected in under dispersion of distances between pairs of susceptible lines.

While this is true for most of the loci with small effects on necrosis, susceptible alleles

at the major association on chromosome 2 are spatially over dispersed worldwide, and

found throughout unrelated lines (Fig. 4). These patterns are difficult to explain via

genetic drift. Second, it may be that the genomic instability in the region surrounding

this association leads to a high turnover of new structural mutations that decrease viral

resistance. This is implausible because this would not generate linkage with individual

SNPs that could be detected by GWA. Moreover, we found a striking concordance

between a single *Copia*-element insertion into *AT2G14080* and necrosis, suggesting that

only a single variant is responsible for increased necrosis in this region. Neither genetic

drift nor mutation-selection balance can thus explain the persistence of susceptible

alleles at the major association for necrosis.

An alternative explanation is that susceptible alleles are beneficial in the absence

of the virus. This would cause a fitness trade-off between alleles in this region, as has

been previously demonstrated for many other pathogen-resistance systems in plants

(Bergelson et al., 2001; Todesco et al., 2010). TuMV outbreaks are common in natural

A. thaliana populations, but ephemeral (Pagán et al., 2010), causing susceptible alleles

to decrease in frequency during outbreaks, but to increase at other times. This would

lead to negative frequency-dependent selection maintaining the susceptible allele at a

frequency in proportion to the frequency of TuMV outbreaks. Despite being at low

frequency worldwide, the susceptible allele is found across the geographic and

genealogical distribution of *A. thaliana*, indicating that the polymorphism must either be

very old and has been maintained by balancing selection over long periods, or that

alleles have spread by adaptive introgression during viral outbreaks. Our data do not

allow us to distinguish between these hypotheses, nor are they mutually exclusive, but

both are consistent with a fitness trade-off at this locus. Our results indicate that the

worldwide geographic distribution of susceptible alleles is therefore consistent with

negative frequency-dependent selection maintaining the susceptible allele at low, but

non-zero, frequency.

434   Materials and methods

435   Viruses and inoculation procedure

We obtained the ancestral virus from infected *Nicotiana benthamiana* DOMIN plants

inoculated with a transcript product from a p35STunos infectious plasmid that contains

TuMV genome cDNA (GenBank line AF530055.2), corresponding to YC5 isolate from

calla lily (Chen et al., 2003). This cDNA was under the control of the cauliflower

mosaic virus 35S promoter and a NOS terminator. We obtained the evolved virus via 12

passages of experimental evolution on *A. thaliana* line Col-0 (González et al., 2019) for

full details. In both cases, we pooled symptomatic tissues, froze them with liquid N_2 and

homogenized them into a fine powder using a Mixer Mill MM400 (Retsch GmbH,

Haan, Germany). For inoculations, we mixed 100 mg of powdered liquid-nitrogen-

frozen infected tissue in 1 mL of phosphate buffer (50 mM phosphate pH 7.0, 3%

PEG6000, 10% Carborundum) and rubbed 5 μL of this mixture into three central rosette

leaves per plant. To minimize the noise that significant variations in vegetative

development could produce, we infected all the lines when they reached growth stage

3.2 - 3.5 in the Boyes et al. (2001) scale, approximately 21 days after germination.

Disease phenotypes in 1,050 accessions

We screened 1,050 lines from the Arabidopsis 1001 Genomes Project (Alonso-Blanco

et al., 2016) for disease phenotypes (Appendix S1). We grew selected lines in climatic

chambers under a long-day regime (16 hours light/8 hours darkness) with 24 ℃ day and

20 ℃ night, 45% relative humidity and light intensity of 125 μmol m^{-2} s^{-1} using a 1:3

mixture of 450 nm blue and 670 nm purple LEDs. Due to limited growth-chamber

space, we inoculated and phenotyped lines in four independent cohorts of 4,800 plants

each. Five lines in the fourth cohort that did not reach the proper size on the day of the

inoculation were inoculated four days after the other lines. We inoculated eight plants

per line with the ancestral virus and another eight with the evolved virus, as well as two

mock-inoculated plants that served as negative controls. We placed four lines inoculated

with each viral isolate and the corresponding mock-inoculated plants in the same tray

and randomized tray positions every day.

We measured four disease-related traits daily for 21 days post inoculation (dpi), at

which point infection phenotypes reached a steady plateau. We measured the proportion

of infected plants ('infectivity'), AUDPS (a measure of disease progression using the

distribution of infected plants over the 21 dpi; Simko & Piepho, 2012), severity of

symptoms (based on a scale shown in Fig. 1A), and necrosis (systemic necrosis or the

most severe stage of disease). Necrosis was considered present in a line if at least one plant showed full-leaf systemic necrosis on all leaves, as shown with number 5 in Fig. 1A.

To test for statistically significant differences in disease phenotypes in response to the two viruses we constructed linear models using the phenotype as the response variable, and viral isolate and cohort as explanatory variables. For infectivity and necrosis, we used a Binomial generalized linear model (GLM) with a logit link function. For AUDPS, we log transformed the data to improve normality, and fitted a GLM with a Gaussian link function. For severity of symptoms, we applied a cumulative link model with a logit link function using the R package "ordinal" (Christensen, 2022). Because P-value estimation is sensitive to departures from normality on the scale of the link function, we calculated a null distribution for the difference due to viral isolate by permutation. For each phenotype we randomly shuffled the response variable 1000 times and refitted the linear model. The (one-sided) P-value is then the proportion of permuted samples with a greater difference in phenotype between the two viral isolates than was found for the observed data. Since it was previously found that the evolved virus induced a stronger response in the reference line Col-0 (Butković et al., 2020), we report one-sided P-values.

Genetic associations

We used the Python package LIMIX 3.0.3 (Lippert et al., 2014) in Python 3.7.3 to perform multi-trait genome-wide association analyses of disease-related traits based on the statistical framework developed in (Korte et al., 2012; Segura et al., 2012). We used the multi-trait GWA analysis implemented in the software package LIMIX to identify individual genetic loci that correlate with the response to each virus (Lippert et al., 2014). LIMIX assesses the response to each virus jointly, and identifies loci associated with (*i*) a shared response to both viruses, and (*ii*) specific responses to individual viral isolates.

We looked for associations with the 10,709,949 SNPs published by Alonso-Blanco et al. (2016). We used a liberal minor-allele-frequency cut-off of 3% to allow us to detect rare variants associated with necrosis, since only 3.7% and 4.9% of lines showed necrosis in response to the ancestral and evolved TuMV isolates respectively, leaving 2,257,599 SNPs. We accounted for cohort effects by including these as fixed

cofactors, and for population structure by modeling the covariance between lines due to

relatedness as a random effect (Bergelson et al., 2001; Kang et al., 2008). In a second

analysis, we repeated this GWA for necrosis including the genotype at chromosome 2

position 5,927,469 as a covariate to account for the confounding effects of this locus

(Korte et al., 2012). For all GWA, we used a significance threshold of $-\log P = 7.65$,

corresponding to a Bonferroni-corrected P-value of 0.05.

We estimated narrow-sense SNP heritability using the R package "Sommer" 4.1.4

(Covarrubias-Pazaran, 2016) by regressing phenotypes onto the matrix of pairwise

relatedness between individuals (Yang et al., 2010). To estimate the variance explained

by the major association for necrosis, and to assess the sensitivity of GWA and

heritability to the presence of this locus we also repeated both GWA and heritability

estimates including genotypes at the SNP most strongly associated with necrosis as a

covariate.

Validation of the major genetic association

Replication cohort

To validate the association between necrosis and symptom severity with the candidate

region on chromosome 2 (see results), we tested an additional cohort including all 51

lines that had previously shown necrosis in response to one or both viruses together

with 67 randomly chosen non-necrotic lines from the previous screen. This

confirmation cohort was inoculated and grown in the same way as the four previous

cohorts.

Knock-out lines

We assessed disease resistance for knock-out (KO) mutants for two candidate genes

within the region associated with necrosis and severity of symptoms. *AT2G14080*

encodes for a TIR-NBS-LRR protein, and *AT2G14120* for the DRP3B protein. We

ordered T-DNA knock-out (KO) mutants from the NASC stock center

(http://arabidopsis.info/BrowsePage). We chose mutants that (*i*) were to be in the Col-0

background, (*ii*) had be T-DNA inserts that cause gene knock-outs, (*iii*) were be

homozygous, and (*iv*) had already been related to virus infection and selected SALK

homozygous line SALK_105230C (NASC ID: N656836) to study *AT2G14080* and

SALK_112233C (N667049) for *DRP3B*. We grew mutant and wild-type plants as

previously described with 10 replicates per mutant/control, as well as two replicates of

mock-inoculated controls. We visually inspected inoculated plants daily for 21 dpi

noting the number of infected plants and the severity of their symptoms. We assess the

statistical significance of differences between genotypes or viral treatment with Mann-

Whitney *U* tests.

539   Structural variation around locus *AT2G14080*

Confirming the insertion using long-read genome assemblies

We examined haplotype structure in the 225 Kb surrounding the region associated with

necrosis using data from 161 full genomes sequenced using PacBio long reads

(Arabidopsis 1001 Genomes Consortium, pers. comm.), and assembled using the

methods previously described for the 1001 Genomes Plus Project (Jaeger et al., 2023).

We extracted sequences around the peak of association, close to *AT2G14080*. We

identified the precise location of that gene, as well as eight annotated genes and

transposable elements on each side from the Araport 11 database

(https://www.araport.org/), using BLAST (Camacho et al., 2009) if they showed 70%

identity and were at least 70% of the length of the gene or transposable element in the

reference genome. When comparing whole genomes, it is not meaningful to plot SNP

positions with the coordinate system of a reference genome, so to visualize the results

we plotted haplotypes on a relative scale centered at the midpoint between *AT2G14080*

and *AT2G14120* (*DRP3B*), since these genes were largely syntenic between lines and

flanked most of the region associated with necrosis. Finally, we sorted lines based on

the distance between those two genes (Fig. 4).

To confirm the insertions observed in the necrotic/susceptible lines 351, 870,

7346 and 9102, we mapped the raw PacBio reads to the new assembly using Minimap2

(Li, 2021) with the option "-ax map-pb" and checked if any breaks coincided with the

insertion position that would suggest an artifact of the assembly. This was not the case.

Genotyping the insertion in the 1001 Genomes

We attempted to confirm the presence of the insertion in the full sample using existing

short-read sequencing data (Alonso-Blanco et al., 2016). We used multiple approaches,

all based on the flanking sequence of the insertion. First, using two different mappers,

bwa-mem (Li and Durbin, 2009) and bowtie (Langmead et al., 2009), we mapped all the

short-read data to the insertion sequence plus 5 Kb of flanking sequence (on either side)

from line 870, filtering the bam files using samtools "rmdup" (Bonfield et al., 2021) to

remove duplicates. For each line, the number of paired reads mapping to each side of

the insertion borders were extracted and counted. We assumed that the presence of such

reads would confirm the presence of the insertion, whereas lines without the insertion

would only have paired reads flanking the whole insertion. However, the mapping

quality at the flanking regions was too low for this approach to work. Many flanking

regions had no coverage from short-read sequencing (even when duplicated reads were

kept). The reasons for this are not clear, but presumably reflect high levels of

repetitiveness and high levels of polymorphism in the region. In a second approach, we

use bbmap (Bushnell et al., 2017) to look for the exact sequence at the insertion site

(±50 bp) within the raw fastq file. However, when we used the four lines with a

confirmed insertion from *de novo* long-read assemblies as a control, this approach only

worked in two cases. Again, the reasons are not clear, but the approach is clearly too

unreliable for genotyping.

Geographic distribution of major and minor necrosis alleles

We plotted the worldwide distribution of necrosis phenotypes and genotypes at the most

strongly-associated SNP using R packages "maps" and "mapdata" (Becker et al., 2018,

2021). To test whether the lines harboring minor alleles are more or less geographically

dispersed than would be expected by chance, we identified 10,000 loci with similar

minor allele counts (between 33 and 178) to the SNPs associated with necrosis (Table

S2), and the lines that harbor those alleles. We then compared the mean distance

between lines harboring the SNP associated with necrosis to the distribution of

distances between lines with the randomly chosen alleles.

Data availability

Phenotype data and code to reproduce the analyses presented here are given at GitHub

(https://github.com/ellisztamas/tumv_ms) and Zenodo (DOI: 10.5281/zenodo.7829724).

Sequence   data   are   available   from   the   1001   Genomes   Project   website

(https://1001genomes.org/; Alonso-Blanco et al., 2016). Unless otherwise stated

analyses and plotting were done in R 4.0.3 (R Core Team, 2021) under RStudio Server

Pro 1.3.1093-1 (RStudio Team, 2021).

Supporting information

Fig. S1. Manhattan plots for the association between SNPs and necrosis in the replicate dataset of 118 lines.

Fig. S2. Manhattan plots for associations with severity and symptoms and necrosis in a model conditioned on the most strongly associated SNP (5,927,469 in chromosome 2).

Fig. S3. Linkage disequilibrium between SNPs associated with necrosis.

Fig. S4. Manhattan plots for common and virus-specific associations with AUDPS and infectivity.

Table S1. Significant SNPs in the GWA analysis of the ancestral and evolved isolates.

Table S2. Conditional GWA analysis on the significant SNP on the chromosome 2.

Appendix S1. Full list of *A. thaliana* lines for the GWAS analysis. (EXCEL).

Acknowledgements

We thank Paula Agudo, Francisca de la Iglesia and Joanna Gunis for excellent technical support. Work was supported by grant PID2019-103998GB-I00 funded by MCIN/AEI/10.13039/501100011033 and Generalitat Valenciana grants GRISOLIAP/2018/005 and PROMETEO/2019/012 to SFE. RG were supported by contract BES-2016-077078 funded by MCIN/AEI/10.13039/501100011033 and "ESF investing in your future". MN was funded by ERC AdG 789037 - EPICLINES, ERA-CAPS (FWF I 3684-B25) and by the GMI.

Author contributions

MN and SFE planned and designed the research. AB and RG performed experiments. AB, TJE, RG, SFE and BJ analyzed data. AB, TJE, RG, MN, and SFE wrote the manuscript. AB, TJE and RG contributed equally.

References

Agudelo-Romero P, Carbonell P, Pérez-Amador MA, Elena SF. Virus adaptation by
manipulation of host's gene expression. PLoS ONE. 2008;3: e2397.

Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J. 1,135 Genomes reveal
the global pattern of polymorphism in *Arabidopsis thaliana*. Cell. 2016;166: 481–
491.

Arabidopsis 1001 Genomes Consortium. In preparation.

Bartoli C, Roux F. Genome-wide association studies in plant pathosystems: toward an
ecological genomics approach. Front Plant Sci. 2017;8: 763.

Becker RA, Wilks AR, Brownrigg R, Minka T, Deckmyn A. maps: draw geographical
maps. 2021. Available: https://CRAN.R-project.org/package=maps.

Becker RA, Wilks AR, Brownrigg R. mapdata: Extra map databases. 2018. Available:
https://CRAN.R-project.org/package=maps

Bergelson J, Kreitman M, Stahl EA, Tian D. Evolutionary dynamics of plant *R* genes.
Science. 2001;292: 2281–2285.

Bhattacharjee S, Zamora A, Azhar MT, Sacco MA, Lambert LH, Moffett P. Virus
resistance induced by NB-LRR proteins involves *Argonaute4*-dependent translational
control. Plant J. 2009;58: 940–951.

Bomblies K, Lempe J, Epple P, Warthmann N, Lanz C, Dangl J, *et al*. Autoimmune
response as a mechanism for Dobzhansky-Muller-type incompatibility syndrome in
plants. PLoS Biol. 2007;5: 1962–1972.

Bonfield JK, Marshall J, Danecek P, Li H, Ohan V, Whitwham A, *et al*. HTSlib: C
library for reading/writing high-throughput sequencing data. GigaScience. 2021;10:
giab007.

Bosque G, Folch-Fortuny A, Picó J, Ferrer A, Elena SF. Topology analysis and
visualization of *Potyvirus* protein-protein interaction network. BMC Syst Biol.
2014;8: 129.

Boyes DC, Zayed AM, McCaskill AJ, Hoffman NE, Davis KR. Growth stage-based
phenotypic analysis of Arabidopsis: a model for high throughput functional
genomics in plants. Plant Cell. 2001;13: 1499–1510.

Brosseau C, Bolaji A, Roussin□Léveillée C, Zhao Z, Biga S, Moffett P. Natural
variation in the Arabidopsis *AGO2* gene is associated with susceptibility to potato
virus X. New Phytol. 2020;226: 866–878.

Bushnell B, Rood J, Singer E. BBMerge-accurate paired shotgun read merging via
overlap. PLoS ONE. 2017;12: e0185056.

Butković A, González R, Cobo I, Elena SF. Adaptation of turnip mosaic potyvirus to a
specific niche reduces its genetic and environmental robustness. Virus Evol. 2020;6:
veaa041.

Butković A, González R, Rivarez MPS, Elena SF. A genome-wide association study
identifies *Arabidopsis thaliana* genes that contribute to differences in the outcome of
infection with two turnip mosaic potyvirus strains that differ in their evolutionary
history and degree of host specialization. Virus Evol. 2021;7: veab063.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J. BLAST+: architecture
and applications. BMC Bioinformatics. 2009;10: 421.

Caparrós-Martín JA, McCarthy-Suárez I, Culiáñez-Macià FA. HAD hydrolase function
unveiled by substrate screening: enzymatic characterization of Arabidopsis thaliana
subclass I phosphosugar phosphatase AtSgpp. Planta. 2013;237: 943–954.

Chen CC, Chao CH, Chen CC, Yeh SD, Tsai HT. Identification of turnip mosaic virus
isolates causing yellow stripe and spot on calla lily. Plant Dis. 2003;87: 901–905.

Covarrubias-Pazaran G. Genome-assisted prediction of quantitative traits using the R
package sommer. PLoS ONE. 2016;11: e0156744.

Christensen, R. H. B. (2022). ordinal - Regression Models for Ordinal Data. R package
version 2022.11-16. https://CRAN.R-project.org/package=ordinal.

Deng P, Wu Z, Wang A. The multifunctional protein CI of potyviruses plays interlinked
and distinct roles in viral genome replication and intercellular movement. Virol J.
2015;12: 141.

Ellis J, Dodds P, Pryor T. Structure, function and evolution of plant disease resistance
genes. Curr Opin Plant Biol. 2000;3: 278–284.

Fujimoto M, Arimura S, Mano S, Kondo M, Saito C. Arabidopsis dynamin-related
proteins DRP3A and DRP3B are functionally redundant in mitochondrial fission, but
have distinct roles in peroxisomal fission. Plant J. 2009;58: 388–400.

Gallegos JE, Rose AB. Intron DNA sequences can be more important than the proximal
promoter in determining the site of transcript initiation. Plant Cell. 2017;29: 843–
853.

Garcia-Moreno M, Järvelin AI, Castello A. Unconventional RNA-binding proteins step
into the virus-host battlefront. WIREs RNA. 2018;9: e1498.

González R, Butković A, Elena SF. Role of host genetic diversity for susceptibility-to-
infection in the evolution of virulence of a plant virus. Virus Evol. 2019;5: vez024.

González R, Butković A, Escaray FJ, Martínez-Latorre J, Melero I, Pérez-Parets E, *et*

al. Plant virus evolution under strong drought conditions results in a transition from

parasitism to mutulism. Proc Natl Acad Sci USA. 2021;118: e2020990118.

Hillung J, Cuevas JM, Valverde S, Elena SF. Experimental evolution of an emerging

plant virus in host genotypes that differ in their susceptibility to infection. Evolution.

2014;68: 2467–2680.

Jaegle B, Pisupati R, Soto-Jiménez LM, Burns R, Rabanal FA, Nordborg M. Extensive

sequence duplication in Arabidopsis revealed by pseudo-heterozygosity. Genome

Biol. 2023;24: 44.

Jiménez I, López L, Alamillo JM, Valli A, García JA. Identification of a plum pox virus

CI-interacting protein from chloroplast that has a negative effect in virus infection.

Mol Plant Microbe Interact. 2006;19: 350–358.

Jones RAC. Global plant virus disease pandemics and epidemics. Plants. 2021;10.

Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, *et al*. Efficient

control of population structure in model organism association mapping. Genetics.

2008;178: 1709–1723.

Kobayashi M, Seo S, Hirai K, Yamamoto-Katou Y, Katou S, Seto H, Meshi T,

Mitsuhara I, Ohashi Y. Silencing of WIPK and SIPK mitogen-activated protein

kinases reduces tobacco mosaic virus accumulation but permits systemic viral

movement in tobacco possessing the *N* resistance gene. Mol Plant-Microbe Interact.

2010;23: 1032–1041.

Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q. A mixed-model approach for

genome-wide association studies of correlated traits in structured populations. Nat

Genet. 2012;44: 1066–1071.

Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol. 2009;10:

R25.

Lalić J, Agudelo-Romero P, Carrasco P, Elena SF. Adaptation of tobacco etch potyvirus

to a susceptible ecotype of *Arabidopsis thaliana* capacitates it for systemic infection

of resistant ecotypes. Phil Trans R Soc B 2010; 365:1997-2007.

Lalić J, Elena SF. Epistasis between mutations is host-dependent for an RNA virus. Biol

Lett. 2013;9: 20120396.

Leister D. Tandem and segmental gene duplication and recombination in the evolution

of plant disease resistance genes. Trends Genet. 2004;20: 116–122.

Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics.
2021;37: 4572–4574.

Li H, and Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
Transform. Bioinformatics. 2009;25: 1754–1760.

Li X, Clarke JD, Zhang Y, Dong, X. Activation of an EDS1-mediated *R*-gene pathway
in the *snc1* mutant leads to constitutive, NPR1-independent pathogen resistance. Mol
Plant-Microbe Interact. 2001;14: 1131–1139.

Lippert C, Casale FP, Rakitsch B, Stegle O. LIMIX: genetic analysis of multiple traits.
bioRxiv. 2014; doi: 10.1101/003905.

Lisch D. How important are transposons for plant evolution? Nat Rev Genet. 2013;14:
49–61.

Liu S, Chen M, Li R, Li, WX, Gal-On A, Jia Z, Ding SW. Identification of positive and
negative regulators of antiviral RNA interference in *Arabidopsis thaliana*. Nat
Commun. 2022;13: 2994.

Luan H, Niu H, Luo J, Zhi H. Soybean cytochrome b5 is a restriction factor for soybean
mosaic virus. Viruses. 2019;11: 546.

Marone D, Russo M, Laidò G, De Leonardis A, Mastrangelo A. Plant nucleotide
binding site-leucine-rich repeat (NBS-LRR) genes: active guardians in host defense
responses. Int J Mol Sci. 2013;14: 7302–7326.

Martínez F, Carrasco JL, Toft C, Hillung J, Giménez-Santamarina S, Yenush L,
Rodrigo G, Elena SF. A binary interaction map between turnip mosaic virus and
Arabidopsis thaliana proteomes. Commun Biol. 2023;6: 28.

McDonald BA, Stukenbrock EH. Rapid emergence of pathogens in agro-ecosystems:
global threats to agricultural sustainability and food security. Phil Trans R Soc B.
2016;371: 20160026.

Melero I, González R, Elena SF. Host developmental stages shape the evolution of a
plant RNA virus. Phil Trans R Soc B. 2023;378: 20220005.

Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. Genome-wide analysis of
NBS-LRR–encoding genes in Arabidopsis. Plant Cell. 2003;15: 809–834.

Monnot S, Desaint H, Mary-Huard T, Moreau L, Schurdi-Levraud V, Boissot N.
Deciphering the genetic architecture of plant virus resistance by GWAS, state of the
art and potential advances. Cells. 2021;10: 3080.

Montes N, Cobos A, Gil-Valle M, Caro E, Pagán I. *Arabidopsis thaliana* genes
associated with cucumber mosaic virus virulence and their link to virus seed
transmission. Microorganisms. 2021;9: 692.

Navarro R, Ambrós S, Butković A, Carrasco JL, González R, Martínez F, *et al*. Defects
in plant immunity modulate the rates and patterns of RNA virus evolution. Virus
Evol. 2022;8: veac059.

Oerke EC. Crop losses to pests. J Agric Sci. 2006;144: 31–43.

Ouibrahim L, Caranta C. Exploitation of natural genetic diversity to study plant-virus
interactions: what can we learn from *Arabidopsis thaliana*? Natural virus resistance
in *Arabidopsis thaliana*. Mol Plant Pathol. 2013;14: 844–854.

Pagán I, Fraile A, Fernandez-Fueyo E, Montes N, Alonso-Blanco C, García-Arenal F.
Arabidopsis thaliana as a model for the study of plant–virus co-evolution. Phil Trans
R Soc B. 2010;365: 1983–1995.

R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing; 2021.

Revers F, García JA. Molecular biology of potyviruses. Adv Virus Res. 2015;92: 101–
199.

RStudio Team. RStudio: Integrated development environment for R. Boston, MA:
RStudio, PBC; 2021.

Rubio B, Cosson P, Caballero M, Revers F, Bergelson J. Genome-wide association
study reveals new loci involved in *Arabidopsis thaliana* and turnip mosaic virus
(TuMV) interactions in the field. New Phytol. 2019;224: 2026–2038.

Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, *et al*. An efficient
multi-locus mixed-model approach for genome-wide association studies in structured
populations. Nat Genet. 2012;44: 825–830.

Simko I, Piepho HP. The area under the disease progress stairs: calculation, advantage,
and application. Phytopathology. 2012;102: 381–389.

Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M. Natural allelic variation
underlying a major fitness trade-off in *Arabidopsis thaliana*. Nature. 2010;465: 632–
636.

Tomlinson JA. Epidemiology and control of virus diseases of vegetables. Ann Appl
Biol. 1987;110: 661–681.

Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, *et al*. A

species-wide inventory of NLR genes and alleles in *Arabidopsis thaliana*. Cell.

2019;178: 1260-1272.e14.

Wu G, Cui X, Chen H, Renaud JB, Yu K. Dynamin-like proteins of endocytosis in

plants are coopted by potyviruses to enhance virus infection. J Virol. 2018;92:

e01320–18.

Xu J, Wang X, Guo W. The cytochrome P450 superfamily: Key players in plant

development and defense. J Integ Agric. 2015;14: 1673–1686.

Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, *et al.* Common

SNPs explain a large proportion of the heritability for human height. Nat Genet.

2010;42: 565–569.

Table 1. Significant SNPs and their corresponding genomic regions on chromosome 2 detected for necrosis in the multiple trait analysis for both viral isolates.

SNP	Locus	$-\log_{10}P$	Function of significant genes	Reference
5,923,326	200 bp upstream of *AT2G14070*	20.55	Wound-responsive protein-like protein	-
5,928,864	*AT2G14080*	20.43	Disease-resistance protein of the TIR-NBS-LRR class lead to hypersensitive response and resistance	Meyers et al. (2003); Battacharjee et al. (2009); Marone et al. (2013); van de Weyer et al. (2019)
5,931,085	1.2 Kb upstream of *AT2G14095*	14.39	Hypothetical protein	-
5,933,529	*AT2G14095* and upstream of *AT2G14100*	8.42	A member of the cytochrome P450 family; plays roles in plant defense against pathogens and resistance.	Xu et al. (2015); Luan et al. (2019)
5,952,757	*AT2G14110*	11.32	Haloacid dehalogenase-like hydrolase (HAD) proteins are involved in phosphate homeostasis	Caparrós-Martín et al. (2013)
5,956,602	*AT2G14120*	14.62	Dynamin-related protein 3b suppresses TuMV replication and movement-	Wu et al. (2018)
5,975,892	200 bp downstream of *AT2G14160*	13.69	RNA-binding (RRM/RBD/RNP motifs) family protein; involved in inhibition of RNA virus replication, movement, and translation.	García-Moreno et al. (2018)

	Alive	**Necrotic**
Major	987	27
Minor	10	26