Genetic basis of *Arabidopsis thaliana* responses to infection with naïve and adapted isolates of turnip mosaic virus

Anamarija Butković[1,†,§], Thomas James Ellis[2,†], Rubén González[1,†,#], Benjamin Jaegle[2], Magnus Nordborg[2,*], Santiago F. Elena[1,3,*]

[1] Instituto de Biología Integrativa de Sistemas (I²SysBio), CSIC-Universitat de València, Parc Científic UV, Catedrático Agustín Escardino 9, Paterna, 46980 València, Spain,

[2] Gregor Mendel Institute (GMI), Austrian Academy of Sciences, Vienna BioCenter, Doktor-Bohr-Gasse 3, 1030 Vienna, Austria,

[3] The Santa Fe Institute, Santa Fe, 1399 Hyde Park Road, NM 87501, USA.

[†]These authors contributed equally

[§]Current address: Archaeal Virology Unit, Institut Pasteur, 75015 Paris, France.

[#]Current address: Institut de Biologie de l'Ecole Normale Supérieure, CNRS, INSERM, 75005 Paris, France.

[*]**Corresponding authors' e-mail:** magnus.nordborg@gmi.oeaw.ac.at and santiago.elena@csic.es

ORCIDs:
A.B.:   0000-0002-1435-0912
T.J.E.:   0000-0002-8511-0254
R.G.:   0000-0001-9588-1127
B.J.:   0000-0002-2679-1328
M.N.:   0000-0001-7178-9748
S.F.E.: 0000-0001-8249-5593

Summary

Plant viruses account for enormous agricultural losses worldwide, and the most effective way to combat them is to identify genetic material conferring plant resistance to these pathogens. Aiming to identify genetic associations with responses to infection, we screened a large panel of *Arabidopsis thaliana* natural inbred lines for four disease-related traits caused by infection by *A. thaliana*-naïve and -adapted isolates of the natural pathogen turnip mosaic virus (TuMV). We detected a strong, replicable association in a 1.5 Mb region on chromosome 2 with a 10-fold increase in relative risk of systemic necrosis. The region contains several plausible causal genes as well as abundant structural variation, including an insertion of a *Copia* transposon into an TIR-NBS-LRR coding gene, that could be either a driver or a consequence of the disease-resistance locus. When inoculated with TuMV, loss-of-function mutant plants of this gene exhibited different symptoms than wild-type plants, being the magnitude and sign of the difference dependent on the degree of adaptation of the viral isolate to *A. thaliana*. This increase in symptoms severity was specific for infections with the naïve isolate. Necrosis-associated alleles are found worldwide, and their distribution is consistent with a trade-off between resistance during viral outbreaks and a cost of resistance otherwise, leading to negative frequency-dependent selection.

Keywords: emerging viruses; experimental evolution; frequency-dependent selection; GWAS; *Potyvirus*; resistance genes; virus evolution

Introduction

Plant viruses represent an enormous threat to crop yields and food security (Tomlinson, 1987; Oerke, 2006; Jones, 2021). Infected plants are difficult to treat, so it is of vital importance to identify genetic material which is resistant to infection (Monnot et al., 2021). In spite of this, the genetic architecture of plant responses to viral infections has received much less attention than the response to bacterial and fungal pathogens (Bartoli & Roux, 2017; Monnot et al., 2021). In agricultural settings, plants predominantly form monocultures, thus allowing for more virulent and specialized viruses that cause more detrimental effects on the host (McDonald & Stukenbrock, 2016; González et al., 2019). Through their specialization in one host species, viruses also evolve better counter defenses than their naive counterparts (Brosseau et al., 2020).

Among plant viruses, members of *Potyviridae* family are of special interest. Potyviruses affect a wide variety of crops, especially from the families *Brassicaceae* and *Solanaceae*, and are among the most widespread crop pathogens worldwide (Revers & García, 2015). Turnip mosaic virus (TuMV species *Turnip mosaic virus*, genus *Potyvirus*, family *Potyviridae*) is among the most damaging of the potyviruses (Tomlinson, 1987), and also has a high incidence in wild populations of *Arabidopsis thaliana* (L.) Heynh (Pagán et al., 2010). *A. thaliana* is a perfect model organism for investigating the mechanisms behind viral resistance in plants because of the genetic resources available. Also, because of its near genetic resemblance to the agriculturally significant brassicas, this system permits the study of a variety of crop viruses, thereby broadening the scope of research (Pagán et al., 2010; Ouibrahim & Caranta, 2013). Viral outbreaks are frequent in natural populations of *A. thaliana*, and there is substantial genetic variation in resistance, indicating that viral coevolution represents a meaningful selection pressure in this species (Pagán et al., 2010). Several previous genome-wide association (GWA) studies of the response of *A. thaliana* to viral infection have been carried out. Rubio et al. (2019) used TuMV and 317 lines grown under field conditions, while Butković et al. (2021) also used TuMV but 450 lines kept in laboratory conditions. Montes et al. (2021) and by Liu et al. (2022) used 156 and 496 inbred lines respectively to measure the response of *A. thaliana* to infection by cucumber mosaic virus under controlled conditions. These studies have demonstrated that genetic variation for virus response exists, and that individual loci with large effects on virus response segregate in *A. thaliana*.

Here, we report the results of GWA studies using two isolates of TuMV and one of the largest (1,050) set of *A. thaliana* inbred lines studied so far. We compare an "ancestral" isolate of TuMV that was obtained from a calla lily (*Zantedeschia* sp.) and was naïve to *A. thaliana* (Chen et al., 2003)*,* to its "evolved" descendant that had been submitted to12 serial passages of experimental evolution on *A. thaliana* Col-0 line (González et al., 2019). The two virus isolates differ in their adaptation to Col-0 and their infection phenotypes, where the evolved virus induces stronger symptoms and faster disease progression. We aim to identify plant genes that play a role in TuMV infection and whether some of these genes may respond differentially to each viral isolate.

Materials and methods

Viruses and inoculation procedure

We obtained the ancestral virus from infected *Nicotiana benthamiana* DOMIN plants inoculated with a transcript product from a p35STunos infectious plasmid that contains TuMV genome cDNA (GenBank line AF530055.2), corresponding to YC5 isolate from calla lily (Chen et al., 2003). This cDNA was under the control of the cauliflower mosaic virus 35S promoter and a NOS terminator. We obtained the evolved virus via 12 passages of experimental evolution on *A. thaliana* line Col-0; see González et al. (2019) for full details. The evolved virus has two mutations (CI/T1293I and VPg/N2039H) not present in its antecessor. In both cases, symptomatic tissues were pooled, frozen with liquid N_2 and homogenized into a fine powder using a Mixer Mill MM400 (Retsch GmbH, Haan, Germany). For inoculations, we mixed 100 mg of powdered liquid-nitrogen-frozen infected tissue in 1 mL of phosphate buffer (50 mM phosphate pH 7.0, 3% PEG6000, 10% Carborundum) and rubbed 5 μL of this mixture into three central rosette leaves per plant. To minimize the noise that significant variations in vegetative development could produce, we infected all the lines at roughly the same growth stage; when they reached growth stage 3.2 - 3.5 in the Boyes et al. (2001) scale. In our growing conditions the lines used in this study reached that stage approximately 21 days after germination.

Plant growth and phenotyping

We screened a total of 1,050 lines from the Arabidopsis 1001 Genomes Project (Alonso-Blanco et al., 2016) for disease phenotypes (Appendix S1). We grew selected lines in

climatic chambers under a long-day regime (16 hours light/8 hours darkness) with 24 ºC

day and 20 ºC night, 45% relative humidity and light intensity of 125 μmol m^{-2} s^{-1} using

a 1:3 mixture of 450 nm blue and 670 nm purple LEDs. Due to limited growth-chamber

space and labor force, we inoculated and phenotyped lines in four independent cohorts of

4,800 plants each. Five lines in the fourth block that did not reach the proper size on the

day of the inoculation were inoculated four days after the other lines. We inoculated eight

plants per line with the ancestral virus and another eight with the evolved virus, as well

as two mock-inoculated plants that served as negative controls. We placed four lines

inoculated with each viral isolate and the corresponding mock-inoculated plants in the

same tray and randomized tray positions every day.

We measured four disease-related traits daily for 21 days post inoculation (dpi), at

which point infection phenotypes reached a steady plateau. We measured infectivity,

AUDPS (an analysis of the disease progression using the number of infected plants during

the 21 dpi (Simko & Piepho, 2012), severity of symptoms (based on a scale showed in

Fig. 1A), and necrosis (systemic necrosis or the most severe stage of disease). Necrosis

was considered present in a line if at least one plant showed full-leaf systemic necrosis,

as shown with number 5 in Fig. 1A.

To test for statistically significant differences in disease phenotypes in response to

the two viruses we constructed linear models using the phenotype as the response

variable, and viral isolate and block as explanatory variables. For infectivity and necrosis,

we used a Binomial generalized linear model (GLM) with a logit link function. For

AUDPS, we log transformed the data to improve normality, and fitted a GLM with a

Gaussian link function. For severity of symptoms, we applied a cumulative link model

with a logit link function using the R package "ordinal"

(https://github.com/runehaubo/ordinal). Because *P*-value estimation is sensitive to

departures from normality on the scale of the link function, we calculated a null

distribution for the difference due to viral isolate by permutation. For each phenotype we

randomly shuffled the response variable 1000 times and refitted the linear model. The

(one-sided) *P*-value is then the proportion of permuted samples with a greater difference

in phenotype between the two viral isolates than was found for the observed data. Since

it was previously found that the evolved virus induced a stronger response in the reference

line Col-0 (Butković et al., 2020), we report one-sided *P*-values.

To validate the association between necrosis and symptom severity with the

candidate region on chromosome 2, we tested an additional cohort including all 51 lines

that had previously shown necrosis in response to one or both viruses together with 67

randomly chosen non-necrotic lines from the previous screen. This confirmation cohort

was inoculated and grown in the same way as the four previous cohorts.

In addition, we selected two candidate genes within this region, *AT2G14080*

(encoding for a nucleotide-binding-site leucine-rich repeat disease-resistance protein with

an N-terminal Toll/interleukin receptor (TIR-NBS-LRR)) and *AT2G14120* (encoding for

DYNAMIN RELATED PROTEIN 3B (*DRP3B*)) and ordered T-DNA knock-out (KO)

mutants from the NASC stock center (http://arabidopsis.info/BrowsePage). Mutants

were chosen on the following criteria: (*i*) have to be in the Col-0 background, (*ii*) must

be T-DNA inserts that cause gene knock-outs, (*iii*) must be homozygous, and (*iv*) last but

not least, they must have already been related to virus infection. We used the SALK

homozygous line SALK_105230C (NASC ID: N656836) to study *AT2G14080* and

SALK_112233C (N667049) for *DRP3B*. Visual inspection of inoculated plants was done

daily for 21 dpi, annotating the number of infected plants and the severity of their

symptoms. Symptoms severity data were fitted to a generalized linear mixed model

(GLMM) with a Poisson probability distribution and a logarithmic link function. Plant

genotype and virus isolate were treated as orthogonal inter-individual factors and dpi as

intra-individual factor (repeated measures). The main effects as well as their interactions

were evaluated using type III sum of squares. These analyses were done with SPSS

version 28.0.1.0 (IBM, Armonk NY, USA).

Genetic associations

We used the Python package LIMIX 3.0.3 (Lippert et al., 2014) in Python 3.7.3 to

perform multi-trait genome-wide association analyses of disease-related traits based on

the statistical framework developed in (Korte et al., 2012; Segura et al., 2012). We looked

for associations with the 10,709,949 SNPs published by Alonso-Blanco et al. (2016). We

used a liberal minor-allele-frequency cut-off of 3% to allow us to detect rare variants

associated with necrosis, since only 3.7% and 4.9% of lines showed necrosis in response

to the ancestral and evolved viruses respectively, leaving 2,257,599 SNPs. We accounted

for cohort effects by including these as fixed cofactors, and for population structure by

modeling the covariance between lines due to relatedness as a random effect (Bergelson

et al., 2001; Kang et al., 2008). In a second analysis, we repeated this GWA for necrosis

including the genotype at chromosome 2 position 5,927,469 as a covariate to account for

the confounding effects of this locus (Korte et al., 2012). For all GWA, we used a

significance threshold of $-\log P = 7.65$, corresponding to a Bonferroni-corrected P-value

of 0.05.

We estimated narrow-sense SNP heritability (h^2_{SNP}) using the R package "Sommer"

4.1.4 (Covarrubias-Pazaran, 2016) by regressing phenotypes onto the matrix of pairwise

relatedness between individuals (Yang et al., 2010). To estimate variance explained by

the major association for necrosis, and to assess the sensitivity of GWA and heritability

to the presence of this locus we also repeated both GWA and heritability estimates

including genotypes at the SNP most strongly associated with necrosis as a covariate.

Structural variation around locus *AT2G14080*

Confirming the insertion using long-read genome assemblies

We examined haplotype structure in the 225 Kb surrounding the region associated with

necrosis using data from 53 full genomes sequenced using PacBio long reads

(Arabidopsis 1001 Genomes Consortium, pers. comm.), and assembled using the

methods previously described for the 1001 Genomes Plus Project (Koren et al., 2017).

We extracted sequences around the peak of association close to *AT2G14080*. We

identified the precise location of that gene, as well as eight annotated genes and

transposable   elements   on   each   side   from   the   Araport   11   database

(https://www.araport.org/), using BLAST (Camacho et al., 2009) if they showed 70%

percent identity and were at least 70% of the length of the gene or transposable element

in the reference genome. When comparing whole genomes, it is not meaningful to plot

SNP positions with the coordinate system of a reference genome, so to visualize the

results we plotted haplotypes on a relative scale centered between *AT2G14050* and

AT2G14120 (*DRP3B*), since these genes were largely syntenic between lines and flanked

most of the region associated with necrosis. Finally, we sorted lines based on the distance

between those two genes (Fig. 4).

To confirm the insertions observed in the necrotic/susceptible lines 351, 870, 7346

and 9,102, we mapped the raw PacBio reads to the new assembly using Minimap2 (Li,

2021) with the option "-ax map-pb" and checked if any breaks coincided with the

insertion position that would suggest an artifact of the assembly. This was not the case.

Genotyping the insertion in the 1001 Genomes

We attempted to confirm the presence of the insertion in the full sample using existing short-read sequencing data (Alonso-Blanco et al., 2016). We used multiple approaches, all based on the flanking sequence of the insertion. First, using two different mappers, bwa-mem (http://bio-bwa.sourceforge.net) as well as bowtie (Langmead et al., 2009), we mapped all the short-read data to the insertion sequence plus 5 Kb of flanking sequence (on either side) from line 870, filtering the bam files using samtools "rmdup" (Bonfield et al., 2021) to remove duplicates. For each line, the number of paired reads mapping to each side of the insertion borders were extracted and counted. We assumed that the presence of such reads would confirm the presence of the insertion, whereas lines without the insertion would only have paired reads flanking the whole insertion. However, the mapping quality at the flanking regions was too low for this approach to work. Many flanking regions had no coverage from short-read sequencing (even when duplicated reads were kept). The reasons for this are not clear, but presumably reflect high levels of repetitiveness and high levels of polymorphism in the region. In a second approach, we use bbmap (https://sourceforge.net/projects/bbmap) to look for the exact sequence at the insertion site (±50 bp) within the raw fastq file. However, when we used the four lines with a confirmed insertion from *de novo* long-read assemblies as a control, this approach only worked in two cases. Again, the reasons are not clear, but the approach is clearly too unreliable for genotyping.

Geographic distribution of major and minor necrosis alleles

We plotted the worldwide distribution of necrosis phenotypes and genotypes at the most strongly-associated SNP using R packages "maps" and "mapdata" (Becker et al., 2018, 2021). We defined "resistant" and "susceptible" genotypes at each associated SNP based on the allele most strongly associated with systemic necrosis. To test whether the lines harboring susceptible alleles are more or less geographically dispersed than would be expected by chance, we first calculated the mean distances between all pairs of lines with the susceptible alleles, and compared this to mean distances between lines with the minor allele at 10,000 randomly chosen lines with minor allele counts between 10 and 200 (minor allele counts at significant ranged from 33 to 178).

Results

Increased disease phenotypes in response to the evolved virus

We inoculated 1,050 lines of *A. thaliana* (Alonso-Blanco et al., 2016) with the ancestral (Chen et al., 2003) or evolved (González et al., 2019) isolates of TuMV and characterized the response to each virus based on four phenotypes: (1) infectivity (proportion of plants showing symptoms), (2) AUDPS (area under the disease progress stairs, a measure of disease progress (Simko & Piepho, 2012)), (3) severity of symptoms on a semi-quantitative scale from 0 - 5 (Fig. 1A) (Butković et al., 2020), and (4) necrosis (a binary trait reflecting presence/absence of systemic necrotic symptoms). Note that along the manuscript we refer to necrosis as the most severe stage of infection: systemic necrosis that results in the death of the plant (Fig. 1A). We do not refer to the hypersensitive response that induces local necrotic spots and limits the spread of a pathogen.

Lines infected with the evolved virus showed more severe disease symptoms than those infected with the ancestral virus. On average, we found a 27% increase in infectivity (Fig. 1B; $P < 0.001$), 37% increase in AUDPS (Fig. 1C; $P < 0.001$), and 23% increase in severity of symptoms (Fig. 1D; $P < 0.001$). We also found a 31% increase in necrosis (Fig. 1E), although this is not statistically significant when cohort effects are taken into account ($P = 0.08$). None of the 2,100 mock-inoculated plants showed any symptoms.

Although the severity of disease phenotypes increased in response to the evolved virus, the nature of that increase varied between phenotypes (Fig. 1B - 1E). Off-diagonal elements of the correlation matrix between phenotypes were largely symmetric, with strong correlations between AUDPS and infectivity, and weak correlations between necrosis and other traits, indicating that the overall pattern of correlations between phenotypes in response to a single virus were very similar (Fig. 1F). However, although necrosis was highly repeatable between viruses, we found only moderate correlations for AUDPS, infectivity and severity of symptoms in response to the two viruses (Fig. 1F, diagonal elements). This suggests that disease phenotypes in response to each virus varied between *A. thaliana* lines. AUDPS and infectivity showed low to moderate h^2_{SNP} for both isolates, with substantially higher heritability for necrosis (> 50%) and severity of symptoms (> 40%) in response to the evolved virus (Fig. 1G).

Figure 1. Disease phenotypes. (A) Illustration of the scale used to evaluate the severity of symptoms: (0) no symptoms or healthy plant (1) mild symptoms without chlorosis (2) visible chlorosis, (3) advanced chlorosis, (4) strong chlorosis and incipient necrosis, (5) necrosis and death of the plant. (B - E) Infection phenotypes across lines in response to the ancestral (blue) and evolved (red) TuMV isolates: infectivity (B), AUDPS (C), intensity of symptoms (D), and necrosis (E). (F) Correlation matrix between disease phenotypes in response to the ancestral (upper left) and evolved (lower right) isolates. The diagonal shows correlations between the same phenotype in response to each viral isolate. (G) SNP heritability (h^2_{SNP}) for each disease-related trait in response to each viral isolate.

When considering the way each line responds to infection with the ancestral or evolved TuMV isolate (Fig. 2A), we observed that the evolved isolate was significantly more infective (Fig. 2B; Friedman test: $\chi^2 = 528.122$, 2 d.f., $P < 0.001$), which means that adaptation to the Col-0 line leads to a virus isolate often more infective in other lines.

287   Indeed, the number of lines that did not show any symptoms after inoculation with each

288   viral isolate are largely different: 91 lines in case of the ancestral and only 11 for the

289   evolved. Twenty-two lines show no symptoms of infection regardless the viral isolate.

Figure 2. (A) The progression of disease (measured as AUDPS) was compared at different time points between the ancestral and evolved viral isolates in multiple *A. thaliana* lines. Each data point represents a different line, and the segments connecting the same line are color-coded green if the AUDPS has increased for the evolved virus, or purple if it has decreased. (B) Number of *A. thaliana* lines exhibiting increased, decreased, or nonsignificant changes in susceptibility when inoculated with the evolved or ancestral isolates.

292   **A major locus associated with necrosis and severity of symptoms**

We used the multi-trait GWA analysis implemented in the software package LIMIX to identify individual genetic loci that correlate with the response to each virus (Lippert et al., 2014). LIMIX assesses the response to each virus jointly, and identifies loci associated with (1) a shared response to both viruses, and (2) specific responses to individual viral isolates.

We found a single strong peak of association for necrosis in response to both viruses on chromosome 2 (Figs. 3A - 3D). The region of association encompasses five annotated genes, and is within 300 bp of two more, as well as at least six annotated transposable elements (Fig. 3, Table 1). 56% and 49% of the lines showing necrosis in response to the ancestral and evolved viruses respectively harbor the susceptible allele at the most strongly associated SNP at position 5,927,469. To validate this association, we repeated the experimental procedure using 51 lines that showed systemic necrosis in the initial experiment and 65 that did not, and recovered the association in the same region on chromosome 2, with a peak at position 5,927,469 (Fig. S1). These results demonstrate that a major association on chromosome 2 explains approximately half the cases of necrosis.

The peak of association with necrosis overlaps with a peak found for severity of symptoms (Fig. S2). Since necrosis is the final stage of disease progress in the semi-quantitative scale used to score symptoms, this overlap in genetic architecture likely reflects the same biological signal. For this reason, and because ordinal data are hard to handle statistically, we focus on necrosis for the rest of this paper. We did not identify any significant associations with AUDPS or infectivity (Figs. S2 and S3), consistent with the low heritability for these traits (< 30%; Fig. 1B).

The presence of a strong association can interfere with the detection of other associations (Lippert et al., 2014). To identify additional loci associated with systemic necrosis we therefore repeated the GWA analysis including the genotype at the most strongly associated SNP as a covariate. This revealed four additional loci showing a significant association with the shared response to both TuMV isolates (Figs. 3A and 3C), and eight loci with an isolate-specific response (Figs. 3B and 3D). Linkage disequilibrium between these loci is weak (Fig. S4). Necrotic alleles at these loci were rare (3.1 - 17.0% allele frequency; median = 3.6%) and showed up to 9.7-fold increase in the relative risk of developing systemic necrosis compared to the non-necrotic alleles (Table S1, Figs. 3E and 3F). Where loci displayed a virus-specific effect, this was

 associated with increased relative risk of systemic necrosis in response to the evolved

 viral isolate compared to that of the ancestral virus one (Fig. 3F). Overall, all but two

 necrosis-developing lines harbored the necrotic allele in at least one of the loci identified.

Figure 3. Genetic associations with systemic necrosis. (A), (C) and (E) show SNPs associated with a common response to both viruses. (B), (D) and (F) show SNPs associated with a differential response to each viral isolate. (A) and (B) show associations with raw necrosis data. (C) and (D) show associations in a model that includes the genotype at the most strongly associated SNP in (A) as a cofactor. (E) and (F) show relative risk of necrosis in response to the ancestral (closed symbols) and evolved (open symbols) TuMV isolates for the susceptible alleles at (E) the five loci associated with the response to both isolates and (F) the eight loci associated with an isolate-specific response. Note the log scale of the ordinate axis. Colors in (E) and (F) correspond to those in the Manhattan plots above.

Loss-of-function mutants of *AT2G14080* and *DRP3B* show TuMV isolate-dependent differences in the severity of symptoms

We assessed the impact of associated loci on response to virus infection by analyzing the severity of symptoms in two KO mutants mapped within 1.5 Mb of the candidate region of chromosome 2: the TIR-NBS-LRR protein encoded by *AT2G14080* and *DRP3B*. We fitted a GLMM to assess the effect of infection with each viral isolate in the severity of symptoms shown by the wild-type and both mutant plants. Focusing in the intra-individual effects, the duration of infections shows a highly significant effect ($F_{2,162}$ = 40.481, $P < 0.001$), increasing a 328.6% ±82.2 (±1 SEM) in severity between 7 and 21 dpi (Fig. 4). No other intra-individual effect was found significant (*i.e.*, interaction of the factors with dpi; in all cases $P \geq 0.430$). Regarding the inter-individual effects, all three were significant. First, differences between the three plant genotypes exist in the average severity of symptoms ($F_{2,162}$ = 5.379, P = 0.006), with the plants *drp3b* showing symptoms 30.0% ±5.4 milder than wild-type plants but *at2g14080* showing 16.6% ±13.3 stronger symptoms than wild-type plants (Fig. 4). The average stronger symptoms in *at2g14080* plants are consistent with the antiviral role of the encoded TIR-NBS-LRR protein, while the average weaker symptoms in *drp3b* is consistent with a proviral activity. Second, the two viral isolates also induce symptoms that significantly differ in severity ($F_{1,162}$ = 9.001, P = 0.003), with the evolved isolate inducing 45.6% ±15.9 more severe symptoms, on average, than the ancestral one (Fig. 4). Finally, and more interestingly, a significant interaction between plant genotype and viral isolate exists ($F_{2,162}$ = 3.907, P = 0.022). This effect means that the magnitude of the difference in symptoms induced by the evolved and ancestral TuMV isolates depends on the host genotype (Fig. 4). Whereas in wild-type plants, the evolved isolate induces symptoms that are 124.9% ±47.3 stronger than those induced by the ancestral, in *at2g14080* plants the difference is largely reduced to a non-significant 1.53% ±13.8. By contrast, the symptoms induced by the evolved isolate in *drp3b* plants are 35.3% ±27.7 stronger than what the ancestral isolate does.

Figure 4. Symptoms severity at different stages of infection for the ancestral and the evolved TuMV isolates inoculated in wild-type, *at2g14080* and *drp3b* plants. Error bars represent ±1 SEM.

A *Copia* element insertion in the primary candidate gene

We explored haplotypes in the region associated with necrosis in 53 genomes assembled

de novo from long-read PacBio sequencing data as part of a different study

(https://1001genomes.org/). We aligned the sequences in the region around the strong

association for necrosis and severity of symptoms. To assess patterns of synteny, we

looked for homology with the two most strongly associated genes (*AT2G14080* and

DRP3B) and the eight annotated genes or transposable elements from the TAIR10

genome annotation on either side of these genes using BLAST. This analysis revealed

many structural polymorphisms (Fig. 5), including abundant intergenic indel variation,

duplications and a large presence/absence polymorphism downstream of *AT2G14080*

containing *AT2G14100* and a block of transposable elements. At least one line (9470)

shows a large-scale inversion for the entire region. The region around the strong

association for necrosis and severity of symptoms appears to be a hotspot for structural

variation.

Figure 5. Multiple sequence alignment around the major association with necrosis on chromosome 2. (A) Enlarged view of the peak in Fig. 3A, with the TAIR10 annotation below. (B) A summary of structural variation in 53 assembled genomes. Transposable elements are shown in orange and coding genes in gray, with candidate genes *AT2G14080* and *DRP3B* highlighted in purple and pink respectively. Black lines connect homologous features. Each genome is labeled (by squares on the right) according to its necrosis phenotype and genotype at the most associated SNP. The lines without boxes have not been phenotyped.

We next looked for structural polymorphism that might be causal for the disease

phenotypes. We categorized lines as "alive" *vs.* "systemic necrosis" based on whether

they showed systemic necrosis in response to either virus, and as having the necrotic *vs.*

non-necrotic alleles at the most strongly associated SNP (Fig. 3).   Unfortunately, only
four lines of the 53 sequenced both showed systemic necrosis and had the necrotic allele
at the major association, but all of them harbored a *Copia*-element inside the first intron
of *AT2G14080* that was present in only one other line.   We tried to genotype the
presence/absence of this element in the full sample using short-read sequencing data, but
this turned out to be unachievable due to the repetitiveness of the sequence (see Materials
and methods).   Thus, this element remains a promising candidate for a causal
polymorphism.

The major susceptibility allele is spatially overdispersed
390
The susceptible allele at the major association with necrosis on chromosome 2 is spread
throughout the global range of the GWA panel (Fig. 6A).   To investigate whether this
broad spatial distribution could be due to chance, we compared the mean geographic
distance between lines with the susceptible alleles at each of the top SNPs identified by
GWA (Fig. 3) to the mean distances between lines with the minor allele at 10,000
randomly chosen loci with similar minor allele counts.   For most associated loci, distances
between lines with susceptible alleles are well within the distribution of randomly chosen
loci, consistent with genetic drift allowing minor alleles to reach modest frequency on a
local spatial scale.   In contrast, lines harboring the susceptible alleles at the major
association on chromosome 2 are further apart on average than 98.5% of randomly chosen
alleles at similar frequencies (Fig. 5C).   This allele is found at a similar frequency in all
but three admixture groups identified by reference (Alonso-Blanco et al., 2016), so this
overdispersion cannot be accounted for by an association with a particular ancestry with
a high dispersal rate (Fig. 6B).   We found similar spatial overdispersion for the associated
loci on chromosome 4 at positions 273,465 and 7,917,355.   Alleles associated with
systemic necrosis at the major association, thus seem to be spatially overdispersed, in a
way which is independent of genetic background, and which is unlikely to be due to
genetic drift alone.

Figure 6. Global distribution of necrotic alleles. (A) Global distribution of genotypes (non-necrotic/necrotic alleles at the SNP most strongly associated with systemic necrosis), and phenotypes (whether lines showed systemic necrosis or not). The inset table shows the numbers of each genotype/phenotype; colors correspond to those on the map. (B) Frequencies of necrotic alleles at chromosome 2 position 5,927,469 within each admixture group. The horizontal line indicates global frequency. (C) Mean distances between pairs of loci harboring the necrotic allele at the 13 associated loci (red) and between minor alleles at 10,000 randomly chosen loci.

Discussion

We report results from the largest GWA study done so far aiming to identify *A. thaliana*

genes involved in the infection response to one of its natural pathogens, TuMV, using an

ancestral isolate naïve to *A. thaliana*, and a second isolate that had been experimentally

evolved on *A. thaliana* line Col-0. The comparison of the results for the two isolates has

pinpointed candidate host targets for virus adaptation.

Increased virulence of the evolved virus

Lines infected with the evolved virus showed increased disease symptoms compared to

plants infected with the ancestral virus (Fig. 1D - 1G). Systemic necrosis was strongly

correlated between virus treatments (Fig. 1C), and there was no evidence for a virus-

specific effect of alleles at the major association locus on chromosome 2 (Fig. 3E). The

increase in necrosis can thus probably be attributed to increased virulence of the evolved

virus. In contrast, the increase in AUDPS, infectivity and severity of symptoms was

associated with changes in the rank order of lines, reflected in only moderate correlations

between phenotypes in response to the two viral isolates (Fig. 1C). This suggests that the

change in these phenotypes reflects a more complex line-specific response to naïve and

adapted TuMV isolates.

The increase in virulence of the evolved isolate used in this study is associated with

two non-synonymous mutations acquired during adaptation to *A. thaliana* (Navarro et al.,

2022). First, mutation T1293I was found in the cylindrical inclusion protein (CI) that is

involved in viral genome replication and cell-to-cell movement (Deng et al., 2015), and

has been shown to interact with the photosystem I PSI-K protein in *A. thaliana* (Jiménez

et al., 2006). Second, mutation N2039H was found in the viral genome-linked protein

(VPg) that is involved in replication and intracellular movement (Wu et al., 2018), and is

a hub for interactions with other viral proteins (Bosque et al., 2014). Mutations in VPg

have been pervasively observed in evolution experiments of potyviruses in *A. thaliana*,

in all cases resulting in increased infectivity, virulence and viral load (Agudelo-Romero

et al., 2018; Hillung et al., 2014; González et al., 2021; Navarro et al, 2022; Melero et al.,

2023). Together, these observations indicate that genome replication and viral movement

underlie the increased virulence of the evolved virus.

Our study confirms that the adaptation to one susceptible *A. thaliana* line, in this

case Col-0, enables potyviruses to perform better in otherwise poorly susceptible lines.

This observation was first made by Lalić et al. (2010) after evolving a tobacco etch virus

isolate from tobacco in the susceptible line L*er*-0. Upon inoculation of 18 other lines,

they observed that the evolved isolate exhibited greater infectivity compared to the

ancestral one.

The genetic architecture of TuMV resistance

We identified a 1.5 Mb region on chromosome 2 that was strongly associated with the

severity of symptoms, and accounted for approximately half the cases of systemic

necrosis. This region encompasses several genes (Table 1), two of which are especially

strong candidates for causing the phenotype. First, *AT2G14080* encodes a known TIR-

NBS-LRR family protein. These proteins recognize and block translation of viral

transcripts in the cytoplasm, induce hypersensitive response leading to cell death, act as

a second line of defense if the initial antiviral response fails to restrict virus accumulation,

and are involved in signaling pathways (Li et al., 2001; Meyers et al., 2003; Bhattacharjee

et al., 2009; Kobayashi et al., 2010; Marone et al., 2013; van de Weyer et al., 2019). TIR-

NBS-LRR genes are the most numerous disease-resistance genes in the *A. thaliana*

genome and are under diversifying selection to respond to as broad a spectrum of

pathogens as possible (Ellis et al., 2003). This gene appears to have a significant role in

development of systemic necrosis during infection with both TuMV strains. Second,

DRP3B encodes a self-assembling GTPase involved in fission and fusion of membranes

in mitochondria and peroxisomes (Fujimoto et al., 2009; Wu et al., 2018). Its analogue,

DRP2B, has been shown to be co-opted by TuMV to enhance virulence, and treatment

with dynamin-specific inhibitors suppress growth by interfering with VPg (Wu et al.,

2018). As previously noted, one of the two amino-acid differences between the ancestral

and evolved TuMV isolates is in VPg (Navarro et al., 2022). These observations make

AT2G14080 and *DRP3B* strong candidate genes for a role in viral resistance, either

independently or in tandem. To test this possibility, the ancestral and evolved isolates

were inoculated into KO mutants *at2g14080* and *drp3b*. We observed that, compared to

wild-type plants, *at2g14080* plants showed different responses for each viral isolate:

while the ancestral isolate induced more severe symptoms, those induced by the evolved

virus were similar or weaker in the wild-type plants (Fig. 4). These findings demonstrate

a difference in the host genes involved in the response to the two viral isolates and suggest

that *AT2G14080* is efficient for the defense against the ancestral but not for the evolved

isolate. The evolved viral isolate may be able to overcome the response imposed by

AT2G14080 due to the mutations acquired during their adaptation to Col-0, most probably

because of the changes in the VPg viral protein that allow it to evade detection by the

TIR-NBS-LRR protein encoded by *AT2G14080*. In the case of the ancestral isolate, we

found no meaningful differences in symptomatology between wild-type and *drp3b* plants.

In sharp contrast, *drp3b* plants show milder symptoms than wild-type plants, confirming

a possible proviral effect of DRP3B that has been optimized during the previous

evolutionary history in Col-0. The possibility of epistatic effects between these two

candidate genes is highly likely considering the complex interplay between genetic factors and environmental cues in determining the phenotypic outcomes of viral infections.

Twelve additional loci (besides the major association on chromosome 2) were significantly associated with the response to one (5) or both (8) viruses (Fig. 3). Lines with the susceptible alleles at these loci were associated with up to 9.7-fold increased risk of systemic necrosis compared to lines with the resistant allele (Figs. 2E - 2F), and the low linkage disequilibrium between these loci indicates that they segregate independently of the major association on chromosome 2, and of each other (Fig. S4). However, these alleles are also at very low global frequencies (Table S1). On the one hand, this is consistent with the expectation that selection against such deleterious alleles keeps susceptible alleles rare. On the other hand, there is a substantial risk that alleles at such low frequency should be associated with a binary phenotype at similar frequency just by chance. As such, caution is warranted in interpreting a causative role for these associations without further evidence.

We found very little overlap with associations found in two previous GWA studies of TuMV resistance in *A. thaliana*. First, Butković et al. (2021) identified genome-wide associations for AUDPS, severity of symptoms and infectivity under very similar experimental conditions to those used here, but using two viral isolates that differed in their host range (one evolved as a specialist in more permissive *A. thaliana* lines and the other evolved as a generalist able to successfully infect more resistant lines) and fewer host lines. That analysis recovered the association with *AT2G14080* for both isolates, and identified eleven additional loci associated with the response to one or both viruses. These eleven loci identified were not recovered in our larger dataset. This lack of consistency can either indicate that these associations were spurious artifacts of a limited sample size or, alternatively, a true biological effect due to the different TuMV isolates used by Butković et al. (2021) and in this study. Second, we did not detect any of the genes associated with viral load or infectivity in 317 lines grown under field conditions identified by Rubio et al. (2019), and nor did that study detect the association on chromosome 2 reported here. In this case, the lack of overlap may be explained not only by the differences in sample sizes, but also by the differences in traits measured, viral isolate and environmental conditions.

The *Copia*-element insertion is a candidate causative variant

We identified a *Copia*-element polymorphism in the first intron of *AT2G14080,* which is found in all the four lines with long-read genome assemblies that show systemic necrosis in response to either virus (and only one other line). Given this strong association, and the fact that first introns often harbor regulatory elements, this insertion is a strong candidate for a causative mutation. Whether it turns out to be casual or not, the abundance of structural variation in this region highlights the need for restraint when trying to identify any causal polymorphisms from SNP data alone, and more generally the limitation of reference-biased polymorphism data.

It is also interesting to consider whether structural variation is driven by selection for diversification of resistance genes, or whether particular structural variants themselves cause variation in disease resistance. It has been previously observed that plant disease-resistance genes often form clusters of duplicated copies within the genome, which would contribute to structural variation between genomes (Leister, 2004). In particular, NBS-LRR genes, of which *AT2G14080* is an example, are known to have undergone expansion by numerous large- and small-scale duplication and translocation events (Leister, 2004). However, we did not observe widespread duplications of *AT2G14080* or its domains, nor evidence of synteny with other regions of the genome that would indicate translocations, so in this case a simple transposon insertion remains the most likely explanation.

The distribution of susceptible alleles is consistent with frequency-dependent selection

Given that potyvirus outbreaks are common in nature (Pagán et al., 2010) and susceptibility is unquestionably deleterious, it is curious that susceptible alleles should be at sufficiently high frequency to be detectable. There are three explanations for this. First, selection may be sufficiently weak or the virus geographically restricted, such that alleles can arise in one location and increase in frequency by genetic drift. If this were true, we would expect to see a clustered geographic distribution of susceptible alleles, reflected in under dispersion of distances between pairs of susceptible lines. While this is true for most of the loci with small effects on necrosis, susceptible alleles at the major association on chromosome 2 are spatially over dispersed worldwide, and found throughout unrelated lines (Fig. 6). These patterns are difficult to explain via genetic drift. Second, it may be that the genomic instability in the region surrounding this

association leads to a high turnover of new structural mutations that decrease viral resistance. This is implausible because this would not generate linkage with individual SNPs that could be detected by GWA. Moreover, we found a striking concordance between a single *Copia*-element insertion into *AT2G14080* and necrosis, suggesting that only a single variant is responsible for increased necrosis in this region. Neither genetic drift nor mutation-selection balance can thus explain the persistence of susceptible alleles at the major association for necrosis.

An alternative explanation is that susceptible alleles are beneficial in the absence of the virus. This would cause a fitness trade-off between alleles in this region, as has been previously demonstrated for many other pathogen-resistance systems in plants (Bergelson et al., 2001; Todesco et al., 2010). Potyvirus outbreaks are common in natural *A. thaliana* populations, but ephemeral (Pagán et al., 2010), causing susceptible alleles to decrease in frequency during outbreaks, but to increase at other times. This would lead to negative frequency-dependent selection maintaining the susceptible allele at a frequency in proportion to the frequency of TuMV outbreaks. Despite being at low frequency worldwide, the susceptible allele is found across the geographic and genealogical distribution of *A. thaliana*, indicating that the polymorphism must either be very old and has been maintained by balancing selection over long periods, or that alleles have spread by adaptive introgression during viral outbreaks. These data do not allow us to distinguish these hypotheses, nor are they mutually exclusive, but both are consistent with a fitness trade-off at this locus. We conclude that the worldwide geographic distribution of susceptible alleles is therefore consistent with negative frequency-dependent selection maintaining the susceptible allele at low, but non-zero, frequency.

Data availability

Phenotype data and code to reproduce the analyses presented here are given at GitHub (https://github.com/ellisztamas/tumv_ms) and Zenodo (doi: 10.5281/zenodo.5789488). Sequence data are available from the 1001 Genomes Project website (https://1001genomes.org/) (Alonso-Blanco et al., 2016). Unless otherwise stated analyses and plotting were done in R 4.0.3 (R Core Team, 2021) under RStudio Server Pro 1.3.1093-1 (RStudio Team, 2021).

Acknowledgements

We thank Paula Agudo, Francisca de la Iglesia and Joanna Gunis for excellent technical
support. Work was funded by Spain's Agencia Estatal de Investigación-FEDER grant
PID2019-103998GB-I00 and Generalitat Valenciana grants GRISOLIAP/2018/005 and
PROMETEO/2019/012 to SFE. RG were supported by Agencia Estatal de Investigación-
FEDER contract BES-2016-077078. MN was funded by ERC AdG 789037 -
EPICLINES, ERA-CAPS (FWF I 3684-B25) and by the GMI.

Authors contributions

608 MN and SFE planned and designed the research. AB and RG performed experiments.
609 AB, TJE, RG, SFE and BJ analyzed data. AB, TJE, RG, MN, and SFE wrote the
610 manuscript. AB, TJE and RG contributed equally.

References

Agudelo-Romero P, Carbonell P, Pérez-Amador MA, Elena SF. Virus adaptation by
manipulation of host's gene expression. PLoS ONE. 2008;3: e2397.

Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J. 1,135 Genomes reveal the
global pattern of polymorphism in *Arabidopsis thaliana*. Cell. 2016;166: 481–491.
Arabidopsis 1001 Genomes Consortium. In preparation.
Bartoli C, Roux F. Genome-wide association studies in plant pathosystems: toward an
ecological genomics approach. Front Plant Sci. 2017;8: 763.
Becker RA, Wilks AR, Brownrigg R, Minka T, Deckmyn A. maps: draw geographical
maps. 2021. Available: https://CRAN.R-project.org/package=maps.
Becker RA, Wilks AR, Brownrigg R. mapdata: Extra map databases. 2018. Available:
https://CRAN.R-project.org/package=maps
Bergelson J, Kreitman M, Stahl EA, Tian D. Evolutionary dynamics of plant *R* genes.
Science. 2001;292: 2281–2285.
Bhattacharjee S, Zamora A, Azhar MT, Sacco MA, Lambert LH, Moffett P. Virus
resistance induced by NB-LRR proteins involves *Argonaute4*-dependent translational
control. Plant J. 2009;58: 940–951.
Bonfield JK, Marshall J, Danecek P, Li H, Ohan V, Whitwham A, *et al*. HTSlib: C library
for reading/writing high-throughput sequencing data. GigaScience. 2021;10: giab007.
Bosque G, Folch-Fortuny A, Picó J, Ferrer A, Elena SF. Topology analysis and
visualization of *Potyvirus* protein-protein interaction network. BMC Syst Biol.
2014;8: 129.
Boyes DC, Zayed AM, McCaskill AJ, Hoffman NE, Davis KR. Growth stage-based
phenotypic analysis of Arabidopsis: a model for high throughput functional genomics
in plants. Plant Cell. 2001;13: 1499–1510.
Brosseau C, Bolaji A, Roussin-Léveillée C, Zhao Z, Biga S, Moffett P. Natural variation
in the Arabidopsis *AGO2* gene is associated with susceptibility to potato virus X. New
Phytol. 2020;226: 866–878.
Butković A, González R, Cobo I, Elena SF. Adaptation of turnip mosaic potyvirus to a
specific niche reduces its genetic and environmental robustness. Virus Evol. 2020;6:
veaa041.
Butković A, González R, Rivarez MPS, Elena SF. A genome-wide association study
identifies *Arabidopsis thaliana* genes that contribute to differences in the outcome of
infection with two turnip mosaic potyvirus strains that differ in their evolutionary
history and degree of host specialization. Virus Evol. 2021;7: veab063.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J. BLAST+: architecture and
applications. BMC Bioinformatics. 2009;10: 421.

Caparrós-Martín JA, McCarthy-Suárez I, Culiáñez-Macià FA. HAD hydrolase function
unveiled by substrate screening: enzymatic characterization of Arabidopsis thaliana
subclass I phosphosugar phosphatase AtSgpp. Planta. 2013;237: 943–954.

Chen CC, Chao CH, Chen CC, Yeh SD, Tsai HT. Identification of turnip mosaic virus
isolates causing yellow stripe and spot on calla lily. Plant Dis. 2003;87: 901–905.

Covarrubias-Pazaran G. Genome-assisted prediction of quantitative traits using the R
package sommer. PLoS ONE. 2016;11: e0156744.

Deng P, Wu Z, Wang A. The multifunctional protein CI of potyviruses plays interlinked
and distinct roles in viral genome replication and intercellular movement. Virol J.
2015;12.

Ellis J, Dodds P, Pryor T. Structure, function and evolution of plant disease resistance
genes. Curr Opin Plant Biol. 2000;3: 278–284.

Fujimoto M, Arimura S, Mano S, Kondo M, Saito C. Arabidopsis dynamin-related
proteins DRP3A and DRP3B are functionally redundant in mitochondrial fission, but
have distinct roles in peroxisomal fission. Plant J. 2009;58: 388–400.

Garcia-Moreno M, Järvelin AI, Castello A. Unconventional RNA-binding proteins step
into the virus-host battlefront. WIREs RNA. 2018;9: e1498.

González R, Butković A, Elena SF. Role of host genetic diversity for susceptibility-to-
infection in the evolution of virulence of a plant virus. Virus Evol. 2019;5: vez024.

González R, Butković A, Escaray FJ, Martínez-Latorre J, Melero I, Pérez-Parets E, *et al*.
Plant virus evolution under strong drought conditions results in a transition from
parasitism to mutulism. Proc Natl Acad Sci USA. 2021;118: e2020990118.

Hillung J, Cuevas JM, Valverde S, Elena SF. Experimental evolution of an emerging
plant virus in host genotypes that differ in their susceptibility to infection. Evolution.
2014;68: 2467–2680.

Jiménez I, López L, Alamillo JM, Valli A, García JA. Identification of a plum pox virus
CI-interacting protein from chloroplast that has a negative effect in virus infection.
Mol Plant Microbe Interact. 2006;19: 350–358.

Jones RAC. Global plant virus disease pandemics and epidemics. Plants. 2021;10.

Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, *et al*. Efficient
control of population structure in model organism association mapping. Genetics.
2008;178: 1709–1723.

Kobayashi M, Seo S, Hirai K, Yamamoto-Katou Y, Katou S, Seto H, Meshi T, Mitsuhara
I, Ohashi Y. Silencing of WIPK and SIPK mitogen-activated protein kinases reduces
tobacco mosaic virus accumulation but permits systemic viral movement in tobacco
possessing the *N* resistance gene. Mol Plant-Microbe Interact. 2010;23: 1032–1041.

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res. 2017;27: 722–736.

Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q. A mixed-model approach for
genome-wide association studies of correlated traits in structured populations. Nat
Genet. 2012;44: 1066–1071.

Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biol. 2009;10: R25.

Lalić J, Agudelo-Romero P, Carrasco P, Elena SF. Adaptation of tobacco etch potyvirus
to a susceptible ecotype of *Arabidopsis thaliana* capacitates it for systemic infection
of resistant ecotypes. Phil Trans R Soc B 2010; 365:1997-2007.

Leister D. Tandem and segmental gene duplication and recombination in the evolution of
plant disease resistance genes. Trends Genet. 2004;20: 116–122.

Li H. New strategies to improve minimap2 alignment accuracy. Bioinformatics. 2021;37:
4572–4574.

Li X, Clarke JD, Zhang Y, Dong, X. Activation of an EDS1-mediated *R*-gene pathway in
the *snc1* mutant leads to constitutive, NPR1-independent pathogen resistance. Mol
Plant-Microbe Interact. 2001;14: 1131–1139.

Lippert C, Casale FP, Rakitsch B, Stegle O. LIMIX: genetic analysis of multiple traits.
bioRxiv. 2014; doi: 10.1101/003905.

Liu S, Chen M, Li R, Li, WX, Gal-On A, Jia Z, Ding SW. Identification of positive and
negative regulators of antiviral RNA interference in *Arabidopsis thaliana*. Nat
Commun. 2022;13: 2994.

Luan H, Niu H, Luo J, Zhi H. Soybean cytochrome b5 is a restriction factor for soybean
mosaic virus. Viruses. 2019;11: 546.

Marone D, Russo M, Laidò G, De Leonardis A, Mastrangelo A. Plant nucleotide binding
site-leucine-rich repeat (NBS-LRR) genes: active guardians in host defense responses.
Int J Mol Sci. 2013;14: 7302–7326.

McDonald BA, Stukenbrock EH. Rapid emergence of pathogens in agro-ecosystems:
global threats to agricultural sustainability and food security. Phil Trans R Soc B.
2016;371: 20160026.

Melero I, González R, Elena SF. Host developmental stages shape the evolution of a plant
RNA virus. Phil Trans R Soc B. 2023;378: 20220005.

Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW. Genome-wide analysis of
NBS-LRR–encoding genes in Arabidopsis. Plant Cell. 2003;15: 809–834.

Monnot S, Desaint H, Mary-Huard T, Moreau L, Schurdi-Levraud V, Boissot N.
Deciphering the genetic architecture of plant virus resistance by GWAS, state of the
art and potential advances. Cells. 2021;10: 3080.

Montes N, Cobos A, Gil-Valle M, Caro E, Pagán I. *Arabidopsis thaliana* genes associated
with cucumber mosaic virus virulence and their link to virus seed transmission.
Microorganisms. 2021;9: 692.

Navarro R, Ambrós S, Butković A, Carrasco JL, González R, Martínez F, *et al*. Defects
in plant immunity modulate the rates and patterns of RNA virus evolution. Virus Evol.
2022;8: veac059.

Oerke EC. Crop losses to pests. J Agric Sci. 2006;144: 31–43.

Ouibrahim L, Caranta C. Exploitation of natural genetic diversity to study plant-virus
interactions: what can we learn from *Arabidopsis thaliana*? Natural virus resistance in
Arabidopsis thaliana. Mol Plant Pathol. 2013;14: 844–854.

Pagán I, Fraile A, Fernandez-Fueyo E, Montes N, Alonso-Blanco C, García-Arenal F.
Arabidopsis thaliana as a model for the study of plant–virus co-evolution. Phil Trans
R Soc B. 2010;365: 1983–1995.

R Core Team. R: A language and environment for statistical computing. R Foundation
for Statistical Computing; 2021.

Revers F, García JA. Molecular biology of potyviruses. Adv Virus Res. 2015;92: 101–
199.

RStudio Team. RStudio: Integrated development environment for R. Boston, MA:
RStudio, PBC; 2021.

Rubio B, Cosson P, Caballero M, Revers F, Bergelson J. Genome-wide association study
reveals new loci involved in *Arabidopsis thaliana* and turnip mosaic virus (TuMV)
interactions in the field. New Phytol. 2019;224: 2026–2038.

Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, *et al*. An efficient multi-
locus mixed-model approach for genome-wide association studies in structured
populations. Nat Genet. 2012;44: 825–830.

Simko I, Piepho HP. The area under the disease progress stairs: calculation, advantage,
and application. Phytopathology. 2012;102: 381–389.

Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M. Natural allelic variation
underlying a major fitness trade-off in *Arabidopsis thaliana*. Nature. 2010;465: 632–
636.

Tomlinson JA. Epidemiology and control of virus diseases of vegetables. Ann Appl Biol.
1987;110: 661–681.

Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, *et al*. A
species-wide inventory of NLR genes and alleles in *Arabidopsis thaliana*. Cell.
2019;178: 1260-1272.e14.

Wu G, Cui X, Chen H, Renaud JB, Yu K. Dynamin-like proteins of endocytosis in plants
are coopted by potyviruses to enhance virus infection. J Virol. 2018;92: e01320–18.

Xu J, Wang X, Guo W. The cytochrome P450 superfamily: Key players in plant
development and defense. J Integ Agric. 2015;14: 1673–1686.

Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, *et al.* Common
SNPs explain a large proportion of the heritability for human height. Nat Genet.
2010;42: 565–569.

Table 1. Significant SNPs and their corresponding genomic regions on chromosome 2 detected for necrosis in the multiple trait analysis for both viral isolates.

SNP	Locus	$-\log P$	Function of significant genes	Reference
5,923,326	200 bp upstream of *AT2G14070*	20.55	Wound-responsive protein-like protein	-
5,928,864	*AT2G14080*	20.43	Disease-resistance protein of the TIR-NBS-LRR class lead to hypersensitive response and resistance	Meyers et al. (2003); Marone et al. (2013);
5,931,085	1.2 Kb upstream of *AT2G14095*	14.39	Hypothetical protein	-
5,933,529	*AT2G14095* and upstream of *AT2G14100*	8.42	A member of the cytochrome P450 family; plays roles in plant defense against pathogens and resistance.	Xu et al. (2015); Lua
5,952,757	*AT2G14110*	11.32	Haloacid dehalogenase-like hydrolase (HAD) proteins are involved in phosphate homeostasis	Caparrós-Martín et a
5,956,602	*AT2G14120*	14.62	Dynamin-related protein 3b suppresses TuMV replication and movement-	Wu et al. (2018)

| 5,975,892 | 200 bp downstream of *AT2G14160* | 13.69 | RNA-binding (RRM/RBD/RNP motifs) family protein; involved in inhibition of RNA virus replication, movement, and translation. | García-Moreno et al. |

Table S1. Loci associated with necrosis, indicating whether the association was with a common response to both TuMV isolates, or with an isolate-specific response.

Effect	Chromosome	Position	Locus	Allele frequency	−log$P$	Function of significant genes
Isolate-specific	1	6,167,132	*AT1G17930*	0.036	10.96	*MAIN*, involved in silencing of transposable elements. Interacts and functions with *MAIL1* and *PP7L* in gene silencing
Isolate-specific	1	12,086,029	*AT1G33330*	0.031	8.70	Class I peptide chain release factor involved in translational termination
Isolate-specific	1	22,508,950	*AT1G61100*	0.310	9.61	Disease resistance protein of the TIR class
Common	2	5,927,469	*AT2G14080*	0.034	74.28	Disease resistance protein of the TIR-NBS-LRR family
Isolate-specific	2	8,198,266	*AT2G18917*	0.045	8.71	Anti-sense long-noncoding RNA overlaps with *AT2G18920* (hypothetical protein) and *ADAGIO2* (regulator of circadian rhythm)
Common	2	8,570,307	*AT2G19850*	0.054	7.68	Transcription repressor
Isolate-specific	3	10,661,084	-	0.031	10.16	Non-coding intergenic region
Isolate-specific	4	273,465	*AT4TE01410*	0.170	8.20	Transposable element in 5' of *AT4G00651* (Fe superoxide dismutase)
Isolate-specific	4	3,920,265	*AT4TE17125*	0.039	10.30	Gypsy-like retrotransposon family (Athila)
Common	4	7,917,355	*AT4TE34810*	0.070	8.01	Transposable element in 5' of *MATERNAL EFFECT EMBRIO ARREST 57*, a DNA methyltransferase regulated by Polycomb group complex during endosperm development
Common	4	16,556,249	*AT4G34690*	0.032	9.23	Transmembrane protein
Common	5	2,386,477	*GRP16*	0.037	7.93	*GLYCINE-RICH PROTEIN 16* is expressed in flowering during specific developmental stage
Isolate-specific	5	16,238,180	*AT5G40540*	0.035	8.21	Cytoplasmic protein kinase superfamily protein

Figure S1. Manhattan plots for the association between SNPs and necrosis in the replicate dataset of 116 lines.

Figure S2. Manhattan plots for associations with common disease responses to both TuMV isolates.

Figure S3. Manhattan plots for associations with disease responses specific to individual

TuMV isolates.