1 **Pantothenate Kinase 4 Governs Lens Epithelial Fibrosis by Negatively**

2 **Regulating Pyruvate Kinase M2-Related Glycolysis**

3 Xue Li [1], Lin-Lin Luo [1], Rui-Feng Li [1], Chun-Lin Chen [1], Min Sun [1,*], Sen Lin [2,*3]

[3] Part of the work was done in Army Medical Center, Chongqing, 400042, China

[1]Department of Ophthalmology, Daping Hospital, Army Medical Center of PLA, Army Medical University, Chongqing, 400042

[2]Department of Neurology, Xinqiao Hospital, The Second Affiliated Hospital, Army Medical University, Chongqing, 400037

Corresponding authors: Sen Lin, sam_lin_amc@163.com; Min Sun, minmins@126.com

Running title: PANK4 deficiency attenuate LEC EMT

Abstract

Lens fibrosis is one of the leading causes of cataract in the elderly population. The primary energy substrate of the lens is glucose from the aqueous humor, and the transparency of mature lens epithelial cells (LECs) is dependent on glycolysis for ATP.

Therefore, the deconstruction of reprogramming of glycolytic metabolism can

contribute to further understanding of LEC epithelial–mesenchymal transition (EMT).

In the present study, we found a novel pantothenate kinase 4 (PANK4)-related

glycolytic mechanism that regulates LEC EMT. The PANK4 level was correlated with

aging in cataract patients and mice. Loss of function of PANK4 significantly

contributed to alleviating LEC EMT by upregulating pyruvate kinase M2 isozyme

(PKM2), which was phosphorylated at Y105, thus switching oxidative

phosphorylation to glycolysis. However, PKM2 regulation did not affect PANK4,

demonstrating the downstream role of PKM2. Inhibition of PKM2 in $Pank4^{-/-}$ mice

caused lens fibrosis, which supports the finding that the PANK4–PKM2 axis is

required for LEC EMT. Glycolytic metabolism-governed hypoxia inducible factor

(HIF) signaling is involved in PANK4–PKM2-related downstream signaling.

However, HIF-1α elevation was independent of PKM2 (S37) but PKM2 (Y105) when

PANK4 was deleted, which demonstrated that PKM2 and HIF-1α were not involved

in a classic positive feedback loop. Collectively, these results indicate a

PANK4-related glycolysis switch that may contribute to HIF-1 stabilization and

PKM2 phosphorylation at Y105 and inhibit LEC EMT. The mechanism elucidation in

our study may also shed light on fibrosis treatments for other organs.

Keywords: PANK4; Fibrosis; Lens epithelial cell; Epithelial-Mesenchymal Transition;

PKM2; Glycolysis; HIF

Introduction

Lens fibrosis, which is characterized by hyperproliferation, migration, and

epithelial–mesenchymal transition (EMT) of lens epithelial cells (LECs), causes

anterior subcapsular cataract (ASC) and posterior capsule opacification (PCO),

damaging vision by changing light scattering[1]. Because of the unique lens

biological properties, lens fibrosis can act as an ideal model for investigating the

molecular mechanisms of tissue and organ fibrosis, such as liver cirrhosis and

idiopathic pulmonary and renal fibrosis, which account for more than 45% mortality[2,

3]. EMT is a major cause of the progression of organ and ocular fibrosis, including the

eye diseases PCO, ASC, pterygium, and glaucoma, and is a part of wound healing

after ocular surgery[4-9]. In fibrotic cataract, the lens epithelium not only loses its

integrity and normal function but also is disorganized with excessive fibronectin and

collagen in the extracellular matrix. Lens epithelium EMT is regulated by various

mechanisms with the classic prominent roles of transforming growth factor

(TGFβ)/Smad signaling[10-13] and glycolytic metabolism[14]. The lens is an

avascular tissue relying on limited supply of oxygen and glucose[15], and the lens

epithelium can utilize oxidative phosphorylation (OXPHOS) for ATP generation by

mitochondria; however, the lens fiber cells are reliant on glycolysis to support their

energy demands[14]. Hence, the regulation of glycolytic metabolism of LECs is a

potential strategy to prevent lens fibrosis. However, findings on the relationship of

cataract to statins (HMG-CoA reductase inhibitors) have been inconsistent[16-18].

Acetyl-CoA is the precursor of HMG-CoA, a CoA product produced during the

second step of aerobic cellular respiration and pyruvate decarboxylation, which

occurs in the mitochondrial matrix. Pantothenate kinase (PANK) 1, PANK2, and

PANK3 are key regulatory enzymes involved in the CoA biosynthetic pathway[19],

and   catalyze   the   phosphorylation   of   the   vitamin   pantothenate   to

4′-phosphopantothenate, subsequently leading to CoA formation. However, PANK4,

the fourth member of the PANK family, lacks enzyme activity because an important

catalytic glutamate site is missing, and it is thus a peculiarity[19-21]. In a previous

study, we found a PANK4 mutation in congenital cataract formation[22], which

results in LEC apoptosis and the aberrant expression patterns of crystallin. However,

the function and mechanism of PANK4 are still under consideration.

Studies in Han Chinese have shown that gene polymorphisms in PANK4 are

associated with type 2 diabetes[23], and functional experiments have shown that

PANK4 can affect the apoptosis of pancreatic β cells[24]. In diabetic cataract

formation, aldose reductase (AR)-mediated polyol accumulation has been shown to

lead to EMT, identified by osmotic imbalance and oxidative stress that result in fiber

cell swelling, liquefaction, and eventually cataract development[25, 26] [27].

Eventually, AR leads to pyruvate formation, which is limited by pyruvate kinase M2

isozyme (PKM2) in the last step of glycolysis[28, 29]. Rat Pank4 is associated with

PKM2 of KOG2201 and KOG4584 domains under both *in vitro* and *in vivo*

conditions[30]. PKM2 is expressed in fetal tissues and cancers, and is a part of EMT

in human colorectal cancer cells[31]. Its expression is enhanced in cancers and favors

the glycolytic pathway and the metastatic potential such as in pancreatic ductal

adenocarcinoma tissues and cell lines[32]. PKM2-mediated glycolysis is also

involved in inflammasome activation by modulating EIF2AK2 phosphorylation in

macrophages[33]. Moreover, PKM2 activation is a potential treatment against diabetic

nephropathy by increasing the glucose metabolic flux and inhibiting the production of

toxic glucose metabolites[34]. PKM2 is recruited by phosphorylating histone H3 at

T11 that results in H3-K9 acetylation and transcription of glycolytic genes, including

SLC2A1, *MYC*, *CCND1*, and lactate dehydrogenase A, which enhance the Warburg

effect, whereas PKM2 is a Cdc25A substrate that forms a positive feedback

loop[35-38]. Unlike many other organs, the eye lens grows throughout life, although

growth is slower in adults than in the young, depending on glycolytic activities[39].

However, it is unclear whether PKM2-dependent glycolytic metabolism is one of the

major activities in lens epithelium maintenance and EMT. Nonetheless, the

relationship between PANK4 and PKM2 in cataract and lens epithelium metabolism

remains unclear.

In the present study, we investigated the role of PANK4 in lens epithlium EMT

and fibrosis. We show that PANK4 was specifically expressed in human and mice lens

epithelium, as well as negatively regulated LEC EMT and functioned in a PKM2

phosphorylation-dependent glycolytic metabolism manner. Without PANK4,

age-related or injury-related lens epithelium fibrosis was attenuated. The glycolytic

metabolism balance was also perturbed. The suppression of the metabolic switch in

LEC by PKM2 activation or by PANK4 loss of function promoted LEC EMT. In

contrast, the preemptive amplification of the metabolic response by the inhibition of

PKM2 promoted lens epithlium EMT, which sheds light on the potential of clinical

drugs in preventing PCO.

Materials and Methods

Pank4 transgenic mice generation

Pank4$^{-/-}$ mice were generated by Nanjing Biomedical Research Institute of Nanjing University, China, as previously described, with 5-bp mutation, in a C57BL/6J background[40]. Briefly, *pank4* knockout mice were generated using the CRISPR/Cas9 system. Cas9 mRNA and sgRNA were co-injected into zygotes. sgRNA direct Cas9 endonuclease caused a cleavage in exon 3 creating a double-strand break. Such breaks were repaired by non-homologous end joining, and resulted in the destruction of *Pank4* gene. The Pank4 targeting sequence was *gacgtggagcaggatcatgagccaccctatgagatctc*

agtccaggagagatcacagctcgcctgcatttcatcaagtttgagaatacctacatggaagcctgcctggacttcatcag

agaccacctagtcaacactgagaccaaggtcatccaggccactgggggtggagcctacaagttcaaggacctcatcg

aggagaagctgcgtctgaa. Offspring were screened for correct genotype by PCR analysis of tail DNA. Mice were housed in cages in a room with a 12-h light-dark cycle with *ad libitum* access to water and rodent chow diet. Animal protocols were approved by the Institutional Animal Care and Use Committee (IACUC) of Army Medical Center, Army Medical University (No. AMUWEC20201055).

Lens epithelial tissue collection from cataract patients

A total of 73 cataract patients (aged 17 to 89 yrs) with unilateral cataract presenting to the Department of Ophthalmology, Army Medical Center, Army Medical University, participated in this study. In particular, 17-year-old patients suffered from traumatic cataract, while other patients had age-related cataract (ARC). All patients provided

informed consent to participate in the study, and all samples were collected with

written informed consent from participants. The study protocol was approved by the

Ethics Committee of the Army Medical Center (No. 2018-31-32 and No. 2020-131).

The inclusion criteria of ARC included: (1) age more than 50 years; (2) no history of

congenital cataract, secondary cataracts, or any other eye diseases; (3) no history of

systemic diseases, such as autoimmune disease and cancer, chronic inflammatory

diseases, and kidney or hepatic diseases; (4) ARC was considered the presence of any

type of ARC (including C-ARC, N-ARC, and PS-ARC) in either eye; (5) ARC

recorded using the Lens Opacities Classification System III (LOCS III). All

procedures involving human material were performed in accordance with the ethical

standards of the institutional and national research committee and the Declaration of

Helsinki. All patients underwent detailed examination that included photography and

slit-lamp microscopy of the lens. Lens epithelium samples were only from healthy

patients. Abnormal coagulation function and etiological examinations for hepatitis B,

hepatitis C, syphilis, human immunodeficiency virus, diabetes, hypertension, and

hyperlipidemia were excluded from the study. The lens epithelium was dissected and

delaminated from the lens of patients by a surgeon. After immediately rinsing with

sterile saline solution, some samples were transferred to dry ice for further molecular

biological experiments, while the other samples were transferred to 4%

paraformaldehyde (PFA) for further immunostaining experiments.

Cell culture and EMT induction

The human lens epithelial cell line (SRA01/04) was presented by Hao-Tian Lin from

the Zhong Shan Ophthalmic Center, Sun Yat-Sen University, as previously

described[41]. SRA01/04 cells were grown in Dulbecco's modified Eagle medium

(DMEM) supplemented with 10% (v/v) fetal bovine serum, 100 U/mL streptomycin,

and 100 U/mL penicillin, and grown at 37°C in a humidified incubator with 5% CO_2.

LECs were seeded in 12-well cell culture plates at 5.0×10^4 cells/mL and incubated in

an incubator (37°C and 5% CO_2) in the standard medium for 24 h. LECs were treated

with TGF-β2 (1 μg/mL) for 12 h until the cell density increased to 60%. Next, LECs

were divided into groups to deal with PKM-IN-1 or DASA-58 separately for 24 h.

Lens culture and treatment

All experimental procedures conformed to the Association for Research in Vision and

Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and Vision

Research. Briefly, the lens culture protocol followed a previous report with some

modification[42-44]. Briefly, lenses from one-month C57BL/6J mice were carefully

removed using a forceps and a pair of fine scissors, and maintained in 4 mL

serum-free DMEM/F12 medium, containing 0.1 mg/mL L-glutamine, 50 IU/mL

penicillin, and 50 mg/mL streptomycin. TGF-β2 was added to the culture medium at a

final concentration of 5 ng/mL. The culture medium was renewed every second day

throughout the culture period. Lenses were cultured for up to 5 days, and the images

were captured using a dissecting microscope (Olympus, Tokyo, Japan).

Immunofluorescence staining

For mice, the eyeballs were removed and fixed in 4% PFA for 3 h, and dehydrated in

a sucrose solution (30%) for 3 h. Eyeballs were embedded in OCT Tissue Tek

Medium, and cut into 10-µm-thick sections adhered onto anti-slip slides using a

frozen section machine. LECs were grown on 13-mm coverslips in 24-well plates and

fixed in 4% PFA for 15 min at room temperature (RT). Eyeball sections and fixed

LECs were washed with phosphate-buffered saline (PBS), and samples were

permeabilized and blocked in 0.3% Triton X-100 and 3% bovine serum albumin in

PBS for 30 min at RT. Samples were incubated with appropriate primary antibodies

(PANK4 was purchased from Cell Signaling Technology (CST), Cat.12055, rabbit

anti-mouse, 1:100 for IF; PKM2 from CST, Cat. 4053, rabbit anti-mouse, 1:100 for IF;

PKM2 from Bioss, Cat. bs-0102M, mouse anti-mouse, 1:200 for IF; p-PKM2^{Y105}

from Abcam, Cat. ab156856, 1:100 for IF; p-PKM2^{ser37} from Thermo Fisher Scientific,

Cat. PA5-37684, 1:200 for IF) (Table 1) overnight at 4°C. PANK4 antibody

specificity was identified in tissues from genetic knockout, heterozygous and WT

mice from the same littermate as described previously [40]. After these steps, the

samples were washed with PBS and incubated with fluorescent secondary antibodies

(Alexa Fluor® 594-AffiniPure Goat Anti-Rabbit IgG (H+L), Cat. 111-585-003, Alexa

Fluor® 488-AffiniPure Goat Anti-Rabbit IgG (H+L), Cat. 111-545-003 and Alexa

Fluor® 594-AffiniPure sheep Anti-mouse IgG (H+L), Cat. 515-585-003 were

purchased from Jackson ImmunoResearch, 1:500 for 2nd antibodies conjugations)

(Table 1) for 0.5 h at RT. Samples were mounted after DAPI staining. In this process,

samples were washed with PBS three times for 3 min between each step. Isotype

antibody controls (Rabbit IgG, polyclonal - Isotype Control, Cat. ab172730 and

mouse IgG2b, kappa monoclonal [7E10G10] - Isotype Control, Cat. ab170192 were

purchased from Abcam) and secondary antibody only controls were also performed to

validate antibody specificity.   Images were captured with a confocal microscope

(Olympus FV3000, Japan).

Protein extraction and Western blot

The protein was obtained from lens tissues in *in vivo* experiments or primary cultured

SRA01/04 cells *in vitro*. Samples were lysed in ice-cold lysis buffer, which consisted

of RIPA buffer and the protease inhibitor cocktail. The protein was then clarified by

Heraeus Sepatech, and the supernatant was collected in new tubes. The protein

concentration was determined using the bicinchoninic acid (BCA) protein assay kit.

The loading buffer was added to protein, and the mixture was boiled for 10 min and

stored at −80°C.

For Western blot, 20 μg of total protein was used to separate on 10% sodium

dodecyl sulfate (SDS)-polyacrylamide gels, and transferred to polyvinylidene fluoride

membranes. The membranes were blocked for 2 h in Tris-buffered saline with 5%

non-fat skimmed milk powder at RT. The membranes were then probed with primary

antibodies (PKM1, Cat. 7067; PKM2, Cat. 4053; N-cadherin, Cat. 14215S; α-SMA,

Cat. 48938; HIF-1α, Cat. 36169; HIF-2α, Cat. 59973; HIF-1β, Cat. 5537 and GAPDH

Cat. 5174 were all rabbit anti-mouse antibodies and purchased from CST, with

working concentration of 1:1000. p-PKM2^{Y105} from Abcam, Cat. ab156856, 1:1000

for WB; p-PKM2^{ser37} from Thermofisher, Cat. PA5-37684, 1:2000 for WB) (Table 1)

at 4°C overnight and probed by the secondary antibodies at RT for 1.5 h. Finally, the

images were captured by the Western blot automatic exposure apparatus.

Co-immunoprecipitation (Co-IP)

Co-immunoprecipitation was performed as described in the manual. Briefly, after

thorough resuspension, magnetization, and washing, SureBeads (Biorad, 1614013)

were resuspended with 5 µg PKM1 (CST, D30G6, Cat. No. 7067), PKM2 (CST,

D78A4, Cat. No. 4053), and PANK4 (CST, D8N2C, Cat. No. 12055) antibodies in a

final volume of 200 µl, respectively, and IgG as the control. Rotate for 10 min at RT.

Supernatant was discarded, and the beads were washed three times with 1,000 µl

PBS-T. Total lysate (300 µg) from WT and *Pank4$^{-/-}$* mice lens epithelium was diluted

to a final volume of 500 µL NP40 lysis buffer (Solarbio, N8032), and

phenylmethylsulfonyl fluoride (PMSF, Solarbio, P0100). Rotate for 1 h at RT.

Magnetized beads were removed and the supernatant was discarded. The beads were

washed with 1,000 µl PBS-T, resuspend the beads thoroughly, centrifugated,

magnetized beads and supernatant was discarded; this process was repeated three

times. Before the last magnetization, the resuspended beads were transferred to a new

tube. All tubes were spun down for several seconds. Magnetized beads were removed,

and the residual buffer was aspirated from the tubes, and 40 µl 1× Laemmli buffer

(biorad,1610747) was added and incubated for 10 min at 70°C. Magnetized beads

were removed, and the eluent was moved to a new vial. Equal amounts of extracts

were separated by SDS-PAGE, and then they were transferred onto nitrocellulose

membranes and then blotted with specific antibodies.

Wound healing assay

SRA01/04 cells were cultured in a 12-well plate. After cells were grouped and

received the corresponding treatment. When cell density increased to 90%, a 10-μL

pipette tip was used to make a straight scratch through the monolayers. In the

following period, cells migrated to the wound surface. The scratch width was used to

represent the migration ability of SRA01/04 cells. After 24 h, the scratch width was

measured in five randomly selected fields. Data presented are means of triplicates.

ATP measurement

Intracellular ATP was determined using an ATP assay kit as previously described[45].

Briefly, cells were lysed for 5 min and centrifuged at 12,000 g at 4°C for 5 min. The

supernatant was added to test plates pre-treated with the ATP detection reagent.

Luminescence was measured using a microplate reader. The remaining cell lysate was

measured with the BCA protein assay kit to determine the protein concentration.

Finally, the ATP concentration was expressed as μmol/g protein.

mRNA sequencing and bioinformatics data analysis

Total RNA extraction, mRNA library construction, and RNA sequencing

(RNA-seq)

Total RNA was extracted from the lens epithelial tissues using TRIzol (Invitrogen,

Carlsbad, CA, USA) according to previously described protocols[46, 47]. Briefly,

after routine RNA extraction, 25–100 µL of diethyl pyrocarbonate-treated water was

added to dissolve RNA. First-strand cDNA was generated using random

hexamer-primed reverse transcription, followed by second-strand cDNA synthesis.

A-tailing mix and RNA index adapters were added for incubating for end repair.

cDNA fragments were amplified using PCR, and the products were purified using

Ampure XP beads (Beckman Coulter) dissolved in EB solution. The product was

confirmed using an Agilent 2100 Bioanalyzer for quality control. The double-stranded

PCR products were heated, denatured, and circularized by the splint oligo sequence to

derive the final library. The single-stranded circular DNA was formatted as the final

library. The final library was amplified with phi29 to prepare a DNA nanoball (DNB),

with more than 300 copies. DNBs were loaded into the patterned nanoarray, and

single-end 50-base reads were generated on an MGISEQ-2000 sequencing platform.

mRNA sequencing was performed by Genetics Biotech Inc. and BioWavelet Ltd. in

Chongqing, China.

Identification of differentially expressed genes

The expression level of genes was calculated using Salmon (v1.4.0). The heatmap

was drawn using pheatmap (v1.0.8). Essentially, differential expression analysis was

performed using DESeq2 (v1.4.5) with Q-value ≤0.05. To gain insight into the change

in phenotype, Gene Ontology (http://www.geneontology.org/) and Kyoto

Encyclopedia of Genes and Genomes (https://www.kegg.jp/) enrichment analyses of

the annotated differentially expressed genes were performed using pHYPER based on

the hypergeometric distribution. The significant terms and pathways were corrected

with Q-value with a rigorous threshold (Q-value ≤0.05) using the Bonferroni method.

Pyruvate detection

The PA levels were determined by the pyruvate assay kit (Pyruvate Assay Kit

(ab65342/K609-100), Abcam). The PA content was evaluated by measuring the

absorbance at 505 nm, and the PA content was calculated using the following formula

(A is absorbance): PA content (µmol/mL) = (A (sample) − A (blank)) / (A (standard) −

A (blank)) × standard concentration (0.2 µmol/mL) / protein concentration of sample

(mg/mL).

Seahorse measurement

Seahorse analysis was performed as previously described[48]. The oxygen

consumption rate (OCR) and extracellular acidification rate (ECAR) were determined

using the Seahorse XF96 Extracellular Flux Analyzer through the Seahorse XF Mito

Stress Test Kit and Seahorse XF Glycolysis Stress Test Kit. It is used to measure cell

dependence on oxidative metabolism or preferentially the glycolytic function in cells.

LECs were counted and seeded in XF96 Seahorse® microplates with 10,000 cells per

well. After cells were grouped and received the corresponding treatment. When cell

density reached 80%, cells were cultured in XF assay medium supplemented with 1

mM glutamine to assess ECAR and in XF assay medium supplemented with 2 mM

glutamine to assess OCR. The cell cultures were placed in a CO_2-free incubator with

37°C for 0.5 h. The plate was left to equilibrate in before being transferred to the

Seahorse XF96 analyzer. The pre-hydrated cartridge was filled with the indicated

compounds and calibrated for 30 min in the Seahorse Analyzer. All experiments were

performed at 37°C. Normalization to protein content was performed after each

experiment. The Seahorse XF Report Generator automatically calculated the

parameters from wave data that were exported to GraphPad Prism software (v8.4,

GraphPad Software).

Statistical analysis

Statistical analysis was performed using GraphPad Prism 8.0 software. Data are

presented as means ± SD. Shapiro-Wilk normality test was performed to assess the

normality of the distribution of data. Due to the small sample sizes, we acknowledge

the assessment of normality may not reliable. For comparison between two groups,

Student's *t*-test was performed. For comparisons among more than two groups,

one-way analysis of variance followed by Bonferroni's post hoc test was performed.

All included data were tested for normality. If normal distribution was not achieved, a

nonparametric test was used. Non-parametric comparisons (Mann-Whitney U Test or

Kruskal-Wallis Test) were used for non-normally distributed data or n < 6. A *P* value

< 0.05 was considered statistically significant.

Results

PANK4 levels increase with age in both rodent and human lens epithelial tissues

PCO-related fibrosis is reported to be associated with lens epithelium aging[12]. In this study, PANK4-positive punctate structures were specifically located in the cytoplasm of LECs of 17-, 47-, and 81-year-old patients (Fig. 1A). The fluorescence intensity of 47- and 81-year-old groups exhibited more intensive gray value in the cytoplasm (Fig. 1B). Furthermore, an increased number of puncta was found in the intracellular cytoplasm of LECs in the aged groups (47 yrs and 81 yrs, respectively) compared with that in younger groups (Fig. 1C; $P < 0.0001$, n = 19). To verify the immune-intensity results, we collected and pooled lens epithelium from patients aged 50–60 yrs, 60–70 yrs, 70–80 yrs, and over 80 yrs groups. The intensities of PANK4 blots were significantly elevated in 70–80 yrs and over 80 yrs female groups compared with the 50–60 yrs female group, respectively (Fig. 1D; P=0.384 *vs* 60-70yrs group, n=6; P=0.0073 *vs* 70-80yrs group, n=6; P=0.0028 *vs* 80yrs+ group, n=6). Similarly, the intensities of PANK4 blots were significantly elevated in 70–80 yrs and over 80 yrs male groups compared with the 50–60 yrs group, respectively (Fig. 1E; P=0.1514 *vs.* 60–70 yrs group, n=6; P=0.0096 *vs.* 70–80 yrs group, n=6; $P <$ 0.0001 *vs.* 80 yrs + group, n=6). In mice, PANK4 was localized in the whole anterior lens epithelium and equator by immunostaining (Fig. 1F). Amplified high power crops showed PANK4-positive signals in cytoplasmic localization (Fig. 1F inner panel), which were consistent with immunostaining, suggesting the promising pattern

of PANK4 in the lens. PANK4 also increased dramatically in 2 Mo, 6 Mo, and 12 Mo

groups rather than in the P15 group (Fig. 1G; *P*=0.0007 *vs.* 2 Mo group, n=6;

P=0.011 *vs.* 6 Mo group, n=6; *P*=0.0003 *vs.* 80 yrs+ group, n=6), which strengthen

the results of PANK4 in humans. Meanwhile, the lens epithelium EMT-related

characteristic molecule, such as N-cadherin, was significantly upregulated in 6 Mo

and 12 Mo groups compared with the P15 group (Fig. 1I, N-cad panel, *P*=0.0317 *vs.*

2 Mo group, n=6; *P*<0.0001 *vs.* 6 Mo group, n=6; *P*<0.0001 *vs.* 12 Mo group, n=6).

The collagen I level was increased in the 2 Mo, 6 Mo, and 12 Mo groups compared

with the P15 group (Fig. 1I, Col I panel, *P*=0.0361 *vs.* 2 Mo group, n=6; *P*=0.0018 *vs.*

6 Mo group, n=6; *P*=0.0005 *vs.* 12 Mo group, n=6). The α-SMA level was elevated in

2 Mo and 12 Mo groups compared with the P15 group. No significant difference was

found in the 6 Mo group compared with the P15 group (Fig. 1I; α-SMA panel,

P=0.0218 *vs.* 2 Mo group, n=6; *P*=0.0088 *vs.* 6 Mo group, n=6; *P*=0.0005 *vs.* 12 Mo

group, n=6). Thus, PANK4 was correlated with age and age-dependent lens

epithelium fibrosis.

PANK4 is required for lens epithelium fibrosis and EMT

We next considered whether PANK4 modification alters EMT. Dramatic decrease was

observed in levels of N-cadherin (*P*=0.0065, n=6), collagen I (*P*=0.0026, n=6), and

α-SMA (*P*=0.005, n=6) in the lens epithelium of 2-Mo *Pank4*$^{-/-}$ mice (Fig. 2A and B).

To determine whether PANK4 is required for age-related lens fibrosis, PANK4

expression was analyzed in 1-year-old WT and *Pank4*$^{-/-}$ mice. Similarly, N-cadherin

(P=0.0198, n=6), collagen I (P=0.038, n=6), and α-SMA (P=0.0008, n=6) were also

inhibited when PANK4 was deficient (Fig. 2C and D). TGF-β2, the major isoform in

the aqueous humor, acts as a predominant pathogenic factor in the development of

lens epithlium EMT[49, 50]. To determine the effect of loss of function of PANK4 in

EMT alleviation, we next established a lens fibrosis model by TGF-β2 injection into

mice eyes. Compared with the WT group, $Pank4^{-/-}$ mice showed a dramatic decrease

in N-cadherin (P=0.0422 in WT vs WT+TGFβ2 group, n=6; P=0.0075 in WT vs

$Pank4^{-/-}$ group, n=6; P<0.0001 in $Pank4^{-/-}$ vs $Pank4^{-/-}$+ TGFβ2 group, n=6), collagen I

(P=0.0189 in WT vs WT+TGFβ2 group, n=6; P=0.0336, in WT vs $Pank4^{-/-}$ group,

n=6; P<0.0001 in $Pank4^{-/-}$ vs $Pank4^{-/-}$+ TGFβ2 group, n=6), and α-SMA (P<0.0011 in

WT vs WT+TGFβ2 group, n=6; P=0.0242, in WT vs $Pank4^{-/-}$ group, n=6; P<0.0001

in $Pank4^{-/-}$ vs $Pank4^{-/-}$+ TGFβ2 group, n=6) levels with and without TGF-β2 injection

(Fig. 2E and F). The alleviation of lens fibrosis was either found in TGF-β2-treated

groups or lens puncture groups (Fig. 2G) compared with WT mice. Thus, PANK4 was

required for lens epithlium EMT. To confirm the effect of loss of function of PANK4,

siRNA^{Pank4} was used to knock it down in SRA01/04 cells. Here, we assessed three

siRNA sequences—#274, #962, and #1114, of which #1114 showed almost 80%

inhibition of PANK4, whereas the other two siRNAs exhibited attenuated inhibitory

effect of nearly 30% (Fig. S1A). After 48-h interference of $Pank4$ mRNA, the

collagen I (P=0.0038 in NC vs siPank4 group, n=6; P=0.0048 in siControl vs siPank4

group, n=6), N-cadherin (P=0.0381 in NC vs siPank4 group, n=6; P=0.0369 in

siControl vs siPank4 group, n=6) and α-SMA (P<0.0001 in NC vs siPank4 group,

n=6; P=0.0002 in siControl vs siPank4 group, n=6) levels remarkably decreased (Fig.

2H and I). These results were consistent with the decrease in the SRA01/04 cell

migration test (Fig. 2J and K) (P<0.0001 in NC vs siPank4 group, n=7; P<0.0001 in

siControl vs siPank4 group, n=7; 24h group), in which siRNA^{Pank4} inhibited the

migration rate of SRA01/04 cells. PANK4 overexpression by Lentivirus–$Pank4$

$(Pank4$ OE$)$ accelerated the migration (Fig. 2J and K) (P<0.0001 in NC vs $Pank4$ OE

group, n=7; P<0.0001 in siControl vs $Pank4$ OE group, n=7; 12h and 24h groups,

respectively), which suggested that PANK4 was implicated in LEC EMT.

PANK4 negatively regulates PKM2 in LECs

PKM2 is considered as an associated protein that interacts with rat PANK4, which is

isolated from the rat muscle cDNA library by yeast two-hybrid screening[30]. In

addition, by STRING protein–protein interaction networks functional enrichment

analysis (https://string-db.org/), we found a potential relationship between PANK4

and PKM with a score of 0.705. The Co-IP blot results showed that PKM2

significantly increased on immunoprecipitation with PANK4, whereas PKM1 did not.

To verify their binding relationship, PKM2 antibody was used to bait the protein,

which showed significant PANK4 immunoprecipitation (Fig. 3A), indicating the

binding relationship between PANK4 and PKM2. It clearly indicated the specific

location of PKM2 in the lens epithelium, and the co-staining pattern of PKM2 and

PANK4 in the LEC cytoplasm (Fig. 3B). We then investigated their regulatory

mechanism using $Pank4^{-/-}$ mice, which demonstrated a significantly elevated level of

PKM2 in the lens epithelium by immunostaining (Fig. 3C and D) as well as Western

blotting results (Fig. 3E and F; PANK4: $P<0.0001$, n=6; PKM2: $P<0.0001$, WT vs

Pank4$^{-/-}$, n=6), but with no significant difference in PKM1 expression between WT

and *Pank4*$^{-/-}$ groups ($P=0.805$, WT *vs. Pank4*$^{-/-}$, n=6). By siRNA^{Pank4} knockdown of

Pank4 mRNA in SRA01/04, a remarkable increase was observed in PKM2 expression

in the cytoplasm (Fig. 3G), which was consistent with the results of the genome

deficiency of *Pank4* in the lens epithelium, followed by a Western blot confirmation

of the elevated PKM2 protein level (Fig. 3H and I, $P=0.0002$, siCont *vs.* siPank4;

$P<0.0001$, NC *vs.* siPank4, n=6) without a significant change in PKM1 (Fig. 3H and I;

$P=0.9742$, siCont *vs.* siPank4, n=6). TGF-β2 injection induced lens epithelium

fibrosis, and *Pank4* deletion increased the PKM2 level as well (Fig. 3J and K;

$P=0.0069$, WT *vs. Pank4*$^{-/-}$; $P=0.0036$, WT+TGFβ *vs. Pank4*$^{-/-}$+TGFβ, n=6),

suggesting the negative regulation between PANK4 and PKM2. In the conversion of

phosphoenolpyruvate (PEP) to pyruvic acid, PEP is alternatively catalyzed by

PKM2[53]. Strikingly, pyruvic acid increased when PANK4 was inhibited in LECs

(Fig. 3L; $P < 0.0001$, siControl *vs.* siPank4, n=6). These data suggest a role for the

PANK4–PKM2 axis in glycolysis in the lens epithelium (Fig. 3M).

Inhibition of PKM2 attenuates lens epithelium EMT by promoting PKM2

phosphorylation at Y105 residue

The inhibition and activation of PKM2 by PKM-IN-1 and DASA-58, which are

respectively typical antagonist and agonist of PKM2, did not significantly change the

PANK4 level, suggesting a downstream role for PKM2. In the presence of PKM2

inhibition or activation by PKM-IN-1 or DASA-58, the N-cadherin levels (P=0.0033,

Veh *vs.* DASA-58 groups, n=6), collagen I (P=0.0002, Veh *vs.* DASA-58 groups, n =

6), and α-SMA (P=0.0007, Veh *vs.* DASA-58 groups, n=6) significantly decreased

when activated by DASA-58. However, only N-cadherin and α-SMA levels increased

when treated by PKM-IN-1. (N-cad: P=0.0138, Veh *vs.* PKM-IN-1 groups, n=6;

Col-1: P=0.1066, Veh *vs.* PKM-IN-1 groups, n=6; α-SMA: P=0.0019, Veh *vs.*

PKM-IN-1 groups, n=6;) (Fig. 4A and B). These results indicate an important role for

PKM2 downstream of PANK4 in regulating LEC EMT.

To confirm the downstream role of PKM2 that was negatively regulated by

PANK4, PKM-IN-1 was used to inhibit PKM2 elevation when PANK4 was ablated.

As expected, PKM2 expression was decreased but that of phosphor-PKM2 was

elevated, as well as those of EMT markers, such as α-SMA, collagen I, and

N-cadherin (Fig. 4C). Because PKM2 activation attenuates the lens EMT phenotype,

PKM-IN-1 was injected into the anterior chamber in the TGF-β2-induced lens

epithelium fibrosis model. By slit-lamp observation, an opaque anterior lens (fibrosis)

was observed in the PKM-IN-1-treated group at 3 d, but a transparent lens was

observed in the Veh group (Fig. 4D), which revealed PANK4 deficiency-dependent

activation of PKM2.

Since we found PANK4 deficiency-related PKM2 elevation, we ask further

mechanisms. A previous study indicated that cytoplasmic PKM2 functions with

tetramer formation, whereas nuclear PKM2 functions with dimer/monomer formation

that may provide protein-binding ability[54]. Here, we found increased expression of

the 60-kDa band and 120-kDa band, corresponding to monomeric and dimeric PKM2

(Fig. 4E and F; P=0.0286, WT $vs.$ $Pank4^{-/-}$, respectively, n=4). Further, p-PKM2

(Y105) showed no significant difference between WT and $Pank4^{-/-}$ groups, but

p-PKM2 (S37) indeed declined (Fig. 4G and H; P =0.6683, WT $vs.$ $Pank4^{-/-}$, n=6, in

the Y105 group; P =0.0022, WT $vs.$ $Pank4^{-/-}$, n=6, in the S37 group), which

suggested that all relative increase in t-PKM2 was due to phosphorylation of the Y105

residue. To investigate the cellular location of p-PKM2 (Y105 and S37),

immunofluorescence staining was performed in WT and $Pank4^{-/-}$ LECs. p-PKM2

(Y105) signals were mainly located in the LEC nucleus in the WT group but in the

LEC cytoplasm in the $Pank4^{-/-}$ group (Fig. 4I). Plot profile of immunofluorescence

signal described the p-PKM2 (Y105) level in the lens epithelium of WT and $Pank4^{-/-}$

groups (Fig. 4J). To investigate the cellular location of p-PKM2 (Y105), a transection

of the single cellular profile was delineated (Fig. 4K), supporting the

immunofluorescence results that p-PKM2 (Y105) was mainly located in the

cytoplasm and nucleus. Meanwhile, p-PKM2 (S37) immunofluorescence signals were

punctate and located in the cytoplasm and nucleus of WT LECs; however, few

punctate signals were found in $Pank4^{-/-}$ mice (Fig. 4L). Plot profile analysis indicated

lower levels of p-PKM2 (S37) optical density values in both cytoplasm and nucleus

(Fig. 4M). For a single LEC, the curve indicated a typical lower level of p-PKM2

(S37) in the LEC nucleus in $Pank4^{-/-}$ mice (Fig. 4N). These results indicate that

PANK4 deficiency contributes to the increase in monomeric and dimeric forms of

p-PKM2 (Y105) but not of p-PKM2 (S37), as indicated in the schematic diagram (Fig.

4O).

PANK4 deficiency is responsible for PKM2 phosphorylation at Y105 and HIF-1α

activation

PKM2 was previously reported as a PHD3-stimulated coactivator for hypoxia

inducible factor-1 (HIF-1) to favor the genes associated with glycolysis[55]. Tyrosine

phosphorylation of PKM2 at Y105 inhibits the formation of a PKM2 tetramer by

dislodging fructose 1,6-bisphosphate (FBP) binding, which was demonstrated to be

important for the Warburg effect in hypoxia and cancer[56]. Thus, we speculated the

relationship between PKM2 and HIF-1 signaling may contribute to lens EMT

attenuation. We performed Co-IP assay to investigate why PANK4 deletion triggered

LEC to a relatively hypoxic condition. Notably, binding activity in the lens epithelium

was higher in WT (Fig. 5A), which demonstrated that PKM2 (Y105) physically

interacted with HIF-1α. However, no binding activity of PANK4 with HIF-1 was

found in *Pank4$^{-/-}$* mice (Fig. 5A). To verify the glycolysis status of PANK4-deficient

lens epithelium, levels of HIF-1α, HIF-1β, and HIF-2α, which are typical cellular

oxygen sensors, were assessed. As expected, all three hypoxic sensors were

dramatically elevated in *Pank4$^{-/-}$* mice (Fig. 5B; Hif-1α, $P=0.0004$, WT *vs. Pank4$^{-/-}$*,

n=6; Hif-1β, $P=0.0022$, WT *vs. Pank4$^{-/-}$*, n=6; Hif-2α, $P=0.0075$, WT *vs. Pank4$^{-/-}$*,

n=6), supporting the conclusion that PANK4 was a negative regulator in the shift of

OXPHOS and glycolysis. Interestingly, the lactate concentration was also increased in

the lens epithelium but not in blood (Fig. 5C; $P=0.0001$, WT *vs. Pank4$^{-/-}$* in the lens

group, n=4). Although PANK4 is a putative kinase in the PANK family, all amniote

PANK4s, including human PANK4, encode Glu138Val and Arg207Trp substitutions

that might deactivate kinase activity[57]; thus, its exact role is still under

consideration. To investigate the causative role and mechanisms of PANK4 in EMT,

we screened the mRNAs of the lens epithelium by RNA-seq in WT and $Pank4^{-/-}$ mice

(n=4) and performed gene set enrichment analysis (GSEA). We found that "electron

transport chain", "respiratory electron transport", and "oxidative phosphorylation"

were enriched in the WT group, which demonstrates the potential role of PANK4 in

negatively regulating glycolysis. In addition, GSEA showed an enrichment of the

"Glycolysis" gene cluster in the $Pank4^{-/-}$ group (Fig. 5D). To assess the

transformation of OXPHOS to glycolysis by PANK4, Seahorse real-time detection

was used to measure OCR and ECAR when Pank4 was knocked down by siRNA$^{Pank4.}$

No significant difference in basal respiration was found between the two groups;

however, spare respiratory capacity and maximum respiration were decreased in the

siRNA^{Pank4} group. Non-mitochondrial oxygen consumption was dramatically

increased (Fig. 5E and F; spare respiratory capacity, P=0.0286, siCont $vs.$ siPank4,

n=4; non-mitochondrial oxygen consumption, P=0.0286, siCont $vs.$ siPank4, n=4;

maximum respiration, P=0.0286, siCont $vs.$ siPank4, n=4). The ratios of glycolytic

rate, glycolytic capacity, and glycolysis reserve were all significantly increased when

$Pank4$ was knocked down (Fig. 5G and H; glycolytic rate, P=0.0317, siCont $vs.$

siPank4, n=5; glycolysis capacity, P=0.0317, siCont $vs.$ siPank4, n=5; glycolysis

reserve, P=0.0079, siCont $vs.$ siPank4, n=5). These evidences suggest the essential

role of PANK4 in negatively regulating glycolysis. To verify this potential, the

pyruvic acid and ATP levels were assessed in the control and siRNA^{Pank4} groups.

Dramatic increase in pyruvic acid was found when *Pank4* was knocked down, but

reduction in ATP was observed compared with the siControl group (Fig. 5I and J;

pyruvic acid concentration, *P*=0.0286, siCont *vs.* siPank4, n=4; concentration of ATP,

P=0.0211, siCont *vs.* siPank4, n=4). To confirm the possible role of PANK4 in

catalyzing CoA, we assessed CoA production under WT and Pank4 knockout

conditions. Similarly, as previously reported[57], no significant difference in CoA was

observed between the two groups in lens (Fig. S2A), suggesting a putative alternative

role of PANK4 in regulating EMT. To investigate how deficiency of PANK4

contributes to the incline of glycolysis or OXPHOS, we assessed acetyl CoA

carboxylase (ACC) and found an unexpected elevation (Fig. S2B). Because ACC is

the key enzyme in catalyzing the step of acetyl-CoA+ATP to malonyl-CoA+ADP/Pi,

and the evidence showed that the ATP level decreased in the *Pank4*$^{-/-}$ group (Fig. 5J),

we speculated the step exhaust amount of ATP.

To understand OXPHOS to glycolysis with EMT by PKM2 facilitation, Seahorse

real-time detection was used to measure mitochondrial stress during the procedure

(from 0 to 80 min). An increase in ECAR after adding glucose indicated the cell

glycolysis level, whereas a decrease in OCR after adding oligomycin indicated the

level of ATP production or OXPHOS of cells. By PKM-IN-1 treatment, glycolysis

rate, glycolysis capacity and glycolysis reserve ECAR values were all decreased

compared with the *Pank4* knockdown group (Fig. 5K and L; glycolysis, *P*<0.0001,

PKM-IN-1+siPank4 *vs.* siPank4, n=6; glycolysis capacity, *P*<0.0001,

PKM-IN-1+siPank4 *vs.* siPank4, n=6; *P*<0.0001, PKM-IN-1+siPank4 *vs.* siPank4,

n=6). DASA-58 inhibited the Pank4 knockdown-related ECAR value elevation in

glycolysis rate and glycolysis capacity (Fig. 5K and L; glycolysis, *P*<0.0001,

DASA-58+siPank4 *vs.* siPank4, n=6; glycolysis capacity, *P*<0.0001,

DASA-58+siPank4 *vs.* siPank4, n=6; *P*=0.1972, DASA-58+siPank4 *vs.* siPank4, n=6).

Conversely, no difference of OCR in spare respiratory capacity, basal respiration,

maximum respiration, proton leak and ATP production were found between the

siPank4 and DASA-58+sipank4 groups, suggesting the glycolysis regulation of

PANK4 by PKM2 (Fig. 5M, N and O; spare respiratory capacity, *P* = 0.0002,

PKM-IN-1 *vs.* siPank4, n = 6; *P* = 0.1972, siPank4 *vs.* DASA-58+siPank4, n = 6; *P* =

0.0006, PKM-IN-1+siPank4 *vs.* siPank4, n = 6. Basal respiration, *P* < 0.0001,

PKM-IN-1 *vs.* siPank4, n = 6; *P* = 0.9596, siPank4 *vs.* DASA-58+siPank4, n = 6.

Maximum respiration, *P* < 0.0001, PKM-IN-1 *vs.* siPank4, n = 6; *P* < 0.0001,

PKM-IN-1+siPank4 *vs.* siPank4, n = 6. Proton leak, *P* = 0.0022, PKM-IN-1 *vs.*

siPank4, n = 6; *P*=0.0055, PKM-IN-1+siPank4 *vs.* siPank4, n =6. ATP production, *P* <

0.0001, PKM-IN-1 *vs.* siPank4, n=6). The ATP level strikingly decreased in the

DASA-58 treatment group but not in the PKM-IN-1 group (Fig. 5P; *P*=0.0341, Veh *vs.*

DASA-58, n = 3), which suggested that DASA-58 was effective in shifting

PKM2-dependent OXPHOS and glycolysis. These results directly reflect the lower

level of glycolytic activity of SRA01/04 cells by DASA-58 treatment, which could be

rescued by *Pank4* knockdown. However, PKM-IN-1 did not effectively function in

both ECAR and OCR analyses.

Overall, the results provide evidence that PANK4 is a regulator of EMT in lens

fibrosis. PANK4 deficiency contributed to the increase in LEC glycolysis, with ATP

exhaustion and thus HIFs elevation. These results indicate that PANK4 deletion may

contribute to the Warburg effect by alleviating pyruvate generation. We also found a

relationship between PANK4 and PKM2, in which PANK4 negatively regulated

PKM2 elevation and monomer phosphorylation at Y105. Both inhibition of PANK4

and activation of PKM2 by DASA-58 resulted in EMT attenuation, which provides a

potential target for lens fibrosis and PCO treatment, as indicated in the schematic

diagram (Fig. 5Q).

Discussion

Cataract is the leading cause of blindness worldwide, although it can be effectively

cured by surgery. EMT is central to fibrotic cataract[58]. In the present study, PANK4,

which is a PANK family member, played a vital role in LEC EMT by regulating the

balance between OXPHOS and glycolysis. We also found a potential relationship

between PANK4 and PKM2, and glycolysis-related HIF elevation. Consequently, we

propose a model for the regulation of PKM2 in response to PANK4. PANK4 was

strong enough to control LEC EMT; when it was knockout or knockdown, the

expression levels of typical EMT markers, such as N-cadherin, collagen I, and α-SMA,

were decreased.

PANK4 is located on chromosome 1p36.32[59] and was first described as a

putative causative gene in type 2 diabetes in the Chinese population[23]. High glucose

could upregulate PANK4 in rat muscle [30]. However, till now limited studies have

investigated the mechanism further. Blood of diabetic patients was collected to

examine the putative relationship between PANK4 and diabetes. Unfortunately, no

significant difference between normal and diabetic individuals was observed in the

PANK4 mRNA level (Fig. S3), which suggested that PANK4 is not a candidate

causative gene for diabetes. Notably, Liu et al. reported that high PANK4 expression

high PANK4 expression had shorter event-free survival and overall survival than

patients with low PANK4 expression, and was indicative of poor prognosis[60], both

of which need further investigation. Because of the anti-apoptotic effect of PANK4,

the high level of PANK4 might inhibit pro-caspase-9 in pancreas, thus resulting in

low patient survival. A *Pank4*$^{-/-}$ mouse line was generated using the CRISPR/Cas9

system that showed an increased number of apoptotic LECs than the WT group[22].

Therefore, PANK4 may have a prominent anti-apoptotic role not only in pancreatic

and myeloid cells but also in LECs.

The human genome encodes three identified PANKs (i.e., PANK1–3) plus a

putative bifunctional protein, PANK4, with a predicted amino-terminal type II PANK

domain and a carboxy-terminal phosphatase domain (DUF89) that is large, diverse,

and present in all domains of life[20, 61, 62]. Members of the DUF89 protein family

(Pfam01937) occur as C-terminal fusions with PANK in plants, animals, and chytrid

fungi. Human PANK4 has activity against 4′-phosphopantetheine, and its S-sulfonate

or sulfonate is comparable to that against nitrophenyl phosphate[61]. However, Yao et

al. reported that all amniote PANK4s, including human PANK4, encode Glu138Val

and Arg207Trp substitutions that may lack PANK activity[57]. Nonetheless,

minimally characterized but highly conserved PANK4 is a rate-limiting suppressor of

CoA synthesis by its metabolite phosphatase activity[63], demonstrating an essential

role for PANK4 in the regulation of CoA-related metabolism. To inhibit the function

of PANK4, *Pank4$^{-/-}$* mice that have a 5-bp mutation before exon 3 were first designed

and generated. Notably, no significant difference in concentration of acetyl-CoA was

found in the lens between WT and *pank4$^{-/-}$* mice, which demonstrated that PANK4

may be not involved in the CoA biosynthetic in lens. But we could not exclude the

role of PANK4 in CoA biosynthetic in other tissues. Nonetheless, GSEA revealed a

noteworthy tendency of OXPHOS, electron transport chain, respiratory electron

transport, and ATP synthesis enriched in WT mice rather than in *Pank4$^{-/-}$* mice (Fig.

5D), suggesting the ability of PANK4 in redox regulation.

In a yeast two-hybrid system, PKM2 was associated with PANK4[30]. In the

present study, we found a regulatory relationship between PANK4 and PKM2. On

PANK4 loss of function, PKM2 significantly increased, suggesting a negative

relationship between them, but PKM1 was unchanged (Fig. 3E). PKM2 controls the

balance between energy production and metabolic synthesis precursors, and has both

metabolic and non-metabolic functions, which are essential in the cytoplasm and

nucleus, respectively. Tetrameric PKM2 functions as pyruvate kinase, whereas

nuclear PKM2 forms a dimer and functions as protein kinase[54], which interacts

with STAT3 to control downstream gene expression in SW480 cells. However, STAT3

and p-STAT3 were unchanged in the lens epithelium of *Pank4*$^{-/-}$ mice in Western

blotting test (Fig. S4), which suggested that the nuclear dimeric PKM2 regulated

non-metabolic function independent of classic STAT3 signaling. PKM2 activity is

inhibited after phosphotyrosine binding by the release of FBP from the PKM2

allosteric pocket by Y105 phosphorylation that disrupts tetrameric PKM2[56, 64].

Monomeric PKM2 increased under PANK4 deletion, which indicates that the

glycolytic metabolic function was not likely because of tetrameric PKM2. Thus,

PKM2 might not contribute to the glycolysis of LEC by tetramer formation under

Pank4 ablation but is involved in p-PKM2-related signaling transduction.

PKM2-dependent histone H3 modifications are instrumental in epidermal growth

factor (EGF)-induced expression of cyclin D1 and c-Myc, tumor cell proliferation,

cell cycle progression, and brain tumorigenesis in its non-metabolic functions of

histone modification[65]. In this study, monomeric PKM2 were increased, suggesting

that the metabolic activity of PKM2 is attenuated during LEC EMT. Thus, it is

conceivable that the molecular interaction of PKM2 is highly context dependent, with

cell fate determined by the form of PKM2 in regulating gene expression. In addition,

PKM2 plays an essential role in Warburg regulation. Methylation of PKM2 activates

aerobic glycolysis to promote tumorigenesis[29]. In transformed tumor cells, PKM2 is

phosphorylated at Y105 by oncogenic kinases and dissociated to PKM2 (Y105)

dimers that promote yes-associated protein (YAP) nuclear translocation for tumor cell

growth and transformation[66]. In this study, although PKM2 Y105 was dramatically

elevated with PANK4 loss of function, the total PKM2 also increased simultaneously,

demonstrating the consistent ratio of Y105 phosphorylation to total PKM2 protein, in

other words, the elevated PKM2 protein was phosphorylated at Y105. Moreover, the

ratio of PKM2 (S37) to total PKM2 decreased, but no significant difference was

observed in the ratio of PKM2 (S37) to the inner control (Fig. 4G and H). In

hepatocellular carcinoma cell lines, PKM2 could be phosphorylated at Ser37 by EGF

receptor-activated extracellular signal-regulated kinase when stimulated with EGF,

which promoted its nuclear translocation. By acting as both serine/threonine protein

kinase and tyrosine protein kinase, PKM2 binds to and phosphorylates multiple

transcription factors such as β-catenin and STAT3 in the nucleus to promote

tumor-related gene transcription. H3-Thr11 phosphorylation is dependent on the

phosphorylation of PKM2 at Ser37[67]. Herein, to clarify whether PKM2 (S37)

phosphorylates STAT3 under PANK4 deletion, STAT3 and p-STAT3 were both

detected, but no change was observed (Fig. S4). Immunostaining of PKM2 (S37) in

the lens epithelium was attenuated in the $Pank4^{-/-}$ group, which strengthens the

conclusion that S37 residue-related PKM2 transnuclear function may not be involved

in PANK4-unlocked PKM2 elevation.

The lens is sustained in a hypoxic environment in the eye without any vascular

supply. The oxygen partial pressure at the lens surface is 2.9 mmHg, which is lower

than 24.1 mmHg at the surface of the eye[68]. Thus, PKM2 activity contributes to a

high glycolytic potential to support lens metabolism[69-71]. PKM2 can catalyze the

final rate-limiting step of glycolysis, and it functions as a dimer when translocating to

the nucleus, where it stimulates HIF-1α transactivation domain function and

recruitment of p300 to hypoxia response[55]. In the present study, we found that

HIF-1α and HIF-1β increased dramatically in parallel with total PKM2 and PKM2

(Y105) elevation under PANK4 deletion. The current results showed a physical

relationship between PKM2 and HIF-1α in WT mice but not in $Pank4^{-/-}$ mice.

Because the lens epithelium was in a hypoxic environment in the WT condition,

PKM2 and HIF-1α might contribute to the homeostasis of LEC survival and the

balance of metabolism. After PANK4 was absent, more PKM2 was phosphorylated at

Y105 and dissociated with HIF-1α, along with favoring glycolysis, indicating a

negative regulation loop of PANK4 on hypoxia. Although $Pank4^{-/-}$ mice were housed

in a high-level oxygen (70%) condition for 7 continuous days, HIF-1α was still

elevated in the lens epithelium of $Pank4^{-/-}$ mice (Fig. S5, P=0.0021, WT(O2) vs

$Pank4^{-/-}$(O2), n=4), which suggested that HIF-1α elevation is caused not only by

hypoxic conditions but also by PANK4 regulation. In normoxic conditions, HIF-1α

can be hydroxylated in its C-terminal transcription activation domain by the arginyl

hydroxylase factor-inhibiting HIF[72]. Oxygen-dependent HIF-1α degradation is

mediated by the hydroxylation of prolines 402 and 564 in human HIF-1α. However,

the Warburg effect can activate PI3K and Akt signaling to stabilize HIF-1α in an

aerobic condition, which promotes pyruvate-related glycolysis as well[73]. Therefore,

the consistent elevation of HIF-1α is dependent on PANK4 deletion, irrespective of

normoxic, hypoxic, or hyperoxic conditions. Concerning the uncoupling result of

PKM2 and HIF-1α with PANK4 loss of function (Fig. 5A), we speculate that the

elevation of PKM2 and HIF-1α may function in different ways rather than the classic

PKM2-HIF-1-positive feedback loop. PANK4 may function as a coordinator that

organizes PKM2 and HIF-1α in cellular OXPHOS and glycolysis activities. In

addition, we cannot exclude the possibility that HIF-1α promotes PKM2 gene

transcription or PKM2 promotes HIF-1α gene transcription, which needs evidence. In

older animals, HIF-1α is necessary to maintain a high level of p27KIP1 that inhibits

LEC proliferation. Because we previously found an apoptotic phenotype of LECs in

$Pank4^{-/-}$ mice, the conclusion that PANK4 deficiency-dependent HIF-1α elevation

may inhibit the proliferation and EMT of LECs seems plausible.

Acknowledgement

We thank those heroes combating the 2019-nCov, especially Miss Ying Li and Juan

Yu in our department, with respect. We also thank Prof. Lei Li from Shanghai Tech

University for data evaluation. We thank our team members (Especially Jiao Hu, Xi

Chen and Qiu-Mei Hu) for their diligent work. We also thank Cyagen Inc. for mice

embryo cryopreservation and breeding. The project is supported by National Natural

Science Foundation of China (No. 82070945), Science and Technology Innovation

Enhancement Project of Army Medical Center, Army Medical University (No.

2019CXJSB009 and No. 2019CXJSB021), Army Medical University Outstanding

Talent (Chun-Lin Chen) and grants from Basic Research and Scientific Frontier

Foundation of Chongqing (cstc2019jcyj-msxmX0010).

References

[1]   Saika S, Yamanaka O, Sumioka T, Miyamoto T, Miyazaki K, Okada Y, *et al.*

(2008). Fibrotic disorders in the eye: targets of gene therapy. Prog Retin Eye

Res, 27:177-196.

[2]   Eldred JA, Dawes LJ, Wormstone IM (2011). The lens as a model for fibrotic

disease. Philos Trans R Soc Lond B Biol Sci, 366:1301-1319.

[3]   Piera-Velazquez S, Mendoza FA, Jimenez SA (2016). Endothelial to

Mesenchymal Transition (EndoMT) in the Pathogenesis of Human Fibrotic

Diseases. J Clin Med, 5.

[4]   Wormstone IM (2002). Posterior capsule opacification: a cell biological

perspective. Exp Eye Res, 74:337-347.

[5]   Lovicu FJ, McAvoy JW (1989). Structural analysis of lens epithelial explants

induced to differentiate into fibres by fibroblast growth factor (FGF). Exp Eye

Res, 49:479-494.

[6]   Saika S (2004). Relationship between posterior capsule opacification and

intraocular lens biocompatibility. Prog Retin Eye Res, 23:283-305.

[7]   Marcantonio JM, Syam PP, Liu CS, Duncan G (2003). Epithelial

transdifferentiation and cataract in the human lens. Exp Eye Res, 77:339-346.

[8]   Nagamoto T, Eguchi G (1997). Morphologic compatibility or intraocular lens

haptics and the lens capsule. J Cataract Refract Surg, 23:1254-1259.

[9]   Nagamoto T, Eguchi G (1997). Effect of intraocular lens design on migration

of lens epithelial cells onto the posterior capsule. J Cataract Refract Surg,

23:866-872.

[10]   Shu DY, Wojciechowski M, Lovicu FJ (2019). ERK1/2-mediated

EGFR-signaling is required for TGFbeta-induced lens epithelial-mesenchymal

transition. Exp Eye Res, 178:108-121.

[11]   Shu DY, Lovicu FJ (2019). Enhanced EGF receptor-signaling potentiates

TGFbeta-induced lens epithelial-mesenchymal transition. Exp Eye Res,

185:107693.

[12]   Raghavan CT, Smuda M, Smith AJ, Howell S, Smith DG, Singh A, *et al.*

(2016). AGEs in human lens capsule promote the TGFbeta2-mediated EMT of

lens epithelial cells: implications for age-associated fibrosis. Aging Cell,

15:465-476.

[13]   Roberts AB, Tian F, Byfield SD, Stuelten C, Ooshima A, Saika S, *et al.* (2006).

Smad3 is key to TGF-beta-mediated epithelial-to-mesenchymal transition,

fibrosis, tumor suppression and metastasis. Cytokine Growth Factor Rev,

17:19-27.

[14]   Swarup A, Bell BA, Du J, Han JYS, Soto J, Abel ED, *et al.* (2018). Deletion

of GLUT1 in mouse lens epithelium leads to cataract formation. Exp Eye Res,

172:45-53.

[15]   Dahm R, van Marle J, Quinlan RA, Prescott AR, Vrensen GF (2011).

Homeostasis in the vertebrate lens: mechanisms of solute exchange. Philos

Trans R Soc Lond B Biol Sci, 366:1265-1277.

[16]   Casula M, Soranna D, Corrao G, Merlino L, Catapano AL, Tragni E (2016).

Statin use and risk of cataract: A nested case-control study within a healthcare

database. Atherosclerosis, 251:153-158.

[17] Leuschen J, Mortensen EM, Frei CR, Mansi EA, Panday V, Mansi I (2013).

Association of statin use with cataracts: a propensity score-matched analysis.

JAMA Ophthalmol, 131:1427-1434.

[18] Dobrzynski JM, Kostis JB (2015). Statins and Cataracts--a visual insight. Curr

Atheroscler Rep, 17:477.

[19] Zhang YM, Chohnan S, Virga KG, Stevens RD, Ilkayeva OR, Wenner BR, *et*

al. (2007). Chemical knockout of pantothenate kinase reveals the metabolic

and genetic program responsible for hepatic coenzyme A homeostasis. Chem

Biol, 14:291-302.

[20] Hortnagel K, Prokisch H, Meitinger T (2003). An isoform of hPANK2,

deficient in pantothenate kinase-associated neurodegeneration, localizes to

mitochondria. Hum Mol Genet, 12:321-327.

[21] Hong BS, Senisterra G, Rabeh WM, Vedadi M, Leonardi R, Zhang YM, *et al.*

(2007). Crystal structures of human pantothenate kinases. Insights into

allosteric regulation and mutations linked to a neurodegeneration disorder. J

Biol Chem, 282:27984-27993.

[22] Sun M, Chen C, Hou S, Li X, Wang H, Zhou J*, et al.* (2018). A novel mutation

of PANK4 causes autosomal dominant congenital posterior cataract. Hum

Mutat.

[23] Li Y, Wu GD, Zuo J, Meng Y, Fang FD (2005). [Screening susceptibility genes

818   of type 2 diabetes in Chinese population by single nucleotide polymorphism

819   analysis]. Zhongguo Yi Xue Ke Xue Yuan Xue Bao, 27:274-279.

820   [24]   Xiang RL, Yang YL, Zuo J, Xiao XH, Chang YS, De Fang F (2007). PanK4

821   inhibits pancreatic beta-cell apoptosis by decreasing the transcriptional level

822   of pro-caspase-9. Cell Res, 17:966-968.

823   [25]   Zablocki GJ, Ruzycki PA, Overturf MA, Palla S, Reddy GB, Petrash JM

824   (2011). Aldose reductase-mediated induction of epithelium-to-mesenchymal

825   transition (EMT) in lens. Chem Biol Interact, 191:351-356.

826   [26]   Yadav UC, Ighani-Hosseinabad F, van Kuijk FJ, Srivastava SK, Ramana KV

827   (2009). Prevention of posterior capsular opacification through aldose

828   reductase inhibition. Invest Ophthalmol Vis Sci, 50:752-759.

829   [27]   Kinoshita JH, Nishimura C (1988). The involvement of aldose reductase in

830   diabetic complications. Diabetes Metab Rev, 4:323-337.

831   [28]   WJ I, TL D, SM D, BP F, AM H, G B, *et al.* (2013). PKM2 isoform-specific

832   deletion reveals a differential requirement for pyruvate kinase in tumor cells.

833   Cell, 155:397-409.

834   [29]   F L, F M, Y W, L H, H Z, C J, *et al.* (2017). PKM2 methylation by CARM1

835   activates aerobic glycolysis to promote tumorigenesis. Nature cell biology,

836   19:1358-1370.

837   [30]   Li Y, Chang Y, Zhang L, Feng Q, Liu Z, Zhang Y, *et al.* (2005). High glucose

838   upregulates pantothenate kinase 4 (PanK4) and thus affects M2-type pyruvate

839   kinase (Pkm2). Mol Cell Biochem, 277:117-125.

[31] Wong N, Ojo D, Yan J, Tang D (2015). PKM2 contributes to cancer

metabolism. Cancer Lett, 356:184-191.

[32] Li C, Zhao Z, Zhou Z, Liu R (2016). PKM2 Promotes Cell Survival and

Invasion Under Metabolic Stress by Enhancing Warburg Effect in Pancreatic

Ductal Adenocarcinoma. Dig Dis Sci, 61:767-773.

[33] M X, Y Y, R K, S Z, L Y, L Z, *et al.* (2016). PKM2-dependent glycolysis

promotes NLRP3 and AIM2 inflammasome activation. Nature

communications, 7:13280.

[34] Qi W, Keenan HA, Li Q, Ishikado A, Kannt A, Sadowski T, *et al.* (2017).

Pyruvate kinase M2 activation may protect against the progression of diabetic

glomerular pathology and mitochondrial dysfunction. Nat Med, 23:753-762.

[35] Liang J, Cao R, Zhang Y, Xia Y, Zheng Y, Li X, *et al.* (2016). PKM2

dephosphorylation by Cdc25A promotes the Warburg effect and tumorigenesis.

Nat Commun, 7:12431.

[36] Yang W, Zheng Y, Xia Y, Ji H, Chen X, Guo F, *et al.* (2012).

ERK1/2-dependent phosphorylation and nuclear translocation of PKM2

promotes the Warburg effect. Nat Cell Biol, 14:1295-1304.

[37] Yang W, Xia Y, Ji H, Zheng Y, Liang J, Huang W, *et al.* (2011). Nuclear

PKM2 regulates beta-catenin transactivation upon EGFR activation. Nature,

480:118-122.

[38] Yang W, Xia Y, Hawke D, Li X, Liang J, Xing D, *et al.* (2012). PKM2

phosphorylates histone H3 and promotes gene transcription and tumorigenesis.

Cell, 150:685-696.

[39]   Lovicu FJ, McAvoy JW, de Iongh RU (2011). Understanding the role of

growth factors in embryonic development: insights from the lens. Philos Trans

R Soc Lond B Biol Sci, 366:1204-1218.

[40]   Sun M, Chen C, Hou S, Li X, Wang H, Zhou J, *et al.* (2019). A novel mutation

of PANK4 causes autosomal dominant congenital posterior cataract. Hum

Mutat, 40:380-391.

[41]   Wang R, Li J, Zhang X, Zhang X, Zhang X, Zhu Y, *et al.* (2021). Extracellular

vesicles promote epithelial-to-mesenchymal transition of lens epithelial cells

under oxidative stress. Exp Cell Res, 398:112362.

[42]   Chen X, Xiao W, Chen W, Luo L, Ye S, Liu Y (2013). The epigenetic modifier

trichostatin A, a histone deacetylase inhibitor, suppresses proliferation and

epithelial-mesenchymal transition of lens epithelial cells. Cell Death Dis,

4:e884.

[43]   Hales AM, Chamberlain CG, McAvoy JW (1995). Cataract induction in lenses

cultured with transforming growth factor-beta. Invest Ophthalmol Vis Sci,

36:1709-1713.

[44]   Hales AM, Chamberlain CG, McAvoy JW (2000). Susceptibility to

TGFbeta2-induced cataract increases with aging in the rat. Invest Ophthalmol

Vis Sci, 41:3544-3551.

[45]   Choi J, Seo BJ, La H, Yoon SH, Hong YJ, Lee JH, *et al.* (2020). Comparative

analysis of the mitochondrial morphology, energy metabolism, and gene

expression signatures in three types of blastocyst-derived stem cells. Redox

Biol, 30:101437.

[46] Zhou JX, Liu YJ, Chen X, Zhang X, Xu J, Yang K, *et al.* (2018).

Low-Intensity Pulsed Ultrasound Protects Retinal Ganglion Cell From Optic

Nerve Injury Induced Apoptosis via Yes Associated Protein. Front Cell

Neurosci, 12:160.

[47] Zhang Z, Liu W, Huang Y, Luo L, Cai X, Liu Y, *et al.* (2020). NLRP3

Deficiency Attenuates Secondary Degeneration of Visual Cortical Neurons

Following Optic Nerve Injury. Neurosci Bull, 36:277-288.

[48] Sun H, Yang X, Liang L, Zhang M, Li Y, Chen J, *et al.* (2020). Metabolic

switch and epithelial-mesenchymal transition cooperate to regulate

pluripotency. EMBO J, 39:e102961.

[49] Chen X, Xiao W, Chen W, Liu X, Wu M, Bo Q, *et al.* (2017). MicroRNA-26a

and -26b inhibit lens fibrosis and cataract by negatively regulating

Jagged-1/Notch signaling pathway. Cell Death Differ, 24:1431-1442.

[50] Allen JB, Davidson MG, Nasisse MP, Fleisher LN, McGahan MC (1998). The

lens influences aqueous humor levels of transforming growth factor-beta 2.

Graefes Arch Clin Exp Ophthalmol, 236:305-311.

[51] Dawes LJ, Elliott RM, Reddan JR, Wormstone YM, Wormstone IM (2007).

Oligonucleotide microarray analysis of human lens epithelial cells: TGFbeta

regulated gene expression. Mol Vis, 13:1181-1197.

[52] Lee JM, Dedhar S, Kalluri R, Thompson EW (2006). The

epithelial-mesenchymal transition: new insights in signaling, development,

and disease. J Cell Biol, 172:973-981.

[53] Israelsen WJ, Vander Heiden MG (2015). Pyruvate kinase: Function,

regulation and role in cancer. Semin Cell Dev Biol, 43:43-51.

[54] X G, H W, JJ Y, X L, ZR L (2012). Pyruvate kinase M2 regulates gene

transcription by acting as a protein kinase. Molecular cell, 45:598-609.

[55] Luo W, Hu H, Chang R, Zhong J, Knabel M, O'Meally R, *et al.* (2011).

Pyruvate kinase M2 is a PHD3-stimulated coactivator for hypoxia-inducible

factor 1. Cell, 145:732-744.

[56] Hitosugi T, Kang S, Vander Heiden MG, Chung TW, Elf S, Lythgoe K, *et al.*

(2009). Tyrosine phosphorylation inhibits PKM2 to promote the Warburg

effect and tumor growth. Sci Signal, 2:ra73.

[57] Yao J, Subramanian C, Rock CO, Jackowski S (2019). Human pantothenate

kinase 4 is a pseudo-pantothenate kinase. Protein Sci, 28:1031-1047.

[58] Kalluri R, Neilson EG (2003). Epithelial-mesenchymal transition and its

implications for fibrosis. J Clin Invest, 112:1776-1784.

[59] Zhou B, Westaway SK, Levinson B, Johnson MA, Gitschier J, Hayflick SJ

(2001). A novel pantothenate kinase gene (PANK2) is defective in

Hallervorden-Spatz syndrome. Nat Genet, 28:345-349.

[60] Liu Y, Cheng Z, Li Q, Pang Y, Cui L, Qian T, *et al.* (2019). Prognostic

significance of the PANK family expression in acute myeloid leukemia. Ann

Transl Med, 7:261.

[61]   Huang L, Khusnutdinova A, Nocek B, Brown G, Xu X, Cui H, *et al.* (2016). A

family of metal-dependent phosphatases implicated in metabolite

damage-control. Nat Chem Biol, 12:621-627.

[62]   Tilton GB, Wedemeyer WJ, Browse J, Ohlrogge J (2006). Plant coenzyme A

biosynthesis: characterization of two pantothenate kinases from Arabidopsis.

Plant Mol Biol, 61:629-642.

[63]   Dibble CC, Barritt SA, Perry GE, Lien EC, Geck RC, DuBois-Coyne SE, *et al.*

(2022). PI3K drives the de novo synthesis of coenzyme A from vitamin B5.

Nature, 608:192-198.

[64]   HR C, MG VH, N W, JM A, LC C (2008). Pyruvate kinase M2 is a

phosphotyrosine-binding protein. Nature, 452:181-186.

[65]   W Y, Y X, D H, X L, J L, D X, *et al.* (2012). PKM2 phosphorylates histone H3

and promotes gene transcription and tumorigenesis. Cell, 150:685-696.

[66]   Zhou Z, Li M, Zhang L, Zhao H, Sahin O, Chen J, *et al.* (2018). Oncogenic

Kinase-Induced PKM2 Tyrosine 105 Phosphorylation Converts Nononcogenic

PKM2 to a Tumor Promoter and Induces Cancer Stem-like Cells. Cancer Res,

78:2248-2261.

[67]   Niu J, Li W, Liang C, Wang X, Yao X, Yang RH, *et al.* (2020). EGF promotes

DKK1 transcription in hepatocellular carcinoma by enhancing the

phosphorylation and acetylation of histone H3. Sci Signal, 13.

[68]   Helbig H, Hinz JP, Kellner U, Foerster MH (1993). Oxygen in the anterior

chamber of the human eye. Ger J Ophthalmol, 2:161-164.

[69]   Cheng HM, Chylack LT, Jr., Chien J (1978). Control pyruvate kinase activity

951   in the lens. Exp Eye Res, 27:39-44.

952   [70]   Cheng HM, Chylack LT, Jr., von Saltza I (1981). Supplementing glucose

953   metabolism in human senile cataracts. Invest Ophthalmol Vis Sci, 21:812-818.

954   [71]   Banroques J, Gregori C, Schapira F (1978). Aging of pyruvate kinase

955   isozymes in rabbit lens. Exp Eye Res, 27:427-434.

956   [72]   Hewitson KS, McNeill LA, Riordan MV, Tian YM, Bullock AN, Welford RW,

957   *et al.* (2002). Hypoxia-inducible factor (HIF) asparagine hydroxylase is

958   identical to factor inhibiting HIF (FIH) and is related to the cupin structural

959   family. J Biol Chem, 277:26351-26355.

960   [73]   Semenza GL (2010). HIF-1: upstream and downstream of cancer metabolism.

961   Curr Opin Genet Dev, 20:51-56.

Figure Legends

Figure 1. PANK4 elevated with age and related to lens fibrosis and EMT.

(**A**) Immunostaining of PANK4 in lens epithelium from patients aged at 17yrs, 47yrs

and 81yrs. The scale bar = 10μm. (**B**) Plot profile analysis of immunostaining in lens

epithelium of patients aged from 17yrs, 47yrs and 81yrs. (**C**) Quantification of

PANK4 immunostaining positive puncta per cell in LECs, from patients aged at 17yrs,

47yrs and 81yrs. n=19, $P<0.0001$. (**D**) The levels of PANK4 protein in lens epithelium

from female cataract patients (50-60yrs, 60-70yrs, 70-80yrs and more than 80yrs

groups) were determined by Western blotting assay and quantified by densitometry.

Data were analyzed by one way ANOVA plus Bonferroni post hoc test. All data

shown are mean ± SD. Relative expression is the fold changes relative to the 50-60yrs

group. *P*=0.384 *vs* 60-70yrs group, n=6; *P*=0.0073 *vs* 70-80yrs group, n=6; *P*=0.0028

vs 80yrs+ group, n=6. (**E**) The levels of PANK4 protein in lens epithelium from male

cataract patients (50-60yrs, 60-70yrs, 70-80yrs and more than 80yrs groups) were

determined by Western blotting assay and quantified by densitometry. Data were

analyzed by one way ANOVA plus Bonferroni post hoc test. All data shown are mean

± SD. Relative expression is the fold changes relative to the 50-60yrs group.

P=0.1514 *vs* 60-70yrs group, n=6; *P*=0.0096 *vs* 70-80yrs group, n=6; *P*<0.0001 *vs*

80yrs+ group, n=6. (**F**) Immunostaining pattern of PANK4 in the lens epithelium of

adult mice. The scale bar = 200μm. HE image presented an anatomy structure of lens.

Inner panel: High power images of PANK4 immunostaining results in anterior lens

epithelium and equator of adult WT mice. The scale bar = 10μm. (**G**) The levels of

PANK4 protein in lens epithelium of from P15, 2Mo, 6Mo and 12Mo mice, detected

by Western blotting assay and quantified by densitometry. Data were analyzed by one

way ANOVA plus Bonferroni post hoc test. All data shown are mean ± SD. Relative

expression is the fold changes relative to the P15 group. *P*=0.0007 vs 2Mo group, n=6;

P=0.011 vs 6Mo group, n=6; *P*=0.0003 vs 80yrs+ group, n=6. (**H** and **I**) The levels of

N-cadherin, collagen I and α-SMA protein in lens epithelium of from P15, 2Mo, 6Mo

and 12Mo mice, detected by Western blotting assay and quantified by densitometry.

Data were analyzed by one way ANOVA plus Bonferroni post hoc test. Relative

expression is the fold changes relative to the P15 group. Results are expressed as

means ± SD. N-cad panel, *P*=0.0317 *vs* 2Mo group, n=6; *P*<0.0001 *vs* 6Mo group,

n=6; *P*<0.0001 *vs* 12Mo group, n=6. Col I panel, *P*=0.0361 *vs* 2Mo group, n=6;

P=0.0018 *vs* 6Mo group, n=6; *P*=0.0005 *vs* 12Mo group, n=6. α-SMA panel,

P=0.0218 *vs* 2Mo group, n=6; *P*=0.0088 *vs* 6Mo group, n=6; *P*=0.0005 *vs* 12Mo

group, n=6.

1019

1020

Figure 2. PANK4 is required for lens epithelium fibrosis and EMT

(**A**) and (**B**) The levels and quantifications of N-cadherin, collagen I and α-SMA

protein in 2-month-old (2Mo) WT and *Pank4*$^{-/-}$ lens epithelium. anti-PANK4 as

positive control. Data were analyzed by student's *t* test. Relative expression is the fold

changes relative to the 2Mo WT group. N-cadherin, *P*=0.0065, n=6; collagen I,

P=0.0026, n=6; α-SMA, *P*=0.005, n=6. (**C**) and (**D**) The levels and quantifications of

N-cadherin, collagen I and α-SMA protein in 12-month-old (12Mo) WT and *Pank4*$^{-/-}$

lens epithelium. anti-PANK4 as positive control. Data were analyzed by student's *t*

test. Relative expression is the fold changes relative to the 12Mo WT group.

N-cadherin, *P*=0.0198, n=6; collagen I, *P*=0.038, n=6; α-SMA, *P*=0.0008, n=6. (**E**)

and (**F**) The levels and quantifications of N-cadherin, collagen I and α-SMA protein in

2-month-old (2Mo) WT and *Pank4*$^{-/-}$ lens epithelium, with and without TGF-β2

injection. Data were analyzed by one way ANOVA plus Bonferroni post hoc test.

Relative expression is the fold changes to comparable groups. N-cadherin: *P*=0.0422

in WT vs WT+TGFβ2 group, n=6; *P*=0.0075 in WT vs *Pank4*$^{-/-}$ group, n=6; *P*<0.0001

in *Pank4*$^{-/-}$ vs *Pank4*$^{-/-}$+ TGFβ2 group, n=4. collagen I: *P*=0.0189 in WT vs

WT+TGFβ2 group, n=6; *P*=0.0336, in WT vs *Pank4*$^{-/-}$ group, n=6; *P*<0.0001 in

Pank4$^{-/-}$ vs *Pank4*$^{-/-}$+ TGFβ2 group, n=6. α-SMA: *P*=0.0011 in WT vs WT+TGFβ2

group, n=6; *P*=0.0242, in WT vs *Pank4*$^{-/-}$ group, n=6; *P*<0.0001 in *Pank4*$^{-/-}$ vs

Pank4$^{-/-}$+ TGFβ2 group, n=6. (**G**) Cultured lens with TGF-β2 (5.0ng/mL) addition for

2 and 4 days (solid arrow indicated fibrosis in the lens). In vivo lens fibrosis

phenotype, 7 days after puncture experiment in WT and *Pank4*$^{-/-}$ mice. Observed and

recorded by slit light microscope. (**H**) and (**I**) The levels and quantifications of

N-cadherin, collagen I, α-SMA and PANK4 protein after siRNA administration. Data

were analyzed by one way ANOVA plus Bonferroni post hoc test. Relative expression

is the fold changes relative to the NC group. N-cadherin: *P*=0.0381 in NC vs siPank4

group, n=6; *P*=0.0369 in siControl vs siPank4 group, n=6; Collagen I: *P*=0.0038 in

NC vs siPank4 group, n=6; *P*=0.0048 in siControl vs siPank4 group, n=6; α-SMA:

P<0.0001 in NC vs siPank4 group, n=6; *P*=0.0002 in siControl vs siPank4 group, n=6.

PANK4: *P*<0.0001 in NC vs siPank4 group, n=6; *P*<0.0001 in siControl vs siPank4

group, n=6. (**J**) and Lens epithelial cell migration assay at 0h, 12h and 24h, with Lenti

virus-*Pank4* overexpression and siRNA interference treatments. (**K**) Quantifications

of the wound healing rate (%) at 0h, 12h and 24h. Results are expressed as means ±

SD. *P*<0.0001 in NC vs *Pank4* OE group, n=7; *P*<0.0001 in siControl vs *Pank4* OE

group, n=7; 12h and 24h groups, respectively.

1056

1057

Figure 3. PANK4 negatively regulating PKM2 in lens epithelium

(**A**) Co-IP results of PANK4 and PKM2. (**B**) Co-Immunofluorescence staining of

PKM2 and PANK4 in lens epithelium. The scale bar = 10μm. (**C**)

Immunofluorescence staining of PKM2 in lens epithelium of WT and *Pank4^-/-* lens.

The scale bar = 10μm. (**D**) Plot profile curve of PKM2 in WT and *Pank4^-/-* lens. (**E**)

and (**F**) Protein levels and quantifications of PANK4, PKM1, PKM2 in WT and

Pank4^-/- lens. GAPDH as inner control. Data were analyzed by student's *t* test.

Relative expression is the fold changes relative to the WT group. PANK4: *P*<0.0001,

n=6; PKM1: *P*=0.805 PKM2: *P*<0.0001, WT vs *Pank4^-/-*, n=6. (**G**)

Immunofluorescence staining results of PKM2 in LECs in siControl group and

si*Pank4* group. The scale bar = 10μm. (**H**) and (**I**) Protein levels and quantifications

of PANK4, PKM1, PKM2 in si-control group and siPank4 group. Data were analyzed

by one way ANOVA plus Bonferroni post hoc test. Relative expression is the fold

changes relative to the siPank4 group.   PKM2: *P*<0.0001, NC vs siPank4; *P*=0.0002,

siCtrl vs siPank4, n=6. PKM1: *P*=0.9437, NC vs siPank4, n=6; *P*=0.9742, siCtrl vs

siPank4, n=6. PANK4: *P*<0.0001, NC vs siPank4; *P*<0.0001, siCtrl vs siPank4, n=6.

(**J**) and (**K**) Protein levels and relative expression quantifications of PKM2 in lens

epithelium of WT, WT with TGF-β2 injection, *Pank4^-/-*, *Pank4^-/-* with TGF-β2

injection groups. Data were analyzed by one way ANOVA plus Bonferroni post hoc

test. Relative expression is the fold changes relative to the comparable

groups.*P*=0.0069, WT vs *Pank4^-/-*; *P*=0.0036, WT+TGFβ vs *Pank4^-/-*+TGFβ, n=6. (**L**)

Concentrations of pyruvic acid (μg/ml) in LECs of negative control group, si-control

group and siPank4 group. Data were analyzed by one way ANOVA plus Bonferroni

post hoc test. Relative expression is the fold changes relative to the siPank4 group.

P<0.0001, siControl vs siPank4, n=6; *P*<0.0001, NC vs siPank4, n=6. (**M**) Schematic

illustration of current results.

1084

1085

Figure 4. **PANK4 deficiency is responsible for PKM2 Y105 phosphorylation**

(**A**) and (**B**) Protein levels and quantifications of N-cadherin, α-SMA and collagen-I

under PKM-IN-1 and DASA-58 treatments. Data were analyzed by one way ANOVA

plus Bonferroni post hoc test. Relative expression is the fold changes relative to the

Veh group. N-cadherin: *P*=0.0138, Veh vs PKM-IN-1 groups, n=6; *P*=0.0033, Veh vs

DASA-58 groups, n=6; collagen I: *P*=0.1066, Veh vs PKM-IN-1 groups, n=6;

P=0.0002, Veh vs DASA-58 groups, n=6; α-SMA(*P*=0.0019, Veh vs PKM-IN-1

groups, n=6; *P*=0.0007, Veh vs DASA-58 groups, n=6). GAPDH as inner control. (**C**)

Protein levels and quantifications of p-PKM2, PKM2, N-cadherin, α-SMA and

collagen-I under PKM-IN-1 inhibition in *Pank4*$^{-/-}$ mice. (**D**) TGF-β2 injection with

vehicle and PKM-IN-1 treatment in 2Month old *Pank4*$^{-/-}$ mice. (**E**) Represented blot

image of mice lens epithelium samples to show PKM2 monomer, dimer and tetramer.

(**F**) Relative expression of t-PKM2 vs GAPDH. Data were analyzed by

Mann-Whitney U test. All data shown are median±interquartile range. *P*=0.0067, WT

vs *Pank4*$^{-/-}$ in 60kD group, *P*=0.1 vs *Pank4*$^{-/-}$ in 120kD group, *P*=0.8498, WT vs

Pank4$^{-/-}$ in 240kD group, n=3. (**G**) and (**H**) Levels of proteins and relative

quantifications of p-PKM2(Y105), p-PKM2(S37) and t-PKM2 in WT and *Pank4*$^{-/-}$

mice lens epithelium. Data were analyzed by student's *t* test. *P*=0.6683, WT vs

Pank4⁻ᐟ⁻, n=6, in Y105 group; *P*=0.0022, WT vs *Pank4⁻ᐟ⁻*, n=6, in S37 group. (**I**)

Immunofluorescence staining of p-PKM2(Y105) in lens epithelium of WT and

Pank4⁻ᐟ⁻ mice. (**J**) Plot profile analysis described the level of p-PKM2(Y105) in WT

and *Pank4⁻ᐟ⁻* mice lens epithelium. (**K**) Plot profile analysis of represented single lens

epithelial cell immunofluorescence stained with p-PKM2(Y105). (**L**)

Immunofluorescence staining of p-PKM2(S37) in lens epithelium of WT and *Pank4⁻ᐟ⁻*

mice. (**M**) Plot profile analysis described the level of p-PKM2(S37) in WT and

Pank4⁻ᐟ⁻ mice lens epithelium. (**N**) Plot profile analysis of represented single lens

epithelial cell immunofluorescence stained with p-PKM2(S37). (**O**) Schematic

illustration of current results in figure 4.

1114

Figure 5. Negatively regulation of PKM2 by PANK4 in a HIF dependent way

attenuating OXPHOS of lens epithelium but promoting the glycolysis related

Warburg effect

(**A**) Co-IP was performed in WT and *Pank4⁻ᐟ⁻* mice lens epithelium. HIF-1α was

physically interacted with PKM2, but dramatically dissociated when PANK4 ablated.

(**B**) The Western blot results and the quantifications indicated the protein levels of

HIF-1α, HIF-1β and HIF-2α. Data were analyzed by student's *t* test. Relative

expression is the fold changes relative to the WT group. Hif-1α: *P*=0.0004, WT vs

Pank4⁻ᐟ⁻, n=6; Hif-1β: *P*=0.0022, WT vs *Pank4⁻ᐟ⁻*, n=6; Hif-2α:*P*=0.0075, WT vs

Pank4⁻ᐟ⁻, n=6. (**C**) concentrations of lactate (ng/μL) in blood and lens epithelium of

WT and *Pank4⁻ᐟ⁻* mice. Data were analyzed by Mann-Whitney U test. All data shown

are median±interquartile range. *P*=0.0001, WT vs *Pank4*[-/-] in lens group, n=4. (**D**)

GSEA analysis enriched in electron transport chain, oxidative phosphorylation and

respiratory electron transport, showed positive correlation with *Pank4*. GSEA analysis

enriched in electron transport chain, oxidative phosphorylation and respiratory

electron transport, showed negative correlation with *Pank4*. (**E**) and (**F**) Energy

metabolism was analyzed on day 2 during reprogramming with the Seahorse

instrument. Decrease in OCR after si*Pank4* was considered as the ATP production

ability of the cells. Spare respiratory capacity and maximum respiration were

decreased, while non mitochondrial oxygen consumption was increased after si*Pank4*

treated. Data were analyzed by one way ANOVA plus Bonferroni post hoc test.

Relative expression is the fold changes relative to the siControl group. Spare

respiratory capacity: *P*<0.0001, siCont vs siPank4, n=6; Non-mitochondrial oxygen

consumption: *P*<0.0001, siCont vs siPank4, n=6; Maxium respiration: *P*<0.0001,

siCont vs siPank4, n=6. (**G**) and (**H**) Increase in extracellular acidification rate

(ECAR) after si*Pank4* treatment was considered as glycolysis ability of the cells.

Energy metabolism was analyzed on day 2 during Pank4 knockdown with the

Seahorse instrument. Glycolytic rate, glycolysis capacity and glycolysis reserve were

all increased, respectively. Data were analyzed by one way ANOVA plus Bonferroni

post hoc test. Relative expression is the fold changes relative to the siControl group.

Glycolytic rate, *P*=0.0065, siCont vs siPank4, n=6; Glycolysis capacity, *P*<0.0001,

siCont vs siPank4, n=6; Glycolysis reserve, *P*=0.0002, siCont vs siPank4, n=6. (**I**) and

(**J**) Concentrations of pyruvic acid and ATP in siControl and si*Pank4* groups in lens

epithelial cells. Data were analyzed by Mann-Whitney U test. All data shown are

median±interquartile range. Concentration of pyruvic acid: $P=0.0286$, siCont vs

siPank4, n=4; Concentration of ATP: $P<0.0211$, siCont vs siPank4, n=4. (**K**) and (**L**)

Glycolysis, glycolysis capacity and glycolysis reserve were all increased, respectively,

in siPank4 group rather than PKM-IN-1, DASA-58 and PKM-IN-1+siPank4. Data

were analyzed by one way ANOVA plus Bonferroni post hoc test. Relative expression

is the fold changes relative to the siPank4 group. Glycolysis: $P<0.0001$, PKM-IN-1 vs

siPank4, n=6; $P<0.0001$, DASA-58 vs siPank4, n=6; $P<0.0001$, PKM-IN-1+siPank4

vs siPank4, n=6; $P<0.0001$, DASA58+siPank4 vs siPank4, n=6. Glycolysis capacity:

$P<0.0001$, PKM-IN-1 vs siPank4, n=6; $P<0.0001$, DASA-58 vs siPank4, n=6;

$P<0.0001$, PKM-IN-1+siPank4 vs siPank4, n=6; $P<0.0001$, DASA58+siPank4 vs

siPank4, n=6; Glycolysis reserve: $P<0.0001$, PKM-IN-1 vs siPank4, n=6; $P<0.0001$,

DASA-58 vs siPank4, n=6; $P<0.0001$, PKM-IN-1+siPank4 vs siPank4, n=6;

$P<0.0001$, DASA58+siPank4 vs siPank4, n=6. (**M**), (**N**) and (**O**) Decrease in OCR

after si*Pank4* was considered as the ATP production ability of the cells. Spare

respiratory capacity elevated in si*Pank4* compared with PKM-IN-1, DASA-58 and

PKM-IN-1+si*Pank4* groups. Maximum respiration increased after si*Pank4* treatment

compared with PKM-IN-1 and PKM-IN-1+si*Pank4* groups, while Proton leak

increased after si*Pank4* treatment compared with PKM-IN-1 and PKM-IN-1+si*Pank4*

groups. ATP production was elevated in si*Pank4* treatment compared with PKM-IN-1

group. Spare respiratory capacity: $P=0.0002$, PKM-IN-1 vs siPank4, n=6; $P<0.0001$,

DASA58 vs siPank4, n=6; $P=0.0006$, PKM-IN-1+siPank4 vs siPank4, n=6; $P=0.1972$,

siPank4 vs DASA-58+siPank4, n=6. Basal respiration: $P<0.0001$, PKM-IN-1 vs

siPank4, n=6; $P<0.0001$, PKM-IN-1+siPank4 vs siPank4, n=6; $P=0.9596$, siPank4 vs

DASA-58+siPank4, n=6. Maximum respiration: $P<0.0001$, PKM-IN-1 vs siPank4,

n=6; $P<0.0001$, PKM-IN-1+siPank4 vs siPank4, n=6. Proton leak, $P=0.0022$,

PKM-IN-1 vs siPank4, n=6; $P=0.0055$, PKM-IN-1+siPank4 vs siPank4, n=6. ATP

production: $P<0.0001$, PKM-IN-1 vs siPank4, n=6; $P<0.0001$, PKM-IN-1+siPank4

vs siPank4, n=6. (**P**) Concentration of ATP (%) elevated both in Vehicle and

PKM-IN-1 groups rather than that in DASA-58 group. Data were analyzed by one

Kruskal-Wallis test. Relative expression is the fold changes relative to the Veh group.

$P=0.0341$, Veh vs DASA-58, n=3. (**Q**) Scheme summarizing of the results.

1180

Supplementary Figure Legends

Figure 1. Screen assay of 3 types of designed siPank4 by Western blot.

Figure 2. (**A**) Concentrations of acetyl-CoA in WT and $Pank4^{-/-}$ mice lens epithelium,

brain cortex and retina. Data were analyzed by Mann-Whitney U test. All data shown

are median±interquartile range. Lens: $P=0.7$, WT vs $Pank4^{-/-}$, n=3; Retina: $P=0.1$, WT

vs $Pank4^{-/-}$, n=3; Brain: $P=0.4$, WT vs $Pank4^{-/-}$, n=3. (**B**) The protein level and

quantification of acetyl CoA carboxylase (ACC) in WT and $Pank4^{-/-}$ lens. Data were

analyzed by Mann-Whitney U test. All data shown are median±interquartile range.

$P=0.0286$, WT vs $Pank4^{-/-}$, n=4.

Figure 3. Relative mRNA of Pank4 in healthy and diabetes patients. Data were

analyzed by Mann-Whitney U test. All data shown are median±interquartile range.

P=0.8413, WT vs *Pank4$^{-/-}$*, n=5.

Figure 4. Protein levels of total-Stat3 and phosphor-stat3 in WT and *Pank4$^{-/-}$* mice

lens epithelium. Data were analyzed by Mann-Whitney U test. All data shown are

median±interquartile range. *P*>0.9999, WT vs *Pank4$^{-/-}$*, n=3.

Figure 5. Protein levels of PKM2 and HIF-1α in WT and *Pank4$^{-/-}$* mice lens with and

without 70% oxygen treatment for continuous 7 days. Data were analyzed by

Mann-Whitney U test. All data shown are median±interquartile range. *P*=0.0286, WT

vs. Pank4$^{-/-}$, n=4; *P*=0.0286, WT(O2) *vs. Pank4$^{-/-}$*(O2), n=4.

1200

1201

1202   **Conflict of Interest Statement**

1203   All the authors declared that there is no conflict of interest.

1204