A novel *in silico* method employs chemical and protein similarity algorithms to accurately identify chemical transformations in the human gut microbiome

Annamarie Bustion[1,2], Ayushi Agrawal[2], Peter J. Turnbaugh[3,4], Katherine S. Pollard[2,4-7*]

[1]Pharmaceutical Sciences and Pharmacogenomics Graduate Program, University of California, San Francisco, CA, USA.
[2]Institute of Data Science and Biotechnology, Gladstone Institutes, San Francisco, CA 94158, USA
[3]Department of Microbiology & Immunology, University of California San Francisco, San Francisco, CA 94143, USA
[4]Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
[5]Department of Epidemiology & Biostatistics, University of California, San Francisco, San Francisco, CA, 94158, USA
[6]Institute for Human Genetics, University of California, San Francisco, CA 94143, USA
[7]Bakar Computational Health Sciences Institute, University of California, San Francisco, CA 94143, USA

*Correspondence to:
Katherine Pollard
1650 Owens Street
San Francisco, CA 94158
kpollard@gladstone.ucsf.edu
415-734-2711

Abstract

Bacteria within the gut microbiota possess the ability to metabolize a wide array of human drugs, foods, and toxins, but the responsible enzymes for these chemical events remain largely uncharacterized due to the time consuming nature of current experimental approaches. Attempts have been made in the past to computationally predict which bacterial species and enzymes are responsible for chemical transformations in the gut environment, but with low accuracy due to minimal chemical representation and sequence similarity search schemes. Here, we present an *in silico* approach that employs chemical and protein Similarity algorithms that Identify MicrobioMe Enzymatic Reactions (SIMMER). We show that SIMMER predicts the chemistry and responsible species and enzymes for a queried reaction with high accuracy, unlike previous methods. We demonstrate SIMMER use cases in the context of drug metabolism by predicting previously uncharacterized enzymes for 88 drug transformations known to occur in the human gut. Bacterial species containing these enzymes are enriched within human donor stool samples that metabolize the query compound. After demonstrating its utility and accuracy, we chose to make SIMMER available as both a command-line and web tool, with flexible input and output options for determining chemical transformations within the human gut. We present SIMMER as a computational addition to the microbiome researcher's toolbox, enabling them to make informed

hypotheses before embarking on the lengthy laboratory experiments required to
characterize novel bacterial enzymes that can alter human ingested compounds.
Introduction
Humans consume a large array of foods, therapeutics, and other xenobiotics that are
processed, in part, by enzymes of bacteria residing within the gut. While some bacterial
enzymes are orthologous to the human metabolism repertoire, many bacteria possess
metabolic capabilities distinct from our own (Zimmermann et al., 2019a). It is important
to ascertain the extent of microbial capacity for chemical transformation because it has
implications for the bioavailability, toxicity, and efficacy of the compounds humans
ingest (Koppel et al., 2017; Spanogiannopoulos et al., 2016). Additionally, because the
human gut microbiome differs from individual to individual, knowledge of the prevalence
and abundance of bacterial enzymes must be determined before beneficial clinical and
dietary decisions can be made (Javdan et al., 2020).
59
While experimental methods can be employed to expand what we know of bacterial
enzymatic capabilities in the gut, the scientific community lacks genetic tools for nearly
all bacterial species of the human microbiota, and heterologous expression in model
organisms can fail for a plethora of reasons (Bisanz et al., 2020; Patel et al., 2022).
When experimental methods are tractable, the time required is often so extended that
knowledge is gained in a low-throughput manner. For these reasons, attention should
turn to the employment of *in silico* computational methods that can guide
experimentalists in their hypothesis-building process by aiding in the prioritization of
substrates, species, and genes worth studying.
69
Recent attempts have been made to create computational descriptions of chemical
transformation by human gut bacteria, but none can be expanded to predict the
metabolic capabilities of bacterial proteins with unknown function or to explore the
capacity of microbial enzymes to degrade novel substrates. Two previously published
methods aimed to predict known drug metabolism events within the human gut
microbiome, but the accuracy of their predictions were limited due to the fact that both
tools only consider substrates, rather than a full chemical description of substrate(s),
cofactor(s), and product(s) formed in a reaction. Both tools were also limited by the use
of small databases that do not fully capture the diversity of the human gut microbiome
(Guthrie et al., 2019; Sharma et al., 2017).
80
To address these gaps in accurate predictive software for bacterial chemical
transformations, we present SIMMER, a tool that combines chemoinformatics and
metagenomics approaches to accurately predict bacterial enzymes capable of
metabolism events. Given an input reaction, SIMMER predicts an Enzyme Commision
(EC) code that describes the chemical nature of the query. SIMMER additionally
predicts specific bacterial enzyme sequences, functions, prevalence, and abundance for
the reaction. Our key innovations are the use of full chemical representations that
include cofactors employed and products produced in a reaction, the use of statistically
informed sequence searches of a comprehensive human gut microbiome gene catalog,
and the development of a novel EC predictor based on reaction rather than gene
sequence. As a use-case, we present evidence that SIMMER provides high accuracy

predictions of bacterial enzymes responsible for known drug metabolism events, and we
identify the likely bacterial enzyme for 88 drugs known to be metabolized by the gut
microbiome for which the enzyme was previously unknown.
95
Results
SIMMER pipeline to predict xenobiotic metabolizing enzymes
There are many desiderata for a bacterial drug metabolism predictor (Table 1). Such a
tool must be able to, based on quantified chemical similarity, predict EC annotations,
specific enzyme sequences, and the prevalence and abundance across human
samples of those predicted sequences. We developed SIMMER, a tool that leverages
chemical and protein similarity to identify enzymes in the human microbiome that could
perform a queried chemical reaction (Figure 1). Given input substrate(s), metabolite(s),
and any known cofactors, SIMMER predicts bacterial enzymes capable of performing
the reaction and quantifies their prevalence and abundance in the human gut. SIMMER
accomplishes this by chemically fingerprinting an input reaction, and then comparing it
to all reactions in MetaCyc. Enzyme annotations from the most similar MetaCyc
reactions are then used as queries for a protein similarity search to find homologs in the
genomes of gut bacteria. To decrease the runtime of a SIMMER query, we
precomputed chemical descriptions and protein similarity searches for all reactions in
MetaCyc.
112

113
Figure 1. SIMMER architecture. (A) Precomputation on 8,914 gene annotated bacterial
reactions downloaded from MetaCyc. Chemical fingerprints representing each MetaCyc reaction
were created from SMILES descriptors. A latent chemical space was then created via a pairwise

reaction similarity matrix based on Tanimoto coefficients. For each reaction, relevant gene
sequences were retrieved from UniProt and Entrez database linkouts and used to create
multiple sequence alignments and subsequent pHMMs using ClustalO and HMMER3
respectively. pHMMs were used to retrieve homologs in a catalog of human gut microbiome
genes. **(B)** Running a SIMMER query. After receiving a reaction query (input compound, co-
factors, products), SIMMER fingerprints the reaction and compares it to the precomputed
chemical space from 1A. MetaCyc reactions are sorted by similarity to the query. Significantly
enriched EC identities are reported, and from the most similar reaction, human gut microbiome
enzymes are reported along with their abundance and prevalence in gut microbiomes.
126
SIMMER's underlying data
was drawn from the MetaCyc
reaction database because its
small-molecule reaction
descriptions each possess at
least one experimentally
validated enzyme annotation
and often include a
description of the reaction
type via EC code. To build a
precomputed chemical
search space for SIMMER
queries (Figure 1A), we
created two-dimensional
fingerprint representations for
8,914 enzyme driven
reactions in MetaCyc (Caspi
et al., 2008; Schneider et al.,
2015). Using these
fingerprints, we estimated the
similarity between all pairs of
reactions based on Tanimoto
coefficients. To build the
enzyme backbone of

Table 1. Desired elements of a microbiome chemical transformation predictor.

		DrugBug (Sharma et al., 2017)	MicrobeFDT (Guthrie et al., 2019)	**SIMMER**
Input types	Accepts novel SMILES			✓
	User options for different chemical fingerprints	✓		✓
Output types	Reaction similarity measure			✓
	EC predictions	✓	✓	✓
	Enzyme and species predictions	✓		✓
	Function predictions			✓
	Prevalence/abundance		✓	✓
Usability	Web server	✓		✓
	Command line tool			✓
	Docker container		✓	
Accuracy	Prediction of previously characterized enzymes	3%	NA	**87%**
	Prediction of previously characterized EC numbers	39%	14%	**93%**

SIMMER, we compiled the Uniprot and/or Entrez gene identifiers linked to each
MetaCyc transformation into a profile hidden Markov model (pHMM) that represents the
diversity of the enzyme family for a respective reaction. The resulting pHMMs were then
used to query the Unified Human Gastrointestinal Genome (UHGG) collection of
286,997 isolate genomes and metagenome assembled genomes from the human gut
environment (Almeida et al., 2020). Additionally, prevalence and abundance of all
pHMM search hits were assessed in stool metagenomes from the PREDICT human
cohort using MIDAS2, an implementation of Metagenomic Intra-Species Diversity
Analysis System (MIDAS) designed for use with the UHGG catalog (Almeida et al.,
2020; Nayfach et al., 2016; Zhao et al., 2022).
161
After creating SIMMER's precomputed chemical and pHMM search space, we next
made SIMMER queryable (Figure 1B). When queried with a chemical transformation,

SIMMER computes the chemical similarity of the input to all precomputed MetaCyc
reactions, and sorts all MetaCyc reaction fingerprints according to their ascending
Euclidean distance from the query. From this sorted list, SIMMER outputs enzymes (i.e.
the precomputed pHMM search hits for the closest reactions) responsible for the query
reaction and an EC code (i.e. reaction type) prediction. We implemented and validated
a novel method to predict EC codes by extending a common approach to gene set
enrichment analysis (GSEA) (Supplementary Figure 1A-C) (Subramanian et al., 2005).
With this enrichment method, SIMMER predicted reaction types for queries with high
recall, precision, and accuracy for EC classes, sub-classes, and sub-sub-classes
(Figure 2B, Supplementary Table 1, Supplementary Figure 1D).
174
We hypothesized that SIMMER accurately predicted chemical transformations due to its
use of a full reaction that includes reactants, cofactors, and products, rather than just
substrates. We assessed this hypothesis by demonstrating that SIMMER groups similar
reactions together in chemical space. MetaCyc reactions possess EC annotations that
describe the chemical class of a reaction (e.g. oxidoreduction, hydrolysis, intramolecular
rearrangement, etc). We queried SIMMER with all EC annotated MetaCyc reactions and
demonstrated that queries group significantly closer to other reactions within their EC
class than they do to reactions of a different class (Figure 2A). We determined that
SIMMER's ability to group similar reactions in chemical space is resilient to different
fingerprinting methods (Supplementary Figure 2), but not to loss of products created
and cofactors consumed in a reaction (Supplementary Figure 3). Thus we showed that
similar reactions only cluster together in chemical space when a full reaction description
(i.e. SIMMER's representation method) is employed.
188

189
Figure 2. SIMMER's chemical representations capture information relevant to enzymatic
reactions. (A) SIMMER clusters similar reactions together in chemical space. To analyze
SIMMER's ability to group chemically similar reactions, we examined reaction similarity within
versus between EC classes using the procomputed MetaCyc reaction dataset. A silhouette-like
euclidean distance score was created by determining for each reaction its euclidean distance to
all reactions within its EC class versus outside its EC class. For all EC classes, scores were
smaller within versus between EC classes using SIMMER's chemical representation, indicating
that SIMMER can detect reaction similarity within EC classes. From the pairs of distributions we

computed a Kolmogorov-Statistic to determine if the distributions significantly (p<0.05) differed.
(B) The F1-score, or harmonic mean of SIMMER's precision and recall, when predicting EC
numbers. The score is high for EC classes, and it generally decreases as an EC number's
resolution increases.
202
Because SIMMER was created with the assumption that chemically similar reactions
are mediated by sequence similar enzymes, we next ensured that similarity within
SIMMER's chemical space could be used to find shared, responsible enzymes. First, for
all MetaCyc enzymes associated with multiple reactions, one reaction was used as a
SIMMER query, and the second reaction searched for in the ordered reaction list output.
As a negative control, these reaction similarity results were then compared to all
possible pairwise combinations of reactions not conducted by the same enzyme.
SIMMER predicted high similarity between reactions conducted by a shared enzyme,
and low similarity for those reactions without a shared enzyme, (Figure 3A). We also
found a negative association between chemical reaction distance and sequence
similarity of MetaCyc enzyme annotations, indicating that reactions with similar
chemistry are conducted by sequence similar enzymes, though there is much variation
in this relationship (Figure 3B). This reflects the known association between sequence
similarity and similarity in chemical function, as well as reports that this relationship can
often be overestimated (Tian and Skolnick, 2003). Together, these analyses
demonstrated that sequence similar enzymes do indeed mediate chemically similar
reactions, strengthening the logic of combining chemical and protein similarity in a
microbiome enzyme prediction tool.
221

222
Figure 3. SIMMER's chemical representations can be used to find shared, responsible
enzymes. (A) When SIMMER was queried with a MetaCyc reaction, other reactions driven by
the same enzyme are returned as the most similar. As a contrast, reactions driven by a different
enzyme yield a more uniform rank distribution. Solid lines of the violin plots depict median
reaction similarity rank and dashed lines represent lower and upper quartile ranges. **(B)**
Similarly, there is a negative association between pairwise reaction euclidean distance and
pairwise protein identity, demonstrating that SIMMER can capture the known, albeit weak
relationship between sequence identity and similar reaction chemistry (Tian and Skolnick,
2003).

232
An expanded list of gut bacterial enzymes relevant to known cases of drug
metabolism
To assess SIMMER's prediction accuracy for previously characterized reactions and to
mount a comparison to existing methods, we used drug metabolism as a use case.
First, we curated 298 drug-metabolism events associated with the human gut
microbiome from the literature (Supplementary Table 2). For 31 of these reactions the
responsible bacterial enzyme, characterized metabolite(s), and associated EC
annotation are known (Supplementary Table 2). These 31 reactions are conducted by
18 enzymes. Due to orthology and proclivity for genetic transfer between even distantly
related bacteria, however, there are likely many as yet undiscovered homologs of these
drug-metabolizing enzymes that can catalyze identical drug metabolism events (Pollet
et al., 2017). To account for this, we created an expanded database (Supplementary
Table 3, Supplementary Figure 4) of the 18 characterized enzymes from pHMM and
phmmer searches of the UHGG database (Almeida et al., 2020), yielding 52,849 total
candidate homologs (a median of 1,087 candidates per enzyme). After filtering enzymes
by hmmer significance, alignment length, presence in data from the human jejunum
(Zmora et al., 2018) and RNA-sequencing studies (Integrative HMP (iHMP) Research
Network Consortium, 2019), and predicted affinity for the substrate in question using the
Similarity Ensemble Approach (Keiser et al., 2007), our database contained a median of
2 high-confidence homologous sequences per enzyme (range = 0 to 460 across the 18
enzyme families, Supplementary Table 3, Supplementary Figure 4). These 741
additional enzyme sequences for 31 reactions formed our positive control set of known
gut bacterial enzymes.
256
SIMMER captures known gut bacterial enzymes involved in drug metabolism
With our expanded database of drug-metabolizing enzymes from the human gut
microbiome in hand, we next verified that SIMMER can accurately predict reaction types
and responsible enzymes for the 31 known chemical transformations. Only 3 of these
31 reactions are themselves MetaCyc entries (5-ASA, dopamine, and levodopa
degradation); if EC codes and enzymes of reactions not described in MetaCyc were
also accurately predicted, it would show that SIMMER can discover non-identical yet
chemically similar reactions.
265
Of the 31 drug-metabolism events known to occur via human gut bacterial enzymes, EC
annotations exist for 28. SIMMER identified the correct EC class for 26 of these 28
reactions (93%) (Supplementary Table 4). For some queries SIMMER predicted more
than one significant EC code, but again, for 26 out of the 28 reactions, the top EC class
prediction was a match (Supplementary Table 4, Supplementary Figure 5). The two
failed EC predictions were for nicardipine reduction (inappropriately predicted as an
isomerase reaction) and for brivudine transformation (for which SIMMER made no
significant prediction).
274
In addition to accurate EC (i.e. reaction type) identification, SIMMER also accurately
predicted the specific enzymes from the human gut microbiome that conduct the 31
query reactions (Supplementary Table 4). This enzyme list was populated by the results

of the precomputed pHMM searches of human microbiome catalogs with annotated
gene sequences from MetaCyc reactions (Figure 1A). In 27 cases (87%), the
characterized (i.e. positive control) enzyme(s) for a reaction was found in the output
enzyme list for the top 20 of the ranked MetaCyc reactions (Supplementary Figure 4).
Since the positive controls span four EC classes (EC1 oxidoreductases, EC2
transferases, EC3 hydrolases, and EC4 lyases) this result demonstrates SIMMER's
ability to accurately predict microbiome based enzymes for a diversity of reaction types.
Also, despite inaccurate EC predictions for nicardipine reduction and brivudine
transformation, SIMMER was able to respectively predict AzoR and BT_4554 enzymes
as responsible for the reactions (Supplementary Table 4, Supplementary Figure 5).
288
SIMMER outperforms existing methods
To mount a comparison to the other *in silico* methods that, in part, aimed to describe
microbiome drug metabolism, we next queried the 31 positive control reactions using
MicrobeFDT and DrugBug (Table 1) both of which rely solely on substrate chemical
similarity rather than information from a whole reaction (Guthrie et al., 2019; Sharma et
al., 2017). For the 28 EC annotated positive control reactions, DrugBug had 39%
accuracy in predicting EC classes (in comparison to SIMMER's 93%), and predicted the
correct enzyme for a single reaction, SN38 glucuronide deconjugation, despite the
presence of chemically similar reactions metabolized by the same enzyme amongst the
positive controls (Supplementary Table 5). We additionally queried SIMMER with the
four positive controls (ginsenoside Rb1, quercetin-3-glucoside, cycasin, and sorivudine)
associated with characterized bacterial enzymes from the original DrugBug publication
(Supplementary Table 5). Both DrugBug and SIMMER were able to predict EC classes
for sorivudine, but only SIMMER was able to accurately predict the specific enzyme
(BT_4554) responsible for the drug's degradation. For ginsenoside Rb1 (3.2.1.192),
quercetin-3-glucoside (3.2.1.21), and cycasin (3.2.1.21), SIMMER was able to
accurately predict EC codes out to sub-sub-class (3.2.1.-), serial designation (3.2.1.21),
and sub-sub-class (3.2.1.-) respectively, which was a resolution improvement over
DrugBug (Supplementary Table 5).
308
We next queried the 28 EC annotated drug-metabolism positive controls against
MicrobeFDT (which uses a chemical graph to predict EC codes, but not enzymes) in
two ways: first by looking for direct enzyme metabolism events, and second, by looking
for enzyme metabolism of compounds that overlap chemically with the positive control
in question. When directly queried, MicrobeFDT produced metabolism predictions for
four of the 28 positive controls, three of which were correct. When queried with
chemically similar compounds according to its graph, MicrobeFDT produced
metabolism predictions for 13 of the 28 positive controls, and four were correct (14%
overall accuracy in comparison to SIMMER's 93%). It is important to note that
MicrobeFDT is reliant on a fixed database that cannot be modified by the user, meaning
that a compound cannot be queried if it is not already present in MicrobeFDT's graph.
We finished the comparison between SIMMER and MicrobeFDT by querying SIMMER
with the metabolism use-case described in the MicrobeFDT publication, altretamine
demethylation. In our hands, there was no Cypher query against the MicrobeFDT
database that resulted in a demethylase EC code (we determined possible demethylase

EC codes by running a query in the Swiss Institute for Bioinformatics Enzyme
Nomenclature Database)(Bairoch, 2000). We performed queries of direct EC annotation
for melamine and altretamine, as well as EC annotation queries for any compound with
either substructure or toxicity overlap with altretamine or melamine. The closest result to
a demethylase enzyme was a cypher query of toxicity overlap with altretamine that
yielded a nitric oxide synthase (EC 1.14.13.39) acting on L-arginine among its results
(Table 5). For its significant EC (reaction type) prediction, SIMMER identified
altretamine demethylation appropriately as an oxidoreductase reaction acting on a CH-
NH group of donors (EC 1.5.-), but not significantly as a demethylation event
(Supplementary Table 5).
334
335 This comparison illustrates SIMMER's enhanced accuracy over other methods for the
336 use case of characterized drug metabolism events by gut bacteria, and also illustrates
337 SIMMER's novel ability to predict chemical transformations not previously described in
338 literature or databases.
339
SIMMER predicts novel drug-metabolizing enzymes
After establishing SIMMER's accuracy in predicting drug-metabolizing enzymes in the
human gut environment, we predicted EC codes, functional annotations, and enzyme
sequences for novel microbiome drug metabolism reactions that do not yet possess a
responsible, characterized enzyme (Supplementary Table 6). From our literature
curation of 298 non-antibiotic therapeutics affected by the microbiome (Supplementary
Table 2), we were confident that 88 are directly metabolized by gut bacteria due to their
association with an identified bacterial metabolite in the literature. We formatted these
88 reactions in SMILES format and input them as queries to SIMMER.
349
Of the 88 reactions queried, SIMMER determined significant EC predictions for 75
reactions (86.2%), and 61 (70.1%) of these were out to the serial designation (i.e.
highest resolution) EC code (Supplementary Table 6). This list of 61 transformations
presents reactions for which we believe enzyme characterization is worth pursuing as
our predictions are significantly similar to enzymes already explored in the literature.
SIMMER's EC predictions resulted in expanded and diversified EC class membership
for drug-transformations known to occur in the microbiome (Figure 4C). Of interest, this
analysis resulted in a large expansion of putative hydrolysis, reduction, and
isomerization reactions in the human gut microbiome. The number of SIMMER
predictions varies widely by reaction, with median output of 372 genes, 286 genomes,
and 10 phyla predicted as responsible across the 88 reactions (Supplementary Table 6,
Figure 4A). Unsurprisingly, many of these reactions are predicted to occur due to
enzymes found in Firmicutes, Bacteroidetes, Actinobacteria, and Proteobacteria, but
there are also SIMMER enzyme predictions in phyla not previously associated with drug
metabolism (Figure 4B).

Figure 4. SIMMER predicts novel drug metabolizing enzymes. (A) Distributions depict the unique number of genes, strains, and phyla predicted to be responsible for 88 reported drug transformation reactions, as well as predicted gene functions. **(B)** A heatmap illustrating the number of phyla from the UHGG database capable of performing 88 drug metabolism events. Color intensity refers to the number of unique drug-metabolizing enzymes for a given phylum conducting a given reaction. **(C)** Enzyme Commission Class representation for bacterial transformations of therapeutics before and after the employment of SIMMER. Our predictions greatly expand the number of characterized reduction (EC1), hydrolysis (EC2), and

374   isomerization (EC5) events and modestly increase the number of transferase (EC2) and lyase
375   (EC4) events.
376
Eight of the 88 novel transformations were among those investigated in a high-
throughput study exploring the metabolism of 571 compounds in *ex vivo* stool samples
(Javdan et al., 2020). This publication demonstrated bacterial degradation of 57
therapeutics in a single pilot donor stool sample (with associated shotgun sequencing),
as well as in 20 human stool samples (with associated 16S rRNA gene sequencing).
While this study greatly expanded the number of drugs known to break down in the
presence of gut bacteria and identified eight metabolite structures, it only identified a
responsible enzyme in two of the 57 drug degradation cases due to the low-throughput
nature of enzyme characterization. To further assess SIMMER's ability to predict novel
enzymes, and to demonstrate the utility of using SIMMER in an experimental context,
we investigated the presence of our predictions in the Javdan, et al. study sequencing
results. Because shotgun metagenomics sequencing for the pilot donor was deposited,
we were able to confirm via tBLASTn searches that SIMMER enzyme predictions were
directly found in the pilot donor stool sample for all eight of the reactions with identified
metabolites (Supplementary Table 7). However, the sequencing data from the 20
human donor study was only 16S profiling, so we were unable to look directly for
SIMMER enzyme predictions. We were able to ensure that genomes found in
metabolizing stool samples contain SIMMER predictions. We found that donors who
could metabolize a given drug possessed a significant enrichment of genomes that
contain enzymes predicted by SIMMER. This was the case for four out of the five
reactions analyzed (Supplementary Table 7, Figure 5A).
398
Among the 88 novel transformations was the side-chain cleavage of dexamethasone to
17-oxodexamethasone. Dexamethasone was recently shown to be metabolized solely
by *Clostridium scindens* (ATCC 35704) out of a collection of 76 isolates representative
of the human gut microbiome (Zimmermann et al., 2019b). When dexamethasone
metabolism was assessed in 28 human stool samples, metabolite formation varied
substantially by individual, but could not be explained by *C. scindens* species
abundance. We sought to understand this lack of correlation. To do so, we assessed
the abundance of *C. scindens* SIMMER predictions across the 28 samples (i.e., the
amount of enzyme in genomes from different strains, not the amount of the species).
We found a significant association between metabolite formation and number of
SIMMER enzymes, and also a significant association between parent compound
consumption and number of SIMMER enzymes (Figure 5B).
411
It came to our attention while preparing this manuscript that recombinant steroid-17,20-
desmolase (DesAB) enzymes from *C. scindens* were shown to perform side-chain
cleavage on prednisone, but also to a lesser extent on dexamethasone. DesAB's
reduced activity for dexamethasone was assumed to be due to the compound's
potentially inhibitory 16α-methyl group (Ly et al., 2020). To ensure that SIMMER's
enzyme prediction for dexamethasone cleavage was not enriched in metabolizing stool
samples due to co-occurrence with already known DesAB, we next assessed the
abundance of *desAB* reads across the 28 samples, and found no significant correlation

between number of reads and either metabolite formation or dexamethasone
consumption slopes (Supplementary Figure 6).
422
423 These results indicate that species level information alone is not enough to predict
424 chemical transformations in a microbiome sample, but with SIMMER, knowledge of
425 responsible enzymes can recapitulate a sample's potential for therapeutic degradation.
426

427
Figure 5. SIMMER predicted enzymes explain inter-individual variations in drug
metabolism. (A) Donors from the Javdan, et al. 16S rRNA gene sequencing study (Javdan et
al., 2020) possessed an enrichment of genomes harboring SIMMER enzyme predictions when
metabolism of a given drug was observed. Violin plot curves were made using a seaborn
package that performs a kernel density estimation of the underlying datapoint distribution.
Chemical transformations were drawn using ChemDraw software. Asterisks denote p-values ≤
0.05. **(B)** There was a significant correlation between a human stool sample's ability to consume
dexamethasone (consumption slope, a.u.), to produce 17-oxodexamethasone (production
slope, a.u.), and the number of aligned SIMMER predicted sequences for side chain cleavage
of dexamethasone. Patient conversion slopes and metagenomics data were accessed from the
original study (Zimmermann et al., 2019).
439
440

SIMMER software
In addition to providing SIMMER (https://github.com/aebustion/SIMMER) as a command
line tool that quickly generates enzyme sequence predictions (fasta and tab-separated-
value files), EC predictions (tab-separated-value file), and MetaCyc reactions ranked by
similarity (tab-separated-value file) based on a user's input reaction, SIMMER is also
available as a user-friendly website (https://simmer.pollard.gladstone.org/). The user
can either input one query reaction at a time, or upload multiple reactions in tsv file
format (Supplementary Figure 7). All output types available with the SIMMER command
line tool are likewise retrievable via the SIMMER website.
450
Discussion
In this work, we created a tool that appropriately describes reaction chemistry and
harnesses all current information on gut bacterial sequences, both from isolates and
metagenome assembled genomes. This advances our ability to discover chemical
transformations in the human microbiome, because previous methods for *in silico*
metabolism prediction had several key limitations, including low accuracy. Here, we
demonstrated SIMMER's ability to recover known drug-metabolizing enzymes in the
human gut, to extend previous experimental findings for multiple drug metabolism
events by identifying candidate enzymes, and to add clarity to the genetic component of
dexamethasone metabolism by *C. scindens*.
461
To describe chemical reactions, we were initially influenced by recent research that
employed substrate and product chemistry to compare bacterial-drug metabolism
events to primary reactions in the MetaCyc database, but without the end-goal of EC
and enzyme identity prediction (Mallory et al., 2018). From a reaction description
standpoint, the published method was still limited in that it only included a description of
one substrate and one product per reaction, precluding it from utilizing cofactors and
from accurately describing anything other than intramolecular rearrangements (EC class
5, Supplementary Figure 3). For this reason, we employed a chemical representation
technique that can describe multiple inputs and outputs for a single reaction (Schneider
et al., 2015).
472
To connect these chemical descriptions to bacterial proteins in the human gut, we knew
it was important not to rely on EC codes (as previous methods have done) to find
relevant sequences. While EC codes are helpful for describing reaction types, from an
enzyme perspective they contain no information about substrate specificity for a
particular compound. We instead chose to create sequence searches of large genome
databases directly from enzymes known to conduct chemically similar reactions,
whether or not they have been fully annotated with an EC code. For example, enzyme
BT_4096 responsible for diltiazem deacetylation (Zimmermann et al., 2019b) is not yet
annotated by the EC, yet SIMMER was able to accurately predict the deacetylase
enzyme responsible for diltiazem metabolism because it does not require EC annotation
for its enzyme predictions. Indeed, instead of relying on EC codes for sequence
searches, we harnessed EC annotations in the MetaCyc database to create a novel EC
predictor. Using this tool, we accurately predicted diltiazem deacetylation as an EC3
hydrolysis reaction. While EC prediction methods based on sequence exist, to our

knowledge, this is the first instance of an EC prediction method based solely on the
chemical description of a reaction.
489
SIMMER achieved high accuracy when applied to known drug-metabolism events in the
gut microbiome. Correct EC designations and enzyme sequences were recovered for
31 drug metabolism events previously characterized in microbiome literature. These
reactions span multiple EC classes, and were described by multiple publications,
demonstrating the wide application and accuracy of SIMMER. While SIMMER provides
high accuracy (i.e. true positive) enzyme predictions for chemical transformations in the
human gut, the potential for false positives may be high, as its enzyme lists are not
filtered by biologically relevant metrics like substrate affinity or flux consistency in a
microbial community. To the former point, users may wish to employ tools like Similarity
Ensemble Approach to narrow in on hits most likely to interact with compounds of
interest (Keiser et al., 2007). To the latter, a user could choose to further analyze their
SIMMER output for flux-balance if the predicted SIMMER bacterial species are
described in current metabolic reconstruction models (Heinken et al., 2020;
Magnúsdóttir et al., 2017).
504
Due to its high accuracy predictions for previously described drug-metabolism events,
we also used SIMMER to predict novel drug metabolism chemistry in the human gut,
and expanded what we know of the bacterial enzymes at play in identified drug
transformations by gut bacteria. Recent high-throughput experimental research has
greatly increased our knowledge of the number of drugs altered by bacteria in the
human gut, but has led to a bottleneck in identifying the responsible bacterial enzymes.
While direct experimentation is a necessary component to elucidating the bacterial
players responsible, *in silico* methods like SIMMER are needed to help prioritize which
of the many bacterial species and enzymes to assess. Here we showed that SIMMER
both corroborates previous high throughput experimental data, and also adds increased
clarity to the findings. While a previous experimental study was able to elucidate the
importance of an isolate *C. scindens* in the metabolism of dexamethasone, the
abundance of *C. scindens* in human samples did not correlate with metabolism. When
assessed with SIMMER, however, a significant correlation between metabolite
production and amount of SIMMER predictions was observed. This finding
demonstrates that species identity alone is not enough to explain bacterial chemical
transformation, and that responsible genetic elements must be interrogated as well.
522
Two previous computational tools exist for describing non-antibiotic microbial drug
metabolism. MicrobeFDT groups thousands of compounds based on their similarity to
one another and annotates compound groups based on any known links to EC
numbers, and subsequently, microbes known to contain such EC codes (Guthrie et al.,
2019). This network approach was an important addition to the exploration of
microbiome metabolism, but its use is limited to a fixed database of chemicals and EC
annotations which prevents the user from exploring novel chemistry and also from
utilizing hypothetical protein data gathered from metagenomic sequencing studies.
Furthermore, MicrobeFDT's accuracy within its database of substrates is limited by its
exclusive description of substrates rather than full reactions. DrugBug, a tool that

533 employs Random Forests rather than a network approach, also exhibits limited power
534 and accuracy due to its sole reliance on substrate chemistry and relatively small
535 database of only 491 isolated bacterial genomes from the human gut (Sharma et al.,
536 2017). Of note, our comparison of SIMMER's performance to existing methods
537 necessitated downloading and analyzing our positive control list against the other tools,
538 as none of the previous publications provided any computational validation or accuracy
539 metrics.
540
541 One user pitfall of SIMMER in comparison to previous methods, is that a reaction's
542 product(s) and cofactor(s) identity is required to achieve the high accuracy enzyme
543 prediction described here. This is a limitation, as a growing amount of LC-MS/MS data
544 in microbiome research only reports whether or not a compound is depleted in the
545 microbiome and the mass/charge ratio of the product formed, not the product identity.
546 While it is technically possible for the user to submit a SIMMER query that only consists
547 of a substrate, or uses a compound identity as both substrate and product, we do not
548 recommend this due to the previously discussed lack of accuracy when only considering
549 substrates (Supplementary Figure 3, Supplementary Table 5). For users wishing to
550 utilize SIMMER with a compound of interest and its either unknown or uncharacterized
551 products, additional tools such as BioTransformer could be used in tandem to create
552 product template predictions before querying (Djoumbou-Feunang et al., 2019). Lastly,
553 if the user does not hold a certain level of knowledge in chemistry, appropriate cofactors
554 (such as water employed in a hydrolysis event) might be omitted from a query, leading
555 to lower accuracy predictions. If a user is unsure which cofactors may be at play in their
556 reaction of interest, reaction rules tools such as RetroRules could be employed (Duigou
557 et al., 2019).
558
559 Another limitation of SIMMER is that its underlying protein data is solely metagenomics
560 data from the human gastrointestinal tract, but some compounds, such as the vaginal
561 gel tenofovir, are known to be altered by bacteria in non-GI tract settings (Klatt et al.,
562 2017). That being said, for transformations in the human gut, SIMMER employs the
563 largest available database of relevant bacterial sequences, and the tool could easily be
564 expanded in the future to include other human body sites as well as non-host
565 associated environments. Further related to database constraints, while SIMMER is
566 novel in its ability to query reactions not previously described in chemistry databases, its
567 search space is still limited to reactions that broadly relate to those captured in
568 MetaCyc. As MetaCyc expands, or additional databases get employed, SIMMER will
569 likely be able to make increasingly fine-tuned predictions.
570
571 SIMMER enters microbiome biotransformation research at an important point: while
572 there are hundreds of microbiome altered compounds which are in need of enzyme
573 identification, there are also a sufficient number with characterized enzymes to enable
574 us to test the tool's accuracy. Its ability to predict these known enzymes accurately
575 builds confidence for its predictions of yet unknown enzymes. With this tool in hand,
576 microbiome researchers can make informed hypotheses before embarking on the
577 lengthy laboratory experiments required to characterize novel bacterial enzymes that
578 can alter human ingested compounds. Continued refinement of SIMMER and other

579 computational tools will accelerate microbiome research, providing data-driven
580 hypotheses for experimental testing and a first step towards understanding the full
581 scope of metabolism by the human microbiome.
582
Materials and methods
Preparation of SIMMER's underlying chemical data
13,387 gene annotated bacterial reactions were downloaded from MetaCyc (Caspi et
al., 2008). Each reaction from the database contained Simplified Molecular Input Line
Entry System strings (SMILES) of reactant(s) and product(s), EC code if available, and
UniProt or Entrez identifiers for sequences that catalyze the reaction. All MetaCyc
compounds were then protonated based on the pH environment of 7.4 in the human
small intestine, where most oral drug absorption occurs. Protonation states were
calculated using ChemAxon's cxcalc majorms software ("cxcalc calculator functions,"
n.d.).
593
RDKit's rdChemReactions module was employed to create chemical fingerprints
representing each MetaCyc reaction. Chemical reaction objects were constructed from
reaction SMILES arbitrary target specification (SMARTS) strings. Fingerprints for these
reactions were then created using the resulting difference of product(s) and reactant(s)
Atom-Pair fingerprints (Schneider et al., 2015). SIMMER users can also opt to use
Topological Torsion, Pattern, or RDKit fingerprints, but unless otherwise stated, all
analyses in this manuscript use Atom-Pair difference fingerprints. Of the 13,387
MetaCyc reactions, 8,914 were able to be fingerprinted using this method. Failed
fingerprints were due to ambiguous SMILES identifiers or presence of non-small
molecule compounds in a reaction, such as peptides.
604
605 After creating fingerprint vectors for all MetaCyc reactions, an 8,914 by 8,914 pairwise
606 similarity matrix of Tanimoto coefficients was created. These Tanimoto vectors make up
607 SIMMER's underlying chemical data.
608
Preparation of SIMMER's underlying protein data
For each of the 8,914 fingerprinted MetaCyc reactions, all relevant gene sequences
were retrieved from the MetaCyc reaction's UniProt and Entrez database linkouts. If at
least two genes, with a median pairwise sequence similarity greater than or equal to
27%, were linked to a given MetaCyc reaction, the sequences were used to create a
multiple sequence alignment and subsequent profile hidden Markov model (pHMM)
using Clustal Omega and HMMER3 (version 3.2.1) software respectively (Eddy, 2009;
Sievers and Higgins, 2014). This similarity cutoff was chosen based on previous protein
family literature (Mi et al., 2021). If fewer than two genes, or genes with less than 27%
global similarity, were associated with a given MetaCyc reaction, a pHMM of the
MetaCyc gene(s) PANTHER subfamily was retrieved via InterPro linkouts (Mi et al.,
2021). MetaCyc derived and PANTHER subfamily pHMMS were then queried against a
Unified Human Gastrointestinal Genome (UHGG) collection of 286,997 isolate genomes
and metagenome assembled genomes from the human gut environment using the
HMMER3 hmmsearch module (Almeida et al., 2020; Eddy, 2009). In the case of
MetaCyc reactions with too few sequences, too low a median pairwise sequence

identity, *and* a missing PANTHER database subfamily pHMM, single sequence protein
queries were conducted against the UHGG databse using HMMER3's phmmer module,
which internally created protein profiles for the single query sequences based on a
position-independent scoring system. Resulting enzyme hit lists were filtered to only
include high significance hits (e-value < 1E−5, and hit length >= half of the input pHMM
alignment or single sequence length). In sum, for each MetaCyc reaction, a profile
representing the diversity of the enzyme family for that chemical transformation was
used to find sequence similar hits in the human gut microbiome that can mediate
chemically similar reactions.
634
Each human gut microbiome hit was further described by the identity, prevalence, and
abundance of the bacterial strain in which it resides. To establish prevalence and
abundance of UHGG strains, metagenomic analysis was performed on the Predict
(Personalised Responses to Dietary Composition Trial) cohort due to its high number of
samples and favorable sequencing depth (Asnicar et al., 2021). Shotgun metagenomic
reads were analyzed with MIDAS2 an implementation of Metagenomic Intra-Species
Diversity Analysis Subcommands (MIDAS) (Nayfach et al., 2016; Zhao et al., 2022)
designed for the UHGG database. Presence of a SIMMER predicted species in a given
sample was established when reads mapped (HS-BLASTN) to 15 single-copy universal
genes for that species (Chen et al., 2015), with at least 75% alignment coverage. To
assess the gene content of a sample, shotgun metagenomic reads were aligned to a
MIDAS2 created pangenome of the SIMMER species' genes clustered at 99%
nucleotide identity. Copy number of a SIMMER gene prediction was established by
dividing aligned prediction reads by the full length of the prediction. This number was
then normalized by the read coverage of 15 single-copy universal genes in the same
sample to estimate copy number per cell. Presence of a SIMMER enzyme was
established if at least 0.35 gene copies per cell were present in a sample.
652
Phylogenetic trees were also constructed for each hmmsearch and phmmer result. For
each set of MetaCyc reaction human gut microbiome enzyme hits, CD-HIT was used to
cluster results at 95% identity (Fu et al., 2012). Then MUSCLE was used to create a
multiple sequence alignment for input to FastTree (Edgar, 2004; Price et al., 2009).
Compact tree visualizations were made in R using ggtree and ggtreeExtra (Xu et al.,
2021; Yu et al., 2017). All tree tips were colored by phylum, and surrounded by circle
annotators describing a given hit's Prokka predicted function, genome type (i.e. from an
isolate or metagenome assembled genome), and prevalence/abundance in the Predict
cohort (Seemann, 2014).
662
Query functionality of SIMMER
The query functionality of SIMMER was designed similarly to the precomputed
chemistry data. After receiving an input chemical transformation (or tsv describing
multiple input reactions) in the form of SMILES, SIMMER fingerprints the reaction(s)
and compares it to the precomputed chemical space by computing the Tanimoto
coefficients between the input(s) and all precomputed reactions. The 8,914
precomputed MetaCyc reaction Tanimoto vectors are then sorted by ascending
euclidean distance to the query Tanimoto vector. SIMMER by default ranks reactions'

euclidean distances based directly on the Tanimoto vectors, but if a user's inputs
require a decrease in computational burden, PCA can be employed after similarity
matrix creation and before euclidean distance rankings. The number of PCs to be used
depends on the fingerprint style employed, and was determined by the Kaiser criterion
(Supplementary Figure 8). Unless otherwise stated, all analyses in this manuscript
employed the full Tanimoto similarity matrix with no PCA reduction. Human gut
microbiome enzymes that may conduct the input reaction are reported from the
precomputed UHGG hmmsearch or phmmer results of the closest euclidean distance
MetaCyc reaction. Significantly enriched EC identities (i.e. reaction types) are also
reported.
681
Reaction type predictions
SIMMER predicts an EC code (i.e. reaction type) for a query reaction if there is an
enrichment of a particular EC at the top of the reaction list. Enrichment was determined
in a manner similar to gene set enrichment analysis (GSEA).(Subramanian et al., 2005)
For each EC code associated with MetaCyc reactions, an enrichment score (ES) was
calculated by walking down the ranked list of reactions. Starting with a score of zero,
each time the given EC is encountered the score increases by one, and each time a
different EC is encountered the score decreases by one. At the end of this process,
each EC receives an ES that is the score's maximum distance from zero after walking
through the list (Supplementary Figure 1A). Because the MetaCyc database of
reactions is unbalanced in its EC code representation, ES scores for a given EC type
are divided by the number of times the EC in question occurs in the database. This
yields a normalized ES (NES) for SIMMER reporting. Significance is established by
comparing the true NES to the NES achieved from 1000 permutations of a shuffled
reaction list (Supplementary Figure 1B-C). When multiple EC codes are predicted as
significant, they are ranked in ascending order of where in the list of 8,914 reactions the
NES occurs. This method was verified by subsampling the database of MetaCyc
reactions to equal numbers (N=96) of reactions for each EC class, the broadest
resolution level of an EC code. Each of these subsampled reactions was then queried
with SIMMER against the entire MetaCyc reaction database to create sorted reaction
lists for each query. SIMMER predicted an EC code(s) for each reaction based on the
most highly enriched EC. SIMMER's recall, precision, and accuracy are high for EC
class, sub-class, and sub-sub-class level resolution (Figure 2B, Supplementary Table
1). For the serial designation of an EC code (the most granular description of an EC
code), however, SIMMER's performance diminished, potentially because enrichment
calculations suffer from increased uniqueness in the ranked list and therefore reduced
power to determine a match (Supplementary Table 1). This indeed appears to be the
case; when the database is subsampled to ensure at least three of each unique serial
designation, F1-scores (the harmonic mean of precision and recall) and accuracy
remain high despite the increased EC resolution (Supplementary Table 1,
Supplementary Figure 1D).
713
Euclidean distance silhouette scores
To analyze SIMMER's resilience to different reaction chemistry representations, we
created a silhouette-like euclidean distance score. For the precomputed MetaCyc

chemical dataset of 8,914 reactions (i.e. the Tanimoto pairwise similarity matrix), we
split all reactions into their top-level EC codes (i.e. EC class) and determined for each
reaction its euclidean distance to all reactions within its EC class versus outside its EC
class. From the two distributions (within EC and without EC distances) created, we
computed a Kolmogorov-Statistic to determine if the distributions significantly ($p<0.05$)
differed. We repeated this process for finer resolution EC classifications (sub-class, sub-
sub-class, and serial designation). Euclidean distance silhouette scores were used to
compare different chemical representations, such as fingerprint style, inclusion of
products, and inclusion of cofactors.
726
Creating a compendium of drug-metabolism use cases from the human gut
To analyze SIMMER under the use-case of drug metabolism, we created a
compendium of drug degradations that occur in the human gut microbiome. The
compendium of reactions is based on a literature curation of hundreds of papers, and is
organized by reactions producing known/unknown metabolites and driven by
known/unknown bacterial enzymes. The drug-metabolism positive controls used to
assess SIMMER's accuracy were drawn from the list of reactions possessing a
structurally elucidated metabolite and driven by a characterized bacterial enzyme.
735
We further expanded the positive control list to include sequence-similar enzymes that
likely perform the same function. For this expansion, we performed pHMM searches
(when a positive control reaction had been characterized with multiple sequence-similar
enzymes) and phmmer searches (when a positive control reaction had been
characterized with only one sequence) of the UHGG database using HMMER3 software
(Almeida et al., 2020; Eddy, 2009). High significance (e-value $< 1E-5$) hits were kept
when the resulting alignment was at least 50% of the input pHMM or sequence length.
This list of significant hits was filtered by presence in human ileum or jejunum (the site
of human drug absorption) via DIAMOND searches against metagenomic reads from a
published study that employed jejunum and ileum endoscopy (Buchfink et al., 2021;
Zmora et al., 2018). The hits were also filtered for presence in RNAsequencing data via
DIAMOND searches of rnaSPAdes assembled reads from HMP2 metatranscriptomics
control patient samples (Bushmanova et al., 2019; Integrative HMP (iHMP) Research
Network Consortium, 2019). The hits were lastly filtered by predicted affinity for their
substrates using the Similarity Ensemble Approach (Keiser et al., 2007).
751
Corroboration of previous high-throughput experimental findings
For the first experimental validation of SIMMER, we analyzed results from sequencing
studies (NCBI BioProject: PRJNA593062) described in a previously published high-
throughput investigation of bacterial drug metabolism in human stool samples (Javdan
et al., 2020). The first sequencing set in this publication was a deep metagenomic
sequencing of one pilot individual's *ex vivo* stool originally evaluated for its ability to
degrade hundreds of therapeutics. We used MetaSPAdes with default settings to
assemble the metagenomics reads into scaffolds (Nurk et al., 2017). We then queried
SIMMER with eight reactions that were structurally elucidated (via nuclear magnetic
resonance) by the previous publication, and ensured via TBLASTN searches that
SIMMER predicted hits were found in the assembled metagenomic reads. The second

sequencing set was a 16S rRNA sequencing experiment of twenty human donor stool
samples originally evaluated for their inter-individual variation in bacterial drug
degradation. We queried SIMMER with five of these reactions possessing structurally
elucidated metabolites, and evaluated enrichment of SIMMER predicted bacterial
species in metabolizing versus non-metabolizing donors. Species matches between
SIMMER predictions and the 16S study were made using the SequenceMatcher class
from the difflib python module set to an 80% ratio cutoff. Enrichment of SIMMER
predicted bacterial genomes was then assessed by performing a generalized linear
model with poisson distribution for number of SIMMER genomes in non-metabolizers
versus metabolizers for a given reaction.
773
For experimental corroboration of dexamethasone metabolism, we accessed shotgun
sequencing data (PRJEB31790) from a cohort of 28 human stool samples shown to
metabolize dexamethasone to varying degrees (Zimmermann et al., 2019b). Shotgun
reads were assembled using MetaSpades with default settings. Presence of SIMMER
enzyme predictions was established via search with DIAMOND and normalized by
sample read depth. Significance was established with a Pearson's correlation coefficient
and P value calculated using a student's t-distribution.
781
Web tool creation
We used the python web framework Flask (https://flask.palletsprojects.com/en/2.1.x/) to
make SIMMER available as a user-friendly website. The website accepts either a single
query reaction or multiple query reactions via a file upload and provides the same
outputs as the SIMMER command-line tool. The website also allows the user to
download the outputs of interest. Keeping in mind the privacy and security of the data
that a user might upload to the website, the website is designed to delete all uploaded
data within 24 hours from the server. This will ensure security of the uploaded data.
790
Acknowledgements

We thank Russ Altman and Emily Mallory for their work that first inspired the
comparison of MetaCyc reactions to known gut microbiome chemistry (Mallory et al.,
2018), and Russ Altman for further insightful conversation. We thank Peter Karp and
Peter Midford of MetaCyc for providing us with an expanded set of links between
MetaCyc reactions and Uniprot or Entrez sequences. We thank Ben Guthrie and
Renuka Nayak of the Turnbaugh lab for collaboration and guidance in the ongoing
experimental validation of SIMMER enzyme predictions. We thank Daniela Arce and
Xiaofan Jin of the Pollard lab for guidance on assessing enzyme presence in the
jejunum and ileum of the human intestinal tract. We thank Chunyu Zhao and Jason Shi
of the Pollard lab for guidance on the appropriate use of MIDAS2 and the UHGG
database. We thank Patrick Bradley for guidance on the use of HMMER3 software in a
microbiome drug-metabolism context. We thank members of the Pollard and Turnbaugh
laboratories for their suggestions during the research process and writing of this
manuscript. Annamarie Bustion was supported by a trainee pilot award from the UCSF
Benioff Center for Microbiome Medicine, a predoctoral fellowship in informatics from the
PhRMA foundation, an Achievement Rewards for College Scientists (ARCS)
Scholarship from the ARCS foundation, and Training Grants from the NIGMS of NIH

(5T32GM007175-42 and 2T32GM007175-41). This work was also supported by
Gladstone Institutes. P.J.T. was supported by the National Institutes of Health
(R01HL122593). P.J.T. and K.P. are Chan Zuckerberg Biohub Investigators.
812

813 **Competing interests**
814 KSP is a consultant for Phylagen Inc. The other authors declare no competing financial
815 interests.
816

References

1. Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, Pollard KS,
Sakharova E, Parks DH, Hugenholtz P, Segata N, Kyrpides NC, Finn RD. 2020. A
unified catalog of 204,938 reference genomes from the human gut microbiome. *Nat
Biotechnol*. doi:10.1038/s41587-020-0603-3
2. Asnicar F, Berry SE, Valdes AM, Nguyen LH, Piccinno G, Drew DA, Leeming E,
Gibson R, Le Roy C, Khatib HA, Francis L, Mazidi M, Mompeo O, Valles-Colomer
M, Tett A, Beghini F, Dubois L, Bazzani D, Thomas AM, Mirzayi C, Khleborodova A,
Oh S, Hine R, Bonnett C, Capdevila J, Danzanvilliers S, Giordano F, Geistlinger L,
Waldron L, Davies R, Hadjigeorgiou G, Wolf J, Ordovás JM, Gardner C, Franks
PW, Chan AT, Huttenhower C, Spector TD, Segata N. 2021. Microbiome
connections with host metabolism and habitual diet from 1,098 deeply phenotyped
individuals. *Nat Med*. doi:10.1038/s41591-020-01183-8
3. Bairoch A. 2000. The ENZYME database in 2000. *Nucleic Acids Res* **28**:304–305.
doi:10.1093/nar/28.1.304
4. Bisanz JE, Soto-Perez P, Noecker C, Aksenov AA, Lam KN, Kenney GE, Bess EN,
Haiser HJ, Kyaw TS, Yu FB, Rekdal VM, Ha CWY, Devkota S, Balskus EP,
Dorrestein PC, Allen-Vercoe E, Turnbaugh PJ. 2020. A Genomic Toolkit for the
Mechanistic Dissection of Intractable Human Gut Bacteria. *Cell Host Microbe*
27:1001–1013.e9. doi:10.1016/j.chom.2020.04.006
5. Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life
scale using DIAMOND. *Nat Methods* **18**:366–368. doi:10.1038/s41592-021-01101-x
6. Bushmanova E, Antipov D, Lapidus A, Prjibelski AD. 2019. rnaSPAdes: a de novo
transcriptome assembler and its application to RNA-Seq data. *Gigascience* **8**.
doi:10.1093/gigascience/giz100
7. Caspi R, Foerster H, Fulcher CA, Kaipa P, Krummenacker M, Latendresse M, Paley
S, Rhee SY, Shearer AG, Tissier C, Walk TC, Zhang P, Karp PD. 2008. The
MetaCyc Database of metabolic pathways and enzymes and the BioCyc collection
of Pathway/Genome Databases. *Nucleic Acids Res* **36**:D623–31.
doi:10.1093/nar/gkm900
8. Chen Y, Ye W, Zhang Y, Xu Y. 2015. High speed BLASTN: an accelerated
MegaBLAST search tool. *Nucleic Acids Res* **43**:7762–7768.
doi:10.1093/nar/gkv784
9. cxcalc calculator functions. n.d. https://docs.chemaxon.com/display/docs/cxcalc-
calculator-functions.md
10. Djoumbou-Feunang Y, Fiamoncini J, Gil-de-la-Fuente A, Greiner R, Manach C,
Wishart DS. 2019. BioTransformer: a comprehensive computational tool for small

molecule metabolism prediction and metabolite identification. *J Cheminform* **11**:2.
doi:10.1186/s13321-018-0324-5

11. Duigou T, du Lac M, Carbonell P, Faulon J-L. 2019. RetroRules: a database of
reaction rules for engineering biology. *Nucleic Acids Res* **47**:D1229–D1235.
doi:10.1093/nar/gky940

12. Eddy SR. 2009. A NEW GENERATION OF HOMOLOGY SEARCH TOOLS BASED
ON PROBABILISTIC INFERENCEGenome Informatics 2009. PUBLISHED BY
IMPERIAL COLLEGE PRESS AND DISTRIBUTED BY WORLD SCIENTIFIC
PUBLISHING CO. pp. 205–211. doi:10.1142/9781848165632_0019

13. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Res* **32**:1792–1797. doi:10.1093/nar/gkh340

14. Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**:3150–3152.
doi:10.1093/bioinformatics/bts565

15. Guthrie L, Wolfson S, Kelly L. 2019. The human gut chemical landscape predicts
microbe-mediated biotransformation of foods and drugs. *Elife* **8**.
doi:10.7554/eLife.42866

16. Heinken A, Acharya G, Ravcheev DA, Hertel J, Nyga M, Okpala OE, Hogan M,
Magnúsdóttir S, Martinelli F, Preciat G, Edirisinghe JN, Henry CS, Fleming RMT,
Thiele I. 2020. AGORA2: Large scale reconstruction of the microbiome highlights
wide-spread drug-metabolising capacities. *bioRxiv*. doi:10.1101/2020.11.09.375451

17. Integrative HMP (iHMP) Research Network Consortium. 2019. The Integrative
Human Microbiome Project. *Nature* **569**:641–648. doi:10.1038/s41586-019-1238-8

18. Javdan B, Lopez JG, Chankhamjon P, Lee Y-CJ, Hull R, Wu Q, Wang X, Chatterjee
S, Donia MS. 2020. Personalized Mapping of Drug Metabolism by the Human Gut
Microbiome. *Cell* **181**:1661–1679.e22. doi:10.1016/j.cell.2020.05.001

19. Keiser MJ, Roth BL, Armbruster BN, Ernsberger P, Irwin JJ, Shoichet BK. 2007.
Relating protein pharmacology by ligand chemistry. *Nat Biotechnol* **25**:197–206.
doi:10.1038/nbt1284

20. Klatt NR, Cheu R, Birse K, Zevin AS, Perner M, Noël-Romas L, Grobler A,
Westmacott G, Xie IY, Butler J, Mansoor L, McKinnon LR, Passmore J-AS, Abdool
Karim Q, Abdool Karim SS, Burgener AD. 2017. Vaginal bacteria modify HIV
tenofovir microbicide efficacy in African women. *Science* **356**:938–945.
doi:10.1126/science.aai9383

21. Koppel N, Maini Rekdal V, Balskus EP. 2017. Chemical transformation of
xenobiotics by the human gut microbiota. *Science* **356**.
doi:10.1126/science.aag2770

22. Ly LK, Rowles JL 3rd, Paul HM, Alves JMP, Yemm C, Wolf PM, Devendran S,
Hudson ME, Morris DJ, Erdman JW Jr, Ridlon JM. 2020. Bacterial steroid-17,20-
desmolase is a taxonomically rare enzymatic pathway that converts prednisone to
1,4-androstanediene-3,11,17-trione, a metabolite that causes proliferation of
prostate cancer cells. *J Steroid Biochem Mol Biol* **199**:105567.
doi:10.1016/j.jsbmb.2019.105567

23. Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, Greenhalgh
K, Jäger C, Baginska J, Wilmes P, Fleming RMT, Thiele I. 2017. Generation of

genome-scale metabolic reconstructions for 773 members of the human gut
microbiota. *Nat Biotechnol* **35**:81–89. doi:10.1038/nbt.3703

24. Mallory EK, Acharya A, Rensi SE, Turnbaugh PJ, Bright RA, Altman RB. 2018.
Chemical reaction vector embeddings: towards predicting drug metabolism in the
human gut microbiome. *Pac Symp Biocomput* **23**:56–67.
doi:10.1142/9789813235533_0006

25. Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD.
2021. PANTHER version 16: a revised family classification, tree-based
classification tool, enhancer regions and extensive API. *Nucleic Acids Res*
49:D394–D403. doi:10.1093/nar/gkaa1106

26. Nayfach S, Rodriguez-Mueller B, Garud N, Pollard KS. 2016. An integrated
metagenomics pipeline for strain profiling reveals novel patterns of bacterial
transmission and biogeography. *Genome Res* **26**:1612–1625.
doi:10.1101/gr.201863.115

27. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new
versatile metagenomic assembler. *Genome Res* **27**:824–834.
doi:10.1101/gr.213959.116

28. Patel JR, Oh J, Wang S, Crawford JM, Isaacs FJ. 2022. Cross-kingdom expression
of synthetic genetic elements promotes discovery of metabolites in the human
microbiome. *Cell* **185**:1487–1505.e14. doi:10.1016/j.cell.2022.03.008

29. Pollet RM, D'Agostino EH, Walton WG, Xu Y, Little MS, Biernat KA, Pellock SJ,
Patterson LM, Creekmore BC, Isenberg HN, Bahethi RR, Bhatt AP, Liu J,
Gharaibeh RZ, Redinbo MR. 2017. An Atlas of β-Glucuronidases in the Human
Intestinal Microbiome. *Structure* **25**:967–977.e5. doi:10.1016/j.str.2017.05.003

30. Price MN, Dehal PS, Arkin AP. 2009. FastTree: computing large minimum evolution
trees with profiles instead of a distance matrix. *Mol Biol Evol* **26**:1641–1650.
doi:10.1093/molbev/msp077

31. Schneider N, Lowe DM, Sayle RA, Landrum GA. 2015. Development of a novel
fingerprint for chemical reactions and its application to large-scale reaction
classification and similarity. *J Chem Inf Model* **55**:39–53. doi:10.1021/ci5006614

32. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*
30:2068–2069. doi:10.1093/bioinformatics/btu153

33. Sharma AK, Jaiswal SK, Chaudhary N, Sharma VK. 2017. A novel approach for the
prediction of species-specific biotransformation of xenobiotic/drug molecules by the
human gut microbiota. *Sci Rep* **7**:9751. doi:10.1038/s41598-017-10203-6

34. Sievers F, Higgins DG. 2014. Clustal Omega, Accurate Alignment of Very Large
Numbers of Sequences In: Russell DJ, editor. Multiple Sequence Alignment
Methods. Totowa, NJ: Humana Press. pp. 105–116. doi:10.1007/978-1-62703-646-
7_6

35. Spanogiannopoulos P, Bess EN, Carmody RN, Turnbaugh PJ. 2016. The microbial
pharmacists within us: a metagenomic view of xenobiotic metabolism. *Nat Rev*
Microbiol **14**:273–287. doi:10.1038/nrmicro.2016.17

36. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. *Proc Natl Acad Sci U S A* **102**:15545–15550.
doi:10.1073/pnas.0506580102

37. Tian W, Skolnick J. 2003. How well is enzyme function conserved as a function of
pairwise sequence identity? *J Mol Biol* **333**:863–882. doi:10.1016/j.jmb.2003.08.057

38. Xu S, Dai Z, Guo P, Fu X, Liu S, Zhou L, Tang W, Feng T, Chen M, Zhan L, Wu T,
Hu E, Jiang Y, Bo X, Yu G. 2021. ggtreeExtra: Compact Visualization of Richly
Annotated Phylogenetic Data. *Mol Biol Evol* **38**:4039–4042.
doi:10.1093/molbev/msab166

39. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. Ggtree : An r package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol Evol* **8**:28–36. doi:10.1111/2041-210x.12628

40. Zhao C, Dimitrov B, Goldman M, Nayfach S, Pollard KS. 2022. MIDAS2:
Metagenomic Intra-species Diversity Analysis System. *bioRxiv*.
doi:10.1101/2022.06.16.496510

41. Zimmermann M, Zimmermann-Kogadeeva M, Wegmann R, Goodman AL. 2019a.
Separating host and microbiome contributions to drug pharmacokinetics and
toxicity. *Science* **363**. doi:10.1126/science.aat9931

42. Zimmermann M, Zimmermann-Kogadeeva M, Wegmann R, Goodman AL. 2019b.
Mapping human microbiome drug metabolism by gut bacteria and their genes.
Nature. doi:10.1038/s41586-019-1291-3

43. Zmora N, Zilberman-Schapira G, Suez J, Mor U, Dori-Bachash M, Bashiardes S,
Kotler E, Zur M, Regev-Lehavi D, Brik RB-Z, Federici S, Cohen Y, Linevsky R,
Rothschild D, Moor AE, Ben-Moshe S, Harmelin A, Itzkovitz S, Maharshak N,
Shibolet O, Shapiro H, Pevsner-Fischer M, Sharon I, Halpern Z, Segal E, Elinav E.
2018. Personalized Gut Mucosal Colonization Resistance to Empiric Probiotics Is
Associated with Unique Host and Microbiome Features. *Cell* **174**:1388–1405.e21.
doi:10.1016/j.cell.2018.08.041

Supplementary data

Supplementary Table 1 (attached). SIMMER EC predictions for all MetaCyc reactions

Supplementary Table 2 (attached). Literature curated list of drug-metabolism events in the
human gut microbiome

Supplementary Table 3 (attached). Numbers of homologs of characterized bacterial drug-
metabolizing enzymes

Supplementary Table 4 (attached). SIMMER enzyme predictions for positive controls

Supplementary Table 5 (attached). Comparison to existing methods

Supplementary Table 6 (attached). SIMMER EC and enzyme predictions for uncharacterized
reactions

Supplementary Table 7 (attached). Experimental evidence corroboration

Supplementary Table 8 (attached). All SMILES used as SIMMER input in this study

Supplemental figures

Supplementary Figure 1. SIMMER predicts an EC code (i.e. reaction type) for a query
reaction if there is an enrichment of a particular EC at the top of the reaction list. (A) For
each EC code associated with MetaCyc reactions, an enrichment score (ES) was calculated by
walking down the ranked list of reactions. Starting with a score of zero, each time the given EC
is encountered the score increases by one, and each time a different EC is encountered the
score decreases by one. Panel A is an example of such a walk with MetaCyc reaction RXN-
6763. **(B)** Significance is established by comparing the true ES to the ES achieved from 1000
permutations of a shuffled reaction list. Panel B is an example of how RXN-6763 loses its EC
enrichment structure after shuffling. **(C)** Because the MetaCyc database of reactions is
unbalanced in its EC code representation, ES scores for a given EC type are divided by the
number of times the EC in question occurs in the database. This yields a normalized ES (NES)
for SIMMER reporting. This is also performed for the shuffled distributions. **(D)** For the serial
designation of an EC code (the most granular description of an EC code), SIMMER's
performance diminished (Figure 2B), because enrichment calculations suffer from increased
uniqueness in the ranked list and therefore reduced power to determine a match. This is proven
here, as when the database is subsampled to ensure at least three of each unique serial
designation, F1-scores (the harmonic mean of precision and recall) and accuracy remain high
despite the increased EC resolution.

1013

Supplementary Figure 2. **Euclidean distance distributions and silhouette scores for top-level EC codes are resilient to fingerprint type.** Distributions are computed as described in Figure 2A.

1033
Supplementary Figure 3. Euclidean distance distributions and silhouette scores for top-
level EC codes are sensitive to chemical representation type. Distributions are computed as
described in Figure 2A. The resulting score distributions were used to compare SIMMER's
resilience to different chemical representations, such as removal of products and cofactors (the
representation methods employed by DrugBug (Sharma et al., 2017) and MicrobeFDT (Guthrie
et al., 2019)) or the inclusion of exactly one substrate and exactly one product (the Mallory, et
al. method (Mallory et al., 2018)). For all EC classes, scores were smaller within versus
between EC classes using SIMMER's chemical representation, indicating that SIMMER can
detect reaction similarity within EC classes. However, this was not consistently true when using
only the input compound without cofactors or products (substrate), or when using only one
substrate and product (one_substrate_one_product). From the pairs of distributions, we
computed a Kolmogorov-Statistic to determine if the distributions significantly (p<0.05) differed.
This showed that the differences were statistically significant for SIMMER chemical
representations and reduced, or in the wrong direction, when using a reduced representation.
X's in the KS test heatmap indicate an incorrect direction of difference (without grouping closer
than within).

Supplementary Figure 4. An expanded list of gut bacterial enzymes relevant to known cases of drug metabolism. (A) Eleven of the 18 enzymes responsible for positive control drug-metabolism events have high confidence homologs that we gathered by filtering for biological

1054 significance. **(B)** 52,849 total candidate homologs (a median of 1,087 candidates per enzyme)
1055 were obtained via pHMM searches with sequences from the literature. These putative homologs
1056 were filtered for presence in the human jejunum/ileum, presence in RNA-sequencing studies,
1057 and predicted affinity for the substrate in question using SEA. This resulted in 741 additional
1058 enzyme sequences for 31 positive control reactions.
1059
1060

Distance from SIMMER query transformation to all MetaCyc reactions

Distance from SIMMER input (normalized)

1061
Supplementary Figure 5. Distributions of all MetaCyc reactions' euclidean distances to
the positive control list queries. For this analysis, each of the 31 positive control reactions
was queried to SIMMER. Distributions were created based on all MetaCyc reactions' distance to
a given input metabolism event, and colored by their EC class annotation in MetaCyc
(oxidoreductases, transferases, etc.). The dashed blue line depicts the MetaCyc reaction that
yielded a gut microbiome homolog matching the known positive control sequence. For example,

when queried with 5-ASA acetylation, SIMMER outputs a most-similar MetaCyc reaction (RXN-
13871) that SIMMER linked (via sequence similarity) to an N-acetyltransferase known from
previous literature to drive 5-ASA metabolism in the human gut.
1071
1072
1073
1074

1075
Supplementary Figure 6. There was not a significant correlation between a human stool
sample's ability to consume dexamethasone (consumption slope, a.u.) or to produce 17-
oxodexamethasone (production slope, a.u.), and the number of aligned *desAB* sequences.
Patient conversion slopes and metagenomics data were accessed from the original study
(Zimmermann et al., 2019). This adds confidence to the finding described in Figure 6B, as it
means SIMMER predictions were not correlated with dexamethasone metabolism due to co-
occurrence in *C. scindens* with a gene previously reported to underlie bacterial side-chain
cleavage of steroids (Ly et al., 2020).
1084

1085
1086 **Supplementary Figure 7.** SIMMER webtool. The landing page for the SIMMER website
1087 (https://simmer.pollard.gladstone.org/) allows the user to upload a TSV file of queries or add a
1088 single query manually to run SIMMER on. It is recommended to use the command-line tool
1089 (https://github.com/aebustion/SIMMER) for more than 10 input queries.
1090

1091
Supplementary Figure 8. Numbers of principal components used for each fingerprint option in
the precomputed chemical space. All analyses in this manuscript employ a full reaction vector
based on computed Tanimoto coefficients described in Figure 1A. Should a user require a
speed-up in running SIMMER, however, precomputed PCA reductions for the precomputed
chemical space can be employed. Number of components for each type of SIMMER fingerprint
option were chosen based on the Kaiser criterion (i.e. all principal components with eigenvalues
greater than 1.0).

Supplement references

1. Guthrie L, Wolfson S, Kelly L. 2019. The human gut chemical landscape predicts microbe-mediated biotransformation of foods and drugs. *Elife* **8**. doi:10.7554/eLife.42866

2. Javdan B, Lopez JG, Chankhamjon P, Lee Y-CJ, Hull R, Wu Q, Wang X, Chatterjee S, Donia MS. 2020. Personalized Mapping of Drug Metabolism by the Human Gut Microbiome. *Cell* **181**:1661–1679.e22. doi:10.1016/j.cell.2020.05.001

3. Ly LK, Rowles JL 3rd, Paul HM, Alves JMP, Yemm C, Wolf PM, Devendran S, Hudson ME, Morris DJ, Erdman JW Jr, Ridlon JM. 2020. Bacterial steroid-17,20-desmolase is a taxonomically rare enzymatic pathway that converts prednisone to 1,4-androstanediene-3,11,17-trione, a metabolite that causes proliferation of prostate cancer cells. *J Steroid Biochem Mol Biol* **199**:105567. doi:10.1016/j.jsbmb.2019.105567

4. Mallory EK, Acharya A, Rensi SE, Turnbaugh PJ, Bright RA, Altman RB. 2018. Chemical reaction vector embeddings: towards predicting drug metabolism in the human gut microbiome. *Pac Symp Biocomput* **23**:56–67. doi:10.1142/9789813235533_0006

5. Sharma AK, Jaiswal SK, Chaudhary N, Sharma VK. 2017. A novel approach for the prediction of species-specific biotransformation of xenobiotic/drug molecules by the human gut microbiota. *Sci Rep* **7**:9751. doi:10.1038/s41598-017-10203-6

6. Tian W, Skolnick J. 2003. How well is enzyme function conserved as a function of pairwise sequence identity? *J Mol Biol* **333**:863–882. doi:10.1016/j.jmb.2003.08.057

7. Zimmermann M, Zimmermann-Kogadeeva M, Wegmann R, Goodman AL. 2019. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature*. doi:10.1038/s41586-019-1291-3