1

An adhesion signaling axis involving Dystroglycan, β1-Integrin and Cas adaptor

proteins regulates the establishment of the cortical glial scaffold

Running title: Adhesion signaling in glial scaffold development

Wenny Wong [1,*], Jason A. Estep [2,*], Alyssa M. Treptow [2], Niloofar Rajabli [2], Jennifer N.

Jahncke [3], Teresa Ubina [1], Kevin M. Wright [3], Martin M. Riccomagno [1, 2]

[1] Neuroscience Graduate Program,

[2] Cell, Molecular and Developmental Biology Graduate Program, Department of

Molecular, Cell & Systems Biology, University of California, Riverside, CA, USA

[3] Neuroscience Graduate Program, Vollum Institute, Oregon Health & Science

University, Portland, OR 97239, USA.

* These authors contributed equally to this work

Corresponding Author: Martin M. Riccomagno, martinmr@ucr.edu.

Abstract

The mature mammalian cortex is composed of six architecturally and functionally distinct layers. Two key steps in the assembly of this layered structure are the initial establishment of the glial scaffold and the subsequent migration of post-mitotic neurons to their final position. These processes involve the precise and timely regulation of adhesion and detachment of neural cells from their substrates. Although much is known about the roles of adhesive substrates during neuronal migration and the formation of the glial scaffold, less is understood about how these signals are interpreted and integrated within these neural cells. Here, we provide *in vivo* evidence that Cas proteins, a family of cytoplasmic adaptors, serve a functional and redundant role during cortical lamination. *Cas* triple conditional knock-out (*Cas* TcKO) mice display severe cortical phenotypes that feature cobblestone malformations. Molecular epistasis and genetic experiments suggest that Cas proteins act downstream of transmembrane Dystroglycan and β1-Integrin in a radial glial cell-autonomous manner. Overall, these data establish a new and essential role for Cas adaptor proteins during the formation of cortical circuits and reveal a signaling axis controlling cortical scaffold formation.

Keywords: Radial glial cell; Cortical scaffold; Cortical lamination; Cell adhesion; Cas proteins; cobblestone cortex; Type II lissencephaly

Introduction

The cortex is an exquisitely organized structure and provides an excellent model

for understanding the cellular and molecular mechanisms that direct neuronal migration,

lamination and stratification [1,2,3,4,5,6,7,8,9]. The mammalian neocortex is composed

of six architecturally and functionally distinct layers [10]. Layers II to VI of the cortex

develop in an inside first-outside last manner, i.e. Layer VI is born first and Layer II is

born last [5,6,11,12,13]. Excitatory neurons that form these layers are born in the

ventricular zone (VZ) of the dorsal telencephalon and migrate radially towards the pial

surface [2,6,14,15]. After an initial phase of glial-independent migration, the majority of

these newly born neurons will switch from a multipolar to a bipolar morphology near the

intermediate zone (IZ), and associate with Radial Glial Cell (RGC) processes [10,16

17,18,19,20]. The RGC basal process dynamically changes from highly branched to

club-like during early development to establish a scaffold for neuronal migration

[10,21,22]. After this initial remodeling event, RGCs have a bipolar shape with an oval

nucleus, a short apical process at the VZ, and a long basal process contacting the

cortical surface [21]. Neurons use these glial processes as tracks to migrate into the

expanding cortical plate (CP) by glia-guided locomotion [3,4,23,24]. Once neurons

reach the cortical marginal zone (MZ), they detach from the glial fibers and perform

somal translocation to their final location [4,24,25,26]. Thus, two key steps in the

assembly of the cortical layers are the establishment of the glial scaffold and the

migration of post-mitotic neurons to their final positions [5,21,27].

During these events, neurons and RGCs actively interact with each other and the

extracellular matrix (ECM). These processes involve the precise and timely regulation of

adhesion and detachment of neural cells from their substrates [28,29,30]. Newborn

excitatory neurons constantly remodel their Integrin Adhesion Complexes (IAC) as they

migrate radially [29,30,31,32,33,34,35]. In addition, these neurons change adhesive

preferences during their migratory path when switching between glial-independent

translocation and glial-dependent locomotion [4,19,24]. RGCs remodel their points of

contacts and actively engage the ECM as the cortex grows [3,5,21,36,37,38,39,40,41].

Thus, interactions between the basal lamina and the RGC endfeet also play a key role

during migration and lamination. In support of this, β1-Integrin and Dystroglycan (Dag1),

two important transmembrane regulators of IAC establishment and remodeling, have

been shown to be required for cortical histogenesis by acting primarily during the

formation and maintenance of the glial scaffold [42,43,44,45,46,47]. Furthermore,

mutations that disrupt the expression or glycosylation of Dystroglycan cause

cobblestone cortex phenotypes in both human and mouse [46,47,48,49,50].

Although much is known about the roles of adhesive substrates and their

cognate receptors during neuronal migration and the formation of the glial scaffold, less

is understood about how these signals are interpreted and integrated within these

neural cells. The Cas family of cytosolic adaptor proteins are known to participate in

IAC/focal adhesion turnover, and accumulating evidence suggests that Cas protein

phosphorylation and ensuing recruitment of downstream effectors can be modulated by

permissive and adhesive cues *in vitro* [51,52,53]. Cas family members have been

shown to mediate a variety of biological processes including cell migration and changes

in cell morphology *in vitro* [54,55,56,57]. Three of the four mammalian *Cas* genes are

expressed in the developing rodent CNS, including the dorsal telencephalon

[58,59,60,61,62]. Using a conditional gene targeting approach in the mouse to inactivate these three *Cas* genes, we previously uncovered a cell-autonomous role for Cas adaptor proteins acting redundantly to mediate Integrin signaling during the organization of the retinal ganglion cell layer (GCL) [61]. Our findings suggest that inactivation of Cas proteins disrupts normal ganglion cell migration and cell positioning [61]. Interestingly, recent genetic evidence from our lab demonstrates that Cas proteins also act cell-autonomously to regulate the adhesive preferences of mammalian Dorsal Root Ganglia (DRG) sensory axons [62], and a similar role has also been shown in epidermal cells [63]. Taken together, these raise the intriguing possibility that Cas proteins could be acting to regulate cortical stratification.

Here, we show that Cas proteins are essential for normal cortical migration and lamination. We also provide evidence suggesting that these adaptor proteins act in a glial-autonomous manner downstream of β1-Integrin and Dystroglycan. These data thus uncover an adhesive signaling pathway essential for cortical glial scaffold formation and lamination.

Results

As mentioned above, three *Cas* paralogs are expressed during vertebrate development: *p130Cas/BCar1*, *CasL/Nedd9* and *Sin/Efs* [58,59,60,61,62]. A fourth family member, *Cass4*, initially believed to be a pseudogene, appears to only be expressed in the adult [58,64]. We started by undertaking expression analysis of *p130Cas*, *CasL* and *Sin* during cortical development using multiplex RNAscope *in situ*

hybridization [65]. Examination of the RNAscope probes in wild-type (WT) mice at

Embryonic day (E) 12.5 (**Fig 1A**) indicated *p130Cas* transcripts are strongly

concentrated in the developing preplate (PP), with weaker expression in other areas of

the neocortex. *CasL* transcripts are strongest in the cortical hem (midline) and fainter,

but distinctly expressed on the dorsolateral neocortex. In comparison to *p130Cas*, *CasL*

transcripts are less prominent in the PP and are found evenly distributed in the

subventricular zone (SVZ), and ventricular zone (VZ). Based on transcript expression

localization, *p130Cas* puncta are likely found in postmitotic migrating neurons, while

both *p130Cas* and *CasL* are expressed in the proliferative pool of cells in the SVZ and

VZ. *Sin* is undetectable at this stage as indicated by the absence of signal on coronal

sections of the cerebral hemispheres, even at higher magnification. *Cas* gene

expression at E14.5 appears somewhat different from their expression at E12.5 (**Fig

1B**). Noticeably, detection of *p130Cas* transcripts at E14.5 is strongest in the IZ instead

of the cortical plate (CP). Across the two timepoints, *p130Cas* and *CasL* expression

remained consistent in the SVZ and VZ (**Fig 1A and 1B**). The previously undetectable

mRNA transcripts for *Sin* are now clearly localized to the SVZ and VZ. The overlap of

p130Cas, CasL, and *Sin* transcripts in the proliferative zone suggest a possible function

for these genes during cortical neurogenesis.

       We then performed a complementary analysis of Cas expression by

immunohistochemistry on a GENSAT BAC transgenic mouse line that expresses

Enhanced Green Fluorescent Protein (EGFP) under the control of *p130Cas* regulatory

sequences [66,67]. This *p130Cas::BacEGFP* transgenic line allows for the detection of

cells expressing the *BCar1/p130Cas* gene [61,62]. Coronal sections from E12.5 and

E15.5 *p130Cas::BacEGFP* mice were immunostained for EGFP, and co-stained for

Tbr1 to help delineate the PP and CP (**Fig 2A-2C**). No signal is detected when using

the same EGFP antibodies on WT tissue (**S1 Fig**). At E12.5, EGFP was strongly

expressed in the PP, with moderate expression present in the SVZ and VZ (**Fig 2A**). At

E15.5 robust expression is observed in the IZ and CP (**Fig 2B, 2C**). EGFP is also

abundantly expressed in the cortical white matter and blood vessels. Weaker but

specific expression of EGFP was found in the VZ and SVZ. EGFP expression in

p130Cas::BacEGFP is thus mostly consistent with endogenous *p130Cas* mRNA

expression in WT animals, as detected by RNAscope. While p130Cas transcript is

moderately expressed in the CP at E14.5, the strong expression of EGFP in the CP at

E15.5 likely reflects perdurance of EGFP reporter due to accumulation of the

fluorescent protein. Taken together, these expression patterns in the CP and

proliferative regions indicate that *Cas* genes might be involved in several aspects of

cortical development, including cortical neurogenesis, migration, and lamination.

To begin to explore whether Cas adaptor proteins play functional roles during

cortical circuit assembly, we generated a cortical-specific *p130Cas* ablation in

a *CasL$^{-/-}$;Sin$^{-/-}$* double null mutant genetic background (we refer to *p130Cas$^{flox/-}$*

;*CasL$^{-/-}$;Sin$^{-/-}$* mice as triple conditional knock-outs: "TcKO") [61,68,69]. The highly

overlapping expression and redundant roles played by Cas proteins make using the

Cas TcKO mice of vital importance to understand Cas function (**Fig 1A and 1B, 2A-2C**)

[61,62]. To drive *Cre* recombinase expression we used the *Emx1Cre* line, which

expresses *Cre* in early cortical neural progenitors and radial glial cells (RGCs) in the

forebrain [70], resulting in recombination in RGCs themselves and the excitatory

neurons in the cortex and hippocampus (**S2A Fig**). RNAscope on E14.5

Emx1Cre;CasTcKO coronal sections validated that mutant animals do not produce

functional *Cas* transcripts (**S2B Fig**). To test the requirement for *Cas* genes during

cortical histogenesis we started by performing an overall examination of

Emx1Cre;CasTcKO cortical structure using pan-neural markers. When compared to

littermate controls (*TcKO*s without *Cre* and *Emx1Cre; p130Cas$^{+/\,flox}$;CasL$^{-/-}$; Sin$^{-/-}$*

animals), the *Emx1Cre;CasTcKO* adult cortical phenotype appears dramatically

different (**Fig 3A**). The smooth cortical surface, typical for a mouse, is replaced with a

bumpy or cobblestone surface. Nissl stain, which highlights the contrast between axons

and Nissl bodies, shows empty pockets in the cortical plate (**Fig 3A**). A more in-depth

analysis of the cortex with a mature neuronal marker (NeuN) indicated the cortex has

similar empty pockets as seen in the Nissl stain (**Fig 3B**). Interestingly, NeuN+ cells

appear to organize in a wave-like pattern in *Emx1Cre;CasTcKO* animals. In some areas

of these cortices, there are cell clusters forming outside of the pial surface and

infiltrating into the subarachnoid space of the meninges. This displacement of neurons

in the cortex and the cobblestone appearance suggest a possible disruption in the

laminar organization of the cortical plate.

The six-layered cortical plate contains distinct populations of projection neurons

that can be distinguished based on cell morphology, connectivity, or expression of

specific transcription factors [9,13,71]. To test whether the laminar structure of the

cortex is affected in *Emx1Cre;CasTcKO* animals, we performed immunohistochemistry

with a variety of cortical layer markers. Cux1, Ctip2, Rorβ, and Tbr1 were chosen from a

number of transcription factors that maintained their cell lineage identity throughout

development [9,13]. Superficial layer neurons (II/III and IV) are identifiable with Cux1

[72], while Rorβ is one of the few markers that will exclusively label layer IV neurons

[73]. The deep layer neurons can be identified with Ctip2 (high expression in layer V &

low in layer VI) [74] and Tbr1 (layer VI) [75,76]. To precisely quantify any cortical

stratification defects, the lamination patterns of control and *Emx1Cre;CasTcKO* animals

were analyzed by binning the cortex into ten regions of equal-width as previously

described [77,78,79,80,81]. Histological analyses of *Emx1Cre;CasTcKO* animals at P7

and P14 (**Fig 3C-3E; S3A-C Fig**) indicated notable lamination defects for all of the

different populations examined in comparison to control animals. Significant differences

were observed in the localization of Cux1+, Ctip2+, Rorβ+, and Tbr1+ between P7

Emx1Cre;CasTcKO and control animals (**Fig 3D and 3E; S3A-C Fig;** Mann-Whitney *U*

test with Bonferroni correction, *p<0.005 two-tailed test for several bins for each

marker). The observed cortical dysplasia involves the appearance of Ctip2+/Tbr1+ cell

clusters in the upper cortical plate and Cux1+ neurons in the lower cortical plate (**Fig**

3C; S3A and S3C Fig). Furthermore, cortices missing *Cas* genes have a wave-like

appearance for layer IV, with Rorβ+ cells invading the superficial cortical plate and

meninges (**S3B Fig**). This shows that abnormal positioning of both deep- and

superficial-layer neurons contribute to the cobblestone phenotype. This cobblestone

phenotype shows a remarkably strong expressivity and 100% penetrance in all

postnatal *Emx1Cre;CasTcKO* animals analyzed either by laminar markers or

NeuN/Nissl staining, while it was never observed in littermate controls (**Table 1**; n=48

controls, n=46 *Emx1Cre;CasTcKO* animals, Fisher exact test p<0.00001). The number

of focal dysplasia per section in P7 *Emx1Cre;CasTcKO* somatosensory cortex averaged

1.47 ± 0.20 dysplasia/mm^2, while the average was 0 for control animals (**S4 Fig**; Mann-

Whitney *U* test with Bonferroni correction, **p<0.0005 n=8 for each genotype). Similar

phenotypes are also observed in *Emx1Cre;p130Cas$^{flox/-}$;CasL$^{-/-}$;Sin$^{+/-}$* animals, but not

in any other combination of *Cas* family alleles (**S4 Fig,** Mann-Whitney *U* test with

Bonferroni correction, *p<0.005 *Emx1Cre;p130Cas$^{flox/-}$;CasL$^{-/-}$;Sin$^{+/-}$* vs control animals**,**

n=3-8 animals per genotype). Despite the misplacement of these laminar markers, the

overall density of Ctip2+ and Cux1+ cells were not significantly different between control

and *Emx1Cre;CasTcKO* animals (**S3A Fig**; n=5 for each genotype; two tailed t-test *ns*:

p=0.12 for Ctip2+ cells; p=0.09 for Cux1+ cells). Taken together these data revealed

that *Cas* genes are required for cortical lamination.

The misplacement of layer-specific markers observed in *Emx1Cre;CasTcKO*

animals could be due to defects in migration/neural positioning, or defects in neuronal

fate specification [9,82,83,84]. To distinguish between these possibilities, newly born

cells were tracked during the course of radial migration by performing Ethynyl

deoxyuridine (EdU) pulse-chase experiments. This thymidine analog is incorporated in

cells undergoing the S-phase of the cell cycle [85] and can be used to observe the final

position of cells born at a specific timepoint. EdU was administered to pregnant dams at

E12.5 (**Fig 4A**) or E15.5 (**Fig 4B**) to label newly born deep and superficial layer

neurons, respectively. Collection at seven days post-intraperitoneal injection for each

timepoint (E12.5->P0, E15.5->P3) is sufficient for these populations of excitatory

neurons to finish radial migration [7,9,26,39,75,86,87]. Sporadic columns of EdU+ cells

are a distinct feature of *Emx1Cre;CasTcKO* animals compared to controls.

Quantification of the position of E12.5 pulsed cells shows a clear trend for EdU+ cells to

locate closer to the pial surface (Bin 1) in *Emx1Cre;CasTcKO* cortices than in controls,

where they tend to settle in layer VI (**Fig 4A;** Mann-Whitney *U* test with Bonferroni

correction, *p<0.005 two-tailed test for Bin 7, n=5 animals per genotype). More

noticeable differences are observed when animals are pulsed at E15.5 and collected at

P3: prospective superficial neurons that are EdU+ are spread across the ten bins in

Emx1Cre;CasTcKO animals, whereas in control mice these EdU+ cells are primarily

positioned closer to the pial surface (**Fig 4B;** Mann-Whitney *U* test with Bonferroni

correction, *p<0.005 two-tailed test for Bins 2, 3, 5, 6 and 7, n=5). This misplacement of

cells appears to be caused, at least in part, by columns of ectopic EdU cells in the CP.

These data suggest that the cortical dysplasia observed in *Emx1Cre;CasTcKO* animals

is likely due to a neuronal mispositioning defect.

To further confirm that the lamination phenotypes are caused by a migration

defect and not due to cell fate specification errors, we repeated the pulse-chase

experiments but co-labeled the EdU+ neurons with layer specific markers. It is expected

that if the cells born at a particular timepoint are misplaced without changing their fate,

they should maintain expression of the appropriate cortical layer marker. In control

animals 57.0±5% neurons labeled by EdU at E12.5 also co-express the deep layer

marker Tbr1 at P0 (**Fig 4C**). A comparable proportion of EdU+ neurons was co-labeled

by EdU and Tbr1 in P0 *Emx1Cre;CasTcKO* animals that were treated in the same way

(**Fig 4C**; 57.4±8%, Mann-Whitney *U* test with Bonferroni correction, p=1.0 two-tailed

test; n=5 for each genotype). When animals were pulsed with EdU at E15.5 and the

brains were collected at P3 to label superficial layer neurons, the percentage of EdU+

neurons that co-expressed the superficial layer marker Cux1 was also very similar

11

between control and *Emx1Cre;CasTcKO* mice (**Fig 4D;** 78.7±6.4% vs. 75.9±6%

respectively; Mann-Whitney *U* test with Bonferroni correction, p=0.69 two-tailed test;

n=5 for each genotype). These results confirm that the EdU+ cells in

Emx1Cre;CasTcKO mice are properly specified, and thus are most likely mispositioned

due to a migration defect.

When is the cortical dysplasia phenotype first observed in the

Emx1Cre;CasTcKO mutants? By P0 *Emx1Cre;*CasTcKO, animals already display an

overt disruption of cortical organization (**Fig 4C**). This disruption of the laminar

architecture of the cortex by P0 was confirmed by staining *Emx1Cre;*CasTcKO and

control cortices with the cortical marker FoxP2 and the axonal marker L1CAM,

suggesting a possible embryonic onset of this phenotype (**Fig 5A**). To visualize

dysplasia and ectopias at embryonic stages we labeled E12.5 and 15.5 control and

Emx1Cre;CasTcKO cortices for Tbr1, which at those stages labels the PP or majority of

the CP, respectively. In E12.5 *Emx1Cre;CasTcKO* cortices, localization of Tbr1+

neurons appear indistinguishable from controls (**Fig 5B**). However, by E15.5 ectopic

neurons breeching the pial surface are clearly observable in *Emx1Cre;CasTcKO*

embryos but are never observed in control animals (**Fig 5C**). By this stage

Emx1Cre;CasTcKO animals already show ectopic Tbr1+ and Ctip2+ cells positioned

outside of the compromised basal lamina (stained with Laminin) (**Fig 5C-5E**). On

average, at this stage we observed 2.84± 0.3 ectopias/mm of cortical surface length in

Emx1Cre;CasTcKO embryos, but observed 0 in controls (**Fig 5F**; Mann-Whitney *U* test,

*p<0.05 two-tailed test, n=4 animals per genotype). These results establish the

developmental onset of the cobblestone phenotype in *Emx1Cre;CasTcKO* cortices

between embryonic days 12.5 and 15.5.

One possible contributing factor for the migration defects observed in animals is

the improper splitting of the PP [88,89,90]. This involves the establishment of layer VI

within a plexus of pioneer neurons [91,92,93]. Layer VI splits the PP to form the MZ on

the surface of the cortex and the subplate (SP) as the ventral boundary. Successive

waves of neuronal migration and lamination of cortical neurons form the CP in between

these two boundaries. Thus, disruption of this early developmental event can change

laminar organization [13,75,94]. To better understand when and how the migration

defects observed *Emx1Cre;CasTcKO* animals arise, PP splitting was examined in these

mutants and control mice. The use of selected SP and MZ markers at E15.5 indicated a

defective PP split in *Emx1Cre;CasTcKO* cortices compared to age-matched littermate

controls (**S5 Fig**). Immunostaining for Microtubule Associated Protein 2 (MAP2), which

labels differentiating neurons and the cell bodies of SP neurons [75,89], revealed

disorganized SP cells underneath the CP. MAP2+ subplate neurons also form ectopic

clusters that invade the breached MZ (**S5 Fig**). Staining for Chondroitin sulfate

proteoglycan (CSPG), which labels the SP [95], also demonstrated an abnormal

distribution of SP neurons, with columns of CSPG+ cells extending dorsally from the SP

to contact the pial surface in *Emx1Cre;CasTcKO* cortices. To better characterize PP

split phenotypes, we then stained for calcium binding proteins Calretinin and Calbindin

(**S5 Fig**). Calretinin is normally expressed by Cajal-Retzius (CR) cells in the MZ and

thalamocortical projections in the IZ [96]. Calretinin+ cells in the MZ appear

disorganized, forming aggregates on the surface of the cortex, and thalamocortical

afferents appear to invade the cortical plate prematurely in *Emx1Cre;CasTcKO* mice

compared to controls. This early disorganization of the SP, CP, IZ and MZ was

confirmed using Calbindin. Calbindin also labels the MZ and migrating interneurons

[97,98]. Calbindin staining shows a disorganized tangential stream of interneurons

originating from the medial ganglionic eminence and abnormal positioning of CR cells in

the MZ of *Emx1Cre;CasTcKO* cortices. Overall, these suggest that the splitting of the

PP and the subsequent organization of the MZ and SP are severely affected in

Emx1Cre;CasTcKO cortices.

The cobblestone phenotype could be partially caused or compounded by

changes in programed cell death during early developmental timepoints. We examined

the levels of cleaved (active) Capase-3 in control and *Emx1Cre;CasTcKO* cortices at

different developmental stages to visualize the levels of apoptosis before (E12.5), right

after (E15.5) and several days after (P0 and P3) the onset of the cortical dysplasia

phenotype (**S6A-S6D Fig;** Mann-Whitney *U* two-tailed test, p≥0.7 for all stages, n=3

animals per genotype). The density of caspase-3 activation in control and

Emx1Cre;CasTcKO mice was not significantly different at any of these stages. These

data suggest that programmed cell death is not severely disrupted in

Emx1Cre;CasTcKO cortices and is unlikely to be a primary cause of the ectopias and

cortical dysplasia phenotypes observed in these mice.

The use of the *Emx1Cre* mice precludes us from determining whether *Cas* genes

act in a neuronal-autonomous or non-neuronal-autonomous manner during cortical

migration and lamination: recombination in these mice occurs as early as E10 in RGCs,

the neural progenitors that generate all excitatory neurons and macro glial cells in the

dorsal telencephalon [70] (**S2 Fig**). To tease apart the neuronal-autonomous and non-

neuronal-autonomous requirements for *Cas* genes, and their relative contributions to

the cobblestone phenotypes observed in *Emx1Cre;CasTcKO* animals, we generated

NexCre;CasTcKO mice*. NexCre* is expressed only in pyramidal cells, but not in RGCs,

targeting early postmitotic, premigratory excitatory neurons [99,100]. This pattern of

Cre activity was confirmed by crossing the *NexCre* mice to the Cre reporter line *Ai14*

[101]. Analysis of *NexCre;Ai14* cortices at E13.5 revealed that tdTomato expression

was found in the IZ and CP (**S7A Fig**) and E15.5 included the SVZ [99,100]. Validation

of gene inactivation by RNAscope on E16.5 *NexCre;CasTcKO* coronal sections

revealed that mutant animals indeed produce next to no *p130Cas* transcripts in the IZ

and CP, and indicated a moderate but significant reduction of *p130Cas* transcript levels

in the SVZ+VZ (**S7B Fig;** ****$P<0.0001$ one-way ANOVA w/ Tukey Honestly Significant

Difference (HSD) post-hoc test; n=5-6 samples). As expected for the *CasL$^{-/-}$;Sin$^{-/-}$*

background that these mice are bred into, no mRNA is detected for *CasL* or *Sin* even in

controls (**S7B Fig**) [61,68,69].

To elucidate the neuronal-autonomous requirement for *Cas* genes during radial

migration and cortical stratification, the laminar organization of *NexCre;CasTcKO*

cortices was examined (**Fig 6A; S8A-S8B Fig**). *NexCre;CasTcKO* mutants and control

littermates were immunostained with the same layer-specific markers as

Emx1Cre;CasTcKO animals (**Fig 3C; S3A-C Fig**). A prediction is that, if subpopulations

of cortical excitatory neurons are mispositioned in *NexCre;CasTcKO* mice, this will lend

evidence to a neuronal-autonomous requirement for *Cas* genes during cortical migration

and lamination. There are no apparent differences in lamination between control and

NexCre;CasTcKO cortices when these are immunostained with a battery of cortical layer markers (**Fig 6A; S8A-S8B Fig**). The transcription factors are expressed in the appropriate laminar layer at P7, where superficial layer (Cux1+), layer IV (Rorβ+), and deep layer (Ctip2+/Tbr1+) subpopulations are spatially distinct. No significant differences in cell positioning were observed for any of the markers examined in control and *NexCre;CasTcKO* animals (p>0.05, Mann-Whitney *U* test with Bonferroni correction). These results strongly suggest that Cas adaptor proteins are not required in a neuronal-autonomous manner during cortical migration.

To further confirm the lack of migration defects in *NexCre;CasTcKO*, we performed the same unbiased pulse-chase experiment to track migration of early post-mitotic cells that we used on *Emx1Cre;CasTcKO* animals (**Fig 4A and 4B**). EdU administration into *NexCre;CasTcKO* animals at E12.5 (**Fig 6B**) or at E15.5 (**Fig 6C**) resulted in labeling that was indistinguishable from that of control littermates: both deep and superficial layer neurons exhibit a uniform band of EdU+ cells in the cortical plate (Mann-Whitney *U* test with Bonferroni correction, p>0.05 for all bins examined). Thus, the migration defects observed in *Emx1Cre;CasTcKO* animals are not recapitulated in *NexCre;CasTcKO* mutants. This further demonstrates that *Cas* genes act in a non-neuronal autonomous manner during cortical migration, and that they are likely functioning in RGCs to direct cortical lamination.

As the cortex expands in size, radial glial cells create a scaffold for neurons to migrate [10,9,102]. Interactions between the embryonic pial basement membrane and radial glial endfeet are essential for development of the cortex [21,40]. The absence of the cobblestone appearance and intact basement membrane in *NexCre;CasTcKO*

cortices (**Fig 6; S8 Fig**), suggest that the major laminar disruptions observed in the

Emx1Cre;CasTcKO cortices might be a result of radial glial cell dysfunction. Based on

this information, we reexamined the expression of *p130Cas*, *CasL* and *Sin* in the

ventricular zone. To facilitate the labeling of RGC cell bodies and processes we

performed *in utero* electroporation (IUE) of a plasmid that drives the expression of

EGFP under the Radial Glia-specific *Blbp* promoter [103]. The Blbp-EGFP construct

was transfected into WT embryos at E13.5 and tissue was collected at E16.5 to achieve

sparse labeling of RGCs. RNAscope analysis of these brains showed that the

expression of all *Cas* family members was very similar to that observed at E14.5: *CasL*

and *Sin* transcripts were highly enriched in the VZ and SVZ, while *p130Cas* was more

broadly expressed from VZ to CP (**Fig 7A** and **7B**). More importantly, RNAscope

followed by EGFP immunofluorescence confirmed high levels of expression for

p130Cas and *Sin* (**Fig 7A**), and *CasL* (**Fig 7B**) mRNA in the cell bodies of RGCs. Little

expression, if any, was observed in the RGC endfeet. We next sought to independently

validate the expression of *p130Cas* by taking a closer look at EGFP expression in the

p130Cas::BacEGFP animals*.* Coronal sections of E12.5 and E15.5 *p130Cas::BacEGFP*

brains were immunostained for the RGC marker Nestin, the ECM component Laminin

and the intermediate progenitor marker Tbr2 (**Fig 8A-8G**). As mentioned above, EGFP

is broadly expressed at E12.5, with moderate expression in the VZ and SVZ that

overlaps with Nestin and Tbr2, respectively (**Fig 8A-C)**. At this stage, strongest

expression is detected in the PP and basement membrane that are co-labeled by Tbr1

and Laminin, respectively (**Fig 2A** and **Fig 8C**). At E15.5, although EGFP is clearly

present in the VZ co-localizing to cells that express Nestin, reporter expression is

highest in the IZ, CP and around the basement membrane (Fig **2B** and **8D-G**). Taken

together this analysis demonstrates moderate to strong gene expression of *Cas* family

members in RGCs, in addition to other cortical cell types.

To test whether *Cas* genes are required for the embryonic pial basement

membrane and radial glial endfeet integrity, we performed histological assessment of

the cortical scaffold of *Emx1Cre;CasTcKO* at E15.5 (**Fig 9A**). While no obvious defects

were observed in control animals, there are several disruptions to the glial-pial interface

in *Emx1Cre;CasTcKO* cortices. There is widespread rupture of the basal lamina as

indicated by regions with breached Laminin staining. The Nestin+ RGCs do not make

proper adhesion contact with the basal lamina at the exposed Laminin sites, where

basal processes extend into the subarachnoid space. These results suggest that *Cas*

function is required for the maintenance of the embryonic pial basement membrane-

RGC interactions.

To determine whether the disrupted basement membrane-RGC interaction

caused mispositioning of proliferating progenitor cells as observed in other models of

cortical heteropia and dysplasia [47,104], EdU was again administered to

Emx1Cre;CasTcKO animals (**Fig 9B; S9 Fig**). However, collection was 0.5 Hrs.

following injection at E12.5 (**S9 Fig**) or E15.5 (**Fig 9B**) for these experiments. This short

EdU pulse labels the proliferating cells during neurogenesis of deep layer and

superficial layer neurons, respectively. We then stained these brains for EdU and the

intermediate progenitor marker Tbr2. At E12.5, EdU+ proliferating neural progenitors

are positioned in the SVZ and VZ (**S9A Fig**). At this stage, a few proliferative cells are

also observed in the PP of both controls and *Emx1Cre;CasTcKO* animals (**S9A and**

S9B Fig). At E15.5, proliferative cells are mainly restricted to the SVZ and VZ of control

animals, with a few cells labeled by EdU in the CP and MZ. In *Emx1Cre;CasTcKO*

animals, EdU+ cells are also primarily observed in the SVZ and VZ. While occasionally

a few EdU+ cells were localized to the MZ and upper CP regions, no significant

differences were observed between the density of proliferative cells in the CP of

Emx1Cre;CasTcKO and control animals (**Fig 9C**). This indicates that the breakage in

the basement membrane and disruption of radial glial endfeet observed in

Emx1Cre;CasTcKO cortices does not result in ectopic positioning of progenitor cells like

in *Eml1* or *Dystroglycan* mutants [47,104]. To test whether the overall levels of

proliferation and neurogenesis were affected, we measured the density of EdU+ cells in

E12.5 and E15.5 control and *Emx1Cre;CasTcKO* cortices (**Fig 9D** and **S9C Fig**).

Importantly, the density of proliferating cells is not significantly different between these

mutants and controls (Mann-Whitney *U* two-tailed test, p=0.34 for E12.5 and p=0.89 for

E15.5, n=4 animals per genotype). To further examine possible defects in neurogenesis

and to test whether the transition from RGC to intermediate progenitor is disrupted in

Emx1Cre;CasTcKO cortices*,* we quantified the proportion of EdU+ cells that co-

expresses Tbr2. This proportion is not significantly different between control and

Emx1Cre;CasTcKO animals at E12.5 (**S9B and S9D Fig**) or E15.5 (**Fig 9B and 9E;**

Mann-Whitney *U* two-tailed test, p=0.49 for both stages, n=4 animals per genotype),

suggesting that the disruptions in cortical lamination observed in these mutants are also

unlikely to be caused by an early transition from RGC to intermediate progenitor fate.

Dystroglycan (Dag1) provides a structural link between the basement membrane

and components that remodel the actin cytoskeleton [46,105]. Glycosylation of the

extracellular alpha subunit of Dag1 is responsible for maintaining the pial basement

membrane integrity [46,47]. Previous analysis of *NestinCre;Dag1*$^{flox/flox}$ animals at E14.5

to P0.5 showed a heterotopic distribution of cortical neurons [47], and an intriguing

resemblance to the *Emx1Cre;CasTcKO* cortical phenotype. We thus re-examined these

cortical phenotypes at P7 using the same layer markers and the same cortical Cre

driver (*Emx1Cre*) that we used to study the *CasTcKO* mice. Genetic ablation of *Dag1*

using *Emx1Cre* (*Emx1Cre;Dag1*$^{flox/-}$) indeed caused cobblestone malformations, as

suggested by previous studies using a pan-neural Cre [46,47] (**Fig 10A and 10B**).

Mispositioning of the different neuronal subpopulations in *Emx1Cre;Dag1*$^{flox/-}$ mice is

strikingly similar to that of *Emx1Cre;CasTcKO* animals, where superficial layer neurons

under-migrate and deep layer neurons over-migrate (**Fig 10A and 10B**). As in

Emx1Cre;CasTcKO animals*,* the cobblestone cortex phenotype displays strong

expressivity and is 100% penetrant (**Table 1**; n=18 for *Emx1Cre;Dag1*$^{flox/-}$ and controls;

Fisher exact test p<0.0001). Interestingly, these phenotypes are also likely RGC-

autonomous, as deletion of *Dag1* in postmitotic neurons using *NexCre* does not disrupt

cortical lamination [46,106]. In addition to supporting the reproducibility of previous

reports, this provides definitive evidence that the *Emx1Cre;CasTcKO* cortical

organization phenocopies multiple aspects of the defects observed when *Dag1* is

conditionally ablated in early cortical progenitors.

As mentioned above, Dag1 is an adhesion receptor that provides a direct link

between the ECM and pathways involved in cytoskeletal remodeling [107]. While the

extracellular domain of α-Dag1 interacts with ECM components, the cytoplasmic

domain of β-Dag1 can function to regulate ERK/MAPK and Cdc42 pathway activation,

and even directly interact with actin binding proteins [108,109,110,111]. Given the

phenotypic similarities observed in *Emx1Cre;CasTcKO and Emx1Cre;Dag1$^{flox/-}$*, we

asked whether Dag1 could be modulating Cas activity by promoting/enhancing its

phosphorylation to effect downstream signaling. We started by overexpressing myc-

tagged full-length Dystroglycan (myc-Dag1) in the neural cell line Neuro2A and

compared the levels of tyrosine phosphorylated p130Cas (pY165Cas) to that of cells

transfected with a myc-tagged empty vector alone. Transfection of myc-Dag1 increased

pY165Cas levels by over 2.5 fold compared to cells transfected with control vector (**Fig**

10C; 1.03±0.16 au vs. 2.85±0.25 au; One-way ANOVA p<0.001, Tukey HSD test

***p<0.0001). Based on the ability of Dag1 to serve as a scaffold for ERK signaling

components, we hypothesized that the intracellular domain of Dag1 could be required

for Dag1-induced Cas phosphorylation. To test this hypothesis we overexpressed the

extracellular domain of Dag1 (myc-Dag1$_{ecto}$) and compared p130Cas phosphorylation

levels vs control- and full-length Dag1-transfected cells. Surprisingly, myc-Dag1$_{ecto}$

increased pY165Cas levels notably and significantly compared to control (**Fig 10C**;

2.74±0.17 au; One-way ANOVA p<0.001, Tukey HSD test ***p<0.0001). The increase

in pY165Cas was not significantly different than the one observed after full-length myc-

Dag1 transfection (Tukey HSD test ns, p>0.05). This suggests that both full-length and

a truncated form of Dag1 lacking a cytoplasmic domain are sufficient to induce an

increase in p130Cas phosphorylation in a neural cell line.

The intracellular domain of Dag1 appears to be dispensable for the Dag1-

dependent increase in Cas phosphorylation in vitro. Is the cytoplasmic domain of Dag1

required for cortical lamination? To answer this question we examined the cortical

lamination pattern of a knock-in mouse line in which the endogenous *Dag1* coding

sequence was replaced with a truncated form of Dag1 that lacks the intracellular

domain (*Dag1^{βcyto}*), rendering it unable to bind dystrophin/utrophin or initiate ERK/MAPK

or Cdc42 signaling [112]. We performed this analysis using similar markers to those

employed for characterizing the *Emx1Cre;CasTcKO* and *Emx1Cre;Dag1$^{flox/-}$* cortical

phenotypes. Surprisingly, in *Dag1$^{βcyto/-}$* all the layer markers are expressed in the

appropriate pattern in all postnatal animals examined, where superficial layers (Cux1+),

layer IV (Rorβ+), and deep layer neurons (Ctip2+ and Tbr1+) are clearly delineated (**Fig**

11A-C). Furthermore, no evidence of ectopias or cortical dysplasia was observed in

these mutants compared to littermate controls (**Fig 11A-C**; **Table 1**; n=10 postnatal

animals for both genotypes; Fisher Exact test p=1). These data suggest that the

intracellular domain of Dag1 is dispensable for cortical lamination.

　　　　Several studies using cell-based assays have shown that Dag1 is required for

the initial clustering of Laminin-1 on cells [113,114,115,116]. Integrins subsequently

bind to clustered Laminin through a distinct interaction site to transduce outside-in

signaling [116,117,118]. A previous study showed that Cas phosphorylation in the retina

is highly dependent on β1-Integrin function [61]. Interestingly, pan-neural *β1-Integrin*

ablation [43] leads to heterotopias similar to the ones observed in *Emx1Cre;Dag1$^{flox/-}$*

mice and *Emx1Cre;CasTcKO*. These phenotypes also appear to be RGC-autonomous,

since *NexCre*-driven ablation of *β1-Integrin* does not cause these types of defects [45].

The laminar organization of *NestinCre;β1-Integrin* mice has not been fully characterized

[43,45]. To begin to probe whether *β1-Integrin* might be acting together with *Cas* genes

and *Dag1* during cortical scaffold formation, we revisited the cortical phenotypes of *β1-*

Integrin knock-outs using layer-specific markers and *Emx1Cre* as a driver. Staining for

layer-specific transcription factors confirmed a distinct cobblestone phenotype in

Emx1Cre;β1-Integrin$^{flox/flox}$ cortices, as compared to control littermates (*Emx1Cre;β1-*

Integrin$^{flox/+}$) (**Fig 11D-13F**). Cux1, Ctip2, Rorβ, and Tbr1 labeling revealed that cortical

layer misplacement occurred for all the examined subpopulations in *Emx1Cre;β1-*

Integrin$^{flox/flox}$ animals (**Fig 11D-11F**). The cobblestone phenotype shows a high level of

expressivity and 100% penetrance in *Emx1Cre;β1-Integrin$^{flox/flox}$* animals but is never

observed in control littermates (**Table 1**; n=17 postnatal *Emx1Cre;β1-Integrin$^{flox/flox}$*

animals, n=20 control littermates; Fisher Exact Test p<0.0001). On the other hand,

NexCre;β1-Integrin$^{flox/flox}$ and control P7 cortices are indistinguishable from each other:

cortical layers are uniform and defined, as previously shown [45] (**S10A-S10C Fig**).

More importantly, no evidence of cortical dysplasia was observed in these mice (**S10**

Fig; Table 1; 0% for both control and *NexCre;β1-Integrin$^{flox/flox}$*, n=10 and n=8 postnatal

animals respectively; Fisher Exact Test p=1). This reinforces that *β1-Integrin* acts in an

RGC-autonomous manner similar to *Dag1* and *Cas* genes to control cortical lamination.

Based on the fact that Dag1 is sufficient to increase pY165Cas in neural cells

even in the absence of its cytoplasmic domain, the previously established role for β1-

Integrin in regulating Cas function during retina development [61], and the close

resemblance between the *Emx1Cre;CasTcKO, Emx1Cre;Dag1$^{flox/-}$* , and *Emx1Cre;β1-*

Integrin$^{flox/flox}$ cortical phenotypes, we hypothesized that β1-Integrin could be acting as a

signal transducing receptor in RGC for this Dag1-dependent increase in Cas

phosphorylation. To begin to assess this possibility, we first established whether Dag1

expression can modulate Cas tyrosine phosphorylation in WT RGCs as it did in

Neuro2A cells. Mixed primary cultures containing radial glial cells and neurons were

transfected at two days *in vitro* with either an empty vector or myc-Dag1. The presence

of radial glial cells was confirmed using previously validated developmental markers

[119,120] (**S11 Fig**). Subsequently, RGCs were identified by co-labeling with Nestin.

Nestin[+] WT RGCs transfected with an empty vector (pcDNA3.1-myc/his) showed basal

endogenous pY165Cas, where puncta are distributed broadly in cell bodies (**Fig 12A).**

In Nestin[+] myc-Dag1-transfected WT cells, we observed a significant accumulation of

pY165Cas in the glial endbulbs when compared to control transfected WT cells (**Fig**

12A-12C, Mann-Whitney *U* test, ***p<0.0001 two-tailed test). This finding suggests that

Dag1 can modulate p130Cas phosphorylation and may be acting in the same pathway

to regulate RGC endbulb interaction with the pial basement membrane.

To test whether Dag1-dependent phosphorylation of Cas proteins requires β1-

Integrin as a signal transducing receptor, *Emx1Cre;β1-Integrin^{flox/flox}* cultures were

transfected with the empty vector or full-length Dystroglycan (**Fig 12A and 12B**).

Analysis using the Kruskal-Wallis test provided very strong evidence of a difference

(p=0.002) between the mean ranks of at least one pair of treatments, where a series of

Mann-Whitney *U* tests indicated a significant difference between WT RGCs transfected

with myc-Dag1 and all of the other groups (**Fig 12C**; ***p<0.001 two-tailed test).

Importantly, *Emx1Cre;β1-Integrin^{flox/flox}* cells overexpressing myc-Dag1 yielded weak

pY165Cas expression in radial glial endbulbs, in comparison to WT cultures

overexpressing myc-Dag1 (**Fig 12A-12C,** Mann-Whitney *U* test, ***p<0.001 two-tailed

test). Furthermore, no significant difference was observed between *Emx1Cre;β1-*

Integrin^{flox/flox} RGCs transfected with the control vector or myc-Dag1 (**Fig 12A-C)**. These

data indicate that Dag1-dependent phosphorylation or recruitment of pYCas to the

endbulbs requires β1-Integrin function.

These tissue culture experiments and the resemblance between the cortical

phenotypes of the *Dag1*, *β1-Integrin* and *Cas* pan-cortical knock-outs, while suggestive

of a possible interaction, are in no way definitive proof of an epistatic relationship

between these genes. To directly test whether Cas adaptor proteins act downstream of

β1-Integrin to regulate cortical lamination we performed a rescue experiment. We

reasoned that if β1-Integrin-dependent signaling is critically mediated by Cas protein

phosphorylation to regulate cortical migration and lamination, forcing p130Cas tyrosine

phosphorylation into RGC might be able to rescue a β1-Integrin deficiency. With this in

mind, we took advantage of the functional interaction trap (FIT) system [121,122]. FIT

constructs allow for the tethering of a kinase to its target through an engineered, highly

specific binding interface, using a pair of complementary synthetic amphipathic helices

(coiled coils) [121,122]. This highly specific binding interface is used to replace their

natural interaction sites, and results in efficient phosphorylation of the substrate. To test

whether forced p130Cas phosphorylation could rescue the formation of cortical

dysplasia in *Emx1Cre;β1-Integrin^{flox/flox}*, we co-electroporated FIT constructs that prompt

the interaction between Src and Cas into the cortices of these embryos and littermate

controls at E14.5 and collected them at P0. For Src, we used a construct lacking the

substrate binding SH2 and SH3 domains, but containing a coiled-coiled domain (ZipA)

and an intact membrane targeting sequence (Src-ZipA) (**Fig 13A**). The target p130Cas

was fused to ZipB, the coiled-coil segment complementary to ZipA (p130Cas-ZipB) (**Fig**

13A) [121,122]. Due to the difficulty of these experiments and the possibility of the

plasmid titer diluting out inside RGCs as they divide, we opted for driving the expression

of these constructs with a strong constitutive promoter (*EF-1α*) rather than weaker

RGC-specific ones. As negative control, we co-electroporated a similar deletion

construct for Src lacking the ZipA coiled-coil domain (ΔSrc) and p130Cas-ZipB. These

combinations of constructs were co-electroporated with pCAGGS-EGFP to visualize the

electroporated area. Coronal sections of P0 electroporated brains were then stained for

Laminin to label the basement membrane, and Ctip2 to visualize the cortical lamination

pattern. Co-electroporation of the negative control constructs (ΔSrc + p130Cas-ZipB)

into *Emx1Cre;β1-Integrin$^{flox/+}$* or *β1-Integrin$^{flox/flox}$* control animals did not affect cortical

lamination or basement membrane stability (**Fig 13B and 13C**). Importantly, neither did

co-electroporation of Src-ZipA + p130Cas-ZipB into the same control animals (**Fig 13B**).

When ΔSrc + p130Cas-ZipB were electroporated into *Emx1Cre;β1-Integrin$^{flox/flox}$* the

characteristic cobblestone cortex with obvert dysplasia was observed, even in the

EGFP+ region (**Fig 13B and 13C**). This was comparable to the presence of dysplasia

on the uninjected contralateral side (**S12 Fig**). However, introducing Src-ZipA +

p130Cas-ZipB into the *Emx1Cre;β1-Integrin$^{flox/flox}$* mutant cortices notably rescued the

cobblestone phenotype within the EGFP+ electroporated region (**Fig 13B and 13C**):

Ctip2+ cells are positioned in their normal laminar location, with little to no evidence of

dysplasia or ectopias breaking the basement membrane (One-Way ANOVA, p=0.0003

two-tailed test; Tukey HSD **p<0.01 Src + p130Cas-ZipB IUE into *Emx1Cre;β1-*

Integrin$^{flox/flox}$ vs. Src-ZipA + p130Cas-ZipB IUE into *Emx1Cre;β1-Integrin$^{flox/flox}$*; **p<0.01

Src + p130Cas-ZipB IUE into *Emx1Cre;β1-Integrin$^{flox/flox}$* vs. both IUEs into control

animals; *ns:* Src-ZipA + p130Cas-ZipB IUE into *Emx1Cre;β1-Integrin$^{flox/flox}$* vs. both

control IUEs; 6-9 animals per treatment per genotype). This rescue was specific to the

electroporated region since the contralateral side still displayed noticeable dysplasia

(**S12 Fig**). These results unequivocally place Cas protein tyrosine phosphorylation

downstream of β1-Integrin during cortical lamination. Furthermore, the combination of

these genetic and tissue culture experiments strongly suggest that Dag1, β1-Integrin

and Cas act in the same pathway and in a non-neuronal autonomous manner to

regulate basement membrane integrity and cortical lamination.

Discussion

Lamination and stratification of neuronal cell bodies are widespread early

developmental events that organize many regions of the CNS [5,12,13,16,123,124].

These foundational processes are thought to provide a structure upon which neural

circuits can form [125,126,127]. During many of these stratification events, migrating

neurons utilize RGCs to reach the proper layer [10,128,129,130]. In this study we

describe a molecular mechanism by which the glial scaffold is assembled to facilitate

cortical migration and lamination. Our data identify an adhesion signaling axis that acts

in RGCs to maintain glial endfeet-pial interactions and establish a novel and essential

role for Cas proteins in these developmental processes.

Assembly and maintenance of the glial scaffold is particularly important during

neocortical lamination [10,47,131]. The formation of this scaffold requires proper

attachment of the glial endfeet to the pial basement membrane. Disruption of these

interactions in mouse and human results in severe lamination defects like cortical

dysplasia and cobblestone cortex [46,47,50,132]. Why is constant engagement of the

basal lamina by RGCs so critical? While the cortical scaffold is traditionally portrayed as

a somewhat rigid structure, the cortex expands vastly during neurogenesis

[133,134,135]. RGC processes must grow and remodel to accommodate this rapid

expansion of the cortical plate as new layers are born and incorporated [102,131]. RGC

endfeet must actively adhere to the basement membrane to maintain the structure of

the scaffold [40,131,136]. Thus, dynamic regulation of adhesion signaling and

cytoskeletal remodeling are essential for these glial-pial interactions.

Our data support a role for Cas adaptor proteins acting as signal transducers

downstream of the β1-Integrin and Dystroglycan transmembrane proteins to regulate

RGC-basement membrane interactions and maintain the integrity of the glial scaffold. *In

vitro* data further suggests that the Dag1-dependent phosphorylation of Cas proteins is

mediated by β1-Integrin (**Fig 10C** and **Fig 12**) and data from rescue experiments

reinforce this idea (**Fig 13**). This signaling axis is likely modulated by other transducers

of Integrin signaling known to act in non-neuronal-autonomous manner to establish or

maintain the RG scaffold [8,137,138]. While there is a clear disruption of the glial

scaffold in the *Emx1Cre;CasTcKO* animals, the alterations to the dorsal and ventral

boundaries of the CP and clustering of Cajal-Retzius cells observed in these animals

also likely contribute to the lamination defects [36,97,139,140,141], similar to what

occurs in *Dag1* mutant cortices [46,47,142,143,144]. Based on phenotypic analyses,

genetic rescue, and molecular epistasis data we propose a working model whereby the

Dag1-Integrin-Cas adhesion axis acts in at least two ways (**Fig 14**). In cis, it allows

RGCs to anchor and remodel their endfeet as the cortex grows, promoting the

establishment and maintenance of the glial scaffold by interacting with the ECM. Dag1

and Integrin in the RGC endfeet also act in trans through their extracellular domains to

organize and stabilize the basement membrane by nucleating or interacting with ECM

molecules like Laminin. Proper establishment and maintenance of the glial scaffold and

basement membrane allow for normal neuronal migration and lamination. While this

model is supported by genetic experiments by others and us

[8,43,45,46,47,137,138,145], further studies will be needed to test this at the molecular

level and to probe whether other genes with similar roles in glial scaffold formation act in

parallel or in the same signaling pathway [168, 169].

The findings presented here add to a growing list of studies in a variety of animal

models that have demonstrated the evolutionarily conserved requirement for Cas

adaptor proteins during cell migration and axon pathfinding [59,60,61,62]. In the cortex,

the aforementioned Integrin-Cas axis appears to act primarily in a glial-autonomous

manner [43,45]. Genetic evidence and gene expression data strongly support this

function of Cas family members in RGC. However, since we were unable to detect

specific phospho-Cas signal *in vivo*, it is still possible that Cas proteins influence

endfeet attachment and glial scaffold organization by acting in other subcellular

compartments of RGC. Contrary to its glial-autonomous function in the cortex, previous

work demonstrated a neuronal-autonomous function for a similar signaling pathway in

mouse and fly peripheral axon pathfinding, and mammalian retina development

[59,61,62]. During retina lamination and stratification, neurons initiate migration without

the aid of radial glia and the Laminin-rich basement membrane provides an instructive

cue that defines the position where the single-cell ganglion cell layer will form

[61,146,147,148,149,150]. In the eye the Integrin-Cas signaling axis acts in migrating

retinal ganglion cells to sense the inner limiting membrane and is necessary for the

reorganization of the ganglion cell layer [61]. Of note, even though Crk-Associated

Substrates (Cas) are essential for RGC process and endfeet organization, histological

and biochemical evidence support the idea that the Crk proteins themselves function in

a neuronal-autonomous manner downstream of reelin signaling to regulate cortical

lamination [41,90,151].

Cobblestone or type II lissencephaly is a debilitating neurodevelopmental

disorder caused by breaks in the basement membrane (BM) at the pial surface

[50,152,153,154]. These defects have been associated with RGC endfeet that are not

well attached to the ECM [131,155]. The BM breakages result in the over-migration of

neurons and a dramatic loss of normal cortical lamination [37,46,47,142,156]. The

genetic causes of cobblestone lissencephaly have largely been linked to autosomal

recessive mutations in genes that encode for proteins that participate in the post-

translational modification of alpha-Dystroglycan: POMT1, POMT2, POMGNT1,

FCMD/FKTN, FKRP, TMEM5/RXYLT1, ISPD/CRPPA and LARGE1

[46,50,107,157,158]. Mutations in one of these genes are found in 32–66% of patients

with Type II lissencephaly [50,158,159,160].However, despite intensive research in this

area, more than one-third of the cases have an unknown genetic etiology [50].The

striking resemblance of the phenotypes observed in cortical *CasTcKOs* and patients

with cobblestone lissencephaly, in addition to the molecular epistasis experiments

suggesting that Cas proteins act downstream of Dag1 and β1-Integrin in radial glial cells

to maintain basement membrane integrity, pinpoint cytoplasmic effectors of adhesion

signaling as possible contributing factors to the etiology of these neurodevelopmental

disorders. A recent study revealed that several genes involved in regulating cell-matrix

adhesion and IAC assembly are somatically mutated in patients with Focal Cortical

Dysplasia [161]. Patients with FCD type I show disruptions of the cortical laminar

structure that parallel those observed in Type II lissencephaly, but as the name implies,

these dysplasia are more focal in nature, i.e. restricted to a small region of the cortex

[154,162]. Putting our study in the context of these data, our findings open new and

intriguing avenues in the study of the genetic causes of cobblestone lissencephaly and

possibly FCD.

Materials & Methods

Animals

The day of vaginal plug observation was designated as embryonic day 0.5 (E0.5) and

the day of birth was postnatal day 0 (P0). The ICR mouse strain was purchased from

Taconic. The following animals were purchased from The Jackson Laboratory:

Ai14/R26$^{LSL-TdTomato}$ (stock 007914), *β1-Integrin^{flox}* (stock 004605), *Emx1Cre* (stock

005628), and *Dag1^{flox}* (stock 009652). *NexCre* mice were kindly provided by Drs. Nave

and Goebbels [99,100]. Generation of the *p130Cas$^{flox/flox}$*, *CasL$^{-/-}$* and *Sin$^{-/-}$* transgenic

mouse lines has been described previously [61,68,69]. The *Dag1^{βcyto}* mice were kindly

provided by Dr. Kevin Campbell and previously described in [112]. Generation and

characterization of *p130Cas::BacEGFP* animals was described in [61, 62, 66, 67]. All

animal procedures were performed according to the University of California, Riverside's

Institutional Animal Care and Use Committee (IACUC) guidelines.

Immunofluorescence

Embryonic brains were dissected in cold PBS and postnatal mice were perfused then dissected. The brain samples were post-fixed in 4% PFA for 2 Hrs., 4 Hrs., or overnight at 4°C and washed with PBS the next day. The samples were either kept in PBS or sunk in 30% sucrose overnight at 4°C prior to cryopreservation with Optimum Cutting Temperature (Tissue-Tek). Brains were either embedded in 3% agarose for coronal sectioning at 150 μm thickness on a vibrating microtome (VT1000S; Leica) or cryoprotected samples and sectioned coronally at 20 μm on a Leica CM3050 cryostat. Cryosections were blocked for 1 hour in RT in blocking buffer (0.1% PBST and 10% goat serum) in a humidified chamber and free- floating slices were blocked in permeabilization solution overnight at 4°C as previously described [163]. Primary antibodies were diluted in antibody dilution buffer (cryosections; 0.1% PBST, 1% goat serum) or permeabilization solution (free-floating slices) with the following antibodies: rabbit anti-Cux1 (1:250, ProteinTech# 11733-1-AP), rat anti-Ctip2 (1:500, Abcam# ab18465), rabbit anti-Tbr1 (1:500, Abcam# ab31940), rabbit anto-Tbr2 (1:500, Abcam# ab23345), rabbit anti-Rorβ (1:250, Proteintech# 17635-1-AP), rabbit anti-Calretinin (1:500, Swant# 7697), rabbit anti-Calbindin (1:500, Swant# CB38), chicken anti-MAP2 (1:500, Aves MAP), rat anti-NCAM-L1 (1:500, Millipore# MAB5272), rabbit anti FoxP2 (1:500, Abcam# ab16046), chicken anti-EGFP (1:1000, Aves Labs# GFP-1020), rabbit anti-EGFP (1:1000, Thermo Fisher# A-6455) and rabbit anti-CSPG (1:1000; Sigma# AB5320). The samples incubated in the primary antibody mix overnight at 4°C. Cryosections were washed with PBS 15 mins x 4 in a slide mailer and incubated in secondary antibody (goat anti-chicken 647, Themo Fisher Scientific Cat# A-21449 and

goat anti-rabbit 546, Thermo Fisher Scientific Cat# A-11035) with DAPI (1 μg/ml) and antibody dilution buffer at RT for 1 hour in a humidified chamber. Slides were washed with PBS 15 mins x 4 in a slide mailer and sealed with fluorogel. Free floating sections were washed with PBS 15 mins x 8 and incubated in secondary antibody at 1:1000 dilution in permeabilization solution overnight at 4°C. Sections were washed with PBS 15 mins x 4 and sealed with fluorogel. Samples were imaged with a confocal microscope.

Ethynyl deoxyuridine (EdU) Labeling

EdU was dissolved in DMSO and diluted with sterile water to 15 mg/ml at the time of injection. EdU was injected into the intraperitoneal cavity of pregnant mice at a dosage of 150 mg/kg. Animals were collected 30 minutes or 7 days after administration. Embryonic brains were dissected in cold PBS and postnatal animals were perfused then dissected. The brains were fixed for 2 hours in 4% PFA, dehydrated in 30% sucrose overnight, then cryopreserved with Optimum Cutting Temperature (Tissue-Tek). Cryosections were stained according to the instructions in the Click-iT EdU Cell Proliferation Kit for Imaging, Alexa Fluor 488 dye, Invitrogen Cat# C10337. Slides were incubated with 3% BSA for 30 mins at RT, then blocked in 0.5% PBST at RT for 1 hour, washed with 3% BSA in PBS x 2, and incubated in Click-iT Reaction Cocktail (1x Click-iT Reaction Buffer, $CuSO_4$, Alexa Fluor Azide 488, and Reaction Buffer Additive) for 1 hour at RT. Cryosections were washed with 3% BSA in PBS x 4 and immunohistochemistry proceeded as described [163]. Primary antibodies chicken anti-Nestin (1:500, Aves Nes) and rabbit anti-Laminin (1:500, Sigma# L9393) were diluted in

antibody dilution buffer (0.1% PBST, 1% goat serum). Slides incubated in the primary

antibody mix in a humidified chamber at 4°C overnight. Slides were washed with PBS

15 mins x 4 in a slide mailer and incubated in secondary antibody at 1:1000 dilution

(goat anti-chicken 647, Themo Fisher Scientific Cat# A-21449 and goat anti-rabbit 546,

Thermo Fisher Scientific Cat# A-11035) at RT for 1 hour in a humidified chamber.

Slides were washed with PBS 15 mins x 4 in a slide mailer. Fluorogel was applied to the

slide and sealed with a coverslip. Samples were imaged with a confocal microscope.

Mixed Primary Cell Culture

Primary mixed radial glial cells and cortical neurons were isolated from E13.5 embryos

essentially as described [164] using Complete Hank's Balanced Salt Solution (HBSS,

2.5 mM HEPES pH 7.4, 2.5 mM D-glucose, 1 mM $CaCl_2$, 1 mM $MgSO_4$, 2.5 mM

$NaHCO_3$) [163]. Cells were dissociated as described [165] and grown in Radial Glia

Media modified from [165] (Modified DMEM/Eagle's Media, 1x Penicillin/Streptomycin,

0.5 mM Glutamax, 20 mM D-glucose, and 5% horse serum). 12 mm circular coverslips

were treated with 12 M HCL overnight, neutralized with deionized water, and stored in a

glass petri dish with 70-100% ethanol. The coverslips were fire-polished into a 24-well

plate and coated with Laminin (8.3 µg/ml)) and poly-D-lysine/poly-L-lysine (83.3 µg/ml)

[163] in sterile water overnight in a tissue culture incubator at 37°C in 5% CO_2. The

coating media was aspirated and the coverslips were air-dried in an airflow cabinet prior

to plating cells. Dissociated cells were plated in a 24-well plate at a density of 1 x 10^5

cells/well with 500 µl of cell suspension and Radial Glia Media. One-third of the media

was replaced daily and wells were transfected two days *in vitro* (2DIV) with Metafectene

PRO (Biontex Cat# T040-1.0), according to the manufacturer's suggested protocol of 1

µg DNA to 2 µl Metafectene ratio. Cells were transfected with either pcDNA3.1-myc/his

(control vector) or pcDNA3.1-myc/his-Dag1-FL (myc-Dag1) plasmids previously

described [166]. At 48 hours after transfection (4DIV), the coverslips were fixed in 4%

PFA for 10 minutes at RT and washed with PBS with 0.03% NaN_3. Cells incubated in

blocking buffer (0.1% PBST, 10% goat serum) for 30 minutes at RT. Un-transfected

cells were stained with the following primary antibodies: rabbit anti-Blbp (1:500, Abcam#

ab32423), mouse anti-β3-Tubulin (1:500, CST# 4466), rat anti-GFAP (1:500,

Invitrogen# 13-0300), rabbit anti-Tbr2 (1:500, Abcam# ab23345), and chicken anti-

Nestin (1:500, Aves Nes) in antibody dilution buffer (0.1% PBST, 1% goat serum)

overnight at 4°C. Transfected cells incubated in primary antibodies: mouse anti-c-myc

9E10 (1:1000, Invitrogen# 13-2500), rabbit anti-Tyr165 (1:250, CST# 4015), and

chicken anti-Nestin (1:500, Aves Nes) in antibody dilution buffer overnight at 4°C. Cells

were then washed with PBS with 0.03% NaN_3 for 15 mins RT x 4 and incubated in

secondary antibodies at 1:1000 dilution (goat anti-mouse 488, Thermo Fisher Scientific#

A-28175, goat anti-rabbit 647, Thermo Fisher Scientific# A-27040, goat anti-chicken

546, Thermo Fisher Scientific# A-11040, goat anti-rat 647, Thermo Fisher Scientific# A-

21247) and DAPI (1 µg/ml) in antibody dilution buffer at RT for 2 hours. Cells were

washed with PBS with 0.03% NaN_3 for 15 mins x 4 and coverslips were mounted on

microscope slides with fluorogel and imaged with a Leica confocal microscope.

Neuro2A Cell Culture. Neuro2A cells were obtained from ATCC and incubated at 5%

CO_2 at 37°C. The growth media was DMEM with pen/strep (Gibco Cat#15140-122)

supplemented with 10% fetal bovine serum. Cells were visually inspected for

contamination daily and routinely tested for Mycoplasma infection using ATCC

Universal Mycoplasma Detection Kit (ATCC# 30-1012K). Cells were plated on Laminin

and poly-D-lysine (8.3μg/ml) coated 12 mm coverslips in a 24-well plate at 7.8×10^4

cells/cm^2. Cells were transfected 24 hours later with Metafectene PRO (Biontex Cat#

T040-1.0) with according to the manufacturer's suggested protocol with a 1:4 DNA to

Metafectene ratio. Cells were transfected with either pcDNA3.1-myc/his (control vector)

or pcDNA3.1-myc/his-Dag1-FL (myc-Dag1) or pcDNA3.1-myc/his-Dag1-FC (myc-

Dag1$_{ecto}$) plasmids previously described [166]. Cells were fixed 48 hours after

transfection with 4% PFA for 10 minutes at RT, rinsed with PBS, and immunostained as

described above for primary cultures.

Quantification of Layer Markers, EdU and ectopias, and Statistical Analysis

Cells were counted using the RapID Cell Counter [167]. The thickness of the cortex was

divided into 10 bins of equal width and the number of cells labeled with the layer

markers (Cux1, Ctip2, Rorβ, Tbr1) and/or EdU were counted for each bin. Bin 1 is the

marginal zone (MZ) and bin 10 is ventral to layer VI. The proportion of cells in each bin=

number of cells labeled in bin divided by total number of labeled cells. The graph values

are mean ± standard error of mean, n=3-5 independent animals per group depending

on the experiment, and on average 3 sections per sample. For experiments quantifying

density, area or length were quantified using ImageJ. Cell counts for neurons labeled by

one or more markers were performed using RapID Cell Counter [167]. Number of

dysplasia or ectopias were counted manually using the ImageJ multi-point tool. The

datasets were then tested for normality using the Shapiro-Wilk test and QQ plot. For

normally distributed datasets, two tailed t-test was performed for comparing two

samples, and one-way ANOVA followed by Tukey HSD was used for multiple

comparisons. For datasets that didn't fit a normal distribution the Mann-Whitney U test

was performed with Bonferroni correction (e.g. $p<0.005$ was required for significance for

binning experiments with 10 pairwise comparisons).

Mean Fluorescent Intensity of immunofluorescence

ImageJ (NIH) was used to measure the fluorescent intensity of anti-Nestin and anti-

Tyr165 in radial glial endbulb. The Nestin channel was used to trace the borders for the

radial glial endbulb. The following selections were used to determine the fluorescent

intensity of Nestin and pY165Cas channels using ImageJ: Analyze -> Set

Measurements (Area, Min & max gray value, Integrated density, Mean gray value)->

Measure mean gray value. Relative fluorescent intensity is mean fluorescent intensity of

pY165Cas divided by mean fluorescent intensity of Nestin. The graph values are mean

± standard error of mean, n=6-7 independent samples per group, and 9-18 cells per

sample. The dataset was tested for normality using the Shapiro-Wilk test and QQ plot.

For the mixed cultures Kruskal-Wallis test was performed for multiple-comparisons,

followed by a series of Mann-Whitney U tests to examine which pairs are significantly

different. For Neuro2A cells, pY165Cas signal was normalized to transfected cells on

the same image, since expression of myc in the absence of a fusion protein is very low.

For these experiments, datasets were normally distributed and thus ANOVA followed by

Tukey HSD test was performed.

***In utero* electroporation.** Pregnant female mice were anesthetized with isoflurane. An

abdominal incision was made to expose the uterus and intrauterine electroporation was

then conducted as described in [170]. Briefly, pups were visualized through the uterine

wall. Plasmids diluted in fast green and sterile PBS were injected through sharpened

glass capillary needles into the lateral ventricle. 5mm paddles were used to deliver five

35V (E13.5) or 37V (E14.5) pulses of 50 ms each with 950 ms intervals across the

hemispheres. After electroporation, uterine horns were repositioned inside the female

and the abdominal cavity was filled with warm 5x penicillin/streptomycin (pen/strep) in

sterile PBS. For labeling with EGFP under the *BLBP* promoter, E13.5 WT ICR embryos

were injected and electroporated with 3ug pBLBP-EGFP. Pups were collected at E16.5

and processed for RNAscope as described above. For rescue experiments, E14.5

embryos from a *β1-Integrin^{flox/flox}* dam crossed to an *Emx1Cre;β1-Integrin^{flox/+}* sire were

electroporated with a 1:1:1 ratio ΔSrc + p130Cas-ZipB + pCAGGS-EGFP mix or Src-

ZipA + p130Cas-ZipB + pCAGGS-EGFP using 2 μg of each plasmid/IUE. Brains were

collected at P0 after transcardial perfusion with PBS and 4% paraformaldehyde (PFA),

and fixed in 4% PFA for four hours at 4°C. Brain we then rinsed with PBS, and

sectioned coronally at 150 μm on a vibrating microtome (VT100S; Leica) and processed

for immunofluorescence. BLBP-GFP was a gift from Bart De Strooper (Addgene

plasmid # 63174).

Fluorescent *in situ* Hybridization

Fluorescent *in situ* hybridization was performed on 20 μm E12.5, E14.5 or E16.5

cryosections using the RNAScope Multiplex Fluorescent Detection Kit v2 from

Advanced Cell Diagnostics according to manufacturer instructions (ACD, 323110) [65].

Briefly, target RNAs are hybridized to ssDNA "z-probes" complementary to RNA of

interest. Oligos bind to the tail region of the z-probe, which are then bound to amplifiers

labeled with horse radish peroxidase (HRP) and fluorophores. *p130Cas/BCAR1* (ACD,

806961), *CasL/NEDD9* (ACD, 446311), and *Sin/EFS* (ACD, 806951) probes generated

by Advanced Cell Diagnostics were uniquely amplified with Opal dyes (Akoya

Biosciences, FP1487001KT, FP1488001KT, FP1496001KT) and then counterstained

with DAPI (1 µg/ml). Slides were mounted using Fluoro-Gel mounting medium (Electron

Microscopy Sciences, 1798510) and imaged on a Leica SPE confocal microscope.

Quantification was performed in ImageJ: the different cortical areas were outlined based

on the DAPI channel, and the fluorescent channel corresponding to the RNA was then

made binary. The watershed function was then applied, and particles were analyzed

within the ROI using the "analyze particles" function. All counts are reported with ±

standard error of the mean [171].

Acknowledgements

We would like to thank Drs. Sachiko Haga-Yamanaka and Edward Zagha for critically

reading the manuscript and providing helpful comments. We would like to thank Drs.

Nave and Goebbels for the *NexCre* mice. We would also like to thank Drs. Sachiko Seo

and Mineo Kurokawa, and Dr. Konstantina Alexandropoulos for sharing the *CasL*$^{-/-}$ and

Sin$^{-/-}$ mouse lines, respectively. We also would like to thank Bruce Mayer for the FIT

constructs and Bart De Strooper for the Blbp-EGFP plasmid.

Competing interests

The authors declare no competing financial interests.

Availability of data and materials

All data analyzed during this study are included in this article. The data underlying all

the graphs is presented in **S1 Data**-**S14 Data.**

Authors Contributions

WW and JNJ designed and performed experiments, analyzed data and wrote the

manuscript. AMT, NR, JAE, TU, and KMW designed and performed experiments, and

analyzed data. JAE and KMW also provided comments and suggestions for the

manuscript. MMR conceived the project, designed experiments, analyzed data and

wrote the manuscript.

Ethics approval

All animal procedures presented here were performed according to the University of

California, Riverside's Institutional Animal Care and Use Committee (IACUC)-approved

guidelines, under protocol A-20200021

Figure Legends:

Fig. 1. Expression analysis of *Cas* family members. (A-B) Expression analysis of *p130Cas* (magenta), *CasL* (**A**, blue; **B**, green), and *Sin* (**A**, green: **B**, blue) mRNA on coronal sections of E12.5 (**A**) or E14.5 (**B**) WT cortex using RNAscope. PP: preplate; CP: cortical plate; SVZ: subventricular zone; VZ: ventricular zone. Scale bars for A and B (top): 500 µm; A and B (bottom): 100 µm.

Figure 2. Expression analysis of *p130Cas::BacEGFP* during cortical development. (A-C) Coronal sections of E12.5 (**A**) and E15.5 (**B, C**) *p130Cas::BacEGFP* cortices stained for the preplate/cortical plate marker Tbr1 (magenta) and EGFP (green). Nuclei were counterstained with DAPI (blue). PP: preplate; CP: cortical plate; IZ: intermediate zone; SVZ: subventricular zone; VZ: ventricular zone. Scale bars for A: 25 µm; B: 250 µm; C: 75 µm.

Fig. 3. *Cas* genes are required for cortical lamination and neuronal migration. (**A**) Nissl staining of adult control and *Emx1Cre;CasTcKO* coronal cortical sections. Bottom: higher magnification images of Nissl-stained cortices. White arrows mark ectopias. (**B**) Immunostaining of control and *Emx1Cre;CasTcKO* adult coronal sections using the postmitotic neuronal marker NeuN (green) in adult cortices shows a disruption in laminar organization in Emx1Cre;CasTcKO animals. This pan-neuronal marker indicated empty pockets of cells in Emx1Cre;CasTcKO cortices (white arrows). (**C**) Coronal sections of P14 control and *Emx1Cre;CasTcKO* cortices stained for the layer markers Cux1 (green, layer II/III) and Ctip2 (magenta, layer V & VI), counterstained with DAPI (blue). (**D-E**) Quantification of the proportion of Cux1+ (**D**) and Ctip2+ (**E**) cells per bin. Values provided are Mean ± S.E.M., n=3 animals per group, 3 sections per animal, Mann-Whitney U test with Bonferroni correction, *p<0.005 two-tailed test. For data plotted in graphs, see S1 Data. Scale bar for A (top) and B: 1 mm; A (bottom) and C: 500 µm.

Fig. 4. Genetic ablation of *Cas* genes in cortical progenitors results in

misplacement of neurons. (**A-D**) Detection of prospective deep layer- (**A, C**) and

superficial layer-neurons (**B, D**) labeled by EdU (green) at E12.5 and E15.5,

respectively, on coronal sections of control and *Emx1Cre;CasTcKO* cortices. Sections

were also stained for laminin (**A, B**), deep layer marker Tbr1 (**C**) or superficial layer

marker Cux1 (**D**) shown in magenta, and counterstained with DAPI (blue). (**A, B**) In

Emx1Cre;CasTcKO animals ectopic deep layer cells are found in the upper CP (**A,**

white arrows). EdU also revealed distinct columns of cells extending from the superficial

layer into the deep layers (**B,** white arrows) of *Emx1Cre;CasTcKO* animals. Right

panels: quantification of proportion of EdU+ cells per bin. n=5 independent samples per

group, 2-5 sections per sample, Mann-Whitney U test with Bonferroni correction,

*$p<0.005$ two-tailed test for Bin 7 (**A**) and (**B**) Bins 2, 3, 5, 6 and 7. Bin 1 is the MZ and

Bin 10 is ventral to layer VI. (**C, D**) Cells labeled at different timepoints still express the

appropriate laminar marker. Right panels: quantification of proportion of EdU+ cells that

co-express Tbr1 (**C**) or Cux1 (**D**). n=5 animals per genotype, 3-5 section per animal.

Mann-Whitney U Two-tailed test: *ns* p=1 (**C**), *ns* p=0.69 (**D**). Values provided are Mean

± S.E.M. CP: cortical plate; IZ: intermediate zone; SVZ: subventricular zone; VZ:

ventricular zone. For data plotted in graphs, see S4 Data. Scale bars for lower

magnification panels in A, B: 500 µm; higher magnification panels in A, B: 100 µm. C, D:

75 µm.

Fig. 5. Developmental onset of cortical dysplasia in pancortical *Cas* mutants. (A-

E) Coronal sections of control and *Emx1Cre;CasTcKO* cortices at P0 (**A**), E12.5 (**B**) and

E15.5 (**C-E**) stained for FoxP2 (green) and the axonal marker L1CAM (magenta) (**A**),

Tbr1 (green) (**B, C, D**), or Ctip2 (green) and Laminin (magenta) (**E**). While there are no

obvious differences observed between *Emx1Cre;CasTcKO* and controls at E12.5 (**B**),

by E15.5 there are clear ectopias breeching through the pial surface of

Emx1Cre;CasTcKO cortices (**C-E**, white arrows). (**F**) Quantification of the number of

ectopias per length of cortical surface in mm. Values provided are Mean ± S.E.M.

*$p<0.05$, Mann-Whitney U Two-tailed test, n=4 animals per genotype, 3-5 section per

animal. CP: cortical plate; IZ: intermediate zone; SVZ: subventricular zone; VZ:

ventricular zone. For data plotted in graphs, see S5 Data. Scale bars: A: 1 mm; B: 250

µm ; C: 500 µm; D and E: 75 µm.

Fig. 6. Cas adaptor proteins are required in a non-neuronal-autonomous manner

for cortical lamination. (**A**) Coronal cortical sections of P7 control and

NexCre;CasTcKO mice stained for the layer markers Cux1 (green) and Ctip2

(magenta), counterstained with DAPI (blue). Normal positioning of Cux1+ superficial

layer- and Ctip2+ deep layer-neurons. Bottom: quantification of proportion of Cux1+ and

Ctip2+ cells per bin. (**B**) Detection of deep layer neurons born at E12.5 and stained for

EdU (green; pulsed at E12.5 and collected at P0) in coronal sections of control and

NexCre;CasTcKO cortices. (**C**) EdU labeling (green) of superficial layer neurons in

coronal sections of control and *NexCre;CasTcKO* cortices (pulsed at E15.5 and

collected at P3). (**B-C**) Sections were also stained for laminin (magenta) and DAPI

(blue). Right panel: quantification of proportion of EdU+ cells per bin. Migration is not

affected in *NexCre;CasTcKO* mice. Values given are Mean ± S.E.M., n=3 animals per

group, 3 sections per animal, Mann-Whitney *U* test with Bonferroni correction, no

significant differences were observed (p>0.05). Bin 1 is the MZ and Bin 10 is ventral to

layer VI. MZ: marginal zone; CP: cortical plate; IZ: intermediate zone; VZ: ventricular

zone. For data plotted in graphs, see S8 Data. Scale bar for A, lower mag images in B,

C: 500 µm; high magnification panels in B, C: 100 µm.

Fig. 7. Cas genes are highly expressed in RGCs. (A-B) RNAscope for *p130Cas*

(yellow) and *Sin* (cyan) mRNA (**A**), or *CasL* (cyan) (**B**) on coronal sections of E16.5 WT

embryos electroporated with Blbp:EGFP at E13.5. Sections were also immunostained

for EGFP (magenta). Dotted squares in the left panels highlight the regions displayed at

higher magnification in the middle and right panels. All *Cas* genes are highly expressed

in RGC cell bodies (lower panels), but are less abundant in endfeet (upper panels).

Scale bars: 50 µm.

Figure 8. Expression of *p130Cas::BacEGFP* in the VZ and CP. (A-G) Coronal sections of E12.5 (**A-C**) and E15.5 (**D-G**) *p130Cas::BacEGFP* cortices stained for the RGC marker Nestin (**A, D, E, F**), Tbr2 (**B**) or laminin (**C, G**) (magenta), and EGFP (green). Moderate expression is observed in the VZ/SVZ, but expression is highest in CP. DAPI was used to counterstain nuclei (blue). PP: preplate; CP: cortical plate; IZ: intermediate zone; SVZ: subventricular zone; VZ: ventricular zone. Scale bars: A, B, C, E,F,G: 25 µm ; D: 250 µm

Fig. 9. A requirement for Cas adaptor proteins during glial scaffold formation. (**A**) Laminin (green) and Nestin (magenta) antibody staining on E15.5 control and *Emx1Cre;CasTcKO* coronal sections. Radial glial endfeet do not make proper contact with the basal lamina in *Emx1Cre;CasTcKO* animals at exposed laminin sites (arrows). (**B**) Detection of proliferative cells using EdU (green) at E15.5, co-labeled with Tbr2 (magenta). Proliferating cells are occasionally observed within the CP of control and *Emx1Cre;CasTcKO* animals (white arrows). (**C**) Quantification of density of EdU+ cells in CP. Mann-Whitney *U* two-tailed test, p=1, n=4, 4 sections per animal. (**D**) Density of EdU+ cells per area of cortex. Mann-Whitney *U* two-tailed test, p=0.89, n=4 animals per genotype, 3 sections per animal. (**E**) Proportion of EdU+ cells that co-express Tbr2. Mann-Whitney *U* two-tailed test, p=0.49, n=4 animals per genotype, 3-5 sections per animal. Values given are Mean ± S.E.M. CP: cortical plate; MZ: marginal zone; VZ: ventricular zone. For data plotted in graphs, see S10 Data. Scale bars: 100 µm.

Fig. 10. *Dag1* is required for cortical lamination and sufficient to increase Cas phosphorylation in neural cells. (**A-B**) Coronal cortical sections of P7 control and *Emx1Cre;Dag1^{flox/-}* animals stained for the layer markers Cux1 (green) and Ctip2 (magenta) (**A**) or Tbr1 (red) (**B**), counterstained with DAPI (blue). *Emx1Cre;Dag1^{flox/-}* mice display a similar cortical phenotype to *Emx1Cre;CasTcKO* mutants with disorganized distribution of superficial and deep layer neurons. Scale bars for A: 70 µm; B: 500 µm. (**C**) The extracellular domain of Dag1 is sufficient to increase Cas phosphorylation. Immunofluorescence for Myc (magenta) and pY165Cas (cyan) of Neuro2A cells transfected with control vector (myc), full-length Dag1 (myc-Dag1) or the

extracellular domain of Dag1 (myc-Dag1$_{ecto}$). DAPI was used to stain nuclei (yellow). Both myc-Dag1 and myc-Dag1$_{ecto}$ cause a significant increase in pY165Cas. Values provided are Mean ± S.E.M, n=11-23 cells from 3 independent experiments; ANOVA, p<0.0001. Tukey HSD *** p<0.0001 myc-Dag1 vs myc, and myc-Dag1$_{ecto}$ vs. myc. No significant difference was observed between myc-Dag1 and myc-Dag1$_{ecto}$. For data plotted in graphs, see S12 Data. Scale bars for A: 70 μm; B: 500 μm; C: 25 μm.

Fig. 11. The cytoplasmic domain of Dag1 is not required for cortical lamination, but *β1-Integrin* is essential for this process. (A-C) The cytoplasmic domain of Dag1 is dispensable for cortical lamination. Coronal sections of P7 control and *Dag1$^{βcyto/-}$* cortices immunostained for Cux1 (green) and Ctip2 (magenta) (**A**), Rorβ (red) (**B**) or Tbr1 (red) (**C**). DAPI was used as nuclear counterstain (blue). No obvert lamination defects were observed in these animals (n=10). **(D-F)** *β1-Integrin* ablation results in Cobblestone cortex phenotypes. Immunostaining on coronal sections of P7 control and *Emx1Cre;β1-Integrin$^{flox/flox}$* cortices for Ctip2 (magenta) and Cux1 (green) (**D**), RORβ (red) (**E**) or Tbr1 (red) (**F**). Similar laminar disorganization and cobblestone malformation were observed in *Emx1Cre;CasTcKO* animals. Scale bars: 500 μm.

Fig. 12. Dag1 and *β1*-Integrin act upstream of Cas phosphorylation in RGC. (A-B) Immunofluorescence for pY165Cas (cyan), c-Myc (magenta) and Nestin (blue) of WT and *Emx1Cre;β1-Integrin$^{flox/flox}$* mixed cortical cultures transfected with a control vector (myc) or a construct overexpressing myc-tagged full-length Dag1 (myc-Dag). Counterstain in (**A**) is DAPI (yellow). (**B**) is a high magnification view of a representative endbulb boxed in (**A**). Dag1 recruited pY165Cas to the radial glial endbulbs. In the absence of *β1-Integrin*, overexpression of full-length Dag1 does not result in an increase in p130Cas phosphorylation. **(C)** Quantification of the relative fluorescent intensity of pY165Cas at RGC endbulbs. Values given are Mean ± S.E.M., n=6-7 independent samples per group; 9-18 cells per sample, Mann-Whitney *U* test, ***p=0.0001 two-tailed test. For data plotted in graphs, see S13 Data. Scale bars for A: 25 μm; B: 5 μm

Fig. 13. Forced phosphorylation of p130Cas effectively rescues the cobblestone cortex in *β1-Integrin* mutants. (A) Schematic representation of Src-ZipA and p130Cas-ZipB constructs used for IUE. The ΔSrc construct is very similar to Src-ZipA, but missing the ZipA domain. (**B**) Coronal cortical sections of Control or *Emx1Cre;β1-Integrin$^{flox/flox}$* P0 animals that were co-electroporated with ΔSrc + p130Cas-ZipB (negative control) or Src-ZipA + p130Cas-ZipB (FIT rescue). These combinations of plasmids were electroporated with pCAGGS-EGFP. Sections were immunostained for EGFP (yellow), Ctip2 (cyan) and laminin (magenta). Nuclei were counterstained with DAPI (blue). Note the evenly layered distribution of Ctip2+ cells in *Emx1Cre;β1-Integrin$^{flox/flox}$* cortices after IUE with Src-ZipA + p130-ZipB. (**C**) Quantification of rescue experiments. Values given are Mean ± S.E.M., n=6-9 independent samples per group; ANOVA p<0.01. Tukey HSD ** p<0.01 for *Emx1Cre;β1-Integrin$^{flox/flox}$* + ΔSrc + p130Cas-ZipB vs. all other groups. There is no significant difference between *Emx1Cre;β1-Integrin$^{flox/flox}$* + Src-ZipA + p130Cas-ZipB (FIT rescue) and Control animals electroporated with either combination of plasmids. For data plotted in graphs, see S14 Data. Scale bar: 250 µm.

Fig. 14. A working model for an adhesion signaling axis regulating RGC scaffold formation and basement membrane maintenance. A Dag1-Integrin-Cas axis acts in *cis* in RGCs to maintain attachment to the basement membrane as the cortex expands**.** In addition, Dag1 and β1-Integrin participate in the maintenance of the basement membrane by acting in *trans* through their extracellular domains. Additional cytoplasmic IAC proteins that appear in grey have been showed to participate in cortical lamination based on genetic evidence and we speculate that they could be acting in the same pathway [29,30,31,32,33,34,35].

References

1. Ramón y Cajal S. Histologie du Système Nerveux de l'Homme et des Vertébrés. 1911(2). doi: 10.5962/bhl.title.48637

2. Caviness VS, Takahashi T. Proliferative events in the cerebral ventricular zone. Brain Dev. 1995;17(3): 159–163. doi: 10.1016/0387-7604(95)00029-B

3. Hatten ME. Central nervous system neuronal migration. Annu. Rev. Neurosci. 1999;22: 511–539. doi: 10.1146/annurev.neuro.22.1.511

4. Nadarajah B, Brunstrom JE, Grutzendler J, Wong RO, Pearlman AL. Two modes of radial migration in early development of the cerebral cortex. Nat. Neurosci. 2001;4(2): 143–150. doi: 10.1038/83967

5. Marín O, Rubenstein JLR. Cell migration in the forebrain. Annu. Rev. Neurosci. 2003;26: 441–483. doi: 10.1146/annurev.neuro.26.041002.131058

6. Kriegstein AR, Noctor SC. Patterns of neuronal migration in the embryonic cortex. Trends Neurosci. 2004;27(7): 392–399. doi: 10.1016/j.tins.2004.05.001

7. Ayala R, Shu T, Tsai LH. Trekking across the Brain: The Journey of Neuronal Migration. Cell. 2007;128(1): 29-43. doi: 10.1016/j.cell.2006.12.021

8. Valiente M, Marín O. Neuronal migration mechanisms in development and disease. Curr. Opin. Neurobiol. 2010;20(1): 68–78. doi: 10.1016/j.conb.2009.12.003

9. Kwan KY, Sestan N, Anton ES. Transcriptional co-regulation of neuronal migration and laminar identity in the neocortex. Development. 2012;139(9): 1535-1546. doi: 10.1242/dev.069963

10. Rakic P. Mode of cell migration to the superficial layers of fetal monkey neocortex. J. Comp. Neurol. 1972;45(1): 61–83. doi: 10.1002/cne.901450105

11. Marin-Padilla M. Dual origin of the mammalian neocortex and evolution of the cortical plate. Anat. Embryol. 1978;152(2): 109–126. doi: 10.1007/BF00315920

12. O'Leary DD, Koester SE. Development of projection neuron types, axon pathways, and patterned connections of the mammalian cortex. Neuron. 1993;10(6): 991–1006. doi: 10.1016/0896-6273(93)90049-w

13. Molyneaux BJ, Arlotta P, Menezes JRL, Macklis JD. Neuronal subtype specification in the cerebral cortex. Nat. Rev. Neurosci. 2007;8: 427-437. doi: 10.1038/nrn2151

14. Angevine Jr JB, Sidman RL. Autoradiographic study of cell migration during histogenesis of cerebral cortex in the mouse. Nature. 1961;192: 766–768. doi: 10.1038/192766b0

15. Luskin MB. Restricted proliferation and migration of postnatally generated neurons derived from the forebrain subventricular zone. Neuron. 1993;11(1): 173–189. doi: 10.1016/0896-6273(93)90281-u

16. Rakic P. Specification of cerebral cortical areas. Science. 1988;241(4862): 170–176. doi: 10.1126/science.3291116

17. Gadisseux JF, Evrard P, Mission JP, Caviness VS. Dynamic structure of the radial glial fiber system of the developing murine cerebral wall. An immunocytochemical analysis. Brain Res. Dev. 1989;50(1): 55–67. doi: 10.1016/0165-3806(89)90126-0

18. Noctor SC, Flint AC, Weissman TA, Dammerman RS, Kriegstein AR. Neurons derived from radial glial cells establish radial units in neocortex. Nature. 2001;409(6821): 714–720. doi: 10.1038/35055553

19. Tabata H, Nakajima K. Multipolar Migration: The Third Mode of Radial Neuronal Migration in the Developing Cerebral Cortex. J. Neurosci. 2003;23(31): 9996-10001. doi: 10.1523/JNEUROSCI.23-31-09996.2003

20. Loturco JJ, Bai J. The multipolar stage and disruptions in neuronal migration. Trends Neurosci. 2006;29(7): 407–413. doi: 10.1016/j.tins.2006.05.006

21. Rakic P. Developmental and evolutionary adaptations of cortical radial glia. Cereb. Cortex. 2003;13(6): 541–549. doi: 10.1093/cercor/13.6.541

22. Yokota Y, Eom TY, Stanco A, Kim WY, Rao S, Snider WD, Anton ES. Cdc42 and Gsk3 modulate the dynamics of radial glial growth, inter-radial glial interactions and polarity in the developing cerebral cortex. Development. 2010;137(23): 4101–4110. doi: 10.1242/dev.048637

23. Hatten ME. Riding the glial monorail: A common mechanism for glialguided neuronal migration in different regions of the developing mammalian brain. Trends Neurosci. 1990;13(5): 179-184. doi: 10.1016/0166-2236(90)90044-B

24. Nadarajah B, Parnavelas J. Modes of neuronal migration in the developing cerebral cortex. Nat. Rev. Neurosci. 2002;3: 423–432. doi: 10.1038/nrn845

25. Miyata T, Kawaguchi A, Saito K, Kawano M, Muto T, Ogawa M. Asymmetric production of surface-dividing and non-surface-dividing cortical progenitor cells. Development. 2004;131(13): 3133-3145. doi: 10.1242/dev.01173

26. Nadarajah B, Alifragis P, Wong ROL, Parnavelas JG. Neuronal migration in the

developing cerebral cortex: Observations based on real-time imaging. Cerebr. Cortex.

2003;13(6): 607-611. doi: 10.1093/cercor/13.6.607

27. Noctor SC, Martínez-Cerdeño V, Ivic L, Kriegstein AR. Cortical neurons arise in

symmetric and asymmetric division zones and migrate through specific phases. Nat.

Neurosci. 2004;7(2): 136–144. doi: 10.1038/nn1172

28. Barros CS, Franco SJ, Müller U. Extracellular matrix: functions in the nervous

system. Cold Spring Harb. Perspect. Biol. 2001;3(1): a005108. doi:

10.1101/cshperspect.a005108

29. Kawauchi T. Regulation of cell adhesion and migration in cortical neurons: Not only

Rho but also Rab family small GTPases. Small GTPases. 2011;2(1): 36–40. doi:

10.4161/sgtp.2.1.15001

30. Kawauchi T. Cell adhesion and its endocytic regulation in cell migration during

neural development and cancer metastasis. Int. J. Mol. Sci. 2012;13(4): 4564–4590.

doi: 10.3390/ijms13044564

31. Rutishauser U, Jessell TM. Cell adhesion molecules in vertebrate neural

development. Physiol. Rev. 1988;68(3), 819–857. doi: 10.1152/physrev.1988.68.3.819

32. Sanes JR. Extracellular matrix molecules that influence neural development. Annu.

Rev. Neurosci. 1989;12: 491–516. doi: 10.1146/annurev.ne.12.030189.002423

33. Rakic RS, Cameron H, Komuro H. Recognition, adhesion, transmembrane

signaling, and cell motility in guided neuronal migration. Curr. Opin. Neurobiol.

1994;4: 63-69. doi: 10.1016/0959-4388(94)90033-7

34. Anton ES, Kreidberg JA, Rakic P. Distinct functions of α3and α(v) Integrin receptors

in neuronal migration and laminar organization of the cerebral cortex. Neuron.

1999;22(2): 277–289. doi: 10.1016/S0896-6273(00)81089-2

35. Ridley AJ, Schwartz MA, Burridge K, Firtel RA, Ginsberg MH, Borisy G, et al. Cell

Migration: Integrating Signals from Front to Back. Science. 2003;302(5651): 1704-1709.

doi: 10.1126/science.1092053

36. Hartfuss E, Förster E, Bock HH, Hack MA, Leprince P, Lugue JM, et al. Reelin

signaling directly affects radial glia morphology and biochemical maturation.

Development. 2003;130(19): 4597–4609. doi: 10.1242/dev.00654

37. Haubst N, Georges-Labouesse E, Arcangelis AD, Mayer U, Go□tz M. Basement

membrane attachment is dispensable for radial glial cell fate and for proliferation, but

affects positioning of neuronal subtypes. Development. 2006;133(16): 3245–3254. doi:

10.1242/dev.02486

38. Lu P, Takai K, Weaver VM, Werb Z. Extracellular matrix degradation and
remodeling in development and disease. Cold Spring Harb. Perspect. Biol. 2011;3(12):
a005058. doi: 10.1101/cshperspect.a005058

39. Lambert de Rouvroit, C,Goffinet, AM. Neuronal migration. Mech. Dev. 2001;105(1-
2): 47–56. doi: 10.1016/s0925-4773(01)00396-3

40. Halfter W, Dong S, Yip YP, Willem M, Mayer U. A critical function of the pial
basement membrane in cortical histogenesis. J. Neurosci. 2002;22(14): 6029–6040.
doi: 10.1523/JNEUROSCI.22-14-06029.2002

41. Franco SJ, Müller U. Extracellular matrix functions during neuronal migration and
lamination in the mammalian central nervous system. Dev. Neurobiol. 2011;71(11):
889–900. doi: 10.1002/dneu.20946

42. Georges-Labouesse E, Mark M, Messaddeq N, Gansmüller A. Essential role of
alpha 6 Integrins in cortical and retinal lamination. Curr. Biol. 1998;8(17): 983–986. doi:
10.1016/s0960-9822(98)70402-6

43. Graus-Porta D, Blaess S, Senften M, Littlewood-Evans A, Damsky C, Huang Z, et
al. Beta1-class Integrins regulate the development of laminae and folia in the cerebral

and cerebellar cortex. Neuron. 2001;31(3): 367–379. doi: 10.1016/s0896-6273(01)00374-9

44. Schmid RS, Anton ES. Role of Integrins in the Development of the Cerebral Cortex. Cereb. Cortex. 2003;13(3): 219–224. doi: 10.1093/cercor/13.3.219

45. Belvindrah R, Graus-Porta D, Goebbels S, Nave K-A Muller U. B1 Integrins in Radial Glia But Not in Migrating Neurons Are Essential for the Formation of Cell Layers in the Cerebral Cortex. J. Neurosci. 2007;27(50): 13854–13865. doi: 10.1523/JNEUROSCI.4494-07.2007

46. Satz, JS, Ostendorf AP, Hou S, Turner A, Kusano H, Lee JC, et al. Distinct functions of glial and neuronal dystroglycan in the developing and adult mouse brain. J. Neurosci. 2010;30(43): 14560–14572. doi: 10.1523/JNEUROSCI.3247-10.2010

47. Myshrall TD, Moore SA, Ostendorf AP, Satz JS, Kowalczyk T, Nguyen H, et al. Dystroglycan on radial glia end feet is required for pial basement membrane integrity and columnar organization of the developing cerebral cortex. J. Neuropathol. Exp. 2012;71(12): 1047–1063. doi: 10.1097/NEN.0b013e318274a128

48. Endo T. Dystroglycan glycosylation and its role in alpha-dystroglycanopathies. Acta Myol. 2007;26(3): 165–170.

49. Muntoni F, Brockington M, Godfrey C, Ackroyd M, Robb, S, Manzur A. Muscular dystrophies due to defective glycosylation of dystroglycan. Acta Myol. 2007;26(3): 129–135.

50. Devisme L, Bouchet C, Gonzals M, Alanio E, Bazin A, Bessires B, et al. Cobblestone lissencephaly: Neuropathological subtypes and correlations with genes of dystroglycanopathies. Brain. 2012;135(2): 469-382. doi: 10.1093/brain/awr357

51. Vuori K, Ruoslahti E. Tyrosine phosphorylation of p130Cas and cortactin accompanies Integrin-mediated cell adhesion to extracellular matrix. J. Biol. Chem. 1995;270(38): 22259–22262. doi: 10.1074/jbc.270.38.22259

52. Harte MT, Hildebrand JD, Burnham MR, Bouton, AH, Parsons. p130Cas, a substrate associated with v-Src and v-Crk, localizes to focal adhesions and binds to focal adhesion kinase. J. Biol. Chem. 1996;271(23): 13649–13655. doi: 10.1074/jbc.271.23.13649

53. Nakamoto T, Sakai R, Honda H, Ogawa S, Ueno H, Suzuki T, et al. Requirements for localization of p130cas to focal adhesions. Mol. Cell. Biol. 1997;17(7): 3884–3897. doi: 10.1128/MCB.17.7.3884

54. Burnham MR, Harte MT, Richardson A, Parsons JT, Bouton AH. The identification

of p130cas-binding proteins and their role in cellular transformation. Oncogene.

1996;12(11): 2467–2472.

55. Klemke RL, Leng J, Molander R, Brooks, PC, Vuori K, Cheresh DA. CAS/Crk

coupling serves as a "molecular switch" for induction of cell migration. J. Cell Biol.

1998;140(4): 961-972. doi: 10.1083/jcb.140.4.961

56. Cho, SY, Klemke, RL. Extracellular-regulated kinase activation and CAS/Crk

coupling regulate cell migration and suppress apoptosis during invasion of the

extracellular matrix. J. Cell Biol. 2000;149(1): 223-236. doi: 10.1083/jcb.149.1.223

57. Tikhmyanova N, Little JL, Golemis EA. CAS proteins in normal and pathological cell

growth control. Cell. Mol. Life Sci. 2010;67(7): 1025–1048. doi: 10.1007/s00018-009-

0213-1

58. Merrill RA, See AWM, Wertheim ML, Clagett-Dame M. Crk-associated substrate

(Cas) family member, NEDD9, is regulated in human neuroblastoma cells and in the

embryonic hindbrain by all-trans retinoic acid. Dev. Dyn. 2004;231(3): 564-575. doi:

10.1002/dvdy.20159

59. Huang Z, Yazdani U, Thompson-Peer KL, Kolodkin AL,Terman JR. Crk-associated

substrate (Cas) signaling protein functions with Integrins to specify axon guidance

during development. Development. 2007;134(12): 2337–2347. doi: 10.1242/dev.004242

60. Liu G, Li W, Gao X, Li X, Jürgensen C, Park HT, et al. p130CAS is required for

netrin signaling and commissural axon guidance. J. Neurosci. 2007;27(4): 957–968. doi:

10.1523/JNEUROSCI.4616-06.2007

61. Riccomagno MM, Sun LO, Brady CM, Alexandropoulos K, Seo S, Kurokawa M, et

al. Cas adaptor proteins organize the retinal ganglion cell layer downstream of Integrin

signaling. Neuron. 2014;81(4): 779-786. doi: 10.1016/j.neuron.2014.01.036

62. Vahedi-Hunter TA, Estep JA, Rosette KA, Rutlin ML, Wright KM, Riccomagno MM.

Cas Adaptor Proteins Coordinate Sensory Axon Fasciculation. Sci. Rep. 2018;8: 5996.

doi: 10.1038/s41598-018-24261-x

63. Camacho Leal M, Costamagna A, Tassone B, Saoncella S, Simoni M, Natalini D, et

al. Conditional ablation of p130Cas/BCAR1 adaptor protein impairs epidermal

homeostasis by altering cell adhesion and differentiation. Cell Commun. Sig.

2018;16(1): 73. doi: 10.1186/s12964-018-0289-z

64. Deneka A, Korobeynikov V, Golemis, EA. Embryonal Fyn-associated substrate

(EFS) and CASS4: The lesser-known CAS protein family members. Gene. 2015;570(1):

25–35. doi: 10.1016/j.gene.2015.06.062

65. Wang F, Flanagan J, Su N, Wang LC, Bui S, Nielson A, et al. RNAscope: a novel in

situ RNA analysis platform for formalin-fixed, paraffin-embedded tissues. J. Mol. Diagn.

2012;14(1): 22–29. doi: 10.1016/j.jmoldx.2011.08.002

66. Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, et al. A gene

expression atlas of the central nervous system based on bacterial artificial

chromosomes. Nature. 2003;425(6961): 917-25. doi: 10.1038/nature02033

67. Heintz N. Gene Expression Nervous System Atlas (GENSAT). Nat. Neurosci.

2004;7: 483. doi: 10.1038/nn0504-483

68. Donlin LT, Danzl NM, Wanjalla C, Alexandropoulos K. Deficiency in Expression of

the Signaling Protein Sin/Efs Leads to T-Lymphocyte Activation and Mucosal

Inflammation. Mol. Cell. Bio. 2005;25(24): 11035-11046. doi:

10.1128/MCB.25.24.11035-11046.2005

69. Seo S, Asai T, Saito T, Suzuki T, Morishita Y, Nakamoto T, et al. Crk-associated

substrate lymphocyte type is required for lymphocyte trafficking and marginal zone B

cell maintenance. J. Immunol. 2005;175(6): 3492–501. doi:

10.4049/jimmunol.175.6.3492

70. Gorski JA, Talley T, Qiu M, Puelles L, Rubenstein JLR, Jones KR. Cortical

excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-

expressing lineage. J. Neurosci. 2002;22(15): 6309–6314. doi:

10.1523/JNEUROSCI.22-15-06309.2002

71. Hevner RF. Layer-specific markers as probes for neuron type identity in human

neocortex and malformations of cortical development. J. Neuropathol. Exp. 2007;66(2):

101–109. doi: 10.1097/nen.0b013e3180301c06

72. Ferrere A, Vitalis T, Gingras H, Gaspar P, Cases O. Expression of Cux-1 and Cux-2

in the developing somatosensory cortex of normal and barrel-defective mice. Anat. Rec.

2006;288(2): 158–165. doi: 10.1002/ar.a.20284

73. Nakagawa Y, O'Leary DD. Dynamic patterned expression of orphan nuclear

receptor genes RORalpha and RORbeta in developing mouse forebrain. Dev.

Neurosci. 2003;25(2-4): 234–244. doi: 10.1159/000072271

74. Leid M, Ishmael JE, Avram D, Shepherd D, Fraulob V, Dollé P. CTIP1 and CTIP2

are differentially expressed during mouse embryogenesis. Gene Expr. Patterns.

2004;4(6): 733–739. doi: 10.1016/j.modgep.2004.03.009

75. Hevner RF, Shi L, Justice N, Hsueh Y, Sheng M, Smiga S, et al. Tbr1 regulates

differentiation of the preplate and layer 6. Neuron. 2001;29(2): 353–366. doi:

10.1016/s0896-6273(01)00211-2

76. Bedogni F, Hodge RD, Elsen GE, Nelson BR, Daza RA, et al. Tbr1 regulates

regional and laminar identity of postmitotic neurons in developing neocortex. PNAS.

2010;107(29): 13129–13134. doi: 10.1073/pnas.1002285107

77. Rosário M, Schuster S, Jüttner R, Parthasarathy S, Tarabykin V, Birchmeier W.

Neocortical dendritic complexity is controlled during development by NOMA-GAP-

dependent inhibition of Cdc42 and activation of cofilin. Genes Dev. 2012;26(15): 1743–

1757. doi: 10.1101/gad.191593.112

78. Franzoni E, Booker SA, Parthasarathy S, Rehfeld F, Grosser S, Srivatsa S, et al.

miR-128 regulates neuronal migration, outgrowth and intrinsic excitability via the

intellectual disability gene Phf6. eLife. 2015;4: e04263. doi: 10.7554/eLife.04263

79. Tonchev AB, Tuoc TC, Rosenthal EH, Studer M, Stoykova, A. Zbtb20 modulates the

sequential generation of neuronal layers in developing cortex. Mol. Brain. 2016;9: 65.

doi: 10.1186/s13041-016-0242-2

80. Umemura M, Kaneko Y, Tanabe R, TakahashiY. ATF5 deficiency causes abnormal cortical development. Sci. Rep. 2021;11: 7295. doi: 10.1038/s41598-021-86442-5

81. Sokpor G, Kerimoglu C, Nguyen H, Pham L, Rosenbusch J, Wagener R, et al. Loss of BAF Complex in Developing Cortex Perturbs Radial Neuronal Migration in a WNT Signaling-Dependent Manner. Front. Mol. Neurosci. 2021;14: 687581. doi: 10.3389/fnmol.2021.687581

82. Guillemot F. Cell fate specification in the mammalian telencephalon. Prog. Neurobiol. 2007;83(1): 37–52. doi: 10.1016/j.pneurobio.2007.02.009

83. Pramparo T, Youn YH, Yingling J, Hirotsune S, Wynshaw-Boris A. Novel embryonic neuronal migration and proliferation defects in Dcx mutant mice are exacerbated by Lis1 reduction. J. Neurosci. 2010;30(8): 3002–3012. doi: 10.1523/JNEUROSCI.4851-09.2010

84. Wang H, Ge G, Uchida Y, Luu B, Ahn S. Gli3 is required for maintenance and fate specification of cortical progenitors. J. Neurosci. 2011;31(17): 6440–6448. doi: 10.1523/JNEUROSCI.4892-10.2011

85. Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of DNA synthesis in vivo. PNAS.2008;105(7): 2415-2420. doi: 10.1073/pnas.0712168105

86. Frantz GD, McConnell, SK. Restriction of late cerebral cortical progenitors to an upper-layer fate. Neuron. 1996;17(1): 55–61. doi: 10.1016/s0896-6273(00)80280-9

87. Kwan KY, Lam MM, Krsnik Z, Kawasawa YI, Lefebvre V, Sestan N. SOX5 postmitotically regulates migration, postmigratory differentiation, and projections of subplate and deep-layer neocortical neurons. PNAS. 2008;105(41): 16021–16026. doi: 10.1073/pnas.0806791105

88. Sheppard AM, Pearlman AL. Abnormal reorganization of preplate neurons and their associated extracellular matrix: an early manifestation of altered neocortical development in the reeler mutant mouse. J. Comp. Neurol. 1997;378(2): 173–179. doi: 10.1002/(sici)1096-9861(19970210)378:2<173::aid-cne2>3.0.co;2-0

89. Magdaleno S, Keshvara L, Curran T. Rescue of ataxia and preplate splitting by ectopic expression of Reelin in reeler mice. Neuron. 2002;33(4): 573–586. doi: 10.1016/s0896-6273(02)00582-2

90. Park T-J, Curran T. Crk and Crk-Like Play Essential Overlapping Roles Downstream of Disabled-1 in the Reelin Pathway. J. Neurosci. 2008;28(50): 13551-13562. doi: 10.1523/JNEUROSCI.4323-08.2008

91. Kostovic I, Rakic P. Developmental history of the transient subplate zone in the

visual and somatosensory cortex of the macaque monkey and human brain. J. Comp.

Neurol. 1990;297(3): 441–470. doi: 10.1002/cne.902970309

92. Nichols AJ, Olson EC. Reelin promotes neuronal orientation and dendritogenesis

during preplate splitting. Cereb. Cortex. 2010;20(9): 2213–2223. doi:

10.1093/cercor/bhp303

93. Olson EC. Analysis of preplate splitting and early cortical development illuminates

the biology of neurological disease. Front. Pediatr. 2014;2: 1–9. doi:

10.3389/fped.2014.00121

94. Takahashi T, Goto T, Miyama S, Nowakowski RS, Caviness VS Jr. Sequence of

neuron origin and neocortical laminar fate: relation to cell cycle of origin in the

developing murine cerebral wall. J. Neurosci. 1999;19(23): 10357–10371. doi:

10.1523/JNEUROSCI.19-23-10357.1999

95. Bicknese AR, Sheppard AM, O'Leary DD, Pearlman AL. Thalamocortical axons

extend along a chondroitin sulfate proteoglycan-enriched pathway coincident with the

neocortical subplate and distinct from the efferent path. J Neurosci. 1994;14(6): 3500–

3510. doi: 10.1523/JNEUROSCI.14-06-03500.1994

96. del Río JA, Martínez A, Fonseca M, Auladell C, Soriano E. Glutamate-like immunoreactivity and fate of Cajal-Retzius cells in the murine cortex as identified with calretinin antibody. Cereb. Cortex, 1995;5(1): 13–21. doi: 10.1093/cercor/5.1.13

97. Meyer G, Soria JM, Martínez-Galán JR, Martín-Clemente B, Fairén A. Different origins and developmental histories of transient neurons in the marginal zone of the fetal and neonatal rat cortex. J. Comp. Neurol. 1998;397(4): 493–518. doi: 10.1002/(SICI)1096-9861(19980810)397:4<493::AID-CNE4>3.0.CO;2-X

98. Ang Jr. ESBC, Haydar TF, Gluncic V, Rakic P. Four-dimensional migratory coordinates of GABAergic interneurons in the developing mouse cortex. J. Neurosci. 2003;23(13): 5805–5815. doi: 10.1523/JNEUROSCI.23-13-05805.2003

99. Wu SX, Goebbels S, Nakamura K, Nakamura K, Kometani K, Minato N, et al. Pyramidal neurons of upper cortical layers generated by NEX-positive progenitor cells in the subventricular zone. PNAS. 2005;102(47): 17172–17177. doi: 10.1073/pnas.0508560102

100. Goebbels S, Bormuth I, Bode U, Hermanson O, Schwab MH, Nave K. Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice. Genesis. 2006;44(12): 611–621. Doi: 10.1002/dvg.20256

101. Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala, HA, Gu H. A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat. Neurosci. 2006;13(1): 133–140. doi: 10.1038/nn.2467

102. Nowakowski TJ, Pollen AA, Sandoval-Espinosa C, Kriegstein, AR. Transformation of the Radial Glia Scaffold Demarcates Two Stages of Human Cerebral Cortex Development. Neuron. 2016;91(6): 1219–1227. Doi: 10.1016/j.neuron.2016.09.005

103. Shariati, S. A., Lau, P., Hassan, B. A., Muller, U., Dotti, C. G., De Strooper, B., & Gartner, A. (2013). APLP2 regulates neuronal stem cell differentiation during cortical development. Journal of Cell Science, 126(5), 1268-77. Doi: 10.1242/jcs.122440

104. Kielar M, Tuy FPD, Bizzotto S, Lebrand C, Romero CDJ, Poirier K, et al. Mutations in Eml1 lead to ectopic progenitors and neuronal heterotopia in mouse and human. Nat. Neurosci. 2014;17: 923–933. Doi: 10.1038/nn.3729

105. Waite A, Brown SC, Blake DJ. The dystrophin-glycoprotein complex in brain development and disease. Trends Neurosci. 2012;35(8): 487–496. Doi: 10.1016/j.tins.2012.04.004

106. Miller DS, Wright KM. Neuronal Dystroglycan regulates postnatal development of CCK/cannabinoid receptor-1 interneurons. Neural Dev. 2021;16(1): 4. Doi: 10.1186/s13064-021-00153-1

107. Moore CJ, Winder SJ. Dystroglycan versatility in cell adhesion: a tale of multiple motifs. Cell Commun. Signal. 2010;8(3). doi: 10.1186/1478-811X-8-3

108. Batchelor CL, Higginson JR, Chen YJ, Vanni C, Eva A, Winder SJ. Recruitment of Dbl by ezrin and dystroglycan drives membrane proximal Cdc42 activation and filopodia formation. Cell Cycle. 2007; 6(3): 353–363. doi: 10.4161/cc.6.3.3819

109. Ervasti JM, Campbell KP. A role for the dystrophin-glycoprotein complex as a transmembrane linker between Laminin and actin. J. Cell Biol. 1993;122(4): 809-23. doi: 10.1083/jcb.122.4.809

110. James M, Man NT, Wise CJ, Jones, GE, Morris GE. Utrophin-dystroglycan complex in membranes of adherent cultured cells. Cell Motil. 1996;33(3): 163-174. doi: 10.1002/(SICI)1097-0169(1996)33:3<163::AID-CM1>3.0.CO;2-C

111. Spence HJ, Dhillon AS, James M, Winder SJ. Dystroglycan, a scaffold for the ERK-MAP kinase cascade. EMBO Rep. 2004;5(5): 484–489. doi: 10.1038/sj.embor.7400140

112. Satz JS, Philp AR, Nguyen H, Kusano H, Lee J, Turk R, et al. Visual impairment in the absence of dystroglycan. J. Neurosci. 2009;29(42): 13136–13146. doi: 10.1523/JNEUROSCI.0474-09.2009

113. Cohen MW, Jacobson C, Yurchenco PD, Morris GE, Carbonetto, S. Laminin-induced clustering of dystroglycan on embryonic muscle cells: comparison with agrin-induced clustering. J. Cell. Bio. 1997;136(5): 1047–1058. doi: 10.1083/jcb.136.5.1047

114. Durbeej M, Henry MD, Ferletta M, Campbell KP, Ekblom P. Distribution of dystroglycan in normal adult mouse tissues. J. Histochem. Cytochem. 1998;46(4): 449–457. doi: 10.1177/002215549804600404

115. Montanaro F, Gee SH, Jacobson C, Lindenbaum MH, Froehner SC, Carbonetto S. Laminin and alpha-dystroglycan mediate acetylcholine receptor aggregation via a MuSK-independent pathway. J. Neurosci. 1998;18(4): 1250–1260. doi: 10.1523/JNEUROSCI.18-04-01250.1998

116. Henry MD, Cohen MB, Campbell KP. Reduced expression of dystroglycan in breast and prostate cancer. Hum. Pathol. 2001;32(8: 791–795. doi: 10.1053/hupa.2001.26468

117. Colognato H, Yurchenco PD. Form and function: the Laminin family of heterotrimers. Dev. Dyn. 2000;218(2): 213–234. doi: 10.1002/(SICI)1097-0177(200006)218:2<213::AID-DVDY1>3.0.CO;2-R

118. Cabodi S, del Pilar Camacho-Leal M, Di Stefano P, Defilippi P. Integrin signalling

adaptors: not only figurants in the cancer story. Nat. Rev. Cancer. 2010;10(12): 858–

870. doi: 10.1038/nrc2967

119. Malatesta P, Hack MA, Hartfuss E, Kettenmann H, Klinkert W, Kirchhoff F, Götz M.

Neuronal or glial progeny: regional differences in radial glia fate. Neuron. 2003;37(5):

751–764. doi: 10.1016/s0896-6273(03)00116-8.

120. Markó K, Köhidi T, Hádinger N, Jelitai, M, Mezö G, Madarász E. Isolation of radial

glia-like neural stem cells from fetal and adult mouse forebrain via selective adhesion to

a novel adhesive peptide-conjugate. PLoS ONE. 2011;6(12): e28538. doi:

10.1371/journal.pone.0028538

121. Sharma A, Antoku S, Fujiwara K, Mayer BJ. Functional interaction trap: a strategy

for validating the functional consequences of tyrosine phosphorylation of specific

substrates in vivo. Mol. Cell. Proteomics. 2003;2(11): 1217-1224. doi:

10.1074/mcp.M300078-MCP200

122. Sharma A, Mayer BJ. Phosphorylation of p130Cas initiates Rac activation and

membrane ruffling. BMC Cell Biol. 2008;9: 50. doi: 10.1186/1471-2121-9-50

123. Monuki ES, Walsh CA. Mechanisms of cerebral cortical patterning in mice and

humans. Nat. Neurosci. 2001;4: 1199–1206. doi: 10.1038/nn752

124. Rash BG, Grove EA. Area and layer patterning in the developing cerebral cortex. Curr. Opin. Neurobiol. 2006;16(1): 25–34. doi: 10.1016/j.conb.2006.01.004

125. Huang ZJ, Zeng H. Genetic approaches to neural circuits in the mouse. Annu. Rev. Neurosci. 2013;36: 183–215. doi: 10.1146/annurev-neuro-062012-170307

126. Motta A, Berning M, Boergens KM, Staffler B, Beining M, Loomba S, et al. Dense connectomic reconstruction in layer 4 of the somatosensory cortex. Science. 2019;366(6469): eaay3134. doi: 10.1126/science.aay3134

127. Sanes JR, Zipursky SL. Synaptic Specificity, Recognition Molecules, and Assembly of Neural Circuits. Cell. 2020;181(3): 536–556. doi: 10.1016/j.cell.2020.04.008

128. Rakic P. Guidance of neurons migrating to the fetal monkey neocortex. Brain Res. 1971;33(2): 471–476. doi: 10.1016/0006-8993(71)90119-3

129. Campbell K, Götz M. Radial glia: multi-purpose cells for vertebrate brain development. Trends Neurosci. 2002;25(5): 235–238. doi: 10.1016/s0166-2236(02)02156-2

130. Marín O, Valiente M, Ge X, Tsai LH. Guiding neuronal cell migrations. Cold Spring

Harb. Perspect. Biol. 2010;2(2): a001834. doi: 10.1101/cshperspect.a001834

131. Ferent J, Zaidi D, Francis F. Extracellular Control of Radial Glia Proliferation and

Scaffolding During Cortical Development and Pathology. Front. Cell Dev. Biol. 2020;8:

578341. doi: 10.3389/fcell.2020.578341

132. Subramanian L, Calcagnotto ME, Paredes MF. Cortical Malformations: Lessons in

Human Brain Development. Front. Cell. Neurosci. 2020;13: 576. doi:

10.3389/fncel.2019.00576

133. Molnár Z, Métin C, Stoykova A, Tarabykin V, Price DJ, Francis F, et al.

Comparative aspects of cerebral cortical development. Eur. J. Neurosci. 2006;23(4):

921–934. doi: 10.1111/j.1460-9568.2006.04611.x

134. Rakic P. Evolution of the neocortex: a perspective from developmental

biology. Nat. Rev. Neurosci. 2009;10(10): 724–735. doi: 10.1038/nrn2719

135. Cadwell CR, Bhaduri A, Mostajo-Radji MA, Keefe MG, Nowakowski TJ.

Development and Arealization of the Cerebral Cortex. Neuron. 2019;103(6): 980–1004.

doi: 10.1016/j.neuron.2019.07.009

136. Yurchenco PD. Basement membranes: cell scaffoldings and signaling

platforms. Cold Spring Harb. Perspect. Biol. 2011;3(2): a004911. doi:

10.1101/cshperspect.a004911

137. Beggs HE, Schahin-Reed D, Zang K, Goebbels S, Nave, KA, Gorski J, et al. FAK

deficiency in cells contributing to the basal lamina results in cortical abnormalities

resembling congenital muscular dystrophies. Neuron. 2003;40(3): 501–514. doi:

10.1016/s0896-6273(03)00666-4

138. Niewmierzycka A, Mills J, St-Arnaud R, Dedhar S, Reichardt LF. Integrin-linked

kinase deletion from mouse cortex results in cortical lamination defects resembling

cobblestone lissencephaly. J. Neurosci. 2003;25(30): 7022–7031. doi:

10.1523/JNEUROSCI.1695-05.2005

139. Hevner RF, Daza RA, Rubenstein JL, Stunnenberg H, Olavarria JF, Englund C.

Beyond laminar fate: toward a molecular classification of cortical projection/pyramidal

neurons. Dev. Neurosci. 2003;25(2-4): 139–151. doi: 10.1159/000072263

140. Voss AK, Britto JM, Dixon MP, Sheikh BN, Collin C, Tan SS, et al. C3G regulates

cortical neuron migration, preplate splitting and radial glial cell

attachment. Development. 2008;135(12): 2139–2149. doi: 10.1242/dev.016725

141. Gil-Sanz C, Franco SJ, Martinez-Garay I, Espinosa A, Harkins-Perry S, Müller, U. Cajal-Retzius cells instruct neuronal migration by coincidence signaling between secreted and contact-dependent guidance cues. Neuron. 2013; 79(3): 461-477. doi: 10.1016/j.neuron.2013.06.040

142. Nakagawa N, Yagi H, Kato K, Takematsu H, Oka S. Ectopic clustering of Cajal–Retzius and subplate cells is an initial pathological feature in Pomgnt2-knockout mice, a model of dystroglycanopathy. Sci. Rep. 2015;5: 11163. doi: 10.1038/srep11163

143. Booler HS, Pagalday-Vergara V, Williams JL, Hopkinson M, Brown SC. Evidence of early defects in Cajal-Retzius cell localization during brain development in a mouse model of dystroglycanopathy. Neuropathol. Appl. Neurobiol. 2017; 43(4): 330–345. doi: 10.1111/nan.12376

144. Booler HS, Williams JL, Hopkinson M, Brown SC. Degree of Cajal-Retzius Cell Mislocalization Correlates with the Severity of Structural Brain Defects in Mouse Models of Dystroglycanopathy. Brain path. 2016;26(4): 465–478. doi: 10.1111/bpa.12306

145. Radner S, Banos C, Bachay G, Li YN, Hunter DD, Brunken WJ, et al. β2 and γ3 Laminins are critical cortical basement membrane components: ablation of Lamb2 and Lamc3 genes disrupts cortical lamination and produces dysplasia. Dev. Neurobiol. 2013;73(3): 209–229. doi: 10.1002/dneu.22057

146. Zolessi FR, Poggi L, Wilkinson CJ, Chien CB, Harris WA. Polarization and orientation of retinal ganglion cells in vivo. Neural Dev. 2006;1: 2. doi: 10.1186/1749-8104-1-2

147. Edwards MM, Mammadova-Bach E, Alpy F, Klein A, Hicks WL, Roux M, et al. Mutations in Lama1 disrupt retinal vascular development and inner limiting membrane formation. J. Biol. Chem. 2010;285(10): 7697–7711. doi: 10.1074/jbc.M109.069575

148. Pinzón-Duarte G, Daly G, Yong L, Hunter DD, Koch M, Brunken WJ. Defective Formation of the Inner Limiting Membrane in Laminin β2- and γ3-null Mice Produces Retinal Dysplasia. Investig. Ophthalmol. Vis. Sci. 2010;51(3): 1773–1782. doi: 10.1167/iovs.09-4645

149. Sanes JR, Zipursky SL. Design principles of insect and vertebrate visual systems. Neuron. 2010;66(1): 15–36. doi: 10.1016/j.neuron.2010.01.018

150. Randlett O, Poggi L, Zolessi FR, Harris WA. The oriented emergence of axons from retinal ganglion cells is directed by Laminin contact in vivo. Neuron. 2011;70(2): 266–280. doi: 10.1016/j.neuron.2011.03.013

151. Huang Y, Magdaleno S, Hopkins R, Slaughter C, Curran T, Keshvara L. Tyrosine phosphorylated Disabled 1 recruits Crk family adapter proteins. Biochem. Biophys. Res. Commun. 2004;318(1): 204–212. doi: 10.1016/j.bbrc.2004.04.023

152. Olson EC, Walsh CA. Smooth, rough and upside-down neocortical development. Curr. Opin. 2002;12(3): 320–327. doi: 10.1016/s0959-437x(02)00305-2

153. Muntoni F, Voit T. The congenital muscular dystrophies in 2004: a century of exciting progress. Neuromuscul. Disord. 2004;14(10): 635–649. doi: 10.1016/j.nmd.2004.06.009

154. Loturco JJ, Booker AB. Neuronal migration disorders. Cellular Migration and Formation of Axons and Dendrites. 2013: 481-494. doi: 10.1016/B978-0-12-397266-8.00038-7

155. Nickolls AR, Bönnemann CG. The roles of dystroglycan in the nervous system: insights from animal models of muscular dystrophy. Dis. Model Mech. 2018;11(12): dmm035931. doi: 10.1242/dmm.035931

156. Hu H, Yang Y, Eade A, Xiong Y, Qi Y. Breaches of the pial basement membrane and disappearance of the glia limitans during development underlie the cortical lamination defect in the mouse model of muscle-eye-brain disease. J. Comp. Neurol. 2007;501(1): 168–183. doi: 10.1002/cne.21238

157. Manzini MC, Gleason D, Chang BS, Hill RS, Barry BJ, Partlow JN, et al. Ethnically diverse causes of Walker-Warburg syndrome (WWS): FCMD mutations are a more

common cause of WWS outside of the Middle East. Hum. Mutat. 2008;29(11): E231–E241. doi: 10.1002/humu.20844

158. Vuillaumier-Barrot S, Bouchet-Séraphin C, Chelbi M, Devisme L, Quentin S, Gazal S, et al. Identification of mutations in TMEM5 and ISPD as a cause of severe cobblestone lissencephaly. Am. J. Hum. Genet. 2012;91(6): 1135–1143. doi: 10.1016/j.ajhg.2012.10.009

159. Godfrey C, Clement E, Mein R, Brockington M, Smith J, Talim B, et al. Refining genotype phenotype correlations in muscular dystrophies with defective glycosylation of dystroglycan. Brain. 2007;130(10): 2725–2735. doi: 10.1093/brain/awm212

160. Mercuri E, Messina S, Bruno C, Mora M, Pegoraro E, Comi GP, et al. Congenital muscular dystrophies with defective glycosylation of dystroglycan: a population study. Neurol. 2009;72(21): 1802–1809. doi: 10.1212/01.wnl.0000346518.68110.60

161. Chung C, Yang X, Bae T, Vong KL, Mittal S, Donkels, C, et al. Comprehensive multi-omic profiling of somatic mutations in malformations of cortical development. Nat. Genet. 2023;55: 209–220. doi: 10.1038/s41588-022-01276-9

162. Iffland II PH, Crino PB. Focal Cortical Dysplasia: Gene Mutations, Cell Signaling, and Therapeutic Implications. Annu. Rev. Pathol. 2017:12(1): 547-571. doi: 10.1146/annurev-pathol-052016-100138

163. Polleux F, Ghosh A. The Slice Overlay Assay: A Versatile Tool to Study the

Influence of Extracellular Signals on Neuronal Development. Sci. Signal.

2002;2002(136): pl9. doi: 10.1126/stke.2002.136.pl9

164. Kim HJ, Magrané J. Isolation and culture of neurons and astrocytes from the

mouse brain cortex. Methods Mol. Biol. 2011;793: 63–75. doi: 10.1007/978-1-61779-

328-8_4

165. Culican SM, Baumrind NL, Yamamoto M, Pearlman AL. Cortical radial glia:

identification in tissue culture and evidence for their transformation to astrocytes. J.

Neurosci. 1990;10(2): 684–692. doi: 10.1523/JNEUROSCI.10-02-00684.1990

166. Lindenmaier LB, Parmentier N, Guo C, Tissir F, Wright KM. Dystroglycan is a

scaffold for extracellular axon guidance decisions. ELife. 2019;8: 1–26. doi:

10.7554/eLife.42143

167. Sekar A, Sanches TM, Hino K, Kumar M, Wang J, Ha, E, et al. RapID Cell

Counter□: Semi-automated and mid- throughput estimation of cell density within diverse

cortical layers. Eneuro. 2021;8(6). doi: 10.1523/ENEURO.0185-21.2021

168. Li S, Jin Z, Koirala S, Bu L, Xu L, Hynes RO, Walsh CA, Corfas G, Piao X. GPR56 regulates pial basement membrane integrity and cortical lamination. J Neurosci. 2008; 28(22):5817-26. doi: 10.1523/JNEUROSCI.0853-08.2008.

169. D'Arcy BR, Lennox AL, Manso Musso C, Bracher A, Escobar-Tomlienovich C, Perez-Sanchez S, Silver DL. Non-muscle myosins control radial glial basal endfeet to mediate interneuron organization. PLoS Biol. 2023;21(2):e3001926. doi: 10.1371/journal.pbio.3001926

170. Ubina T, Vahedi-Hunter T, Agnew-Svoboda W, Wong W, Gupta A, Santhakumar V, Riccomagno MM. ExBoX – a simple Boolean exclusion strategy to drive expression in neurons. J Cell Sci 2021; 134 (20): jcs257212. doi: 10.1242/jcs.257212

171. Agnew-Svoboda W, Ubina T, Figueroa Z, Wong YC, Vizcarra EA, Roebini B, Wilson EH, Fiacco TA, Riccomagno MM. A genetic tool for the longitudinal study of a subset of post-inflammatory reactive astrocytes. Cell Rep Methods 2022; 2(8):100276. doi: 10.1016/j.crmeth.2022.100276.

Supplemental Figure Legends

S1 Fig. EGFP antibodies are specific for EGFP. (A-B) Immunostaining of cortical

sections with the same chicken-anti EGFP (**A**) and rabbit-anti EGFP (**B**) antibodies

used to detect it in *p130Cas::BacEGFP* animals produce negligible signal on WT

sections at E12.5 and E15.5. Magenta channel in (**A**) is Tbr1 and in (**B**) is Nestin. Dapi

counterstain is shown in blue. n=3 animals. PP: preplate, CP: cortical plate; IZ:

intermediate zone; SVZ: subventricular zone; VZ: ventricular zone. Scale bars for A and

B (top): 25 µm; A and B (bottom): 75 µm.

S2 Fig. *Emx1Cre* efficiently recombines in the developing cortex. (**A**) tdTomato

staining of E13.5 *Emx1Cre;Ai14* coronal cortical sections. *Emx1Cre* drives Cre

recombinase expression in cortical neural progenitors and radial glial cells that generate

excitatory neurons in the forebrain and hippocampus. (**B**) RNAscope at E14.5 shows

that *Emx1Cre;CasTcKO* cortices do not produce functional *p130Cas* transcripts. Control

littermates express *p130Cas* transcripts strongly in the IZ and broadly throughout the

neocortex. Neither controls, nor *Emx1Cre;CasTcKO* animals express *Sin* or *CasL.* n=3

animals per genotype. CP: cortical plate; IZ: intermediate zone; SVZ: subventricular

zone; VZ: ventricular zone. Scale bars for A: 500 µm; D: 100 µm.

S3 Fig. Characterization of cortical phenotypes in *Emx1Cre;CasTcKO* mice. (**A-C**)

Immunostaining of P7 control and *Emx1Cre;CasTcKO* coronal sections using the layer

markers Cux1 (green, layer II/III) and Ctip2 (magenta, layer V & VI) (**A**), Rorβ+ (**B**, red,

layer IV) and Tbr1+ (**C**, red, layer VI). Neurons appear misplaced in *Emx1Cre;CasTcKO*

mice (white arrows). DAPI (blue) was used to counterstain. (**A**) Right panels:

quantification of Ctip+ and Cux1+ neuron density. two tailed t-test *ns*: p=0.12 for Ctip2+

cells; p=0.09 for Cux1+ cells; n=5 independent animals for each genotype, 3 section per

animal. (**B, C**) Right panels: quantification of the proportion of Rorβ+ and Tbr1+ cells

per bin. Bin 1 is the MZ and bin 10 is ventral to layer VI. n=3 animals per group, 3

sections per animal, Mann-Whitney *U* test & Bonferroni correction, *p<0.005. Values

given are Mean ± S.E.M. For data plotted in graphs, see S2 Data. Scale bars: 500 μm.

S4 Fig. Analysis of for *CasL, Sin* and *p130Cas* null allelic series. Coronal sections

of P7 *Emx1Cre;p130Cas$^{flox/+}$;CasL$^{+/-}$;Sin$^{+/-}$* , *Emx1Cre; p130Cas$^{flox/+}$;CasL$^{-/-}$;Sin$^{-/-}$* ,

Emx1Cre; p130Cas$^{flox/-}$;CasL$^{+/-}$;Sin$^{-/-}$, *Emx1Cre; p130Cas$^{flox/-}$;CasL$^{-/-}$;Sin$^{+/-}$* and

Emx1Cre; p130Cas$^{flox/-}$;CasL$^{-/-}$;Sin$^{-/-}$ (*Emx1Cre;CasTcKO*) cortices stained for the layer

markers Ctip2 (green, layer V & VI) and Tbr1 (red, Layer VI), and the mature neuron

marker NeuN (grey), counterstained with DAPI (blue). Only *Emx1Cre; p130Cas$^{flox/-}$*

;CasL$^{-/-}$;Sin$^{+/-}$ and *Emx1Cre; p130Cas$^{flox/-}$;CasL$^{-/-}$;Sin$^{-/-}$* showed the cobblestone cortex

phenotype. Bottom Right panel: quantification of the # of dysplasia/mm^2 in the different

transgenic animals. Values given are Mean ± S.E.M., n=3-8 independent samples per

group, 3 sections per sample, Mann-Whitney *U* test & Bonferroni correction, *p<0.005

vs. control; ** p<0.0005 vs. control. For data plotted in graphs, see S3 Data. Scale bar:

1 mm.

S5 Fig. Defective preplate split in *Emx1Cre;CasTcKO* mice. Immunostaining of

control and *Emx1Cre;CasTcKO* coronal sections for marginal zone (MZ) and subplate

(SP; white arrows) markers at E15.5. MAP2, CSPG, Calretinin, and Calbindin show

clustering of cells (yellow arrows). CSPG+ cells form a column connecting the SP with

the MZ (yellow arrow). Thalamocortical afferents labeled with Calretinin show ectopic

bundles extending from the intermediate zone (IZ) (white arrowheads). DAPI was used

as counterstain (blue). n=3 animals per genotype. Scale bars: 100 μm.

S6 Fig. Caspase3 activation is not affected in *Emx1Cre;CasTcKO* mice at key

developmental stages. (A-D) Immunostaining for activated Caspase3 (aCasp3, green)

on E12.5 (**A**), E15.5 (**B**), P0 (**C**) or P3 (**D**) coronal sections of Control and

Emx1Cre;CasTcKO cortices. Sections were counterstained with DAPI (blue). No

notable differences are observed between genotypes. Right panels: quantification of

density of activated Caspase 3. Values given are Mean ± S.E.M., n=3 independent

samples per group, 3-5 sections per sample, Mann-Whitney *U* test & Bonferroni

correction, ns for all stages (p≥0.7). For data plotted in graphs, see S6 Data. Scale bars

for A: 100 μm; B, C, D:500 μm.

S7 Fig. *NexCre* efficiently ablates *p130Cas^flox* in postmitotic cortical neurons. (**A**)

tdTomato staining of E13.5 *NexCre;Ai14* coronal cortical sections. *NexCre* targets

postmitotic, premigratory neurons in the neocortex. (**B**) RNAscope at E16.5 indicates

that *NexCre;CasTcKO* cortices express negligible levels of functional *p130Cas*

transcripts in the IZ and CP, and display significantly less transcript in the VZ and SVZ.

Top Right panel: quantification puncta per micron DAPI area in cortical plate +

intermediate zone (CP+IZ) was 0.114±0.009 for *Nex-Cre⁺;CasTcKO^flox/+* and

0.116±0.005 for ventricular + subventricular zones (VZ+SVZ). For *Nex-*

Cre⁺;CasTcKO^{flox/Δ} the average *p130Cas* puncta per micron DAPI area in CP+IZ was 0.013±0.003 and 0.056±001 for VZ and SVZ. Values given are mean ±SEM, n=5-6, ***p<0.0001 vs. Control (One-way ANOVA; Tukey post-hoc test). For data plotted in graphs, see S7 Data. Scale bars: 100 μm.

S8 Fig. Characterization of cortical phenotypes in *NexCre;CasTcKO* mice. (A-B) Immunostaining of coronal sections for P7 control and *NexCre;CasTcKO* animals with layer makers Rorβ (**A**, red, layer IV) and Tbr1 (**B**, red, layer VI). DAPI was used to counterstain nuclei (blue). No obvert lamination phenotypes are observed. Right panels: quantification of proportion of Rorβ+ (**A**) and Tbr1+ (**B**) cells per bin. Values given are Mean ± S.E.M., n=3 animals per group, 3 sections per animal, Mann-Whitney *U* test & Bonferroni correction, no significant differences (p>0.05). For data plotted in graphs, see S9 Data. Scale bar: 500 μm.

S9 Fig. Progenitor phenotype in E12.5 *Emx1Cre;CasTcKO* cortices. (A-B) Detection of proliferative cells in Control and *Emx1Cre;CasTcKO* coronal sections using short-pulsed EdU at E12.5 (**A,B**, green), colabeled with the intermediate progenitor marker Tbr2 (**B,** magenta). Nuclei were counterstained with DAPI (blue). (**C**) Density of EdU+ cells/mm^2 of cortex. p=0.34, two-tailed Mann-Whitney *U* test, n=4 independent samples per group, 3 sections per sample. (**D**) Proportion of Edu+ cells that co-express Tbr2. p=0.49, two-tailed Mann-Whitney *U* test, n=4 independent samples per group, 2-3 sections per sample. Values given are Mean ± S.E.M. For data plotted in graphs, see

S11 Data. Scale bars: Lower magnification panel for A: 500 µm; Higher mag panels in

A: 100 µm; B: 75 µm.

S10 Fig. No lamination phenotypes are observed in *NexCre;β1-Integrin*^{loxp/loxp}

animals. (**A-C**) Immunostaining for cortical layer markers on P7 cortical sections from

Control and *NexCre;β1-Integrin*^{loxp/loxp} animals. Staining for Cux1 (**A**, green, layers II-III)

and Ctip2 (**A**, magenta, layers V and VI), or Tbr1 (**B**, red, layer VI), or Rorβ (**C,** red,

layer IV), are shown. Sections were counterstained with DAPI (blue). n=3 animals per

genotype for each marker. Scale bars: 500 µm.

S11 Fig. Some cells in mixed primary cortical cultures express RGC markers.

Example of E13.5 4DIV (four days in vitro) mixed primary cortical cultures after staining

for neural markers to identify distinct cell types. Left panel: Nestin (green,

neuroepithelial and radial glial cells), GFAP (magenta, astrocytes); Middle panel: Blbp

(red, radial glial cells); Right panel: Tuj1 (green, postmitotic neurons), Tbr2 (magenta,

intermediate progenitors). DAPI was used to counterstain nuclei (blue). n=3

independent cultures. Scale bar: 25 µm.

S12 Fig. Cortical phenotypes on the contralateral hemispheres to the *in utero*

electroporated lateral ventricles. Immunostaining of P0 control and

Emx1Cre;CasTcKO coronal sections of the contralateral hemispheres to the sides co-

electroporated with ∆Src + p130Cas-ZipB (negative control) or Src-ZipA + p130Cas-

ZipB (FIT rescue). Sections were stained for the layer V & VI marker Ctip2 (green) and

laminin (magenta), and counterstained with DAPI (blue). In utero electroporation of one side does not affect the expected phenotype on the opposite hemisphere: i.e. cobblestone phenotypes are still observed in the contralateral side of ΔSrc + p130Cas-ZipB or Src-ZipA + p130Cas-ZipB electroporated *Emx1Cre;CasTcKO* cortices. n=6 animals per genotype. Scale bar: 250 μm.

S1 Data. Numerical values for graphs in Fig 3 including Fig 3D and Fig 3E

S2 Data. Numerical values for graphs in S3 Fig including S3A Fig, S3B Fig and S3C Fig.

S3 Data. Numerical values for graph in S4 Fig.

S4 Data. Numerical values for graphs in Fig 4 including Fig 4A, Fig 4B, Fig 4C and Fig 4D.

S5 Data. Numerical values for graph in Fig 5.

S6 Data. Numerical values for graphs in S6 Fig.

S7 Data. Numerical values for graph in S7 Fig.

S8 Data. Numerical values for graphs in Fig 6 including Fig 6A, Fig 6B and Fig 6C.

S9 Data. Numerical values for graphs in S8 Fig including S8A Fig and S8B Fig.

S10 Data. Numerical values for graphs in Fig 9 including Fig 9C, Fig 9D and Fig 9E.

S11 Data. Numerical values for graphs in S9 Fig including S9C Fig and S9D Fig

S12 Data. Numerical values for graph in Fig 10.

S13 Data. Numerical values for graphs in Fig 12.

S14 Data. Numerical values for graphs in Fig 13.

A *p130Cas::BacEGFP*

EGFP Tbr1 DAPI | EGFP

PP
SVZ
VZ
E12.5

B
E15.5

C
CP
IZ
SVZ
VZ
E15.5

A Control   *Emx1Cre;CasTcKO*

FoxP2 L1CAM DAPI
P0

B Tbr1 DAPI
E12.5

C Tbr1 DAPI
E15.5

D Control   *Emx1Cre;CasTcKO*

Tbr1 DAPI
CP
IZ
SVZ
VZ
E15.5

E Laminin Ctip2 DAPI
CP
IZ
SVZ
VZ
E15.5

F

Tbr1+ ectopias/ mm cortical surface

*

Control   *Emx1Cre;TcKO*

A

Control   *Emx1Cre;CasTcKO*

Laminin   Nestin

MZ

CP

VZ

E15.5

MZ

CP

VZ

B

EdU   Tbr2   EdU   Tbr2
DAPI

EdU e15.5 0.5 Hrs.

Control

MZ

CP

VZ

Emx1Cre;CasTcKO

MZ

CP

VZ

E15.5

C

EdU+ cells in CP/mm²

Control   *Emx1Cre;TcKO*

D

EdU+ cells/mm²

Control   *Emx1Cre;TcKO*

E

Tbr2+, EdU+ cells/EdU+ cells

Control   *Emx1Cre;TcKO*

A

Zip A — Src Kinase Domain

Zip B — p130Cas

Zip A — Src Kinase Domain
Zip B — p130Cas — P P

B

Ctip2 EGFP
Laminin DAPI | Ctip2 | EGFP | Laminin

Control

ΔSrc + p130Cas-ZipB

Src-ZipA + p130Cas-ZipB

Emx1Cre;β1^{flox/flox}

ΔSrc + p130Cas-ZipB

Src-ZipA + p130Cas-ZipB

C

# of dysplasia/IUE area

**

- Control + Src+p130ZipB
- Control + SrcZipA+p130ZipB
- *Emx1Cre;B1^{f/f}* +Src+p130ZipB
- *Emx1Cre;B1^{f/f}* + SrcZipA+p130ZipB

Contingency tables for cortical dysplasia penetrance

Emx1Cre; Cas TcKO

	Control	*Emx1Cre; Cas TcKO*
No Dysplasia	52	0
Dysplasia	0	48

Fisher Exact test
The two-tailed P value is less than 0.0001

NexCre; Cas TcKO

	Control	*NexCre; Cas TcKO*
No Dysplasia	21	17
Dysplasia	0	0

The two-tailed P value equals 1.0000

Emx1Cre; B1$^{flox/flox}$

	Control	*Emx1Cre; B1*$^{flox/flox}$
No Dysplasia	20	0
Dysplasia	0	17

The two-tailed P value is less than 0.0001

NexCre; B1$^{flox/flox}$

	Control	*NexCre; B1*$^{flox/flox}$
No Dysplasia	10	8
Dysplasia	0	0

The two-tailed P value equals 1.0000

Emx1Cre; Dag$^{flox/-}$

	Control	*Emx1Cre; Dag*$^{flox/-}$
No Dysplasia	18	0
Dysplasia	0	18

The two-tailed P value is less than 0.0001

Dag$^{Cyto/-}$

	Control	*Dag*$^{Cyto/-}$
No Dysplasia	10	10
Dysplasia	0	0

The two-tailed P value equals 1.0000