SNP discovery by exome capture and resequencing in a pea genetic resource collection

G. Aubert[1], J. Kreplak[1], M. Leveugle[2,8], H. Duborjal[2,8], A. Klein[1], K. Boucherot[1], E. Vieille[1], M. Chabert-Martinello[1], C. Cruaud[3], V. Bourion[1], I. Lejeune-Hénaut[4], M.L. Pilet-Nayel[5], N. Francillonne[6,7], N. Tayeh[1], J.P. Pichon[2,8], N. Rivière[2,8], J. Burstin[1]

[1] Agroécologie, INRAE, Institut Agro, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France

[2] Biogemma, F-63720 Chappes, France

[3] Genoscope, Institut François Jacob, CEA, Université Paris-Saclay, 2 Rue Gaston Crémieux, 91057 Evry, France

[4] BioEcoAgro, INRAE, Univ. Liège, Univ. Lille, Univ. Picardie Jules Verne, 2, Chaussée Brunehaut, F-80203, Estrées-Mons, France

[5] IGEPP, INRAE, Institut Agro, Univ Rennes, 35653, Le Rheu, France

[6] Université Paris-Saclay, INRAE, URGI, 78026, Versailles, France

[7] Université Paris-Saclay, INRAE, BioinfOmics, Plant bioinformatics facility, 78026, Versailles, France

[8] Present address: Limagrain F-63720 Chappes, France

Background & Summary

In addition to being the model plant used by Mendel[1] to establish genetic laws, pea (*Pisum sativum* L.) is a major pulse crop cultivated in many temperate regions of the world. In order to face new challenges imposed particularly by global climate change and new regulations targeted at reducing chemical inputs, pea breeders have to take advantage of the genetic diversity present in the *Pisum* genepool to develop improved, resilient varieties. The aim of this study was to assess the genetic diversity of a pea germplasm collection and allow genome-wide association studies using this collection.

To be able to perform genome-wide association approaches with high resolution, genotyping with a large set of genetic markers such as Single Nucleotide Polymorphism (SNP) markers well-spread over the genome is required. Rapid advances in second-generation sequencing technologies and the development of bioinformatic tools have revolutionized the access to and the characterization of available genetic diversity. High-density, high-throughput genotyping has been possible for a large number of species, including those with large and complex genomes[2] such as pea which genome size is estimated to be 4.45 Gb[3]. In this study, which is part of the PeaMUST project[4], we used a target capture technology based on pea transcriptome sequences to generate exome-enriched genomic libraries that were further subjected to Illumina sequencing in paired-end mode. This methodology was chosen because whole-genome resequencing is relatively expensive for species with large genomes and because capturing genetic variations in repeated non-coding regions is difficult to achieve or to interpret[5]. Whole-exome sequencing represented an interesting alternative that focused on coding regions only[6,7]. Mapping the obtained reads on the reference pea genome sequence enabled the discovery of an abundant set of SNPs. The development of this resource is a crucial cornerstone in research and breeding projects towards boosting the improvement of pea production and quality.

Methods
Plant material and DNA extraction

A set of 240 *Pisum* accessions was selected, including 220 accessions originated from a larger panel structured into 16 genetic groups[8] and 20 additional accessions chosen for their phenotypes. The 240-accession collection is referred to as Architecture and Multi-Stress (AMS) collection, since the accessions represent a broad diversity for root and shoot architecture and for biotic and abiotic stress responses. This collection contains cultivars, landraces, and wild types (including some *Pisum fulvum* and *Pisum sativum* subspecies accessions) with diverse geographical origins (Supplementary Table 1). Leaves were collected from 10 plants per accession, flash frozen in liquid nitrogen and stored at -80°C prior to DNA extraction. Tissues were then ground in liquid nitrogen using a pestle and a mortar. Genomic DNA extraction was performed using Nucleospin PlantII minikit (Macherey-Nagel, Hoerdt, France) following the manufacturer's instructions.

Probe design

As the pea genome sequence was not available at the time the probe design was made, two pea transcriptome datasets[9,10] were used to build a reference set (refset) of expressed genes. After redundancy was removed, 67,161 unique contigs were kept, 20,972 of them being common to the two sequence datasets. The first exome capture design based on the refset was undertaken after predicting putative exon/intron junctions, masking repetitive sequences as well as excluding putative mitochondrial and chloroplastic sequences. A first probe design was performed by Roche[Tm] (Madison, WI, USA) targeting 68.3 Mb. The analysis of the first capture results with the original probes demonstrated that a minority of 10-15% of the contigs retained the majority of the sequencing efforts, resulting in insufficient coverage for the remaining contigs. Sequencing data were used to identify the repetitive regions and a new probe design was performed after masking them to target a final capture space of 41.3Mb, representing 51,225 cDNA contigs.

Library preparation and target enrichment

DNA samples were normalized before being fragmented with Adaptive Focused Acoustics® Technology (Covaris Inc., Massachusetts, USA). A 250-bp target size was obtained by using a Covaris E220 system, according to the manufacturer's instructions. Then DNA fragments underwent a NGS library preparation procedure consisting in end repair and Illumina adaptor ligation using the KAPA HTP kit (Roche, Basel, Switzerland). Individual index sequences were added to each library for identifying reads and sorting them according to their initial origins. The Sequence Capture was performed using SeqCap EZ Developer kit[11] according to the manufacturer's instructions (Roche[Tm]). The sequence capture reaction efficiency was evaluated by measuring, using quantitative PCR, a relative fold enrichment and loss of respectively targeted and non-targeted regions before and after the sequence capture reaction.

Targeted resequencing, sequence alignment and SNP calling

The captured samples were sequenced on HiSeq 2000 sequencing platform (Illumina, California, USA) with a Paired-End sequencing strategy of 2 reads of 100 bases. The sequenced reads were trimmed for adaptor sequences using cutadapt 1.8.3[12]. Low-quality nucleotides with quality value<20 were removed from both ends. The longest sequence without adapters and low-quality bases was kept. Sequences between the second unknown nucleotide (N) and the end of the read were also trimmed. Reads shorter than 30 nucleotides after trimming were discarded. These trimming steps were achieved using fastx_clean (http://www.genoscope.cns.fr/fastxtend), an internal software based on the FASTX library (http://hannonlab.cshl.edu/fastx_toolkit/index.html). The reads were then aligned on the targeted regions with Novoalign V3.09(http://www.novocraft.com, Selangor, Malaysia). We took advantage of the v1 genome sequence of cv. Cameor published meanwhile[13] to perform SNP detection. Single nucleotide variants were detected on all the samples using samtools mpileup, followed by bcftools call and bcftools filter[14] with a minimum genotype quality of 20 and a minimum coverage of 5 reads per sample. SNP variants for which more than 10 percent of the samples in the panel were heterozygous were then filtered out.

Phylogenetic analysis

A subset of 206,474 SNPs was selected from the Dataset by applying filters on missing data (<20%) and linkage disequilibrium (LD pruning based PLINK). This subset was used to build a phylogenetic tree. A maximum likelihood phylogenetic tree was calculated using IQtree[15], version 2.1.2, with model of substitution GTR+F+ASC. Tree visualizations were generated using R package ggtree[16].

Structure analysis

Structure within the collection was calculated using the Bayesian clustering program FastStructure[17] using a logistic prior for K ranging from 1 to 10. The script chooseK.py (part of the FastStructure distribution) was used to determine the best K that explained the structure in the collection based on model complexity. In addition, discriminant analysis of principal components (DAPC) was applied using DAPC function from adegenet package (version 2.1.5)[18] in order to describe the genetic structure of the panel. Both FastStructure and DAPC groups were visualized on the phylogenetic tree using R package ggtreeExtra[16].

Data Records

The collection of 240 pea accessions was genotyped using an original set of probes for exome capture. Sequence data are available on NCBI (Bioproject PRJEB56612). Using the pea Cameor genomic sequence v1[13] as a reference, 2,285,342 SNPs were identified. The full set of variants has been recorded as a VCF file and deposited at URGI (https://doi.org/10.15454/3QRIPA). The variant statistics per accession are reported in Supplementary Table 2. In average, 183,170 homozygous variants (compared to Cameor genome sequence) were detected per accession. As expected, the genotyping data from Cameor (DCG0251) used as an accession in the AMS panel were overall conform to the reference alleles from the reference genome sequence. Differences were only seen for 121 positions (0,005%). Among the detected SNPs, 647,220 were singletons (detected in only one accession). Excluding the reference Cameor (DCG0251), the number of singletons per accession ranged between 2 for VKL0176, a cultivated winter fodder pea to 113,086 for VSD0034, a *Pisum sativum* subsp. *abyssinicum* accession. The set of identified polymorphisms spans the seven pseudomolecules of the Cameor genome assembly at a frequency varying from 1 variant every 1831 bases for chromosome 2 to one every 1345 bases for chromosome 1 (Supplementary Table 3).

Variant effects.

We used the snpEff program[19] in order to categorize the detected SNPs according to their predicted effects or their locations (Supplementary Table 3). The vast majority (76,71%) were labelled as "Modifier" (placed upstream or downstream of genes) and 14.32%, 8.76%, and 0.22% of the SNPs had a low, moderate, and high predicted impact on gene functions, respectively. In fact, out of the total SNPs detected in coding regions (23.29%), 57.52% were predicted to be silent, 41.95% were predicted to induce an amino acid change in the coded protein and 0.53% were predicted to have a disruptive nonsense effect (premature stop codon, splicing junction modification or start codon missing).

Our data provide insights into exome genetic variation and highlight mutations with functional effects. This polymorphism inventory is valuable to explain the phenotypic diversity in the *Pisum* species.

Phylogeny and structure analysis.

Three different complementary approaches (Discriminant analysis of principal components (DAPC), FastSTRUCTURE, Neighbor-Joining Phylogenetic trees) have been used to study the structure of the germplasm panel.

DAPC led to a clustering of the accessions in seven groups, as the most likely structure according to Bayesian information criterion. Clustering (Table 1, Figure 1) tended to separate accessions according to crop evolution and cultivation types. Cluster 1 consisted in 14 accessions, mainly landraces or primitive germplasm. Cluster 2 comprised 87 accessions with an important proportion of spring cultivars including garden peas. Cluster 3 is composed of 27 accessions with a majority of spring-type lines coming from an Aphanomyces root rot breeding programme from Groupement des Sélectionneurs de Protéagineux (GSP, France) and Cluster 4 was a mix of spring-type and winter-type field peas with three primitive or landrace accessions. Cluster 5 grouped 5 wild-type accessions including *Pisum fulvum*, and Clusters 6 and 7 included winter-type field pea and fodder pea cultivars, respectively.

FastStructure, on the other hand, inferred that the panel was divided into five ancestral subpopulations (numbered A to E, Figure 1). Subpopulation E corresponded to the DAPC cluster of 5 wild accessions with very little admixture with the other clusters while subpopulation C corresponded to Cluster 1 (landraces/primitive germplasm). Subpopulation D corresponded to the fodder pea cluster 7. Clusters 3,4 and 6 (field pea cultivars mainly) seemed to derive from the ancestral subpopulation A. Some admixture between subpopulations A and D was observed for the winter field pea accessions from Cluster 6, and between subpopulations A, B and D for some garden pea cultivars.

The Phylogenetic tree confirmed the DAPC and FastStructure observations with only few placement differences that could be due to kinship.

Conclusion

In conclusion, this dataset is a large marker resource that can be used for different purposes, including the development of targeted genotyping tools for molecular identification, genetic mapping or genomic selection in pea. It provides insights into pea diversity and helps to investigate selection processes in this species. The SNP resource also empowers Genome-wide Association Studies targeted at revealing the genetic architecture of important traits and highlighting alleles to be used in pea breeding programmes. Indeed, the collection has been evaluated for different traits including plant architecture, phenology and resistance or tolerance to a range of biotic and abiotic stresses, as exemplified by Ollivier et al[20] who deciphered the genetic determinism of resistance to two aphid biotypes.

Acknowledgements

This study is part of the PeaMUST project, that was funded by the French government through the Investment for the Future program (project ANR-11-BTBR-0002) and Ministère de l'Enseignement supérieur, de la Recherche et de l'Innovation.
The authors wish to thank the reviewers and the recommender for very useful comments on the manuscript.

Author Contributions

ILH, MLPN, VB and JB designed the diversity panel. Experiments were conceived and designed by GA, NR, JPP and JB, and performed by HD, AK, KB, EV, MCM and CC. Data were analysed by ML, JK and GA and NF organised the data access. The manuscript was written by GA and JK and all authors read and improved it.

Competing interests

The authors declare they have no conflict of interest relating to the content of this article

References

1. Mendel, G. Versuche über Pflanzenhybriden. in *Versuche über Pflanzenhybriden* (1866). doi:10.1007/978-3-663-19714-0_4.

2. Hill, C. B. *et al.* Targeted enrichment by solution-based hybrid capture to identify genetic sequence variants in barley. *Sci. Data* **6**, (2019).

3. Doležel, J. *et al.* Plant genome size estimation by flow cytometry: Inter-laboratory comparison. *Ann. Bot.* **82**, (1998).

4. Burstin, J. *et al.* PeaMUST (Pea MultiStress Tolerance), a multidisciplinary French project uniting researchers, plant breeders, and the food industry. *Legume Sci.* **3**, (2021).

5. Ku, C. S. *et al.* Exome versus transcriptome sequencing in identifying coding region variants. *Expert Rev. Mol. Diagn.* **12**, (2012).

6. Ng, S. B. *et al.* Targeted capture and massively parallel sequencing of 12 human exomes. *Nature* **461**, (2009).

7. Ng, S. B. *et al.* Exome sequencing identifies the cause of a mendelian disorder. *Nat. Genet.* **42**, (2010).

8. Siol, M. *et al.* Patterns of genetic structure and linkage disequilibrium in a large collection of pea germplasm. *G3 Genes Genomes Genet.* **7**, (2017).

9. Duarte, J. *et al.* Transcriptome sequencing for high throughput SNP development and genetic mapping in Pea. *BMC Genomics* **15**, (2014).

10. Alves-Carvalho, S. *et al.* Full-length *de novo* assembly of RNA-seq data in pea (*Pisum sativum* L.) provides a gene expression atlas and gives insights into root nodulation in this species. *Plant J.* **84**, 1–19 (2015).

11. Bainbridge, M. N. *et al.* Whole exome capture in solution with 3 Gbp of data. *Genome Biol.* **11**, (2010).

12. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, (2011).

13. Kreplak, J. *et al.* A reference genome for pea provides insight into legume genome evolution. *Nat. Genet.* **51**, (2019).

14. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

15. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

16. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

17. Raj, A., Stephens, M. & Pritchard, J. K. FastSTRUCTURE: Variational inference of population structure in large SNP data sets. *Genetics* **197**, (2014).

18. Jombart, T., Devillard, S. & Balloux, F. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genet.* **11**, 94 (2010).

19. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* **6**, (2012).

20. Ollivier, R. *et al.* A major-effect genetic locus, ApRVII, controlling resistance against both adapted and non-adapted aphid biotypes in pea. *Theor. Appl. Genet.* (2022) doi:10.1007/s00122-022-04050-x.

Figure1: Phylogenetic analysis, DAPC grouping, and FastStructure composition of the 240-accession pea AMS collection. Colour codes indicate the different groups as inferred by DAPC and FastStructure.