Faster detection of "darks" than "brights" by monkey superior colliculus neurons

Tatiana Malevich[1, 2*], Tong Zhang[1, 2*], Matthias P. Baumann[1, 2*], Amarender R. Bogadhi[1, 2, 3], and Ziad M. Hafed[1, 2]

1. Werner Reichardt Center for Integrative Neuroscience,
University of Tübingen, Tübingen, Germany 72076
2. Hertie Institute for Clinical Brain Research,
University of Tübingen, Tübingen, Germany 72076
3. Central Nervous System Diseases Research, Boehringer Ingelheim Pharma GmbH & Co. KG,
Biberach, Riß, Germany 88400

* Co-first authors

Abbreviated title:

Visual processing of darks by superior colliculus

Correspondence to:

Tatiana Malevich (tatiana.malevich@cin.uni-tuebingen.de)
and
Ziad M. Hafed (ziad.m.hafed@cin.uni-tuebingen.de)
Werner Reichardt Center for Integrative Neuroscience
Otfried-Müller Str. 25
Tübingen, Germany, 72076

Number of pages: 53

Number of figures: 11

Number of words in Abstract: 247

Number of words in Introduction: 661

Number of words in Discussion: 1434

Conflict of interest statement: The authors declare no competing interests.

Acknowledgements: We were funded by the Deutsche Forschungsgemeinschaft (DFG): (1) BO5681/1-1; (2) EXC307; (3) HA6749/3-1; (4) HA6749/4-1; and (5) SFB 1233, Robust Vision: Inference Principles and Neural Mechanisms, TP 11, project number: 276693517.

Abstract

Visual processing is segregated into ON and OFF channels as early as in the retina, and the superficial (output) layers of the primary visual cortex are dominated by neurons preferring dark stimuli. However, it is not clear how the timing of neural processing differs between "darks" and "brights" in general, especially in light of psychophysical evidence; it is also equally not clear how subcortical visual pathways that are critical for active orienting represent stimuli of positive (luminance increments) and negative (luminance decrements) contrast polarity. Here, we recorded from all visually-responsive neuron types in the superior colliculus (SC) of two male rhesus macaque monkeys. We presented a disc (0.51 deg radius) within the response fields (RF's) of neurons, and we varied, across trials, stimulus Weber contrast relative to a gray background. We also varied contrast polarity. There was a large diversity of preferences for darks and brights across the population. However, regardless of individual neural sensitivity, most neurons responded significantly earlier to dark than bright stimuli. This resulted in a dissociation between neural preference and visual response onset latency: a neuron could exhibit a weaker response to a dark stimulus than to a bright stimulus of the same contrast, but it would still have an earlier response to the dark stimulus. Our results highlight an additional candidate visual neural pathway for explaining behavioral differences between the processing of darks and brights, and they demonstrate the importance of temporal aspects in the visual neural code for orienting eye movements.

Significance statement

Objects in our environment, such as birds flying across a bright sky, often project shadows

(or images darker than the surround) on our retina. We studied how primate superior

colliculus (SC) neurons visually process such dark stimuli. We found that the overall

population of SC neurons represented both dark and bright stimuli equally well, as

evidenced by a relatively equal distribution of neurons that were either more or less

sensitive to darks. However, independent of sensitivity, the great majority of neurons

detected dark stimuli earlier than bright stimuli, evidenced by a smaller response latency for

the dark stimuli. Thus, SC neural response latency can be dissociated from response

sensitivity, and it favors the faster detection of dark image contrasts.

Introduction

Early visual processing is segregated into parallel pathways conveying information about either luminance increments or decrements in visual scenes (Hartline, 1938; Schiller et al., 1986). Such segregation starts in the retina and persists in the early retino-geniculate visual pathway (Hubel and Wiesel, 1961; Schiller et al., 1986). Interestingly, such segregation is also accompanied by asymmetries with which dark and bright stimuli are processed. For example, primate retinal ganglion cells possess asymmetric spatial and temporal properties depending on whether they are part of the ON or OFF pathway (Chichilnisky and Kalmar, 2002). Similarly, in the primate lateral geniculate nucleus (LGN), neurons with OFF-center response fields (RF's) are more sensitive to their preferred stimuli (dark contrasts) than neurons with ON-center RF's experiencing bright contrasts (Jiang et al., 2015). OFF-center neurons also have higher spontaneous activity and more sustained responses during visual stimulation (Jiang et al., 2015). Ultimately, signals reach the primary visual cortex (V1), where ON/OFF asymmetries are amplified. For example, primate V1 is dominated by "black" responses, especially in the superficial cortico-cortical output layers (Yeh et al., 2009).

Asymmetries in the processing of dark versus bright stimuli might make ecological sense. For example, the incidence of dark contrasts in natural scenes is not necessarily uniform. Instead, there is a coincidence of dark contrasts with regions of low spatial frequency, high contrast, and far binocular disparities in natural images (Cooper and Norcia, 2015). As a result, rhesus macaque V1 neurons having far preferred binocular disparities tend to also prefer dark contrasts (Samonds et al., 2012). Similarly, in cat V1, there is a systematic contrast-dependent OFF-dominance, matching natural scene statistics (Liu and Yao, 2014),

and cat V1 neurons are more strongly driven by luminance decrements than increments at

low spatial frequencies (Kremkow et al., 2014). Interestingly, cat studies revealed that ON

and OFF domains in the LGN also exist in the V1 projections (Jin et al., 2008), with area

centralis representations being dominated by dark preferences. Moreover, OFF-dominated

LGN (Jin et al., 2011) and V1 (Komban et al., 2014) neurons respond earlier than ON-

dominated ones. These last observations on OFF and ON channel timing are consistent with

a large body of psychophysical literature for better and faster processing of dark stimuli (e.g.

Komban et al., 2011; Komban et al., 2014).

Having said that, whether monkey superior colliculus (SC) neurons differentially process dark

stimuli remains unclear. In the mouse SC, the majority of superficial layer neurons prefer

dark stimuli (De Franceschi and Solomon, 2018), consistent with the RF subfield structure of

these neurons (Wang et al., 2010). Yet, it is not clear whether such dark preference still

exists in the deeper SC layers, and whether it is accompanied by differences in visual

response latencies. Moreover, differences in the ecological environments and

neuroanatomical organizations of mice and other species do not trivially predict how

primate SC neurons might behave with respect to luminance contrast polarity. Therefore, we

exhaustively characterized all visually-responsive rhesus macaque monkey SC neurons (that

is, also including intermediate and deeper layer neurons). We were particularly motivated by

our recent observations of differential effects of contrast polarity on microsaccades

(Malevich et al., 2021).

In contrast to LGN, V1, and SC results from other species, we did not find a dominant

preference for dark stimuli in the primate SC. Rather, there was significant diversity, with

approximately half of the neurons being more sensitive to bright stimuli. Moreover, at high

contrasts, SC neurons tended to prefer bright rather than dark stimuli, with this trend

disappearing for the lowest contrasts. Despite such diversity, what we did find was that the

majority of SC neurons had significantly shorter visual response latencies to dark stimuli.

Thus, there was a dissociation between visual response latency and visual response

sensitivity, reminiscent of a similar dissociation that we observed in the case of spatial

frequency tuning (Chen et al., 2018). Such a dissociation was sufficient to account for at least

some saccadic reaction time dependencies on stimulus luminance polarity in our

experiments.

Materials and methods

Experimental animals and ethics approvals

We recorded superior colliculus (SC) neural activity from two adult, male rhesus macaque monkeys (M and A) aged 9 and 10 years, and weighing 9.5 kg and 10 kg, respectively. We also measured saccadic reaction times from the same two animals plus a third one (F; aged 11 years and weighing 14 kg). The experiments were approved by ethics committees at the regional governmental offices of the city of Tübingen.

Laboratory setup and animal preparation

The experiments were conducted in the same laboratory as that described in our recent studies (Bogadhi et al., 2020; Bogadhi and Hafed, 2022). Briefly, the monkeys were seated in a darkened booth approximately 72 cm from a calibrated and linearized CRT display spanning approximately 31 deg horizontally and 23 deg vertically. For monkey F only, the display was an LCD device running at 138 Hz (AOC AG273QX2700, 27"), as in (Malevich et al., 2021). Data acquisition and stimulus control were managed by a custom-made system based on PLDAPS (Eastman and Huk, 2012). The system integrated a DataPixx display control device (VPixx Technologies, Inc.) with the Psychophysics Toolbox (Brainard, 1997; Pelli, 1997; Kleiner et al., 2007) and an OmniPlex neural data processor (Plexon, Inc.).

The monkeys were prepared for behavioral training and electrophysiological recordings earlier (Tian et al., 2018; Buonocore et al., 2019; Skinner et al., 2019; Malevich et al., 2020).

Specifically, each monkey was implanted with a head-holder, and monkeys M and A were

also implanted with a scleral search coil in one eye. The search coil allowed tracking eye

movements using the magnetic induction technique (Fuchs and Robinson, 1966; Judge et al.,

1980), and the head-holder comfortably stabilized head position during the experiments.

Eye movements in monkey F were recorded with a video-based eye tracker (EyeLink1000;

desktop mount; 1 KHz sampling rate). For the present experiments, monkeys M and A also

each had a recording chamber centered on the midline and tilted 38 deg posterior of

vertical, allowing access to both the right and left SC (Bogadhi and Hafed, 2022).

Behavioral tasks

For the recording data in monkeys M and A, we employed a gaze fixation task in which we

presented static disc of 0.51 deg radius within the visual response field (RF) of a recorded

neuron. Each trial started with the onset of a black (0.11 cd/m^2) fixation spot at screen

center. After 550-800 ms of stable fixation on the spot, the disc appeared and remained on

for at least ~500 ms. In each trial, the disc could have a Weber contrast of 5%, 10%, 20%,

50%, or 100%. We defined Weber contrast as $|I_s\text{-}I_b|/I_b$, where I_s is the disc's luminance value

and I_b is the gray background's luminance value. We often described the contrast as a

percentage for convenience (e.g. 5% contrast). Importantly, across trials, the disc could have

either positive or negative luminance polarity relative to the gray background, meaning that

I_s could be either higher (positive polarity) or lower (negative polarity) than I_b. The gray

background had a luminance (I_b) of 25.09 cd/m^2. We collected approximately 50 trials per

condition per neuron.

For some neurons in both monkeys (sometimes in the very same sessions as in the above

task), we also ran an immediate orienting version of the stimulus polarity task. That is, at the

time of disc onset, we extinguished the fixation spot (which was now white instead of black)

simultaneously. This instructed the monkeys to generate an immediate orienting saccade

towards the disc. We used this task to confirm that initial visual responses in the main task

above were not dictated by the black fixation spot at display center, since the current task

had a white fixation spot and showed similar observations (see Results), and also to obtain

saccadic reaction time data for additional behavioral analyses (see below). Also note that,

for neurophysiological analysis purposes, we only analyzed the initial visual response in this

task. Saccade-related responses were deferred to another unrelated project focusing on SC

motor bursts, and they are not described here. Finally, to reduce trial counts in this task, we

only tested three contrast levels (10%, 50%, and 100%). We collected approximately 50 trials

per condition per neuron.

For exploring a potential behavioral consequence of faster detection of dark stimuli by SC

neurons (which we describe in Results), we tested our three monkeys on the saccadic

reaction time version of the task, which we just described above. For monkeys M and A, we

analyzed reaction times from the same sessions as those collected during neurophysiological

recordings. Stimulus locations were, thus, dictated by recorded neurons' response field (RF)

locations. For monkey F, we ran behavior-only sessions. In this case, we randomly varied

stimulus locations across 4 diagonals, with an eccentricity of 8.9 deg. We analyzed a total of

457-773 saccades per condition per monkey for our behavioral reaction time analyses.

Neurophysiological procedures

For most experiments, we recorded SC neurons using linear electrode arrays inserted across

the SC depth (24 channel V Probes; Plexon, Inc.). For some experiments, we also used single

tungsten electrodes, in which we targeted and isolated individual neurons online during the

experiments. In all cases, including the single electrode sessions, we also performed offline

sorting to re-isolate neurons for inclusion in the data analysis pipeline (Pachitariu et al.,

2016). Sorting and general data analysis pipeline details were similar to those described

recently (Bogadhi and Hafed, 2022).

Before collecting data from our main tasks, we first identified the SC by running RF mapping

tasks. These included delayed and memory-guided saccades (Chen et al., 2015; Chen and

Hafed, 2017). The mapping tasks allowed us to select the stimulus location for our main

experiments, and also to confirm that our neurons possessed visual responses (or stimulus-

triggered inhibition). For the simultaneous recordings of multiple neurons with electrode

arrays, we picked a disc location that we felt lay within the RF's of most neurons that we

could identify online. This was possible given that our electrodes were penetrating the SC

surface at a quasi-orthogonal angle, meaning that the RF's at different depths generally had

similar locations. Also note that for all analyses, we were always interested in comparing

responses to bright and dark discs at the very same location. That is, our comparison of

interest was the luminance polarity at a given RF location for a given neuron. In separate

experiments, we mapped RF's with positive and negative luminance polarity spots, but these

data will be described in detail separately. For the present purposes, suffice it to say that all

RF's had sensitivity to both black and white targets at their center, justifying our current

comparison of response sensitivity at a single given RF location per neuron.

Across all experiments, we recorded from neurons with extrafoveal eccentricities (e.g. 2.1-

20 deg preferred eccentricity across the population), meaning that we often presented

stimuli far from the fixation spot.

Eye movement data analysis

We detected saccades and microsaccades as described previously (Chen and Hafed, 2013;

Bellet et al., 2019). We used the detections for two primary purposes. First, in the recording

tasks, we excluded all trials in which there were microsaccades occurring within an interval

from -50 ms to +50 ms relative to stimulus onset. This allowed us to measure baseline visual

responses that were not modulated by the known influences of microsaccades on SC activity

(Hafed and Krauzlis, 2010; Chen et al., 2015; Chen and Hafed, 2017). We also performed the

same filtering for behavioral analyses of saccadic reaction times. Second, for the behavioral

analyses, we used saccade detection to measure saccadic reaction times towards the dark

and bright stimuli.

To identify a response saccade and subsequently analyze its reaction time, we required that

it had a latency of 50 to 500 ms from stimulus onset, and that it was directed towards the

stimulus (this latter criterion was easy to achieve because we used computer-controlled

reward windows around the target to allow rewarding the monkeys based on successful

saccade generation towards the target). In all neural and behavioral analyses, we also

excluded trials with blinks or other movement artifacts near stimulus onset. Statistically, we

were interested in whether contrast or luminance polarity affected saccadic reaction times.

Therefore, we performed a 1-way non-parametric ANOVA (Kruskal-Wallis test) in each

monkey testing for the effect of contrast (d.f.: 2), when collapsing across luminance

polarities, on the monkey's reaction times. Similarly, we also performed a Kruskal-Wallis test

exploring the effect of luminance polarity (d.f.: 1), when collapsing across contrasts, on

reaction times.

Neural data analysis

We analyzed a total of 221 SC neurons (109 from monkey M and 112 from monkey A) from

the fixation task. We also analyzed 225 neurons from the immediate saccade version of the

task (113 from monkey M and 112 from monkey A). Ninety of the neurons in the second task

(all in monkey A) were also recorded from the fixation variant of the task.

The bulk of our analyses was on neurons exhibiting a positive visual response to stimulus

onset (that is, an increase in firing rate relative to baseline shortly after stimulus onset). We,

therefore, first tested for the presence of a positive visual response (or burst) after stimulus

appearance. In each neuron, we defined a baseline interval as the final 50 ms before

stimulus onset. We then defined a visual response interval as the time interval 10-200 ms

after stimulus onset. Across all repetitions of a given stimulus condition (e.g. 100% contrast;

positive polarity), we measured average firing rate in the response interval and statistically

compared it to average firing rate in the baseline interval. If the response interval firing rate,

across repetitions of a given condition, was statistically significantly larger (one-tailed, paired

t-test; $p < 0.025$) than baseline firing rate, and if this significance occurred for both polarity

conditions (dark and bright) and with absolute Weber contrasts of 50% and 100%, then we

considered the neuron to have a positive visual response to stimulus onset. We did not

include lower contrast trials (whether positive or negative polarity) in assessing for the

presence of visual responses because some neurons, even when having strong visual bursts

for high contrast stimuli, did not respond to such lower contrasts. Across our population in

the fixation task, we had a total of 172 neurons (92 from monkey M and 80 from monkey A)

exhibiting visual bursts after stimulus onset with the above criteria. For the immediate

saccade version of the task, we only analyzed neurons with a positive visual burst in the

interval 10-180 ms after stimulus onset; this resulted in a total of 213 neurons (109 from

monkey M and 104 from monkey A).

For a subset of neurons, stimulus onset caused a transient decrease in firing rate from

baseline, rather than an increase. We performed analyses of these neurons as well, from the

fixation task only. To assess the neurons as having a transient decrease in firing rate that was

time-locked to stimulus onset, we repeated the same procedure above, but we now checked

for a statistically significant decrease in firing rate in the response interval, rather than an

increase. We analyzed 15 neurons with transiently decreasing firing rates immediately after

stimulus onset (9 from monkey M and 6 from monkey A).

To obtain contrast sensitivity curves from the neurons with visual bursts, we measured the

peak value of the average firing rate curve in a response interval after stimulus onset. Since

visual response latency in the SC varies with stimulus contrast (Li and Basso, 2008; Marino et

al., 2012; Marino et al., 2015), we tailored the measurement interval for each contrast as

follows: 15-105 ms after stimulus onset for 100% contrast; 20-110 ms after stimulus onset

for 50% contrast; 20-115 ms after stimulus onset for 20% contrast; 35-125 ms after stimulus

onset for 10% contrast; and 45-135 ms after stimulus onset for 5% contrast. Note that we

used the same measurement intervals for all neurons and also for both positive and negative

polarity stimuli. Even though we found a difference in response latency between positive

and negative polarity stimuli (as described in Results below), our measurement intervals

were large enough to encompass (and exceed) any such latency differences. Therefore, our

estimates of contrast sensitivity for brights and darks were not biased by using similar

measurement intervals for both types of stimuli (especially because we were searching for

only the peak firing rate). After measuring firing rates in the above intervals for each

contrast, we plotted the measured firing rates as a function of absolute contrast. We then fit

contrast sensitivity curves using the following equation:

$$f(c) = R \frac{c^N}{C_{50}^N + c^N} + B \qquad \text{(Equation 1)}$$

where f is the estimated firing rate, c is stimulus contrast, $C50$ is semi-saturation contrast, R

is the dynamic range of the response, N is the sensitivity/slope of the contrast sensitivity

curve, and B is the baseline firing rate (which we just measured across all trials from the

same baseline interval mentioned above; final 50 ms before stimulus onset). We then

compared the fit parameters R, $C50$, and N for either bright or dark stimuli to assess whether

there were differences in contrast sensitivity between them in the SC. We did this by

computing parameter modulation indices as a function of luminance contrast polarity. For

example, to compare how R was modulated by luminance polarity, we calculated the R

parameter for bright stimuli minus the R parameter for dark stimuli, and we divided this

difference by the sum of R values for bright and dark stimuli. This gave us a value between -1

and +1. We then plotted histograms of parameter modulation indices across the population.

For assessing the time course of changes of contrast sensitivity in the sustained interval long

after stimulus onset (that is, after the initial visual burst), we again obtained fits of equation

1 but now based on measurements after the initial visual burst. To do so, we exploited the

fact that our firing rate estimates were already averaging across time (because of a

convolution of spike times with a gaussian of σ 40 ms). Therefore, for each time after 80 ms

after stimulus onset and before 220 ms, we used instantaneous average firing rate as a

measure to input to the fit of equation 1 for a given contrast. This allowed us to obtain time

courses of semi-saturation contrast ($C50$), sensitivity/slope (N), and dynamic range (R) during

the sustained interval long after stimulus onset. Note that even though the intervals that we

chose for the sustained response analysis slightly overlapped with the intervals that we

picked up for the initial visual burst analyses mentioned above, the latter analyses were

performed on the peak values of the average firing rates, which definitely belonged to the

earlier phases of the neural responses (and typically occurred earlier than 80 ms); that is, the

initial visual burst intervals were just ranges meant to catch the peak response. Also note

that the above contrast sensitivity fits were only performed on the fixation version of the

task because we could obtain a longer period of sustained response than in the immediate

saccade version of the task.

For estimating visual response latency in both tasks, we measured the firing rate of a given

neuron in a baseline interval (final 50 ms before stimulus onset) across all trials. Then, for

each condition (e.g. bright luminance polarity; 20% contrast), we marched forward in time

after stimulus onset until 300 ms (typically, the algorithm converged on a visual burst much

earlier, of course). As soon as the lower bound of the 95% confidence interval around the

average firing rate of the neuron across trial repetitions exceeded the average baseline

activity (and continued to do so for at least 30 ms), we flagged the time as the response

latency of the neuron. Whenever this algorithm failed to detect the response latency in at

least one of the luminance polarities for a given stimulus contrast level, we excluded the

neuron from further analysis in that contrast level. This explains the varying numbers of

neurons reported in some figures (e.g. the different panels of Fig. 4 in Results). We then

compared such response latency across contrasts and stimulus polarities. Note that we

focused on relative latency differences across luminance polarities in our analyses. This is

important to note because firing rate estimates (in our case, convolution of spike times with

a Gaussian kernel) necessarily blurs the exact response onset times of the neurons.

However, our approach of estimating response latencies described above still captured the

latency differences that we were interested in documenting, and it simplified the detection

of visual response latencies for neurons with non-zero baseline firing rates.

To statistically test for differences in latencies between luminance polarities at a given

contrast level, we used non-parametric permutation tests on the pairwise mean latency

differences, with 10000 permutations. That is, we obtained the permutation distribution by

shuffling the polarity labels of the latencies for 10000 times while maintaining their pairwise

relationship and calculating their pairwise difference. Monte Carlo p-values were obtained

by assessing the probability of getting larger than or equal to absolute latency differences in

the permutation distribution than the absolute latency difference of the original data. We

ran the tests separately for each monkey to ensure that our pooling of data in figures for

visualization purposes was justified.

We also used a similar approach to test for statistically significant effects of upper versus

lower visual field RF location on the latency differences between luminance polarities. This

time, we obtained the latency differences between the responses to bright and dark stimuli,

and then subtracted this measurement for the upper visual field neurons from the

measurement for the lower visual field neurons. We defined lower and upper visual field

neurons based on the location of the stimulus (which was placed close to the location of the

RF hotspot location). Thus, negative values in the final measurement would indicate a larger

difference between dark and bright stimuli in the upper visual field than in the lower visual

field. After that, we ran permutation tests by shuffling the labels of the upper and lower

visual field neurons for 10000 times. To assess the significance of the results, we calculated

the Monte Carlo p-value. The same procedure was applied to assess the absolute values of

sensitivity differences (see next paragraph) between dark and bright stimuli in the upper and

lower visual fields.

To compare visual response latency to sensitivity, we also measured peak firing rate in the

initial visual response interval (as defined for each contrast above) of the neuron. First, to

test whether there was an effect of luminance polarity on sensitivity, we used permutation

tests in the same way as we did for the latency analysis described above, but this time on

the pairwise mean peak response differences, separately for each monkey and contrast

level. We then checked whether there was a dissociation between response latency and

sensitivity (i.e. response strength) for black and white stimuli, as we previously saw for

spatial frequency stimuli (Chen et al., 2018). We did so by sorting the neurons according to

the difference in response latencies between brights and darks, and then checking whether

the same sorting applied to the difference in response sensitivities. Further, we pooled the

data across monkeys and calculated Pearson's correlation coefficients between differences

in peak visual responses and differences in visual response latencies, separately for each of

the contrast levels (see Results).

For the immediate saccade version of the task, we only analyzed initial visual bursts (50-130

ms after stimulus onset) and not sustained intervals. This was because the response saccade

occurred too soon after the initial visual bursts. We assessed both response sensitivity and

response latency (as described above) to confirm that we got similar results to those from

the fixation task.

For the neurons with transient decreases in firing rate, we assessed response latency in a

similar way to the neurons with visual bursts, but we looked for statistically significant

decreases in firing rate after stimulus onset, rather than increases.

In some figures, for illustration and visualization purposes, we elected to show example

population firing rates from individual monkeys. For example, we did this in Fig. 8A, B in

Results. To obtain such population firing rates, we obtained the normalized average firing

rate of each neuron, per monkey and condition. That is, for each neuron, we found the peak

visual response in the interval 0-100 ms after stimulus onset for the 100% contrast stimuli,

regardless of the stimulus polarity. Then, for each contrast and polarity, we normalized the

neuron's average firing rate by that peak visual response value. This resulted in a series of

average normalized curves for the neuron across conditions. After that, we averaged all of

the normalized firing rate curves of each monkey's neurons in a given condition. This gave us

a population summary of responses, maintaining the relative changes in responses across

conditions. We used a similar approach in Fig. 11B, D.

Experimental design and statistical analyses

We recorded neurons in an unbiased manner by collecting data in parallel (with linear

electrode arrays) in most sessions and then sorting the neurons offline. This allowed us to

minimize sampling bias. In each variant of the task, we also analyzed >80 neurons per

monkey. This provided a large enough sample to assess the reliability of our interpretations.

Within each neuron, we ensured collecting approximately 35-50 repetitions per condition

(after filtering out bad trials and so on) to allow robust within-neuron statistics. Similarly, in

our behavioral analyses, we collected thousands of saccades. In all cases, we randomly

interleaved stimulus presentations across trials, to avoid any blocking effects.

We provided descriptive statistics in all figures, showing numbers of observations and

measures of variability. Also, in most of our critical analyses (e.g. Figs. 4-6 in Results), we

showed the full distributions of data points that we had.

Since the replicate of interest was neurons, our numbers of sampled neurons were

sufficient. The use of two monkeys in recording was valuable to increase neuron counts, and

to also demonstrate repeatability across individuals. Our results were highly similar in the

two animals (e.g. Fig. 8A, B in Results). When they did differ, we showed each individual

monkey's results separately (e.g. Fig. 11 in Results), and this was highly useful for us to

interpret the behavioral results. Moreover, we collected behavior from a third monkey

exactly to improve our interpretation of our individual monkey behavioral phenomena.

All statistical tests are reported and justified in Results at appropriate points in the text. As

stated above, we statistically analyzed each monkey's data individually, confirming that each

monkey showed the same effects (unless otherwise stated; for example, in Fig. 11 in

Results).

Results

We investigated how monkey superior colliculus (SC) neurons respond to dark and bright visual stimuli. In our primary task, the monkeys fixated while we presented a small disc that was either higher or lower in luminance than the gray background of the display. We varied the contrast of the disc from the background luminance, and we assessed contrast sensitivity curves separately for positive and negative luminance polarities. We first analyzed the neurons that exhibited visual bursts (that is, increases in firing rate) after stimulus onset, and we investigated visual burst strength, visual burst latency, as well as sustained response dynamics for dark and bright stimuli. The results for these neurons are described next, followed by an analysis of a smaller number of neurons for which stimulus onsets caused transient decreases in firing rates, rather than increases.

Diverse preferences for darks and brights across SC neurons

We first asked whether neurons tended to be more sensitive to darks or brights across the population. For each recorded neuron, we plotted firing rate as a function of time from stimulus onset, and we assessed the strength of the visual burst as a function of luminance contrast polarity. Figure 1A-C shows the responses of three example neurons (from the same monkey, A) to a 100% contrast stimulus. The black lines show responses to the negative polarity stimulus (darker than background), and the light gray lines show responses to the positive polarity stimulus (brighter than background). In all cases, the negative and positive polarity stimuli were of the same size and presented at the same location. They also had the same absolute Weber contrast, and their presentation sequence was randomly

counterbalanced across trials. As can be seen, there was a diversity of neural preferences

across the three neurons: neuron 1 (Fig. 1A) was more sensitive to the positive polarity

stimulus than to the negative polarity stimulus; neuron 2 was, more or less, equally sensitive

to the two stimuli (Fig. 1B); and neuron 3 was clearly more sensitive to the dark stimulus

(Fig. 1C). We also plotted full contrast sensitivity curves for the same neurons (Fig. 1D-F) by

relating peak visual response strength to stimulus contrast (Methods). Consistent with Fig.

1A-C, there was a diversity of preferences for darks and brights across the three neurons in

their full contrast sensitivity curves.

These observations held across the population of 172 neurons that we analyzed. For each

neuron, we fit a contrast sensitivity function (equation 1; see Fig. 1D-F for examples) by

optimizing three parameters characterizing how the neuron altered its visual response with

Weber contrast: R reflected the dynamic range of the response, $C50$ characterized the semi-

saturation contrast of the neuron, and N characterized the steepness of the contrast

sensitivity curve (slope parameter). We performed such a fit for either positive or negative

luminance polarity stimuli. We then obtained a parameter modulation index, describing, for

each neuron, to what extent each parameter of the fit was different between positive and

negative luminance polarity stimuli. For example, for dynamic range (parameter R in

equation 1), we obtained the R value for bright stimuli minus the R value for dark stimuli in

each neuron, and we then divided this difference by the sum of R values for the two stimulus

types (Methods). This gave us an index in which 1 meant that the neuron responded

maximally only to bright stimuli and -1 meant that the neuron responded maximally only to

dark stimuli. An R parameter modulation index value of 0, instead, indicated equal visual

response dynamic ranges for bright and dark stimuli. We then plotted histograms of the

parameter modulation indices across the population. As can be seen in Fig. 2, all three

modulation indices of the contrast sensitivity function fits had distributions straddling 0, and

with large diversity across the population. Some neurons clearly preferred bright stimuli,

others clearly preferred dark stimuli, and yet others were equally sensitive to darks and

brights (near 0 in the histograms of Fig. 2). The vertical lines in Fig. 2 indicate the mean

(solid) and median (dashed) parameter modulation index values across neurons, and they

were all close to 0. Approximately half of the neurons were more sensitive to bright stimuli

(whether in terms of dynamic range, semi-saturation contrast, or slope of the contrast

sensitivity function), and the other half were more sensitive to dark stimuli.

Therefore, in the SC, we noticed a substantial diversity of sensitivity preferences for darks

and brights across the population (unlike in LGN and V1). This suggests that stimuli of both

positive and negative luminance polarities can indeed be represented well by SC neural

populations.

Earlier detection of darks by SC neurons, regardless of preference

Unlike response sensitivity, for which we saw diverse preferences for brights and darks (Figs.

1, 2), SC neurons exhibited systematically shorter visual response latencies for dark stimuli,

independently of their visual response strengths at a given contrast. Consider, for example,

the same three neurons of Fig. 1A-C. In each of them, visual responses occurred earlier for

the dark stimuli than for the bright stimuli, as can be visually assessed from the spike rasters

and the firing rate density plots below them. This happened even for neuron 1, which

preferred bright stimuli (Fig. 1A). It also happened at different contrast levels (Fig. 3), even

though stimulus contrast expectedly modulated the response strength and latency of each

neuron. For example, at 20% contrast, all three neurons from Fig. 1 still responded earlier to

dark than bright stimuli, despite the weakened and delayed visual responses relative to the

100% contrast conditions. Thus, at each contrast level, there was an apparent dissociation

between visual response sensitivity and visual response latency in these three neurons, not

unlike what we recently observed when we presented different spatial frequencies to SC

neurons (Chen et al., 2018).

To investigate this dissociation further, we estimated, for each neuron, the onset of the

visual burst as the first time point at which the lower bound of the 95% confidence interval

of the neuron's average firing rate was elevated for a prolonged period of time above its

baseline activity (Methods). Even though estimating neural response latencies from firing

rate measures like we did might blur the actual absolute values of the response latencies,

due to convolution kernels with spike times, this approach was still sufficient to capture the

latency differences across luminance polarities that we were interested in (Methods).

Therefore, for each contrast, we subtracted each neuron's visual response latency for dark

stimuli from its visual response latency for bright stimuli, and we sorted the neurons

according to this difference. An example of such sorting can be seen in the top panel of Fig.

4A for the 100% contrast stimuli. Note how a majority of neurons (76.7%; 128 out of 167)

had an earlier visual response for dark stimuli (evidenced by a positive latency difference in

the figure). This is in contrast to the diversity of preferences for darks and brights seen in

Figs. 1-3. In fact, with the very same sorting of the neurons as in the top panel, we next

plotted (bottom panel of Fig. 4A) the same neurons' differences in peak visual burst

strengths between darks and brights (Methods). The neurons were no longer as properly

ordered as in the top panel, suggesting that the latency effect in the top panel was not

trivially explained by a systematic difference in response sensitivity between darks and

brights. For example, both neurons 126 and 156 (highlighted in Fig. 4A with small diagonal

arrows) possessed clearly stronger responses for bright stimuli than dark stimuli (positive

difference in the bottom panel), but they both had a later response latency for bright stimuli

(positive difference in the top panel). Therefore, visual responses to dark stimuli still

occurred earlier than visual responses to bright stimuli even when neurons preferred bright

stimuli.

We also made similar observations for the other stimulus contrasts that we tested (Fig. 4B-

E). Note that for each panel in Fig. 4, we indicated the total number of neurons included into

each analysis, which varied across panels (that is, across contrast levels). This happened

because some neurons may not have met our inclusion criteria for estimating visual

response latencies, resulting in slightly different neuron counts across the different panels

(Methods). For example, for the particularly low contrast stimuli (e.g. 5% and 10%), some

neurons did not exhibit any significant visual bursts at all (Methods), so they were not

included in the figure. Having said that, in all contrasts, there was a majority of neurons

responding earlier to dark than bright stimuli (top row in each panel of Fig. 4) regardless of

the relative strengths of their visual responses (bottom row).

We confirmed this observation statistically with permutation tests, conducted on pairwise

latency differences separately for each contrast level and for each monkey (Methods). In

monkey M, there were significantly longer latencies for bright stimuli in all contrasts (100%

and 50% contrasts: mean differences = 3.18 ms and 3.56 ms, respectively, Monte Carlo p-

values < 0.0001; 20% contrast: mean difference = 1.62 ms, Monte Carlo p-value = 0.0448;

10% contrast: mean difference = 2.03 ms, Monte Carlo p-value = 0.0152; and 5% contrast:

mean difference = 4.5 ms, Monte Carlo p-value < 0.001). In monkey A, latencies were

significantly longer for bright stimuli in 100%, 50%, and 5% contrasts (mean differences =

2.97 ms, 4.31 ms, and 4.91 ms, respectively; Monte Carlo p-values: < 0.0001, < 0.0001, and

0.0059, respectively); no significant differences were found in 20% and 10% contrasts (mean

differences = 1.39 ms and 0.27 ms, respectively; Monte Carlo p-values: 0.0548 and 0.7587,

respectively), but the same trends were still there (also see Fig. 5E). Thus, faster detection of

dark than bright stimulus contrasts is a general property of SC neurons.

Of course, our results do not deny that high visual response sensitivity is normally associated

with short visual response latencies. For example, with the sorting of neurons shown in Fig. 4

based on their response latency differences (top row), there was still a hint of an additional

trend: neurons with a smaller latency difference between dark and bright stimuli tended to

be the neurons preferring bright stimuli (bottom row). For example, compare the first and

last quartiles in the bottom panel of Fig. 4A: more bright-preferring neurons occurred in the

first quartile (having 0 or negative latency differences) than in the last quartile (having

positive latency differences). This suggests that there were divergent forces influencing

visual response latency: a neuron strongly preferring bright stimuli might have had its high

response strength for bright stimuli (at a given contrast level) counterbalance the normally

earlier detection of dark stimuli. Indeed, when we evaluated response latency as a function

of both stimulus contrast (a proxy for visual response strength in the neurons) and

luminance polarity, we found that both factors clearly influenced the neurons' visual

response latencies. This is shown in Fig. 5A, B for an example neuron, and in Fig. 5C for the

population. In Fig. 5A, high contrast stimuli evoked stronger and, therefore, earlier visual

responses than low contrast stimuli, as expected (Boehnke and Munoz, 2008; Marino et al.,

2012; Marino et al., 2015; Hafed and Chen, 2016; Chen et al., 2018). With high contrast dark

stimuli, the same neuron exhibited even earlier visual bursts than for high contrast bright

stimuli (Fig. 5B). Across the population, cumulative histograms of estimated visual response

latencies (Fig. 5C), as well as their averages and standard errors of the mean (Fig. 5D),

revealed that increasing stimulus contrast systematically decreased response latencies, as

expected (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015; Chen et al.,

2018), but also that visual response latencies were always systematically shorter, at a given

contrast, for dark than bright stimuli (consistent with Fig. 4; also see Fig. 6 below). This

polarity effect on response latencies had an order of magnitude of a few milliseconds

difference between dark and bright response latencies (Fig. 5E), similar to results in the cat

LGN (Jin et al., 2011) and V1 (Komban et al., 2014). Therefore, both stimulus contrast (a

proxy for response sensitivity) and stimulus polarity (conferring a temporal advantage for

darks) dictated our SC neurons' visual response latencies.

Interaction between stimulus contrast and the processing of darks and brights

by SC neurons

The results of Fig. 5E were particularly intriguing to us, in the sense that the lowest contrast

stimuli (5%) were associated with a seemingly bigger effect of visual response latency

difference between darks and brights than the more visible 10% and 20% contrast targets.

One possibility could be that at 5% contrast, there were fewer bright-preferring neurons.

We, therefore, next asked whether there was an interaction between luminance contrast

level and the preference of neurons for darks or brights. To do so, we replotted the data

above as scatter plots of visual response latency for brights versus darks in Fig. 6A-E and as

scatter plots of visual response sensitivity for brights versus darks in Fig. 6F-K. The latency

plots confirmed our earlier observations that there was systematically faster detection of

dark contrasts across all contrasts. For sensitivity, there was an interaction between contrast

level and SC population preference. At high contrasts (e.g. Fig. 6F, G), the population was

biased toward preferring bright stimuli (despite responding faster for dark stimuli), whereas

at 5% contrast (Fig. 6K), this bias disappeared and tended to be in the opposite direction;

neuron 1 in Figs. 1D, 3A also demonstrates this effect: even though the neuron preferred

brights in its plateau firing rate of the contrast sensitivity curve, its (weak) response at 5%

contrast was still higher for dark targets. Thus, there was an interaction between contrast

level and sensitivity to darks in our SC neurons. While this pattern is different from natural

image statistics (Cooper and Norcia, 2015) and cat V1 properties (Liu and Yao, 2014), in the

sense that we found more bright-preferring than dark-preferring neurons at high contrast, it

does suggest that the larger latency effect magnitude in Fig. 5E at 5% contrast might have

been driven by a larger number of dark-preferring neurons at this contrast level.

Statistically, we confirmed that there were contrast-dependent sensitivity preference

differences between brights and darks. We applied the same pairwise latency difference

procedure described above, but now to pairwise peak visual response differences

(Methods). In monkey M, SC visual responses to brights were significantly stronger than

responses to darks in the 100%, 50%, and 20% contrast conditions (mean differences = 11.69

spikes/s, 8.92 spikes/s, and 4.26 spikes/s, respectively; Monte Carlo p-values: < 0.0001, <

0.001, and 0.0187, respectively); the differences were not significant for 10% and 5%

contrasts, and their trends were in the opposite direction (mean differences = -0.09 spikes/s

and -2.35 spikes/s, respectively; Monte Carlo p-values: 0.9611 and 0.1712, respectively). In

monkey A, the neurons were significantly more sensitive for brights in all but the lowest

contrast (100%, 50%, 20%, 10%, and 5% contrasts: mean differences = 6.73 spikes/s, 7.35

spikes/s, 6.52 spikes/s, 6.92 spikes/s, and 0.81 spikes/s, respectively; Monte Carlo p-values:

0.0106, 0.002, 0.0028, < 0.001, and 0.8652 respectively).

In the same vein, for each neuron, we plotted the visual response latency difference against

the peak response difference in Fig. 6L-P. We used the same conventions as in Fig. 4: a

positive response latency difference indicating a faster response to dark stimuli, and a

positive peak response difference meaning higher sensitivity to bright stimuli. If the results

of Figs. 4, 5 were solely determined by response sensitivity at each contrast level, then all

neurons should have occupied the shaded quadrants of these plots. In contrast, only a

minority of neurons occupied these quadrants, particularly at high contrast. For example, in

Fig. 6L, even neurons with >50 spikes/s difference in peak sensitivity in favor of bright stimuli

were still significantly faster to detect dark stimuli. Interestingly, at 5% contrast, there were

significantly fewer bright-preferring neurons, again providing a plausible explanation for the

relatively large effect size in response latency seen in Fig. 5E at 5% contrast versus 10% and

20% contrast.

Interaction between visual field location and the faster detection of darks by SC

neurons

The above results indicate that there is faster detection of dark than bright stimuli by SC neurons, in general. However, it is also known that SC visual responses preferentially process the upper visual field (Hafed and Chen, 2016), consistent with the notion that eye movements support orienting towards or away from extra-personal stimuli largely occupying the upper visual field (Previc, 1990). If that is indeed the case, then it might be expected that differential temporal processing of dark versus bright stimuli might be magnified in the SC's upper visual field representation. For example, birds of prey, or other threats, across a daylight sky would normally cast dark contrasts on retinal images, and they need to be detected efficiently by SC neurons. We, therefore, also asked whether the results of Fig. 4 could depend on the visual field locations of our recorded neurons.

We repeated the analyses of Fig. 4, but this time after separating neurons based on upper and lower visual field RF locations. The results are shown in Fig. 7 (for the highest contrast stimuli only, for simplicity). There was indeed a larger latency difference between dark and bright stimulus responses in the upper visual field neurons than in the lower visual field neurons (top panel). That is, the latency advantage for dark stimuli was magnified in the case of upper visual field SC neurons. We tested this observation statistically by using a permutation test with 10000 shuffles. Here we should note that although we pooled the data of both monkeys for visualization purposes in Fig. 7, we performed the statistical procedures only on data collected from monkey A. This was because there was a strongly unbalanced sampling of neurons in the upper and lower visual fields in monkey M (68 and 21 upper and lower visual field neurons in this monkey, versus a more balanced distribution of

24 and 33 neurons in monkey A). In monkey A, there was a significant difference between

upper and lower visual field effects (mean difference = -4.09 ms, Monte Carlo p-value =

0.0099, Methods). Therefore, a known visual response latency advantage for the upper

visual field in SC neurons (Hafed and Chen, 2016) was accompanied, at least in one monkey,

by a larger difference between dark and bright stimulus responses. This result is consistent

with the ecological likelihood of dark contrasts in natural environments (Liu and Yao, 2014;

Cooper and Norcia, 2015), and also with the role of the SC's visual processing machinery in

supporting the sampling of extra-personal visual space by orienting eye movements (Previc,

1990; Hafed and Chen, 2016; Fracasso et al., 2022).

Note also that the same dissociation between response latency and response sensitivity

occurred in Fig. 7 as in Fig. 4: the bottom panel in Fig. 7 shows that with the same ordering

of the neurons as in the top panel, response sensitivity was not systematically ordered in

either the upper or lower visual fields, consistent with the results of Fig. 4. Interestingly, the

absolute value of the difference in response strength between dark and bright stimuli was

also higher in the upper visual field neurons than in the lower visual field neurons (monkey

A; mean difference = -13.3 spikes/s; Monte Carlo p-value = 0.0166, permutation test). This

suggests that both latency differences (top panel) and absolute values of sensitivity

differences (bottom panel) between dark and bright stimuli were amplified in the upper

visual field representation of the SC, adding to a growing body of evidence of visual field

asymmetries in the primate SC (Hafed and Chen, 2016; Hafed, 2021; Fracasso et al., 2022).

Independence of the faster SC detection of darks from the luminance polarity at

fixation

Finally, we wondered whether the black fixation spot at display center (Methods) might have

dictated our results above. Such an effect would be unlikely because our neurons were

extra-foveal; our stimuli were, therefore, generally far from the fixation spot (Methods).

However, to unambiguously rule such an effect out, we repeated the same experiment with

two slight modifications. First, the fixation spot was now white instead of black (Methods). If

the black fixation spot was the reason for the faster detection of dark stimuli in the results of

Figs. 1, 3-7 above, then this effect should be altered with a white fixation spot. Second, the

fixation spot was now removed at the same time as stimulus onset, allowing the monkeys to

generate immediate, visually-guided saccades. We analyzed 213 neurons recorded with this

variant of the task (81 were also recorded in the original fixation task). We will describe

saccadic reaction times as a function of contrast and luminance polarity in more detail

below. However, for now, our aim was to replicate the visual burst results shown above. For

each neuron, we normalized the neuron's average firing rate by the peak visual response for

100% contrast stimuli in the interval 0-100 ms after stimulus onset (Methods). We then

averaged all of the normalized firing rate curves of each monkey's neurons (Fig. 8A, B). We

separated the neurons of each monkey in this analysis to demonstrate the repeatability of

our results across the animals, and also because subsequent saccadic behavior later in the

trials differed between them, as we clarify in more detail below.

Both animals had clear visual responses in the task, consistent with the results of the fixation

variant (Figs. 1, 3-7). Most importantly, these responses were also clearly still occurring

earlier for dark stimuli than for bright stimuli (Fig. 8A, B). To summarize these results on an

individual neuron basis, we replicated the same analyses of Fig. 4 (Fig. 8C-E; we combined

the neurons of both monkeys here because of how similarly they behaved in Fig. 8A, B). We

did this for all three stimulus contrast levels that we tested in this variant of the task

(Methods). For all contrasts, most neurons still detected dark contrasts earlier than bright

contrasts, irrespective of neural sensitivity (Fig. 8C-E), just as in Figs. 1, 3-7. Permutation

tests run on the latency differences confirmed this observation for all contrasts (in monkey

M, 100% and 50% contrasts: mean differences = 4.97 ms and 4.42 ms, respectively, Monte

Carlo p-values < 0.0001; 10% contrast: mean difference = 1.62 ms, Monte Carlo p-value =

0.0185; in monkey A, 100% and 50% contrasts: mean differences = 3.71 ms and 3.35 ms,

respectively; Monte Carlo p-values < 0.0001; 10% contrasts: mean difference = 2.52 ms;

Monte Carlo p-value < 0.001). Note that the effect sizes were also of the same order of

magnitude as those shown in Fig. 5E. Therefore, the results of Figs. 1-7 were not trivially

caused by the use of a black fixation spot at screen center. Moreover, the results still

persisted in a more reflexive behavioral task, in which prolonged fixation was not enforced

in the face of a salient eccentric stimulus onset.

Different temporal dynamics of firing rates in the sustained interval for darks

and brights

The results so far have focused on initial visual bursts. However, with prolonged fixation (as

in our primary task of Figs. 1-7), we also observed significant differences in SC neural

response dynamics in the sustained interval (long after stimulus onset) for bright and dark

stimuli. In particular, bright stimuli were generally associated with secondary elevations of

firing rate above those of dark stimuli. To illustrate this, Fig. 9A, B shows the responses of

four example neurons to high contrast stimuli (100%). Two neurons are from monkey A (Fig.

9A), and two neurons are from monkey M (Fig. 9B). In the first three neurons (neurons 5-7 in

Fig. 9A, B), after the initial visual bursts, bright stimuli evoked stronger sustained activity

than dark stimuli (see pink intervals highlighting the sustained interval). The stimuli were still

present within the RF's of the neurons in all cases, but there was an altered response

dynamic after the initial visual bursts, particularly for bright stimuli. Even the fourth neuron

(neuron 8 in Fig. 9B), which showed relatively weak sustained activity, still showed a

subdued secondary peak in firing rate after the initial visual burst for bright stimuli (also see

Fig. 11 below for more details on monkey M's secondary bursts for bright stimuli).

To characterize this altered dynamic of neural responses as a function of time in more detail,

we took each firing rate curve after 80 ms from stimulus onset (that is, after the initial visual

bursts). We then estimated contrast sensitivity curves at each time point. Each time sample

of a firing rate curve is already a kind of average over some discrete measurement interval

(due to the convolution of spike times with a gaussian kernel to generate firing rates).

Therefore, we took each sample of the firing rate curve of a neuron in the sustained interval,

and we used it to fit contrast sensitivity curves from equation 1 at each time point. This gave

us a series of contrast sensitivity curves as a function of time. We then plotted the time

courses of the parameters R, $C50$, and N of the fits during the sustained interval, and we did

this for either bright or dark stimuli. The results across the entire population of neurons are

shown in Fig. 9C. As can be seen, all parameters were varying differently between darks and

brights in the interval around approximately 100-200 ms after stimulus onset (that is, during

sustained visual response intervals), consistent with the example neurons of Fig. 9A, B. The

biggest effect was in the R parameter, which was stronger for brights than darks, suggesting

higher sustained firing rates for brights after the ends of the initial visual bursts. Both *C50*

and *N* gradually changed in a manner that was consistent with higher thresholds and

shallower contrast sensitivity functions in the sustained interval. That is, the contrast

sensitivity of the neurons was generally the highest in the initial visual burst intervals, and it

gradually degraded in sustained intervals (other than the *R* parameter elevation for bright

stimuli). This makes sense given that sustained intervals were generally associated with

much lower firing rates than in the initial visual burst intervals (and were therefore less likely

to be strongly differentially modulated by stimulus properties). In any case, during the

sustained interval, and unlike in the initial phases of SC visual responses, there was a

generalized elevation of firing rates for bright stimuli compared to dark stimuli for all

contrasts. As we show later, this effect was strong enough in monkey M, to the extent that it

appeared to dominate this monkey's saccadic reaction time patterns in the immediate,

visually-guided saccade version of the task.

Earlier detection of dark stimuli also by inhibited SC neurons

In all of the above analyses, we focused solely on neurons exhibiting positive visual

responses (that is, increases in firing rates above baseline). However, with our offline neuron

sorting pipelines (Methods), we also isolated a fewer number of neurons that exhibited

transient decreases in activity after stimulus onset rather than increases. These neurons

were obtained from similar recording sites to those from which we isolated neurons with

visual bursts (we used linear electrode arrays primarily orthogonal to the SC surface;

Methods). The neurons were, therefore, from similar topographic locations to those

associated with the neurons reported in Figs. 1-9. When we analyzed these inhibited

neurons in more detail, we found that their transient, stimulus-induced decreases in firing

rates were still sensitive to luminance polarity. For example, in Fig. 10A, B, we show two

example neurons from one of our electrode penetrations in monkey M. The two neurons

showed classic visual responses (to black stimuli); moreover, their RF's (shown in the insets

for data collected with the presentation of black small spots during fixation) were spatially

localized and overlapping with each other. From the very same electrode penetration, Fig.

10C shows a third sample neuron that was recorded simultaneously with the two other

neurons; it was thus in the same SC topographic region as the two neurons of Fig. 10A, B.

The neuron of Fig. 10C was inhibited instead. Most interestingly, this neuron clearly

"responded" to a high contrast dark stimulus earlier than to a bright stimulus of the same

contrast, with the only difference from the results of Figs. 1-9 being that the response in this

case was a transient reduction from baseline activity rather than an increase. Across the

population of such inhibited neurons (n=15 neurons), we repeated the same latency

analyses of Fig. 4 above. That is, we assessed the relative time of "response" between bright

and dark contrasts (Methods). As can be seen from Fig. 10D, the majority of such inhibited

neurons also reacted to dark stimuli earlier than to bright stimuli, just like with the neurons

possessing visual bursts. Similar observations were also made for the lower contrast stimuli.

Interestingly, all 15 inhibited neurons had their "response" to stimulus onset slightly later

than classic visual bursts in other neurons (compare the visual bursts in Fig. 10A, B to the

inhibition time in Fig. 10C; the inhibition occurred slightly later than the bursts). Therefore,

even inhibited neurons in the SC detected dark contrasts faster than bright contrasts.

Saccadic reaction times can be significantly shorter for dark stimuli

Finally, prior work has demonstrated a tight relationship between SC visual response

properties and saccadic reaction times (Boehnke and Munoz, 2008; Marino et al., 2012;

Marino et al., 2015; Hafed and Chen, 2016; Chen et al., 2018). Specifically, both visual

response sensitivity (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015;

Hafed and Chen, 2016; Chen et al., 2018) and visual response latency (Chen et al., 2018) can

predict such reaction times. Therefore, given the faster response latencies of SC neurons for

dark stimuli that we found, we wondered whether this effect was sufficient to be associated

with faster saccadic reaction times to such stimuli (even when response sensitivity was, on

average, similar for darks and brights, as shown in Fig. 2; or even slightly favoring brights at

high contrasts, as shown in Fig. 6). We tested our two monkeys and a third one on the

immediate visually-guided saccade task; we used the same sessions as in Fig. 8 for monkeys

M and A, and we ran separate behavior-only sessions for monkey F. The monkeys simply

generated a saccade as soon as the target appeared (the fixation spot also disappeared at

target onset, as mentioned above for Fig. 8 and in Methods). We measured saccadic reaction

times and plotted them as a function of stimulus contrast and stimulus luminance polarity.

All monkeys showed faster reaction times for higher contrast stimuli, as expected (Marino et

al., 2012; Marino et al., 2015) ($p < 3 \times 10^{-103}$ in each monkey individually, Kruskal-Wallis test

exploring the effect of contrast on reaction time, when collapsing across luminance

polarities). Interestingly, two out of the three monkeys (A and F) also showed consistently

faster reaction times for the darker stimuli, like with the SC visual bursts. These results are

shown in Fig. 11A, C, E; monkeys A and F were faster to react to dark stimuli at all contrasts.

Monkey M, on the other hand, had faster reaction times for the bright stimuli (Fig. 11C). All

of these results (of luminance polarity effects on reaction times) were significant ($p < 2.5 \times 10^8$

in each monkey individually, Kruskal-Wallis test exploring the relationship between

luminance polarity and reaction time, when collapsing across contrasts); the effect sizes

(shown for each condition in Fig. 11A, C, E) were also substantial. In addition, the effect sizes

in monkeys A and F were of the same order of magnitude as the effect sizes of the visual

response latency differences between darks and brights seen in Fig. 5E.

We were particularly intrigued by the discrepancy in the reaction times of monkey M with

respect to dark and bright stimuli. On the one hand, it might suggest that SC visual response

latency (e.g. Figs. 4, 5) is not the only determinant of saccadic reaction times, which is

indeed plausible. For example, we earlier found that SC visual response latency and visual

response sensitivity together provided a better correlate of reaction times than either

parameter alone (Chen et al., 2018). Therefore, since about half of the neurons in our

population were more sensitive to bright stimuli anyway (Fig. 2), despite the faster detection

of darks, it could be that this particular monkey's reaction times were more dictated by SC

visual response sensitivity than by visual response latency. On the other hand, it could

additionally be the case that the later elevation of responses for bright stimuli that we saw in

Fig. 9 was more pronounced in this monkey, potentially suggesting stronger top-down

control for bright stimuli. In that case, bright stimuli could be preferentially processed by this

monkey. Indeed, in a previous behavioral study in which we investigated the properties of

saccadic inhibition as a function of luminance contrast polarity, this monkey reacted

differently to full field white versus black visual flashes from the two other monkeys in the

very initial oculomotor response to flash onset, again reacting faster for bright than dark

flashes (Malevich et al., 2021) (see their Fig. 3). Therefore, we decided to check how this

monkey's neurons, in particular, reacted to white stimuli long after the initial visual bursts,

and we were able to do so from our fixation variant of the task.

We plotted each monkey's population visual responses for dark and bright stimuli in the

fixation variant of the task, allowing us to explore the longer sustained interval. These results

are shown in Fig. 11B, D. Even though monkey M's neurons still detected dark stimuli earlier

than bright stimuli in the initial visual response period (consistent with all of our results

shown earlier), this monkey's elevation of sustained visual activity for the bright stimuli (e.g.

Fig. 9) was particularly pronounced when compared to monkey A (note the secondary peak

in population firing rate for bright stimuli in Fig. 11D for monkey M, which was stronger than

the same peak in monkey A). We also even saw hints of this secondary elevation in Fig. 8B in

the immediate, visually-guided saccade variant of the task, with a sharper elevation for

bright stimuli right after the initial visual burst and leading up to the saccade-related burst;

however, of course, in this task, this sharper elevation for brights was harder to properly

analyze in the saccade task because of how quickly the motor burst came.

Therefore, the results of Fig. 11 suggest that saccadic reaction times can indeed be faster for

dark than bright stimuli, consistent with the faster detection of dark stimuli by SC neurons,

and that even violations of such an observation (as in the case of monkey M) are still related

to the SC visual responses (in this case, the sustained responses after the initial visual bursts

subside).

In all, our results in this study indicate that SC neurons robustly detect dark stimuli faster

than bright stimuli; that sustained visual responses in the SC instead favor bright stimuli; and

913 that saccadic reaction times can reflect the faster detection of dark stimuli in the SC's initial

914 visual bursts and/or the later elevation for bright stimuli.

Discussion

We evaluated the sensitivity of monkey SC neurons to luminance contrast polarity. We found that there was a diversity of preferences for darks and brights across the population (Figs. 1, 2). However, irrespective of preference (as defined by visual neural sensitivity), most neurons detected dark contrasts earlier than bright contrasts (Figs. 3-8, 10). Such earlier detection of dark stimuli was correlated with faster reaction times for such stimuli in 2 out of 3 monkeys (Fig. 11). And, even in the third monkey, this monkey's observed opposite reaction time effect could be related to sustained modulations of SC neural activity, which exhibited a strong secondary elevation particularly for bright stimuli after the initial visual bursts (Figs. 9, 11).

Our results demonstrate that the primate SC does not necessarily exhibit identical ON/OFF sensitivity asymmetries for brights and darks as LGN and V1, refuting the idea that the primate SC simply inherits its visual properties from V1. For example, V1 neurons mostly prefer dark contrasts (Yeh et al., 2009), unlike in our SC population, and it would be interesting to further investigate whether deep V1 layers, projecting to the SC, violate this property or not. In fact, at high contrasts, our SC neurons significantly preferred bright, rather than dark, stimuli even while having faster response latencies to the dark ones (Fig. 6A, F, L).

Our results are additionally interesting because they add to a growing literature demonstrating that the primate SC is as visual a brain structure as the SC in other species, like mice, in which the SC is the primary recipient of retinal projections and, indeed, a

primary visual structure. Consistent with this idea, the monkey SC receives a large amount of

cortical visual input (Kadoya et al., 1971; Fries, 1984; Lui et al., 1995; Lock et al., 2003;

Cerkevich et al., 2014), in addition to direct retinal input (Perry and Cowey, 1984). Thus, the

SC in primates should be viewed as being even more visual than, say, the mouse SC. Such a

rich visual nature of the primate SC matters a great deal for orienting responses, consistent

with how SC visual responses can be linked to various aspects of saccadic behavior, like

reaction time (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015; Hafed

and Chen, 2016; Chen et al., 2018) and landing accuracy (Hafed and Chen, 2016). Such a link

to saccadic behaviors was also clearly still evident in our current study (e.g. Fig. 11). In the

future, it would be important to relate trial-to-trial variability in saccadic reaction times to

trial-to-trial variability in SC, LGN, and V1 visual responses, as was done previously in V1 (Lee

et al., 2010), to better appreciate the different functional specializations that exist in early

visual responses that occur in multiple brain areas at approximately the same time.

Our observation that the primate SC can represent dark contrasts well (e.g. Fig. 2) is also

consistent with earlier observations that SC neurons detect dark "shadows" (Humphrey,

1968; Cynader and Berman, 1972; Updyke, 1974). We are additionally particularly intrigued

by the earlier response latencies for dark stimuli that we observed (e.g. Fig. 4), as well as by

the altered temporal dynamics of responses during the sustained interval long after stimulus

onsets (e.g. Fig. 9). These observations could potentially be used to further interpret earlier

reports in the literature about SC visual and visual-motor modulations. For example, in

investigating color-related responses in the SC, White and colleagues used a high contrast

black target as the comparison stimulus to the colored ones (White et al., 2009). Because of

that black stimulus, we predict that the latency differences that these authors observed

relative to colored targets were slightly amplified than what they would have observed had

they used a white target as the reference non-colored stimulus. Similarly, Churan and

colleagues investigated how SC visual RF's were modified around the time of saccades

(Churan et al., 2011). They found dramatically different effects depending on whether the

saccades were made across a gray background or across a dark background. It is intriguing to

consider whether (and how) our sustained response effects, amplifying responses for bright

stimuli (e.g. Fig. 9), could be related to their observations.

The fact that SC neurons can be strongly sensitive to dark contrasts is also interesting with

respect to spatial frequency tuning in SC neurons. In recent work, we found that SC neurons

can be sensitive to minute phase shifts of spatial frequency gratings, as small as 1 minute of

arc in amplitude (Hafed et al., 2022). It would be fruitful, in light of these observations and

the current work, to investigate RF subfield structure in more detail, for example, to study

phase tuning in SC neurons. Indeed, both the current work and these recent results motivate

a detailed mapping of RF's with both bright and dark stimuli, to assess asymmetries beyond

just visual response sensitivity and visual response latency. Indeed, prior work with reverse

correlation techniques has suggested that there may be informative observations to be

made about RF's mapped with bright versus dark stimuli (Churan et al., 2012). In the near

future, we hope to report SC RF maps for bright and dark stimuli in detail.

The altered long term temporal dynamics of firing rates as a function of luminance polarity

that we observed (e.g. Fig. 9) also motivate modeling how these dynamics emerge. In V1,

various stimulus factors, like contrast, alter not only the initial visual bursts (as might be

expected), but also the sustained responses. Moreover, such alterations can be modeled

using variants of linear/nonlinear filters and divisive normalization (Groen et al., 2021). It

would be valuable to investigate such models in the SC, and to relate them to asymmetries

in ON and OFF channels in both the LGN (Jin et al., 2011) and V1 (Komban et al., 2014).

Related to this, we would like to investigate, in the near future, how scene statistical

regularities, with respect to orienting eye movements, can allow further expansion of our

upper versus lower visual field analyses of Fig. 7. In these analyses, we were motivated by

the theoretical framework of (Previc, 1990), in which he predicted that SC neurons should

over-represent the upper visual field of retinal images because eye movements are relevant

for sampling extra-foveal visual space. Thus, in the analyses of Fig. 7, we were driven by our

earlier discoveries of significant asymmetries between upper and lower visual field SC

neurons and saccadic performance (Hafed and Chen, 2016; Hafed and Goffart, 2020; Hafed,

2021; Fracasso et al., 2022). Indeed, we found that dark versus bright asymmetries were

amplified in the upper visual field (Fig. 7), and this is ecologically sensible. For example, birds

in the sky normally cast shadows on the retina. However, it would be even more intriguing

to go even deeper when assessing such visual field anisotropies. For example, one could

consider studying SC binocularity in more detail, to investigate whether neurons preferring

far disparities would be more prevalent in the upper visual field representation of the SC or

not. And, if so, would these far-preferring neurons also prefer more dark contrasts, like in

the case of V1 (Samonds et al., 2012)? This is important to consider, especially given how our

neurons seemed to prefer bright stimuli in a contrast-dependent manner (Fig. 6F-K) that was

the opposite of what might be predicted from natural scene statistics (Cooper and Norcia,

2015).

1012

Finally, we found a small subset of neurons, in the same topographic location as our bursting

neurons, that were inhibited by stimulus onset (Fig. 10). Interestingly, these neurons still

"detected" dark contrasts (by their transient inhibition of firing rate) earlier than bright

contrasts. It would be important to investigate whether such neurons contribute to saccadic

inhibition (Reingold and Stampe, 2002; Buonocore and McIntosh, 2008; Hafed and

Ignashchenkova, 2013), which we recently found to also depend on stimulus luminance

polarity (Malevich et al., 2021). Our initial intuition with regard to saccadic inhibition,

described in detail in our theoretical proposal elsewhere (Hafed et al., 2021), is that

structures beyond the SC are critical for this phenomenon. However, this does not deny the

potential involvement of SC neurons (particularly those neurons that are transiently

inhibited by stimulus onsets), and future research should investigate the mechanisms of

saccadic inhibition in much more detail, including recording SC neurons with full or localized

flashes of different luminance polarities like in psychophysics. Critical in those studies would

be to quantitatively assess whether the small latency differences in "inhibition" versus

excitatory visual "bursts" that we observed in Fig. 10 are consistent with the timing

properties of saccadic inhibition or not.

1029

1030

References

Bellet ME, Bellet J, Nienborg H, Hafed ZM, Berens P (2019) Human-level saccade detection performance using deep neural networks. J Neurophysiol 121:646-661.

Boehnke SE, Munoz DP (2008) On the importance of the transient visual response in the superior colliculus. Curr Opin Neurobiol 18:544-551.

Bogadhi AR, Hafed ZM (2022) Express detection and discrimination of visual objects by primate superior colliculus neurons. BioRxiv.

Bogadhi AR, Buonocore A, Hafed ZM (2020) Task-irrelevant visual forms facilitate covert and overt spatial selection. J Neurosci 40:9496-9506.

Brainard DH (1997) The Psychophysics Toolbox. Spatial vision 10:433-436.

Buonocore A, McIntosh RD (2008) Saccadic inhibition underlies the remote distractor effect. Exp Brain Res 191:117-122.

Buonocore A, Skinner J, Hafed ZM (2019) Eye Position Error Influence over "Open-Loop" Smooth Pursuit Initiation. J Neurosci 39:2709-2721.

Cerkevich CM, Lyon DC, Balaram P, Kaas JH (2014) Distribution of cortical neurons projecting to the superior colliculus in macaque monkeys. Eye and brain 2014:121-137.

Chen CY, Hafed ZM (2013) Postmicrosaccadic enhancement of slow eye movements. The Journal of neuroscience : the official journal of the Society for Neuroscience 33:5375-5386.

Chen CY, Hafed ZM (2017) A neural locus for spatial-frequency specific saccadic suppression in visual-motor neurons of the primate superior colliculus. J Neurophysiol 117:1657-1673.

Chen CY, Ignashchenkova A, Thier P, Hafed ZM (2015) Neuronal Response Gain Enhancement prior to Microsaccades. Curr Biol 25:2065-2074.

Chen CY, Hoffmann KP, Distler C, Hafed ZM (2019) The Foveal Visual Representation of the Primate Superior Colliculus. Curr Biol 29:2109-2119 e2107.

Chen CY, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity in macaque midbrain. Nat Commun 9:2852.

Chichilnisky EJ, Kalmar RS (2002) Functional asymmetries in ON and OFF ganglion cells of primate retina. J Neurosci 22:2737-2747.

Churan J, Guitton D, Pack CC (2011) Context dependence of receptive field remapping in superior colliculus. J Neurophysiol 106:1862-1874.

Churan J, Guitton D, Pack CC (2012) Spatiotemporal structure of visual receptive fields in macaque superior colliculus. J Neurophysiol 108:2653-2667.

Cooper EA, Norcia AM (2015) Predicting cortical dark/bright asymmetries from natural image statistics and early visual transforms. PLoS Comput Biol 11:e1004268.

Cynader M, Berman N (1972) Receptive-field organization of monkey superior colliculus. J Neurophysiol 35:187-201.

De Franceschi G, Solomon SG (2018) Visual response properties of neurons in the superficial layers of the superior colliculus of awake mouse. J Physiol 596:6307-6332.

Eastman KM, Huk AC (2012) PLDAPS: A Hardware Architecture and Software Toolbox for Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral Control. Front Neuroinform 6:1.

Fracasso A, Buonocore A, Hafed ZM (2022) Peri-saccadic visual sensitivity is higher in the upper visual field. bioRxiv.

1077   Fries W (1984) Cortical projections to the superior colliculus in the macaque monkey: a
1078   retrograde study using horseradish peroxidase. J Comp Neurol 230:55-76.
1079   Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye
1080   movement chronically in the monkey. J Appl Physiol 21:1068-1070.
1081   Groen IIA, Pinantoni G, Montenegro S, Flinker A, Devore S, Devinsky O, Doyle W, Dugan P,
1082   Friedman D, Ramsey N, Petridou N, Winawer J (2021) Temporal dynamics of neural
1083   responses in human visual cortex. BioRxiv.
1084   Hafed ZM (2021) Superior colliculus saccade motor bursts do not dictate movement
1085   kinematics. bioRxiv.
1086   Hafed ZM, Krauzlis RJ (2010) Microsaccadic suppression of visual bursts in the primate
1087   superior colliculus. J Neurosci 30:9542-9547.
1088   Hafed ZM, Ignashchenkova A (2013) On the dissociation between microsaccade rate and
1089   direction after peripheral cues: microsaccadic inhibition revisited. J Neurosci
1090   33:16220-16235.
1091   Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field Representation in
1092   Primate Superior Colliculus. Curr Biol 26:1647-1658.
1093   Hafed ZM, Goffart L (2020) Gaze direction as equilibrium: more evidence from spatial and
1094   temporal aspects of small-saccade triggering in the rhesus macaque monkey. J
1095   Neurophysiol 123:308-322.
1096   Hafed ZM, Chen C-Y, Khademi F (2022) Superior colliculus visual neural sensitivity at the
1097   lower limit of natural self-induced image displacements. bioRxiv.
1098   Hafed ZM, Yoshida M, Tian X, Buonocore A, Malevich T (2021) Dissociable Cortical and
1099   Subcortical Mechanisms for Mediating the Influences of Visual Cues on Microsaccadic
1100   Eye Movements. Front Neural Circuits 15:638429.
1101   Hartline HK (1938) The response of single optic nerve fibers of the vertebrate eye to
1102   illumination of the retina. American Journal of Physiology 121:400-415.
1103   Hubel DH, Wiesel TN (1961) Integrative action in the cat's lateral geniculate body. J Physiol
1104   155:385-398.
1105   Humphrey NK (1968) Responses to visual stimuli of units in the superior colliculus of rats and
1106   monkeys. Exp Neurol 20:312-340.
1107   Jiang Y, Purushothaman G, Casagrande VA (2015) The functional asymmetry of ON and OFF
1108   channels in the perception of contrast. J Neurophysiol 114:2816-2829.
1109   Jin J, Wang Y, Lashgari R, Swadlow HA, Alonso JM (2011) Faster thalamocortical processing
1110   for dark than light visual targets. J Neurosci 31:17471-17479.
1111   Jin JZ, Weng C, Yeh CI, Gordon JA, Ruthazer ES, Stryker MP, Swadlow HA, Alonso JM (2008)
1112   On and off domains of geniculate afferents in cat primary visual cortex. Nat Neurosci
1113   11:88-94.
1114   Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for
1115   measurement of eye position: an improved method. Vision Res 20:535-538.
1116   Kadoya S, Massopust LC, Jr., Wolin LR (1971) Striate cortex-superior colliculus projection in
1117   squirrel monkey. Exp Neurol 32:98-110.
1118   Kleiner M, Brainard D, Pelli DG (2007) What's new in Psychtoolbox-3? (Abstract). Perception
1119   36.
1120   Komban SJ, Alonso JM, Zaidi Q (2011) Darks are processed faster than lights. J Neurosci
1121   31:8654-8658.
1122   Komban SJ, Kremkow J, Jin J, Wang Y, Lashgari R, Li X, Zaidi Q, Alonso JM (2014) Neuronal
1123   and perceptual differences in the temporal processing of darks and lights. Neuron
1124   82:224-234.

Kremkow J, Jin J, Komban SJ, Wang Y, Lashgari R, Li X, Jansen M, Zaidi Q, Alonso JM (2014)
Neuronal nonlinearity explains greater visual spatial resolution for darks than lights.
Proc Natl Acad Sci U S A 111:3170-3175.
Lee J, Kim HR, Lee C (2010) Trial-to-trial variability of spike response of V1 and saccadic
response time. J Neurophysiol 104:2556-2572.
Li X, Basso MA (2008) Preparing to move increases the sensitivity of superior colliculus
neurons. The Journal of neuroscience : the official journal of the Society for
Neuroscience 28:4561-4577.
Liu K, Yao H (2014) Contrast-dependent OFF-dominance in cat primary visual cortex
facilitates discrimination of stimuli with natural contrast statistics. Eur J Neurosci
39:2060-2070.
Lock TM, Baizer JS, Bender DB (2003) Distribution of corticotectal cells in macaque. Exp Brain
Res 151:455-470.
Lui F, Gregory KM, Blanks RH, Giolli RA (1995) Projections from visual areas of the cerebral
cortex to pretectal nuclear complex, terminal accessory optic nuclei, and superior
colliculus in macaque monkey. J Comp Neurol 363:439-460.
Malevich T, Buonocore A, Hafed ZM (2020) Rapid stimulus-driven modulation of slow ocular
position drifts. eLife 9.
Malevich T, Buonocore A, Hafed ZM (2021) Dependence of the stimulus-driven microsaccade
rate signature in rhesus macaque monkeys on visual stimulus size and polarity. J
Neurophysiol 125:282-295.
Marino RA, Levy R, Munoz DP (2015) Linking express saccade occurance to stimulus
properties and sensorimotor integration in the superior colliculus. J Neurophysiol
114:879-892.
Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response
properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738-
1752.
Pachitariu M, Steinmetz NA, Kadir SN, Carandini M, Harris KD (2016) Fast and accurate spike
sorting of high-channel count probes with KiloSort. Advances in Neural Information
Processing Systems (NIPS 2016) 29.
Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spatial vision 10:437-442.
Perry VH, Cowey A (1984) Retinal ganglion cells that project to the superior colliculus and
pretectum in the macaque monkey. Neuroscience 12:1125-1137.
Previc FH (1990) Functional Specialization in the Lower and Upper Visual-Fields in Humans -
Its Ecological Origins and Neurophysiological Implications. Behavioral and Brain
Sciences 13:519-575.
Reingold EM, Stampe DM (2002) Saccadic inhibition in voluntary and reflexive saccades. J
Cogn Neurosci 14:371-388.
Robinson DA (1972) Eye movements evoked by collicular stimulation in the alert monkey.
Vision Res 12:1795-1808.
Samonds JM, Potetz BR, Lee TS (2012) Relative luminance and binocular disparity
preferences are correlated in macaque primary visual cortex, matching natural scene
statistics. Proc Natl Acad Sci U S A 109:6313-6318.
Schiller PH, Sandell JH, Maunsell JH (1986) Functions of the ON and OFF channels of the
visual system. Nature 322:824-825.

Skinner J, Buonocore A, Hafed ZM (2019) Transfer function of the rhesus macaque
oculomotor system for small-amplitude slow motion trajectories. J Neurophysiol
121:513-529.
Tian X, Yoshida M, Hafed ZM (2018) Dynamics of fixational eye position and microsaccades
during spatial cueing: the case of express microsaccades. J Neurophysiol 119:1962-
1980.
Updyke BV (1974) Characteristics of unit responses in superior colliculus of the Cebus
monkey. J Neurophysiol 37:896-909.
Wang L, Sarnaik R, Rangarajan K, Liu X, Cang J (2010) Visual receptive field properties of
neurons in the superficial superior colliculus of the mouse. J Neurosci 30:16573-
16584.
White BJ, Boehnke SE, Marino RA, Itti L, Munoz DP (2009) Color-related signals in the
primate superior colliculus. J Neurosci 29:12159-12166.
Yeh CI, Xing D, Shapley RM (2009) "Black" responses dominate macaque primary visual
cortex v1. J Neurosci 29:11753-11760.

1186

1187

1188

1189

1190

1191 **Figure legends**

1192

Figure 1 Diversity of preferences, in terms of visual burst strength, for dark and bright stimuli by SC neurons.
(A-C) Visual responses of three example neurons from monkey A to a 100% contrast stimulus appearing within their response fields (RF's). Black indicates that the stimulus was negative in luminance polarity (darker than the background); gray indicates that the stimulus was brighter than the background. Each row of tick marks indicates a single trial, and each tick mark indicates the time of an action potential. The firing rate plots below the raster plots summarize the neurons' firing rates. Neuron 1 had a higher sensitivity to bright stimuli, whereas neuron 2 was equally sensitive to dark and bright stimuli. Neuron 3, on the other hand, clearly preferred dark stimuli. Note that response latency (that is, when the stimulus-evoked action potentials first appeared) was always shorter for dark stimuli (see subsequent analyses). Error bars denote s.e.m. across trials. **(D-F)** For each neuron, we measured peak average firing rate after stimulus onset (individual symbols), and we plotted it as a function of stimulus Weber contrast. We also fit the data with continuous curves (Methods). Neuron 1 had higher contrast sensitivity for bright stimuli, evidenced by the higher plateau firing rate at maximal contrast. Neuron 2 plateaued at the same firing rate for both dark and bright stimuli, and neuron 3 was more sensitive to dark stimuli. Dashed vertical lines indicate *C50*, the semi-saturation contrast of each neuron (Methods).

1207

1208

1209

Figure 2 Diversity of contrast sensitivity curve parameters for dark and bright stimuli across SC neurons. (A) For each neuron, we compared the neuron's visual response dynamic range (parameter *R* in equation 1; Methods) between dark and bright stimuli. A modulation index value of 1 indicates maximal responsiveness to only bright stimuli, and a value of -1 indicates maximal responsiveness only to dark stimuli. Most neurons responded to both stimuli (note the rarity of +/-1 modulation index values), but with varying degrees of sensitivity; some neurons clearly preferred dark stimuli, whereas others clearly preferred bright stimuli (similar to the examples in Fig. 1). The vertical lines show mean (solid) and median (dashed) modulation index values across the population, and they were both close to 0: across the population, SC visual responses were equally sensitive to dark and bright stimuli. **(B)** Similar observations for the semi-saturation contrast (*C50*) parameter. Neurons with positive modulation indices in this case are neurons with higher semi-saturation contrasts for bright stimuli (that is, they were less sensitive to bright than dark stimuli). Again, the population average and median semi-saturation contrasts were largely similar between darks and brights (vertical lines), but with large variability across individual neurons. **(C)** Similar observations for the slope parameter of the contrast sensitivity curves. These results are consistent with the example neurons of Fig. 1.

Figure 3 Earlier visual bursts for dark than bright stimuli at different stimulus contrasts. (A-C) Visual responses of the three example neurons of Fig. 1 at each of our tested contrasts (different columns). Black curves indicate dark stimuli, and gray curves indicate bright stimuli. Reducing contrast expectedly weakened and delayed visual responses for both dark and bright stimuli (compare firing rates across columns; also see Fig. 1D-F). Note, however, that at each contrast level, visual bursts occurred slightly earlier for dark than bright stimuli, even for neuron 1, which was more sensitive to bright stimuli. Also note that at the weakest contrast level (5%), both neuron 1 (**A**) and neuron 3 (**C**) were more sensitive to light decrements than light increments. We quantify these observations in subsequent figures and analyses. Error bars denote s.e.m. across trials.

Figure 4 Faster detection of darks than brights by monkey superior colliculus neurons. (A) For the highest stimulus contrast, we estimated visual response latency (Methods) separately for dark and bright stimuli in each neuron. We then subtracted, for each neuron, the visual response latency for dark stimuli from the visual response latency for bright stimuli (positive would indicate earlier responses for dark stimuli). We then sorted all neurons based on this difference (top panel). In the bottom panel, we used the very same sorting, but we now plotted the difference in peak visual response strength between bright and dark stimuli (Methods). Most neurons had a shorter visual response latency for dark than bright stimuli (top panel; vertical line shows the sorted neuron index at which response latency differences flipped sign from negative to positive). This happened independently of response strength; the bottom panel (with the same sorting) did not show a systematic ordering. For example, the neurons highlighted with diagonal arrows preferred bright stimuli (bottom panel) but still detected dark stimuli earlier (top panel). **(B-E)** Similar results for lower contrasts. Of course, with lower and lower contrasts, the earlier detection of darks was less and less prevalent (see the crossover points in the top row). However, this was because lower contrasts were already associated with delayed and weakened visual responses (see Fig. 5). Also note that the lower row shows a decreasing likelihood of bright-preferring neurons as contrast level decreases, suggesting a contrast-dependent processing of darks and brights in SC neurons (see Fig. 6).

Figure 5 Interaction between contrast and luminance polarity in SC visual response latencies. (A) Visual responses of an example SC neuron from monkey M for high (100%) and low (5%) contrast bright stimuli (that is, with luminance higher than the background luminance). Each row of tick marks represents a single trial repetition, and each tick mark indicates the time of an action potential. The neuron responded earlier to the high contrast stimulus. **(B)** In the same neuron, the visual responses to a dark 100% contrast stimulus (that is, with luminance darker than the background luminance) occurred even earlier than the responses to a bright 100% contrast stimulus. Thus, both contrast and luminance polarity affected the neuron's visual response latency. **(C)** Cumulative histograms of our estimates of visual response latency across all of our neurons, for both stimulus contrast (different colors) and stimulus luminance contrast polarity (solid versus dashed lines). High contrasts were associated with earlier visual response latencies in the SC. In addition, at each contrast, dark stimuli were systematically associated with earlier visual response latencies. **(D)** Average visual response latencies for brights and darks across contrast levels, demonstrating consistently faster responses for dark stimuli. **(E)** Average differences in visual response latencies between responses for bright and dark stimuli per contrast level. All contrast levels were associated with faster detection of dark than bright stimuli. The effect increased in strength with increasing contrast from 10% to 100%. At the 5% contrast condition, the effect was the strongest, likely because there were more dark-preferring neurons than at higher contrasts (see the bottom row of Fig. 4 and Fig. 6K, P). Error bars in **D, E** denote s.e.m. across neurons.

1279
1280
Figure 6 Contrast-dependent SC processing of dark and bright stimuli. (A-E) Scatterplots of response latencies
to dark versus bright stimuli at each contrast level. The proportion of neurons responding faster to darks
(above the unity line) increased with increasing contrast level. The proportion of neurons responding faster to
darks was also substantially larger at the lowest contrast level (**E**). (**F-K**) Scatterplots of peak visual responses to
darks versus brights at each contrast level. With increasing contrast, the proportion of neurons responding
stronger to brights (above the unity line) also increased. (**L-P**) Correlations between differences in visual
response latencies and differences in peak visual responses for brights and darks. The differences were
obtained in the same way as in Fig. 4: a positive response latency difference indicates a faster response to dark
stimuli, and a positive peak response means higher sensitivity to bright stimuli. The gray quadrants indicate the
regions where preferences in terms of sensitivity and latency coincided: neurons in the upper left quadrant
responded both stronger and faster for brights, and neurons in the lower right quadrant responded both
stronger and faster for darks. There were weak negative correlations between response latency and sensitivity
differences across contrasts, consistent with a known relationship between response latency and sensitivity.
Critically, however, earlier responses to darks were not dictated by sensitivity preferences; if this was the case,
most neurons would have occupied the shaded gray quadrants. Instead, at the highest contrast levels (e.g. **L**,
M, **N**), the majority of neurons, which responded earlier for darks, responded stronger for brights (occupying
the upper right quadrants), suggesting that dark stimuli were potent in expediting visual bursts despite the
bursts being non-preferred by the neurons.
1299
1300

1301
1302
Figure 7 Magnification of response latency and sensitivity differences between dark and bright stimuli in the
upper visual field. We repeated the same analyses of Fig. 4A, but now separating the neurons according to
whether they represented upper or lower visual field locations. The top panel shows that the faster detection of
dark stimuli by SC neurons that we saw in Fig. 4 was amplified for upper visual field neurons. The lower panel
again shows that the effects of the top panel were dissociated from visual response sensitivity (all figure
conventions are similar to Fig. 4). Interestingly, the lower panel shows that the absolute value of visual response
sensitivity difference (between brights and darks) was also higher for upper visual field neurons than for lower
visual field neurons (compare the dynamic ranges of the two curves). Therefore, both visual response latency
and visual response sensitivity effects, in terms of luminance contrast polarity, were magnified in the upper visual
field.

1313
1314

1315
1316

Figure 8 Faster detection of darks than brights in the SC, independently of central fixation spot appearance.
(A, B) We repeated the same experiment as in Figs. 1-7, but this time with a white fixation spot rather than a
black fixation spot (Methods). Also, here, we removed the fixation spot at stimulus onset, to allow monkeys to
generate visually-guided saccades to the appearing stimuli. In each panel, we averaged each monkey's stimulus-
aligned firing rates (after normalizing each neuron's firing rate curve to its peak visual response at high contrast;
Methods). The numbers of neurons are shown in each panel, and error bars denote s.e.m. across neurons. As
can be seen, visual responses still occurred earlier for dark stimuli than for bright stimuli in this task, and in both
monkeys. Note that at around 100 ms from stimulus onset, there was an elevation of firing rates, which was the
beginning of the saccade bursts for the triggered eye movements. Nonetheless, in monkey M, the elevation
looked to be sharper than in monkey A, an observation that we discuss in more detail in Figs. 9, 11 below with
respect to saccadic reaction times. **(C-E)** We replicated the analyses of Fig. 4 for the three contrasts that we
tested in this task variant. The same conclusions were reached. Most neurons detected dark stimuli earlier than
bright stimuli in their visual response latencies (top row), and this effect was dissociated from individual neuron
sensitivity to either darks or brights (bottom row). All other conventions are similar to Fig. 4.
1331
1332

1333
1334
Figure 9 Preference for bright stimuli in later sustained intervals of visual neural SC responses. (A) Two example
neurons from monkey A showing how later sustained responses (after the initial visual bursts) were elevated for
bright more than dark stimuli. Responses for 100% contrast stimuli are shown, but similar observations were also
made across contrasts (see **C**). **(B)** Two example neurons from monkey M showing similar observations. Note
how neuron 8 had a weaker sustained response for bright stimuli, but it still exhibited a secondary burst (of small
amplitude) for bright stimuli (also see Fig. 11D for this monkey's population response summary in the sustained
interval). The pink rectangles denote our interval of choice when analyzing sustained visual responses. **(C)** In such
interval, we obtained millisecond-by-millisecond fits of contrast sensitivity curves for bright and dark stimuli. The
dynamic range parameter of equation 1, *R*, showed a clear and significant elevation for bright stimuli relative to
dark stimuli across the population (top panel). This was also the case in each monkey individually. The middle
and lower panels show that the thresholds (middle panel) and slopes (lower panel) of contrast sensitivity curves
were getting progressively worse in the sustained interval (relative to initial visual bursts), as expected, but with
little differences between dark and bright stimuli. Error bars in all panels denote s.e.m. (across trials in **A**, **B**, and
across neurons in **C**).
1349

Figure 10 Faster detection of darks than brights even in SC neurons inhibited by stimulus onset. (A, B) Firing rates of two example neurons from a single linear electrode array penetration into the SC of monkey M. Both neurons had visual responses to dark stimuli in the upper left quadrant, with their RF's (obtained by presenting small black spots at different locations; insets) being well localized in space, consistent with the SC topographic representation (Robinson, 1972; Chen et al., 2019). **(C)** A third neuron recorded simultaneously with the neurons in **A**, **B**. This neuron was inhibited by stimulus onset (also see the RF map in the inset). Nonetheless, the inhibition was still stimulus-dependent: there was earlier inhibition for dark than bright stimuli, consistent with our earlier results (e.g. Figs. 1, 4-8). Error bars denote s.e.m. across trials in **A-C**. **(D)** Replication of the analysis of Fig. 4A (top) for all neurons that were inhibited by stimulus onset. Now, we estimated visual response latency by checking when the neural activity was significantly decreased from baseline. Most neurons were still modulated earlier by dark than bright stimuli, consistent with our earlier results for visual bursts (e.g. Figs. 1, 4-8). All other conventions are similar to Fig. 4A.

1367
1368
Figure 11 Relationship between saccadic reaction times and SC visual response properties for darks and brights
in the SC. (A) Saccadic reaction times as a function of stimulus contrast (x-axis) and luminance polarity (different
lines) in monkey A from the saccade version of our task. Error bars denote s.e.m. across trials. Reaction times
were significantly shorter for dark than bright stimuli at all contrasts, suggesting a potential role for the earlier
SC visual responses for dark stimuli in triggering earlier saccades for such stimuli. **(B)** This monkey's neural
responses during the fixation task showed clearly earlier visual bursts for dark stimuli, with equal visual burst
strengths for darks and brights (consistent with Fig. 2). The figure was obtained similarly to Fig. 8A, B. Note also
that the population sustained response (from the peak of the visual burst onward) was larger for brights than
darks (consistent with Fig. 9). Error bars denote s.e.m. across neurons. **(C)** Monkey M showed the opposite
reaction time effects from Monkey A. **(D)** In this monkey's neurons, the secondary burst for bright stimuli in the
fixation variant of the task was particularly prominent (compare to the monkey A neural responses). This suggests
that in this monkey, this secondary preference for bright stimuli might have dominated the monkey's reaction
times in the saccade task. **(E)** We tested a third monkey behaviorally, and we replicated the monkey A results.
Therefore, in 2 out of the 3 monkeys, saccadic reaction times were earlier for dark than bright stimuli, consistent
with the neural results of Figs. 1, 4-8, 10.
1384