Dynamics of transposable element accumulation in the non-recombining regions of mating-type chromosomes in anther-smut fungi

Marine Duhamel[1,2], Michael E. Hood#[3], Ricardo C. Rodríguez de la Vega#[1], Tatiana Giraud#[1]

\# These authors co-supervised the study

[1] Ecologie Systématique Evolution, IDEEV, CNRS, Université Paris-Saclay, AgroParisTech, Bâtiment 680, 12 route RD128, 91190 Gif-sur-Yvette, France.

[2] Evolution der Pflanzen und Pilze, Ruhr-Universität Bochum, Universitätsstraße 150, 44780, Bochum, Germany.

[3] Department of Biology, Amherst College, 01002-5000 Amherst, Massachusetts, United State of America.

*Corresponding author: Marine Duhamel

IDEEV, Bâtiment 680, 12 route RD128, 91190 Gif-sur-Yvette, France

phone: +33 1 69 15 56 69   fax: +33 1 69 15 46 97

marine.c.duhamel@gmail.com

Abstract

Background: Transposable element (TE) activity is generally deleterious to the host fitness, such that TE copies are often purged by selection, which is facilitated when meiotic recombination reshuffles variation among loci. In the absence of recombination, the number of TE insertions usually increases, but the dynamics of such TE accumulations is unknown.

Results: In this study, we investigated the temporal dynamics of TE accumulation in the non-

recombining genomic regions of 15 *Microbotryum* species, leveraging on a unique dataset of

21 independent evolutionary strata of recombination cessation of different ages. We show that

TEs rapidly accumulated in regions lacking recombination, but that the TE content reached a

plateau at ca. 50% of occupied base pairs by 1.5 MY following recombination suppression. The

same TE superfamilies have repeatedly expanded in independently evolved non-recombining

regions, in particular rolling-circle replication elements (*Helitrons*), despite being scarce before

recombination suppression. The most abundant elements, long-terminal repeat (LTR)

retrotransposons of the *Copia* and *Ty3* superfamilies, expanded through transposition bursts

affecting both the non-recombining regions of mating-type chromosomes and autosomes, thus

suggesting that non-recombining regions constitute a reservoir of TEs that transpose to

recombining regions. Based on LTR sequence divergence within and among copies, we could

distinguish bursts of transposition from gene conversion.

Conclusion: Our study supports the TE reservoir hypothesis, by showing that TE accumulation

in non-recombining has a genome-wide impact. TEs accumulated through bursts, and following

a non-linear, decelerating dynamics, thus improving our knowledge on genome evolution,

particularly in association with sex or mating-type chromosomes.

Key words: genomic degeneration, degeneration tempo, recombination suppression, sex

chromosomes, centromeres, gene conversion

Background

Transposable elements (TEs) are autonomously replicating DNA sequences ubiquitous in eukaryotic genomes (Hua-Van *et al.*, 2005; Wicker *et al.*, 2007; Bourque *et al.*, 2018). TEs have occasionally been recruited by their host genomes, *e.g.,* as regulatory sequences, for centromere maintenance or other functions (Kaessmann *et al.*, 2009; Martin *et al.*, 2009; Hof *et al.*, 2016; Dallery *et al.*, 2017; Lanciano & Mirouze, 2018). However, TE replication and insertion are generally deleterious to their host genome (Payer & Burns, 2019; Andrenacci *et al.*, 2020; Burns, 2020). Their insertions can indeed disrupt gene coding or regulatory sequences (Hancks & Kazazian, 2016), induce epigenetic changes (Castanera *et al.*, 2016; Choi & Lee, 2020) or promote chromosomal rearrangements via ectopic recombination (Lim & Simmons, 1994; Han *et al.*, 2008; Ade *et al.*, 2013; Bennetzen & Wang, 2014; Balachandran *et al.*, 2022). TEs can also incur a substantial replication cost as they can represent a large proportion of the genome (Schnable *et al.*, 2009). TE proliferation is therefore often controlled by the host genome, by inactivating TE copies through epigenetic mechanisms such as RNAi silencing (Saito & Siomi, 2010) and DNA methylation (Zemach *et al.*, 2010; Jones, 2012; Bewick *et al.*, 2019; Deniz *et al.*, 2019).

Deleterious TE insertions can also be purged *a posteriori* by recombination between homologous chromosomes carrying different TE insertions. In the absence of recombination, the average number of TE insertions in the population increases with time (Dolgin & Charlesworth, 2008). TEs therefore tend to accumulate in regions with lower recombination rates (Kent *et al.*, 2017; Rifkin *et al.*, 2022), particularly in non-recombining regions such as in centromeres (Plohl *et al.*, 2014), pericentromeric regions (Wright *et al.*, 2003; Peng & Karpen, 2008; Charlesworth & Campos, 2014), and the non-recombining regions of sex chromosomes in plants and animals (Bachtrog, 2003, 2005; Marais *et al.*, 2008; Ahmed *et al.*, 2014) and of

mating-type chromosomes in fungi (Lengeler *et al.*, 2002; Bakkeren *et al.*, 2006; Badouin *et*

al., 2015; Branco *et al.*, 2018). The TE copies accumulating in the non-recombining regions of

sex and mating-type chromosomes may form a reservoir for the rest of the genome, with TE

copies transposing from these regions onto autosomes when epigenetic defenses are insufficient

(Hood *et al.*, 2004; Wei *et al.*, 2020; Peona *et al.*, 2021).

The accumulation of TEs has been extensively documented, but the dynamics of their

proliferation in non-recombining regions, and whether their accumulation in non-recombining

regions has genome-wide impacts as a source of TEs, are unknown (Charlesworth, 2021).

Comparing TE proportions in non-recombining regions of different ages in closely related

species would allow assessing the dynamics of TE accumulation, *i.e.*, the rate at which TEs

increase in copy number in the absence of recombination. The cessation of recombination leads

to degeneration in a variety of ways, which can have non-linear, decelerating dynamics as

suggested for gene losses on theoretical grounds (Bachtrog, 2013; Charlesworth, 2021) and for

optimal codon usage as shown by genomic data (Carpentier *et al.*, 2022). Alternatively, other

components of degeneration can have a linear dynamics of accumulation, as shown for changes

in protein sequences (Carpentier *et al.*, 2022). As TEs are a major degeneration factor, it is

important to assess whether TEs accumulate rapidly in young non-recombining regions,

whether the accumulation slows down over time, for example because of the evolution of more

efficient control mechanisms, whether the number of TE copies grows exponentially out of

control, versus linearly or by occasional bursts of proliferation. The curve shape for TE

accumulation could thus inform our understanding of how their proliferation is controlled in the

absence of recombination. In addition, comparing the genealogies of TE copies within the

genomes using sequence identity and genomic localisation would allow testing whether TE

insertion events following recombination suppression are restricted to the non-recombining

regions or whether TE copies from non-recombining regions can transpose to the rest of the

genome, thus having a genome-wide impact.

Chromosomes carrying the genes for reproductive compatibility are particularly

interesting and useful models to address the temporal dynamics of TE accumulation in non-

recombining regions. Closely related species can have independently acquired sex

chromosomes with non-recombining regions of different sizes and ages, which have often

extended in a stepwise manner, providing independent events of recombination suppression

within and across species (Lahn & Page, 1999; Bergero & Charlesworth, 2009; Charlesworth,

2017). Such stepwise extensions of recombination suppression produce evolutionary strata of

different levels of genetic differentiation between sex chromosomes, as first observed on the

human sex chromosomes (Lahn & Page, 1999), and later on various sex chromosomes in other

animals and plants (Handley *et al.*, 2004; Bergero & Charlesworth, 2009).

The accumulation of TEs has even been suggested to play a role in the extension of

recombination cessation along sex chromosomes (Li *et al.*, 2016; Kent *et al.*, 2017). TE-induced

loss-of-function mutations constitute recessive deleterious mutations in the recombining

regions adjacent to the non-recombining zone, called pseudo-autosomal regions. The

segregation of deleterious mutations near non-recombining regions can foster the expansion of

recombination suppression. For example, inversions carrying fewer deleterious mutations than

average are selected when rare but suffer from their exposed genetic load when frequent enough

to occur as homozygotes; if such less-loaded inversions occur near a permanently heterozygous

allele, such as near mating-type loci or Y-like sex-determining alleles, they remain

heterozygous and can reach maximal frequencies, thereby expanding the non-recombining

region (Jay *et al.*, 2022; Tezenas *et al.*, 2022). TEs can themselves promote inversions at the

margin of non-recombining regions (Ironside, 2010; Zhang *et al.*, 2011). Additionally,

epigenetic modifications that silence TEs, such as DNA methylation and heterochromatin

formation, can also in themselves block recombination (Maloisel & Rossignol, 1998; Ben-

Aroya *et al.*, 2004; Yelina *et al.*, 2012, 2015; Bachtrog, 2013; Li *et al.*, 2016). If TEs accumulate

in the margin of non-recombining regions, they may thus drive the extension of recombination

suppression, either as ultimate or proximate causes, or both (Li *et al.*, 2016; Kent *et al.*, 2017).

Similarly to sex chromosomes, fungal mating-type chromosomes can display large non-

recombining regions, with evolutionary strata of different ages as well as footprints of

degeneration (Menkis *et al.*, 2008; Whittle *et al.*, 2011; Fontanillas *et al.*, 2015; Idnurm *et al.*,

2015; Branco *et al.*, 2018; Hartmann *et al.*, 2021; Carpentier *et al.*, 2022). Fungal mating-type

chromosomes are particularly useful models for studying sex-related chromosome evolution as

they often display young non-recombining regions, with many independent events of

recombination suppression, which allows studying early stages of degeneration (Whittle *et al.*,

2011; Fontanillas *et al.*, 2015; Carpentier *et al.*, 2022). Fungi with evolutionary strata are most

often automictic, i.e. undergoing intra-tetrad selfing (Hartmann *et al.*, 2020; Vittorelli *et al.*,

2022). Theoretical models have suggested that this may be because automictic fungi are

dikaryotic for most of their lifecycle, even in ascomycetes, thus allowing the sheltering of

deleterious alleles (Jay *et al.*, 2022), and because automixis can prevent diploid selfing to purge

deleterious mutations near mating-type loci (Antonovics & Abrams, 2004; Tezenas *et al.*,

2022). When non-recombining regions occur on fungal mating-type chromosomes, they are

relatively young and some genera display multiple independent events of recombination

cessation across closely related species, allowing to study the early stages of genomic

degenerescence with independent data points (Carpentier *et al.*, 2022). In addition, fungi

generally have small genomes and can be cultivated at the haploid phase, facilitating genomic

assemblies and comparisons. The reduced TE load in fungal genomes compared to plants and

animals (Amselem *et al.*, 2015) further facilitates assemblies of their chromosomes, including

the TE-enriched non-recombining regions (Badouin *et al.*, 2015; Branco *et al.*, 2018). Fungal

defenses against TE activity include several forms of RNAi-mediated post-transcriptional

inactivation and heterochromatic silencing (Gladyshev, 2017), in particular DNA methylation

(Castanera *et al.*, 2016; Bewick *et al.*, 2019) and targeted C-to-T mutation in repetitive

sequences, such as repeat-induced point mutation (RIP), targeting specific di- or tri-nucleotides

(Galagan & Selker, 2004; Hood *et al.*, 2005; Amselem *et al.*, 2015).

Microbotryum fungi form a complex of closely related plant-castrating species, mostly

infecting Caryophyllaceae plants, and multiple species have independently evolved non-

recombining mating-type chromosomes (Branco *et al.*, 2017, 2018; Carpentier *et al.*, 2019;

Duhamel *et al.*, 2022). In *Microbotryum* fungi, as in most other basidiomycetes, mating only

occurs between haploid cells carrying alternative alleles at two mating-type loci: the

pheromone-receptor (PR) locus controlling gamete fusion, encompassing genes coding for a

pheromone-receptor and pheromones, and the homeodomain (HD) locus controlling dikaryotic

hyphal growth, encompassing two homeodomain genes (Kronstad & Staben, 1997). Most

Microbotryum fungi display only two mating types, called a_1 and a_2 (Day, 1979), which is due

to multiple independent events of recombination suppression having linked the two mating-type

loci across the *Microbotryum* genus (Badouin *et al.*, 2015; Branco *et al.*, 2017, 2018; Duhamel

et al., 2022; Carpentier *et al.*, 2022) (Figure 1); a few of the studied *Microbotryum* species still

have unliked mating-type loci, but often with linkage of each mating-type locus to its

centromere (Carpentier *et al.*, 2019) (Figure 1). These two types of genomic linkage, *i.e.,* of PR

and HD mating-type loci to each other or to their centromeres, are beneficial under intra-tetrad

mating (automixis), as they increase the odds of compatibility among gametes of a single

meiosis during automixis (Carpentier *et al.*, 2019), the common mating system in *Microbotryum*

fungi (Hood & Antonovics, 2000; Giraud *et al.*, 2008).

Furthermore, patterns of decreasing divergence between the alleles on alternative

mating-type chromosomes when farther from the mating-type loci in the ancestral gene order

have revealed that recombination suppression extended in a stepwise manner beyond mating-

type loci in several *Microbotryum* fungi, forming evolutionary strata of different ages (Branco

et al., 2017, 2018; Duhamel *et al.*, 2022; Carpentier *et al.*, 2022) (Figure 1 and Supplementary

Table 1). In contrast, the recombining regions of the mating-type chromosomes, called pseudo-

autosomal regions, are collinear and highly homozygous in *Microbotryum* fungi (Branco *et al.*,

2017, 2018; Duhamel *et al.*, 2022). The non-recombining regions of the mating-type

chromosomes have accumulated TEs compared to the pseudo-autosomal regions and the

autosomes (Fontanillas *et al.*, 2015; Branco *et al.*, 2018). However, whether specific TE

superfamilies have accumulated and what is the temporal dynamics of TE accumulation across

these genomic compartments remain unresolved questions.

At the whole-genome scale, *Microbotryum lychnidis-dioicae* carries long-terminal

repeat (LTR) retroelements from the *Copia* and *Ty3* (also known as *Gypsy*, but see Wei *et al.*,

2022) superfamilies (Hood *et al.*, 2005; Amselem *et al.*, 2015) as well as, to a lesser extent,

Helitron DNA transposons, *i.e.*, rolling-circle replicating TEs (Kapitonov & Jurka, 2001). In

the *Microbotryum* species parasitizing *Dianthus carthusianorum*, a burst of *Ty3* retroelements

has caused a massive accumulation of this element in a short period of time, in spite of RIP-like

activity (Horns *et al.*, 2017). In *Microbotryum* genomes, a RIP-like activity targets TCG

trinucleotides (CGA on the reverse complement strand) rather than di-nucleotides as in

ascomycetes (Hood *et al.*, 2005; Johnson *et al.*, 2010; Horns *et al.*, 2012).

In this study, we investigated the temporal dynamics of TE accumulation in the non-

recombining regions of 15 *Microbotryum* species, leveraging on the existence of 21

independent evolutionary strata of different ages, within and across species (Carpentier *et al.*,

2022) (Figure 1). We show that TEs rapidly accumulate in young non-recombining regions but

that their abundance reached a plateau by 1.5 MY following recombination suppression at ca.

50% occupied base pairs. *Helitron* DNA transposons repeatedly expanded in non-recombining

regions despite being rare in species with recombining mating-type chromosomes. The most

abundant TE superfamilies, *i.e.*, *Copia* and *Ty3*, were also over-represented in non-recombining

regions and accumulated through multiple bursts that impacted both the non-recombining

regions of the mating-type chromosomes and the autosomes at the same time, supporting the

TE reservoir hypothesis. TEs, however, did not preferentially accumulate at the margin of non-

recombining regions.

Results

Transposable elements in *Microbotryum* genomes

We identified transposable elements (TEs) in the a_1 and a_2 haploid genomes deriving from a

single diploid individual of each of 15 *Microbotryum* species and of a genome of the red yeast

Rhodothorula babjavae, an outgroup with the ancestral condition of the PR and HD mating-

type loci on separate chromosomes. We detected no significant difference in TE content

between the two haploid genomes within species (Wilcoxon paired test, V=5284, *p-*

value=0.208, Supplementary Figure S1). In the following analyses, we therefore calculated the

mean from the a_1 and a_2 genomes for each species. *Microbotryum* genomes, where there was

an average of 18.3% base pairs occupied by TEs, displayed a higher density of TEs than in *R.*

babjavae (Supplementary Figure S1). *Microbotryum intermedium*, with fully recombining

mating-type chromosomes, had the lowest TE content (6.57%), closely followed by *M. v.*

lateriflora (7.02%) with a very young non-recombining region (0.304 My, Duhamel *et al.*,

2022), while *M. violaceum paradoxa* with the largest and among the oldest non-recombining

regions had the highest TE load (30.33%) (Figure 2a and Supplementary Figure S2 and

Supplementary Table 2; detailed comparison below).

All *Microbotryum* genomes contained a substantial fraction of long-terminal repeat

(LTR) retrotransposons from the *Copia* and *Ty3* superfamilies (6.04% and 4.97% of occupied

base pairs on average for *Copia* and *Ty3* retroelements, respectively, Supplementary Table 4).

These retroelements were particularly abundant in the *Microbotryum* genomes with non-

recombining mating-type chromosomes, *Copia* TEs being for example ten times as abundant in

M. coronariae (10.27% of occupied base pairs) as in *M. intermedium* (1.08% of occupied base

pairs). Taken together, *Copia* and *Ty3* superfamilies represented 76.93 to 94.01% of the

centromeric TEs across species (Figure 2B). The trimodal distributions of *Copia* and *Ty3*

retroelement length (Figure 2C and Supplementary Figure S3a) show that, while most copies

are shorter than 1 kb and are likely degenerated and fragmented, a high proportion of these

retroelements, being 5 kb and 7 kb length on average, are likely intact TE copies (González &

Deyholos, 2012). TE copies with a length greater than 10 kb likely represent nested TE copies

and were particularly abundant in the young stratum of *M. v. paradoxa,* called the white stratum

(Supplementary Figure S3a).

Because retrotransposons such as LTRs can be common in plants (Piegu *et al.*, 2006;

González & Deyholos, 2012), we investigated whether horizontal transfers of TEs could have

occurred between the host plant and *Microbotryum* genomes. We compared the sequences from

the plant Repbase database to the annotated TE sequences in *Microbotryum* genomes by blast.

We obtained several BLAST hits with high identity and low e-value scores, but none exceeded

6% of the annotated TE sequence lengths (Supplementary Table 3). Therefore, no putative inter-

kingdom horizontal TE transfer was identified.

DNA transposons were mostly represented by *Helitrons* and terminal inverted repeats

(TIR) elements in most *Microbotryum* genomes (1.44 and 0.42% of occupied base pairs on

average, respectively), while these were nearly absent in *M. intermedium* (0.04% of base pairs

occupied by *Helitrons* and no TIR elements) and in *M. scabiosae* (Figure 2a). The proportion

of TEs annotated as "*unclassified*" was generally low (0.83 to 3.25% of occupied base pairs),

being highest in genomes with high TE content, such as *M. v. paradoxa* and *M. v. gracilicaulis*

(7.4 and 10.18% of occupied base pairs, respectively). The low average size of "*unclassified*"

TE copies (263.13 bp, Supplementary Figure S3b) suggests that they represent highly

fragmented TE relics. Such "*unclassified*" elements were predominant in subtelomeric regions

(Figure 2B).

Higher content of transposable element and of RIP-like footprints in non-recombining

regions

TEs represented significantly higher proportions of base pairs in non-recombining regions

compared to pseudo-autosomal regions and autosomes (ANOVA post-hoc Tukey test, adj. *p-*

value < 0.01), being on average 41.68% and 40.31% higher in the non-recombining regions

compared to pseudo-autosomal regions and autosomes, respectively (Supplementary Table 4,

Supplementary Figure S4). No significant difference was observed between pseudo-autosomal

regions and autosomes (ANOVA post-hoc Tukey test, adj. *p-value* = 0.922, Supplementary

Table 4), despite tendencies in some species for lower content of some elements in pseudo-

autosomal regions (Figure 2C and Supplementary Figure S4). The centromeres of all species

displayed a high density of TEs (73% of base pairs occupied by TEs on average, Supplementary

Table 5). Plotting the density of TEs along mating-type chromosomes further illustrated the

higher values in the non-recombining regions than in the pseudo-autosomal regions, and that

the TE-rich regions corresponded to gene-poor regions (see Figure 3 for *M. lychnidis-dioicae*

and Supplementary Figure S5A to S5O for the other *Microbotryum* species).

TE density was generally higher in older evolutionary strata such as the stratum around

the PR locus, called the purple stratum (Supplementary Figure S5H, S5J and S5O), and lower

in young strata such as the pink and white strata of *M. v. paradoxa* (Supplementary Figure S5G).

In part because of this variation in TE content related to age of the non-recombining region, the

size of the mating-type chromosomes varied greatly across *Microbotryum* species (Figure 1B),

from 1,177,022 bp in *M. scabiosae* to 10,060,889 bp in *M. v. paradoxa* (3,076,521 bp on

average); the size of the mating-type chromosomes also varies because different chromosomal

rearrangements were involved in the origin of HD-PR locus linkage (Duhamel *et al.*, 2022). For

the same reasons, the size of the non-recombining regions was variable, ranging from 630,774

bp in *M. scabiosae* to 9,461,162 bp in *M. v. paradoxa*, thus spanning almost the full length of

the mating-type chromosome in this later species (Figure 1B), while being absent in *M.*

intermedium.

We found no evidence that TEs have particularly accumulated in the flanking regions

of the non-recombining regions when comparing the proportion of base pairs occupied by TEs

within the 100 kb of the pseudo-autosomal regions flanking the non-recombining regions to the

distribution of the TE proportions in regions of the same size in fully recombining autosomes

(Supplementary Figure S6). Indeed, the probability of having a greater TE load in a 100 kb

window on autosomes than at the margin of the non-recombining region was ≤ 0.05 in a single

species (Supplementary Figure S6). We compared the TE content in the flanking region of the

non-recombining region to that in autosomes because the pseudo-autosomal regions were too

small for building a distribution.

The non-recombining regions had more RIP-like signatures in TEs than autosomes and

pseudo-autosomal regions in most species (Figure 3 and Supplementary Figure S5). The

youngest strata, such as the pink and white strata of *M. v. paradoxa* and the young light blue

stratum in *M. v. caroliniana,* were in contrast less affected by RIP in TEs than the older non-

recombining regions (Supplementary Figure S5G and S5L, respectively).

Particular TE family expansions in non-recombining regions

Some TE families have particularly expanded in non-recombining regions. *Helitron* elements,

for example, have expanded in non-recombining regions, while they are relatively rare in the

recombining mating-type chromosomes of *M. intermedium* and in autosomes in all species

(ANOVA post-hoc Tukey test, adj. *p-value* < 0.01, see Supplementary Table 4 and Figure 2B

and 2C). TEs of the *Copia* and *Ty3* superfamilies displayed higher proportions in the non-

recombining regions compared to autosomes and pseudo-autosomal regions in almost all

species (ANOVA post-hoc Tukey test, adj. *p-value* < 0.01, see Figure 2C and Supplementary

Table 4). The content of *Helitrons*, *Copia* and *Ty3* elements increased on average by 5.42, 6.56,

and 4.77% in non-recombining regions compared to autosomes, while the content in other TEs

increased by 1.48% in non-recombining regions on average across species. The proportion of

TIR DNA transposons was higher in the non-recombining regions compared to pseudo-

autosomal regions but not compared to autosomes (ANOVA post-hoc Tukey test, adj. *p-value*

< 0.01, Supplementary Table 4 and Figure 2C).

Genome-wide bursts of transposable elements following recombination suppression

We found a significant and strongly positive correlation between the proportions of base pairs occupied by TEs on autosomes and non-recombining regions (r = 0.67; *p-value* = 0.006; Supplementary Figure S7), which is consistent with the reservoir hypothesis, *i.e.*, that the TE copies accumulating in non-recombining regions have a genome-wide impact by transposing to autosomes.

We inferred TE genealogies of the *Copia* and *Ty3* superfamilies independently, using the alignment of their 5'-LTR sequences. Because the genealogies rely on the identification of the 5'-LTR sequences of the retrotransposons, only a subset of the *Copia* and *Ty3* retroelements were used to reconstruct the genealogies (on average 12 and 9.65% of the annotated *Copia* and *Ty3* superfamily retroelements, respectively, Supplementary Figure S8) and genealogies were built for the species having sufficiently large sets of retained *Copia* and *Ty3* sequences.

The *Copia* and *Ty3* genealogies revealed the existence of many copies clustering within large clades having very low divergence, suggesting multiple bursts of transposition of these retroelements that impacted all genomic compartments (Figure 4 and additional species shown in Supplementary Figures S9 and S10), including in the species with old non-recombining regions (Figure 4B and Supplementary Figures S9G-H and S10B). Clusters of similar TE copies could however not only be recent bursts of transpositions (that we define here as a sudden accumulation of copies, resulting from either an increase in intrinsic TE activity or from less efficient purge of new insertions) but also gene conversion events among old copies. We tested these alternative hypotheses by considering that conversion events among old copies would generate multiple similar copies with an old age while recent bursts would generate multiple similar copies with a similar age of insertion as their time of divergence. The age of a copy can be estimated from the divergence between 5'-LTR and 3'-LTR sequences as they are identical

at the time of insertion and then diverge gradually with time. Similarly, the 5'-LTR sequences

of a parental copy and its progeny copy are identical at the time of transposition and then diverge

gradually (Figure 5A). Therefore, copies resulting from a burst of transposition events should

have accumulated roughly the same number of substitutions between their 5'- and 3'-LTR

sequences as between their 5'-LTR sequence and the 5'-LTR sequence of their parental copy,

i.e., their most similar TE copy. We therefore expect that, in case of transposition bursts without

conversion events, plotting the number of substitutions between the 5'-LTR sequences of the

most similar copies against the number of substitutions between the 5'-LTR - 3'-LTR sequences

within copies would result in a cloud of points around the first bisector, with some stochasticity.

We estimated the variance around the first bisector expected without gene conversion by

simulating a cloud of points under a Poisson distribution. The points outside the expected cloud,

at the bottom right, likely represent conversion events (low number of substitutions between 5'-

LTRs from different copies but elevated number of substitutions between LTRs within copies).

In all species, we found that only a minority of points fell at the bottom right outside of

the Poisson-simulated cloud of points, for the clusters of similar *Copia* or *Ty3* copies (Figures

5B and 5C, Supplementary Figures S11). This suggests that conversion events are rare and

indicates that the clusters of similar copies correspond mostly to bursts of transposition, due to

increased activity or less efficient selection against insertions, rather than gene conversion

among old copies. For several species, a few points fell on the top left corner outside of the

Poisson-simulated cloud of points, likely corresponding to copies for which the closest related

copy could not be identified. The colors indicating the compartment of origin of the pairs of

copies are scattered on the plots, showing that conversion is not more frequent in any genomic

compartment (Figures 5B and 5C, Supplementary Figures S11).

The recent bursts of proliferation show that *Copia* and *Ty3* retrotransposons are active

in *Microbotryum* genomes. In several species, the centromeric TE copies tended to cluster

together (Figure 4 and Supplementary figure S9 and S10). In a few species, such as *M. v.*

paradoxa (Figure 4B), we identified *Copia* TE bursts specific to non-recombining regions.

However, the clustering of copies from autosomes and non-recombining regions in the same

TE clade with low divergence indicates that TEs transpose from one compartment to the other.

We found similar distributions of pairwise divergence times between TE copies in autosomes

and in the non-recombining regions of the mating-type chromosomes (Supplementary Figure

S12). This indicates that bursts occurred at the same time in the two genomic compartments,

thus suggesting that non-recombining regions serve as reservoirs from which TEs can transpose

in autosomes. Further supporting the reservoir hypothesis, we found fewer pairs of most closely

related copies with both copies in the non-recombining regions than expected by chance given

the number of copies located in the non-recombining regions in the genealogies (decreasing by

67% compared to random distribution on average, ranging from 95.62% to 40.6% across

species, Supplementary Figure S13).

In order to assess whether these TE bursts occurred following recombination

suppression, we inferred TE genealogies for *Copia* and *Ty3* retroelements with the sequences

of all species. The TE copies resulting from the recent TE bursts often clustered by species and

by group of species deriving from the same recombination suppression event, supporting the

inference that TE accumulation has followed recombination suppression (Figures 6 and

Supplementary Figure S14). The bursts of retrotransposon accumulation occurred at different

times across lineages, as shown by the different phylogenetic distances of the clades with short

branches to the root (Supplementary Figure S15A and S15B). The only exception was the

occurrence of three *Copia* bursts at the same time, in the three species of the *M. v. gracilicaulis*

clade, as the three bursts appeared equidistant to the root (Supplementary Figure S15A, pointed

by arrows and aligned on the green circle).

Temporal dynamics of TE accumulation in non-recombining regions on mating-type
chromosomes

The multiple independent events of recombination suppression across the *Microbotryum*

phylogeny, with a range of ages, provides a unique opportunity to investigate the temporal

dynamics of TE accumulation. The large non-recombining regions of different ages formed by

the stepwise recombination suppression between the HD and PR loci were highly rearranged in

M. v. viscidula, *M. v. melanantha*, *M. v. gracilicaulis* and *M. v paradoxa,* rendering difficult the

assignment of some TE copies to a given age of recombination suppression. In order to be

conservative, we therefore only considered TE copies that could be assigned to a stratum with

high confidence, by taking into account only blocks of at least 80 kb (size of the pink stratum,

the smallest stratum of the dataset) and encompassing at least two genes of a given stratum. The

resulting blocks represented 66.52 to 90.52% of the total non-recombining region sizes

(Supplementary Figure S16), thus likely yielding good estimates of their TE content. We

excluded from analyses the orange stratum (Figure 1) shared by all or most of the species

(Branco *et al.*, 2017, 2018; Carpentier *et al.*, 2019, 2022; Duhamel *et al.*, 2022), as it is small

and might not represent independent events of recombination suppression. We also excluded

from analyses the small strata that were too much fragmented (for example the red stratum in

M. lychnidis-dioicae and *M. silenes-dioicae;* Figure 1; Branco *et al.*, 2018). To avoid

pseudoreplication, we considered the mean of the percentages of base pairs occupied by TEs in

the a_1 and a_2 genomes for each species. Similarly, for the strata shared by multiple species, we

took the mean values across the species derived from the same recombination suppression

event. We used the TE content in the autosomes of *M. intermedium* as the point at time zero,

i.e., before recombination suppression, as its mating-type chromosomes still recombine over

almost their entire length, thereby precluding any potential TE reservoir effect.

We tested whether the percentage of base pairs occupied by TEs in the non-recombining

regions could be explained by the time since recombination suppression and the ancestral size

of the non-recombining region. We compared generalized additive models assuming linear or

logarithmic relationships with the time since recombination suppression, with or without

smoothing splines (*i.e.*, phases), as well as a non-linear negative exponential model. The

percentage of base pairs occupied by TEs in the non-recombining regions was best explained

by the negative exponential model with the age of the stratum as an explanatory variable of the

percentage of TEs in this stratum (Figure 7 and Supplementary Table 6). We found that TEs

rapidly accumulated in the first 1.5 million years following recombination suppression and then

reached relatively abruptly a plateau at around 50% occupancy of TEs. A similar curve was

obtained when fitting TE accumulation with the average synonymous divergence (d_S) between

a_1 and a_2 associated alleles of the genes within strata, as expected considering the quasi-linear

relationship between d_S and stratum age (Supplementary Figure S17). The ancestral size of the

non-recombining region did not significantly impact TE content in the retained negative

exponential model.

We found that the relative abundance of RIP-like footprints (as assessed by the

normalized RIP-index metric, see Methods) was highly variable in the young strata and

remained constant with time since recombination suppression, with normalized RIP-index

values greater than zero, suggesting that a process similar to RIP efficiently mutates TE copies

in young non-recombining regions (Supplementary Figure S18). There was indeed no

significant correlation between the age since recombination suppression and the abundance of

RIP-like footprints in the TEs of the strata (two-sided Pearson's correlation test, *p-value* = 0.9).

The fraction of detected intact *Helitron* copies decreased with time since recombination

suppression and even reached zero after 4 MY (Supplementary Figure S19). The proportion of

intact copies of *Copia* and *Ty3* retroelements decreased rapidly following recombination

suppression, from 20% down to 10%, and then remained stable at around 10% (Supplementary

Figure S19).

Discussion

We found a large degree of variation among *Microbotryum* species in their TE load, and the

key to understanding this variation seems to be the evolution of the recombination landscape

associated with regions determining reproductive compatibility. Moreover, there appears to be

a significant feedback of the TE accumulation on non-recombining mating-type chromosomes

where the buildup is rapid, and the rest of the genome is then burdened with greater repetitive

element insertions. Leveraging on a unique dataset with 21 regions having independently

evolved recombination suppression, we found that TEs accumulated rapidly following

recombination cessation but then reached a plateau at ca. 50% of occupied base pairs, after 1.5

MY.

We indeed revealed more details on the degree of variation in genomic TE loads among

Microbotryum genomes than previously recognized, ranging from 6.5 to 30% base pairs

occupied by TEs, with an intermediate TE content in the most studied species, *M. lychnidis-*

dioicae (19.4%). We were also able to annotate a large proportion of the detected TE copies,

e.g. 89% in *M. lychnidis-dioicae*. In comparison, a previous study using Illumina paired-end

sequencing estimated the genomic TE content of *M. lychnidis-dioicae* at 14.1% and only 60%

of the TE copies were annotated (Amselem *et al.*, 2015). Relative to other fungi, *Microbotryum*

genomes carry intermediate TE contents, generally lower than their close relatives in the

Pucciniomycotina (Raffaele & Kamoun, 2012). We found that LTR retrotransposons, especially

from the *Copia* and *Ty3* superfamilies, represented the majority of the base pairs occupied by

TEs in *Microbotryum* genomes. These LTR retrotransposons summed to 87% of the base pairs

occupied by TEs in *M. intermedium* and 70% in *M. lychnidis-dioicae*, which was much higher

than previously estimated and among the highest LTR retrotransposon content in fungi, relative

to other TE categories (Amselem *et al.*, 2015).

Our results also indicate that the lack of recombination has allowed the accumulation of

TEs by a lower efficacy of selection against deleterious insertions, as shown by the higher TE

loads in non-recombining regions of mating-type chromosomes compared to recombining

autosomal or pseudo-autosomal regions. The accumulation of TEs has been widespread in non-

recombining regions of sex chromosomes of plants and animals (Erlandsson *et al.*, 2000;

Bachtrog, 2003, 2005; Marais *et al.*, 2008; Ahmed *et al.*, 2014; Śliwińska *et al.*, 2016;

Mawaribuchi *et al.*, 2017). In the absence of recombination, the average number of TE copies

can only increase with time in the population, because chromosomes devoid of new TE copies

cannot be generated by recombination and the chromosomes carrying fewer TE copies can be

randomly lost, a phenomenon known as Muller's ratchet (Muller, 1918; Charlesworth &

Charlesworth, 2000; Bachtrog, 2003; Dolgin & Charlesworth, 2008). In *Drosophila miranda*,

TEs played an early and important role in the degeneration of the neo-Y chromosome (Bachtrog,

2005; Bachtrog *et al.*, 2008). We detected RIP-like footprints in *Microbotryum* genomes, but

this genome defense mechanism was not sufficient to control TE proliferation in the absence of

recombination, as previously noted in some *Microbotryum* species (Johnson *et al.*, 2010; Horns

et al., 2017).

Transposable elements from the *Copia*, *Ty3* and *Helitron* superfamilies were particularly

involved in the repeat accumulation in non-recombining regions. *Copia* and *Ty3*

retrotransposons proliferated independently and repeatedly following the multiple independent

events of recombination suppression across *Microbotryum* lineages. *Copia* and *Ty3*

superfamilies are predominant in *M. intermedium* and in autosomes of the species with non-

recombining mating-type chromosomes, suggesting that they were already abundant before

recombination suppression, and then proliferated due to transposition and Muller's ratchet.

Helitron elements were in contrast rare in *M. intermedium* and in autosomes of all species, and

yet have also accumulated in the multiple non-recombining regions, as observed in the non-

recombining regions of the sex chromosomes of *Drosophila pseudoobscura* and *D. affinis*

(Nguyen *et al.*, 2022). *Helitrons* transpose by a mechanism similar to rolling-circle replication

via a single-stranded DNA intermediate and have been suggested to preferentially insert in AT-

rich regions; RIP and RIP-related mechanisms increase the percentage in AT and are more

abundant in non-recombining regions, which could have facilitated the accumulation of

Helitrons (Grabundzija *et al.*, 2016). It is striking that the same superfamilies preferentially

expanded repeatedly in multiple non-recombining regions of *Microbotryum* fungi. Their

specific mechanisms of transpositions may be particularly efficient in regions where

recombination is suppressed, perhaps due to the chromatin landscape (Quadrana *et al.*, 2019).

LTR retroelements for example preferentially target heterochromatin (Marsano & Dimitri,

2022).

Through the inferred TE copy genealogies of the most abundant superfamilies in non-

recombining regions (*Copia* and *Ty3* superfamilies), we found recent accumulation of TE copies

through bursts occurring in all genomic compartments, *i.e.*, the non-recombining regions, the

autosomes and the centromeres, many of them being shared by species deriving from the same

recombination suppression event. This insight into the dynamics of TE accumulation in the

different genomic compartments suggests that TE accumulation in *Microbotryum* genomes

followed recombination suppression, not only in the non-recombining regions, but also in the

rest of the genome. TE accumulation in non-recombining regions therefore have genome-wide

impacts. Our TE genealogies were based on a fraction of the *Copia* and *Ty3* retrotransposon

copies, due to the necessity to identify the 5'-LTR sequences. However, the 5'-LTR sequences

were sometimes lost in degraded TE copies or could not be distinguished from the 3'-LTR

sequence when the coding sequences were degenerated and could not be oriented. Nevertheless,

considering the very low divergence between the TE copies within the genealogies, the burst

patterns should not be affected by the sub-sampling of TE copies. Moreover, we found

independent TE bursts across multiple *Microbotryum* species, indicating that the sampling of

the TE copies used for the genealogies was good enough to capture accumulation events older

than species divergence. Furthermore, the comparison of LTR sequence divergence within and

between copies indicated that the clusters of similar copies were due mostly due to recent bursts

of transposition (due to increased TE activity or less efficient selection against their insertion),

with only rare conversion events among copies. This analysis thus informs on TE dynamics in

genomes of species with a large range of ages of recombination suppression. Supporting the TE

reservoir hypothesis (Hood *et al.*, 2004; Wei *et al.*, 2020), we found a similar pattern of copy

near identity between the non-recombining regions and the autosomes of the *Copia* and *Ty3*

retroelements and a strong significant positive correlation between TE load in the non-

recombining regions and the autosomes. The TE bursts likely result from relaxed purifying

selection following recombination suppression rather than from increased intrinsic TE activity,

as shown also following polyploidization (Baduel *et al.*, 2019).

We found similar TE contents in pseudo-autosomal regions compared to autosomes,

without preferential accumulation of TE copies at the margin of the non-recombining regions.

TEs have nevertheless accumulated everywhere in genomes following the formation of a non-

recombining region acting as a TE reservoir. TE copies thereby accumulating at the margin of

the non-recombining regions may impact their evolution (Maloisel & Rossignol, 1998; Ben-

Aroya *et al.*, 2004; Yelina *et al.*, 2012; Kent *et al.*, 2017). TEs can trigger chromosomal

rearrangements (Balachandran *et al.*, 2022), and they recruit silencing marks, such as DNA

methylation and heterochromatin formation, that can suppress recombination (Maloisel &

Rossignol, 1998; Ben-Aroya *et al.*, 2004; Yelina *et al.*, 2012, 2015; Bachtrog, 2013; Li *et al.*,

2016; Kent *et al.*, 2017). TEs can also induce deleterious mutations, which can select for

recombination suppression that ensures the sheltering of the genetic load they create (Jay *et al.*,

2022). A DNA fragment evolving recombination cessation that captures fewer deleterious

mutations than the population average should be selected for, but this region will have its genetic

load exposed to selection in the homozygous state more often when it rises in frequency unless

it also captures a permanently heterozygous allele such as the haploid self-incompatibility of a

mating-type allele (Jay *et al.*, 2022). The more often that deleterious mutations segregate in

genomes then the more this effect can act: a given fragment evolving recombination cessation

and capturing a permanently heterozygous allele will have a greater chance of having a higher

fitness differential with the average locus if the distribution of the number of deleterious

mutations in this fragment is wider in the population, resulting in stronger selection (Jay *et al.*,

2022). We found that TEs did not accumulate preferentially in the margin of the non-

recombining regions, in contrast to our prediction based on a recent theoretical model showing

that, in automictic species such as *Microbotryum* fungi, a mating-type locus can shelter

deleterious mutations in its flanking regions (Tezenas *et al.*, 2022). As TE insertions can be

deleterious, one may have expected that selfing would have promoted their purge more easily

farther away from the mating-type locus (Tezenas *et al.*, 2022). Not all TE insertions are

however deleterious and their active spread may counteract the purging effect.

We found that TEs have rapidly accumulated in mating-type chromosomes following

recombination suppression, but TE content reached a plateau relatively abruptly at about 50%

of occupied base pairs after 1.5 MY. The rapid initial accumulation of TEs is expected: under

the lack of recombination, new TE insertions cannot be purged, and each new copy can further

make new copies (Quadrana *et al.*, 2019; Legrand *et al.*, 2019). The deceleration of TE

accumulation after 1.5 MY can be due to several, non-exclusive phenomena: i) selection against

new insertions or for a more efficient control may be stronger when the load becomes higher,

particularly if its cost increases with copy number, or if TE insertions are more deleterious with

time because genes become hemizygous. Indeed, TE insertions in genes would be sheltered in

a heterozygous state early after recombination suppression while there are two copies of each

gene in the diploid (or dikaryotic) state, but importantly not following the loss of function for

one copy or the gene becoming hemizygous, as often occurs in sex chromosomes and in

particular in *Microbotryum* fungi (Bachtrog, 2013; Badouin *et al.*, 2015); ii) extant control

mechanisms against TE proliferation can become more efficient with time by a cumulative

effect, e.g. if methylation accumulates with generations (Teixeira & Colot, 2010), or could even

evolve to become more efficient; iii) new TE insertions nested in more ancient copies can further

contribute to inactivate TEs, as suggested by the decreased proportion of *Helitrons* intact copies

with time since recombination suppression in non-recombining regions; *Helitrons* in particular

often display high proportions of non-autonomous copies (Sweredoski *et al.*, 2008); iv) the

chromatin landscape of non-recombining regions may become less favorable to new insertions,

as for example histone can induce a preferential integration of C*opia* retrotransposons
(Quadrana *et al.*, 2019).

The rate of degeneration for non-recombining regions in *Microbotryum* fungi has been
previously analyzed in terms of non-synonymous substitutions and the frequency of optimal
codons (Carpentier *et al.*, 2022). Purifying selection on non-synonymous substitutions was
lower in non-recombining regions than in autosomes and remained nearly constant over time
(Carpentier *et al.*, 2022). However, the substitutions toward non-optimal codons increased
rapidly following recombination suppression and then decelerated to reach a level below random
codon usage. It is not unexpected to find different temporal dynamics of degeneration for DNA
substitutions and TE insertions, as they correspond to very different mechanisms of evolution.

Conclusion

In conclusion, this study leverages on a unique dataset of 21 independent evolutionary strata of
different ages in closely related species to assess the dynamics of TE accumulation with time
following recombination suppression. We show that TEs have rapidly accumulated following
recombination cessation and that the TE accumulation slowed down abruptly after 1.5 MY. We
further show that some superfamilies repeatedly expanded in independent non-recombining
regions, and in particular *Helitrons* that were not yet abundant before recombination
suppression. *Copia* and *Ty3* superfamilies were also over-represented in non-recombining
regions and have accumulated through bursts of proliferation, both in the non-recombining
regions of the mating-type chromosomes and in the autosomes of *Microbotryum* species at the
same time; this finding supports the TE reservoir hypothesis, i.e., that the TEs accumulated in
non-recombining regions have a genome-wide impact by transposing to recombining regions.

This study thus sheds light on important processes to improve our knowledge on genome

evolution and in particular on the consequences of recombination suppression.

Material and methods

Detection and annotation of transposable elements

Transposable elements (TEs) were detected *de novo* in the a_1 and a_2 haploid genomes of 16

Microbotryum species and the red yeast *Rhodothorula babjavae* using LTRharvest (Ellinghaus

et al., 2008) from GenomeTools 1.5.10, performing long-terminal repeat (LTR)

retrotransposons detection and RepeatModeler 1.0.11 (Smit & Hubley, 2008) combining results

from three other programs: RECON (Bao & Eddy, 2002), RepeatScout (Price *et al.*, 2005) and

Tandem Repeats Finder (Benson, 1999). The TE detection was enriched by BLASTn 2.6.0+

(Altschul *et al.*, 1990) using the genomes as a database and the previously detected TE models

as queries. To fulfill the repeat criterion, a TE sequence detected by RepeatModeler or

LTRHarvest and its BLAST hits were retained only if the query matched at least three sequences

in the same species with an identity ≥ 0.8, a sequence length > 100 bp and a coverage ≥ 0.8

(defined as the query alignment length with removed gaps divided by the query length). When

these criteria were met, we retained the other query matches for the following parameters:

identity ≥ 0.8, sequence length > 100 bp, e-value $\leq 5.3e-33$ (25th percentile of e-value

distribution) and coverage ≥ 0.8.

We performed TE annotation (Wicker *et al.*, 2007) using the fungal Repbase database

23.05 (Bao *et al.*, 2015) based on sequence similarity. We first performed a BLASTn search

and a BLASTx search using the fungal Repbase DNA and protein sequence database,

respectively, with TE sequences as queries. For each of these similarity-based searches, we set

the minimum e-value score at 1e−10 and minimum identity at 0.8. Then, we performed protein

domain detection using pfam_scan.pl (ftp://ftp.ebi.ac.uk/pub/databases/Pfam/Tools/) on TE

sequences and compared them to protein domain detection in the fungal Repbase database,

keeping matches with e-values ≤ 1e−5. We also considered the annotation found by

RepeatModeler. In order to determine the best annotation of a TE sequence, we applied a

majority rule: we chose the most frequent annotation, and, in case of equality, we kept both

annotations, generating a multiple annotation; these multiple annotations can be due to TEs

being inserted within other TE copies, generating nested TEs. When we found no annotation,

we assigned the TE sequence to an "unclassified" category; we discarded such annotations when

overlapping with predicted genes to avoid false positives. We also discarded predicted TE

sequences annotated as rRNA or overlapping *Microbotryum* ribosomal sequences (downloaded

from https://www.arb-silva.de/ and http://combio.pl/rrna/taxId/5272/) and mitochondrial DNA

sequences (NC_020353) with e-value ≤ 1e-10. We used a python script for annotation, and the

TE   detection   and   annotation   pipeline   is   available   at

https://gitlab.com/marine.c.duhamel/microtep.

Phylogenomic species tree reconstruction

626   After removing the genes overlapping with annotated TEs, we inferred a species phylogenomic

627   tree from the coding sequences of the 3,669 single-copy genes present in all *Microbotryum*

628   species and the outgroup *Rhodothorula babjavae*, as done previously (Duhamel *et al.*, 2022).

Evolutionary histories of *Copia* and *Ty3* superfamily retroelements

631   We inferred the evolutionary histories of the most abundant elements, *i.e.,* in the *Copia* and *Ty3*

632   retroelement superfamilies, by reconstructing their genealogies based on the sequence of their

633   long terminal repeat (LTR) region at the 5' end of the elements (5'-LTR), the rationale being

that, at the time of transposition, the LTR sequences of the new copy are identical to the 5'-LTR sequence of its progenitor (Zhang *et al.*, 2014). We only retained sequences harboring one or two LTR sequences identified by LTRHarvest (Ellinghaus *et al.*, 2008), thus avoiding nested copies. We distinguished the 5'- from the 3'-LTR based on the orientation of their coding sequences detected using RepeatProteinMask (Smit & Hubley, 2008). Because some retrotransposon sequences were degenerated, their coding sequences could not be identified and thus oriented, so that only a subset of the retrotransposons were used for building the genealogies. We built multi-species genealogies using the TE copies of *Copia* and *Ty3* retrotransposons in all genomes (2,468 and 694 copies, respectively). We built single-species (*i.e.*, within genome) genealogies, only for species with at least 100 copies remaining after filtering (see proportion of TE copies used for the genealogies in Supplementary Figure S8). We then aligned the 5'-LTR sequences of *Copia* or *Ty3* elements using MAFFT v7.310 (Katoh *et al.*, 2002) (alignment length ranging from 816 to 2,767 bp, 1,678 bp on average). We used IQ-TREE 2.0.4 (Minh *et al.*, 2020) to infer the maximum likelihood tree from the 5'-LTR sequences alignment under the GAMMA + GTR model of substitution. We assessed tree robustness with 1000 replicates for ultrafast bootstrap and SH-aLRT. Trees were plotted using phytools (Revell, 2012) and ggtree (Yu *et al.*, 2017). Trees were displayed as rooted using midpoint rooting (*midpoint.root*, phytools, (Revell, 2012). Pairwise sequence divergence values in trees were obtained using *cophenetic.phylo* from ape 5.0 (Paradis & Schliep, 2019).

We wanted to assess whether the identified clusters of TE copies with low divergence to each other originated from bursts of transposition or conversion events among copies. In case of gene conversion, multiple copies would be similar while their age of insertion would be ancient, if an old copy had copied itself by gene conversion into several other copies. In contrast, in case of recent bursts, all copies should have a young age of insertion in addition to be similar

to each other. Because the 5'-LTR and 3'-LTR sequences are identical at the time of insertion

and then diverge (SanMiguel *et al.*, 1998), we estimated the time since transposition of TE

copies in the putative bursts by computing the number of substitutions between the 5'-LTR and

3'-LTR sequences, in the TE copies for which the 3'-LTR sequence could also be identified.

For clusters of genetically similar copies, we compared the number of substitutions between 5'-

LTR and 3'-LTR sequences per copy to the number of substitutions between the 5'-LTR

sequences of pairs of most similar copies within the same cluster. For both measures, we

computed the gamma-corrected Kimura 2-model parameter (γ-K2P), using MEGA11 (Tamura

et al., 2021), and multiplied the divergence by the number of aligned sites. To compare the two

values, we normalized the number of substitutions between the 5'- and 3'-LTR sequences by

the number of aligned sites in the 5'-LTR sequences alignment. In case of bursts of transposition

without conversion events, the 5'-LTR and 3'-LTR sequences of each copy diverge gradually

and at the same rate as the 5'-LTR sequences between copies from the same burst (Figure 5A).

Plotting the number of substitutions between 5'-LTR sequences of the most similar copies

against the number of substitutions between the 5'-LTR - 3'-LTR sequences of each copy

should therefore give a cloud of points around the first bisector. In case of conversion events,

we expect that old copies (with divergent 5'-LTR and 3'-LTR sequences) have low divergence

between one another while they have a high divergence between their 5'-LTR and 3'-LTR

sequences, and therefore fall at the bottom right of the plot, far from the first bisector (Figure

5A). To assess whether clusters of highly similar copies were caused by conversion events

within clusters, we therefore plotted the number of substitutions between 5'-LTR sequences

between closest copies as a function of the number of substitutions between their 5'- and 3'-

LTR sequences (normalized by the same alignment length), for TE copies within clusters of

genetically similar elements. As the number of substitutions in a sequence are rare discrete

events following a Poisson law (Felsenstein, 2003), we compared for each species this observed

cloud of points to a cloud of points generated by drawing their two coordinates from a Poisson

distribution, with a lambda parameter equal to the observed average number of substitutions

between the 5'- and 3'-LTR sequences within copies, as this value is not affected by potential

conversion events among copies, but only by divergence since insertion. The points outside the

simulated cloud, in the right bottom sector (low number of substitutions between copies but

high number of substitutions within copies) would result from conversion events.

Search for potential horizontal transfers of transposable elements between *Microbotryum*

species and their host plants

To search for potential horizontal transfers of transposable elements between *Microbotryum*

fungi and their host plants, we compared the nucleotide sequences from the Repbase database

23.05 (Bao *et al.*, 2015) of dicotyledones and *Arabidopsis thaliana* with the *Copia* and *Ty3* TE

annotations from the *Microbotryum* genomes using BLASTN 2.6.0+ (Altschul *et al.*, 1990).

Detection of centromeres and telomeres in the autosomes of *Microbotryum* genomes

We identified centromeres (Supplementary Table 7) as in Duhamel *et al.* (2022) by blasting

(BLASTN 2.6.0+, Altschul *et al.*, 1990) the centromeric repeats previously described in

Microbotryum fungi (Badouin *et al.*, 2015) on autosomal contigs larger than 20 kb; the largest

stretch of centromeric repeats on a given contig was annotated as its predicted centromere. We

delimited the predicted centromere by recursively extending the focal region by 1 kb intervals

as long as gene density remained lower than 0.25 in the focal window. We detected telomeres

(Supplementary Table 8) within 100 bp regions at the end of contigs carrying at least five times

the telomere-specific TTAGGG motif (CCCTAA on reverse complementary strand), as done

previously (Badouin *et al.*, 2015). Subtelomeric regions were defined as the 20 kb regions

adjacent to the telomeres, plus the 100 bp telomeres.

RIP-like index

We computed values for a RIP-like index in non-overlapping 1 kb genomic regions along the

mating-type chromosomes and autosomal contigs as previously described (Beckerson *et al.*,

2019). The RIP-like index was calculated as the ratio *t/n*, *t* being the ratio of the RIP-affected

sites (TTG + CAA trinucleotides, forward and reverse; Hood *et al.*, 2005) over the non RIP-

affected sites (TCG + CGA, minus overlapping tetra-nucleotides TGCA) for the RIP target

sites, and *n* being the same ratio for non-RIP target sites ([A,C,G]TG + CA[C,G,T] over

[A,C,G]CG + CG[C,G,T] - [A,C,G]CG[C,G,T]), to control for sequence composition. The RIP

index was normalized as *t/n - 1*, so that a normalized RIP index greater than 0 was indicative of

an excess of RIP-like mutations compared to random expectations. The script used to compute

the RIP index is available at https://gitlab.com/marine.c.duhamel/ripmic. We used the

SlidingWindow function of the *evobiR* package (mean values, 10 kb steps - Blackmon & Adams,

2015) and *ggplot2* (Wickham, 2009) to plot the distribution of RIP along the mating-type

chromosomes and autosomal contigs and identify the RIP-affected regions. We used the same

parameters to plot the density of genes and TEs in the same non-overlapping 1 kb windows. In

order to test the relationship between the abundance of RIP footprint in TEs in non-recombining

regions and the age of evolutionary strata, we calculated the average of the RIP index of the

TEs in each stratum and performed a two-sided Pearson's correlation test.

Estimates of TE content in non-recombining regions and their flanking regions

The distinct evolutionary strata within non-recombining regions have been previously distinguished based on the following criteria: i) the distribution of recombination suppression across the phylogenetic tree to assess when recombination suppression evolved and thereby identifying distinct evolutionary strata, *i.e.*, genomic regions that stopped recombining at different times, and ii) the level of trans-specific polymorphism (*i.e.*, alleles clustering by mating type rather than by species), which is also a strong indicator of the time of recombination suppression (Branco *et al.*, 2017, 2018; Carpentier *et al.*, 2019, 2022; Duhamel *et al.*, 2022) (Supplementary Table 1). Young non-recombining regions, formed after the linkage of the PR and HD loci (*i.e.*, the white and pink strata in *M. v. paradoxa* and the light blue stratum in *M. v. caroliniana*; Figure 1) were not rearranged with other strata, so that it was straightforward to compute their percentages of base pairs occupied by TEs. In contrast, the ancient blue and purple strata (Figure 1), having evolved at the base of the *Microbotryum* clade, were too much rearranged and intermingled with other strata for their specific TEs to be identified in species with large non-recombining regions, but they were very small (*i.e.*, encompassing on average only 12 genes). We therefore pooled the ancient blue and purple strata with the larger strata with which they were intermingled. For the multiple independent evolutionary strata corresponding to the linkage of the HD locus with the PR locus in a single step (black strata; Figure 1), it was straightforward to compute their percentage of base pairs occupied by TEs. In *M. v. viscidula*, *M. v. melanantha*, *M. v. gracilicaulis* and *M. v paradoxa*, recombination suppression extended gradually to link the HD and PR loci and then underwent multiple rearrangements that intermingled their genes and TEs. In order to avoid misassignment of TEs to these intermingled strata, we only considered regions spanning at least 80 kb (corresponding to the size of the smallest stratum from the dataset, the pink stratum in *M. v. paradoxa*) and with

only genes from a single stratum. We defined the limit between two strata as the middle point

between two genes from two distinct rearranged strata. The resulting blocks assigned to strata

overall represented 66.52 to 90.26% of the total non-recombining regions formed by

intermingled strata (Supplementary Figure S15), which should thus give a good estimate of the

stratum TE contents. We used circos (Krzywinski *et al.*, 2009) to visualize the blocks assigned

to strata. For the non-recombining regions linking the HD or PR loci and their respective

centromere (blue-to-centromere and purple-to-centromere respectively, in *M. lagerheimii* and

M. saponariae), the percentage of base pairs occupied by TEs was calculated in the region

between the end of the centromere and the mating-type locus. For each stratum, we took the

mean percentage of TEs across a_1 and a_2 haploid genomes, and for the strata shared across

several species, the mean of the a_1 and a_2 values across all the species sharing the same

recombination suppression event. The ancestral size of the non-recombining regions were

estimated by calculating the average size of the corresponding genomic region in the a_1 and a_2

haploid genomes of *M. intermedium*, *i.e.,* the *Microbotryum* species most distant from the other

species in our dataset, with unlinked mating-type loci and with little recombination suppression

(Duhamel *et al.*, 2022).

To test whether TEs also preferentially accumulate at the margin of the non-recombining

regions, we compared the fraction of base pairs occupied by TEs within the 100 kb of the

pseudo-autosomal regions flanking the non-recombining regions to the distribution of TE load

in 100 kb non-overlapping windows in fully recombining autosomes (100 kb away from the

centromeres and after removal of 20 kb corresponding to subtelomeric regions). We only

considered the species with a length of the pseudo-autosomal region at least equal to 100 kb

(thus excluding *M. v. paradoxa*) and with at least 30 non-overlapping windows of 100 kb on

their autosomes (thus excluding *M. scabiosae* and *M. v. parryi*, for which the genome assemblies

were too much fragmented). We set the window length at 100 kb because this is the DNA

fragment size at which linkage disequilibrium decreases below $r^2 = 0.2$ in *Microbotryum*

populations (Badouin *et al.*, 2017). The distribution of the 100kb-window values on autosomes

were plotted using the Sturges binning method.

We estimated the fraction of intact *Copia*, *Ty3* and *Helitron* copies in each stratum based

on their length. For each stratum, we calculated the ratio of putative intact TE copies (5-7.5 kb

for *Copia* and *Ty3* and 5-11 kb for *Helitron*) over the total number of *Copia* and *Ty3* copies or

Helitron copies. For each stratum, we also calculated the mean fraction of intact TE copies

across a_1 and a_2 haploid genomes and for the strata shared by several species, the mean across

species deriving from the same recombination suppression event.

Estimates of non-recombining region ages

In *Microbotryum* fungi, sexual reproduction occurs before the infection of a new plant, which

will produce spores in its flowers at the next flowering seasons, so that one year corresponds to

one generation for these fungi. We used the dates of recombination suppression estimated in

previous studies (Duhamel *et al.*, 2022; Carpentier *et al.*, 2022), for 21 genomic regions

corresponding to independent events of recombination suppression distributed across 15

Microbotryum species. The *M. scabiosae* non-recombining regions used in a previous study

(Carpentier *et al.*, 2022) was not considered here due to the large confidence interval on its

estimated age. To this dataset of 15 *Microbotryum* species, we added the TE content in the

autosomes of *M. intermedium* as the zero time point; indeed, the autosomes are recombining

and there is little potential for a reservoir effect in *M. intermedium* as the non-recombining

regions are very small, being restricted to close proximity around the separate mating-type loci

(Duhamel *et al.*, 2022); we set the age since recombination suppression at 0.00000003 MY for

M. intermedium autosomes, corresponding to one day since the recombination suppression

event, to avoid the zero value for logarithm computation in temporal dynamics analyses.

Statistical analyses

We tested whether the genomic percentage of base pairs occupied by TEs was different between

the a_1 and a_2 mating-type haploid genomes using a paired two-sided Wicoxon's rank test. To

compare the percentages of base pairs occupied by TEs in the autosomes, the pseudo-autosomal

regions and the non-recombining regions of the mating-type chromosomes, we performed a

variance analysis (ANOVA) after testing the normality of the TE percentage distributions

(Kolmogorov-Smirnov test). We then performed post-hoc Tukey's range tests to identify the

pairs of genomic compartments which displayed significantly different TE loads, using the

TukeyHSD R function providing adjusted p-values for multiple comparisons. We assessed the

correlation between TE loads in autosomes and non-recombining regions using the two-sided

Pearson's test.

In order to assess the shape of the functions for accumulation of TEs in the non-

recombining regions of the *Microbotryum* mating-type chromosomes, we performed

regressions using generalized additive models (*gam* function of R, Wood, 2011) and non-linear

regression models (*drm* function of the drc package, Ritz *et al.*, 2019). Based on the Akaike

information criterion (AIC), we evaluated the model fits for predicting the percentage of base

pairs occupied by TEs in the non-recombining regions as a function of the time since

recombination suppression, with the ancestral size of the non-recombining region as a

covariable, and assuming either a linear or a logarithmic relationship with the time since

recombination suppression, with or without smoothing spline, or a negative exponential model.

In order to assess the relationship between the fraction of intact *Copia*, *Ty3* and *Helitron*

copies and time since recombination suppression, we performed a local regression using the

loess.as function from the fANCOVA package, using the Akaike Information Criterion for

automated parameter selection (Hurvich *et al.*, 1998; Wang & Ji, 2020).

We performed all R analyses in R version 4.2.1.

Supplementary information

829   **Additional file 1.** Supplementary figures S1-S14.

830   **Additional file 2.** Supplementary figures S15-S19.

831   **Additional file 3.** Supplementary tables 1-8.

832   **Additional file 4.** Datasets published on Figshare.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article

(Supplementary Tables 1 - 8). All data generated or analysed during this study are included in

this published article and the supplementary information files. The TE detection pipeline is

available at https://gitlab.com/marine.c.duhamel/microtep. The script used to compute the RIP

index is available at https://gitlab.com/marine.c.duhamel/ripmic.

Competing interests: The authors declare that they have no competing interests.

Funding

This work was supported by the National Institute of Health (NIH) grant R15GM119092 to M.

E. H., the Louis D. Foundation award and the EvolSexChrom ERC advanced grant #832352 to

T. G., and an IDEX Paris-Saclay Bochum University PhD grant to M.D.

Author contributions

M.D., T.G., R.C.R.d.l.V. and M.E.H designed the study. T.G., R.C.R.d.l.V. and M.E.H

supervised the study. M.D. performed the genomic and statistical analyses and produced the

figures. R.C.R.d.l.V. advised in genomic analysis and data visualization. T.G. and M.D. wrote

the original draft. All authors contributed to the manuscript.

Acknowledgements

We thank Dominik Begerow, Sebastian Klenner, Lena Steins and Frederick Witfeld for

providing useful comments on the manuscript. We thank Gilles Fisher, Paul Jay, Jacqui Shykoff

and Olivier Gascuel for useful discussions.

References

Ade C, Roy-Engel AM, Deininger PL. **2013**. *Alu* elements: An intrinsic source of human

genome instability. *Current opinion in virology* **3**: 639–645.

Ahmed S, Cock JM, Pessia E, Luthringer R, Cormier A, Robuchon M, Sterck L, Peters

AF, Dittami SM, Corre E, *et al.* **2014**. A haploid system of sex determination in the brown alga

Ectocarpus sp. *Current biology: CB* **24**: 1945–1957.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. **1990**. Basic local alignment search

tool. *Journal of Molecular Biology* **215**: 403–410.

Amselem J, Lebrun M-H, Quesneville H. **2015**. Whole genome comparative analysis of

transposable elements provides new insight into mechanisms of their inactivation in fungal

genomes. *BMC genomics* **16**: 141.

Andrenacci D, Cavaliere V, Lattanzi G. **2020**. The role of transposable elements activity in

aging and their possible involvement in laminopathic diseases. *Ageing Research Reviews* **57**:

100995.

Antonovics J, Abrams JY. **2004**. Intratetrad mating and the evolution of linkage relationships.

Evolution **58**: 702–709.

Bachtrog D. **2003**. Accumulation of Spock and Worf, two novel non-LTR retrotransposons, on

the neo-Y chromosome of *Drosophila miranda*. *Molecular Biology and Evolution* **20**: 173–181.

Bachtrog D. **2005**. Sex chromosome evolution: Molecular aspects of Y-chromosome

degeneration in Drosophila. *Genome Research* **15**: 1393–1401.

Bachtrog D. **2013**. Y chromosome evolution: emerging insights into processes of Y

chromosome degeneration. *Nature reviews. Genetics* **14**: 113–124.

Bachtrog D, Hom E, Wong KM, Maside X, de Jong P. **2008**. Genomic degradation of a

young Y chromosome in *Drosophila miranda*. *Genome Biology* **9**: R30.

Badouin H, Gladieux P, Gouzy J, Siguenza S, Aguileta G, Snirc A, Prieur SL, Jeziorski C,

Branca A, Giraud T. **2017**. Widespread selective sweeps throughout the genome of model

plant pathogenic fungi and identification of effector candidates. *Molecular Ecology* **26**: 2041–

2062.

Badouin H, Hood ME, Gouzy J, Aguileta G, Siguenza S, Perlin MH, Cuomo CA, Fairhead

C, Branca A, Giraud T. **2015**. Chaos of rearrangements in the mating-type chromosomes of

the anther-smut fungus *Microbotryum lychnidis-dioicae*. *Genetics* **200**: 1275–1284.

Baduel P, Quadrana L, Hunter B, Bomblies K, Colot V. **2019**. Relaxed purifying selection

in autopolyploids drives transposable element over-accumulation which provides variants for

local adaptation. *Nature Communications* **10**: 5818.

Bakkeren G, Jiang G, Warren RL, Butterfield Y, Shin H, Chiu R, Linning R, Schein J,

Lee N, Hu G, *et al.* 2006. Mating factor linkage and genome evolution in basidiomycetous

pathogens of cereals. *Fungal genetics and biology: FG & B* **43**: 655–666.

Balachandran P, Walawalkar IA, Flores JI, Dayton JN, Audano PA, Beck CR. **2022**.

Transposable element-mediated rearrangements are prevalent in human genomes. *bioRxiv*:

2022.06.16.496479.

Bao Z, Eddy SR. **2002**. Automated de novo identification of repeat sequence families in

sequenced genomes. *Genome Research* **12**: 1269–1276.

Bao W, Kojima KK, Kohany O. **2015**. Repbase Update, a database of repetitive elements in

eukaryotic genomes. *Mobile DNA* **6**: 11.

Beckerson WC, Rodríguez de la Vega RC, Hartmann FE, Duhamel M, Giraud T, Perlin

MH. **2019**. Cause and effectors: whole-genome comparisons reveal shared but rapidly evolving

effector sets among host-specific plant-castrating fungi. *mBio* **10**: e02391-19.

Ben-Aroya S, Mieczkowski PA, Petes TD, Kupiec M. **2004**. The compact chromatin structure

of a *Ty* repeated sequence suppresses recombination hotspot activity in *Saccharomyces*

cerevisiae. *Molecular Cell* **15**: 221–231.

Bennetzen JL, Wang H. **2014**. The contributions of transposable elements to the structure,

function, and evolution of plant genomes. *Annual Review of Plant Biology* **65**: 505–530.

Benson G. **1999**. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*

Research **27**: 573–580.

Bergero R, Charlesworth D. **2009**. The evolution of restricted recombination in sex

chromosomes. *Trends in Ecology & Evolution* **24**: 94–102.

Bewick AJ, Hofmeister BT, Powers RA, Mondo SJ, Grigoriev IV, James TY, Stajich JE,

Schmitz RJ. **2019**. Diversity of cytosine methylation across the fungal tree of life. *Nature*

Ecology & Evolution **3**: 479–490.

Blackmon H, Adams RH. **2015**. EvobiR: Tools for comparative analyses and teaching

evolutionary biology.

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M,

Izsvák Z, Levin HL, Macfarlan TS, *et al.* **2018**. Ten things you should know about

transposable elements. *Genome Biology* **19**: 199.

Branco S, Badouin H, Rodríguez de la Vega RC, Gouzy J, Carpentier F, Aguileta G,

Siguenza S, Brandenburg J-T, Coelho MA, Hood ME, *et al.* **2017**. Evolutionary strata on

young mating-type chromosomes despite the lack of sexual antagonism. *Proceedings of the*

National Academy of Sciences of the United States of America **114**: 7067–7072.

Branco S, Carpentier F, Rodríguez de la Vega RC, Badouin H, Snirc A, Le Prieur S,

Coelho MA, de Vienne DM, Hartmann FE, Begerow D, *et al.* **2018**. Multiple convergent

supergene evolution events in mating-type chromosomes. *Nature Communications* **9**.

Burns KH. **2020**. Our Conflict with Transposable Elements and Its Implications for Human

Disease. *Annual Review of Pathology* **15**: 51–70.

Carpentier F, Rodríguez de la Vega RC, Branco S, Snirc A, Coelho MA, Hood ME, Giraud

T. **2019**. Convergent recombination cessation between mating-type genes and centromeres in

selfing anther-smut fungi. *Genome Research* **29**: 944–953.

Carpentier F, Rodríguez de la Vega RC, Jay P, Duhamel M, Shykoff JA, Perlin MH,

Wallen RM, Hood ME, Giraud T. **2022**. Tempo of degeneration across independently evolved

non-recombining regions. *Molecular Biology and Evolution*: msac060.

Castanera R, López-Varas L, Borgognone A, LaButti K, Lapidus A, Schmutz J,

Grimwood J, Pérez G, Pisabarro AG, Grigoriev IV, *et al.*2016. Transposable elements

versus the fungal genome: Impact on whole-genome architecture and transcriptional profiles.

PLOS Genetics **12**: e1006108.

Charlesworth D. **2017**. Evolution of recombination rates between sex chromosomes.

Philosophical Transactions of the Royal Society B: Biological Sciences **372**: 20160456.

Charlesworth D. **2021**. The timing of genetic degeneration of sex chromosomes. *Philosophical*

Transactions of the Royal Society B: Biological Sciences **376**: 20200093.

Charlesworth B, Campos JL. **2014**. The relations between recombination rate and patterns of

molecular variation and evolution in *Drosophila*. *Annual Review of Genetics* **48**: 383–403.

Charlesworth B, Charlesworth D. **2000**. The degeneration of Y chromosomes. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences **355**: 1563–1572.

Choi JY, Lee YCG. **2020**. Double-edged sword: The evolutionary consequences of the

epigenetic silencing of transposable elements. *PLoS Genetics* **16**: e1008872.

Dallery J-F, Lapalu N, Zampounis A, Pigné S, Luyten I, Amselem J, Wittenberg AHJ,

Zhou S, de Queiroz MV, Robin GP, *et al.*2017. Gapless genome assembly of *Colletotrichum*

higginsianum reveals chromosome structure and association of transposable elements with

secondary metabolite gene clusters. *BMC Genomics* **18**: 667.

Day AW. **1979**. Mating type and morphogenesis in *Ustilago violacea*. *Botanical Gazette* **140**:

94–101.

Deniz Ö, Frost JM, Branco MR. **2019**. Regulation of transposable elements by DNA

modifications. *Nature Reviews. Genetics* **20**: 417–431.

Dolgin ES, Charlesworth B. **2008**. The effects of recombination rate on the distribution and

abundance of transposable elements. *Genetics* **178**: 2169–2177.

Duhamel M, Carpentier F, Begerow D, Hood ME, Rodríguez de la Vega RC, Giraud T.

2022. Onset and stepwise extensions of recombination suppression are common in mating-type

chromosomes of *Microbotryum* anther-smut fungi. *Journal of Evolutionary Biology*.

Ellinghaus D, Kurtz S, Willhoeft U. **2008**. LTRharvest, an efficient and flexible software for

de novo detection of LTR retrotransposons. *BMC bioinformatics* **9**: 18.

Erlandsson R, Wilson JF, Pääbo S. **2000**. Sex chromosomal transposable element

accumulation and male-driven substitutional evolution in humans. *Molecular Biology and*

Evolution **17**: 804–812.

Felsenstein J. **2003**. *Inferring Phylogenies*. Sunderland, Mass: Sinauer.

Fontanillas E, Hood ME, Badouin H, Petit E, Barbe V, Gouzy J, de Vienne DM, Aguileta

G, Poulain J, Wincker P, *et al.*2015**. Degeneration of the non-recombining regions in the

mating-type chromosomes of the anther-smut fungi. *Molecular Biology and Evolution* **32**: 928–

943.

Galagan JE, Selker EU. **2004**. RIP: the evolutionary cost of genome defense. *Trends in*

Genetics **20**: 417–423.

Giraud T, Yockteng R, López-Villavicencio M, Refrégier G, Hood ME. **2008**. Mating

system of the anther smut fungus *Microbotryum violaceum*: selfing under heterothallism.

Eukaryotic Cell **7**: 765–775.

Gladyshev E. **2017**. Repeat-induced point mutation and other genome defense mechanisms in

fungi. *Microbiology Spectrum* **5**.

González LG, Deyholos MK. **2012**. Identification, characterization and distribution of

transposable elements in the flax (*Linum usitatissimum L.*) genome. *BMC Genomics* **13**: 644.

Grabundzija I, Messing SA, Thomas J, Cosby RL, Bilic I, Miskey C, Gogol-Döring A,

Kapitonov V, Diem T, Dalda A, *et al.* 2016. A *Helitron* transposon reconstructed from bats
reveals a novel mechanism of genome shuffling in eukaryotes. *Nature Communications* **7**:
10716.

Han K, Lee J, Meyer TJ, Remedios P, Goodwin L, Batzer MA. **2008**. L1 recombination-
associated deletions generate human genomic variation. *Proceedings of the National Academy*
of Sciences of the United States of America **105**: 19366–19371.

Hancks DC, Kazazian HH. **2016**. Roles for retrotransposon insertions in human disease.
Mobile DNA **7**: 9.

Handley L-JL, Ceplitis H, Ellegren H. **2004**. Evolutionary strata on the chicken Z
chromosome: implications for sex chromosome evolution. *Genetics* **167**: 367–376.

Hartmann FE, Duhamel M, Carpentier F, Hood ME, Foulongne-Oriol M, Silar P,
Malagnac F, Grognet P, Giraud T. **2021**. Recombination suppression and evolutionary strata
around mating-type loci in fungi: documenting patterns and understanding evolutionary and
mechanistic causes. *The New Phytologist* **229**: 2470–2491.

Hartmann FE, Rodríguez de la Vega RC, Gladieux P, Ma W-J, Hood ME, Giraud T. **2020**.
Higher gene flow in sex-related chromosomes than in autosomes during fungal divergence.
Molecular Biology and Evolution **37**: 668–682.

Hof AE van't, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, Hall N, Darby
AC, Saccheri IJ. **2016**. The industrial melanism mutation in British peppered moths is a
transposable element. *Nature* **534**: 102–105.

Hood ME, Antonovics J. **2000**. Intratetrad mating, heterozygosity, and the maintenance of
deleterious alleles in *Microbotryum violaceum* (=*Ustilago violacea*). *Heredity* **85 Pt 3**: 231–
241.

Hood ME, Antonovics J, Koskella B. **2004**. Shared forces of sex chromosome evolution in

haploid-mating and diploid-mating organisms. *Genetics* **168**: 141–146.

Hood ME, Katawczik M, Giraud T. **2005**. Repeat-induced point mutation and the population

structure of transposable elements in *Microbotryum violaceum*. *Genetics* **170**: 1081–1089.

Horns F, Petit E, Hood ME. **2017**. Massive expansion of gypsy-like retrotransposons in

Microbotryum fungi. *Genome Biology and Evolution* **9**: 363–371.

Horns F, Petit E, Yockteng R, Hood ME. **2012**. Patterns of repeat-induced point mutation in

transposable elements of basidiomycete fungi. *Genome Biology and Evolution* **4**: 240–247.

Hua-Van A, Le Rouzic A, Maisonhaute C, Capy P. **2005**. Abundance, distribution and

dynamics of retrotransposable elements and transposons: similarities and differences.

Cytogenetic and Genome Research **110**: 426–440.

Hurvich CM, Simonoff JS, Tsai C-L. **1998**. Smoothing parameter selection in nonparametric

regression using an improved Akaike information criterion. *Journal of the Royal Statistical*

Society: Series B (Statistical Methodology) **60**: 271–293.

Idnurm A, Hood ME, Johannesson H, Giraud T. **2015**. Contrasted patterns in mating-type

chromosomes in fungi: hotspots versus coldspots of recombination. *Fungal Biology Reviews*

29: 220–229.

Ironside JE. **2010**. No amicable divorce? Challenging the notion that sexual antagonism drives

sex chromosome evolution. *BioEssays* **32**: 718–726.

Jay P, Tezenas E, Véber A, Giraud T. **2022**. Sheltering of deleterious mutations explains the

stepwise extension of recombination suppression on sex chromosomes and other supergenes.

PLOS Biology **20**: e3001698.

Johnson LJ, Giraud T, Anderson R, Hood ME. **2010**. The impact of genome defense on

mobile elements in *Microbotryum*. *Genetica* **138**: 313–319.

Jones PA. **2012**. Functions of DNA methylation: islands, start sites, gene bodies and beyond.

Nature Reviews Genetics **13**: 484–492.

Kaessmann H, Vinckenbosch N, Long M. **2009**. RNA-based gene duplication: mechanistic

and evolutionary insights. *Nature Reviews Genetics* **10**: 19–31.

Kapitonov VV, Jurka J. **2001**. Rolling-circle transposons in eukaryotes. *Proceedings of the*

National Academy of Sciences of the United States of America **98**: 8714–8719.

Katoh K, Misawa K, Kuma K, Miyata T. **2002**. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.

Kent TV, Uzunović J, Wright SI. **2017**. Coevolution between transposable elements and

recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**:

20160458.

Kronstad JW, Staben C. **1997**. Mating type in filamentous fungi. *Annual Review of Genetics*

31: 245–276.

Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra

MA. **2009**. Circos: an information aesthetic for comparative genomics. *Genome Research* **19**:

1639–1645.

Lahn BT, Page DC. **1999**. Four evolutionary strata on the human X chromosome. *Science*

(New York, N.Y.) **286**: 964–967.

Lanciano S, Mirouze M. **2018**. Transposable elements: all mobile, all different, some stress

responsive, some adaptive? *Current Opinion in Genetics & Development* **49**: 106–114.

Legrand S, Caron T, Maumus F, Schvartzman S, Quadrana L, Durand E, Gallina S,

Pauwels M, Mazoyer C, Huyghe L, *et al.* **2019**. Differential retention of transposable element-

derived sequences in outcrossing *Arabidopsis* genomes. *Mobile DNA* **10**: 30.

Lengeler KB, Fox DS, Fraser JA, Allen A, Forrester K, Dietrich FS, Heitman J. **2002**.

Mating-type locus of *Cryptococcus neoformans*: a step in the evolution of sex chromosomes.

Eukaryotic Cell **1**: 704–718.

Li S-F, Zhang G-J, Yuan J-H, Deng C-L, Gao W-J. **2016**. Repetitive sequences and epigenetic modification: inseparable partners play important roles in the evolution of plant sex chromosomes. *Planta* **243**: 1083–1095.

Lim JK, Simmons MJ. **1994**. Gross chromosome rearrangements mediated by transposable elements in *Drosophila melanogaster*. *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology* **16**: 269–275.

Maloisel L, Rossignol JL. **1998**. Suppression of crossing-over by DNA methylation in Ascobolus. *Genes & Development* **12**: 1381–1389.

Marais G, Nicolas M, Bergero R, Chambrier P, Kejnovsky E, Monéger F, Hobza R, Widmer A, Charlesworth D. **2008**. Evidence for degeneration of the Y chromosome in the dioecious plant *Silene latifolia*. *Current biology : CB* **18**: 545–9.

Marsano RM, Dimitri P. **2022**. Constitutive heterochromatin in eukaryotic genomes: A mine of transposable elements. *Cells* **11**: 761.

Martin A, Troadec C, Boualem A, Rajab M, Fernandez R, Morin H, Pitrat M, Dogimont C, Bendahmane A. **2009**. A transposon-induced epigenetic change leads to sex determination in melon. *Nature* **461**: 1135–1138.

Mawaribuchi S, Takahashi S, Wada M, Uno Y, Matsuda Y, Kondo M, Fukui A, Takamatsu N, Taira M, Ito M. **2017**. Sex chromosome differentiation and the W- and Z-specific loci in *Xenopus laevis*. *Developmental Biology* **426**: 393–400.

Menkis A, Jacobson DJ, Gustafsson T, Johannesson H. **2008**. The mating-type chromosome in the filamentous Ascomycete *Neurospora tetrasperma* represents a model for early evolution of sex chromosomes. *PLoS Genetics* **4**.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. **2020**. IQ-TREE 2: New models and efficient methods for phylogenetic inference

in the genomic era. *Molecular Biology and Evolution* **37**: 1530–1534.

Muller HJ. **1918**. Genetic variability, twin hybrids and constant hybrids, in a case of balanced

lethal factors. *Genetics* **3**: 422–499.

Nguyen A, Wang W, Chong E, Chatla K, Bachtrog D. **2022**. Transposable element

accumulation drives size differences among polymorphic Y Chromosomes in Drosophila.

Genome Research: gr.275996.121.

Paradis E, Schliep K. **2019**. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics (Oxford, England)* **35**: 526–528.

Payer LM, Burns KH. **2019**. Transposable elements in human genetic disease. *Nature*

Reviews. Genetics **20**: 760–772.

Peng JC, Karpen GH. **2008**. Epigenetic regulation of heterochromatic DNA stability. *Current*

opinion in genetics & development **18**: 204.

Peona V, Palacios-Gimenez OM, Blommaert J, Liu J, Haryoko T, Jønsson KA, Irestedt

M, Zhou Q, Jern P, Suh A. **2021**. The avian W chromosome is a refugium for endogenous

retroviruses with likely effects on female-biased mutational load and genetic incompatibilities.

Philosophical Transactions of the Royal Society B: Biological Sciences **376**: 20200186.

Piegu B, Guyot R, Picault N, Roulin A, Sanyal A, Saniyal A, Kim H, Collura K, Brar DS,

Jackson S, *et al.*2006. Doubling genome size without polyploidization: dynamics of

retrotransposition-driven genomic expansions in *Oryza australiensis*, a wild relative of rice.

Genome Research **16**: 1262–1269.

Plohl M, Meštrović N, Mravinac B. **2014**. Centromere identity from the DNA point of view.

Chromosoma **123**: 313–325.

Price AL, Jones NC, Pevzner PA. **2005**. De novo identification of repeat families in large

genomes. *Bioinformatics (Oxford, England)* **21 Suppl 1**: i351-358.

Quadrana L, Etcheverry M, Gilly A, Caillieux E, Madoui M-A, Guy J, Bortolini Silveira A, Engelen S, Baillet V, Wincker P, *et al.* 2019. Transposition favors the generation of large effect mutations that may facilitate rapid adaptation. *Nature Communications* **10**: 3421.

Raffaele S, Kamoun S. 2012. Genome evolution in filamentous plant pathogens: why bigger can be better. *Nature Reviews Microbiology* **10**: 417–430.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things): *phytools: R package*. *Methods in Ecology and Evolution* **3**: 217–223.

Rifkin JL, Hnatovska S, Yuan M, Sacchi BM, Choudhury BI, Gong Y, Rastas P, Barrett SCH, Wright SI. 2022. Recombination landscape dimorphism and sex chromosome evolution in the dioecious plant *Rumex hastatulus*. *Philosophical Transactions of the Royal Society B*.

Ritz C, Jensen SM, Gerhard D, Streibig JC. 2019. *Dose-Response Analysis Using R*. New York.

Saito K, Siomi MC. 2010. Small RNA-mediated quiescence of transposable elements in animals. *Developmental Cell* **19**: 687–697.

SanMiguel P, Gaut BS, Tikhonov A, Nakajima Y, Bennetzen JL. 1998. The paleontology of intergene retrotransposons of maize. *Nature Genetics* **20**: 43–45.

Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J, Fulton L, Graves TA, *et al.* 2009. The B73 maize genome: complexity, diversity, and dynamics. *Science (New York, N.Y.)* **326**: 1112–1115.

Śliwińska EB, Martyka R, Tryjanowski P. 2016. Evolutionary interaction between W/Y chromosome and transposable elements. *Genetica* **144**: 267–278.

Smit AFA, Hubley R. 2008. RepeatModeler Open-1.0.

Sweredoski M, DeRose-Wilson L, Gaut BS. 2008. A comparative computational analysis of

nonautonomous *Helitron* elements between maize and rice. *BMC Genomics* **9**: 467.

Tamura K, Stecher G, Kumar S. **2021**. MEGA11: Molecular Evolutionary Genetics Analysis

Version 11. *Molecular Biology and Evolution* **38**: 3022–3027.

Teixeira FK, Colot V. **2010**. Repeat elements and the Arabidopsis DNA methylation

landscape. *Heredity* **105**: 14–23.

Tezenas E, Giraud T, Véber A, Billiard S. **2022**. The fate of recessive deleterious or

overdominant mutations near mating-type loci under partial selfing. *bioRxiv*:

2022.10.07.511119.

Vittorelli N, Snirc A, Levert E, Gautier V, Lalanne C, Filippo ED, Vega RCR de la,

Gladieux P, Guillou S, Zhang Y, *et al.* **2022**. Stepwise recombination suppression around the

mating-type locus in the fungus *Schizothecium tetrasporum* (Ascomycota, Sordariales).

bioRxiv: 2022.07.20.500756.

Wang X, Ji X. **2020**. fANCOVA: Nonparametric Analysis of Covariance.

Wei K, Aldaimalani R, Mai D, Zinshteyn D, Prv S, Blumenstiel JP, Kelleher ES, Brooks

E. **2022**. Rethinking the "*gypsy*" retrotransposon: A roadmap for community-driven

reconsideration of problematic gene names.

Wei KH-C, Gibilisco L, Bachtrog D. **2020**. Epigenetic conflict on a degenerating Y

chromosome increases mutational burden in Drosophila males. *Nature Communications* **11**:

5537.

Whittle CA, Sun Y, Johannesson H. **2011**. Degeneration in codon usage within the region of

suppressed recombination in the mating-type chromosomes of *Neurospora tetrasperma*.

Eukaryotic Cell **10**: 594–603.

Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P,

Morgante M, Panaud O, *et al.* **2007**. A unified classification system for eukaryotic

transposable elements. *Nature Reviews Genetics* **8**: 973–982.

Wickham H. **2009**. *ggplot2: Elegant graphics for data analysis*. New York: Springer-Verlag.

Wood SN. **2011**. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **73**: 3–36.

Wright SI, Agrawal N, Bureau TE. **2003**. Effects of recombination rate and gene density on transposable element distributions in Arabidopsis thaliana. *Genome Research* **13**: 1897–1903.

Yelina NE, Choi K, Chelysheva L, Macaulay M, de Snoo B, Wijnker E, Miller N, Drouaud J, Grelon M, Copenhaver GP, *et al.***2012**. Epigenetic remodeling of meiotic crossover frequency in Arabidopsis thaliana DNA methyltransferase mutants. *PLoS Genetics* **8**: e1002844.

Yelina NE, Lambing C, Hardcastle TJ, Zhao X, Santos B, Henderson IR. **2015**. DNA methylation epigenetically silences crossover hot spots and controls chromosomal domains of meiotic recombination in Arabidopsis. *Genes & Development* **29**: 2183–2202.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. **2017**. ggtree: an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* **8**: 28–36.

Zemach A, McDaniel IE, Silva P, Zilberman D. **2010**. Genome-wide evolutionary analysis of eukaryotic DNA methylation. *Science (New York, N.Y.)* **328**: 916–919.

Zhang L, Yan L, Jiang J, Wang Y, Jiang Y, Yan T, Cao Y. **2014**. The structure and retrotransposition mechanism of LTR-retrotransposons in the asexual yeast *Candida albicans*. *Virulence* **5**: 655–664.

Zhang J, Yu C, Krishnaswamy L, Peterson T. **2011**. Transposable elements as catalysts for chromosome rearrangements. *Methods in molecular biology (Clifton, N.J.)* **701**.

Figures

 Figure 1: Multiple evolutionary strata formed by independent events of mating-type locus

 linkage and stepwise extension of recombination suppression in *Microbotryum* fungi. A)

 The mating-type chromosomes of *Microbotryum* species with linked mating-type loci have had

 many rearrangements but are depicted here according to the gene order at the moment of the

 fusion between the homeodomain (HD) and pheromone-receptor (PR) chromosomes, the fusion

 site being indicated by a dashed red line. The ancestral gene order was inferred from

 Microbotryum lagerheimii and *M. intermedium* which have unlinked mating-type loci and

 collinear mating-type chromosomes. The old blue and purple strata have evolved before

 diversification of the *Microbotryum* clade. In *M. saponariae* and *M. lagerheimii*, recombination

 cessation linked each mating-type locus to their respective centromeres, forming the HD- and

PR-to-centromere strata. The black strata correspond to the events of recombination suppression having linked the two mating-type loci, which occurred independently at least nine times in *Microbotryum*, indicated by red dots on the phylogeny. In some species (*M. violaceum viscidula*, *M. v. melanantha*, *M. v. gracilicaulis)*, recombination between the mating-type loci was gradually suppressed following the chromosomal fusion event, and formed several strata called black, light gray and dark gray strata. In *M. v. paradoxa*, recombination was partially restored by introgression before being suppressed again, forming the brown stratum. In many species, recombination cessation further extended beyond the mating-type loci, capturing different sets of genes and forming strata that were named as indicated in the figure key. **B)** Average (a_1-a_2) size of the mating-type chromosomes in *Microbotryum* fungi. The average (a_1-a_2) size of the pseudo-autosomal region, non-recombining region and centromere is represented in gray, black and yellow, respectively.

1203

Figure 2: Transposable element (TE) composition of *Microbotryum* genomes. A) Whole-

genome TE load (as percentage of occupied base pairs) across the *Microbotryum* phylogeny,

each color representing a TE category. Numeric values correspond to the a_1-a_2 average. **B)**

Relative proportions of the various TE categories in the different genomic compartments of

Microbotryum genomes. Each color represents a TE category, as in A). **C)** Distribution of the

length of *Copia* and *Ty3* retroelements in the autosomes, the pseudo-autosomal regions and the

non-recombining regions of *Microbotryum* fungi, in red, green and black, respectively. **D)** TE

load per genomic compartment, *i.e.*, autosomes (without the non-recombining centromeres),

pseudo-autosomal regions and non-recombining regions for the main TE categories (*Copia,*

Ty3, Helitron, terminal inverted repeats - TIR). The lines connect the values in the same species,

to facilitate comparisons. Each color represents a *Microbotryum* species.

Figure 3: Density of transposable elements (TEs), genes and repeat-induced point mutation (RIP) index along the mating-type chromosome and the autosomes of *Microbotryum lychnidis-dioicae*. Density of TEs, genes and RIP-like index were calculated in 20 kb non-overlapping windows and plotted along the mating-type chromosomes and the largest autosomes, using the same X scale. A RIP-like index greater than zero indicates that the region is affected by a mechanism leading to C to T transitions resembling RIP. Predicted centromeres are indicated in yellow. Recombining regions of the mating-type chromosomes (pseudo-autosomal regions) and an autosome (the largest autosomal contig in the genome harboring a predicted centromere) are highlighted in pale green. The non-recombining region is delimited by red lines. Pheromone-receptor (PR) and homeodomain (HD) loci are indicated by purple and

1226   blue diamonds, respectively. Identified telomeres are indicated by brown triangles. The plots

1227   for the other species are shown in Figure S5.

A) **M. lychnidis-dioicae**
on *Silene latifolia*

0.5

B) **M. v. paradoxa**
on *Silene paradoxa*

0.5

Genomic location

- Non-recombining regions
  of the mating-type chromosomes
- Pseudo-autosomal
  regions
- Autosomes
- Centromeres

1228

Figure 4: Single-species genealogies of *Copia* retroelement copies in *Microbotryum*

genomes. Single-species genealogies of *Copia* copies in *Microbotryum lychnidis-dioicae* **(A)**

and *Microbotryum violaceum paradoxa* **(B)**. The color of the dots at the tip of the branches

corresponds to the genomic location of the transposable element (TE) copies. Other single-

species *Copia* and *Ty3* genealogies are shown in Supplementary Figures S10 and S11,

respectively.

1235

1236

Figure 5: Investigation of the possible occurrence of conversion events within clusters of highly similar LTR-retrotransposon copies. (A) Illustration of LTR (long-terminal repeat) sequence divergence processes for retrotransposons: at the time of a new LTR-retrotransposon insertion, the 5'-LTR and 3'-LTR sequences are identical within the progeny copy and identical to the 5'-LTR of the parental copy (1). The LTR sequences then gradually diverge because they accumulate distinct mutations (2). The divergence between the 5'-LTR sequences of the two copies (N1) should thus be equal to the divergence between the 5'- and 3'-LTR sequences of the progeny copy (N2), the two measures being proportional to the age of the new copy. Following a conversion event, the 5'-LTR sequences between copies will be identical (3a) while

the divergence between the 5'- and 3'-LTR sequences within copies remain high (3b). **(B)** and

(C) Comparison of the observed number of different substitutions, between LTR sequences

within (N2) and between copies (N1), in color, to the expected distribution under a Poisson

model, in gray, with a lambda parameter equal to the average number of substitutions between

5'- and 3'-LTR sequences within copies, for the *Copia* retroelements in *Microbotryum*

lychnidis-dioicae **(B)** and *M. violaceum paradoxa* **(C)**. If no gene conversion occurred, we

expect a cloud of points around the first bisector (y=x, dashed line) when plotting the number

of substitutions between 5'-LTR sequences of the pairs of most similar copies within putative

bursts against their 5' - 3'-LTR sequence number of substitutions. In case of gene conversion,

we expect points at the bottom right (low divergence between copies but old transposition ages).

The points on the top left likely correspond to copies for which the closest related copy could

not be identified. The colors of the points match the genomic location of the copy pairs. The

observed distributions of the number of substitutions between 5'-LTR sequences of most similar

copies and between 5' - 3' LTR sequences within copies are shown at the top and at the right,

respectively.

Genomic location (a)

- ■ Non-recombining regions of the mating-type chromosomes
- ■ Pseudo-autosomal regions)
- ■ Centromeres
- ■ Autosomes

Species (branch color)

- *M. lychnidis-dioicae* on *Silene latifolia*
- *M. silenes-dioicae* on *Silene dioica*
- *M. coronariae* on *Silene flos-cuculi*
- *M. v. viscidula* on *Silene viscidula*
- *M. v. melanantha* on *Silene melanantha*
- *M. v. gracilicaulis* on *Silene gracilicaulis*
- *M. silenes-acaulis* on *Silene acaulis*
- *M. v. paradoxa* on *Silene paradoxa*
- *M. saponariae* on *Saponaria officinalis*
- *M. v. lateriflora* on *Moehringia lateriflora*
- *M. lagerheimii* on *Silene vulgaris*
- *M. v. tatarinowii* on *Silene tatarinowii*
- *M. v. caroliniana* on *Silene caroliniana*
- *M. v. parryi* on *Silene parryi*
- *M. scabiosae* on *Knautia arvensis*
- *M. intermedium* on *Salvia pratensis*

Recombination suppression event group (b)

Figure 6: Multi-species genealogies of *Copia* retroelement copies in *Microbotryum*

genomes. Multi-species *Copia* tree of all *Microbotryum* species of this study. The branch color

corresponds to the species, the first inner track (a) corresponds to the genomic location of the

TE copies, the second outer track (b) corresponds to the linkage event group of the species

carrying the TE copy. Brackets highlight bursts of TEs. Multi-species *Ty3* tree of all

Microbotryum species is shown in Supplementary Figure S14. Multi-species *Copia* and *Ty3*

genealogies with branch lengths are shown in Supplementary Figure S1.

1270

1271

Figure 7: Tempo of transposable element (TE) accumulation in the non-recombining regions of the *Microbotryum* mating-type chromosomes. Percentage of base pairs occupied by TEs in non-recombining regions (means in a_1 and a_2 genomes and means across species sharing the same strata) as a function of their age in million years (MY) with confidence intervals. The red curve shows the prediction of the best model corresponding to a negative exponential model. Each dot corresponds to an independent evolutionary stratum.

Supplementary figure S1: Percentage of base pairs occupied by transposable elements (TEs) in *Microbotryum* genomes, both for a1 and a2 mating-type chromosomes. Whole-genome TE content across the *Microbotryum* phylogeny, for each mating-type genome, each color representing a TE category.

1283

Supplementary figure S2: Relationship between the percentage of base pairs occupied by TEs and the total size of non-recombining regions. Each color represents one species, one point per mating type.

1287

Supplementary figure S3: Distribution of transposable element (TE) copy length in

Microbotryum **genomes. A)** Distribution of the *Copia* and *Ty3* retrotransposon copy length in

the different genomic compartments in various *Microbotryum* species. Note that the Y axis is

interrupted by parallel bars. **B)** Distribution of the length of individual TE copies for each TE

annotation, across all *Microbotryum* genomes. Each color corresponds to a specific annotation.

The dashed line indicates the 5 kb expected mean length of *Copia* and *Ty3* retroelements.

Similar annotations representing a small proportion of the *Microbotryum* genomes were pooled

in all the other figures; e.g., *ClassII:TIR* with two inverted repeats corresponds to all the *TIR*

elements, i.e., *CACTA, Harbinger, hAT, MuDR, MuLE* and *Tc1-Mariner*.

1297

1298 **Supplementary Figure S4: Transposable element (TE) proportions in the different**

1299 **genomic compartments of *Microbotryum* fungi.** Total TE load per genomic compartment,

1300   i.e., autosomes (without the centromeres), pseudo-autosomal regions and non-recombining

1301   regions. The lines connect the values in the same species to facilitate comparisons. Each color

1302   represents a *Microbotryum* species.

A)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

1303

B)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

C)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- Transposable elements (TEs)
- Genes
- Repeat-induced point mutation (RIP) index

Key to symbols
- PR locus
- HD locus
- Telomere

Key to boundaries
- Non-recombining region (center) / pseudo-autosomal region (flanking)
- Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- Predicted centromere
- Recombining region

D)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- ■ Transposable elements (TEs)
- ■ Genes
- ■ Repeat-induced point mutation (RIP) index

Key to symbols
- ◆ PR locus
- ◆ HD locus
- ▶ Telomere

Key to boundaries
- | Non-recombining region (center) / pseudo-autosomal region (flanking)
- ┊ Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- ■ Predicted centromere
- ■ Recombining region

1306

E)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

MvSilra-01299-0D_C031

MvSilra-01299-0T_C001

Key to lines
- Transposable elements (TEs)
- Genes
- Repeat-induced point mutation (RIP) index

Key to symbols
- PR locus
- HD locus
- Telomere

Key to boundaries
- Non-recombining region (center) / pseudo-autosomal region (flanking)
- Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- Predicted centromere
- Recombining region

1307

F) M. silenes-acaulis on Silene acaulis

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- ■ Transposable elements (TEs)
- ■ Genes
- ■ Repeat-induced point mutation (RIP) index

Key to symbols
- ◆ PR locus
- ◆ HD locus
- ▶ Telomere

Key to boundaries
- | Non-recombining region (center) / pseudo-autosomal region (flanking)
- ┊ Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- ■ Predicted centromere
- ■ Recombining region

1308

G)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
■ Transposable elements (TEs)
■ Genes
■ Repeat-induced point mutation (RIP) index

Key to symbols
◆ PR locus
◆ HD locus

Key to boundaries
| Non-recombining region (center) / pseudo-autosomal region (flanking)
⁞ Contig junction (not fully assembled mating-type chromosome)

Key to boxes
■ Predicted centromere
■ Recombining region
■ Young pink stratum
□ Young white stratum

1309

I)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- ■ Transposable elements (TEs)
- ■ Genes
- ■ Repeat-induced point mutation (RIP) index

Key to symbols
- ◆ PR locus
- ◆ HD locus
- ▶ Telomere

Key to boundaries
- | Non-recombining region (center) / pseudo-autosomal region (flanking)
- ┆ Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- ■ Predicted centromere
- ■ Recombining region
- ■ Old purple stratum
- ■ Old blue stratum

K)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

L)

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

MvCa-1250-A1-R1_MC01-1

MvCa-1250-A2-R1_MC01

1314

M) *M. v. parryi* *
on *Silene parryi*

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- ■ (blue) Transposable elements (TEs)
- ■ (green) Genes
- ■ (black) Repeat-induced point mutation (RIP) index

Key to symbols
- ◆ (purple) PR locus
- ◆ (blue) HD locus
- ▶ (red) Telomere

Key to boundaries
- | (red) Non-recombining region (center) / pseudo-autosomal region (flanking)
- ⁞ (black) Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- ■ (yellow) Predicted centromere
- ■ (green) Recombining region

1315

N) **M. scabiosae** on *Knautia arvensis*

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- ■ Transposable elements (TEs)
- ■ Genes
- ■ Repeat-induced point mutation (RIP) index

Key to symbols
- ◆ PR locus
- ◆ HD locus
- ▶ Telomere

Key to boundaries
- | Non-recombining region (center) / pseudo-autosomal region (flanking)
- ⋮ Contig junction (not fully assembled mating-type chromosome)

Key to boxes
- ■ Predicted centromere
- ■ Recombining region

1316

O) M. intermedium on Salvia pratensis

Mating-type chromosomes

Largest autosomal contigs with predicted centromere

Key to lines
- Transposable elements (TEs)
- Genes
- Repeat-induced point mutation (RIP) index

Key to symbols
- PR locus
- HD locus
- Telomere

Key to boundaries
- Non-recombining region (center) / pseudo-autosomal region (flanking)

Key to boxes
- Predicted centromere
- Old purple stratum
- Old blue stratum
- Recombining region

1317

Supplementary Figure S5: Density of transposable elements (TEs), genes and repeat-

induced point mutations (RIP) along the mating-type chromosomes and the autosomes of

***Microbotryum* genomes.** Density of TEs, genes and RIP index were calculated in 20 kb non-

overlapping windows. A RIP index greater than one indicates that the region is RIP-affected.

Predicted centromeres are indicated in yellow. Recombining regions of the mating-type

chromosomes (pseudo-autosomal regions) and the autosomes (largest autosomal contig

harboring a predicted centromere in each genome) are highlighted in pale green. Non-

recombining regions are delimited by red lines. Non-rearranged evolutionary strata are

displayed in their corresponding color (see legend). Non-fully assembled mating-type

chromosomes have been joined and the junctions are indicated by black dashed lines.

Pheromone-receptor (PR) and homeodomain (HD) loci are indicated by purple and blue

diamonds, respectively. Identified telomeres are indicated by brown triangles.

1330

Supplementary Figure S6: Transposable element (TE) load at the margin of the non-recombining regions compared to the distribution in recombining autosomal regions. The average values of TE content in the regions flanking the non-recombining region are indicated by red lines and the ones in autosomal windows constitute the grey distribution. The probability of finding a higher TE content in a 100 kb window on autosomes than in the 100 kb region directly flanking the non-recombining region ($P[TE_{wAUT}>TE_{NRRf}]$) is reported in the top right corner of each panel. The panels correspond to different species, and there are four or eight red lines (individual flanking regions of non-recombining regions) per species: one a_1 and one a_2 genome per species, and for each haploid genome one or two mating-type chromosomes, so four or eight flanking regions. Three species are not represented because their pseudo-autosomal regions were smaller than 100 kb (*Microbotryum violaceum paradoxa*) or their autosome assemblies were too fragmented (*M. v. parryi* and *M. scabiosae*).

1343

1344 **Supplementary Figure S7: Correlation between the proportion of base pairs occupied by**

1345 **transposable elements (TEs) in the non-recombining regions and the autosomes.**

Supplementary Figure S8: Relative percentage and number of transposable element (TE) sequences per genomic compartment used for the genealogies based on their 5'-LTR sequences, for *Copia* (A) and *Ty3* (B) elements. The total number of 5'-LTR sequences available per species is indicated in brackets under the species name. Single-species genealogies were only built for species with grey filled rectangles around their names.

Genomic location

- Non-recombining regions of the mating-type chromosomes
- Pseudo-autosomal regions
- Autosomes
- Centromeres

1352

C) **M. v. viscidula** on *Silene viscidula*

0.5

D) **M. v. melanantha** on *Silene melanantha*

0.5

Genomic location

- ● Non-recombining regions of the mating-type chromosomes
- ● Pseudo-autosomal regions
- ● Autosomes
- ● Centromeres

1353

E) **M. v. gracilicaulis** on *Silene gracilicaulis*

0.5

F) **M. silenes-acaulis** on *Silene acaulis*

0.5

Genomic location

● Non-recombining regions of the mating-type chromosomes
● Pseudo-autosomal regions
● Autosomes
○ Centromeres

1354

G) **M. v. caroliniana** on *Silene caroliniana*

H) **M. v. parryi** on *Silene parryi*

0.5

Genomic location

● Non-recombining regions of the mating-type chromosomes ● Pseudo-autosomal regions ● Autosomes ○ Centromeres

Supplementary Figure S9: Single-species genealogies of *Copia* retroelement copies in

1357   ***Microbotryum* genomes.** Single-species genealogies of *Copia* copies in *Microbotryum silenes-*

1358   *dioicae* **(A)**, *M. coronariae* **(B)**, *M. violaceum viscidula* **(C)**, *M. v. melanantha* **(D)**, *M. v.*

1359   *gracilicaulis* **(E)**, *M. silenes-acaulis* **(F)**, *M. v. caroliniana* **(G)** and *M. v. parryi* **(H)**. The color

1360   of the dots at the tip of the branches corresponds to the genomic location of the transposable

1361   element (TE) copies.

Genomic location

- ● Non-recombining regions
  of the mating-type chromosomes
- ● Pseudo-autosomal
  regions
- ● Autosomes
- ○ Centromeres

1362

1363 **Supplementary Figure S10: Single-species genealogies of *Ty3* retroelement copies in**

1364   ***Microbotryum* genomes.** Single-species genealogies of *Ty3* copies in *Microbotryum silenes-*

1365   *dioicae* **(A)** and *M. v. paradoxa* **(B)**. The color of the dots at the tip of the branches corresponds

1366   to the genomic location of the transposable element (TE) copies.

Supplementary Figure S11: Investigation of the possible occurrence of conversion events within clusters of highly similar LTR-retrotransposon copies in *Microbotryum* fungi. We looked for footprints of conversion events as explained on Figure 5 and in the method section, for the *Copia* (**A**) and *Ty3* (**B**) retrotransposon copies of *Microbotryum silenes-dioicae*, for the *Copia* retrotransposons of *M. coronariae* (**C**), *M. violaceum viscidula* (**D**), *M. v. melanantha* (**E**), *M. v. gracilicaulis* (**F**), *M. silenes-acaulis* (**G**), for the *Ty3* retrotransposon copies of *M. v. paradoxa* (**H**), and for the *Copia* retrotransposon copies of *M. v. caroliniana* (**I**) and *M. v. parryi*

(J). In colors are shown the number of substitutions between the 5'-LTR sequences of pairs of

most similar copies (N2) within clusters of highly similar retroelement copies as a function of

the number of substitutions between the 5'- and 3'-LTR sequences within copies (N1). The

Poisson simulated numbers of substitutions (with lambda equal to the average number of

substitutions between the 5'- and 3'-LTR sequences within copies) are plotted in gray. The

colors of the points indicate the genomic location of the copy pairs. The observed distributions

of the number of substitutions between 5'-LTR sequences and between 5' - 3' LTR sequences

are shown at the top and at the right of each scatterplot.

1384

Supplementary Figure S12: Distribution of pairwise divergence for retroelements in the

non-recombining regions of the mating-type chromosomes and the autosomes in each

species, plotted separately for the different independent events of recombination suppression.

(A) *Copia* retrotransposons, **(B)** *Ty3* retrotransposons.

Supplementary Figure S13: Percentage of pairs of most closely related long-terminal

repeat (LTR) sequences with at least one copy in the non-recombining regions compared

to their representation in their transposable element (TE) genealogy. Each colored point

corresponds to a TE genealogy (*Copia* or *Ty3*) of a *Microbotryum* species (Figure 4 and

Supplementary Figure S9 and S10), according to the species color code from Figure 1. From

left to right are represented the percentages of pairs of most closely related copies i) involving

one copy from the non-recombining regions and from another location (*e.g.*, the autosomes), ii)

with both copies in the non-recombining regions, and the percentage of copies located in the

non-recombining regions, in their TE genealogy.

1399

Supplementary Figure S14: Multi-species genealogies of *Ty3* retroelement copies in

***Microbotryum* genomes.** Multi-species *Ty3* trees of all *Microbotryum* species analysed in this

study. The branch color corresponds to the species, the first inner track (a) corresponds to the

genomic location of the transposable element (TE) copies, the second outer track (b)

corresponds to the linkage event group of the species carrying the TE copy. Brackets highlight

bursts of TEs.

A)

a

1406

B)

a

Species
(branch color)

Recombination
suppression
event group (a)

M. lychnidis-dioicae
on *Silene latifolia*

M. silenes-dioicae
on *Silene dioica*

M. coronariae
on *Silene flos-cuculi*

M. v. viscidula
on *Silene viscidula*

M. v. melanantha
on *Silene melanantha*

M. v. gracilicaulis
on *Silene gracilicaulis*

M. silenes-acaulis
on *Silene acaulis*

M. v. paradoxa
on *Silene paradoxa*

M. saponariae
on *Saponaria officinalis*

M. v. lateriflora
on *Moehringia lateriflora*

M. lagerheimii
on *Silene vulgaris*

M. v. tatarinowii
on *Silene tatarinowii*

M. v. caroliniana
on *Silene caroliniana*

M. v. parryi
on *Silene parryi*

M. scabiosae
on *Knautia arvensis*

M. intermedium
on *Salvia pratensis*

1407

 Supplementary Figure S15: Multi-species genealogies of retroelement copies in

1409 *Microbotryum* **genomes.** The phylogenies of *Copia* **(A)** and *Ty3* **(B)** are the same as those

1410 displayed on Figure 5 and Supplementary Figure S14, respectively, but here plotted with branch

1411 lengths. The branch color corresponds to the species, the track (a) corresponds to the linkage

1412 event group of the species carrying the TE copy. The dashed circles are a help to visualize the

1413 timing of the bursts. The green dashed circle highlights the *Copia* bursts which are almost

1414 synchronized in the clade of *M. v. gracilicaulis*, pointed by green arrows.

1415

Supplementary Figure S16: Delimitation of blocks for the rearranged evolutionary strata

in mating-type chromosomes. The PR, HD and pheromone genes are represented by purple,

blue and small light purple circles, respectively. The outer track represents contigs, with length

ticks every 200 kb and centromeres indicated in yellow. Non-recombining regions are displayed

in dark grey and pseudo-autosomal regions in light gray. The young, non-rearranged white and

pink strata of *M. v. paradoxa* are highlighted on the outer track. Genes from the different strata

are displayed by bars on the inner track (see Figure 1 for the stratum formation history). Large

blocks (greater than 80 kb) of genes from a single stratum are indicated on the inner track by

rectangles. The light grey stratum is displayed in green to facilitate visualization. The

percentage of non-recombining regions retained in the blocks of strata are indicated next to the

mating type, on each side of the circos plots. Blue and orange lines link alleles, the latter

corresponding to inversions. The link width is proportional to the corresponding gene length.

Supplementary Figure S17: Relationship between synonymous divergence (d_S), transposable element (TE) load and stratum age in *Microbotryum* mating-type chromosomes. A) Percentage of base pairs occupied by TEs in non-recombining regions (means in a_1 and a_2 genomes and means across species sharing the same strata) as a function of the mean d_S of the genes present in the stratum. The red curve shows the prediction of the best model corresponding to a negative exponential model. **B)** Relationship between the stratum age in million years (MY) and the mean d_S of the genes contained in this stratum. Each dot corresponds to an independent evolutionary stratum in A) and B).

Supplementary Figure S18: Relationship between the abundance of transposable element (TE) inactivation footprints by repeat-induced point mutation (RIP) in the non-recombining regions of the *Microbotryum* mating-type chromosomes and time since recombination suppression in million years (MY). Mean normalized RIP index on TEs in non-recombining regions as a function of their age with confidence intervals. The red dashed line separates the RIP index values suggesting RIP mutations (> 0) or no RIP activity (≤ 0). The two-sided Pearson correlation test was not significant (*p-value* = 0.9).

Supplementary Figure S19: Relationship between the fraction of intact *Copia, Ty3* and

***Helitron* copies and stratum age in *Microbotryum* mating-type chromosomes.** Fraction of

intact *Copia* and *Ty3* copies (5-7.5 kb for intact copies) **(A)** and *Helitron* copies (5-11 kb for

intact copies) **(B)** in non-recombining regions (means in a_1 and a_2 genomes and means across

species sharing the same strata) as a function of the time since recombination suppression in

million years (MY). Each dot corresponds to an independent evolutionary stratum in A) and B).

The red curve shows the smooth curve obtained by local regression.

1454