Title: Sex and *APOE* Genotype Influence Respiratory Function Under Hypoxic and Hypoxic-Hypercapnic Conditions

Authors: Chase E. Taylor[1,2], Laura E. Mendenhall[2], Michael D. Sunshine[2], Jessica N. Wilson[1,2], Chris M. Calulot[2], Ramon C. Sun[3,4], Lance A. Johnson[5,6], Warren J. Alilain*[1,2]

Affiliations: [1]Department of Neuroscience, University of Kentucky; [2]Spinal Cord and Brain Injury Research Center, University of Kentucky; [3]Department of Biochemistry & Molecular Biology, College of Medicine, University of Florida, Department of Biochemistry, University of Florida; [4]Center for Advanced Spatial Biomolecule Research, University of Florida; [5]Department of Physiology, University of Kentucky; [6]Sanders-Brown Center on Aging, University of Kentucky

Corresponding author email*: warren.alilain@uky.edu

CET, LEM, and MDS contributed equally to manuscript

Journal: Journal of Neurophysiology

Submission date: 29 June 2023

ABSTRACT

The apolipoprotein *(APOE)* gene has been studied due to its influence on Alzheimer's disease (AD)

development and work in an *APOE* mouse model recently demonstrated impaired respiratory motor plasticity

following spinal cord injury (SCI). Individuals with AD often co-present with obstructive sleep apnea (OSA)

characterized by cessations in breathing during sleep. Despite the prominence of *APOE* genotype and sex as

factors in AD progression, little is known about the impact of these variables on respiratory control. Ventilation

is tightly regulated across many systems, with respiratory rhythm formation occurring in the brainstem but

modulated in response to chemoreception. Alterations within these modulatory systems may result in

disruptions of appropriate respiratory control and ultimately, disease. Using mice expressing two different

humanized *APOE* alleles, we characterized how sex and the presence of *APOE3* or *APOE4* influences

ventilation during baseline breathing (normoxia) and during respiratory challenge. We show that sex and *APOE*

genotype influence breathing during hypoxic challenge, which may have clinical implications in the context of

AD and OSA. Additionally, female mice, while responding robustly to hypoxia, were unable to recover to

baseline respiratory levels, emphasizing sex differences in disordered breathing.

INTRODUCTION

Apolipoprotein E (ApoE) has 3 distinct alleles: *APOE2*, *E3*, and *E4* each coding for a different lipid transport

protein variant [1]. The *APOE4* allele is a key genetic determinant for late-onset Alzheimer's disease (AD); it is

estimated that heterozygous carriage of the *APOE4* allele results in a 4-fold increased risk of developing AD,

while homozygous carriers exhibit a 15 fold increase [2]. Approximately one in five people are carriers of the

APOE4 allele. Critically, in the context of our work, the *E4* Alzheimer's connection has been linked to the

presentation of cognitive decline and inflammation.

Breathing, the rhythmic inspiratory and expiratory patterns that facilitate gas exchange in the lungs, is essential

to life and conditions impacting respiratory function and neural respiratory control are critical in the context of

aging related disease [3,4,5-7]. Obstructive sleep apnea (OSA) is a disease state which is characterized by

cessations in breathing leading to periods of hypoxia [8]. Repeated exposure to hypoxia leads to

neuroinflammation and cognitive decline which can magnify the impacts of AD [9].

Considering the importance of *APOE4* as the key genetic determinant for AD and hypoxia as a major driver of

cognitive decline, it is critical to understand the impact and relationship of the *APOE4* allele on neural

respiratory control and the body's innate response to hypoxia, i.e. the hypoxic ventilatory response (HVR) [10].

We tested the hypothesis that *APOE4* negatively impacts breathing, particularly under conditions that would

induce a response to respiratory stress. To test this hypothesis, we utilized whole body plethysmography

(WBP) to measure ventilation in a humanized *APOE* mouse model under normoxia (21% O_2), and two different

gas challenges (hypoxia alone and hypoxic hypercapnia).

METHODS

Animals

All experiments were approved by the Institutional Animal Care and Use Committee at the University of

Kentucky. Mice expressing human *APOE* isoforms under control of the endogenous mouse *APOE* promotor

(targeted replacement mice) were backcrossed for at least 10 generations to the C57BL/6 background [11,12].

Mice were group-housed by sex and genotype on a 12/12 light/dark cycle and fed normal chow diet *ad libitum*.

Mice in cohort 1 (hypoxic challenge; n=33) weighed 36.9-48.6 g with ages ranging from 367-441 days. Mice in

cohort 2 (hypoxic-hypercapnic challenge; n=16) weighed 32.02-52.7 g with ages ranging from 472-477 days.

The average age and weight by sex, genotype, and cohort are listed in Table 1.

Cohort	Sex	Genotype	n	Average weight (g) (SD)	Average age (d) (SD)
Hypoxia challenge	Male	E3	9	48.8 (4.7)	410 (16.6)
		E4	8	45.5 (4.6)	441 (65.3)
	Female	E3	8	36.9 (7.5)	367 (10.1)
		E4	8	37.5 (5.3)	377 (9.0)
Hypoxic-hypercapnia challenge	Male	E3	4	52.7 (2.8)	473 (0)
		E4	4	47.5 (6.0)	472.5 (4.0)
	Female	E3	3	51.2 (5.9)	477 (0)
		E4	5	32.02 (4.2)	476 (3.8)

Table 1. Number, average weight, and average age of mice used in hypoxia and hypoxic-hypercapnia experiments. g = grams; SD = standard deviation; d = days

Hypoxia and hypoxic-hypercapnia experiments

Whole body plethysmography (WBP,1 liter per minute flow) (DSI, Buxco FinePointe) was used to measure the

ventilatory response during hypoxic (11% O_2 balance nitrogen) and hypoxic-hypercapnic (7% CO_2, 10.5% O_2,

balance nitrogen) challenge in adult mice. We calculated respiratory rate (breaths/min), tidal volume

(mL/breath/g), and minute ventilation (mL/min/g) from the WBP flow traces. Whole body plethysmography

recordings lasted approximately 90 minutes and occurred during daylight hours between 0930 and 1600 with

food and water restricted during the 90-minute measurement period. To ensure consistency across all

measures, the WBP system was recalibrated prior to every trial. Immediately prior to baseline readings, a 40-

minute period of in-chamber acclimation occurred to reduce the prevalence of non-ventilatory respiratory

behaviors associated with exploring the chamber during the baseline period. Acclimation was followed by a 30-

minute period of normoxia (21% O_2, balance nitrogen), then 10 minutes of respiratory challenge concluding

with a 5- or 10-minute period of normoxia. The challenge design can be seen in Figure 1a and the number of

animals per group is shown in Table 1. Following trials, mice were returned to their home cage in the vivarium.

WBP chambers were thoroughly cleaned and dried between each group.

Data analysis and statistical analysis

The normality and variance of the data was determined to ensure appropriate assumptions were met. If data

within a particular experiment did not meet these assumptions non-parametric tests were used. Specific

statistical tests for each figure are listed in the figure legends. Ventilatory measures were calculated in DSI

FinePointe software and exported, breath by breath. The breath by breath data was used to calculate binned

respiratory rate and average tidal volume, these metrics were multiplied to generate minute ventilation. This

analysis and statistical testing was performed using MATLAB (MathWorks, Version 2021b). For hypoxic

ventilatory challenge, hypoxic-hypercapnic ventilatory challenge, and post challenge measures, data were

normalized to percent change relative from baseline. For statistical analysis, an ANOVA (or Kruskal-Wallis for

non-parametric analyses) was used for statistical comparisons between sex and genotype. Tukey-Kramer

post-hoc test was used to test for differences within sex or genotype. Results were considered statistically

significant if $p < 0.05$. Investigators were not blinded to genotype or sex.

Figure 1. ***APOE4*** **animals across sexes have increased tidal volume to maintain respiratory homeostasis under normoxic conditions while maintaining the same respiratory rate as *E3* mice. a)** Timeline of WBP hypoxic and hypoxic hypercapnic challenges. **b)** Plethysmography airflow traces from a representative *APOE3* (green) or *APOE4* (gray) mouse during baseline (normoxia). **c)** Quantification of plethysmography during baseline (normoxia) in five-minute bins, testing for an overall effect of genotype prior to assessing sex differences (Two-way ANOVA mean ± SEM; n= 25 (*APOE4*, 13F/12M), n=24 (*APOE3*, 11F/13M)).

RESULTS

Both genotype and sex impact normoxic breathing with female mice showing the greatest tidal volume

To determine differences in baseline breathing, *APOE3* and *APOE4* male and female mice underwent a 25-

minute period of normoxia prior to a 10-minute respiratory challenge. Representative flow traces from an

APOE3 and *E4* mouse (Figure 1B) illustrate larger tidal volume during baseline (normoxia) in the *APOE4*

mouse. Quantification of these data showed an effect of genotype with *E4* mice having a larger tidal volume

throughout the normoxic baseline (Figure 1C). Representative flow traces were shown from each sex and

genotype (Figure 2A). Sex differences were present in tidal volume and minute ventilation when corrected for

body weight (Figure 2B-D). Female mice had to breath at a deeper and greater weight corrected tidal volume

relative to male mice to maintain baseline breathing ($p<0.002$, $F_{(1)} = 11.96$). Additionally, female mice had a

larger weight corrected minute ventilation ($p<0.03$, $F_{(1)} = 5.05$). These baseline measures show a sex-

dependent difference present that may affect how male and female mice will respond to a ventilatory

challenge. Finally, *E4* mice had larger weight corrected tidal volume compared to *E3* mice ($p<0.03$, $F_{(1)} = 5.50$).

Figure 2: Female *APOE* mice breathe deeper than males to establish normoxic ventilatory patterns under stress-free conditions. a) Representative airflow traces during baseline (normoxia) from *APOE3* and *APOE4* male and female mice. Metrics of ventilation at baseline **b)** tidal volume (V_T (ml)) and weight corrected tidal volume (V_T (ml/g)), **c)** minute ventilation (V_E (ml)) and weight corrected minute ventilation (V_E (ml/g)), and **d)** respiratory rate (RR (bpm)). Male mice exhibit lower tidal volume and minute ventilation in weight corrected values, and *APOE4* animals exhibit greater tidal volume than *E3* counterparts across sexes. (Two-way ANOVA, mean + SD; n = 11 (*APOE3* female), n = 13 (*APOE4* female), n = 13 (*APOE3* male), n = 12 (*APOE4* male)).

Male APOE3 mice exhibit a traditional HVR while female mice and APOE4 males fail to mount a complete
response

The hypoxic ventilatory response is traditionally described by an increase in minute ventilation and tidal volume

that peaks and is maintained for the duration of hypoxic challenge while frequency increases at first and

followed by a slight decrease prior to leveling out at a rate higher than baseline frequency [13]. To investigate

sex- and genotype-dependent respiratory differences, we challenged the animals to a 10-minute hypoxic (11%

O_2) ventilatory challenge. Figure 3a shows airflow traces from animals undergoing hypoxic ventilatory

challenge. Airflow data from all animals in the hypoxia cohort is quantified in Figure 3b, showing the hypoxic

challenge period, split into two 5-minute bins. Within the first five minutes of hypoxia exposure, there is a clear

effect of sex in respiratory rate and minute ventilation. During this initial period of hypoxia, female mice had a

reduced respiratory rate relative to baseline and male mice had an elevated rate ($p < 0.002$, $F_{(1)} = 12.82$).

During the final five minutes of the challenge, the respiratory rate of male mice returned towards baseline,

while the respiratory rate of female mice declined further ($p < 0.02$, $F_{(1)} = 6.57$). All groups increased their tidal

volume relative to baseline in the first five minutes of the hypoxic challenge. However, male *APOE4* mice had a

less robust response compared to their *APOE3* counterparts ($p < 0.05$, Tukey-Kramer post-hoc), supporting

previous work showing that male *APOE4* mice have impaired ventilatory response under hypoxic conditions

after SCI [14]. All groups increased their minute ventilation during the first 5 minutes, however, only the *APOE3*

male mice were able to maintain an elevated minute ventilation over the entire challenge period. Interestingly,

female mice remained at or below their baseline minute ventilation during the latter portion of the challenge

while *E4* male mice could not maintain tidal volume increases in response to hypoxia.

Response and recovery from hypoxia is impaired in female APOE3 and APOE4 mice.

Post-hypoxia airflow traces show a return towards baseline level (Figure 3C). Following cessation of hypoxic

challenge, there is a female genotype minute ventilation difference in the second 5-minute period ($p<0.04$

Figure 3D). When considering tidal and respiratory rate, female animals show a decline relative to baseline,

with greater deficit seen in *APOE4* female mice. Interestingly, the female *E4* mice do have a slight initial tidal

volume increase relative to baseline following the hypoxic challenge followed by a plummet below baseline

levels and the levels of their female *E3* counterparts.

Figure 3: Male *APOE3* mice demonstrate a traditional HVR stronger than *APOE4* males and return to baseline levels following challenge while female mice fail to increase respiratory rate and *APOE4* females plummet to sub-baseline levels following challenge. a) Representative airflow traces during and after hypoxic ventilatory challenge. **b)** Comparison graphs between sex and genotype during hypoxic. **c)** Post-challenge airflow traces. **d)** Ventilatory metrics during the post-challenge normoxic period. Initial studies conducted in male mice did not include 5-10 minute post challenge period. Asterisks indicate $p<0.05$, Tukey-Kramer post-hoc. Two-way ANOVA, mean + SD; n = 8 (*APOE3* female), n = 8 (*APOE4* female), n = 9 (*APOE3* male), n = 8 (*APOE4* male).

During hypoxic hypercapnic challenge APOE4 animals are able to robustly increase respiratory output relative
144

to APOE3 animals regardless of sex although sex plays a role during the post challenge period
145

As only the *APOE3* male group was able to produce and maintain a pronounced increase in minute ventilation
146

during hypoxic challenge, we wanted to test if the other groups were limited in their ability to produce increased
147

minute ventilation or if hypoxia alone simply did not trigger an increase. To accomplish this, in a separate
148

cohort of animals, following the normoxic baseline period we challenged utilizing a hypoxic-hypercapnic (10.5%
149

O_2, 7% CO_2) mixture. As the male *APOE4* animals during hypoxia did not increase their tidal volume to the
150

same degree as the male *APOE3* animals, we were specifically interested in the possibility of a male *APOE4*
151

ceiling effect (i.e., inability to generate larger tidal volume). Figure 4a shows a robust hypoxic-hypercapnic
152

ventilatory response (HHVR) across all mice, regardless of sex or genotype. This suggests that there was not
153

a ceiling effect within the *APOE4* male mouse hypoxia cohort and increases across all measures could be
154

solicited via a more severe respiratory challenge indicative of impaired chemoreception in male *APOE4* mice.
155

Figure 4b shows the entirety of the hypoxic-hypercapnic challenge and the corresponding response. The
156

APOE3 mice of both sexes had a lower respiratory rate compared to *APOE4* counterparts during the first five
157

minutes ($p < 0.04$, $F_{(1)} = 5.50$) and the final five minutes ($p < 0.04$, $F_{(1)} = 5.61$). Minute ventilation mirrors this
158

effect throughout hypoxic-hypercapnic challenge, driven by genotypic differences in both respiratory rate and
159

tidal volume, indicating that *E3* mice have lowered respiratory capacity relative to *E4* mice during hypoxic-
160

hypercapnic challenge.
161

Airflow traces from the period after the hypoxic-hypercapnic challenge (Figure 4C) show a differential recovery
162

based on sex and genotype. The male mice increased their respiratory rate in the first five-minutes after the
163

challenge, while the female mice maintained the same respiratory rate as during the challenge. The tidal
164

volume of all *E3* mice and *E4* female mice returned towards baseline, while the tidal volume of the *E4* male
165

mice remained elevated.
166

The post-HHVR (Figure 4D) male mice have a higher tidal volume than female mice across both genotypes in
167

the first ($p < 0.03$, $F_{(1)} = 6.93$) 5 minutes with a trend in the second ($p < 0.08$, $F_{(1)} = 3.88$) 5 minutes. This data may
168

be indicative of impaired respiratory control in the male mice, as they should return toward baseline levels
169

following exposure to hypoxic-hypercapnic conditions.
170

Figure 4: Under hypoxic hypercapnic conditions, *APOE4* animals respond robustly to challenge however *APOE4* males remain elevated relative to baseline following challenge. a) Flow traces show a robust increase in tidal volume during challenge. **b)** Comparison of ventilatory metrics during hypoxic-hypercapnic challenge, considering for effect of sex and genotype. **c)** Representative airflow traces during post-HHVR challenge. **d)** Respiratory metrics during the post-HHVR period. Two-way ANOVA, mean + SD; n = 3 (*APOE3* female), n = 5 (*APOE4* female), n = 4 (*APOE3* male), n = 4 (*APOE4* male).

Normoxia	Baseline Measures			
Sex effects	• Females have higher weight corrected **tidal volume** • Females have higher weight corrected **minute ventilation**.			
Genotype effects	• E4 animals have higher weigh corrected **tidal volume**.			
Interaction effects	none			
Hypoxia	**First 5 minutes**	**Second 5 minutes**	**Post-hypoxia first 5 minutes**	**Post-hypoxia second 5 minutes**
Sex effects	• Males have increased **respiratory rate** • Males have increased **minute ventilation**	• Males have increased **respiratory rate** • Males have increased **minute ventilation**	none	none
Genotype effects	none	none	none	• E4 females have depressed **minute ventilation** relative to *E3* females
Interaction effects	• E3 Males have increased tidal volume compared to *E3* **females** and *E4* **males**	• E4 females have increased tidal volume compared to *E3* **females** and *E4* **males**	none	none
Hypoxic-Hypercapnia	**First 5 minutes**	**Second 5 minutes**	**Post-hypoxia-hypercapnia first 5 minutes**	**Post-hypoxia-hypercapnia second 5 minutes**
Sex effects	none	none	• Males have increased **tidal volume**	none
Genotype effects	• *E4* animals have increased **tidal volume, respiratory rate** and **minute ventilation**	• *E4* animals have increased **tidal volume, respiratory rate** and **minute ventilation**	• *E4* animals have increased **tidal volume**	none
Interaction effects	none	none	none	none

Table 2. Summary of significant results across baseline measures and challenges.

DISCUSSION

This study builds on previous research elucidating the effects of genetic influences and sex on respiratory

control [14]. Breathing is a tightly regulated system that incorporates both feedback and feedforward control,

however genetic disorders or predispositions can affect respiratory regulation [3,4,5-7]. In prior work completed by

our lab, female mice expressing the *E3* allele maintained diaphragmatic activity following intermittent hypoxia

while *E4* females had significantly depressed activity. Unlike the female mice, males did not exhibit a difference

in diaphragmatic activity based on genotype in our prior work. The results collected in this study continue to

demonstrate that genotype and sex in humanized *APOE* mice plays a role in regulating breathing, specifically

under conditions of respiratory stress. A summary of the individual changes can be referenced in Table 2.

Severe OSA results in frequent periods of hypoxia which have been shown to increase neuroinflammation and

impair cognition [15,16,17]. In recent years, scientists have continued to address the impact of sex and age

differences on OSA development [18]. In addition to obstructive sleep apnea, approximately 1% of the United

States population exhibits central sleep apnea resulting from impairments in central chemoreception [19]. Severe

respiratory impairments, including congenital central hypoventilation syndrome leading to fatal apneas, can

result from genetic mutations [20,21]. It follows that genetic polymorphisms may play a critical role in the onset

and severity of OSA pathology and symptoms [22,23].

Genotype and sex differences seen during hypoxia are reversed or eliminated when hypercapnia is introduced

While *APOE* genotype has been evaluated within sleep disordered breathing across broad datasets in human

subjects, the research investigating animal models of *APOE* under respiratory stress has been limited [24]. To

model an adverse stimulus effect on respiratory pattern between *E3* and *E4* mice, we utilized hypoxic

conditions eliciting the hypoxic ventilatory response investigating tidal volume, respiratory rate, and minute

ventilation.

When exposed to hypoxia, our *APOE4* male mice were unable to maintain their increased frequency or tidal

volume in response to this stimulus. These findings may have important implications in the context of

obstructive sleep apnea for male *APOE4* carriers. While our *APOE4* female mice do generate a hypoxic

ventilatory response specifically when considering tidal volume upon completion of the challenge they

experience a massive drop in minute ventilation relative to *E3* female counterparts and well below their

baseline minute ventilation. These changes point to either differences in respiratory responsiveness or reduced

plasticity when encountering changes in pO2 which would result in an inability of *APOE4* carriers across sexes

to adapt to periods of hypoxia during OSA [25,26,27]. Indeed, the *E4* male carrier hypoxia response deficit may

actually limit their ability to develop one possible compensatory mechanism of plasticity to overcome OSA

impairments known as respiratory -long term facilitation (LTF) [28]. This same interpretation can be applied to

female *APOE4* mice who generate a robust HVR but at too great a cost.

While limited research asserts that *APOE4* may not contribute to OSA and vice versa our research has

continued to show that the ability to attenuate a hypoxic insult such as that seen in OSA is linked to the *APOE4*

genotype [29,30]. As stated prior, hypoxia and the inability to compensate for it has been clearly shown to drive

neuroinflammation and cognitive impairment. Chemoreceptor function under hypoxia is susceptible to genetic

alterations demonstrated in carotid bodies, although models of humanized *APOE* mice have not yet been

utilized [31].

Interestingly, our hypoxic hypercapnic ventilatory challenge data shows that *E4* animals can indeed increase

their respiratory output in a stimulus dependent manner. During hypoxic hypercapnic challenge tidal volume

and minute ventilation increased 50-200% relative to baseline. We show this across both genotypes and sexes

with *E4* animals actually responding more robustly relative to *E3* counterparts. This is particularly important

since it shows that *APOE4* animals have the ability to increase respiratory drive but not under conditions of

hypoxia such as one would primarily experience during OSA. One reason for this may be that hypercapnic

chemoreception is elicited primarily centrally in the retrotrapezoid nucleus while hypoxic chemoreception takes

place peripherally in the carotid and aortic bodies [32,33,34]. It could therefore be surmised that while mechanisms

of peripheral chemoreception may be impaired in *E4* animals, central hypercapnic chemoreception remains

unaffected or highly responsive to changes in pCO_2 content. Since respiratory drive remains intact, pathways

leaving brainstem respiratory centers may also be unaffected by *APOE*. These findings suggest that the

elicitation of pCO2 chemoreceptive drive may represent one possible intervention for OSA individuals

heterozygous or homozygous for *APOE4*.

Sex-dependent responses to hypoxia and hypoxic-hypercapnia have been studied in other animals, although

the most relevant background to our current findings was a study in Fischer rats showing that ventilatory

response differences are age-dependent as opposed to sex-dependent with middle aged and older female rats

having a greater response [35]. Sex differences in respiratory recovery have also been noted previously in

neonatal mouse models where female respiratory rhythm generating centers have shown greater ability to

recover following a period of severe hypoxia [36].

Future research will elucidate *APOE* specific differences in phrenic motor neuron serotonergic signaling [14].

Additionally, future studies should include a cohort of aged matched wild-type male and female controls. The

data presented here serves to fill a critical gap in knowledge as these *APOE* mice are both closer in relative

age and more closely model conditions akin to human sleep apnea physiology [37].

CONCLUSION

This study further clarifies work from our laboratory and others on the effects of *APOE* on respiratory control in

unfavorable respiratory conditions. Our results show that humanized *APOE4* male mice have more difficulty

mounting a response to hypoxic conditions than their *E3* counterparts while female mice of both genotypes

appear to have difficulty mounting any response in terms of respiratory rate and only *E4* females maintain a

respective tidal volume increase. Interestingly, *E4* female mice have prolonged depression following cessation

of hypoxia, not seen in their male counterparts.

In hypoxic-hypercapnic conditions, genotype differences persist in across all measures, however, more

interesting perhaps are the sex differences seen in recovery following hypoxic-hypercapnia. Because sleep

apnea and disordered breathing have been indicated in earlier onset of disease as well as cognitive

impairment this work will serve as a foundation for researching downstream effects of hypoxia and hypoxic-

hypercapnia in genotype and sex allowing for the application of these findings to the heterogenous aging

human population and to elucidating the development of disease [38-42].

REFERENCES

1   Farmer, B. C., Kluemper, J. & Johnson, L. A. Apolipoprotein E4 Alters Astrocyte Fatty Acid Metabolism and Lipid
Droplet Formation. *Cells* **8**, doi:10.3390/cells8020182 (2019).
2   Raber, J., Huang, Y. & Ashford, J. W. ApoE genotype accounts for the vast majority of AD risk and AD pathology.
Neurobiol Aging **25**, 641-650, doi:10.1016/j.neurobiolaging.2003.12.023 (2004).
3   Warren, P. M., Awad, B. I. & Alilain, W. J. Drawing breath without the command of effectors: the control of
respiration following spinal cord injury. *Respir Physiol Neurobiol* **203**, 98-108, doi:10.1016/j.resp.2014.08.005
(2014).
4   Mitchell, G. S. & Babb, T. G. Layers of exercise hyperpnea: modulation and plasticity. *Respir Physiol Neurobiol*
151, 251-266, doi:10.1016/j.resp.2006.02.003 (2006).
5   Barna, B. F., Takakura, A. C. & Moreira, T. S. Acute exercise-induced activation of Phox2b-expressing neurons of
the retrotrapezoid nucleus in rats may involve the hypothalamus. *Neuroscience* **258**, 355-363,
doi:10.1016/j.neuroscience.2013.11.031 (2014).
6   Rourke, K. S., Mayer, C. A. & MacFarlane, P. M. A critical postnatal period of heightened vulnerability to
lipopolysaccharide. *Respir Physiol Neurobiol* **232**, 26-34, doi:10.1016/j.resp.2016.06.003 (2016).
7   Julu, P. O. *et al.* Characterisation of breathing and associated central autonomic dysfunction in the Rett disorder.
Arch Dis Child **85**, 29-37, doi:10.1136/adc.85.1.29 (2001).
8   Gozal, D., Capdevila, O. S., Kheirandish-Gozal, L. & Crabtree, V. M. APOE epsilon 4 allele, cognitive dysfunction,
and obstructive sleep apnea in children. *Neurology* **69**, 243-249, doi:10.1212/01.wnl.0000265818.88703.83
(2007).
9   Liu, X. *et al.* The relationship between inflammation and neurocognitive dysfunction in obstructive sleep apnea
syndrome. *J Neuroinflammation* **17**, 229, doi:10.1186/s12974-020-01905-2 (2020).
10   Serrano-Pozo, A., Das, S. & Hyman, B. T. APOE and Alzheimer's disease: advances in genetics, pathophysiology,
and therapeutic approaches. *Lancet Neurol* **20**, 68-80, doi:10.1016/S1474-4422(20)30412-9 (2021).
11   Sullivan, P. M. *et al.* Targeted replacement of the mouse apolipoprotein E gene with the common human APOE3
allele enhances diet-induced hypercholesterolemia and atherosclerosis. *J Biol Chem* **272**, 17972-17980,
doi:10.1074/jbc.272.29.17972 (1997).
12   Knouff, C. *et al.* Apo E structure determines VLDL clearance and atherosclerosis risk in mice. *J Clin Invest* **103**,
1579-1586, doi:10.1172/JCI6172 (1999).
13   Powell, F. L., Milsom, W. K. & Mitchell, G. S. Time domains of the hypoxic ventilatory response. *Respir Physiol*
112, 123-134, doi:10.1016/s0034-5687(98)00026-7 (1998).
14   Strattan, L. E. *et al.* Novel Influences of Sex and APOE Genotype on Spinal Plasticity and Recovery of Function
after Spinal Cord Injury. *eNeuro* **8**, doi:10.1523/ENEURO.0464-20.2021 (2021).
15   Zinchuk, A. V. *et al.* Polysomnographic phenotypes and their cardiovascular implications in obstructive sleep
apnoea. *Thorax* **73**, 472-480, doi:10.1136/thoraxjnl-2017-210431 (2018).

16 Andaku, D. K. *et al.* Sleepiness, inflammation and oxidative stress markers in middle-aged males with obstructive
sleep apnea without metabolic syndrome: a cross-sectional study. *Respir Res* **16**, 3, doi:10.1186/s12931-015-
0166-x (2015).
17 Moran, M. *et al.* Sleep disturbance in mild to moderate Alzheimer's disease. *Sleep Med* **6**, 347-352,
doi:10.1016/j.sleep.2004.12.005 (2005).
18 Fietze, I. *et al.* Prevalence and association analysis of obstructive sleep apnea with gender and age differences -
Results of SHIP-Trend. *J Sleep Res* **28**, e12770, doi:10.1111/jsr.12770 (2019).
19 Donovan, L. M. & Kapur, V. K. Prevalence and Characteristics of Central Compared to Obstructive Sleep Apnea:
Analyses from the Sleep Heart Health Study Cohort. *Sleep* **39**, 1353-1359, doi:10.5665/sleep.5962 (2016).
20 Solin, P., Roebuck, T., Johns, D. P., Walters, E. H. & Naughton, M. T. Peripheral and central ventilatory responses
in central sleep apnea with and without congestive heart failure. *Am J Respir Crit Care Med* **162**, 2194-2200,
doi:10.1164/ajrccm.162.6.2002024 (2000).
21 Bachetti, T., Bagnasco, S., Piumelli, R., Palmieri, A. & Ceccherini, I. A Common 3'UTR Variant of the PHOX2B Gene
Is Associated With Infant Life-Threatening and Sudden Death Events in the Italian Population. *Front Neurol* **12**,
642735, doi:10.3389/fneur.2021.642735 (2021).
22 Wang, H. *et al.* Variants in angiopoietin-2 (ANGPT2) contribute to variation in nocturnal oxyhaemoglobin
saturation level. *Hum Mol Genet* **25**, 5244-5253, doi:10.1093/hmg/ddw324 (2016).
23 Lo Martire, V. *et al.* CDKL5 deficiency entails sleep apneas in mice. *J Sleep Res* **26**, 495-497,
doi:10.1111/jsr.12512 (2017).
24 Nikodemova, M., Finn, L., Mignot, E., Salzieder, N. & Peppard, P. E. Association of sleep disordered breathing
and cognitive deficit in APOE epsilon4 carriers. *Sleep* **36**, 873-880, doi:10.5665/sleep.2714 (2013).
25 Blackman, J., Love, S., Sinclair, L., Cain, R. & Coulthard, E. APOE epsilon4, Alzheimer's disease neuropathology
and sleep disturbance, in individuals with and without dementia. *Alzheimers Res Ther* **14**, 47,
doi:10.1186/s13195-022-00992-y (2022).
26 Mitchell, G. S. & Johnson, S. M. Neuroplasticity in respiratory motor control. *J Appl Physiol (1985)* **94**, 358-374,
doi:10.1152/japplphysiol.00523.2002 (2003).
27 Huxtable, A. G., Smith, S. M., Peterson, T. J., Watters, J. J. & Mitchell, G. S. Intermittent Hypoxia-Induced Spinal
Inflammation Impairs Respiratory Motor Plasticity by a Spinal p38 MAP Kinase-Dependent Mechanism. *J
Neurosci* **35**, 6871-6880, doi:10.1523/JNEUROSCI.4539-14.2015 (2015).
28 Mahamed, S. & Mitchell, G. S. Is there a link between intermittent hypoxia-induced respiratory plasticity and
obstructive sleep apnoea? *Exp Physiol* **92**, 27-37, doi:10.1113/expphysiol.2006.033720 (2007).
29 Thakre, T. P., Mamtani, M. R. & Kulkarni, H. Lack of association of the APOE epsilon 4 allele with the risk of
obstructive sleep apnea: meta-analysis and meta-regression. *Sleep* **32**, 1507-1511,
doi:10.1093/sleep/32.11.1507 (2009).
30 Turner, A. D. *et al.* Exploring the combined effects of sleep apnea and APOE-e4 on biomarkers of Alzheimer's
disease. *Front Aging Neurosci* **14**, 1017521, doi:10.3389/fnagi.2022.1017521 (2022).
31 Bishop, T. *et al.* Carotid body hyperplasia and enhanced ventilatory responses to hypoxia in mice with
heterozygous deficiency of PHD2. *J Physiol* **591**, 3565-3577, doi:10.1113/jphysiol.2012.247254 (2013).
32 Guyenet, P. G. *et al.* Proton detection and breathing regulation by the retrotrapezoid nucleus. *J Physiol* **594**,
1529-1551, doi:10.1113/JP271480 (2016).
33 Nattie, E. & Li, A. Central chemoreceptors: locations and functions. *Compr Physiol* **2**, 221-254,
doi:10.1002/cphy.c100083 (2012).
34 Prabhakar, N. R. & Peng, Y. J. Peripheral chemoreceptors in health and disease. *J Appl Physiol (1985)* **96**, 359-
366, doi:10.1152/japplphysiol.00809.2003 (2004).
35 Wenninger, J. M., Olson, E. B., Jr., Cotter, C. J., Thomas, C. F. & Behan, M. Hypoxic and hypercapnic ventilatory
responses in aging male vs. aging female rats. *J Appl Physiol (1985)* **106**, 1522-1528,
doi:10.1152/japplphysiol.90802.2008 (2009).
36 Garcia, A. J., 3rd, Rotem-Kohavi, N., Doi, A. & Ramirez, J. M. Post-hypoxic recovery of respiratory rhythm
generation is gender dependent. *PLoS One* **8**, e60695, doi:10.1371/journal.pone.0060695 (2013).

37   Row, B. W., Liu, R., Xu, W., Kheirandish, L. & Gozal, D. Intermittent hypoxia is associated with oxidative stress
and spatial learning deficits in the rat. *Am J Respir Crit Care Med* **167**, 1548-1553, doi:10.1164/rccm.200209-
1050OC (2003).
38   Bubu, O. M. *et al.* Obstructive sleep apnea, cognition and Alzheimer's disease: A systematic review integrating
three decades of multidisciplinary research. *Sleep Med Rev* **50**, 101250, doi:10.1016/j.smrv.2019.101250 (2020).
39   Yaffe, K. *et al.* Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older
women. *JAMA* **306**, 613-619, doi:10.1001/jama.2011.1115 (2011).
40   Zajac, D. J., Green, S. J., Johnson, L. A. & Estus, S. APOE genetics influence murine gut microbiome. *Sci Rep* **12**,
1906, doi:10.1038/s41598-022-05763-1 (2022).
41   Silverstein, A. L. & Alilain, W. J. Intermittent Hypoxia Induces Greater Functional Breathing Motor Recovery as a
Fixed Rather Than Varied Duration Treatment after Cervical Spinal Cord Injury in Rats. *Neurotrauma Rep* **2**, 343-
353, doi:10.1089/neur.2021.0004 (2021).
42   Warren, P. M. *et al.* Rapid and robust restoration of breathing long after spinal cord injury. *Nat Commun* **9**, 4843,
doi:10.1038/s41467-018-06937-0 (2018).

ACKNOWLEDGEMENTS

The study was supported by funding the Craig H. Neilsen Foundation (WJA) and the Kentucky Spinal Cord and

Head Injury Research Trust (WJA).

Additional funding sources include NIH RO1s: AG065220 (LAJ), AG060056 (LAJ), AG80589 (LAJ), AG081421

(LAJ), AG066653 (RCS), CA266004 (RCS), AG078702 (RCS); the Alzheimer's Association (LAJ), and the V-

Scholar Grant (RCS).

AUTHOR CONTRIBUTIONS

C.E.T.: design of experiment, data collection, data analysis and interpretation; initial draft of introduction and

discussion. L.E.M.: data collection, data analysis and interpretation, initial draft of methods and results. M.D.S.:

data analysis, statistical programming, final graphs, manuscript editing and revisions. J.N.W.: technical support,

conceptual feedback. C.M.C.: technical support, conceptual feedback. R.C.S. data analysis and interpretation,

manuscript editing and revisions: L.A.J.: mouse model breeding and support, review of manuscript draft. W.J.A.:

conception and design of the study, experimental design, interpretation of the data, revision and final approval

of the manuscript.

DATA AND CODE AVAILABILITY

The datasets and MATLAB code generated during the current study are available from the corresponding author

on reasonable request.

COMPETING INTERESTS

The authors declare no competing interests.

ETHICAL DECLARATIONS

All experiments were conducted in accordance with the NIH Guidelines Concerning the Care and Use of

Laboratory Animals and were approved by the IACUC committee at the University of Kentucky.