Mitochondrial uncouplers impair human sperm motility without altering ATP content

Short Title: Mitochondrial uncouplers reduce human sperm motility but not ATP content

Will M. Skinner[1], Natalie T. Petersen[2,#3], Bret Unger[4], Shaogeng Tang[5,6], Emiliano Tabarsi[2,#7], Julianna Lamm[2,#8], Liza Jalalian[9], James Smith[10], Ambre M. Bertholet[11,#12], Ke Xu[4,13,14], Yuriy Kirichok[11], Polina V. Lishko[2,15,*].

[1] Endocrinology Graduate Group, University of California, Berkeley, Berkeley, California, United States of America

[2] Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, California, United States of America

[3] Department of Obstetrics and Gynecology, Stanford University School of Medicine, Palo Alto, California, United States of America

[4] Department of Chemistry, University of California, Berkeley, Berkeley, California, United States of America

[5] Department of Biochemistry, Stanford University School of Medicine, Stanford, California, United States of America

[6] Sarafan ChEM-H, Stanford University, Stanford, California, United States of America

[7] Keck School of Medicine, University of Southern California, Los Angeles, California, United States of America

[8] Dewpoint Therapeutics, Boston, Massachusetts, United States of America

[9] Department of Obstetrics and Gynecology, University of California, San Francisco Center for Reproductive Health, San Francisco, California, United States of America

[10] Department of Urology, University of California, San Francisco, San Francisco, California, United States of America

[11] Department of Physiology, University of California San Francisco, San Francisco, California, United States of America

[12] Department of Physiology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, United States of America

[13] California Institute for Quantitative Biosciences, University of California, Berkeley, California,
United States of America

[14] Chan Zuckerberg Biohub, San Francisco, California, United States of America

[15] Center for Reproductive Longevity and Equality, Buck Institute for Research on Aging, Novato,
California, United States of America

*Corresponding Author: Polina V. Lishko

Email: lishko@berkeley.edu

Author Contributions: W.M.S. and P.V.L conceived the project and designed experiments.

W.M.S. conducted most experiments and analyzed most data, except as noted below, and wrote

the manuscript. N.T.P. and W.M.S. performed *in vitro* fertilization experiments. B.U. and W.M.S.

conducted STORM and B.U. analyzed data, under supervision of K.X. S.T. produced Izumo1

antibodies used during acrosome reaction assays. E.T. and J.L. assisted with multiple

experiments. L.J. and J.S. enabled access to and training on computer-aided sperm analysis

machines for motility assays performed by W.M.S. A.M.B. performed mitochondrial patch-clamp

experiments and analyzed that data, under supervision of Y.K. P.V.L and W.M.S. performed

human sperm patch-clamp electrophysiology. All of the authors discussed the results and

commented on the manuscript.

Conceptualization: W.M.S and P.V.L

Data Curation: W.M.S.

Formal Analysis: W.M.S. and A.M.B.

Funding Acquisition: P.V.L. and W.M.S.

Investigation: W.M.S., N.T.P., B.U., E.T., J.L., A.M.B., P.V.L

Methodology: W.M.S., B.U., A.M.B, P.V.L

Project Administration: W.M.S.

Resources: W.M.S., S.T., L.J., J.S., K.X., Y.K., P.V.L.

Supervision: J.S., K.X., Y.K., P.V.L.

Validation: W.M.S.

Visualization: W.M.S., B.U., A.M.B

Writing – Original Draft Preparation: W.M.S.

Writing – Review & Editing: All Authors

Competing Interest Statement: W.M.S., P.V.L, E.T., A.M.B., and Y.K are co-inventors on a

patent related to this work, which UC Berkeley has licensed to YourChoice Therapeutics, for

which P.V.L is a co-founder, and Y.K. is a shareholder. The other authors declare no competing

interests.

Abbreviations:

ANT: Adenine nucleotide translocase
ALH: Amplitude of lateral head displacement
ATP: Adenosine triphosphate
AUC: Area under the curve
BAM15: (N5,N6-bis(2-Fluorophenyl)-[1,2,5]oxadiazolo[3,4-b]pyrazine-5,6-diamine**)**
BAPTA: (1,2-bis(o-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid)
BSA: Bovine serum albumin
CATR**:** Carboxyatractyloside
COXIV: Cytochrome c oxidase subunit 4 isoform 1, mitochondrial
DNP: 2, 4, Dinitrophenol
EGTA: (ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid)
FCCP: Trifluoromethoxyphenylhydrazone
FDA: Food and Drug Administration
HEPES: (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid)
HRP: Horseradish peroxidase
HS: High Saline
HTF: Human Tubal Fluid
ICC: Immunocytochemistry
IMM: Inner mitochondrial membrane
KSOM: Potassium-supplemented simplex optimized medium
LIN: Linearity
MES: 2-(N-morpholino)ethanesulfonate.
MMP: Mitochondrial membrane potential
NEN: Niclosamide Ethanolamine
NMDG: N-methyl-D-glucamine
PI: propidium iodide
PMT: Photomultiplier tube
PVDF: polyvinylidene difluoride
STORM: Stochastic optical reconstruction microscopy
Tris: tris(hydroxymethyl)aminomethane
VCL: Curvilinear velocity
98

Abstract

Sperm motility is necessary for successful fertilization, but there remains controversy about

whether human sperm motility is primarily powered by glycolysis or oxidative phosphorylation.

To evaluate the plausibility of reducing human sperm mitochondrial ATP production as an

avenue for contraceptive development, we treated human sperm with small-molecule

mitochondrial uncouplers, which reduce mitochondrial membrane potential by inducing passive

proton flow, and evaluated the effects on a variety of physiological processes that are critical for

fertilization. We also sought to clarify the subcellular localization of Adenosine Nucleotide

Translocator 4 (ANT4), a gamete-specific protein that has been suggested as a contraceptive

target. We determined that ANT4 is mitochondrially localized, that induced mitochondrial

uncoupling can be partially mediated by the ANT family, and that two uncouplers, Niclosamide

Ethanolamine and BAM15, significantly decreased sperm progressive motility. However, these

uncouplers did not reduce sperm ATP content or impair other physiological processes, implying

that human sperm can rely on glycolysis for ATP production in the absence of functional

mitochondria. Thus, since certain mitochondrial uncouplers impair motility through ATP-

independent mechanisms, they could be useful ingredients in on-demand, vaginally-applied

contraceptives. However, systemically delivered contraceptives that target sperm mitochondria

to reduce their ATP production would need to be paired with sperm-specific glycolysis

inhibitors.

Main Text

Introduction

Human sperm make one of the most remarkable and challenging journeys of any cell in

the human body. To reach the egg, the mature sperm must traverse a distance thousands of

times greater than their cell length, while surviving environments that differ greatly in viscosity,

pH, and oxygen and sugar concentrations (1,2). This journey requires vigorous flagellar motility

(3), and later asymmetrical "hyperactivated" flagellar bending (4), both of which require large

amounts of ATP. However, each human sperm possesses less than a fifty mitochondria (5), less

than almost any other metabolically active cell. Therefore, there has been persistent debate

about the role of mitochondria and oxidative phosphorylation in sperm physiology, with

metabolic studies showing that glycolysis is sufficient for human sperm ATP production (6), but

physiological and clinical studies showing that proper mitochondrial functionality is correlated

with high fertility in humans (7–10). Additionally, sperm contain many unique proteins and

protein isoforms, including metabolic enzymes (11), membrane transporters (12,13) and

receptors (14), and structural proteins (15). Many of these unique proteins have been appealing

targets for development of contraceptives that could disrupt sperm bioenergetics.

One of these unique transporter isoforms is Adenosine Nucleotide Translocator 4

(ANT4), which is gamete-specific (16) and required for spermatogenesis (17), but has an unclear

role in mature human sperm cells. Since its discovery, ANT4 has been seen as an attractive

candidate for the development of a male contraceptive (17–20). Efforts have been made to

develop specific ANT4 inhibitors, which were only able to achieve modest specificity (19,20).

Somatic ANT isoforms (ANT1, 2, and 3 in humans) are essential for transporting ATP and ADP

across the inner mitochondrial membrane (IMM) and have recently been found to be

responsible for mediating mild mitochondrial proton leak (or mitochondrial "uncoupling") across

the IMM in somatic, non-thermogenic tissues (21). It is also possible to activate ANT-mediated

proton leak using exogenous small molecule uncouplers (22). Therefore, we hypothesized that

instead of developing specific ANT4 inhibitors, it may be possible to develop novel small

molecule uncoupler contraceptives that act by specifically activating proton leak through ANT4,

thereby decreasing sperm mitochondrial membrane potential (MMP), ATP production, and

ultimately functionality. To pursue this hypothesis, we first needed to validate the idea that

breakdown of sperm MMP leads to functional dysregulations that could impair fertility. To do

so, we began by testing the physiological effects of various small molecule uncouplers on human

sperm cells.

Results

Representative small molecule uncouplers decrease human sperm mitochondrial membrane

potential and impair basal motility.

158   Four representative small molecule uncouplers were chosen to assess the effects of this

class of compounds on human sperm physiology: 2, 4, Dinitrophenol (DNP); carbonyl cyanide p-

trifluoromethoxyphenylhydrazone (FCCP); BAM 15 (N5,N6-bis(2-Fluorophenyl)-

[1,2,5]oxadiazolo[3,4-b]pyrazine-5,6-diamine**)**; and Niclosamide Ethanolamine (NEN) (Fig. 1A).

Live human sperm cells were treated with varying concentrations of each compound for 30

minutes at 37°C in noncapacitating media at pH 7.4, after which cells were stained with

MitoTracker Red CMX-Ros and mitochondrial membrane potential (MMP) was qualitatively

assessed using flow cytometry (Fig. 1B) and fluorescence microscopy (Fig. 1D). FCCP, BAM15,

and NEN were all able to dose-dependently reduce midpiece fluorescence and therefore MMP,

while DNP was unable to reduce midpiece fluorescence at concentrations up to 300 µM at this

pH level (Fig. S1A). The DMSO and Ethanol vehicles with which compounds were delivered did

not have a significant effect on mitochondrial fluorescence (Fig. S1A) and none of the

compounds decreased sperm viability (Fig. S1D). NEN and BAM15 reduced midpiece

fluorescence values more strongly than FCCP, had lower IC_{50} values, and are known to have

more favorable toxicity profiles (23,24), and so further experiments focused on these two

compounds.

Figure 1: Representative small-molecule uncouplers dose-dependently reduce sperm

mitochondrial membrane potential and basal motility. A) Chemical structures of representative

small molecule uncouplers used. **B)** Effects of SMUs on human sperm mitochondrial potential as

measured by MitoTracker Red fluorescence, normalized to average fluorescence of relevant

vehicle. Nonlinear least squares regression performed using an unweighted [inhibitor] vs.

response three parameter model (Hill Slope = -1.0), identifying "unstable" fits. Shaded areas

indicate 95% confidence bands for true location of each nonlinear fit curve. The nonlinear

regression fit had an Adjusted R^2 value of .8409 for NEN, .8179 for BAM15, and .7143 for FCCP.

Adjusted R^2 value for DNP was less than .1 and so curve is not shown or reported. Asymmetrical

95% CI intervals of IC_{50} for NEN: 1.415 µM to 5.380 µM; BAM15: 3.827 µM to 17.03 µM; FCCP:

9.148 µM to 64.05 µM**. C)** Model showing the possible mechanisms of action of small molecule

uncouplers in the sperm mitochondria and plasma membrane. **D)** Representative fluorescence

micrographs of sperm cells treated with MitoTracker Red CMX-Ros in the absence (top) and

presence (bottom) of 1 µM NEN. **E)** Effects of SMU on human sperm progressive motility as

measured by Computer-Aided Sperm Analysis. Data points represent the average of ≥500 sperm

cells. Nonlinear least squares regression performed and displayed as in B. The nonlinear

regression fit had an Adjusted R^2 value of .6323 for NEN, .2722 for BAM15, and <.1 for DMSO.

Asymmetrical 95% CI interval for NEN: 0.08256 µM to 0.6284 µM. Asymmetrical 95% CI interval

for BAM15: 0.1264 µM to 78.30 µM. **F)** Area Under the Curve quantification for graph E. Error

bars represent SEM, and each data point represents the area under the curve for all

concentrations of one biological replicate. Statistical significance calculated using Dunnett's

multiple comparisons test.

Next, we assessed the effects of NEN and BAM15 on human sperm motility, using

computer-aided sperm analysis (CASA). Both NEN and BAM15 reduced the percent of sperm

showing progressive motility (Fig. 1E,F) and NEN reduced the average curvilinear velocity (Fig.

S2A); NEN reduced sperm progressive motility percent below the 5th percentile of fertile men

(25) and showed a sub-micromolar IC_{50}.

Adenosine Nucleotide Translocators (ANTs) partially mediate NEN-induced mitochondrial

uncoupling, and ANT4 is localized to the human sperm mitochondria

In somatic, non-thermogenic tissues of the body, Adenosine Nucleotide Translocator

(ANT) proteins are known to mediate mild mitochondrial uncoupling under physiological

conditions and in the presence of exogenous uncouplers by inducing H^+ influx across the inner

mitochondrial membrane (21,22). Since sperm cells possess a unique isoform of the family

known as ANT4 (26), we investigated whether NEN-induced uncoupling is also mediated by the

ANT family of proteins. To do this, we performed mitochondrial patch-clamp electrophysiology,

through which it is possible to directly measure current across the inner mitochondrial

membrane. Indeed, in mouse heart mitochondria, which abundantly express ANT1 (27), 500 nM

NEN induced a strong inward H^+ current at physiological pH and negative voltage. This H^+

current was significantly reduced both in the presence of 1 μM Carboxyatractyloside (CATR), a

known ANT inhibitor (Fig. 2B, C), and in heart mitochondria from ANT1-knockout mice (Fig. 2B,

D), indicating that the NEN-induced current is partially mediated by ANT proteins.

Figure 2: NEN induces partially ANT-mediated H^+ current in mouse heart mitoplasts. A)

Diagram of mouse heart mitoplast isolation and preparation for whole-organelle inner

membrane electrophysiology. **B)** Representative whole-organelle electrophysiological traces of

H^+ current density in response to voltage ramp protocols performed on mouse heart mitoplasts.

Left, Wildtype; Right, ANT1 knockout. **C)** Quantification of NEN-induced H^+ current across

mitochondrial inner membrane before and after application of 1 μM CATR, a known ANT

inhibitor. Significances calculated by Student's unpaired t-test. **D)** Quantification of H^+ current

density induced by 500 nM NEN in wildtype and ANT1-knockout mouse heart mitoplasts.

Given this possibility of ANT-mediated uncoupling occurring in sperm, and the presence

of a unique isoform (ANT4) in sperm cells, we investigated whether ANT4 is indeed expressed in

the sperm mitochondria. Other investigators have proposed that ANT4 is expressed throughout

the sperm flagellum, and hypothesized that it may function in collaboration with glycolytic

enzymes to traffic ATP throughout the sperm flagellum (28). However, all other ANT isoforms

are expressed exclusively on the inner mitochondrial membrane, so a flagellar localization would

be surprising. Using immunocytochemistry and confocal microscopy, we found that when

excess primary antibody is used, some ANT4 staining is indeed seen throughout the flagellum

(Fig. S3F), but when primary antibody concentration is titrated down to an optimal

concentration, that staining is restricted to the midpiece in a similar distribution to that of the

inner mitochondrial membrane protein Cytochrome c oxidase (COXIV) (Fig. 3A-C). This ANT4

staining is not seen in non-permeabilized sperm (Fig. S3D), suggesting an exclusively

mitochondrial localization consistent with other ANT isoforms. To confirm this finding, we

conducted stochastic optical reconstruction microscopy (STORM) on human sperm cells similarly

costained for COXIV and ANT4 (Fig. 3D-F, top) and found that upon virtual cross-sectioning of

the midpiece, their distributions formed rings of similar diameter and thickness (Fig. 3D-F,

bottom), consistent with mitochondrial localization.

Figure 3: ANT4 is localized to the human sperm mitochondria. A-C) Confocal microscopy of

human sperm immunocytochemistry showing the localization of COXIV (magenta), ANT4

(green), and DAPI (blue) **D-F)** Stochastic optical reconstruction microscopy (STORM) of human

sperm cells showing the localization of COXIV (magenta) and ANT4 (green). Top panels:

overhead view (XY plane) of sperm cells. Bottom panels: virtual cross section (XZ plane) of

midpiece area indicated in top panels by white dotted box.

Small molecule uncouplers do not reduce human sperm ATP content, and NEN induces

significant proton current across the sperm plasma membrane.

250   To investigate the mechanism of action of NEN and BAM15's effect on sperm basal

motility (Fig. 1E, F) we assessed their effects on sperm ATP levels and found that neither

compound had any effect on ATP concentrations in noncapacitated (Fig. 4A) or capacitated (Fig.

4B) spermatozoa. Consistent with this finding, we observed that NEN and BAM15 had no effect

on other ATP-dependent human sperm physiological processes, including the acrosome

reaction, a necessary pre-fertilization exocytotic event (29,30) that exposes key fertilization

receptors (31,32) (Fig. S4A – C), and human sperm hyperactivated motility (33) (Fig. S4D, E).

NEN also did not impair *in vitro* fertilization between mouse sperm and eggs (Fig. S4F, G).

Figure 4: Effects of small molecule uncouplers on human sperm ATP production and sperm

membrane proton current. A) Left, quantification of ATP content of non-capacitated human

sperm treated with NEN, BAM15, and DMSO vehicle. Right, area under the curve analysis of the

plot to the left. **B)** Left, quantification of ATP content of human sperm capacitated in the

presence of NEN, BAM15, and DMSO vehicle. In both A and B, nonlinear regressions not shown,

because all adjusted R^2 values were <.1. Outliers in both A and B tested using a ROUT coefficient

of 1%, with no outliers detected. Error bars in all plots represent SEM. Statistical significance in

AUC analysis calculated using Dunnett's multiple comparisons test. **C)** Left, Representative

electrophysiological traces showing the effect of vehicles and uncouplers on human sperm

membrane H^+ current density. Right, diagram of sperm electrophysiology setup. Cytoplasmic

droplet is magnified for visibility. **D)** Quantification of maximal current densities in the presence

of 10μM NEN (left) and FCCP (right), at a voltage of -80mV. Error bars represent SEM, and

significance calculated using Tukey's Multiple Comparisons Test.

Since the uncouplers' negative effects on sperm basal motility were not mediated by a

reduced amount of cellular ATP, to further investigate their mechanism of action we assessed

the effects of each compound on sperm plasma membrane H^+-ion current using whole-cell

patch clamp electrophysiology. In this assay, 10 μM NEN and 10 μM FCCP induced a significant

inward transmembrane H^+-ion current at negative voltages (Fig. 4C, D), while 10 μM BAM15,

100 μM DNP, and the Ethanol and DMSO vehicles induced no discernable H^+ current (Fig. 4C,

S5A). The maximum current density of 10 μM FCCP (Fig. 4D) and 100 μM DNP (Fig. S5A) was

consistent with recently reported values in the plasma membrane of HEK293 cells (22).

Additionally, the current density induced by 10 µM NEN across the sperm plasma membrane

(Fig. 4D) was only roughly 1.5 times greater than the current density caused by a 20x lower

concentration of NEN in the inner mitochondrial membrane (IMM) (Fig. 2D). This buttresses the

idea that part of NEN's current is mediated by ANT proteins located exclusively in the IMM,

which increase the IMM's permeability to NEN-induced current.   Although NEN took

significantly longer than FCCP to reach its maximal transmembrane current, their maximal rates

of change of current density were not significantly different, indicating that their mechanisms of

action may be similar (Fig. S5B, C).

Discussion

This study sheds light on several key aspects of sperm energetics and physiology.

There has been an ongoing debate in the field about whether oxidative phosphorylation or

glycolysis is the primary source of ATP for human spermatozoa (34–36), with increasing

evidence supporting the idea that glycolysis is sufficient to support many human sperm

functions (6). Our data support this conclusion, with both NEN and BAM15 failing to reduce

human sperm's ATP content or ability to undergo the acrosome reaction or hyperactivated

motility. Nonetheless, both uncouplers had an inhibitory effect on sperm basal motility, which

in the case of NEN is likely to be related to its induction of current across the sperm plasma

membrane (Fig. 4C, D), which would in turn affect many ion channels and intracellular signaling

pathways that rely on a carefully moderated membrane potential for proper function. Since

BAM15 is known as a more mitochondria-specific uncoupler (22,37–39) and did not induce

plasma membrane current in our sperm electrophysiology assays (Fig. 4C, S5A) the mechanism

of action of its more modest inhibition of basal progressive motility remains to be clarified. The

fact that DNP, a known mitochondrial uncoupler (22), failed to reduce mitochondrial membrane

potential may be related to DNP's low pK_a (23,40) limiting its permeability into living sperm cells

and/or the amount of current produced at pH 7.4, or due to the different complement of ANT

isoforms present in sperm mitochondria versus skeletal muscle mitochondria (16,17,41,42),

which may bind DNP with different affinites.

Although our study found that ANT4 was indeed mitochondrially localized and that the

ANT family of proteins can mediate NEN-induced uncoupling, our data suggest that

contraceptives that solely interfere with mature sperm oxidative phosphorylation will be

unlikely to successfully inhibit key sperm functions. However, pairing an oxidative

phosphorylation inhibitor with an inhibitor of a sperm-specific glycolytic enzyme, like lactate

dehydrogenase C (43), might produce an effective contraceptive product. Additionally, since

niclosamide has been FDA approved for use in humans with a favorable safety profile (24), and

possesses low bioavailability (44), a pK_a close to 6 (45) (close to the vaginal pH), and significant

deleterious effects on sperm motility, it could be an useful ingredient in on-demand vaginal

contraceptive products as an alternative to traditional nonoxynol 9-based spermicides.

Materials and Methods

All experiments were performed in accordance with NIH guidelines for animal research and

approved by UC Berkeley Animal Care and Use Committee (AUP 2015-07-7742), with every

effort made to minimize animal suffering. All described methods are consistent with the

recommendations of the Panel on Euthanasia of the American Veterinary Medical Association

and IACUC committee. All experimental procedures utilizing human derived samples were

approved by the Committee on Human Research at the University of California, Berkeley, IRB

protocol number 2013-06-5395.

Reagents

- Niclosamide Ethanolamine was purchased from AdipoGen Life Sciences,

#AG-CR1-3644

- BAM15 was purchased from Sigma, product number SML1760-5MG

- 2,4, dinitrophenol was purchased from TCI, product D0109-25G

- FCCP was purchased from abcam, product code ab120081.

- Mitotracker Red CMXRos was purchased from ThermoFisher, #M7512

- Polyclonal Antibody to human ANT4 – Novus Biologicals NBP1-89074

raised in Rabbits against human ANT4 peptide

- Mouse Monoclonal Antibody against human COX4l1 – Proteintech

#66110-1-Ig

- Polyvinyl Alcohol was purchased from Fluka, #81366-100G

- A23187 was purchased from MilliporeSigma, #C7522-5MG.

- 10% Goat Serum from Invitrogen, #50-062Z

- Bovine Serum Albumin from Sigma, #A3059-500G

Sperm Preparation

Human sperm cells were collected by masturbation from healthy donors and visually inspected

for normal morphology and motility before use. Spermatozoa were isolated by the swim-up

procedure in HTF or HS solution as previously described (46) and then concentrated by 5 minute

centrifugation at ≤500g and supernatant removal.

Flow Cytometric Assessment of MMP

Human sperm cells were incubated with small molecule uncouplers or vehicles in HS solution for

30 minutes at 37°C , then treated with 50nM MitoTracker Red in the presence of either

uncouplers or vehicles for an additional 30 minutes. Subsequently, cells were washed by

centrifugation for 5 minutes at 500 g, supernatant was removed, and cells were resuspended in

HS solution. Fluorescence was measured by flow cytometry on a BD LSR Fortessa Celeste or BD

LSR Fortessa X20, using gates for size and density to exclude debris and aggregates, and a PE-

Texas Red Filter set (561nm laser excitation, 610/20 nm emission filter) to record fluorescence

(Fig. 1, S1). Fluorescence midpiece location was confirmed by microphotography on an Olympus

IX71 inverted microscope using a Hamamatsu Orca-ER digital camera, and a 60x Olympus

UPlanSApo objective (Fig. 1). Fluorescence statistics were analyzed using Microsoft Excel

version 16 and GraphPad Prism version 9. The subpopulation of cells with the highest gaussian

distribution of fluorescence were manually gated and the mean fluorescence of this population

was normalized to the average of the means of the same population in the relevant vehicle

control samples of each run. Each data point represents the mean fluorescence of the highest

normally peak in an individual flow cytometry run of 10,000 cells. Nonlinear least squares

regression performed using an unweighted [inhibitor] vs. response three parameter model (Hill

Slope = -1.0), identifying "unstable" fits. Outliers tested using a ROUT coefficient of 1%, with no

outliers detected. Residuals were checked for normality using the D'Agostino-Pearson omnibus

normality test, checked for clustering using the Replicates test, and checked for appropriate

weighting using the homoscedasticity test, and passed all tests, except for FCCP which failed the

test for homoscedasticity.

Area Under the Curve Analysis

For all dose-response plots, Area Under the Curve (AUC) analysis was performed using the

trapezoid method ($\Delta X*(Y_1+Y_2)/2$) on the arithmetic means of the measured parameter at each

compound concentration. Each data point represents the area under the curve of all

concentrations tested in one biological replicate. Effects of each compound were then

compared using Dunnett's Multiple Comparison's Test.

Sperm Motility Assessment

Isolated human sperm were incubated with small molecule uncouplers and vehicles at 37°C for

30 minutes, after which they were mixed with 1% Polyvinyl Alcohol to prevent adherence to

glass and loaded into a Leja Standard Count 2-chamber sperm analysis slide (depth 20 micron).

Sperm motility was analyzed on a Hamilton-Thorne IVOS I computer-aided sperm analyzer, at

37°C. Motility was measured for least 500 cells in each condition. Progressive motility was

defined using Hamilton-Thorne's proprietary software module. Nonlinear least squares

regression performed by GraphPad Prism version 9 using an unweighted [inhibitor] vs. response

three parameter model (Hill Slope = -1.0), identifying "unstable" fits. Outliers were tested using

a ROUT coefficient of 1% and none were identified. Residuals were checked for normality using

the D'Agostino-Pearson omnibus normality test, checked for clustering using the Replicates test,

and checked for appropriate weighting using the homoscedasticity test. Residuals for NEN and

BAM15 passed all tests, except that BAM15 failed the test for homoscedasticity with a p-value

of 0.0207.

Human sperm hyperactivated motility was measured on a Hamilton-Thorne IVOS II computer-

aided sperm analyzer, as above, with the definition for hyperactivated motility set as VCL

>150 µm/s AND LIN <50% AND ALH >7.0 µm (33).

Flow Cytometric Assessment of Acrosome Status

Human sperm cells were incubated for 3.5 hours in capacitation media (20% Fetal Bovine Serum

and 25mM Sodium Bicarbonate in HS solution) at 37°C and 5% CO_2. In the final hour of

incubation, custom monoclonal antibodies raised in mouse hybridoma cells against the

ectodomain of human Izumo1 were added, at a concentration of 10µg/mL. Subsequently 50µM

A23187 was added, and cells were incubated at 37°C and 5% CO_2 for an additional 30 minutes,

after which cells were washed by centrifugation 3x (500 g, 5 min) in HS solution, incubated 30

minutes at room temperature with a goat anti-mouse 647 secondary antibody (Invitrogen

A32728, final concentration 4µg/mL) in 10% goat serum. After an additional 3 washes by

centrifugation, propidium iodide (PI) was added to a final concentration of 1µg/mL, and cells

were analyzed by flow cytometry using the same instrumentation as above, except that Izumo-

647 was visualized using an APC filter set (bandpass 670/30 or 670/14nm) and Propidium Iodide

was visualized using a PE-Texas Red filter set (bandpass 610/20nm). Izumo-647 positive cells

were scored as acrosome reacted, and PI negative cells were scored as living.

Generation of IZUMO1 monoclonal antibody using mouse hybridoma

The IZUMO1 monoclonal antibodies were generated from mouse hybridoma and characterized

as previously described (47). Briefly, IZUMO1 antibodies 4E04 and 6F02 belong to the murine

IgG1 isotype with kappa light chain. For antibody production, hIZUMO1-positive hybridomas

were cultured in medium containing 10% FBS and subsequently adapted to an FBS-free medium.

The condition media were harvested, and the IgG antibody were purified by protein G affinity

chromatography and by Superdex-200 gel filtration chromatography in a buffer of 150 mM NaCl,

20 mM HEPES pH 7.4. Purified IgG antibodies were concentrated to 1 mg/mL with the addition

of 10% (v/v) glycerol for use.

ATP Assessment

Human Sperm cells were incubated with uncoupler compounds or vehicle for 1 hour at 37°C in

white 96-well Costar polystyrene assay plates, under sterile conditions. ATP content was

assessed using a PerkinElmer ATPlite Luminescence Assay System Kit, according to manufacturer

instructions, and luminescence was imaged on a Biotek Synergy H4 Hybrid plate reader, set to

endpoint/kinetic setting (one measurement point per well), with 1 second of integration time

and a manual gain setting of 135. ATP content was quantified by interpolating against a

standard curve of ATP solutions, per manufacturer instructions. Nonlinear least squares

regression performed by GraphPad Prism version 9 using an [inhibitor] vs. response three

parameter model (Hill Slope = -1.0), weighted by $1/Y^2$, identifying "ambiguous" fits. Outliers

tested using a ROUT coefficient of 1%, with no outliers detected. Residuals were checked for

normality using the D'Agostino-Pearson omnibus normality test, checked for clustering using the

Replicates test, and checked for appropriate weighting using the homoscedasticity test, and

passed all tests.

Mouse In Vitro Fertilization

All mice were kept in the Animal Facility of the University of California, Berkeley, fed standard

chow diet (PicoLab Rodent diet 20, #5053, LabDiet), and hyper-chlorinated water *ad libitum* in a

room with controlled light (14 hours light, 10 hours darkness) and temperature (23±0.5°C).

Animals were humanely killed by CO_2 asphyxiation and cervical dislocation according to ACUC

guidelines with every effort made to minimize suffering.

C57Bl/6N (Charles River) female mice between 4 and 16 weeks old were superovulated by

intraperitoneal injection of 5 IU pregnant mare serum gonadotropin (PMSG, Sigma, G4877) and

48 hours later, 5 IU human chorionic gonadotropin (hCG, Millipore, 230734). 13 hours after hCG

injection the females were euthanized and cumulus-oocyte complexes were collected from the

oviduct ampulla in HTF medium (Embryomax, Specialty Media, Millipore) and incubated for 30

minutes at 37°C and 5% CO_2 prior to insemination. Simultaneously, mouse sperm were isolated

from the cauda epididymis as previously described[36] and allowed to capacitate at 37°C and 5%

CO_2 for 60-90 minutes in the presence of uncoupler compounds or control vehicle. Sperm and

eggs were then mixed to a final concentration of 210,000 sperm/mL in 600μL of HTF medium

and incubated at 37°C and 5% CO_2 for 4 hours. Afterwards, the eggs were washed by mouth

pipetting in HTF media to remove excess sperm. A final wash in KSOM media (Zenith Biotech)

supplemented with 1mg/mL BSA was then done before dividing the eggs into 10 μl drops of

KSOM/BSA media overlaid with embryo-tested light mineral oil (Millipore) for culture at 37°C,

5% CO_2. Best efforts were made to divide equal numbers of eggs from each mouse into control

and uncoupler-treated conditions. 3.5 days following IVF, the number of fertilized eggs was

determined visually by assessing the percentage of embryos that reached morula or blastula

stage.

Sperm Whole-Cell Electrophysiology

Whole-cell electrophysiology was performed on isolated human sperm cells as described

previously (48,49). Once sperm cell "break in" was achieved in HS solution, cells were subjected

to a voltage ramp protocol in the presence of bath and pipette solutions designed to ensure that

only H^+ current across the membrane was measured. Intracellular (pipette) solution contained:

150mM NMDG, 5mM BAPTA, 100mM HEPES, 1mM Tris Chloride, pH 7.5 with $MeSO_3$.

Extracellular (bath) solution contained: 157mM NMDG, 20mM MES, 1mM $MgCl_2$, 100µM $ZnCl_2$,

pH 6 with $MeSO_3$. Compounds and vehicles were applied by perfusion, and currents were

recorded until they showed a minimal change between subsequent voltage ramps. Then,

compounds were washed out, and subsequent compounds added once currents had returned to

baseline values. Representative traces and time courses created in Origin Pro version 9.0.0 and

statistical analysis performed in GraphPad Prism 9.

Mitochondrial Patch-Clamp Electrophysiology

Mouse heart mitochondria were isolated from wildtype and ANT1[-/-] C57BL/6J2z mice and

mitoplasts prepared and patch-clamped as described previously (21). Both the bath and pipette

solutions were formulated to record H^+ currents and contained only salts that dissociate into

large anions and cations that are normally impermeant through ion channels or transporters.

Pipettes were filled with 130 mM tetramethylammonium hydroxide (TMA), 1.5 mM EGTA, 2 mM

Tris chloride, and 100 mM HEPES (or MES). pH was adjusted to 7.5 with d-gluconic acid, and

tonicity was adjusted to ∼360 mmol/kg with sucrose. Bath solution contained 100 mM HEPES

(or MES) and 1 mM EGTA (pH adjusted to 7.0 with Trizma base, and tonicity adjusted to ∼300

mmol/kg with sucrose).

Immunocytochemistry

Briefly, sperm were adhered onto poly-D-Lysine coated coverslips, and, in some experiments,

incubated 15 minutes at 37°C with 50 nM MitoTracker Red CMXRos, then washed and fixed in

4% PFA for 10 minutes, and washed and additionally fixed in ice-cold methanol for 1 minute.

Cells were permeabilized and blocked in 5% BSA and 0.1% Triton X-100 in PBS for 45 minutes,

then incubated overnight at 4°C with rabbit polyclonal antibody against Human ANT4 (Novus

#NBP1-89074, used at .08 μg/mL) and, in some experiments, a mouse monoclonal antibody

against COXIV (COX4l1) (Proteintech #66110-1-Ig, used at 6.8μg/mL). After washing, samples

were incubated with secondary antibodies (Jackson Alexa Fluor 647 AffiniPure Donkey Anti-

Rabbit IgG (H+L) (code 711-605-152) at a dilution of 1:1000, and Jackson ImmuoResearch anti-

mouse Cy3 or Molecular Probes A10521 goat anti-mouse Cy3 at a dilution of 1:1000) for 45

minutes, then washed and mounted using ProLong Gold Antifade mountant with DAPI.

Cells were imaged on an Olympus FV3000 inverted laser scanning confocal microscope, using a

Olympus UPLSAPO Super Apochromat 60x oil immersion objective (NA 1.35) and Olympus Z-

81226 immersion oil. Laser Scanning was performed by galvanometer, unidirectionally at a rate

of 2μs/pixel with no averaging, with pixel sizes of .207 μm/pixel in the x and y direction.

Multiple colors were imaged in line-by-line sequential scanning mode, and multiple z-planes

were acquired with a thickness of .47 μm, on a linear-encoded stage.   Images displayed are

maximum intensity z-stacks of all relevant z-planes. All colors were imaged with a pinhole

diameter of 233 μM, and laser illumination was passed through a 10% neutral density filter.

DAPI was illuminated by a 405 nm laser set to 3% power, and emitted light was passed through

a 430-470 nm emission filter and detected with a photomultiplier tube (PMT) set to 500V. Cy3

was illuminated by a 561 nm laser set to 4% power, with a 570-620 nm emission filter and a PMT

set to 500V. Alexa 674 was illuminated by a 642 nm laser set to 4% power, with a 650-750 nm

emission filter and a PMT set to 550V.

Super-resolution Microscopy

ICC slides made as described above were imaged using three-dimensional stochastic optical

reconstruction microscopy (3D-STORM) (50,51) performed on a home-built set-up using a Nikon

CFI Plan Apo λ 100× oil immersion objective (NA 1.45), as described previously (52). In brief, the

sample was mounted with an imaging buffer consisting of 5% (w/v) glucose, 100 mM

cysteamine, 0.8 mg ml^{-1} glucose oxidase and 40 μg ml^{-1} catalase in a Tris HCl buffer (pH 7.5). For

two-color imaging of COXIV and ANT4, the two targets were labeled by Alexa Fluor 647 and

CF568, respectively, and were imaged sequentially using 647- and 560-nm excitation lasers.

These lasers were passed through an acousto-optic tunable filter and illuminated a few

micrometers into the sample at around 2 kW cm^{-2}, thus photoswitching most of the labeled dye

molecules in the sample into the dark state while allowing a small, random fraction of molecules

to emit across the wide-field over different camera frames. Single-molecule emission was

passed through a cylindrical lens of focal length 1 m to introduce astigmatism (51), and recorded

with an Andor iXon Ultra 897 EM-CCD camera at a frame rate of 110 Hz, for a total of around

50,000 frames per image. Data acquisition used publicly available software

(https://github.com/ZhuangLab/storm-control). The raw STORM data were analyzed using

Insight3 software (51) according to previously described methods (50,51). Secondary antibodies

used: Alexa Fluor 647-labelled goat anti-mouse (Invitrogen, A21236, 1:400) and donkey anti-

rabbit (Jackson ImmunoResearch, 711-005-152, 1:70) conjugated with CF568 succinimidyl ester

(Biotium, 92131).

Western Blotting

Human Sperm or HEK293 cells were lysed by addition of RIPA^{++} Buffer (composition: 50mM Tris

HCl, pH 7.4; 150mM NaCl; 1% Triton X-100; 0.5% Sodium Deoxylcholate; 0.1% Sodium Dodecyl

Sulfate; 1 mM EDTA; 10% glycerol; with Pierce Protease inhibitor tablet or solution added to

manufacturer's instructions), then homogenized by 25 passes through a 21-gauge or smaller

syringe, then shook on ice for 30 minutes, and centrifuged at 12000 rpm at 4°C for 20 minutes,

after which the supernatant was collected and frozen at -80°C. After thawing, protein

concentration was assessed using a Pierce Rapid Gold BCA Protein Assay Kit, per manufacturer

instructions, and measured on a Biotek Synergy H4 Hybrid plate reader. A target of 30μg of

protein was mixed with 4x Laemmli Buffer and Beta-Mercaptoethanol, heated for 10 minutes at

99°C, then loaded to each well of a precast 4-20% mini-Protean TGX gel (Bio-Rad), and run in an

electrophoresis chamber until ladder bands were well-resolved. Transfer onto a PVDF

membrane was accomplished with an Invitrogen iBlot Dry Blotting System, following

manufacturer instructions, at 20V for 7 minutes. After washing 3x with PBS with 0.1% Tween

(PBST), the membrane was blocked in 3% Bovine Serum Albumin for 15-30 minutes, then

incubated rocking with primary antibody overnight at 4°C. The following day, the membrane

was washed 3x with PBST, and incubated with a 1:15000 dilution of HRP-conjugated Abcam

Ab6721 Goat pAb anti-Rabbit IgG secondary antibody for one hour at room temperature. After

3x PBST washes and 2x PBS washes, bands were visualized using a Prometheus ProSignal Pico

kit, per instructions, and imaged in a ProteinSimple FluorChem M. Afterwards, images were

inverted, rotated, and cropped in ImageJ 1.52a.

Acknowledgments:

Health. The authors also wish to thank Dr. Liliya Gabelev Khasin, Dr. Caroline Williams, and Dr.

Lisa Treidel, and Dr. Erwin Goldberg for their assistance with this project.

This work was supported by an NSF Graduate Research Fellowship and the Irving H.

Wiesenfeld Fellowship administered by the UC Berkeley Center for Emerging and Neglected

Diseases to W.M.S., a Male Contraceptive Initiative Graduate Fellowship to P.V.L. and Liliya

Gabelev Khasin, NIH NICHD grant K99HD104924 to S.T., NIGMS grant R35GM136415 to Y.K. ,

Pew Biomedical Scholars Award to P.V.L. and K.X. and Bakar Fellow Spark Award to P.V.L.

References

1.   Suarez SS, Pacey AA. Sperm transport in the female reproductive tract. Hum Reprod Update.
2006 Jan 1;12(1):23–37.

2.   Suarez SS. Mammalian Sperm Interactions with the Female Reproductive Tract. Cell Tissue
Res. 2016 Jan;363(1):185–94.

3.   Turner RM. Moving to the beat: a review of mammalian sperm motility regulation. Reprod
Fertil Dev. 2006;18(2):25.

4.   Suarez SS. Control of hyperactivation in sperm. Hum Reprod Update. 2008 Nov 1;14(6):647–
57.

5.   Mundy AJ, Ryder TA, Edmonds DK. Asthenozoospermia and the human sperm mid-piece.
Hum Reprod. 1995 Jan 1;10(1):116–9.

6.   Hereng TH, Elgstøen KBP, Cederkvist FH, Eide L, Jahnsen T, Skålhegg BS, et al. Exogenous
pyruvate accelerates glycolysis and promotes capacitation in human spermatozoa. Hum
Reprod. 2011 Dec 1;26(12):3249–63.

7.   Gallon F, Marchetti C, Jouy N, Marchetti P. The functionality of mitochondria differentiates
human spermatozoa with high and low fertilizing capability. Fertil Steril. 2006
Nov;86(5):1526–30.

8.   Paoli D, Gallo M, Rizzo F, Baldi E, Francavilla S, Lenzi A, et al. Mitochondrial membrane
potential profile and its correlation with increasing sperm motility. Fertil Steril. 2011 Jun
1;95(7):2315–9.

9.   Boguenet M, Bouet PE, Spiers A, Reynier P, May-Panloup P. Mitochondria: their role in
spermatozoa and in male infertility. Hum Reprod Update [Internet]. 2021 Feb 8 [cited 2021
Apr 26];(dmab001). Available from: https://doi.org/10.1093/humupd/dmab001

10. Zhang G, Wang Z, Ling X, Zou P, Yang H, Chen Q, et al. Mitochondrial Biomarkers Reflect
Semen Quality: Results from the MARCHS Study in Chongqing, China. PLOS ONE. 2016 Dec
22;11(12):e0168823.

11. Piomboni P, Focarelli R, Stendardi A, Ferramosca A, Zara V. The role of mitochondria in
energy production for human sperm motility: Mitochondria functionality in human
spermatozoa. Int J Androl. 2012 Apr;35(2):109–24.

12. Rodić N, Oka M, Hamazaki T, Murawski MR, Jorgensen M, Maatouk DM, et al. DNA
Methylation Is Required for Silencing of Ant4, an Adenine Nucleotide Translocase Selectively
Expressed in Mouse Embryonic Stem Cells and Germ Cells. STEM CELLS. 2005;23(9):1314–
23.

13. Zheng LP, Wang HF, Li BM, Zeng XH. Sperm-specific ion channels: targets holding the most
potential for male contraceptives in development. Contraception. 2013 Oct;88(4):485–91.

14. Noda T, Lu Y, Fujihara Y, Oura S, Koyano T, Kobayashi S, et al. Sperm proteins SOF1,
TMEM95, and SPACA6 are required for sperm–oocyte fusion in mice. Proc Natl Acad Sci.
2020 May 26;117(21):11493–502.

15. Leung MR, Chiozzi RZ, Roelofs MC, Hevler JF, Ravi RT, Maitan P, et al. In-cell structures of
conserved supramolecular protein arrays at the mitochondria–cytoskeleton interface in
mammalian sperm. Proc Natl Acad Sci [Internet]. 2021 Nov 9 [cited 2021 Nov 9];118(45).
Available from: https://www.pnas.org/content/118/45/e2110996118

16. Lim CH, Brower JV, Resnick JL, Oh SP, Terada N. Adenine Nucleotide Translocase 4 Is
Expressed Within Embryonic Ovaries and Dispensable During Oogenesis. Reprod Sci. 2015
Feb;22(2):250–7.

17. Brower JV, Rodic N, Seki T, Jorgensen M, Fliess N, Yachnis AT, et al. Evolutionarily Conserved
Mammalian Adenine Nucleotide Translocase 4 Is Essential for Spermatogenesis. J Biol
Chem. 2007 Oct 5;282(40):29658–66.

18. Hamazaki T, Leung WY, Cain BD, Ostrov DA, Thorsness PE, Terada N. Functional Expression
of Human Adenine Nucleotide Translocase 4 in Saccharomyces Cerevisiae. PLOS ONE. 2011
Apr 21;6(4):e19250.

19. Leung WY, Hamazaki T, Ostrov DA, Terada N. Identification of adenine nucleotide
translocase 4 inhibitors by molecular docking. J Mol Graph Model [Internet]. 2013 Sep [cited
2018 Jun 12];45. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3821932/

20. Zhang Y, Tian D, Matsuyama H, Hamazaki T, Shiratsuchi T, Terada N, et al. Human Adenine
Nucleotide Translocase (ANT) Modulators Identified by High-Throughput Screening of
Transgenic Yeast. J Biomol Screen. 2016 Apr;21(4):381–90.

21. Bertholet AM, Chouchani ET, Kazak L, Angelin A, Fedorenko A, Long JZ, et al. H + transport is
an integral function of the mitochondrial ADP/ATP carrier. Nature. 2019 Jul;571(7766):515.

22. Bertholet AM, Natale AM, Bisignano P, Suzuki J, Fedorenko A, Hamilton J, et al.
Mitochondrial uncouplers induce proton leak by activating AAC and UCP1. Nature. 2022
Jun;606(7912):180–7.

23. Childress ES, Alexopoulos SJ, Hoehn KL, Santos WL. Small Molecule Mitochondrial
Uncouplers and Their Therapeutic Potential: Miniperspective. J Med Chem. 2018 Jun
14;61(11):4641–55.

24. Kadri H, Lambourne OA, Mehellou Y. Niclosamide, a Drug with Many (Re)purposes.
ChemMedChem. 2018;13(11):1088–91.

25. World Health Organization. WHO laboratory manual for the examination and processing of
human semen [Internet]. 6th ed. Geneva: World Health Organization; 2021 [cited 2021 Nov
11]. Available from: https://apps.who.int/iris/handle/10665/343208

26. Dolce V, Scarcia P, Iacopetta D, Palmieri F. A fourth ADP/ATP carrier isoform in man:
identification, bacterial expression, functional characterization and tissue distribution. FEBS
Lett. 2005 Jan 31;579(3):633–7.

27. Levy SE, Chen YS, Graham BH, Wallace DC. Expression and sequence analysis of the mouse
adenine nucleotide translocase 1 and 2 genes. Gene. 2000 Aug;254(1–2):57–66.

28. Kim YH, Haidl G, Schaefer M, Egner U, Mandal A, Herr JC. Compartmentalization of a unique
ADP/ATP carrier protein SFEC (Sperm Flagellar Energy Carrier, AAC4) with glycolytic
enzymes in the fibrous sheath of the human sperm flagellar principal piece. Dev Biol. 2007
Feb;302(2):463–76.

29. Brucker C, Lipford GB. The human sperm acrosome reaction: physiology and regulatory
mechanisms. An update. Hum Reprod Update. 1995 Jan;1(1):51–62.

30. Hirohashi N, Yanagimachi R. Sperm acrosome reaction: its site and role in fertilization†. Biol
Reprod. 2018 Feb;99(1):127–33.

31. Satouh Y, Inoue N, Ikawa M, Okabe M. Visualization of the moment of mouse sperm–egg
fusion and dynamic localization of IZUMO1. J Cell Sci. 2012 Nov 1;125(21):4985–90.

32. Balestrini PA, Jabloñski M, Schiavi-Ehrenhaus LJ, Marín-Briggiler CI, Sánchez-Cárdenas C,
Darszon A, et al. Seeing is believing: Current methods to observe sperm acrosomal
exocytosis in real time. Mol Reprod Dev. 2020;87(12):1188–98.

33. Mortimer ST, van der Horst G, Mortimer D. The future of computer-aided sperm analysis.
Asian J Androl. 2015;17(4):545–53.

34. du Plessis SS, Agarwal A, Mohanty G, van der Linde M. Oxidative phosphorylation versus
glycolysis: what fuel do spermatozoa use? Asian J Androl. 2015;17(2):230–5.

35. Ford WCL. Glycolysis and sperm motility: does a spoonful of sugar help the flagellum go
round? Hum Reprod Update. 2006 Jun 1;12(3):269–74.

36. Storey BT. Mammalian sperm metabolism: oxygen and sugar, friend and foe. Int J Dev Biol.
2008;52(5–6):427–37.

37. Kenwood BM, Weaver JL, Bajwa A, Poon IK, Byrne FL, Murrow BA, et al. Identification of a
novel mitochondrial uncoupler that does not depolarize the plasma membrane. Mol Metab.
2014 Apr 1;3(2):114–23.

38. Kenwood BM, Calderone JA, Taddeo EP, Hoehn KL, Santos WL. Structure–activity
relationships of furazano[3,4-b]pyrazines as mitochondrial uncouplers. Bioorg Med Chem
Lett. 2015 Nov 1;25(21):4858–61.

39. Firsov AM, Popova LB, Khailova LS, Nazarov PA, Kotova EA, Antonenko YN. Protonophoric
action of BAM15 on planar bilayers, liposomes, mitochondria, bacteria and neurons.
Bioelectrochemistry. 2021 Feb 1;137:107673.

40. Hopfer U, Lehninger AL, Thompson TE. Protonic conductance across phospholipid bilayer
membranes induced by uncoupling agents for oxidative phosphorylation. Proc Natl Acad Sci.
1968 Feb 1;59(2):484–90.

41. Stepien G, Torroni A, Chung AB, Hodge JA, Wallace DC. Differential expression of adenine
nucleotide translocator isoforms in mammalian tissues and during muscle cell
differentiation. J Biol Chem. 1992 Jul 25;267(21):14592–7.

42. Doerner A, Pauschinger M, Badorff A, Noutsias M, Giessen S, Schulze K, et al. Tissue-specific
transcription pattern of the adenine nucleotide translocase isoforms in humans. FEBS Lett.
1997 Sep 8;414(2):258–62.

43. Goldberg E. The Sperm Specific Form of Lactate Dehydrogenase (LDHC4) is Required for
Fertility and is an Attractive Target for Male Contraception (A Review). Biol Reprod
[Internet]. 2020 [cited 2020 Dec 2]; Available from:
https://academic.oup.com/biolreprod/advance-
article/doi/10.1093/biolre/ioaa217/6008540

44. Zhang X, Zhang Y, Zhang T, Zhang J, Wu B. Significantly enhanced bioavailability of
niclosamide through submicron lipid emulsions with or without PEG-lipid: a comparative
study. J Microencapsul. 2015 Jul 4;32(5):496–502.

45. Jurgeit A, McDowell R, Moese S, Meldrum E, Schwendener R, Greber UF. Niclosamide Is a
Proton Carrier and Targets Acidic Endosomes with Broad Antiviral Effects. PLoS Pathog
[Internet]. 2012 Oct 25 [cited 2020 Aug 18];8(10). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3486884/

46. Skinner WM, Mannowetz N, Lishko PV, Roan NR. Single-cell Motility Analysis of Tethered
Human Spermatozoa. Bio-Protoc [Internet]. 2019 Mar 5 [cited 2019 May 7];9(5). Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6482966/

47. Tang S, Lu Y, Skinner WM, Sanyal M, Lishko PV, Ikawa M, et al. Human sperm TMEM95 binds
eggs and facilitates membrane fusion [Internet]. bioRxiv; 2022 [cited 2022 Jul 9]. p.

2022.06.10.495573. Available from:
https://www.biorxiv.org/content/10.1101/2022.06.10.495573v1

48. Liu B, Mundt N, Miller M, Clapham DE, Kirichok Y, Lishko PV. Recording Electrical Currents
across the Plasma Membrane of Mammalian Sperm Cells. JoVE J Vis Exp. 2021 Feb
14;(168):e62049.

49. Kirichok Y, Lishko PV. Rediscovering sperm ion channels with the patch-clamp technique.
MHR Basic Sci Reprod Med. 2011 Aug 1;17(8):478–99.

50. Rust MJ, Bates M, Zhuang X. Sub-diffraction-limit imaging by stochastic optical
reconstruction microscopy (STORM). Nat Methods. 2006 Oct;3(10):793–6.

51. Huang B, Wang W, Bates M, Zhuang X. Three-Dimensional Super-Resolution Imaging by
Stochastic Optical Reconstruction Microscopy. Science. 2008 Feb 8;319(5864):810–3.

52. Wojcik M, Hauser M, Li W, Moon S, Xu K. Graphene-enabled electron microscopy and
correlated super-resolution microscopy of wet cells. Nat Commun. 2015 Jun 11;6(1):7384.

699
700
701 **Figures**
702
703 **Figure 1: Representative small-molecule uncouplers dose-dependently reduce sperm**
704 **mitochondrial membrane potential and basal motility.**

706   **Figure 2: NEN induces partially ANT-mediated H⁺ current in mouse heart mitoplasts.**

708 **Figure 3: ANT4 is localized to the human sperm mitochondria.**

Figure 4: Effects of small molecule uncouplers on human sperm ATP production and sperm
 membrane proton current.

Supplementary Information for

Mitochondrial uncouplers impair human sperm motility without

altering ATP content

Short Title: Mitochondrial uncouplers reduce human sperm motility but not ATP

content

Will M. Skinner[1], Natalie T. Petersen[2,#3], Bret Unger[4], Shaogeng Tang[5,6], Emiliano Tabarsi[2,#7],

Julianna Lamm[2,#8], Liza Jalalian[9], James Smith[10], Ambre M. Bertholet[11,#12], Ke Xu[4,13,14], Yuriy

Kirichok[11], Polina V. Lishko[2,15,*].

[1] Endocrinology Graduate Group, University of California, Berkeley, Berkeley, California, United
States of America

[2] Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley,
California, United States of America

[3] Department of Obstetrics and Gynecology, Stanford University School of Medicine, Palo Alto,
California, United States of America

[4] Department of Chemistry, University of California, Berkeley, Berkeley, California, United States
of America

[5] Department of Biochemistry, Stanford University School of Medicine, Stanford, California, United
States of America

[6] Sarafan ChEM-H, Stanford University, Stanford, California, United States of America

[7] Keck School of Medicine, University of Southern California, Los Angeles, California, United
States of America

[8] Dewpoint Therapeutics, Boston, Massachusetts, United States of America

[9] Department of Obstetrics and Gynecology, University of California, San Francisco Center for
Reproductive Health, San Francisco, California, United States of America

[10] Department of Urology, University of California, San Francisco, San Francisco, California,
United States of America

[11] Department of Physiology, University of California San Francisco, San Francisco, California,
United States of America

[12] Department of Physiology, David Geffen School of Medicine, University of California Los
Angeles, Los Angeles, California, United States of America

[13] California Institute for Quantitative Biosciences, University of California, Berkeley, California,
United States of America

[14] Chan Zuckerberg Biohub, San Francisco, California, United States of America

[15] Center for Reproductive Longevity and Equality, Buck Institute for Research on Aging, Novato, California, United States of America

*Corresponding Author: Polina V. Lishko
Email: lishko@berkeley.edu

Figure S1. Validation of flow cytometric assessment of human sperm mitochondrial membrane potential A) Left: Effect of compound vehicles on MitoTracker fluorescence. **Right:** Effect of the uncoupler 2, 4, DNP on MitoTracker Fluorescence. In both, nonlinear regression performed, but not shown or reported because adjusted R^2 values were <.1. B) Representative gating logic for flow cytometry experiments, isolating individual sperm cells from debris (top) and aggregations (bottom). C) Representative flow cytometry histograms for MitoTracker Red CMX-Ros Mitochondrial Membrane Potential Assays. D) Percent of propidium iodide-excluding cells (membrane intact) after 30-minute incubation with compounds, as measured by flow cytometry. Each data point represents the average of 10,000 cells, and error bars represent SEM. All conditions were compared to DMSO .1% using Dunnett's Multiple Comparisons Test, and no

conditions were significantly different. E) Representative flow cytometry result of propidium iodide vitality assay.

Figure S2: Effects of small molecule uncouplers on human sperm curvilinear velocity as measured by computer-aided sperm analysis. **A)** Curvilinear velocity analysis of noncapacitated human sperm cells. Left: Each data point represents the average curvilinear velocity of at least 200 sperm cells. Nonlinear least squares regression performed using an unweighted [inhibitor] vs. response three parameter model (Hill Slope = -1.0), identifying "unstable" fits. Shaded areas indicate 95% confidence bands for true location of each nonlinear fit curve. The nonlinear regression fit for NEN had an Adjusted R^2 value of 0.1833. BAM15 & DMSO nonlinear regressions not shown because their adjusted R^2 values were below 0.1. NEN IC_{50} Asymmetrical 95% CI interval: 0.07433 µM to 6.950 µM. Outliers tested using a ROUT coefficient of 1%, with two outliers in the DMSO condition excluded. NEN residuals passed the Replicates and homoscedasticity tests, but failed the D'Agostino-Pearson omnibus normality test with a p-value of .0366. Right: Area Under the Curve analysis for data at left. Each data point represents the area under the curve of one biological replicate. **B)** Curvilinear Velocity analysis of

capacitated human sperm cells. Left: Nonlinear Regression performed as in A. The nonlinear regression fit for DMSO had an Adjusted R^2 value of 0.7328 with an IC_{50} 95% Confidence Interval of 0.1126 μM to 1.925 μM. The nonlinear regression fit for NEN had an Adjusted R^2 value of 0.7275 with an IC_{50} 95% Confidence Interval of 0.9469 μM to 7.230 μM. The nonlinear regression fit for BAM15 had an Adjusted R^2 value of 0.3885 with an IC_{50} 95% Confidence Interval of 0.8117 μM to 294309381 μM. All tests performed as in A, with no outliers detected, and with residuals passing all tests, except that the residuals for NEN failed the test for homoscedasticity with a P-value of .006. Right: Area Under the Curve analysis for data at left.

Figure S3: Human ANT4 polyclonal antibody validation. A,B) hANT4 Western blot of HEK293 cells and human sperm. A), unblocked; B) blocked with immunizing peptide. C) ICC of permeabilized human sperm incubated with hANT4 pAb preblocked with immunizing peptide and 50 nM Mitotracker Red CMX-Ros. **D)** ICC of non-permeabilized human sperm labelled as in C. E) ICC of permeabilized HEK293 cells labelled as in C. F) Permeabilized human sperm cells incubated with excess hANT4 pAb primary antibody (.4 µg/mL) and 50 nM MitoTracker Red CMX-Ros. In C-F, Orange – MitoTracker Red CMX-Ros, Blue – Dapi, Green – hANT4. All images are maximum intensity projections of sequential confocal image z-slices.

Figure S4: NEN and BAM15 have no effect on human sperm acrosome reaction competence or human sperm hyperactivated motility, and NEN has no effect on Mouse IVF success A) Percentage of unfixed, capacitated human sperm cells that were acrosome-reacted and membrane-intact (Izumo1$^+$/Propidium Iodide$^-$) as measured by flow cytometry, after incubation with and without the calcium ionophore A23187. Each data point represents the percentage of 100,000 cells showing this fluorescence pattern. Error bars represent SEM, and significance was calculated by Tukey's multiple comparisons test, with individual variances computed for each comparison. All combinations were compared, but for clarity, not all comparisons are shown. All non-shown comparisons were not significant. **B)** Representative data plots from one run of flow cytometry, comparing ionophore A23187-treated (red) and control cells (blue). **C)** Representative confocal microscopy image of A23187-treated sperm cells taken from flow cytometry run, showing Izumo1 staining in red, and DAPI staining in blue. AR: Acrosome Reacted, AI: Acrosome Intact. **D)** Percent of capacitated human sperm cells

displaying hyperactivated motility, as measured by computer-aided sperm analysis. Each data point represents the percent of at least 200 sperm cells showing hyperactivated motility. Nonlinear least squares regression performed using an unweighted [inhibitor] vs. response three parameter model (Hill Slope = -1.0), identifying "unstable" fits. Shaded areas indicate 95% confidence bands for true location of each nonlinear fit curve. The nonlinear regression fit for DMSO had an Adjusted R^2 value of 0.6501, and an IC_{50} 95% CI of 0.1052 μM to 2.131 μM. The nonlinear regression fit for NEN had an Adjusted R^2 value of 0.5062, and an IC_{50} 95% CI of 0.2957 μM to 12.04 μM. The nonlinear regression fit for BAM15 had an Adjusted R^2 value of 0.1667, and an IC_{50} 95% CI of 0.1418 μM to +infinity. **E)** Area under the curve analysis of hyperactivation data from panel E. Each data point represents the area under the curve of one biological replicate, error bars represent SEM, and statistical significance calculated using Dunnett's Multiple Comparisons Test. **F)** Quantification of mouse eggs that developed to morula or blastocyst stage after *In vitro* fertilization (IVF) using SMU-treated sperm. Significance calculated using a one-tailed ratio paired t-test. **G)** Representative micrographs of mouse embryos 3.5 days after IVF. Left, oocytes exposed to control sperm; Right, oocytes exposed to 10 μM NEN-treated sperm. M - Morula, D - Dead, U - Unfertilized.

Figure S5: Supplemental electrophysiology quantifications. A) 10 µM BAM15, 100 µM DNP, 0.1% EtOH**,** and 0.1% DMSO have no effect on sperm plasma membrane current densities. Significance calculated using Tukey's multiple comparisons test. **B)** NEN takes longer to reach maximal uncoupling than FCCP (top), but their maximum rates of current density increase are similar (bottom). Significance calculated using Šídák's multiple comparison's test, with a single pooled variance **C)** Time courses of current density at -80 mV during application of compounds. Colored bars represent time period during which each compound was being applied. Dotted line in NEN time course represents a period where other electrophysiological tests were performed on that cell.