Transient glycolytic complexation of arsenate enhances resistance in the

enteropathogen *Vibrio cholerae*

Emilio Bueno*[1], Víctor Pinedo[1], Dhananjay D. Shinde[2], André Mateus[3] Athanasios

Typas[3], Mikhail M Savitski[3], Vinai C. Thomas[2] and Felipe Cava*[1].

[1]Laboratory for Molecular Infection Medicine Sweden, Department of Molecular

Biology, Umeå Centre for Microbial Research, Umeå University, Umeå SE-90187,

Sweden.

[2] Center for Staphylococcal Research, Department of Pathology and Microbiology,

University of Nebraska Medical Center, Omaha, Nebraska, 68198, USA

[3] Genome Biology Unit, European Molecular Biology Laboratory (EMBL),

Heidelberg, Germany

Correspondence to:

felipe.cava@umu.se

emilio.bueno@umu.se

Keywords: arsenate, arsenite, quorum sensing, Tn-seq, *Vibrio cholerae*

Running title: Arsenate resistance in *Vibrio cholerae*.

Transient glycolytic complexation of arsenate enhances resistance in the

enteropathogen *Vibrio cholerae*

Abstract

The ubiquitous presence of toxic arsenate (As^V) in the environment has virtually

raised mechanisms of resistance in all living organisms. Generally, bacterial

detoxification of As^V relies on its reduction to arsenite (As^{III}) by ArsC, followed by

the export of As^{III} by ArsB. However, how pathogenic species resist this metalloid

remains largely unknown. Here, we found that *V. cholerae,* the etiologic agent of

the diarrheal disease cholera, outcompetes other enteropathogens when grown

on millimolar concentrations of As^V. To do so, *V. cholerae* uses, instead of ArsCB,

the As^V-inducible *vc1068-1071* operon (renamed *var* for *v*ibrio *a*rsenate

*r*esistance)*,* which encodes the arsenate repressor ArsR, an alternative

glyceraldehyde-3-phosphate dehydrogenase, a putative phosphatase, and the

As^V transporter ArsJ. In addition to Var, *V. cholerae* induces oxidative stress-

related systems to counter ROS production caused by intracellular As^V.

Characterization of the *var* mutants suggested these proteins function

independently from one another and play critical roles in preventing deleterious

effects on the cell membrane potential and growth derived from the accumulation

As^V. Mechanistically, we demonstrate that *V. cholerae* complexes As^V with the

glycolytic intermediate 3-phosphoglycerate into 1-arseno-3-phosphoglycerate

(1As3PG). We further show that 1As3PG is not transported outside the cell;

instead, it is subsequently dissociated to enable extrusion of free As^V through

ArsJ. Collectively, we propose the formation of 1As3PG as a transient metabolic

storage of As^V to curb the noxious effect of free As^V. This study advances our

understanding of As^V resistance in bacteria and underscores new points of

vulnerability that might be an attractive target for antimicrobial interventions.

Introduction

Introduction

Arsenic is a toxic metalloid commonly found in aquatic and terrestrial

environments (1–3) as arsenate (As^V) or its reduced form, arsenite (As^{III}). The

potent toxicity of this element makes it one of the best-studied natural poisons

that impact public health (4). The mechanism of As^V toxicity is due to its structural

resemblance to phosphate oxyanions and its uptake into the cell through

phosphate transport channels (5, 6). Once in the cytoplasm, As^V can replace

phosphate in energy-generating reactions resulting in the formation of ADP-As^V

instead of ATP by the ATPase, and through substrate-level phosphorylation

during glycolysis (7, 8). Conversely, cellular toxicity by As^{III} is exerted by a

different mechanism. As^{III} can interact directly with thiol groups of proteins and

molecules, thereby interfering with diverse cellular processes (9, 10). Despite its

risk to public health, arsenic has been used as an antimicrobial agent to treat

infectious diseases (1, 4, 11, 12), increasing the selective pressure for microbes

to acquire arsenic resistance.

Bacterial strategies to resist As^V include chelation by metal-binding

metallothioneins (13), methylation to less toxic and more volatile arsenic

derivatives (14), respiration (15), and extrusion of As^V or As^{III} by specific efflux

pumps (e.g., ArsJ and ArsB, respectively) (9, 16–18). In bacteria, the most

common mechanism of As^V resistance genes occurs via As^V reduction to As^{III}

followed by the extrusion of the latter through an As^{III} transporter (19). As^V

resistance genes are commonly organized in operons, e.g., the *arsRDABC*

operon from *Escherichia coli* R773 (19). In this example, *arsC* encodes a

dedicated arsenate reductase and *arsB* an efflux pump of As^{III} (20). The *arsR*

gene encodes the repressor of the *ars* operon. When As^{III} is present, its

interaction with ArsR releases it from the *ars* promoter, thus increasing

transcription of the operon. *arsA* encodes an ATPase that, together with ArsB

forms the As^{III} extrusion system. Alternatively, As^{III} extrusion via ArsB can be

driven by proton motive force in species lacking ArsA. Finally, *arsD* encodes a

chaperone that enhances arsenic extrusion by transferring As^{III} to ArsAB (9).

Despite being extensively studied in environmental soil and marine bacteria

species, yet the mechanisms underlying As^{V} resistance in human pathogenic

bacteria remain largely unknown. Here, we investigated As^{V} resistance in

representative enteric pathogens and found that *V. cholerae* exhibits remarkably

high resistance to this metalloid. Using genome mining and functional

characterization, we demonstrated that *V. cholerae* does not detoxify As^{V} by

reducing it to As^{III}. Instead, transposon-based functional screens identified the

operon *vc1068-1071* as the primary genetic determinant of As^{V} resistance in *V.*

cholerae. Mechanistic characterization of this system revealed that resistance to

As^{V} in *V. cholerae* is mediated by a transient metabolic complexation of As^{V} with

the glycolytic intermediate 3-phosphoglycerate (3PG) to generate 1-arseno-3-

phosphoglycerate (1As3PG). Our results support that this As^{V}-containing

metabolite is transient in the cell: its formation cushions the cellular damage

caused by free As^V, while its dissociation enables *V. cholerae* to release free As^V

through the ArsJ efflux permease.

Results

High resistance to arsenate in *V. cholerae* is independent of the ArsC

reductase

While studying the tolerance of enteropathogenic bacteria to metalloids we

observed that *Vibrio cholerae*, the causative agent of cholera, was able to grow in

media supplemented with supraphysiological (30 mM) concentrations of arsenate

(As^V) (Fig. S1). This level of resistance was unmatched by other human enteric

pathogens, as lower concentrations of As^V (10 mM) readily caused total or partial

growth inhibition of *Salmonella enterica*, *Citrobacter rodentium, Yersinia*

pseudotuberculosis, Enterohemorragic E. coli (EHEC) *and Shigella flexneri* (Fig.

1A)*. Consistently, *V. cholerae* outcompeted these species by 5-500-fold in co-

culture experiments grown in media supplemented with As^V (Fig. 1B).

To investigate the molecular mechanisms of As^V resistance in *V. cholerae*, we

first searched for canonical determinants for As^V detoxification in its genome. *V.*

cholerae vc2165 encodes a homolog of *E. coli* K12 arsenate reductase ArsC,

including the conserved catalytic cysteine and arginine residues (36, 37) (Fig.

S2A). However, while *E. coli*'s ArsC enzyme is critical for resistance to As^V,

inactivation of *vc2165* did not compromise *V. cholerae* growth in the presence of

this metalloid (Fig 1C). Thus, we reasoned that *V. cholerae*'s ArsC might not be

active or expressed. To measure ArsC activity we quantified extracellular As^V and

As^{III} by inductively coupled plasma mass spectrometry (ICP-MS) and observed

that while *E. coli* reduced ~80% of As^V to As^{III} in an ArsC-dependent manner, *V.*

cholerae produced no As^{III} (Fig. 1C). Interestingly, replacement of *V. cholerae*

arsC allele by that of *E. coli* caused growth inhibition, likely due to toxic As^{III}

formation, as growth was recovered by expressing *E. coli's* As^{III}-transporter ArsB

(Fig. S2B). Altogether, these results indicate that *V. cholerae* ArsC is inactive and

that As^V resistance in this bacterium is independent of the ArsC-AsrB system that

relies on the production and elimination of As^{III}.

Resistance to As^V in *V. cholerae* depends on the *vc1068-vc1071* operon.

To identify the genetic determinants conferring resistance to As^V in *V. cholerae*,

we performed two complementary genome-wide screenings based on: ***i)***

Transposon Insertion Sequencing (TIS), and ***ii)*** the use of a previously described

V. cholerae transposon mutant library [32] (Fig. 2). Both screens permit

assessment of the mutants' relative fitness. However, the first approach is more

suitable for discovering mutants with subtle fitness differences, while the latter can

identify candidates otherwise underappreciated by TIS due to cross-

complementation between mutants within a population. Only four non-essential

Tn mutants (*vc1068::tn, vc1069::tn, vc1070::tn,* and *vc1071::tn*) were mutually

under-represented in both screenings (Fig. 2). We renamed *vc1068-71* as ***var*** for

***vibrio* a**rsenate **r**esistance cluster referred hereafter for reasons described below.

vc1068 encodes an homolog of the ArsR repressor, *vc1069* (renamed as *varG*)

encodes a putative glyceraldehyde-3-phosphate dehydrogenase (GAPDH),

vc1070 (renamed as *varH*) encodes an uncharacterized putative phosphatase,

and *vc1071*, an ArsJ-like As^V transporter (17). As expected, the *arsC* mutant was

not identified in the screens, confirming that *V. cholerae* does not detoxify As^V by

reducing it to As^{III}. Although homologues to VarG and ArsJ were previously

reported in *Pseudomonas aeruginosa*, their function in As^V resistance has not

been studied *in vivo* (17).

To verify the implication of the *var* cluster on As^V resistance, we constructed

individual in-frame marker-less deletion mutants. All mutants were sensitive to As^V

except for Δ*arsR* (Fig. 3A), suggesting that the conditional lethality of the *arsR*::tn

was likely due to the polar effect on the expression of the downstream genes

varG- varH-arsJ. Indeed, transcriptional analysis using fusions of the *lacZ* reporter

to potential promoter regions located upstream of each *var* gene demonstrated

that expression of this cluster depends on $As^{V,}$ and it is driven from a single

promoter upstream of *arsR* (Fig. S3). Inactivation of *arsR* turned transcription from

this promoter constitutive (both in presence and absence of As^V), confirming that

VC1068 is the *V. cholerae*'s ArsR repressor of the *var* operon (Fig. S3). Further

characterization revealed that Δ*arsJ* presents a more dramatic growth defect

compared to the rest of the *var* mutants at low As^V concentrations (1 mM) (Fig. 3).

However, at 10 mM of As^V, the growth of the *varG* and *varH* mutant strains was

also compromised despite the presence of ArsJ. These results suggest that the

ArsJ transporter is the main determinant for As^V resistance in *V. cholerae* but

VarG and VarH also become relevant at higher concentrations of As^V.

Furthermore, the inactivation of *varG, varH* and *arsJ* (double and triple mutants)

aggravated growth defects exhibited by the single mutants, suggesting

independent functions for these proteins in As^V resistance in *V. cholerae* (Fig.

3A).

168 VarG is an As^V-inducible GAPDH that preferentially binds As^V

V. cholerae VarG presents a high degree of protein sequence (E-value: 5^{e-86}) (Fig.

S4A) and structure (AlphaFold2 TM-score=0.96) (Fig. S4B) similarity to the

glycolytic GAPDH VC2000 (Gap), including the conserved catalytic Cys residue

(38). To characterize VarG's activity compared to the glycolytic paralogue Gap

particularly, we purified both proteins from *V. cholerae* and performed *in vitro*

GAPDH activity assays (39) in the presence or absence of As^V. While both

proteins exhibited GAPDH activity, VarG showed preference for As^V over P_i (the

canonical substrate) (Fig. 4A). Conversely, the glycolytic GAPDH Gap performed

better with P_i. Furthermore, while the expression of *Gap* remained low and

constitutive, transcription from the *var* promoter was strongly induced by As^V (Fig.

4B). Therefore, these results suggest that while *V. cholerae* Gap is active on As^V,

this enzyme specific activity and protein levels are likely insufficient to replace

VarG essentiality in *V. cholerae* resistance to As^V.

Arsenate detoxification in *V. cholerae* is facilitated by extrusion of free arsenate instead of complexed as 1-arseno-3-phosphoglycerate

A VarG homolog from *P. aeruginosa* was proposed to function as a GAPDH that complexes As^V with glyceraldehyde-3-phosphate (G3P) into 1-arseno-3-phosphoglycerate (1As3PG), which is further eliminated through ArsJ (17). However, these experiments were performed *in vitro* using a commercial glycolytic GAPDH from rabbit instead of the As^V-specific GAPDH from *P. aeruginosa* DK2. If this model applies to *V. cholerae,* the absence of either VarG or ArsJ will similarly affect AsV export and resistance. However, our data shows that ∆*varG* did not phenocopy the growth defect of ∆*arsJ* (Fig. 3). Furthermore, analysis of intracellular levels of As^V by ICP-MS revealed that, in relation to the WT, while the ∆*varG* strain only showed moderate (ca. 3x) accumulation of As^V, inactivation of ArsJ accumulated this metalloid ca. 20x (Fig. 4C). Although these results suggest that VarG is not essential to export As^V, we cannot discard the possibility of As^V being complexed and extruded as 1As3PG in *V. cholerae*.

Using high-resolution mass spectrometry (HRMS), we detected a mass (*m/z*) of 304.8696 Da, specific to the cultures supplemented with As^V, which was compatible with 1As3PG (*m/z* 304.8674, *m/z*<0.003 Da difference between experimental and theoretical mass) (Fig. 4D and S5). 1As3PG was also detected

in the ∆*varG*, consistent with the ability of the constitutively expressed glycolytic

Gap GAPDH to use As^V (Fig. S5). 1As3PG was detected in intracellular

cytoplasmic fractions but not in the extracellular milieu, suggesting that this

molecule is not expelled in *V. cholerae* (Fig. 4D and S5). As it has been reported

that 1As3PG is rapidly dissociated to As^V and 3PG (40), we reasoned that if

exported, its decomposition should increase extracellular 3PG levels in relation to

cultures without As^V. However, we found no significant differences in the levels of

extracellular 3PG between conditions supporting the idea that As^V detoxification

likely occurs through the release of free As^V rather than of 1As3PG through ArsJ

(Fig. 4C and S5).

VarH is involved in 1As3PG dissociation and arsenate export

To find clues that help us understand the role that 1As3PG plays in *V. cholerae*'s

resistance to As^V we investigated VarH. Structural *in silico* analysis using the

phyre2 tool (www.sbg.bio.ic.ac.uk/phyre2) and AlphaFold2 revealed that VarH is a

cysteine-based protein tyrosine phosphatase (PTP) belonging to the dual-

specificity phosphatase superfamily that presents a very high topology similarity to

the human Kap PTP phosphatase (Fig. S6AB). To determine if VarH has

phosphatase activity, we purified this protein and assessed its capacity to

dephosphorylate the chromogenic substrate pNPP. Our results show that,

compared to the wheat phosphatase or *V. cholerae's* PTP VC1041 positive

controls, VarH presents lower phosphatase activity, which turned off upon

replacement of the putative catalytic cysteine and arginine by glycine residues

(Fig. 5A, left panel). Furthermore, consistent with the importance of these

residues in VarH activity, replacement of the WT *varH* allele by the catalytically

inactive variant *varH* C113G/R119G impaired *V. cholerae* growth in the presence

of As^V (Fig. 5A, right panel). These results suggest that the phosphatase activity

of VarH is required to provide As^V resistance in *V. cholerae*.

Since VarG and VarH are coregulated by As^V and both support resistance to

this metalloid, we hypothesized that VarH could regulate VarG activity by

modulating its phosphorylation state. To assess such a possibility, we purified

VarG from *V. cholerae* WT and Δ*varH* (Fig. S6C) and analyzed VarG

phosphorylation by liquid chromatography–tandem mass spectrometry (LC–

MS/MS). However, even though VarG was phosphorylated on Tyr108 and

Ser204, those phosphorylation levels were independent of VarH (Fig. S6D). Then,

we reasoned that as P$_i$ and As^V are analogs, VarH could potentially act on

1As3PG as a phosphatase to produce 1AsG + Pi or as an "arsenatase" to render

3PG + As^V. Remarkably, we found that 1As3PG concentration were significantly

elevated in Δ*varH* compared to those in the WT strain (Fig. S6E), suggesting that

VarH might use 1As3PG as a substrate. Although we could not confirm VarH's

activity *in vitro,* potentially due to the instability of 1As3PG (41), the fact VarH is

an active phosphatase but 1AsG is not detected in *V. cholerae* suggests that

VarH could dissociate the 1As3PG complex into 3PG and free As^V which would

be released through ArsJ.

Intracellular arsenate impacts the *V. cholerae* proteome and leads to ROS

accumulation and membrane potential defects

To broadly assess global responses triggered by As^V in *V. cholerae*, we analyzed

the pathogen proteome upon exposure to As^V. Interestingly, a relatively low

number of proteins (□20 proteins) were upregulated upon exposure to As^V,

indicating that adaptations to this metalloid might not require extensive proteome

rewiring in *V. cholerae*. As expected, these proteins included the As^V resistance

Var proteins but also amino acids permeases, a Zn/Cd transporter, members of

the PTS-fructose system, the starvation stress response protein RaiA, and the

cysteine synthesis and the peroxiredoxin PrxA (Fig. 6AB). Cysteine is one of the

main targets of reactive oxygen species (ROS) and a functional component of

glutathione, which plays a crucial role in the adjustment of cellular redox potential

(42). In this line, PrxA has been involved in the adaptation to hydrogen peroxide in

V. cholerae (43). Therefore, we reasoned that the *var* mutants, which display

increased As^V intracellular levels, might suffer from higher oxidative stress than *V.*

cholerae WT strain. To test this hypothesis, we measured ROS levels in the *V.*

cholerae WT and *var* mutant strains and observed a direct correlation between

As^V accumulation and ROS level ($\Delta arsJ>\Delta varH>\Delta varG>$WT) (Fig. 6C). These

findings collectively indicate that AsV induces oxidative stress in *V. cholerae*

which is exacerbated in the *var* mutants due to a supraphysiological accumulation

of this toxic metalloid.

In addition to ROS affecting Cys-containing proteins, we cannot discard a

direct effect of As^V on the proteome of *V. cholerae*. To this end, we performed

thermal proteome profiling (TPP), a method that globally analyzes multiple types

of protein interactions (25). TPP data showed an increased thermal stability of

multiple cytosolic proteins that could potentially interact with As^V ($\square$3-fold) in the

var mutants with respect to the WT (Fig. 6D and table S). As^V-interacting protein

candidates included proteins involved in the synthesis of heme groups of

respiratory cytochromes, respiratory ubiquinone, fermentation, and carbon

catabolism, suggesting that accumulation of As^V in Var-defective strains might

interfere with cell bioenergetics. To investigate this possibility, we monitored the

bacterial membrane potential in the presence of As^V using flow cytometry (Fig.

6E). Interestingly, inactivation of the *var* system resulted in a significant

depolarization of the mutant's membrane. Together, these results demonstrate

that AsV can, directly and indirectly, affect the *V. cholerae*'s proteome, leading to

oxidative stress and defects in cellular bioenergetics, both possibly related to the

growth impairment of the *var* mutants.

***V. cholerae var operon* provides As^V resistance to enteric pathogens**

In silico analysis showed that, except for *C. rodentium*, ArsR is the only As^V-

resistance determinant conserved between *V. cholerae* and the As^V-sensitive

enteric pathogens (Fig. 7A). In general, these enteropathogenic species detoxify

As^V via ArsC-dependent reduction to As^{III}, and subsequent extrusion through

ArsB. As *V. cholerae* lacks the As^{III}- production/extrusion tandem ArsCB, we

wondered whether the *var* genes could also augment resistance to As^V in other

species or would rather require additional species-specific elements. To address

this question, we expressed the *var* operon excluding the ArsR repressor in

enteropathogens sensitive to As^V and studied their growth capacity in the

presence of As^V. Strikingly, induced coexpression of *varG-varH-arsJ* in several

enteropathogens boosted their growth on As^V (Fig. 7B), demonstrating that the

Var system is sufficient and compatible with the ArsCB system in providing

resistance against As^V in other bacteria.

DISCUSSION

Arsenic is ubiquitous in the environment and consequently, most living organisms, including bacteria, encode arsenic-resistance proteins (44–49). However, functional characterization of these systems and how they support bacterial fitness in environments with As^V is underappreciated. In this study, we describe that *V. cholerae* exhibits higher resistance to As^V than other enteric pathogens.

The strategy used by *V. cholerae* to resist As^V is different from that used by other gammaproteobacterial enteropathogens. While most of these species encode an ArsC arsenate reductase and an ArsB arsenite efflux pump, *V. cholerae* lacks the latter and has an inactive ArsC. Therefore, *V. cholerae* does not reduce As^V or expel it out of the cell. From an evolutionary perspective, this could be a vestige of a functional ArsCB system that, after losing the ArsB component, has inactivated ArsC to prevent the generation of highly toxic As^{III}. It remains to be investigated whether *V. cholerae* ArsC has undergone functional diversification to fulfill another role in the cell, perhaps related to its neighboring hypothetical genes (*vc2164-vc2167*).

Our screening identified a four-gene operon (*var operon*) conferring resistance to *V. cholerae* that encoded the ArsR repressor, the As^V permease ArsJ, a putative GAPDH (VarG) and a phosphatase (VarH). Interestingly, while most ArsR derepress transcription by sensing As^{III} (50), the lack of As^{III} production in *V. cholerae* suggests that ArsR probably senses As^V in this bacterium. Although more research is required to determine ArsR responses to As^V and As^{III} in *V.*

cholerae, comparative sequence analysis between *V. cholerae* ArsR and that of

As^{III} producing bacteria have revealed variations in conserved residues, including

the E→K and C→A substitutions in the As^{III} binding motif (50) (Fig. S7A).

Although GAPDH enzymes are mostly known for their role in glycolysis, they

have been functionally linked to diverse processes (51–53). A previous study

reported that the GAPDH homolog to *V. cholerae's* VarG detoxified As^{V} by

generating 1As3PG from G3P. However, such conclusions were drawn from *in*

vitro experiments that used commercial glycolytic GAPDH from rabbits instead of

the specific As^{V}-inducible GAPDH, and monitored As^{V} transport instead of direct

1As3PG detection (17). Our results, combining *in vitro* GAPDH activity assays

and HRMS analyses from intracellular samples provided direct evidence that

VarG is a GAPDH that preferentially uses As^{V} over P$_i$ to synthesize 1As3PG.

VarG role in *V. cholerae's* As^{V} resistance adds to this growing list of GAPDH

moonlighting functions.

Although the same study suggested that As^{V} is exported through ArsJ as

1As3PG (17), the more severe growth defect of *V. cholerae's arsJ* compared to

varG mutant suggests that 1As3PG does not play a major role in the export of As^{V}

in this bacterium. Accordingly, the *arsJ* mutant accumulates intracellular As^{V} ca. 6

times more than the *varG* mutant. Even if we cannot wholly discount the transport

of 1As3PG through ArsJ (as this compound can nonetheless be produced in the

absence of VarG, likely by the constitutive glycolytic Gap activity), the fact that we

have not detected extracellular 1As3PG or increased 3PG in cultures

supplemented with As^V suggest that ArsJ expels free As^V.

During glycolysis, the GAPDH enzyme catalyzes the simultaneous

phosphorylation and oxidation of G3P to form 1,3 bi-phosphoglycerate (1,3-BPG),

which is further converted into 3PG by phosphoglycerate kinase coupled to the

transfer of the high-energy phosphate to ADP, making ATP. We reasoned that

while the metabolic complexing of As^V into 1As3PG might decrease its toxicity by

reducing As^V interaction with *V. cholerae*'s proteome (Fig4D), the formation of

1As3PG could be a dead-end product that would affect a major energy-producing

pathway. In line with this reasoning, our results showed that VarH presents low

phosphatase activity compared to canonical phosphatases. As low catalytic rate

has been previously associated with enzymatic promiscuity (54), we hypothesized

that the putative phosphatase VarH would function as an "arsenatase" that

dissociates the 1As3PG complex to clear the glycolytic pathway and free As^V for

its export through ArsJ. Indeed, inactivation of VarH results in increased

intracellular levels of 1As3PG, which correlates with the accumulation of arsenate

in the cell, thereby supporting the idea that ArsJ primarily exports free As^V in *V.*

cholerae. Although synteny analyses suggest coevolution between ArsJ and

VarG, the presence of VarH is rather exclusive to Vibrionaceae (Fig. S7B). It

remains to be investigated whether the amino acid differences in ArsJ between

VarG$^+$ and VarG$^-$ species explain the choice of transporting free or complexed

As^V.

It has been previously reported that arsenic induces the production of ROS in

the cell (55–57). Our results show that in the presence of As^V, *V. cholerae*

induces oxidative stress responses: i.e., elevated levels of the hydrogen peroxide-

tolerance determinant PrxA (58) and multiple proteins involved in the synthesis of

cysteine. As cysteine-containing proteins are targets of ROS (59), and cysteine is

also the building block of glutathione, stimulating its synthesis might help to

alleviate oxidative damage provoked by As^V. Additionally, we found that As^V

directly or indirectly (via ROS) alters the stability of proteins implicated in energy-

generating pathways such as TCA, fermentation, and respiration. Indeed, As^V

accumulation in *var* mutant dissipated cellular membrane potential (a proxy of the

cell's energy status). The correlation between As^V accumulation with ROS and

membrane depolarization exhibited by the *var* mutants suggests that oxidative

stress and energy depletion might account for their growth impairment in As^V.

V. cholerae is a pathogenic bacteria endemic to Bangladesh and India, regions

where the concentrations of As^V in water and soil are the highest worldwide (60).

Hence, *V. cholerae's* superior resistance to As^V could have evolved as an

adaptive strategy to thrive in As^V-rich environments while endowing the cholerae

pathogen with a fitness advantage over free-living and host-associated

neighboring bacteria. Interestingly, despite the phylogenetic distance,

heterologous expression of the *var* operon enhanced As^V resistance in all the

pathogens assayed, demonstrating that this system has a high degree of

versatility to work in combination with other arsenic detoxification systems.

Collectively, this work proposes a novel mechanism for As^V detoxification in *V.*

cholerae that is entirely independent of As^V-reduction and As^{III}-extrusion. Our

results support a model in which the formation of a transient As^V-containing

glycolytic intermediate cushions the impact of free As^V in the cell. While a

metabolic sink might temporarily aid cells in enduring As^V toxicity, this strategy

may exert a high cost in the long term as glycolysis is a major pathway in cell

bioenergetics. We propose "dearsenylation" of 1As3PG by VarH provides a

double benefit by decongesting any jam that may occur in glycolysis and freeing

As^V for extrusion to the extracellular media by ArsJ.

The essentiality of the *V. cholerae's var* operon to survive in environments with

As^V (e.g., such as the host) suggests that this machinery may be considered a

novel class of bacterial targets suitable for therapeutic intervention.

ACKNOWLEDGMENTS.

This work was supported by the Knut and Alice Wallenberg Foundation (KAW), The Laboratory of Molecular Infection Medicine Sweden (MIMS), the Swedish Research Council and the Kempe Foundation. V.C.T and D.D.S. were supported by the National Institutes of Health/ National Institute of Allergy and Infectious Diseases (NIH/ NIAID) grant P01-AI83211 (Metabolomics Core) and R01-AI125588. The metabolomics analyses were performed at the University of Nebraska Medical Center Mass Spectrometry and Proteomics Core Facility administrated through the Office of the Vice-Chancellor for Research and supported by state funds from the Nebraska Research Initiative (NRI). We thank Erik Björk and Laurent Ouerdane for their help with the ICPMS analyses and for insightful discussions, J. J. Mekalanos for the *V. cholerae* C6706 Tn-mutant library and Inigo Ruiz and David Arranz for their technical support

COMPETING INTEREST.

The authors declare that there are not competing interests in relation with the work described.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Strains used in this work are listed in Supplementary Table S2. *V. cholerae*

strains used in this study are derivatives of the El Tor clinical isolate C6706. In

addition to *V. cholerae*, in this study also were used the strains *Escherichia coli*

K12 MG1655 and the *arsC*::Tn mutant strain from the Keio collection (21),

Enterohemorragic Escherichia coli EHEC O157:H7, *Salmonella enterica serovar*

Typhimurium LT2, *Citrobacter rodentium* DBS100, *Shigella flexneri* M90T, *and*

Yersinia pseudotuberculosis YPIII. Strains were grown aerobically at 37⁰C in 15

ml tubes containing 3 ml of complete LB medium (10 g tryptone, 10 g NaCl, and 5

g yeast extract/L). To grow *Shigella flexneri,* TSB medium was used (Tryptone

(Pancreatic Digest of Casein), 17 g Soytone (Peptic Digest of Soybean), 3 g

Glucose, 2.5 g Sodium Chloride 5 g Dipotassium Phosphate 2.5). When used,

casamino acids were supplemented at 1%. To study resistance to As^V, 3 x 10^6

overnight grown cells were inoculated into 200 µl of LB medium with or without

As^V at indicated concentrations in 96-wells plates and OD values were measured

every 10 minutes using a BioTek Epoch2 plate reader. Where appropriate,

antibiotics were added to *V. cholerae* and *E. coli* cultures at the following

concentrations: 200 (streptomycin, Sm), 100 (carbenicillin, Cb), 5

(Chloramphenicol, Cm) and 50µg/mL (kanamycin, Km).

Construction of plasmids to create *V. cholerae* mutants, overexpression

constructs and transcriptional fusions.

V. cholerae mutants were created by allelic exchange with the suicide plasmid

pCVD442 [48]. ca. 1kbp upstream and downstream DNA fragments flanking each

coding regions were PCR-amplified with the primers listed in Table S3. Upstream

and downstream DNA fragments were spliced together by overlapping PCR. The

resulting ca. 2 Kb fragments were digested as indicated in table S3 and cloned

into   pCVD442. For allelic exchange of *V. cholerae arsC* by *E. coli arsC,* similar

protocol was followed, but in this case the *V. cholerae* upstream and downstream

DNA regions of *V. cholerae arsC* were spliced together with the *E. coli arsC* gene

by overlapping PCR (see tables S2 and S3). Constructs were transformed into *E.*

coli DH5α λpir for amplification. They were confirmed by sequencing, transformed

into the *E. coli* donor strain SM10 λpir and conjugated for 6 hours at 37°C with *V.*

cholerae C6706 by mixing equal volumes (1ml) of exponential phase cultures,

and spot-plating. Single crossover *V. cholerae* were selected on LB plates with

Sm and Cb. Re-streaked single colonies were then plated on salt-free LB agar

containing 10% (w/v) sucrose and Sm. Colonies were streaked on carbenicillin

plates to confirm loss of pCVD442 and then checked by PCR for successful

deletion mutants.

Overexpression of *V. cholerae* and *E. coli* genes was carried out by using the

arabinose inducible expression vectors pBAD33 and pBAD18, and the IPTG-

inducible expression vector pHL100 (22) (see table S2 for specific constructs use

for each enteropathogen). To construct each overexpressing plasmid, genes open

reading frames including ribosome binding sites were PCR amplified (see primers

used in table S3), double digested with indicated restriction enzymes and cloned

into each plasmid. Constructs were transformed into the indicated strain by

electroporation.

For construction of the transcriptional reporter fusions, gene promoter regions of

target genes were amplified using the primers pair indicated in table S3. The PCR

product was double digested with HindIII and EcoRI and then ligated into HindIII-

EcoRI-digested pCB192N (23). Cloned plasmids were then transformed into *V.*

cholerae by electroporation.

Purification of recombinant VarG, VC2000, VC1041 and VarH.

V. cholerae VarG, VC1041 and VarH proteins were overexpressed using the *E.*

coli strain BL21 as 6×His-tagged enzymes from a pET28b vector. Overnight

cultures were diluted into 250 mL LB broth and grown until $OD_{600}\square=\square0.5$. Flasks

were supplemented with 1 mM IPTG and shaking at 37^{o} C for 2 h. Cells were

pelleted, and resuspended in equilibration buffer (Tris 50 mM, pH 7.5, 50 mM

NaCl, with protease inhibitor cocktail (Roche)) and disrupted by passaging once

through a French press. Lysates were then centrifuged for 1 h (25,000 rpm,

Beckman Coulter Avanti J26-XP centrifuge, JL-25.50 rotor) at 4°C. Nickel-NTA

resin (0.5 ml resuspended in equilibration buffer) was then added to the

supernatant, followed by incubation at 4°C in a rotating wheel. The lysate was

separated from the resin by centrifugation for 1 minute at 3,220 g and the resin

washed (5×10 ml) with washing buffer (equilibration buffer adjusted to 1 M NaCl)

and eluted with 2ml of washing buffer containing 500mM imidazole. Fractions

were subjected to SDS-PAGE and Coomassie Brilliant Blue staining for purity

assessment. Protein concentration was quantified with a Bradford assay.

Proteomic and thermal proteome profiling (TPP).

We performed thermal proteome profiling to determine protein abundance and

thermal stability changes in WT, and *var* mutant strains upon exposure to As^V

similarly to what was previously described (24, 25). Briefly, WT and mutant cells

grown to an OD578 of 0.5 were incubated with 1 mM arsenate (or a similar

volume of water) for 10 min. Cells were then washed with 10 ml PBS (for cells

treated with arsenate, 1 mM arsenate were included in the wash step) and

aliquoted to a PCR plate. Each aliquot was then exposed to a different

temperature in the range 37ºC-66.3ºC for 3 min. Following a 3 min incubation at

room temperature, cells were lysed with lysis buffer (final concentration: 50 µg

ml^{-1} lysozyme, 0.8% NP-40, 1 × protease inhibitor (Roche), 250 U ml^{-1}

benzonase, and 1 mM MgCl$_2$ in PBS) for 20 min followed by three freeze–thaw

cycles. Lysates were then prepared for analysis by mass spectrometry by

digesting proteins using a modified sp3 protocol (26), labeling peptides with

TMTpro (Thermo Fisher Scientific) and pooling samples from the same

temperature together. These samples were then fractionated to six fractions with

high pH fractionation and injected on an Orbitrap Q-Exactive Plus (Thermo Fisher

Scientific) coupled to liquid chromatography (details on the run conditions and

instrument parameters as in Mateus et al. (2020) (24, 25).

Mass spectrometry raw data was searched against the Vibrio cholerae FASTA file

(UP000000584 downloaded from UniProt) using the Mascot 2.4 (Matrix Science)

search engine and isobarquant (27). Protein abundance and thermal stability

changes were determined using limma (28) using the same algorithm as in

Mateus et al. (2020)(24).

Phosphorylation state of VarG.

To identify VC1069 GAPDH phosphorylation state*, V. cholerae* wild type and

Δ*varH* strains carrying the overexpression vector pHL100 containing the *varG*-HIS

tagged gene cloned were grown in the presence of As^V and with IPTG. Cells were

pelleted, disrupted, load and run in SDS-PAGE gel. VarG-HIS bands were

excised and diluted with NuPAGE LDS sample bu□er (Invitrogen, Carlsbad, CA)

mixed with10 mM dithiotreitol (DTT) and incubated at 56 °C for 20 min followed by

addition of iodoacetamide (IAA) to a final concentration of 20 mM. The sample

was loaded on 4-12% NuPAGE (Invitrogen). The electrophoresis was run in

MOPS bu□er at 180 V for 1 h. The gel was stained with Coomassie Brilliant Blue.

Bands of interest were cut from the SDS-PAGE. The bands were digested with

trypsin, followed by identification using liquid chromatography–tandem mass

spectrometry (LC–MS/MS). MS/MS spectra were searched with Proteome

Discoverer 2.3 (Thermo Fisher Scientific) against the two sequences obtained for

phosphate groups (UniProtKB). The precursor tolerance and fragment tolerance

were set to 10 ppm and 0.05 Da, respectively. Trypsin was selected as enzyme,

methionine oxidation, phosphorylation of serine, tyrosine, and tryptophan, and

deamidation of asparagine and glutamine were treated as dynamic modification

and carbamidomethylation of cysteine as a fixed modification (see supplementary

methods for further information).

Transposon-insertion sequencing (TIS) analysis.

TIS was carried out essentially as previously described (29) but using selection

plates containing LB medium with As^V 1 mM. In brief, ~600,000 transposon

mutants were generated by conjugation of *V. cholerae* C6706 with SM10λ*pir E.*

coli carrying the Himar1 suicide transposon vector pSC189 (30). They were

collected and their genomic DNA was pooled and analyzed on an Illumina MiSeq

bench-top sequencer (Illumina, San Diego, CA). Insertion sites (which included

35% of TA sites) were identified as previously described (29), and significance

was determined using Con-Artist simulation-based normalization as described

(31). Results were visualized using Artemis (32).

As^V resistance screen of arrayed transposon mutant library.

A non-redundant transposon insertion library from the *Vibrio cholerae* strain

C6706 (33) consisting of 3,156 insertion mutants was used to determine genetic

determinants providing resistance to As^V in *V. cholerae. V. cholerae* WT and the

library insertion mutants were first grown aerobically in 200 µl of LB in 96-well

plates at 37ºC for 24 hours. These cultures were then used as inoculum into 96-

well plates containing 200 µl complete LB medium, in the presence of 1 mM of

As^V. The 96-well plates were incubated under agitation at 37ºC for 24 hours and

then resistance to As^V was evaluated by measuring optical density (OD$_{600}$) using

a BioTek Eon plate reader.

Protein homology studies of As^V resistance determinants in enteric

pathogens.

DNA sequences of the *Vibrio cholerae arsR, varG, varH and arsJ var* genes and

complete genomes sequences of the enteric pathogens: *Salmonella typhimurium*

(PRJNA231221), *Citrobacter rodentium* (PRJNA261533), *Shigella flexeneri*

(PRJNA553848), EHEC (PRJNA218110) and *Yersenia pseudotuberculosis*

(PRJEB6403) were downloaded from NCBI database. Complete genomes of the

pathogens were translated by Prodigal ver. 2.6.2 and homologous sequences to

the *V. cholerae* Var proteins were queried into the enteric pathogens by BlastP

ver. 2.9.0. Sequence homology data were represented in a heatmap diagram with

Spearman correlation done by R ver. 3.6.1.

β-Galactosidase activity determination.

β-Galactosidase activity was measured through ONPG cleavage by the product of the *lacZ* reporter gene, and specific activity was calculated in Miller units (34). In brief, cultures of *V. cholerae* carrying the pCB192N plasmid harboring promoter gene insertions (see table S2) were grown at 37ºC overnight in 15 ml falcons tube containing 3 ml of LB complete medium. Overnight cultures were diluted 1:200 and supplemented with 1 mM of As^{V} and incubated at 37ºC until they reached mid-log phase (OD_{600} 0.4-0.7). A 100 µl aliquot of three different subcultures were collected, cells were permeabilized and assayed in triplicate for each strain as previously described (34).

Determination of intracellular and extracellular As^{V} and As^{III} in *V. cholerae*.

To determinate concentrations of As^{V} and As^{III} in supernatants of *V. cholerae* and *E. coli,* bacterial cells were grown in the presence of As^{V} for 5 hours and then pelleted. Supernatants of each strain were speciated by high-performance liquid chromatography (HPCL) (Agilent Infinity II 1290) using a C18 column reverse-phase (130 Å, 1.7 µm, 2.1 mm by 150 mm; Waters, USA). Elution conditions used were flow rate 1 ml min^{-1}; temperature 25 °C; isocratic elution in 5 mM tetrabutylammonium hydroxide, 5 % methanol (v/v) and 3 mM malonic acid. Identification and quantification of As^{V} and As^{III} was performed by inductively coupled plasma mass spectroscopy (ICP-MS) (Agilent 8900 ICP-MS Triple Quad) by comparison to standards of known concentration and peak integration.

To determine intracellular concentrations of As^V and As^{III} in *V. cholerae,* the

pelleted cells were washed in complete LB medium to remove traces of arsenic,

then resuspended in 500 µl of H$_2$O. Glass beads were adding to the resuspension

and cells were lysed by bead-beating. The resulting extract was centrifuged, and

As^V and As^{III} from those supernatants were analyzed by ICP-MS as previously

described. Samples were analyzed in triplicate.

LC-MS/MS analysis using QTRAP6500+

To measure 3-phosphoglycerate (3PG), *V. cholerae* cells grown in the presence

of 5 mM of As^V were pelleted and used for LC-MS/MS analysis. Bacterial pellets

were resuspended in 1 ml ice-cold 60% ethanol containing 2 µM ribitol as the

internal control. The cells were mechanically disrupted using a bead homogenizer

set to oscillate for 3 cycles of 30 sec each at 6800 rpm with a 10 sec pause

between each cycle. Cell debris was separated at 13,000 rpm. An aliquot (200µl)

of the supernatant containing intracellular metabolites was vacuum dried and

resuspended in an equal volume of 7.5 mM ammonium bicarbonate solution

before LC-MS/MS analysis.

A triple-quadrupole-ion trap hybrid mass spectrometer, QTRAP® 6500+

(Sciex, USA) connected with a Waters ultra-performance liquid chromatography I-

class (UPLC) system was used for metabolite analysis. The chromatographic

separation was performed on an XSELECT HSS XP column (150 mm

×⬚2.1⬚mm ID; 2.5⬚µm particle size, Waters, USA) using a binary solvent system

at a flow rate of 0.1⬚ml/min. Mobile phase A was composed of 7.5 mM

ammonium bicarbonate in LC-MS grade water and mobile phase B was 100%

LC-MS grade methanol. The column was maintained at 40°C, and the

autosampler temperature was set to 7°C. The A/B solvent ratio was maintained at

100/0 for 2⬚minutes, followed by a gradual increase of solvent B to 95% for 2

minutes. The solvent B was maintained at 95% over the next 5 minutes. The

gradient was reduced to 100% solvent A within 0.5⬚min and column was

equilibrated for 5.5⬚minutes before the next run. The needle was washed with 1.2

ml of a strong wash solution containing 100% LC-MS-grade acetonitrile followed

by 1 ml of a weak wash solution comprised of 10% aqueous methanol before

each injection. The injection volume was 5µl. The QTRAP® 6500+ (SCIEX) was

operated in negative as well as positive ion mode for targeted quantitation in

Multiple Reaction Monitoring (MRM). MRM parameters for each analyte are listed

in Table 1. The electrospray ionization (ESI) parameters used are as follows:

electrospray ion voltage of -4500V in negative and 5500V in positive mode,

source temperature of 400°C, curtain gas of 35, and gas 1 and 2 of 40 psi each,

respectively. The compound-specific parameters such as declustering potential

(DP) and collision energy (CE) were optimized for each compound using manual

tuning. These values are listed in Table 1.

Table 1: MRM transitions, declustering potentials and collision energies of

metabolites

Analyte	Q1 (m/z)	Q3 (m/z)	DP (V)	CE (V)
3-phospho glycerate	187.2	98.9	+60	19.2
Ribitol	150.8	88.9	-60	-14.2

Detection of 1As3PG.

1As3PG was identified using a Thermo Orbitrap Exploris 480 high resolution

mass spectrometer. The chromatographic method used for this analysis was the

same as that described for the LC-MS/MS performed using the QTRAP6500+.

The MS data were acquired in both positive and negative polarity within mass

range of 140- 400 Da. The precursor ion resolution was maintained at 240,000,

whereas the product ion resolution at 120,000. Deprotonated [M-H]$^-$ ion of

1As3PG was detected at m/z 304.8696 Da compared to the actual mass of

304.8674 Da, thus Δm = 7.149 ppm. The fragmentation pattern of 1As3PG could

not be further resolved under these conditions.

Proteins structural and topological analysis.

Protein alignments were performed by using Vector NTI. Prediction of VarH

secondary structure was performed by using Phyre2

(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index). Structural prediction

of the VarG and VarH proteins was performed with AlphaFold2 on the CoLabFold

publicly accessible interface (35)

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF

old2.ipynb). Sequences were modeled as monomers using mmseqs2 for multiple

sequence alignment. Studies on proteins topological similarity was carried out by

using TM-score (https://zhanggroup.org/TM-score/).

Competition assays.

Aerobic overnight cultures of *V. cholerae* C6706 wild type lacZ⁻ and

Enterohemorragic Escherichia coli EHEC O157:H7 *lacZ⁺*, *Salmonella enterica*

serovar Typhimurium LT2 lacZ⁺, *Citrobacter rodentium* DBS100 *lacZ⁺*, *Shigella*

flexneri M90T *lacZ⁺*, *and Yersinia pseudotuberculosis* YPIII *lacZ⁺,* were collected

and 1 x 10⁷ cells co-inoculated at 1:1 ratio in 15 ml tubes containing 3 ml of

complete LB medium without or with 10 mM As^V. After 8 hours of incubation at

37°C, an aliquot was diluted and plated on LB agar plates containing X-Gal to

enumerate wild type and competing strains. Competitive indices (CIs) were

determined by dividing the ratio mutant or non-wild type strain to C6706 wild type

colonies by the ratio in the inoculum. To calculate CIs among *V. cholerae* strain

C6706 against a different *V. cholerae* strain against enteric pathogens (Fig. 1B),

CIs obtained after the As^V challenge were normalized respect CIs obtained after

competition in the absence of As^V.

in vitro **GAPDH and phosphatase enzymatic assays.**

The GAPDH activity assay was performed in accordance with manufacturer's

instructions (MAK27; Sigma-Aldrich) but using P_i or As^V as the GAPDH substrate.

10 µg of purified *V. cholerae* VarG and VC2000 enzymes were used in the assay.

This assay quantifies GAPDH activity by measuring enzymes capacity to reduce

NAD^+ to NADH during the conversion of glyceraldehyde 3-phosphate to 1,3-

biphosphoglycerate. NADH produced in this reaction results in a colorimetric

(OD_{450}) product proportional to the GAPDH activity of the enzyme. GAPDH

activity is reported as nmole/min/mL = milliunit/mL. One unit of GAPDH is the

amount of enzyme that will generate 1.0 mmole of NADH per minute at pH 7.2 at

37 °C.

Phosphatase activity was quantified using Cayman's Phosphatase

Colorimetric assay kit and 5 µg of purified *V. cholerae* VarH, C113G R119G VarH,

VC1041 (positive control) and BsrV (negative control) enzymes. The assay uses

p-nitrophenylphosphate (pNPP) as substrate for phosphatase enzymes.

Phosphatase enzymes dephosphorylate pNPP, which deprotonates its phenolic

OH-group by increasing the pH of the reaction, deprotonated pNP yields an

intense yellow color detected at OD 405 nm. Enzymatic activities were calculated

as µmol of phosphate released per min/ml.

Flow cytometry analysis

V. cholerae WT and indicated *var* mutant strains were grown for 5 hours in the

presence of 1 mM As^V, then Aliquots (1 ml) from each culture were pelleted,

washed once, and resuspended in PBS (100 µl) to measure membrane potential

and ROS.

To measure *membrane potential* in *V. cholerae* cells, we used the BacLight

bacterial membrane potential kit (Invitrogen). *V. cholerae* samples resuspended in

100 µl PBS were incubated for 15 min at 20 °C in the dark with the dye 30 µM

3,3'-diethyloxacarbocyanine iodide (DiOC$_2$). Next, cells were washed twice with

PBS and resuspended in 2 ml of PBS, then fluorescence emitted from treated

cells was measured with a Bio-Rad S3e cell sorter at an excitation wavelength of

488 nm and an emission wavelength of 525/30 nm (for green) or 655 nm (for

red).

ROS measurements were performed using the CellROX™ Green Reagent, for

oxidative stress detection (ThermoFisher). Washed *V. cholerae* cells were

supplemented with CellROX® Reagent at a final concentration of 5 µM and

incubate them for 30 minutes at 37°C. Next, cells were washed twice with PBS

and resuspended in 2 ml of PBS, then fluorescence emitted was measured with a

Bio-Rad S3e cell sorter at an excitation wavelength of 488 nm and an emission

wavelength of 525 nm.

Statistics and reproducibility

Statistical significance was assessed with the Student's *t*-test where indicated. A p-value of less than 0.05 was considered statistically significant. Assays were performed with three biological replicates unless otherwise indicated.

REFERENCES

1. Kalia K, Joshi DN. 2009. Detoxification of Arsenic. Handb Toxicol Chem Warf Agents 1083–1100.

2. Singh R, Singh S, Parihar P, Singh VP, Prasad SM. 2015. Arsenic contamination, consequences and remediation techniques: A review. Ecotoxicol Environ Saf 112:247–270.

3. Pal C, Asiani K, Arya S, Rensing C, Stekel DJ, Larsson DGJ, Hobman JL. 2017. Metal Resistance and Its Association With Antibiotic Resistance. Microbiol Met Ions, 1st ed. 70:261–313.

4. Hobman JL, Crossman LC. 2015. Bacterial antimicrobial metal ion resistance. J Med Microbiol 64:471–497.

5. Willsky GR, Malamy MH. 1980. Characterization of two genetically separable inorganic phosphate transport systems in Escherichia coli. J

Bacteriol 144:356–365.

6.   Willsky GR, Malamy MH. 1980. Effect of arsenate on inorganic phosphate

transport in Escherichia coli. J Bacteriol 144:366–374.

7.   Dixon HBF. 1996. The Biochemical Action of Arsonic Acids Especially As

Phosphate Analogues. Adv Inorg Chem 44:191–227.

8.   Rosen BP. 2002. Biochemistry of arsenic detoxification. FEBS Lett 529:86–

92.

9.   Yang HC, Fu HL, Lin YF, Rosen BP. 2012. Pathways of Arsenic Uptake and

Efflux, p. 325–358. *In* Current Topics in Membranes. NIH Public Access.

10.   Scott N, Hatlelid KM, MacKenzie NE, Carter DE. 1993. Reactions of Arsenic

(III) and Arsenic (V) Species with Glutathione. Chem Res Toxicol 6:102–

106.

11.   Przygoda G, Feldmann J, Cullen WR. 2001. The arsenic eaters of Styria: A

different picture of people who were chronically exposed to arsenic. Appl

Organomet Chem 15:457–462.

12.   Liu J, Lu Y, Wu Q, Goyer R a, Waalkes MP. 2008. Mineral Arsenicals in

Traditional Medicines: Perspect Pharmacol 326:363–368.

13.   Shahpiri A, Mohammadzadeh A. 2018. Bioaccumulation of Arsenic by

Engineered Escherichia coli Cells Expressing Rice Metallothionein

Isoforms. Curr Microbiol 75:1537–1542.

14.   Thomas DJ, Rosen BP. 2013. Arsenic Methyltransferases, p. 138–143. *In*

Kretsinger, RH, Uversky, VN, Permyakov, EA (eds.), Encyclopedia of

Metalloproteins. Springer New York, New York, NY.

15.   Saltikov CW, Newman DK. 2003. Genetic identification of a respiratory

arsenate reductase. Proc Natl Acad Sci U S A 100:10983–10988.

16.   Oden KL, Gladysheva TB, Rosen BP. 1994. Arsenate reduction mediated

by the plasmid‐encoded ArsC protein is coupled to glutathione. Mol

Microbiol 12:301–306.

17.   Chen J, Yoshinaga M, Garbinski LD, Rosen BP. 2016. Synergistic

interaction of glyceraldehydes-3-phosphate dehydrogenase and ArsJ, a

novel organoarsenical efflux permease, confers arsenate resistance. Mol

Microbiol 100:945–953.

18.   Wu S, Wang L, Gan R, Tong T, Bian H, Li Z, Du S, Deng Z, Chen S. 2018.

Signature arsenic detoxification pathways in Halomonas sp. Strain GFAJ-1.

MBio 9:1–9.

19.   Fekih I Ben, Zhang C, Li YP, Zhao Y, Alwathnani HA, Saquib Q, Rensing C,

Cervantes C. 2018. Distribution of arsenic resistance genes in prokaryotes.

Front Microbiol 9:1–11.

20.   Németi B, Anderson ME, Gregus Z. 2012. Glutathione synthetase promotes
the reduction of arsenate via arsenolysis of glutathione. Biochimie 94:1327–
1333.

21.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA,
Tomita M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12
in-frame, single-gene knockout mutants: The Keio collection. Mol Syst Biol
2:1–11.

22.   Guzman L, Belin D, Carson MJ. 1995. Tight Regulation, Modulation, and
High-Level Expression by Vectors Containing the Arabinose PBAD
Promoter. J Bacteriol 177:4121–4130.

23.   Schneider K, Beck CF. 1986. Promoter-probe vectors for the analysis of
divergently arranged promoters. Gene 42:37–48.

24.   Mateus A, Hevler J, Bobonis J, Kurzawa N, Shah M, Mitosch K, Goemans C
V., Helm D, Stein F, Typas A, Savitski MM. 2020. The functional proteome
landscape of Escherichia coli. Nature 588:473–478.

25.   Mateus A, Kurzawa N, Becher I, Sridharan S, Helm D, Stein F, Typas A,
Savitski MM. 2020. Thermal proteome profiling for interrogating protein
interactions. Mol Syst Biol 16:1–11.

26.   Hughes CS, Moggridge S, Müller T, Sorensen PH, Morin GB, Krijgsveld J.
2019. Single-pot, solid-phase-enhanced sample preparation for proteomics

experiments. Nat Protoc 14:68–85.

27. Franken H, Mathieson T, Childs D, Sweetman GMA, Werner T, Tögel I,

Doce C, Gade S, Bantscheff M, Drewes G, Reinhard FBM, Huber W,

Savitski MM. 2015. Thermal proteome profiling for unbiased identification of

direct and indirect drug targets using multiplexed quantitative mass

spectrometry. Nat Protoc 10:1567–1593.

28. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015.

Limma powers differential expression analyses for RNA-sequencing and

microarray studies. Nucleic Acids Res 43:e47.

29. Chao MC, Pritchard JR, Zhang YJ, Rubin EJ, Livny J, Davis BM, Waldor

MK. 2013. High-resolution definition of the Vibrio cholerae essential gene

set with hidden Markov model-based analyses of transposon-insertion

sequencing data. Nucleic Acids Res 41:9033–9048.

30. Chiang SL, Rubin EJ. 2002. Construction of a mariner-based transposon for

epitope-tagging and genomic targeting. Gene 296:179–185.

31. Pritchard JR, Chao MC, Abel S, Davis BM, Baranowski C, Zhang YJ, Rubin

EJ, Waldor MK. 2014. ARTIST: High-Resolution Genome-Wide Assessment

of Fitness Using Transposon-Insertion Sequencing. PLoS Genet

10:e1004782.

32. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012. Artemis:

An integrated platform for visualization and analysis of high-throughput

sequence-based experimental data. Bioinformatics 28:464–469.

33.   Cameron DE, Urbach JM, Mekalanos JJ. 2008. A Defined Transposon

Mutant Library and Its Use in Identifying Motility Genes in Vibrio cholerae.

Proc Natl Acad Sci U S A 105:8736–8741.

34.   Miller JH. 1972. Experiments in Molecular Genetics. New York, NY: Cold

Spring Harbor Laboratory.

35.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,

Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C,

Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R,

Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger

M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior

AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein

structure prediction with AlphaFold. Nature 596:583–589.

36.   Demel S, Shi JIN, Martin P, Rosen BP, Edwards BFP. 2004. Arginine 60 in

the ArsC arsenate reductase of E . coli plasmid R773 determines the

chemical nature of the bound As (III) product. Protein Sci 2330–2340.

37.   Martin P, Demel S, Shi J, Gladysheva T, Gatti DL, Rosen BP, Edwards

BFP. 2001. Insights into the Structure, Solvation, and Mechanism of ArsC

Arsenate Reductase, a Novel Arsenic Detoxification Enzyme. Structure

9:1071–1081.

38.   Jenkins JL, Tanner JJ. 2006. High-resolution structure of human D-

glyceraldehyde-3-phosphate dehydrogenase. Acta Crystallogr Sect D Biol

Crystallogr 62:290–301.

39.   Velick SF. 1955. Glyceraldehyde-3-phosphate dehydrogenase from muscle.

Methods Enzymol 1:401–406.

40.   Byers LD, She HS, Alayoff A, Gregus Z, Németi B. 2005. Interaction of

Phosphate Analogues with Glyceraldehyde-3-phosphate Dehydrogenase.

Biochemistry 85:859–869.

41.   Byers LD, She HS, Alayoff A. 1979. Interaction of Phosphate Analogues

with Glyceraldehyde-3-phosphate Dehydrogenase. Biochemistry 18:2471–

2480.

42.   Aoyama K, Nakaki T. 2015. Glutathione in cellular Redox homeostasis:

Association with the excitatory amino acid carrier 1 (EAAC1). Molecules

20:8742–8758.

43.   Modulates O, Activity O. 2017. OxyR2 Modulates OxyR1 Activity and Vibrio

cholerae Oxidative Stress Response.

44.   Heidelberg JF et al. 2000. DNA sequence of both chromosomes of the

cholera pathogen Vibrio cholerae. Nature 406:477–483.

45.   Jin Q. 2002. Genome sequence of Shigella flexneri 2a: insights into

855   pathogenicity through comparison with genomes of Escherichia coli K12

856   and O157. Nucleic Acids Res 30:4432–4441.

857   46.   Batzilla J, Höper D, Antonenka U, Heesemann J, Rakin A. 2011. Complete

858   genome sequence of Yersinia enterocolitica subsp. palearctica Serogroup

859   O:3. J Bacteriol 193:2067–2065.

860   47.   Pearson BM, Gaskin DJH, Segers RPAM, Wells JM, Nuijten PJM, Van Vliet

861   AHM. 2007. The complete genome sequence of Campylobacter jejuni strain

862   81116 (NCTC11828). J Bacteriol 189:8402–8403.

863   48.   Tolen TN, Xie Y, Hernandez AC, Everett GFK. 2016. Complete genome

864   sequence of Salmonella enterica serovar TyphimuriumLT2. Nature

865   413:852–856.

866   49.   Popov G, Fiebig-Comyn A, Shideler S, Coombes BK, Savchenko A. 2019.

867   Complete Genome Sequence of Citrobacter rodentium Strain DBS100 .

868   Microbiol Resour Announc 8:1–2.

869   50.   Shi W, Wu J, Rosen BP. 1994. Identification of a putative metal binding site

870   in a new family of metalloregulatory proteins. J Biol Chem 269:19826–

871   19829.

872   51.   Sirover MA. 2011. Biochimica et Biophysica Acta On the functional diversity

873   of glyceraldehyde-3-phosphate dehydrogenase □: Biochemical mechanisms

and regulatory control. BBA - Gen Subj 1810:741–751.

52.   Ferreira E, Giménez R, Alexandra M, Aguilera L, Aguilar J, Badia J,

Baldomà L. 2015. Glyceraldehyde-3-phosphate dehydrogenase is required

for efficient repair of cytotoxic DNA lesions in Escherichia coli. Interntional J

Biochem Cell Biol 60:202–212.

53.   Whitworth DE, Morgan BH. 2015. Synergism between Bacterial GAPDH

and OMVs: Disparate Mechanisms but Co-operative Action. Front Microbiol

6:1–6.

54.   Leveson-Gower RB, Mayer C, Roelfes G. 2019. The importance of catalytic

promiscuity for enzyme design and evolution. Nat Rev Chem 3:687–705.

55.   Shi H, Shi X, Liu KJ. 2004. Oxidative mechanism of arsenic toxicity and

carcinogenesis 67–78.

56.   Liu SX, Athar M, Lippai I, Waldren C, Hei TK. 2001. Induction of oxyradicals

by arsenic: Implication for mechanism of genotoxicity. Proc Natl Acad Sci U

S A 98:1643–1648.

57.   Sukchawalit R, Prapagdee B, Charoenlap N, Vattanaviboon P, Mongkolsuk

S. 2005. Protection of Xanthomonas against arsenic toxicity involves the

peroxide-sensing transcription regulator OxyR. Res Microbiol 156:30–34.

58.   Wang H, Chen S, Zhang J, Rothenbacher FP, Jiang T, Kan B, Zhong Z,

Zhu J. 2012. Catalases Promote Resistance of Oxidative Stress in Vibrio

cholerae. PLoS One 7.

59.   Miki H, Funato Y. 2012. Regulation of intracellular signalling through

cysteine oxidation by reactive oxygen species. J Biochem 151:255–261.

60.   Mondal P, Majumder CB, Mohanty B. 2006. Laboratory based approaches

for arsenic remediation from contaminated water: Recent developments. J

Hazard Mater 137:464–479.

FIGURE LEGENDS.

Fig. 1 *V. cholerae* is the enteric pathogen with the greatest resistance to

As^V. A Growth curves (OD$_{600}$) of the enteric pathogens *V. cholerae*,

Enterohaemorragic Escherichia coli (EHEC), *Salmonella enterica*, *Citrobacter*

rodentium, *Shigella flexneri*, and *Yersinia pseudotuberculosis*. Cultures were

grown in LB medium in the absence (black) and in the presence (blue) of 10 mM

arsenate (As^V). **B** *in vitro* competition among *V. cholerae* and other

enteropathogenic bacteria (EPB) studied in panel A. Cells were incubated in LB

medium supplemented with 10 mM As^V for 8 hours. CI values from competition in

the presence of As^V were normalized with respect to CI values in the absence

As^V. **C** Growth curves (OD_{600}) of WT (blue filled circles) and *arsC* (blue empty

circles) mutant strains from *V. cholerae* and *E. coli* K12. Cultures were grown in

LB medium supplemented with 10 mM As^V. As^V and As^{III} concentrations were

determined by ICP-MS from supernatants of *V. cholerae* and *E. coli* WT and *arsC*

mutant strains after 8 hours of incubation in 1.5 mM of As^V. Data are the mean of

three biological replicates $\pm$ s.e.m.

Fig. 2   Identification of essential genetic determinants for As^V resistance in

V. cholerae. Left: Experimental workflow for an arrayed transposon mutant

screen to identify the genes required for As^V resistance in *V. cholerae* and the list

of mutant strains identified. Right: Experimental workflow for TIS screen in the

presence and absence of 1 mM As^V. Volcano plot depicting the ratio of read

counts mapped to individual genes in transposon libraries of *V. cholerae* plated

onto media supplemented with As^V). Red dotted lines indicate arbitrary thresholds

of fold change (FC) <0.1 and an inverse p-value >20. Genes in red color show loci

also identified in the arrayed screen (see Supplementary Table 1 for an extended

version).

Fig. 3   Phenotypic characterization of the *V. cholerae var* operon. Growth

curves (OD_{600}) of *V. cholerae* WT, single and combinatorial *var* mutant strains.

Cultures were grown in LB medium in the absence or presence of 1 and 10 mM

As^V. Data are the mean of three biological replicates ± s.e.m.

Fig. 4   Characterization of *V. cholerae* VarG. A *in vitro* GAPDH activity (OD_{450})

of purified *V. cholerae* VarG and the glycolytic Gap enzymes using P_i or As^V as

substrates. **B** β-galactosidase activity of strains in which the *lacZ* reporter gene is

cloned under the control of the *var* operon and the *gap* genes promoters. Cultures

were grown in LB medium in the absence or presence of 1 and 10 mM As^V for 5

hours. **C** Intracellular As^V concentrations of WT, *varG, varH* and *arsJ* mutant

strains grown with 1 mM As^V by ICP-MS. **D** Left: As^V detoxification model in *P.*

aeruginosa DK2 [1]**.** Right: Quantification of 1As3PG and 3PG from intracellular

and extracellular samples of *V. cholerae* cells grown with and without 1 mM As^V.

Data are the mean of three biological replicates ± s.e.m.

Fig. 5   Characterization of *V. cholerae* VarH. A Left: *in vitro* phosphatase

activity of purified *V. cholerae* VarH, and catalytic mutant derivative where

cysteine 113 and arginine 119 were replaced by glycine. *V. cholerae* protein-

tyrosine-phosphatase (VC1041) and broad-spectrum racemase (BsrV) were used

as positive and negative controls, respectively. Right: Growth curves (OD_{600}) of *V.*

cholerae WT, and a VarH catalytic mutant derivative . Cultures were grown in LB

medium in the presence of As^V. **B** Left: Intracellular 1As3PG concentrations from

V. cholerae WT and Δ*varH* cells grown with and without 1 mM As^V by HRMS.

Right: Schematic of putative mechanism for As^V detoxification in *V. cholerae*.

Data are the mean of three biological replicates □±□s.e.m.

Fig. 6   Oxidative stress response and membrane potential alterations of *V.*

***cholerae* WT and the *var* mutants. A** Volcano plot of LIMMA analysis output

depicting read fold change (log$_2$) of protein abundance and inverse P-value for

each protein queried in the proteomic screen in response to 1 mM As^V in *V.*

cholerae WT. Red dotted lines indicate arbitrary thresholds of fold change (log$_2$) of

>1 or <1 and an inverse P-value of >20. Var proteins are labelled in blue and

oxidative stress related proteins (Cystein biosynthesis and PrxA are labelled as in

B. **B** Schematic of the systems induced in response to As^V in *V. cholerae*. Non-

coloured arrows depict genes within an operon not detected. **C** Representative

dot plots from *V. cholerae* WT and *var* mutant cells stained with CellRox Green

following growth in the presence of 1 mM As^V for 5 hours. The black dotted gate

indicates an area with higher CellRox fluorescence (higher ROS levels). **D**

Volcano plot of LIMMA analysis output depicting read fold change (log$_2$) of protein

thermal stability and inverse P value for each protein queried in the proteomic

screen in response to 1 mM As^V in *V. cholerae* WT and *var* mutant strains. As^V

interacting proteins related to energy-generating pathways are labeled in blue

color.   PflB,   VC1866   formate   acetyltransferase.   AdhE,   VC2033   alcohol

dehydrogenase/acetaldehyde dehydrogenase. HemN, VC0116 oxygen-

independent coproporphyrinogen III oxidase. Icd, VC1141 isocitrate

dehydrogenase. FrdA, VC2656 fumarate reductase. IspH, 4-hydroxy-3-methylbut-

2-enyl diphosphate reductase. AceF, VC2413 pyruvate dehydrogenase, E2

component. UbiE, VC0083 ubiquinone/menaquinone biosynthesis

methyltransferase. **E** Representative dot plots from *V. cholerae* WT and *var*

mutant cells stained with DiOC2 following growth in the presence of 1 mM As^V for

5 hours. The black dotted gate indicates an area with lowered green fluorescence

(lower membrane potential). Data are the mean of three biological

replicates □±□s.e.m.

Fig. 7 *V. cholerae* **Var enhances As^V resistance in enteropathogenic**

bacteria. A Hierarchical heat map clustering of enteropathogenic bacteria with

respect to *V. cholerae* based on the percentage identity of *var* genes . **B** Growth

curves (OD$_{600}$) of the As^V-sensitive enteropathogenic bacteria clustered in panel

A, carrying an inducible empty plasmid (empty circle) or the same plasmid

expressingthe *V. cholerae var* operon (blue filled circle). Cultures were grown in

LB medium in the presence of 10 mM As^V. Data from panel B are the mean of

three biological replicates □±□s.e.m.

1000

1001

1002 **SUPPLEMENTARY INFORMATION**

1003 **Transient glycolytic complexation of arsenate enhances resistance in the**

1004 **enteropathogen *Vibrio cholerae***

1005 Emilio Bueno*, Víctor Pinedo, Dhananjay Shinde, André Mateus, Athanasios

1006 Typas, Mikhail M Savitski, Vinai Chittezham Thomas, and Felipe Cava*.

1007

1008 **Content – Supplementary Information**

1009 • Supplementary Figures S1 – S7

1010 • Supplementary Tables S1 - S3

1011 • Supplementary references.

1012

1013

1014

1015

1016

1017

1018

1019

Fig. S1 Growth curves (OD$_{600}$) of *V. cholerae* WT in the presence of As^{V}. *V. cholerae* WT was grown in LB medium in the presence of 10, 20 and 30 mM As^{V}. Data are the mean of three biological replicates ± s.e.m.

Fig. S2   Characterization of *V. cholerae* ArsC. A Schematic of the *arsC*-containing operons from *E. coli* K12 and *V. cholerae,* and protein sequence alignments. Black arrows depict ArsC catalytic cysteine and arginine residues implicated in As^{V} reduction to As^{III}. **B** Growth curves (OD$_{600}$) of *V. cholerae* WT, and two derivative strains where *arsC* has been replaced by *E. coli's arsC* (red, *arsC^{ec}*), and *arsC^{ec}* strain where *arsB* is overexpressed in trans (blue, *arsC^{ec}* + *arsB^{ec}*) (see bottom panel, and material and methods for further details). Cultures were grown in LB medium in the presence 10 mM As^{V}. Bottom panel: Explanatory

diagrams illustrating the expected output from each construct. Data are the mean

of three biological replicates ± s.e.m.

Fig. S3 Transcriptional analysis of the *var* operon. β-galactosidase activity

from *V. cholerae* WT and *arsR* mutant (Δ*arsR*) strains carrying DNA regions

upstream of *arsR, varG, varH* and *arsJ* genes fused to the *lacZ* reporter gene.

Each DNA region fused to *lacZ* is depicted by arrows. Cultures were grown in LB

medium in the presence or absence of 10 mM As^V. Data are the mean of three

biological replicates ± s.e.m.

Fig. S4   *V. cholerae* VarG presents a high degree of similarity to the

glycolytic GAPDH Gap. A Sequence alignments of the *V. cholerae* VarG and the

glycolytic GAPDH proteins using Vector NTI software. **B** Left: Predicted structures

of   VarG   using   AlphaFold2

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF

old2.ipynb). The region of the protein structure depicted in blue colour indicate the

least error in prediction. Right: VarG and Gap structures overlapping using

TMscore (https://zhanggroup.org/TM-score/). TM-score value of 1 indicates 100%

structural identity among proteins.

Fig. S5   Quantification of intra- and extracellular metabolites from *V.*

cholerae. Representative plots from identification of 1As3PG and 3PG from

intracellular and extracellular samples from *V. cholerae* WT and Δ*varG* cells were

grown with and without 1 mM As^V. The compounds were separated by UPLC

using an XSELECT HSS XP column. 1As3PG was identified using a Thermo

Orbitrap Exploris 480 high-resolution mass spectrometer and 3PG using a triple-

quadrupole-ion trap hybrid mass spectrometer QTRAP® 6500+ (Sciex, USA)

connected with a Waters ultra-performance liquid chromatography I-class (UPLC)

system.

Fig. S6   VarH characterization. A Sequences alignment of the secondary

structure of the VarH protein with the crystal structure of the human Kap PTP

phosphatase using the software Phyre2 ([www.sbg.bio.ic.ac.uk/phyre2).](www.sbg.bio.ic.ac.uk/phyre2) Human

Kap structure matched VarH with 100% confidence and 99% coverage. Amino

acid sequences within the black box (from the amino acid 113 (H) onwards)

indicate the conserved catalytic loop (C(x)$_5$R) of protein tyrosine phosphatases

PTP. Green and black boxes indicate deletions and insertions, respectively,

respect the template sequence. **B** Predicted structures of VarH with AlphaFold2

([https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF](https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb)

[old2.ipynb](https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb)). The regions of the protein structure depicted in blue colour indicate

no error in prediction. Right: overlapping between VarH protein structure obtained

with AlphaFold2 and crystal structure of the Kap PTP protein using TMscore

(https://zhanggroup.org/TM-score/).   TM-score   value   of   1   represents   100%

structural identity among proteins. **C** SDS-PAGE gel depicting the VarG His-tag

purified protein from *the V. cholerae* WT and *varH* mutant strains (see material

and methods for details). VarG proteins were excised from the gel and its

phosphorylation state was quantified by LC–MS/MS. **D** Phosphorylation state

quantification of VarG from *V. cholerae* WT and *varH* mutant strains. S204, serine

amino acid from VarG in position 204. Y108, tyrosine amino acid from VarG in

position 108. **E** Representative plots from identification of 1As3PG and 1AsG from

intracellular samples of *V. cholerae* WT and Δ*varH* cells grown with and without 1

mM As^V. The compounds were separated by UPLC using a XSELECT HSS XP

column. The detection of 1As3PG and 1AsG were perfomed using a Thermo

Orbitrap Exploris 480 High Resolution Mass Spectrometer. Data are the mean of

three biological replicates □±□s.e.m.

Fig S7 As^{III} binding site of ArsR is not conserved in *V. cholerae* and *varH* is

specific of the Vibrionaceae family. **A** ArsR amino acids sequences alignment

from *V. cholerae* and representative enteropathogenic bacteria using Vector NTI

software. ArsR cysteine-containing As^{III}-binding motives are depicted in the lower

panel with cysteine residues labeled in red color. **B** Genomic organization

comparison of the *var* operon in the Vibrionaceae family and representative

bacteria by using the Patric software (https://www.patricbrc.org/). The presence of

varH is depicted in green color and under dotted rectangle.

1090

1091 **Table S1.** Complete list of genes identified under-represented by TIS in the

1092 presence of 1 mM As^V with thresholds of fold change <0.1 and an inverse p-value

1093 >75

1094 **Table S2.** List of strains and plasmids used in this study.

1095 **Table S3.** List of primers used in this study.

1096

Supplementary references

1.   Chen J, Yoshinaga M, Garbinski LD, Rosen BP. Synergistic interaction of

glyceraldehydes-3-phosphate dehydrogenase and ArsJ, a novel

organoarsenical efflux permease, confers arsenate resistance. *Mol*

Microbiol 2016; **100**: 945–953.

2.   Miller VL, Mekalanos JJ. A novel suicide vector and its use in construction

of insertion mutations: Osmoregulation of outer membrane proteins and

virulence determinants in Vibrio cholerae requires toxR. *J Bacteriol* 1988;

170: 2575–2583.

3.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al.

Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: The Keio collection. *Mol Syst Biol* 2006; **2**: 1–11.

4.   Sansonetti PJ, Kopecko DJ, Formal SB. Involvement of a Plasmid in the

Invasive Ability of Shigella flexneri. 1982; **35**: 852–860.

5.   Cava F, De Pedro MA, Lam H, Davis BM, Waldor MK. Distinct pathways for

modification of the bacterial cell wall by non-canonical D-amino acids.

EMBO J 2011; **30**: 3442–3453.

6.   del Peso Santos T, Alvarez L, Sit B, Irazoki O, Blake J, Warner BR, et al.

BipA exerts temperature-dependent translational control of biofilm-

associated colony morphology in Vibrio cholerae. *Elife* 2021; **10**: 1–23.

7.   Donnenberg MS, Kaper JB. Construction of an eae deletion mutant of

enteropathogenic Escherichia coli by using a positive-selection suicide

vector. *Infect Immun* 1991; **59**: 4310–7.

8.   Chiang SL, Rubin EJ. Construction of a mariner-based transposon for

epitope-tagging and genomic targeting. *Gene* 2002; **296**: 179–185.

9.   Guzman L, Belin D, Carson MJ. Tight Regulation, Modulation, and High-

Level Expression by Vectors Containing the Arabinose PBAD Promoter. *J*

Bacteriol 1995; **177**: 4121–4130.

10.   Kimsey HH, Waldor MK. Vibrio cholerae LexA coordinates CTX prophage

gene expression. *J Bacteriol* 2009; **191**: 6788–6795.

1127

1128

1129

1130

As^V resistance determinants

Arrayed screen

3.156 Tn insertion mutants

-As^V

Aliquot transferred into:

LB +As^V

Output
Growth analysis at OD_600

TIS

V. Cholerae + Himar1 Tn

-As^V

Collection and inoculation into LB:

LB -As^V   LB +As^V

Output
Mutant library recovery

Comparison of Tn-seq profiles

vc1068	ArsR repressor protein
vc1069	GAPDH protein
vc1070	Putative phosphatase protein
vc1071	Membrane transporter protein

arsR

VC1068   VC1069   VC1070   VC1071

A

Phosphatase activity

B

1As3PG

A

Identity percentage

0 40 80

S. flexneri (PRJNA553848)

S. enterica (PRJNA231221)

Y. pseudotuberculosis (PRJEB6403)

EHEC *(PRJNA218110)*

C. rodentium (PRJNA261533)

V. cholerae C6706

ArsR   VarG   VarH   ArsJ

B

○ Empty plasmid   ● +*var* operon

EHEC

Y. pseudotuberculosis

S. enterica

C. rodentium

S. flexneri