Transient glycolytic complexation of arsenate enhances resistance in the

enteropathogen *Vibrio cholerae*

Emilio Bueno*[1], Víctor Pinedo[1], Dhananjay D. Shinde[2], André Mateus[3] Athanasios

Typas[3], Mikhail M Savitski[3], Vinai C. Thomas[2] and Felipe Cava*[1].

[1]Laboratory for Molecular Infection Medicine Sweden, Department of Molecular

Biology, Umeå Centre for Microbial Research, Umeå University, Umeå SE-90187,

Sweden.

[2] Center for Staphylococcal Research, Department of Pathology and Microbiology,

University of Nebraska Medical Center, Omaha, Nebraska, 68198, USA

[3] Genome Biology Unit, European Molecular Biology Laboratory (EMBL),

Heidelberg, Germany

Correspondence to:

felipe.cava@umu.se

emilio.bueno@umu.se

Keywords: arsenate, arsenite, quorum sensing, Tn-seq, *Vibrio cholerae*

Running title: Arsenate resistance in *Vibrio cholerae*.

Transient glycolytic complexation of arsenate enhances resistance in the

enteropathogen *Vibrio cholerae*

Abstract

The ubiquitous presence of toxic arsenate (As^V) in the environment has virtually

raised mechanisms of resistance in all living organisms. Generally, bacterial

detoxification of As^V relies on its reduction to arsenite (As^{III}) by ArsC, followed by

the export of As^{III} by ArsB. However, how pathogenic species resist this metalloid

remains largely unknown. Here, we found that *V. cholerae,* the etiologic agent of

the diarrheal disease cholera, outcompetes other enteropathogens when grown on

millimolar concentrations of As^V. To do so, *V. cholerae* uses, instead of ArsCB, the

As^V-inducible *vc1068-1071* operon (renamed *var* for *v*ibrio *a*rsenate *r*esistance)*,*

which encodes the arsenate repressor ArsR, an alternative glyceraldehyde-3-

phosphate dehydrogenase, a putative phosphatase, and the As^V transporter ArsJ.

In addition to Var, *V. cholerae* induces oxidative stress-related systems to counter

ROS production caused by intracellular As^V. Characterization of the *var* mutants

suggested these proteins function independently from one another and play critical

roles in preventing deleterious effects on the cell membrane potential and growth

derived from the accumulation As^V. Mechanistically, we demonstrate that *V.*

cholerae complexes As^V with the glycolytic intermediate 3-phosphoglycerate into 1-

arseno-3-phosphoglycerate (1As3PG). We further show that 1As3PG is not

transported outside the cell; instead, it is subsequently dissociated to enable

extrusion of free As^V through ArsJ. Collectively, we propose the formation of

1As3PG as a transient metabolic storage of As^V to curb the noxious effect of free

As^V. This study advances our understanding of As^V resistance in bacteria and

underscores new points of vulnerability that might be an attractive target for

antimicrobial interventions.

Introduction

Arsenic is a toxic metalloid commonly found in aquatic and terrestrial environments (1–3) as arsenate (As^V) or its reduced form, arsenite (As^{III}). The potent toxicity of this element makes it one of the best-studied natural poisons that impact public health (4). The mechanism of As^V toxicity is due to its structural resemblance to phosphate oxyanions and its uptake into the cell through phosphate transport channels (5, 6). Once in the cytoplasm, As^V can replace phosphate in energy-generating reactions resulting in the formation of ADP-As^V instead of ATP by the ATPase, and through substrate-level phosphorylation during glycolysis (7, 8). Conversely, cellular toxicity by As^{III} is exerted by a different mechanism. As^{III} can interact directly with thiol groups of proteins and molecules, thereby interfering with diverse cellular processes (9, 10). Despite its risk to public health, arsenic has been used as an antimicrobial agent to treat infectious diseases (1, 4, 11, 12), increasing the selective pressure for microbes to acquire arsenic resistance.

Bacterial strategies to resist As^V include chelation by metal-binding metallothioneins (13), methylation to less toxic and more volatile arsenic derivatives (14), respiration (15), and extrusion of As^V or As^{III} by specific efflux pumps (e.g., ArsJ and ArsB, respectively) (9, 16–18). In bacteria, the most common mechanism of As^V resistance genes occurs via As^V reduction to As^{III} followed by the extrusion of the latter through an As^{III} transporter (19). As^V resistance genes are commonly organized in operons, e.g., the *arsRDABC* operon from *Escherichia coli* R773 (19).

In this example, *arsC* encodes a dedicated arsenate reductase and *arsB* an efflux

pump of As^{III} (20). The *arsR* gene encodes the repressor of the *ars* operon. When

As^{III} is present, its interaction with ArsR releases it from the *ars* promoter, thus

increasing transcription of the operon. *arsA* encodes an ATPase that, together with

ArsB forms the As^{III} extrusion system. Alternatively, As^{III} extrusion via ArsB can be

driven by proton motive force in species lacking ArsA. Finally, *arsD* encodes a

chaperone that enhances arsenic extrusion by transferring As^{III} to ArsAB (9).

Despite being extensively studied in environmental soil and marine bacteria

species, yet the mechanisms underlying As^{V} resistance in human pathogenic

bacteria remain largely unknown. Here, we investigated As^{V} resistance in

representative enteric pathogens and found that *V. cholerae* exhibits remarkably

high resistance to this metalloid. Using genome mining and functional

characterization, we demonstrated that *V. cholerae* does not detoxify As^{V} by

reducing it to As^{III}. Instead, transposon-based functional screens identified the

operon *vc1068-1071* as the primary genetic determinant of As^{V} resistance in *V.*

cholerae. Mechanistic characterization of this system revealed that resistance to

As^{V} in *V. cholerae* is mediated by a transient metabolic complexation of As^{V} with

the glycolytic intermediate 3-phosphoglycerate (3PG) to generate 1-arseno-3-

phosphoglycerate (1As3PG). Our results support that this As^{V}-containing

metabolite is transient in the cell: its formation cushions the cellular damage caused

by free As^V, while its dissociation enables *V. cholerae* to release free As^V through

the ArsJ efflux permease.

Results

High resistance to arsenate in *V. cholerae* is independent of the ArsC reductase

While studying the tolerance of enteropathogenic bacteria to metalloids we observed that *Vibrio cholerae*, the causative agent of cholera, was able to grow in media supplemented with supraphysiological (30 mM) concentrations of arsenate (As^V) (Fig. S1). This level of resistance was unmatched by other human enteric pathogens, as lower concentrations of As^V (10 mM) readily caused total or partial growth inhibition of *Salmonella enterica*, *Citrobacter rodentium, Yersinia pseudotuberculosis, Enterohemorragic E. coli* (EHEC) *and Shigella flexneri* (Fig. 1A)*. Consistently, *V. cholerae* outcompeted these species by 5-500-fold in co-culture experiments grown in media supplemented with As^V (Fig. 1B).

To investigate the molecular mechanisms of As^V resistance in *V. cholerae*, we first searched for canonical determinants for As^V detoxification in its genome. *V. cholerae vc2165* encodes a homolog of *E. coli* K12 arsenate reductase ArsC, including the conserved catalytic cysteine and arginine residues (36, 37) (Fig. S2A). However, while *E. coli*'s ArsC enzyme is critical for resistance to As^V, inactivation of *vc2165* did not compromise *V. cholerae* growth in the presence of this metalloid (Fig 1C). Thus, we reasoned that *V. cholerae*'s ArsC might not be active or expressed. To measure ArsC activity we quantified extracellular As^V and As^{III} by inductively coupled plasma mass spectrometry (ICP-MS) and observed that while

E. coli reduced ~80% of As^V to As^{III} in an ArsC-dependent manner, *V. cholerae*

produced no As^{III} (Fig. 1C). Interestingly, replacement of *V. cholerae arsC* allele by

that of *E. coli* caused growth inhibition, likely due to toxic As^{III} formation, as growth

was recovered by expressing *E. coli's* As^{III}-transporter ArsB (Fig. S2B). Altogether,

these results indicate that *V. cholerae* ArsC is inactive and that As^V resistance in

this bacterium is independent of the ArsC-AsrB system that relies on the production

and elimination of As^{III}.

Resistance to As^V in *V. cholerae* depends on the *vc1068-vc1071* operon.

To identify the genetic determinants conferring resistance to As^V in *V. cholerae*, we

performed two complementary genome-wide screenings based on: ***i)*** Transposon

Insertion Sequencing (TIS), and ***ii)*** the use of a previously described *V. cholerae*

transposon mutant library [32] (Fig. 2). Both screens permit assessment of the

mutants' relative fitness. However, the first approach is more suitable for

discovering mutants with subtle fitness differences, while the latter can identify

candidates otherwise underappreciated by TIS due to cross-complementation

between mutants within a population. Only four non-essential Tn mutants

(*vc1068::tn, vc1069::tn, vc1070::tn,* and *vc1071::tn*) were mutually under-

represented in both screenings (Fig. 2). We renamed *vc1068-71* as ***var*** for ***v****ibrio*

a*rsenate* ***r****esistance cluster referred hereafter for reasons described below. *vc1068*

encodes an homolog of the ArsR repressor, *vc1069* (renamed as *varG*) encodes a

putative glyceraldehyde-3-phosphate dehydrogenase (GAPDH), *vc1070* (renamed

as *varH*) encodes an uncharacterized putative phosphatase, and *vc1071*, an ArsJ-

like As^V transporter (17). As expected, the *arsC* mutant was not identified in the

screens, confirming that *V. cholerae* does not detoxify As^V by reducing it to As^{III}.

Although homologues to VarG and ArsJ were previously reported in *Pseudomonas*

aeruginosa, their function in As^V resistance has not been studied *in vivo* (17).

To verify the implication of the *var* cluster on As^V resistance, we constructed

individual in-frame marker-less deletion mutants. All mutants were sensitive to As^V

except for $\Delta arsR$ (Fig. 3A), suggesting that the conditional lethality of the *arsR::*tn

was likely due to the polar effect on the expression of the downstream genes *varG-*

varH-arsJ. Indeed, transcriptional analysis using fusions of the *lacZ* reporter to

potential promoter regions located upstream of each *var* gene demonstrated that

expression of this cluster depends on As^{V,} and it is driven from a single promoter

upstream of *arsR* (Fig. S3). Inactivation of *arsR* turned transcription from this

promoter constitutive (both in presence and absence of As^V), confirming that

VC1068 is the *V. cholerae*'s ArsR repressor of the *var* operon (Fig. S3). Further

characterization revealed that $\Delta arsJ$ presents a more dramatic growth defect

compared to the rest of the *var* mutants at low As^V concentrations (1 mM) (Fig. 3).

However, at 10 mM of As^V, the growth of the *varG* and *varH* mutant strains was

also compromised despite the presence of ArsJ. These results suggest that the

ArsJ transporter is the main determinant for As^V resistance in *V. cholerae* but VarG

and VarH also become relevant at higher concentrations of As^V. Furthermore, the

inactivation of *varG, varH* and *arsJ* (double and triple mutants) aggravated growth

defects exhibited by the single mutants, suggesting independent functions for these

proteins in As^V resistance in *V. cholerae* (Fig. 3A).

164   VarG is an As^V-inducible GAPDH that preferentially binds As^V

V. cholerae VarG presents a high degree of protein sequence (E-value: 5^{e-86}) (Fig.

S4A) and structure (AlphaFold2 TM-score=0.96) (Fig. S4B) similarity to the

glycolytic GAPDH VC2000 (Gap), including the conserved catalytic Cys residue

(38). To characterize VarG's activity compared to the glycolytic paralogue Gap

particularly, we purified both proteins from *V. cholerae* and performed *in vitro*

GAPDH activity assays (39) in the presence or absence of As^V. While both proteins

exhibited GAPDH activity, VarG showed preference for As^V over P_i (the canonical

substrate) (Fig. 4A). Conversely, the glycolytic GAPDH Gap performed better with

P_i. Furthermore, while the expression of *Gap* remained low and constitutive,

transcription from the *var* promoter was strongly induced by As^V (Fig. 4B).

Therefore, these results suggest that while *V. cholerae* Gap is active on As^V, this

enzyme specific activity and protein levels are likely insufficient to replace VarG

essentiality in *V. cholerae* resistance to As^V.

Arsenate detoxification in *V. cholerae* is facilitated by extrusion of free arsenate instead of complexed as 1-arseno-3-phosphoglycerate

A VarG homolog from *P. aeruginosa* was proposed to function as a GAPDH that complexes As^V with glyceraldehyde-3-phosphate (G3P) into 1-arseno-3-phosphoglycerate (1As3PG), which is further eliminated through ArsJ (17). However, these experiments were performed *in vitro* using a commercial glycolytic GAPDH from rabbit instead of the As^V-specific GAPDH from *P. aeruginosa* DK2. If this model applies to *V. cholerae,* the absence of either VarG or ArsJ will similarly affect AsV export and resistance. However, our data shows that $\Delta varG$ did not phenocopy the growth defect of $\Delta arsJ$ (Fig. 3). Furthermore, analysis of intracellular levels of As^V by ICP-MS revealed that, in relation to the WT, while the $\Delta varG$ strain only showed moderate (ca. 3x) accumulation of As^V, inactivation of ArsJ accumulated this metalloid ca. 20x (Fig. 4C). Although these results suggest that VarG is not essential to export As^V, we cannot discard the possibility of As^V being complexed and extruded as 1As3PG in *V. cholerae*.

Using high-resolution mass spectrometry (HRMS), we detected a mass (*m/z*) of 304.8696 Da, specific to the cultures supplemented with As^V, which was compatible with 1As3PG (*m/z* 304.8674, *m/z*<0.003 Da difference between experimental and theoretical mass) (Fig. 4D and S5). 1As3PG was also detected in the $\Delta varG$, consistent with the ability of the constitutively expressed glycolytic Gap GAPDH to use As^V (Fig. S5). 1As3PG was detected in intracellular cytoplasmic

fractions but not in the extracellular milieu, suggesting that this molecule is not

expelled in *V. cholerae* (Fig. 4D and S5). As it has been reported that 1As3PG is

rapidly dissociated to As^V and 3PG (40), we reasoned that if exported, its

decomposition should increase extracellular 3PG levels in relation to cultures

without As^V. However, we found no significant differences in the levels of

extracellular 3PG between conditions supporting the idea that As^V detoxification

likely occurs through the release of free As^V rather than of 1As3PG through ArsJ

(Fig. 4C and S5).

VarH is involved in 1As3PG dissociation and arsenate export

To find clues that help us understand the role that 1As3PG plays in *V. cholerae*'s

resistance to As^V we investigated VarH. Structural *in silico* analysis using the

phyre2 tool (www.sbg.bio.ic.ac.uk/phyre2) and AlphaFold2 revealed that VarH is a

cysteine-based protein tyrosine phosphatase (PTP) belonging to the dual-

specificity phosphatase superfamily that presents a very high topology similarity to

the human Kap PTP phosphatase (Fig. S6AB). To determine if VarH has

phosphatase activity, we purified this protein and assessed its capacity to

dephosphorylate the chromogenic substrate pNPP. Our results show that,

compared to the wheat phosphatase or *V. cholerae's* PTP VC1041 positive

controls, VarH presents lower phosphatase activity, which turned off upon

replacement of the putative catalytic cysteine and arginine by glycine residues (Fig.

5A, left panel). Furthermore, consistent with the importance of these residues in

VarH activity, replacement of the WT *varH* allele by the catalytically inactive variant

varH C113G/R119G impaired *V. cholerae* growth in the presence of As^V (Fig. 5A,

right panel). These results suggest that the phosphatase activity of VarH is required

to provide As^V resistance in *V. cholerae*.

Since VarG and VarH are coregulated by As^V and both support resistance to this

metalloid, we hypothesized that VarH could regulate VarG activity by modulating its

phosphorylation state. To assess such a possibility, we purified VarG from *V.*

cholerae WT and Δ*varH* (Fig. S6C) and analyzed VarG phosphorylation by liquid

chromatography–tandem mass spectrometry (LC–MS/MS). However, even though

VarG was phosphorylated on Tyr108 and Ser204, those phosphorylation levels

were independent of VarH (Fig. S6D). Then, we reasoned that as P$_i$ and As^V are

analogs, VarH could potentially act on 1As3PG as a phosphatase to produce 1AsG

+ Pi or as an "arsenatase" to render 3PG + As^V. Remarkably, we found that 1As3PG

concentration were significantly elevated in Δ*varH* compared to those in the WT

strain (Fig. S6E), suggesting that VarH might use 1As3PG as a substrate. Although

we could not confirm VarH's activity *in vitro,* potentially due to the instability of

1As3PG (41), the fact VarH is an active phosphatase but 1AsG is not detected in

V. cholerae suggests that VarH could dissociate the 1As3PG complex into 3PG and

free As^V which would be released through ArsJ.

Intracellular arsenate impacts the *V. cholerae* proteome and leads to ROS accumulation and membrane potential defects

To broadly assess global responses triggered by As[V] in *V. cholerae*, we analyzed the pathogen proteome upon exposure to As[V]. Interestingly, a relatively low number of proteins (~20 proteins) were upregulated upon exposure to As[V], indicating that adaptations to this metalloid might not require extensive proteome rewiring in *V. cholerae*. As expected, these proteins included the As[V] resistance Var proteins but also amino acids permeases, a Zn/Cd transporter, members of the PTS-fructose system, the starvation stress response protein RaiA, and the cysteine synthesis and the peroxiredoxin PrxA (Fig. 6AB). Cysteine is one of the main targets of reactive oxygen species (ROS) and a functional component of glutathione, which plays a crucial role in the adjustment of cellular redox potential (42). In this line, PrxA has been involved in the adaptation to hydrogen peroxide in *V. cholerae* (43). Therefore, we reasoned that the *var* mutants, which display increased As[V] intracellular levels, might suffer from higher oxidative stress than *V. cholerae* WT strain. To test this hypothesis, we measured ROS levels in the *V. cholerae* WT and *var* mutant strains and observed a direct correlation between As[V] accumulation and ROS level ($\Delta arsJ$>$\Delta varH$>$\Delta varG$>WT) (Fig. 6C). These findings collectively indicate that AsV induces oxidative stress in *V. cholerae* which is exacerbated in the *var* mutants due to a supraphysiological accumulation of this toxic metalloid.

In addition to ROS affecting Cys-containing proteins, we cannot discard a direct

effect of As^V on the proteome of *V. cholerae*. To this end, we performed thermal

proteome profiling (TPP), a method that globally analyzes multiple types of protein

interactions (25). TPP data showed an increased thermal stability of multiple

cytosolic proteins that could potentially interact with As^V (~3-fold) in the *var* mutants

with respect to the WT (Fig. 6D and table S). As^V-interacting protein candidates

included proteins involved in the synthesis of heme groups of respiratory

cytochromes, respiratory ubiquinone, fermentation, and carbon catabolism,

suggesting that accumulation of As^V in Var-defective strains might interfere with cell

bioenergetics. To investigate this possibility, we monitored the bacterial membrane

potential in the presence of As^V using flow cytometry (Fig. 6E). Interestingly,

inactivation of the *var* system resulted in a significant depolarization of the mutant's

membrane. Together, these results demonstrate that AsV can, directly and

indirectly, affect the *V. cholerae*'s proteome, leading to oxidative stress and defects

in cellular bioenergetics, both possibly related to the growth impairment of the *var*

mutants.

***V. cholerae var operon* provides As^V resistance to enteric pathogens**

In silico analysis showed that, except for *C. rodentium*, ArsR is the only As^V-

resistance determinant conserved between *V. cholerae* and the As^V-sensitive

enteric pathogens (Fig. 7A). In general, these enteropathogenic species detoxify

As^V via ArsC-dependent reduction to As^{III}, and subsequent extrusion through ArsB.

As *V. cholerae* lacks the As^{III}- production/extrusion tandem ArsCB, we wondered

whether the *var* genes could also augment resistance to As^V in other species or

would rather require additional species-specific elements. To address this question,

we expressed the *var* operon excluding the ArsR repressor in enteropathogens

sensitive to As^V and studied their growth capacity in the presence of As^V. Strikingly,

induced coexpression of *varG-varH-arsJ* in several enteropathogens boosted their

growth on As^V (Fig. 7B), demonstrating that the Var system is sufficient and

compatible with the ArsCB system in providing resistance against As^V in other

bacteria.

DISCUSSION

Arsenic is ubiquitous in the environment and consequently, most living organisms, including bacteria, encode arsenic-resistance proteins (44–49). However, functional characterization of these systems and how they support bacterial fitness in environments with As^V is underappreciated. In this study, we describe that *V. cholerae* exhibits higher resistance to As^V than other enteric pathogens.

The strategy used by *V. cholerae* to resist As^V is different from that used by other gammaproteobacterial enteropathogens. While most of these species encode an ArsC arsenate reductase and an ArsB arsenite efflux pump, *V. cholerae* lacks the latter and has an inactive ArsC. Therefore, *V. cholerae* does not reduce As^V or expel it out of the cell. From an evolutionary perspective, this could be a vestige of a functional ArsCB system that, after losing the ArsB component, has inactivated ArsC to prevent the generation of highly toxic As^{III}. It remains to be investigated whether *V. cholerae* ArsC has undergone functional diversification to fulfill another role in the cell, perhaps related to its neighboring hypothetical genes (*vc2164-vc2167*).

Our screening identified a four-gene operon (*var operon*) conferring resistance to *V. cholerae* that encoded the ArsR repressor, the As^V permease ArsJ, a putative GAPDH (VarG) and a phosphatase (VarH). Interestingly, while most ArsR derepress transcription by sensing As^{III} (50), the lack of As^{III} production in *V. cholerae* suggests that ArsR probably senses As^V in this bacterium. Although more research is required to determine ArsR responses to As^V and As^{III} in *V. cholerae*,

comparative sequence analysis between *V. cholerae* ArsR and that of As[III]

producing bacteria have revealed variations in conserved residues, including the

E→K and C→A substitutions in the As[III] binding motif (50) (Fig. S7A).

Although GAPDH enzymes are mostly known for their role in glycolysis, they

have been functionally linked to diverse processes (51–53). A previous study

reported that the GAPDH homolog to *V. cholerae's* VarG detoxified As[V] by

generating 1As3PG from G3P. However, such conclusions were drawn from *in vitro*

experiments that used commercial glycolytic GAPDH from rabbits instead of the

specific As[V]-inducible GAPDH, and monitored As[V] transport instead of direct

1As3PG detection (17). Our results, combining *in vitro* GAPDH activity assays and

HRMS analyses from intracellular samples provided direct evidence that VarG is a

GAPDH that preferentially uses As[V] over P_i to synthesize 1As3PG. VarG role in *V.*

cholerae's As[V] resistance adds to this growing list of GAPDH moonlighting

functions.

Although the same study suggested that As[V] is exported through ArsJ as

1As3PG (17), the more severe growth defect of *V. cholerae's arsJ* compared to

varG mutant suggests that 1As3PG does not play a major role in the export of As[V]

in this bacterium. Accordingly, the *arsJ* mutant accumulates intracellular As[V] ca. 6

times more than the *varG* mutant. Even if we cannot wholly discount the transport

of 1As3PG through ArsJ (as this compound can nonetheless be produced in the

absence of VarG, likely by the constitutive glycolytic Gap activity), the fact that we

have not detected extracellular 1As3PG or increased 3PG in cultures

supplemented with As^V suggest that ArsJ expels free As^V.

During glycolysis, the GAPDH enzyme catalyzes the simultaneous

phosphorylation and oxidation of G3P to form 1,3 bi-phosphoglycerate (1,3-BPG),

which is further converted into 3PG by phosphoglycerate kinase coupled to the

transfer of the high-energy phosphate to ADP, making ATP. We reasoned that while

the metabolic complexing of As^V into 1As3PG might decrease its toxicity by

reducing As^V interaction with *V. cholerae*'s proteome (Fig4D), the formation of

1As3PG could be a dead-end product that would affect a major energy-producing

pathway. In line with this reasoning, our results showed that VarH presents low

phosphatase activity compared to canonical phosphatases. As low catalytic rate

has been previously associated with enzymatic promiscuity (54), we hypothesized

that the putative phosphatase VarH would function as an "arsenatase" that

dissociates the 1As3PG complex to clear the glycolytic pathway and free As^V for its

export through ArsJ. Indeed, inactivation of VarH results in increased intracellular

levels of 1As3PG, which correlates with the accumulation of arsenate in the cell,

thereby supporting the idea that ArsJ primarily exports free As^V in *V. cholerae*.

Although synteny analyses suggest coevolution between ArsJ and VarG, the

presence of VarH is rather exclusive to Vibrionaceae (Fig. S7B). It remains to be

investigated whether the amino acid differences in ArsJ between $VarG^+$ and $VarG^-$

species explain the choice of transporting free or complexed As^V.

It has been previously reported that arsenic induces the production of ROS in

the cell (55–57). Our results show that in the presence of As^V, *V. cholerae* induces

oxidative stress responses: i.e., elevated levels of the hydrogen peroxide-tolerance

determinant PrxA (58) and multiple proteins involved in the synthesis of cysteine.

As cysteine-containing proteins are targets of ROS (59), and cysteine is also the

building block of glutathione, stimulating its synthesis might help to alleviate

oxidative damage provoked by As^V. Additionally, we found that As^V directly or

indirectly (via ROS) alters the stability of proteins implicated in energy-generating

pathways such as TCA, fermentation, and respiration. Indeed, As^V accumulation in

var mutant dissipated cellular membrane potential (a proxy of the cell's energy

status). The correlation between As^V accumulation with ROS and membrane

depolarization exhibited by the *var* mutants suggests that oxidative stress and

energy depletion might account for their growth impairment in As^V.

V. cholerae is a pathogenic bacteria endemic to Bangladesh and India, regions

where the concentrations of As^V in water and soil are the highest worldwide (60).

Hence, *V. cholerae's* superior resistance to As^V could have evolved as an adaptive

strategy to thrive in As^V-rich environments while endowing the cholerae pathogen

with a fitness advantage over free-living and host-associated neighboring bacteria.

Interestingly, despite the phylogenetic distance, heterologous expression of the *var*

operon enhanced As^V resistance in all the pathogens assayed, demonstrating that

this system has a high degree of versatility to work in combination with other arsenic

detoxification systems.

Collectively, this work proposes a novel mechanism for As^V detoxification in *V.*

cholerae that is entirely independent of As^V-reduction and As^{III}-extrusion. Our

results support a model in which the formation of a transient As^V-containing

glycolytic intermediate cushions the impact of free As^V in the cell. While a metabolic

sink might temporarily aid cells in enduring As^V toxicity, this strategy may exert a

high cost in the long term as glycolysis is a major pathway in cell bioenergetics. We

propose "dearsenylation" of 1As3PG by VarH provides a double benefit by

decongesting any jam that may occur in glycolysis and freeing As^V for extrusion to

the extracellular media by ArsJ.

The essentiality of the *V. cholerae's var* operon to survive in environments with

As^V (e.g., such as the host) suggests that this machinery may be considered a novel

class of bacterial targets suitable for therapeutic intervention.

ACKNOWLEDGMENTS.

This work was supported by the Knut and Alice Wallenberg Foundation (KAW), The Laboratory of Molecular Infection Medicine Sweden (MIMS), the Swedish Research Council and the Kempe Foundation. V.C.T and D.D.S. were supported by the National Institutes of Health/ National Institute of Allergy and Infectious Diseases (NIH/ NIAID) grant P01-AI83211 (Metabolomics Core) and R01-AI125588. The metabolomics analyses were performed at the University of Nebraska Medical Center Mass Spectrometry and Proteomics Core Facility administrated through the Office of the Vice-Chancellor for Research and supported by state funds from the Nebraska Research Initiative (NRI). We thank Erik Björk and Laurent Ouerdane for their help with the ICPMS analyses and for insightful discussions, J. J. Mekalanos for the *V. cholerae* C6706 Tn-mutant library and Inigo Ruiz and David Arranz for their technical support

COMPETING INTEREST.

The authors declare that there are not competing interests in relation with the work described.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Strains used in this work are listed in Supplementary Table S2. *V. cholerae* strains used in this study are derivatives of the El Tor clinical isolate C6706. In addition to

V. cholerae, in this study also were used the strains *Escherichia coli* K12 MG1655

and the *arsC*::Tn mutant strain from the Keio collection (21), *Enterohemorragic*

Escherichia coli EHEC O157:H7, *Salmonella enterica serovar Typhimurium LT2,*

Citrobacter rodentium DBS100, *Shigella flexneri* M90T, *and Yersinia*

pseudotuberculosis YPIII. Strains were grown aerobically at 37°C in 15 ml tubes

containing 3 ml of complete LB medium (10 g tryptone, 10 g NaCl, and 5 g yeast

extract/L). To grow *Shigella flexneri,* TSB medium was used (Tryptone (Pancreatic

Digest of Casein), 17 g Soytone (Peptic Digest of Soybean), 3 g Glucose, 2.5 g

Sodium Chloride 5 g Dipotassium Phosphate 2.5). When used, casamino acids

were supplemented at 1%. To study resistance to As^V, 3×10^6 overnight grown cells

were inoculated into 200 µl of LB medium with or without As^V at indicated

concentrations in 96-wells plates and OD values were measured every 10 minutes

using a BioTek Epoch2 plate reader. Where appropriate, antibiotics were added to

V. cholerae and *E. coli* cultures at the following concentrations: 200 (streptomycin,

Sm), 100 (carbenicillin, Cb), 5 (Chloramphenicol, Cm) and 50µg/mL (kanamycin,

Km).

Construction of plasmids to create *V. cholerae* mutants, overexpression

constructs and transcriptional fusions.

V. cholerae mutants were created by allelic exchange with the suicide plasmid

pCVD442 [48]. ca. 1kbp upstream and downstream DNA fragments flanking each

coding regions were PCR-amplified with the primers listed in Table S3. Upstream

and downstream DNA fragments were spliced together by overlapping PCR. The

resulting ca. 2 Kb fragments were digested as indicated in table S3 and cloned into

pCVD442. For allelic exchange of *V. cholerae arsC* by *E. coli arsC,* similar protocol

was followed, but in this case the *V. cholerae* upstream and downstream DNA

regions of *V. cholerae arsC* were spliced together with the *E. coli arsC* gene by

overlapping PCR (see tables S2 and S3). Constructs were transformed into *E. coli*

DH5α λpir for amplification. They were confirmed by sequencing, transformed into

the *E. coli* donor strain SM10 λpir and conjugated for 6 hours at 37°C with *V.*

cholerae C6706 by mixing equal volumes (1ml) of exponential phase cultures, and

spot-plating. Single crossover *V. cholerae* were selected on LB plates with Sm and

Cb. Re-streaked single colonies were then plated on salt-free LB agar containing

10% (w/v) sucrose and Sm. Colonies were streaked on carbenicillin plates to

confirm loss of pCVD442 and then checked by PCR for successful deletion

mutants.

Overexpression of *V. cholerae* and *E. coli* genes was carried out by using the

arabinose inducible expression vectors pBAD33 and pBAD18, and the IPTG-

inducible expression vector pHL100 (22) (see table S2 for specific constructs use

for each enteropathogen). To construct each overexpressing plasmid, genes open

reading frames including ribosome binding sites were PCR amplified (see primers

used in table S3), double digested with indicated restriction enzymes and cloned

into each plasmid. Constructs were transformed into the indicated strain by

electroporation.

For construction of the transcriptional reporter fusions, gene promoter regions of

target genes were amplified using the primers pair indicated in table S3. The PCR

product was double digested with HindIII and EcoRI and then ligated into HindIII-

EcoRI-digested pCB192N (23). Cloned plasmids were then transformed into *V.*

cholerae by electroporation.

Purification of recombinant VarG, VC2000, VC1041 and VarH.

V. cholerae VarG, VC1041 and VarH proteins were overexpressed using the *E. coli*

strain BL21 as 6×His-tagged enzymes from a pET28b vector. Overnight cultures

were diluted into 250 mL LB broth and grown until OD_{600}=0.5. Flasks were

supplemented with 1 mM IPTG and shaking at 37⁰ C for 2 h. Cells were pelleted,

and resuspended in equilibration buffer (Tris 50 mM, pH 7.5, 50 mM NaCl, with

protease inhibitor cocktail (Roche)) and disrupted by passaging once through a

French press. Lysates were then centrifuged for 1 h (25,000 rpm, Beckman Coulter

Avanti J26-XP centrifuge, JL-25.50 rotor) at 4°C. Nickel-NTA resin (0.5 ml

resuspended in equilibration buffer) was then added to the supernatant, followed

by incubation at 4°C in a rotating wheel. The lysate was separated from the resin

by centrifugation for 1 minute at 3,220 g and the resin washed (5×10 ml) with

washing buffer (equilibration buffer adjusted to 1 M NaCl) and eluted with 2ml of

washing buffer containing 500mM imidazole. Fractions were subjected to SDS-

PAGE and Coomassie Brilliant Blue staining for purity assessment. Protein

concentration was quantified with a Bradford assay.

Proteomic and thermal proteome profiling (TPP).

We performed thermal proteome profiling to determine protein abundance and

thermal stability changes in WT, and *var* mutant strains upon exposure to As^V

similarly to what was previously described (24, 25). Briefly, WT and mutant cells

grown to an OD578 of 0.5 were incubated with 1 mM arsenate (or a similar volume

of water) for 10 min. Cells were then washed with 10 ml PBS (for cells treated with

arsenate, 1 mM arsenate were included in the wash step) and aliquoted to a PCR

plate. Each aliquot was then exposed to a different temperature in the range 37°C-

66.3°C for 3 min. Following a 3 min incubation at room temperature, cells were

lysed with lysis buffer (final concentration: 50 µg ml^{-1} lysozyme, 0.8% NP-40,

1 × protease inhibitor (Roche), 250 U ml^{-1} benzonase, and 1 mM MgCl$_2$ in PBS) for

20 min followed by three freeze–thaw cycles. Lysates were then prepared for

analysis by mass spectrometry by digesting proteins using a modified sp3 protocol

(26), labeling peptides with TMTpro (Thermo Fisher Scientific) and pooling samples

from the same temperature together. These samples were then fractionated to six

fractions with high pH fractionation and injected on an Orbitrap Q-Exactive Plus

(Thermo Fisher Scientific) coupled to liquid chromatography (details on the run

conditions and instrument parameters as in Mateus et al. (2020) (24, 25).

Mass spectrometry raw data was searched against the Vibrio cholerae FASTA file

(UP000000584 downloaded from UniProt) using the Mascot 2.4 (Matrix Science)

search engine and isobarquant (27). Protein abundance and thermal stability

changes were determined using limma (28) using the same algorithm as in Mateus

et al. (2020)(24).

Phosphorylation state of VarG.

To identify VC1069 GAPDH phosphorylation state, *V. cholerae* wild type and Δ*varH*

strains carrying the overexpression vector pHL100 containing the *varG*-HIS tagged

gene cloned were grown in the presence of As^V and with IPTG. Cells were pelleted,

disrupted, load and run in SDS-PAGE gel. VarG-HIS bands were excised and

diluted with NuPAGE LDS sample buffer (Invitrogen, Carlsbad, CA) mixed with10

mM dithiotreitol (DTT) and incubated at 56 °C for 20 min followed by addition of

iodoacetamide (IAA) to a final concentration of 20 mM. The sample was loaded on

4-12% NuPAGE (Invitrogen). The electrophoresis was run in MOPS buffer at 180

V for 1 h. The gel was stained with Coomassie Brilliant Blue. Bands of interest were

cut from the SDS-PAGE. The bands were digested with trypsin, followed by

identification using liquid chromatography–tandem mass spectrometry (LC–

MS/MS). MS/MS spectra were searched with Proteome Discoverer 2.3 (Thermo

Fisher Scientific) against the two sequences obtained for phosphate groups

(UniProtKB). The precursor tolerance and fragment tolerance were set to 10 ppm

and 0.05 Da, respectively. Trypsin was selected as enzyme, methionine oxidation,

phosphorylation of serine, tyrosine, and tryptophan, and deamidation of asparagine

and glutamine were treated as dynamic modification and carbamidomethylation of

cysteine as a fixed modification (see supplementary methods for further

information).

Transposon-insertion sequencing (TIS) analysis.

TIS was carried out essentially as previously described (29) but using selection

plates containing LB medium with As^V 1 mM. In brief, ~600,000 transposon mutants

were generated by conjugation of *V. cholerae* C6706 with SM10λ*pir E. coli* carrying

the Himar1 suicide transposon vector pSC189 (30). They were collected and their

genomic DNA was pooled and analyzed on an Illumina MiSeq bench-top sequencer

(Illumina, San Diego, CA). Insertion sites (which included 35% of TA sites) were

identified as previously described (29), and significance was determined using Con-

Artist simulation-based normalization as described (31). Results were visualized

using Artemis (32).

As^V resistance screen of arrayed transposon mutant library.

A non-redundant transposon insertion library from the *Vibrio cholerae* strain C6706

(33) consisting of 3,156 insertion mutants was used to determine genetic

determinants providing resistance to As^V in *V. cholerae*. *V. cholerae* WT and the

library insertion mutants were first grown aerobically in 200 µl of LB in 96-well plates

at 37ºC for 24 hours. These cultures were then used as inoculum into 96-well plates

containing 200 µl complete LB medium, in the presence of 1 mM of As^V. The 96-

well plates were incubated under agitation at 37ºC for 24 hours and then resistance

to As^V was evaluated by measuring optical density (OD$_{600}$) using a

BioTek Eon plate reader.

Protein homology studies of As^V resistance determinants in enteric

pathogens.

DNA sequences of the *Vibrio cholerae arsR, varG, varH and arsJ var* genes and

complete genomes sequences of the enteric pathogens: *Salmonella typhimurium*

(PRJNA231221), *Citrobacter rodentium* (PRJNA261533), *Shigella flexeneri*

(PRJNA553848), EHEC (PRJNA218110) and *Yersenia pseudotuberculosis*

(PRJEB6403) were downloaded from NCBI database. Complete genomes of the

pathogens were translated by Prodigal ver. 2.6.2 and homologous sequences to

the *V. cholerae* Var proteins were queried into the enteric pathogens by BlastP ver.

2.9.0. Sequence homology data were represented in a heatmap diagram with

Spearman correlation done by R ver. 3.6.1.

β-Galactosidase activity determination.

β-Galactosidase activity was measured through ONPG cleavage by the product of

the *lacZ* reporter gene, and specific activity was calculated in Miller units (34). In

brief, cultures of *V. cholerae* carrying the pCB192N plasmid harboring promoter

gene insertions (see table S2) were grown at 37ºC overnight in 15 ml falcons tube

containing 3 ml of LB complete medium. Overnight cultures were diluted 1:200 and

supplemented with 1 mM of As^V and incubated at 37ºC until they reached mid-log

phase (OD_{600} 0.4-0.7). A 100 µl aliquot of three different subcultures were collected,

cells were permeabilized and assayed in triplicate for each strain as previously

described (34).

Determination of intracellular and extracellular As^V and As^{III} in *V. cholerae*.

To determinate concentrations of As^V and As^{III} in supernatants of *V. cholerae* and

E. coli, bacterial cells were grown in the presence of As^V for 5 hours and then

pelleted. Supernatants of each strain were speciated by high-performance liquid

chromatography (HPCL) (Agilent Infinity II 1290) using a C18 column reverse-

phase (130 Å, 1.7 µm, 2.1 mm by 150 mm; Waters, USA). Elution conditions used

were flow rate 1 ml min^{-1}; temperature 25 °C; isocratic elution in 5 mM

tetrabutylammonium hydroxide, 5 % methanol (v/v) and 3 mM malonic acid.

Identification and quantification of As^V and As^{III} was performed by inductively

coupled plasma mass spectroscopy (ICP-MS) (Agilent 8900 ICP-MS Triple Quad)

by comparison to standards of known concentration and peak integration.

To determine intracellular concentrations of As^V and As^{III} in *V. cholerae,* the

pelleted cells were washed in complete LB medium to remove traces of arsenic,

then resuspended in 500 µl of H_2O. Glass beads were adding to the resuspension

and cells were lysed by bead-beating. The resulting extract was centrifuged, and

As^V and As^{III} from those supernatants were analyzed by ICP-MS as previously

described. Samples were analyzed in triplicate.

LC-MS/MS analysis using QTRAP6500+

To measure 3-phosphoglycerate (3PG), *V. cholerae* cells grown in the presence of

5 mM of As^V were pelleted and used for LC-MS/MS analysis. Bacterial pellets were

resuspended in 1 ml ice-cold 60% ethanol containing 2 μM ribitol as the internal

control. The cells were mechanically disrupted using a bead homogenizer set to

oscillate for 3 cycles of 30 sec each at 6800 rpm with a 10 sec pause between each

cycle. Cell debris was separated at 13,000 rpm. An aliquot (200μl) of the

supernatant containing intracellular metabolites was vacuum dried and

resuspended in an equal volume of 7.5 mM ammonium bicarbonate solution before

LC-MS/MS analysis.

A triple-quadrupole-ion trap hybrid mass spectrometer, QTRAP® 6500+ (Sciex,

USA) connected with a Waters ultra-performance liquid chromatography I-class

(UPLC) system was used for metabolite analysis. The chromatographic separation

was performed on an XSELECT HSS XP column (150 mm × 2.1 mm ID; 2.5 μm

particle size, Waters, USA) using a binary solvent system at a flow rate of

0.1 ml/min. Mobile phase A was composed of 7.5 mM ammonium bicarbonate in

LC-MS grade water and mobile phase B was 100% LC-MS grade methanol. The

column was maintained at 40°C, and the autosampler temperature was set to 7°C.

The A/B solvent ratio was maintained at 100/0 for 2 minutes, followed by a gradual

increase of solvent B to 95% for 2 minutes. The solvent B was maintained at 95%

over the next 5 minutes. The gradient was reduced to 100% solvent A within 0.5 min

and column was equilibrated for 5.5 minutes before the next run. The needle was

washed with 1.2 ml of a strong wash solution containing 100% LC-MS-grade

acetonitrile followed by 1 ml of a weak wash solution comprised of 10% aqueous

methanol before each injection. The injection volume was 5μl. The QTRAP® 6500+

(SCIEX) was operated in negative as well as positive ion mode for targeted

quantitation in Multiple Reaction Monitoring (MRM). MRM parameters for each

analyte are listed in Table 1. The electrospray ionization (ESI) parameters used are

as follows: electrospray ion voltage of -4500V in negative and 5500V in positive

mode, source temperature of 400°C, curtain gas of 35, and gas 1 and 2 of 40 psi

each, respectively. The compound-specific parameters such as declustering

potential (DP) and collision energy (CE) were optimized for each compound using

manual tuning. These values are listed in Table 1.

Table 1: MRM transitions, declustering potentials and collision energies of

metabolites

Analyte	Q1 (m/z)	Q3 (m/z)	DP (V)	CE (V)
3-phospho glycerate	187.2	98.9	+60	19.2
Ribitol	150.8	88.9	-60	-14.2

Detection of 1As3PG.

1As3PG was identified using a Thermo Orbitrap Exploris 480 high resolution mass

spectrometer. The chromatographic method used for this analysis was the same

as that described for the LC-MS/MS performed using the QTRAP6500+. The MS

data were acquired in both positive and negative polarity within mass range of 140-

400 Da. The precursor ion resolution was maintained at 240,000, whereas the

product ion resolution at 120,000. Deprotonated $[M-H]^-$ ion of 1As3PG was detected

at m/z 304.8696 Da compared to the actual mass of 304.8674 Da, thus $\Delta m = 7.149$

ppm. The fragmentation pattern of 1As3PG could not be further resolved under

these conditions.

Proteins structural and topological analysis.

Protein alignments were performed by using Vector NTI. Prediction of VarH

secondary structure was performed by using Phyre2

(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index). Structural prediction

of the VarG and VarH proteins was performed with AlphaFold2 on the CoLabFold

publicly accessible interface (35)

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF

old2.ipynb). Sequences were modeled as monomers using mmseqs2 for multiple

sequence alignment. Studies on proteins topological similarity was carried out by

using TM-score (https://zhanggroup.org/TM-score/).

Competition assays.

Aerobic overnight cultures of *V. cholerae* C6706 wild type lacZ⁻ and

Enterohemorragic Escherichia coli EHEC O157:H7 *lacZ⁺*, *Salmonella enterica*

serovar Typhimurium LT2 lacZ⁺, *Citrobacter rodentium* DBS100 *lacZ⁺*, *Shigella*

flexneri M90T *lacZ⁺, and Yersinia pseudotuberculosis* YPIII *lacZ⁺,* were collected

and 1 x 10^7 cells co-inoculated at 1:1 ratio in 15 ml tubes containing 3 ml of complete

LB medium without or with 10 mM As^V. After 8 hours of incubation at 37°C, an

aliquot was diluted and plated on LB agar plates containing X-Gal to enumerate

wild type and competing strains. Competitive indices (CIs) were determined by

dividing the ratio mutant or non-wild type strain to C6706 wild type colonies by the

ratio in the inoculum. To calculate CIs among *V. cholerae* strain C6706 against a

different *V. cholerae* strain against enteric pathogens (Fig. 1B), CIs obtained after

the As^V challenge were normalized respect CIs obtained after competition in the

absence of As^V.

***in vitro* GAPDH and phosphatase enzymatic assays.**

The GAPDH activity assay was performed in accordance with manufacturer's

instructions (MAK27; Sigma-Aldrich) but using P$_i$ or As^V as the GAPDH substrate.

10 µg of purified *V. cholerae* VarG and VC2000 enzymes were used in the assay.

This assay quantifies GAPDH activity by measuring enzymes capacity to reduce

NAD$^+$ to NADH during the conversion of glyceraldehyde 3-phosphate to 1,3-

biphosphoglycerate. NADH produced in this reaction results in a colorimetric

(OD$_{450}$) product proportional to the GAPDH activity of the enzyme. GAPDH activity

is reported as nmole/min/mL = milliunit/mL. One unit of GAPDH is the amount of

enzyme that will generate 1.0 mmole of NADH per minute at pH 7.2 at 37 °C.

Phosphatase activity was quantified using Cayman's Phosphatase Colorimetric

assay kit and 5 µg of purified *V. cholerae* VarH, C113G R119G VarH, VC1041

(positive control) and BsrV (negative control) enzymes. The assay uses *p*-

nitrophenylphosphate (pNPP) as substrate for phosphatase enzymes.

Phosphatase enzymes dephosphorylate pNPP, which deprotonates its phenolic

OH-group by increasing the pH of the reaction, deprotonated pNP yields an intense

yellow color detected at OD 405 nm. Enzymatic activities were calculated as µmol

of phosphate released per min/ml.

Flow cytometry analysis

V. cholerae WT and indicated *var* mutant strains were grown for 5 hours in the

presence of 1 mM As^V, then Aliquots (1 ml) from each culture were pelleted, washed

once, and resuspended in PBS (100 µl) to measure membrane potential and ROS.

To measure *membrane potential* in *V. cholerae* cells, we used the BacLight

bacterial membrane potential kit (Invitrogen). *V. cholerae* samples resuspended in

100 µl PBS were incubated for 15 min at 20 °C in the dark with the dye 30 µM 3,3'-

diethyloxacarbocyanine iodide (DiOC$_2$). Next, cells were washed twice with PBS

and resuspended in 2 ml of PBS, then fluorescence emitted from treated cells was

measured with a Bio-Rad S3e cell sorter at an excitation wavelength of 488 nm and

an emission wavelength of 525/30 nm (for green) or 655 nm (for red).

ROS measurements were performed using the CellROX™ Green Reagent, for

oxidative stress detection (ThermoFisher). Washed *V. cholerae* cells were

supplemented with CellROX® Reagent at a final concentration of 5 µM and

incubate them for 30 minutes at 37°C. Next, cells were washed twice with PBS and

resuspended in 2 ml of PBS, then fluorescence emitted was measured with a Bio-

Rad S3e cell sorter at an excitation wavelength of 488 nm and an emission

wavelength of 525 nm.

Statistics and reproducibility

Statistical significance was assessed with the Student's *t*-test where indicated. A p-

value of less than 0.05 was considered statistically significant. Assays were

performed with three biological replicates unless otherwise indicated.

REFERENCES

1.   Kalia K, Joshi DN. 2009. Detoxification of Arsenic. Handb Toxicol Chem

Warf Agents 1083–1100.

2.   Singh R, Singh S, Parihar P, Singh VP, Prasad SM. 2015. Arsenic

contamination, consequences and remediation techniques: A review.

Ecotoxicol Environ Saf 112:247–270.

3.   Pal C, Asiani K, Arya S, Rensing C, Stekel DJ, Larsson DGJ, Hobman JL.

2017. Metal Resistance and Its Association With Antibiotic Resistance.

Microbiol Met Ions, 1st ed. 70:261–313.

4.   Hobman JL, Crossman LC. 2015. Bacterial antimicrobial metal ion

resistance. J Med Microbiol 64:471–497.

5.   Willsky GR, Malamy MH. 1980. Characterization of two genetically

separable inorganic phosphate transport systems in Escherichia coli. J

Bacteriol 144:356–365.

6.   Willsky GR, Malamy MH. 1980. Effect of arsenate on inorganic phosphate

transport in Escherichia coli. J Bacteriol 144:366–374.

7.   Dixon HBF. 1996. The Biochemical Action of Arsonic Acids Especially As

Phosphate Analogues. Adv Inorg Chem 44:191–227.

8.   Rosen BP. 2002. Biochemistry of arsenic detoxification. FEBS Lett 529:86–

92.

9.   Yang HC, Fu HL, Lin YF, Rosen BP. 2012. Pathways of Arsenic Uptake and

Efflux, p. 325–358. *In* Current Topics in Membranes. NIH Public Access.

10. Scott N, Hatlelid KM, MacKenzie NE, Carter DE. 1993. Reactions of Arsenic

(III) and Arsenic (V) Species with Glutathione. Chem Res Toxicol 6:102–

106.

11. Przygoda G, Feldmann J, Cullen WR. 2001. The arsenic eaters of Styria: A

different picture of people who were chronically exposed to arsenic. Appl

Organomet Chem 15:457–462.

12. Liu J, Lu Y, Wu Q, Goyer R a, Waalkes MP. 2008. Mineral Arsenicals in

Traditional Medicines : Perspect Pharmacol 326:363–368.

13. Shahpiri A, Mohammadzadeh A. 2018. Bioaccumulation of Arsenic by

Engineered Escherichia coli Cells Expressing Rice Metallothionein

Isoforms. Curr Microbiol 75:1537–1542.

14. Thomas DJ, Rosen BP. 2013. Arsenic Methyltransferases, p. 138–143. *In*

Kretsinger, RH, Uversky, VN, Permyakov, EA (eds.), Encyclopedia of

Metalloproteins. Springer New York, New York, NY.

15. Saltikov CW, Newman DK. 2003. Genetic identification of a respiratory

arsenate reductase. Proc Natl Acad Sci U S A 100:10983–10988.

16. Oden KL, Gladysheva TB, Rosen BP. 1994. Arsenate reduction mediated

by the plasmid-encoded ArsC protein is coupled to glutathione. Mol

Microbiol 12:301–306.

17. Chen J, Yoshinaga M, Garbinski LD, Rosen BP. 2016. Synergistic

interaction of glyceraldehydes-3-phosphate dehydrogenase and ArsJ, a

novel organoarsenical efflux permease, confers arsenate resistance. Mol

Microbiol 100:945–953.

18. Wu S, Wang L, Gan R, Tong T, Bian H, Li Z, Du S, Deng Z, Chen S. 2018.

Signature arsenic detoxification pathways in Halomonas sp. Strain GFAJ-1.

MBio 9:1–9.

19. Fekih I Ben, Zhang C, Li YP, Zhao Y, Alwathnani HA, Saquib Q, Rensing C,

Cervantes C. 2018. Distribution of arsenic resistance genes in prokaryotes.

Front Microbiol 9:1–11.

20. Németi B, Anderson ME, Gregus Z. 2012. Glutathione synthetase promotes

the reduction of arsenate via arsenolysis of glutathione. Biochimie 94:1327–

1333.

21. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA,

Tomita M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12

in-frame, single-gene knockout mutants: The Keio collection. Mol Syst Biol

2:1–11.

22. Guzman L, Belin D, Carson MJ. 1995. Tight Regulation, Modulation, and

High-Level Expression by Vectors Containing the Arabinose PBAD

Promoter. J Bacteriol 177:4121–4130.

23.   Schneider K, Beck CF. 1986. Promoter-probe vectors for the analysis of

774   divergently arranged promoters. Gene 42:37–48.

775   24.   Mateus A, Hevler J, Bobonis J, Kurzawa N, Shah M, Mitosch K, Goemans C

776   V., Helm D, Stein F, Typas A, Savitski MM. 2020. The functional proteome

777   landscape of Escherichia coli. Nature 588:473–478.

778   25.   Mateus A, Kurzawa N, Becher I, Sridharan S, Helm D, Stein F, Typas A,

779   Savitski MM. 2020. Thermal proteome profiling for interrogating protein

780   interactions. Mol Syst Biol 16:1–11.

781   26.   Hughes CS, Moggridge S, Müller T, Sorensen PH, Morin GB, Krijgsveld J.

782   2019. Single-pot, solid-phase-enhanced sample preparation for proteomics

783   experiments. Nat Protoc 14:68–85.

784   27.   Franken H, Mathieson T, Childs D, Sweetman GMA, Werner T, Tögel I,

785   Doce C, Gade S, Bantscheff M, Drewes G, Reinhard FBM, Huber W,

786   Savitski MM. 2015. Thermal proteome profiling for unbiased identification of

787   direct and indirect drug targets using multiplexed quantitative mass

788   spectrometry. Nat Protoc 10:1567–1593.

789   28.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015.

790   Limma powers differential expression analyses for RNA-sequencing and

791   microarray studies. Nucleic Acids Res 43:e47.

29.   Chao MC, Pritchard JR, Zhang YJ, Rubin EJ, Livny J, Davis BM, Waldor

MK. 2013. High-resolution definition of the Vibrio cholerae essential gene

set with hidden Markov model-based analyses of transposon-insertion

sequencing data. Nucleic Acids Res 41:9033–9048.

30.   Chiang SL, Rubin EJ. 2002. Construction of a mariner-based transposon for

epitope-tagging and genomic targeting. Gene 296:179–185.

31.   Pritchard JR, Chao MC, Abel S, Davis BM, Baranowski C, Zhang YJ, Rubin

EJ, Waldor MK. 2014. ARTIST: High-Resolution Genome-Wide Assessment

of Fitness Using Transposon-Insertion Sequencing. PLoS Genet

10:e1004782.

32.   Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012. Artemis:

An integrated platform for visualization and analysis of high-throughput

sequence-based experimental data. Bioinformatics 28:464–469.

33.   Cameron DE, Urbach JM, Mekalanos JJ. 2008. A Defined Transposon

Mutant Library and Its Use in Identifying Motility Genes in Vibrio cholerae.

Proc Natl Acad Sci U S A 105:8736–8741.

34.   Miller JH. 1972. Experiments in Molecular Genetics. New York, NY: Cold

Spring Harbor Laboratory.

35.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,

Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C,

Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R,

Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger

M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior

AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein

structure prediction with AlphaFold. Nature 596:583–589.

36.   Demel S, Shi JIN, Martin P, Rosen BP, Edwards BFP. 2004. Arginine 60 in

the ArsC arsenate reductase of E . coli plasmid R773 determines the

chemical nature of the bound As (III) product. Protein Sci 2330–2340.

37.   Martin P, Demel S, Shi J, Gladysheva T, Gatti DL, Rosen BP, Edwards

BFP. 2001. Insights into the Structure, Solvation, and Mechanism of ArsC

Arsenate Reductase, a Novel Arsenic Detoxification Enzyme. Structure

9:1071–1081.

38.   Jenkins JL, Tanner JJ. 2006. High-resolution structure of human D-

glyceraldehyde-3-phosphate dehydrogenase. Acta Crystallogr Sect D Biol

Crystallogr 62:290–301.

39.   Velick SF. 1955. Glyceraldehyde-3-phosphate dehydrogenase from muscle.

Methods Enzymol 1:401–406.

40.   Byers LD, She HS, Alayoff A, Gregus Z, Németi B. 2005. Interaction of

Phosphate Analogues with Glyceraldehyde-3-phosphate Dehydrogenase.

Biochemistry 85:859–869.

41.   Byers LD, She HS, Alayoff A. 1979. Interaction of Phosphate Analogues

833  with Glyceraldehyde-3-phosphate Dehydrogenase. Biochemistry 18:2471–

834  2480.

835 42. Aoyama K, Nakaki T. 2015. Glutathione in cellular Redox homeostasis:

836  Association with the excitatory amino acid carrier 1 (EAAC1). Molecules

837  20:8742–8758.

838 43. Modulates O, Activity O. 2017. OxyR2 Modulates OxyR1 Activity and Vibrio

839  cholerae Oxidative Stress Response.

840 44. Heidelberg JF et al. 2000. DNA sequence of both chromosomes of the

841  cholera pathogen Vibrio cholerae. Nature 406:477–483.

842 45. Jin Q. 2002. Genome sequence of Shigella flexneri 2a: insights into

843  pathogenicity through comparison with genomes of Escherichia coli K12

844  and O157. Nucleic Acids Res 30:4432–4441.

845 46. Batzilla J, Höper D, Antonenka U, Heesemann J, Rakin A. 2011. Complete

846  genome sequence of Yersinia enterocolitica subsp. palearctica Serogroup

847  O:3. J Bacteriol 193:2067–2065.

848 47. Pearson BM, Gaskin DJH, Segers RPAM, Wells JM, Nuijten PJM, Van Vliet

849  AHM. 2007. The complete genome sequence of Campylobacter jejuni strain

850  81116 (NCTC11828). J Bacteriol 189:8402–8403.

851 48. Tolen TN, Xie Y, Hernandez AC, Everett GFK. 2016. Complete genome

sequence of Salmonella enterica serovar TyphimuriumLT2. Nature

413:852–856.

49. Popov G, Fiebig-Comyn A, Shideler S, Coombes BK, Savchenko A. 2019.

Complete Genome Sequence of Citrobacter rodentium Strain DBS100 .

Microbiol Resour Announc 8:1–2.

50. Shi W, Wu J, Rosen BP. 1994. Identification of a putative metal binding site

in a new family of metalloregulatory proteins. J Biol Chem 269:19826–

19829.

51. Sirover MA. 2011. Biochimica et Biophysica Acta On the functional diversity

of glyceraldehyde-3-phosphate dehydrogenase : Biochemical mechanisms

and regulatory control. BBA - Gen Subj 1810:741–751.

52. Ferreira E, Giménez R, Alexandra M, Aguilera L, Aguilar J, Badia J,

Baldomà L. 2015. Glyceraldehyde-3-phosphate dehydrogenase is required

for efficient repair of cytotoxic DNA lesions in Escherichia coli. Interntional J

Biochem Cell Biol 60:202–212.

53. Whitworth DE, Morgan BH. 2015. Synergism between Bacterial GAPDH

and OMVs: Disparate Mechanisms but Co-operative Action. Front Microbiol

6:1–6.

54. Leveson-Gower RB, Mayer C, Roelfes G. 2019. The importance of catalytic

promiscuity for enzyme design and evolution. Nat Rev Chem 3:687–705.

55. Shi H, Shi X, Liu KJ. 2004. Oxidative mechanism of arsenic toxicity and

carcinogenesis 67–78.

56. Liu SX, Athar M, Lippai I, Waldren C, Hei TK. 2001. Induction of oxyradicals

by arsenic: Implication for mechanism of genotoxicity. Proc Natl Acad Sci U

S A 98:1643–1648.

57. Sukchawalit R, Prapagdee B, Charoenlap N, Vattanaviboon P, Mongkolsuk

S. 2005. Protection of Xanthomonas against arsenic toxicity involves the

peroxide-sensing transcription regulator OxyR. Res Microbiol 156:30–34.

58. Wang H, Chen S, Zhang J, Rothenbacher FP, Jiang T, Kan B, Zhong Z,

Zhu J. 2012. Catalases Promote Resistance of Oxidative Stress in Vibrio

cholerae. PLoS One 7.

59. Miki H, Funato Y. 2012. Regulation of intracellular signalling through

cysteine oxidation by reactive oxygen species. J Biochem 151:255–261.

60. Mondal P, Majumder CB, Mohanty B. 2006. Laboratory based approaches

for arsenic remediation from contaminated water: Recent developments. J

Hazard Mater 137:464–479.

FIGURE LEGENDS.

Fig. 1 *V. cholerae* **is the enteric pathogen with the greatest resistance to As^V.**

A Growth curves (OD$_{600}$) of the enteric pathogens *V. cholerae*, *Enterohaemorragic Escherichia coli* (EHEC), *Salmonella enterica*, *Citrobacter rodentium*, *Shigella flexneri*, and *Yersinia pseudotuberculosis*. Cultures were grown in LB medium in the absence (black) and in the presence (blue) of 10 mM arsenate (As^V). **B** *in vitro* competition among *V. cholerae* and other enteropathogenic bacteria (EPB) studied in panel A. Cells were incubated in LB medium supplemented with 10 mM As^V for 8 hours. CI values from competition in the presence of As^V were normalized with respect to CI values in the absence As^V. **C** Growth curves (OD$_{600}$) of WT (blue filled circles) and *arsC* (blue empty circles) mutant strains from *V. cholerae* and *E. coli* K12. Cultures were grown in LB medium supplemented with 10 mM As^V. As^V and As^{III} concentrations were determined by ICP-MS from supernatants of *V. cholerae* and *E. coli* WT and *arsC* mutant strains after 8 hours of incubation in 1.5 mM of As^V. Data are the mean of three biological replicates ± s.e.m.

Fig. 2 **Identification of essential genetic determinants for As^V resistance in** *V. cholerae.* Left: Experimental workflow for an arrayed transposon mutant screen to identify the genes required for As^V resistance in *V. cholerae* and the list of mutant

strains identified. Right: Experimental workflow for TIS screen in the presence and

absence of 1 mM As^V. Volcano plot depicting the ratio of read counts mapped to

individual genes in transposon libraries of *V. cholerae* plated onto media

supplemented with As^V). Red dotted lines indicate arbitrary thresholds of fold

change (FC) <0.1 and an inverse p-value >20. Genes in red color show loci also

identified in the arrayed screen (see Supplementary Table 1 for an extended

version).

Fig. 3   Phenotypic characterization of the *V. cholerae var* operon. Growth

curves (OD_{600}) of *V. cholerae* WT, single and combinatorial *var* mutant strains.

Cultures were grown in LB medium in the absence or presence of 1 and 10 mM

As^V. Data are the mean of three biological replicates ± s.e.m.

Fig. 4   Characterization of *V. cholerae* VarG. A *in vitro* GAPDH activity (OD_{450})

of purified *V. cholerae* VarG and the glycolytic Gap enzymes using P_i or As^V as

substrates. **B** β-galactosidase activity of strains in which the *lacZ* reporter gene is

cloned under the control of the *var* operon and the *gap* genes promoters. Cultures

were grown in LB medium in the absence or presence of 1 and 10 mM As^V for 5

hours. **C** Intracellular As^V concentrations of WT, *varG, varH* and *arsJ* mutant strains

grown with 1 mM As^V by ICP-MS. **D** Left: As^V detoxification model in *P. aeruginosa*

DK2 [1]**.** Right: Quantification of 1As3PG and 3PG from intracellular and

extracellular samples of *V. cholerae* cells grown with and without 1 mM As^V. Data

are the mean of three biological replicates ± s.e.m.

Fig. 5 Characterization of *V. cholerae* VarH. A Left: *in vitro* phosphatase activity of purified *V. cholerae* VarH, and catalytic mutant derivative where cysteine 113 and arginine 119 were replaced by glycine. *V. cholerae* protein-tyrosine-phosphatase (VC1041) and broad-spectrum racemase (BsrV) were used as positive and negative controls, respectively. Right: Growth curves (OD_{600}) of *V. cholerae* WT, and a VarH catalytic mutant derivative . Cultures were grown in LB medium in the presence of As^V. **B** Left: Intracellular 1As3PG concentrations from *V. cholerae* WT and $\Delta varH$ cells grown with and without 1 mM As^V by HRMS. Right: Schematic of putative mechanism for As^V detoxification in *V. cholerae*. Data are the mean of three biological replicates ± s.e.m.

Fig. 6 Oxidative stress response and membrane potential alterations of *V. cholerae* WT and the *var* mutants. A Volcano plot of LIMMA analysis output depicting read fold change ($\log_2$) of protein abundance and inverse P-value for each protein queried in the proteomic screen in response to 1 mM As^V in *V. cholerae* WT. Red dotted lines indicate arbitrary thresholds of fold change ($\log_2$) of >1 or <1 and an inverse P-value of >20. Var proteins are labelled in blue and oxidative stress related proteins (Cystein biosynthesis and PrxA are labelled as in B. **B** Schematic of the systems induced in response to As^V in *V. cholerae*. Non-coloured arrows

depict genes within an operon not detected. **C** Representative dot plots from *V.*

cholerae WT and *var* mutant cells stained with CellRox Green following growth in

the presence of 1 mM As^V for 5 hours. The black dotted gate indicates an area with

higher CellRox fluorescence (higher ROS levels). **D** Volcano plot of LIMMA analysis

output depicting read fold change (log$_2$) of protein thermal stability and inverse P

value for each protein queried in the proteomic screen in response to 1 mM As^V in

V. cholerae WT and *var* mutant strains. As^V interacting proteins related to energy-

generating pathways are labeled in blue color. PflB, VC1866 formate

acetyltransferase. AdhE, VC2033 alcohol dehydrogenase/acetaldehyde

dehydrogenase. HemN, VC0116 oxygen-independent coproporphyrinogen III

oxidase. Icd, VC1141 isocitrate dehydrogenase. FrdA, VC2656 fumarate

reductase. IspH, 4-hydroxy-3-methylbut-2-enyl diphosphate reductase. AceF,

VC2413 pyruvate dehydrogenase, E2 component. UbiE, VC0083

ubiquinone/menaquinone biosynthesis methyltransferase. **E** Representative dot

plots from *V. cholerae* WT and *var* mutant cells stained with DiOC2 following growth

in the presence of 1 mM As^V for 5 hours. The black dotted gate indicates an area

with lowered green fluorescence (lower membrane potential). Data are the mean of

three biological replicates ± s.e.m.

Fig. 7 *V. cholerae* Var enhances As^V resistance in enteropathogenic bacteria.

A Hierarchical heat map clustering of enteropathogenic bacteria with respect to *V.*

cholerae based on the percentage identity of *var* genes . **B** Growth curves (OD$_{600}$)

of the As^V-sensitive enteropathogenic bacteria clustered in panel A, carrying an

inducible empty plasmid (empty circle) or the same plasmid expressingthe *V.*

cholerae var operon (blue filled circle). Cultures were grown in LB medium in the

presence of 10 mM As^V. Data from panel B are the mean of three biological

replicates ± s.e.m.

988   **SUPPLEMENTARY INFORMATION**

989   **Transient glycolytic complexation of arsenate enhances resistance in the**

990   **enteropathogen *Vibrio cholerae***

991   Emilio Bueno*, Víctor Pinedo, Dhananjay Shinde, André Mateus, Athanasios

992   Typas, Mikhail M Savitski, Vinai Chittezham Thomas, and Felipe Cava*.

Content – Supplementary Information

995   • Supplementary Figures S1 – S7

996   • Supplementary Tables S1 - S3

997   • Supplementary references.

1000

1001

1002

1003

1004

1005

1006   **Fig. S1 Growth curves (OD$_{600}$) of *V. cholerae* WT in the presence of As^{V}.** *V.*

1007   *cholerae* WT was grown in LB medium in the presence of 10, 20 and 30 mM As^{V}.

1008   Data are the mean of three biological replicates ± s.e.m.

Fig. S2   Characterization of *V. cholerae* ArsC. A Schematic of the *arsC*-containing operons from *E. coli* K12 and *V. cholerae,* and protein sequence alignments. Black arrows depict ArsC catalytic cysteine and arginine residues implicated in As^V reduction to As^{III}. **B** Growth curves (OD_{600}) of *V. cholerae* WT, and two derivative strains where *arsC* has been replaced by *E. coli's arsC* (red, $arsC^{ec}$), and $arsC^{ec}$ strain where *arsB* is overexpressed in trans (blue, $arsC^{ec}$ + $arsB^{ec}$) (see bottom panel, and material and methods for further details). Cultures were grown in LB medium in the presence 10 mM As^V. Bottom panel: Explanatory diagrams illustrating the expected output from each construct. Data are the mean of three biological replicates ± s.e.m.

Fig. S3 Transcriptional analysis of the *var* operon. β-galactosidase activity from *V. cholerae* WT and *arsR* mutant (Δ*arsR*) strains carrying DNA regions upstream of *arsR, varG, varH* and *arsJ* genes fused to the *lacZ* reporter gene. Each DNA region fused to *lacZ* is depicted by arrows. Cultures were grown in LB medium in the presence or absence of 10 mM As^V. Data are the mean of three biological replicates ± s.e.m.

Fig. S4   *V. cholerae* VarG presents a high degree of similarity to the glycolytic GAPDH Gap. A Sequence alignments of the *V. cholerae* VarG and the glycolytic GAPDH proteins using Vector NTI software. **B** Left: Predicted structures of VarG using   AlphaFold2

(https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF

old2.ipynb). The region of the protein structure depicted in blue colour indicate the least error in prediction. Right: VarG and Gap structures overlapping using TMscore (https://zhanggroup.org/TM-score/). TM-score value of 1 indicates 100% structural identity among proteins.

Fig. S5   Quantification of intra- and extracellular metabolites from *V. cholerae*. Representative plots from identification of 1As3PG and 3PG from intracellular and extracellular samples from *V. cholerae* WT and $\Delta varG$ cells were grown with and without 1 mM As^V. The compounds were separated by UPLC using an XSELECT HSS XP column. 1As3PG was identified using a Thermo Orbitrap Exploris 480 high-resolution mass spectrometer and 3PG using a triple-quadrupole-ion trap hybrid mass spectrometer QTRAP® 6500+ (Sciex, USA) connected with a Waters ultra-performance liquid chromatography I-class (UPLC) system.

Fig. S6   VarH characterization. A Sequences alignment of the secondary structure of the VarH protein with the crystal structure of the human Kap PTP phosphatase using the software Phyre2 (www.sbg.bio.ic.ac.uk/phyre2). Human Kap structure matched VarH with 100% confidence and 99% coverage. Amino acid sequences within the black box (from the amino acid 113 (H) onwards) indicate the conserved catalytic loop (C(x)$_5$R) of protein tyrosine phosphatases PTP. Green and black boxes indicate deletions and insertions, respectively, respect the template sequence. **B** Predicted structures of VarH with AlphaFold2 (https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaF

old2.ipynb). The regions of the protein structure depicted in blue colour indicate no

error in prediction. Right: overlapping between VarH protein structure obtained with

AlphaFold2 and crystal structure of the Kap PTP protein using TMscore

(https://zhanggroup.org/TM-score/). TM-score value of 1 represents 100%

structural identity among proteins. **C** SDS-PAGE gel depicting the VarG His-tag

purified protein from *the V. cholerae* WT and *varH* mutant strains (see material and

methods for details). VarG proteins were excised from the gel and its

phosphorylation state was quantified by LC–MS/MS. **D** Phosphorylation state

quantification of VarG from *V. cholerae* WT and *varH* mutant strains. S204, serine

amino acid from VarG in position 204. Y108, tyrosine amino acid from VarG in

position 108. **E** Representative plots from identification of 1As3PG and 1AsG from

intracellular samples of *V. cholerae* WT and Δ*varH* cells grown with and without 1

mM As^V. The compounds were separated by UPLC using a XSELECT HSS XP

column. The detection of 1As3PG and 1AsG were perfomed using a Thermo

Orbitrap Exploris 480 High Resolution Mass Spectrometer. Data are the mean of

three biological replicates ± s.e.m.

Fig S7 As^{III} binding site of ArsR is not conserved in *V. cholerae* and *varH* is

specific of the Vibrionaceae family. A ArsR amino acids sequences alignment

from *V. cholerae* and representative enteropathogenic bacteria using Vector NTI

software. ArsR cysteine-containing As^{III}-binding motives are depicted in the lower

panel with cysteine residues labeled in red color. **B** Genomic organization

comparison of the *var* operon in the Vibrionaceae family and representative

bacteria by using the Patric software (https://www.patricbrc.org/). The presence of

varH is depicted in green color and under dotted rectangle.

Table S1. Complete list of genes identified under-represented by TIS in the

presence of 1 mM As^V with thresholds of fold change <0.1 and an inverse p-value

>75

Table S2. List of strains and plasmids used in this study.

Table S3. List of primers used in this study.

Supplementary references

1.   Chen J, Yoshinaga M, Garbinski LD, Rosen BP. Synergistic interaction of

glyceraldehydes-3-phosphate dehydrogenase and ArsJ, a novel

organoarsenical efflux permease, confers arsenate resistance. *Mol*

Microbiol 2016; **100**: 945–953.

2.   Miller VL, Mekalanos JJ. A novel suicide vector and its use in construction

of insertion mutations: Osmoregulation of outer membrane proteins and

virulence determinants in Vibrio cholerae requires toxR. *J Bacteriol* 1988;

170: 2575–2583.

3.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al.

Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Mol Syst Biol* 2006; **2**: 1–11.

4. Sansonetti PJ, Kopecko DJ, Formal SB. Involvement of a Plasmid in the Invasive Ability of Shigella flexneri. 1982; **35**: 852–860.

5. Cava F, De Pedro MA, Lam H, Davis BM, Waldor MK. Distinct pathways for modification of the bacterial cell wall by non-canonical D-amino acids. *EMBO J* 2011; **30**: 3442–3453.

6. del Peso Santos T, Alvarez L, Sit B, Irazoki O, Blake J, Warner BR, et al. BipA exerts temperature-dependent translational control of biofilm-associated colony morphology in Vibrio cholerae. *Elife* 2021; **10**: 1–23.

7. Donnenberg MS, Kaper JB. Construction of an eae deletion mutant of enteropathogenic Escherichia coli by using a positive-selection suicide vector. *Infect Immun* 1991; **59**: 4310–7.

8. Chiang SL, Rubin EJ. Construction of a mariner-based transposon for epitope-tagging and genomic targeting. *Gene* 2002; **296**: 179–185.

9. Guzman L, Belin D, Carson MJ. Tight Regulation, Modulation, and High-Level Expression by Vectors Containing the Arabinose PBAD Promoter. *J Bacteriol* 1995; **177**: 4121–4130.

10. Kimsey HH, Waldor MK. Vibrio cholerae LexA coordinates CTX prophage

1111   gene expression. *J Bacteriol* 2009; **191**: 6788–6795.

1112

1113

1114

1115

A

Identity percentage
0 40 80

S. flexneri (PRJNA553848)

S. enterica (PRJNA231221)

Y. pseudotuberculosis (PRJEB6403)

EHEC *(PRJNA218110)*

C. rodentium (PRJNA261533)

V. cholerae C6706

ArsR   VarG   VarH   ArsJ

B

○ Empty plasmid   ● +*var* operon

As^V resistance determinants

As^V resistance determinants

Arrayed screen

3.156 Tn insertion mutants

-As^V

Aliquot transferred into:

LB +As^V

Output
Growth analysis at OD$_{600}$

EON

vc1068	ArsR repressor protein
vc1069	GAPDH protein
vc1070	Putative phosphatase protein
vc1071	Membrane transporter protein

arsR

VC1068 VC1069 VC1070 VC1071

TIS

***V. Cholerae* + Himar1 Tn**

-As^V

Collection and inoculation into LB:

LB -As^V LB +As^V

Output
Mutant library recovery

*

Comparison of Tn-seq profiles

Inverse p-value

10^{20}
10^{15}
10^{10}
10^5
10^0

vc1071

vc1070
vc1069
vc1068

10^{-4} 10^{-3} 10^{-2} 10^{-1} 10^0 10^1 10^2 10^4

Log$_{10}$ (FC) -As^V/+As^V

A

Phosphatase activity

B

1As3PG