Species invasions shift microbial phenology in a two-decade freshwater time series

Robin R. Rohwer[1*], Riley J. Hale[2], Todd R. Miller[3], Katherine D. McMahon[2,4]

[1]Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA.

[2]Department of Civil and Environmental Engineering, University of Wisconsin-Madison; Madison, WI, USA.

[3]Zilber School of Public Health, University of Wisconsin-Milwaukee; Milwaukee, WI, USA.

[4]Department of Bacteriology, University of Wisconsin-Madison; Madison, WI, USA.

*Robin R. Rohwer

Email: Robin.Rohwer@gmail.com

Author Contributions: Conceptualization: RRR, KDM**;** Funding acquisition: RRR, TRM, KDM**;** Methodology: RRR, TRM**;** Investigation: RRR, RJH**;** Data curation: RRR**;** Formal analysis: RRR**;** Visualization: RRR**;** Writing – original draft: RRR**;** Writing – review & editing: RRR, RJH, TRM, KDM

Competing Interest Statement: Authors declare that they have no competing interests.

Keywords: Invasive Species, Microbial Ecology, Phenology, Limnology, Time Series

This PDF file includes:

Main Text
Figures 1 to 4
Supp Table Legends 1 to 4
Supp Tables 1 & 3
Supp Figures 1 to 2

Abstract

Invasive species impart abrupt changes on ecosystems, but their impacts on microbial communities are often overlooked. We paired a 20-year freshwater microbial community time series with zooplankton and phytoplankton counts, rich environmental data, and a 6-year cyanotoxin time series. We observed strong microbial phenological patterns that were disrupted by the invasions of spiny water flea (*Bythotrephes longimanus*) and zebra mussels (*Dreissena polymorpha*). First, we detected shifts in *Cyanobacteria* phenology. After the spiny water flea invasion, *Cyanobacteria* dominance crept earlier into clearwater; and after the zebra mussel invasion, *Cyanobacteria* abundance crept even earlier into the diatom-dominated spring. During summer, the spiny water flea invasion sparked a cascade of shifting diversity where zooplankton diversity decreased and *Cyanobacteria* diversity increased. Second, we detected shifts in cyanotoxin phenology. After the zebra mussel invasion, microcystin increased in early summer and the duration of toxin production increased by over a month. Third, we observed shifts in heterotrophic bacteria phenology. After both invasions, heterotrophic bacteria with putative cyanobacterial associations increased. The *Bacteroidota* phylum and members of the acI *Nanopelagicales* lineage were differentially more abundant. The diverse summer bacterial community was most resistant to change from the zebra mussel invasion despite the observed shifts in diversity and toxicity during the summer; but the spring and clearwater bacterial communities changed most from the spiny water flea invasion, which lessened clearwater duration and intensity. These long-term invasion-mediated shifts in microbial phenology demonstrate the interconnectedness of microbes with the broader food web, and their susceptibility to long-term environmental change.

Significance Statement

Microbial communities are typically studied as part of the microbial loop, separately from the broader food web. Using a two-decade freshwater time series, we explored whether two species invasions that shifted the metazoan food web also impacted the microbial community. We looked for seasonal responses because of strong seasonal patterns within the microbial community. We discovered that *Cyanobacteria* increased in the clearwater and spring seasons, and that *Cyanobacteria* diversity increased in the summer. Cyanotoxins also increased in early summer, and the duration of toxin production increased. In the heterotrophic bacterial community, many organisms that putatively interact with *Cyanobacteria* increased. These findings represent direct observations of long-term bacterial change, and of the interconnectedness of bacteria with the broader food web.

Main Text

Introduction

Invasive species have wide-ranging impacts on ecosystems, often rewiring food webs and disrupting nutrient cycling, and they can be even more disruptive than global abiotic change (1). However, the impact of species invasions on microbial communities are rarely considered. Microbes are usually considered separately from metazoan lake ecology; bacterial communities

70 are thought to interact with the broader food web as part of a microbial loop, where phytoplankton-derived dissolved organic matter is returned to higher trophic levels (2, 3). Bacteria that consume organic matter are in turn consumed by ciliates and nanoflagellates, which are eaten by microzooplankton, thus linking microbial communities with higher trophic levels. Although this microbial food web is often studied separately from the metazoan food web, we

75 hypothesized that metazoan species invasions would change the bacterial community.

We collected a 20-year microbial community time series (2000-2019, 496-samples) from Lake Mendota, a eutrophic temperate lake in WI, USA. This period includes two metazoan species invasions: the mesozooplankton spiny water flea in 2009 (*Bythotrephes longimanus*) and zebra mussels in 2015 (*Dreissena polymorpha*). The spiny water flea invasion triggered a food

80 web cascade that resulted in a one meter loss in water clarity (4), as increased zooplankton predation by spiny water flea led to reduced grazing pressure on phytoplankton. This impact was particularly strong during the spring clearwater phase, which decreased in intensity and duration (5). The invasion of zebra mussels triggered a 300% increase in benthic zooplankton and phytoplankton abundance, and they themselves became abundant enough to filter the epilimnion

85 volume every 45 days (6). Zebra mussels increase filtration, but they can also worsen *Cyanobacteria* blooms via selective retention of eukaryotic algae and release of Cyanobacteria back into the lake (7), and by altering nutrient stoichiometry (8). In Lake Mendota, zebra mussels did not impact the pelagic water clarity in either direction (6). These mid-time series disturbances set up a natural experiment with which we explored the impact of metazoan species invasions on

90 the bacterial community.

Lake Mendota is a temperate, dimictic lake and its ecology follows consistent seasonal patterns (phenology). The lake freezes in the winter, and ice-off is followed by spring mixing and a diatom bloom. As the water warms, zooplankton grazing increases and results in a short-lived period of high water clarity, which ends when planktivorous fish begin eating zooplankton (5).

95 After this clearwater phase, diatoms give way to *Cyanobacteria* which persist throughout the

summer (9). During the summer the water column is thermally stratified, and anoxia develops in the bottom layer while nutrient draw-down occurs in the upper layer. When temperatures drop in the fall, Lake Mendota mixes and remains mixed until ice-on. Seasonal patterns have also been observed in microbial community composition (10), and these seasonal cycles can mask long-

100   term changes (11). Therefore, we split our time series into seasons and looked for changes in bacterial phenology.

Here we identify seasons that described the bacterial phenology (Fig 1), and then identify phenological change in *Cyanobacteria* (Fig 2), cyanotoxins (Fig 3), and heterotrophic bacteria (Fig 4). These observed phenological shifts are a direct observation of long-term microbial

105   change in response to invasive species.

Results

We characterized the microbial community composition via 16S rRNA gene amplicon

110   sequencing performed on duplicate water grabs (961 total samples, 496 sample dates) (Fig 1). The water grabs integrated the top 12 m of the water column, which approximates the mixed upper layer. In addition to the sequencing-based relative abundances, we obtained environmental data available from the North Temperate Lakes Long Term Ecological Research program (NTL-LTER), including water physical and chemical measurements and microscopy-based counts of

115   zooplankton and phytoplankton.

Microbial Phenology

To compare across years, we defined lake seasons using environmental variables and compared samples within the same season. We compared across multiple years by season instead of by Julian date to discriminate shifts in each season's community composition from shifts in seasonal timing. We created multiple potential season definitions based on

120   environmental data, and then selected the definitions that were most representative of the microbial phenology. We applied non-metric multidimensional scaling (NMDS) (12) to sample distance matrices calculated with the phi distance metric for centered log-ratio (CLR)-transformed relative abundances (13). We selected the season definitions most relevant to the microbial

125   community based on the clarity of grouping we observed in NMDS plots of season boundaries (Fig 1). Samples grouped by our selected season definitions were significantly less variable within groups than between groups for every season boundary (ANOSIM < 0.001).

The season definitions that best described the microbial phenology were ice-on, spring, clearwater, early summer, late summer, and fall (Fig 1, Table S1). The ice-on season was

130 determined by the onset and break-up of continuous ice-cover across a central lake transect. Spring began with ice-off and ended with clearwater onset, spanning a period that includes the spring diatom bloom (9). Clearwater was manually defined based on water clarity measurements, and is an annual phase driven by high zooplankton grazing pressure (5). Early summer was a period of growing stratification, bounded by clearwater and the point at which hypolimnetic anoxia

135 extended above 12 m. Late summer was a period of strong stratification and anoxia, which ended with fall mixing. We defined mixing as a difference in average epilimnion and hypolimnion temperatures $< 3°C$. Fall began with mixing and ended with ice-on.

Shifts in *Cyanobacteria* phenology

Given spiny water flea's marked disruption of clearwater phenology, *i.e.* shortened

140 duration and intensity, we first looked for changes in spring phytoplankton phenology. Historically, the end of clearwater phase marks the transition from a diatom-dominated system to a *Cyanobacteria*-dominated system (9). We found that after the spiny water flea invasion, *Cyanobacteria* began appearing during clearwater phase (Fig 2A). We did not see a similar increase in chloroplast relative abundance or diatom counts during clearwater phase, indicating

145 that the transition to *Cyanobacteria* dominance shifted earlier. We further found that after the zebra mussel invasion, *Cyanobacteria* began appearing even before the start of clearwater phase (Fig 2A). Count data confirmed that the biovolume of *Cyanobacteria* increased 2-fold, growing to over 10% of total phytoplankton (Fig S1). This earlier creep of *Cyanobacteria* describes a kingdom-level regime shift from a eukaryotic algae-dominated period to one that includes

150 *Cyanobacteria*.

After identifying these changes in the spring phenology, we next looked for spiny water flea invasion impacts on the summer cyanobacterial phenology. *Cyanobacteria abundance (relative to the microbial community)* during summer and fall did not change with the spiny water flea invasion. However, *Cyanobacteria* diversity during the late summer season increased (Fig

155 2B). This increase in cyanobacterial diversity was observed at fine and coarse-resolution taxonomic levels, including levels as coarse as order (Fig 2B). It is possible that the shift in summer diversity could be explained by successional processes since the timing of the *Cyanobacteria*-dominated season shifted earlier; however, we also observed a concurrent change in zooplankton diversity. After the spiny water flea invasion zooplankton diversity

160 decreased (Fig 2B, Table S2). Additionally, the spring cyanobacterial incursion was comprised of a single amplicon sequence assigned to *Aphanizomenon*, not to a more complex community

typical of early summer. Therefore, we attribute the shift in cyanobacterial summer diversity to a cascade of shifting diversity originating in the zooplankton community. The lower diversity of zooplankton may have resulted in a narrower grazing pressure, allowing a wider range of
165   *Cyanobacteria* to flourish.

Next, we looked for changes in the summer cyanobacterial community stemming from the zebra mussel invasion. We first compared total *Cyanobacteria* abundance before and after zebra mussels but did not see strong and consistent changes in relative abundance or biovolume (Fig S1). We also did not observe strong shifts in cyanobacterial community composition.
170   *Cyanobacteria* can produce potent toxins, but *Cyanobacteria* abundance does not always correlate with cyanotoxin concentrations (14, 15), suggesting that toxin concentrations could yet have changed as a result of the invasion.

Shifts in cyanotoxin phenology

To examine the impact of zebra mussels on cyanobacterial toxicity, we collected a 6-
175   year, 246-sample toxin time series spanning the 3 years preceding and following the zebra mussel invasion. We measured the hepatotoxin microcystin and the bioactive peptide anabaenopeptin using high performance liquid chromatography coupled with tandem mass spectrometry (HPLC-MS/MS) (Table S3) (16). We observed a 3-fold decrease in anabaenopeptin and a concurrent 3-fold increase in microcystin during early summer ($p < 0.05$) (Fig 3A). The
180   family *Microcystaceae* is considered the main microcystin producer in Lake Mendota, although it is also capable of producing anabaenopeptin (14). However, the relative abundance of *Microcystaceae* did not change. This suggests that the switch from anabaenopeptin to microcystin is due to more complex ecological drivers than the abundance of *Microcystaceae*, and illustrates the complexity of invasion impacts. This shift in toxin concentrations has
185   implications for human health, because microcystin is a more potent toxin than anabaenopeptin (17).

After observing this increase in early summer microcystin, we next explored the phenology of microcystin production (Fig 3B). Total toxin concentrations were variable between years, so for each year we normalized total microcystin by the maximum measured concentration
190   within that year. We found that the duration of toxin production, defined as the period during which toxins were at least 15% of maximum annual toxicity, increased after the zebra mussel invasion by 53 ± 9 days ($p < 0.001$) (Fig 3C). This increase was due to a longer late summer season, as well as toxin production extending further into the early summer and fall seasons. We observed these increases in toxicity despite the lack of change in the *Cyanobacteria* community

195   composition. This could be explained by preferential retention of non-toxic strains or individual

cells, with toxic cells released back into the water column. Zebra mussels were previously found

to preferentially retain a non-toxic *Microcystis* strain over a toxic one (18).

In summary, cyanobacterial phenology was profoundly affected by the two metazoan

invasions, with changes in timing, diversity, and toxicity.

200   **Shifts in heterotrophic bacteria phenology**

To explore whether invasion impacts extended further, into the heterotrophic bacterial

community, we compared each season's heterotrophic community before and after invasion

using NMDS plots of their CLR-transformed phi dissimilarities (13). We observed modest

grouping based on invasion status (Fig S2), and an ANOSIM analysis confirmed that within-group

205   differences were smaller than between-group differences in all seasons (ANOSIM significance =

0.001).

To identify which bacteria were responsible for the shift in community composition, we

performed a differential abundance analysis after applying a CLR transformation to account for

the relativized nature of sequencing data (19). We used a generalized linear model to test for

210   organisms that were differentially abundant before and after each invasion (20), and we looked

for changes within each season. We considered an operational taxonomic unit (OTU)

differentially abundant when it had an effect size > 0.5, and we manually confirmed the ecological

significance of this effect size cutoff by inspecting relative abundance over time (examples in Fig

4).

215   To assess which seasonal communities were most susceptible to change, we looked at

the proportion of OTUs within the community that were differentially abundant. To better reflect

the ecological impact of the differentially abundant OTUs, we normalized the OTUs by their mean

relative abundance. The invasions impacted seasonal community composition to different

degrees. For example, spiny water flea had the largest impact on the proportion of the community

220   that changed during spring and clearwater, with 38% of the average community becoming

differentially abundant (Fig 4C, left). This means that in addition to changing the seasonal timing

of clearwater (5), spiny water flea also changed the heterotrophic community composition.

Meanwhile, zebra mussels had the smallest impact on the late summer microbial community, with

only 4% of the average community becoming differentially abundant (Fig 4C, right). The late

225   summer microbial community had the smallest proportion of differentially abundant organisms,

and also the fewest overall differentially abundant OTUs (Fig 4). The late summer community

also has the highest alpha diversity, and biodiversity has been linked to ecosystem stability (21). It is possible that this high diversity makes the late summer ecosystem more robust. These seasonally-distinct invasion impacts emphasize the importance of a phenological context.

230   Among the organisms that were differentially abundant, OTUs in the phylum *Bacteroidota* were zebra mussel invasion "winners" broadly (Fig 4A, Table S4). These winners included abundant OTUs from the bacI, bacII, bacIII, and bacIV lineages (*Chitinophagales*, *Flavobacteriales*, and *Cytophagales* orders, lineage nomenclature from (22, 23)). However, most changes in differential abundance were not consistent within taxonomic lineages or across

235   seasons. For example, within the abundant lineages of order *Frankiales*, *Nanopelagicaceae* (acI) tended to win while acTH1 and acSTL tended to lose, and these changes were detected primarily in spring and early summer seasons (Fig 4B, Table S4).

        The *Bacteroidota* winners are likely particle-associated organisms, and many are known to associate with *Cyanobacteria* (23, 24). They were broadly differentially abundant after both the

240   spiny water flea invasion, which increased *Cyanobacteria* diversity, and the zebra mussel invasion, which increased *Cyanobacteria* toxicity. This suggests that the shifts in cyanobacterial phenology may have benefitted their associated heterotrophic organisms.

        In contrast, the *Frankiales* order is comprised of small, free-living bacteria. Their non-uniform response suggests that they occupy niche spaces that are not changing coherently. For

245   example, the acI lineage can uptake cyanobacterial exudates such as amino acids and polypeptides (25, 26), and acI tended to win, especially during seasons where they already dominated. The acI lineage is oligotrophic (27), and acI-A and acI-B tended to win during clearwater and early summer, while acI-C has been found in closer association with Cyanobacteria (28) and it tended to win during fall, when cyanoblooms still plague Lake Mendota.

250   Like acI, acTH1 and acSTL are small, free-living *Frankiales*, but little is known about their ecology and metabolism. Since they comprised many of the abundant invasion losers, this suggests that their niche is different from their close relatives'.

        In summary, heterotrophic bacteria phenology was shifted by metazoan species invasions, with organisms that may be more dependent on *Cyanobacteria* tending to win. These changes

255   were seasonal, and some seasonal communities were more robust to change than others.

Discussion

260   In addition to invasive species, the interacting driver of climate change is shifting the phenology of northern hemisphere lakes. Winters are shortening (29), summer stratification and anoxia is increasing (30), and extreme weather is becoming more common (31). In Lake Mendota specifically, ice duration (32) and clearwater (5) are shortening, while the stratified (32) and anoxic (33) periods are lengthening. We posit that lake microbial communities will also change in

265   response to climate change because the lake microbial community has phenological patterns that track these shifting seasons. Further, we directly observed that food web-mediated shifts in seasonal timing can change microbial community timing, composition, and toxicity. It follows that the interacting driver of climate change shifting seasons over longer time frames will likely shift microbial phenology too.

270   Moreover, the species invasions themselves were driven by climate change. Previous work on Lake Mendota found that an unusually cool year allowed a sleeper population of spiny water flea to explode (34), and weather extremes like this are expected to increase with climate change (31). It is possible that the spiny water flea-mediated food web shift also paved the way for the zebra mussel invasion in a series of events termed an invasion meltdown (35). Zebra

275   mussels likely had multiple introductions to Lake Mendota based on the large amount of recreational boat traffic on the lake, but they did not establish prior to 2015. We speculate that spiny water flea enabled the establishment of zebra mussels, thus linking the zebra mussel invasion with the same climate anomaly that triggered the spiny water flea population explosion. We found that after the spiny water flea invasion fall eukaryotic algae biovolume increased 20-

280   fold, while after the zebra mussel invasion it returned to pre-spiny water flea levels, suggesting that zebra mussels may have consumed the additional eukaryotic algae (Fig S1). A climate anomaly triggered a population explosion of the invasive spiny water flea sleeper population, and the resulting food web disruptions may have paved the way for the establishment of zebra mussels in an invasion meltdown. The impact of these invasions reached all the way into the

285   bacterial community, illustrating how the interacting drivers of species invasions and climate change can have complex and far-reaching impacts.

        Harmful cyanobacterial blooms are predicted to increase with warmer temperatures, higher CO_2, and increasingly intense rain events (36) We observed an earlier creep of Cyanobacteria into the clearwater and spring seasons. In other lakes, regime changes have been

290   observed as oligotrophic systems switch to a new stable state dominated by *Cyanobacteria* (37). It is possible that we may be observing the beginning of a transition to a new stable state at a seasonal level. In the summer seasons already dominated by *Cyanobacteria*, we found the fewest differentially abundant organisms. This suggests that even if tipping points are reached in

295 other seasons, the community is less likely to shift when it would be most desired: during the summer months plagued by toxic cyanobacterial blooms.

Long-term change is difficult to distinguish from short-term variability and seasonal cycles, a phenomenon termed the "invisible present" (11). Observations of the microbial community for almost twenty years have allowed us to distinguish consistent phenological patterns, and to identify separate long-term, invasion-mediated shifts. The establishment and

300 continuation of time series like ours are the only way to directly observe the long-term impacts of environmental change. Our study found that microbial communities are sensitive to long-term change, specifically to the cascading impacts of invasive species. These impacts included many aspects of phenology, from timing and abundance, to diversity and community composition, to human health and cyanobacterial toxicity.

Materials and Methods

Bacterial Community Data

310 Microbial sampling, sequencing, and data processing was performed as described in detail in Rohwer & McMahon *(submitted)* (38). Briefly, integrated epilimnion samples were collected from the central deep hole of Lake Mendota (WI, USA) using a 12 m tube, and water was well mixed before filtering onto a 0.22 um filter (Pall Corporation). Filters were stored in a -80°C freezer until DNA extraction in 2018-2019. DNA was extracted using a FastDNA Spin Kit

315 (MP Biomedicals) after sample order was randomized. Samples were sent to Argonne National lab for amplification of the 16S rRNA gene V4-V5 region (515F-Y GTGYCAGCMGCCGCGGTAA, 926R CCGYCAATTYMTTTRAGTTT) and sequencing on an Illumina MiSeq instrument.

After sequencing, data was processed using custom R scripts available at

320 https://github.com/rrohwer/limony and described in Rohwer & McMahon (*submitted*) (38). Microbial relative abundances from replicate samples were averaged, and a standard deviation was calculated. This processed data is available as part of an R package at https://github.com/rrohwer/limony. The raw sequencing data is available through the NCBI Sequence Read Archive under accession number PRJNA846788 (Reviewer link:

325 https://dataview.ncbi.nlm.nih.gov/object/PRJNA846788?reviewer=varcj4a5hng2m7n1bcde77oecg).

Environmental Data

We collected lake physical and chemical data alongside microbial samples, including
330   water temperature and dissolved oxygen measured with a YSI ProPlus multiparameter sonde
(Yellow Springs Instruments), and water clarity measured with a Secchi disk. This data is
available through the Environmental Data Initiative (EDI) portal under identifiers knb-lter-ntl.415.2
(39) and knb-lter-ntl.416.1 (40). We combined our measurements with additional water
temperature, dissolved oxygen, water clarity, and ice cover datasets made available by the North
335   Temperate Lakes Long-Term Ecological Research program (NTL-LTER) under EDI identifiers
knb-lter-ntl.29.29 (41), knb-lter-ntl.130.29 (42), knb-lter-ntl.335.1 (43), knb-lter-ntl.400.2 (44),
knb-lter-ntl.31.30 (45), and knb-lter-ntl.33.35 (46).

We defined lake seasons using metrics calculated from this environmental data. We
340   identified the epilimnion, metalimnion, and hypolimnion depths by identifying the inflection points
of temperature profile vectors, calculated the average epilimnion and hypolimnion temperatures
for all complete profiles, and smoothed these values with a 7-day moving average. We defined
mixing as the day when the linearly interpolated difference in epilimnion and hypolimnion
temperature fell below 3°C. We considered depths with < 1 mg/L dissolved oxygen anoxic, and
345   we applied a 7-day moving average to the depth of the anoxic layer. We defined the start of the
late summer season as the day when the linearly interpolated anoxic layer depth reached 12 m.
We manually defined the clearwater season's start and end for each year based on the Secchi
depth measurements.

350   We obtained absolute abundance and biovolume of large *Cyanobacteria* and eukaryotic
algae using microscopy-based phytoplankton counts from the NTL-LTER, and we used
microscopy-based zooplankton counts to calculate zooplankton diversity. This data is available
under EDI identifiers knb-lter-ntl.88.30 (47) and knb.lter.ntl.90.31 (48).

355   **Cyanotoxin Data**

We measured toxins in 246 whole-water samples spanning 6 years (2013-2018). These
samples were taken from the same water grabs used to characterize the microbial community.
We quantified cyanotoxins using methods described in detail by Miller *et al.* (16). Briefly,
lyophilized samples were resuspended in formic acid, subjected to three freeze-thaw cycles, and
360   extracted in methanol and formic acid in a sonicating water bath. The extract supernatant was

analyzed by targeted high performance liquid chromatography coupled with tandem mass spectrometry (HPLC-MS/MS).

Eleven microcystin (MC) congeners and three anabaenopeptin (Apt) congeners were
365   analyzed. Certified reference standards of MC-LR, [Dha⁷]MC-LR and nodularin were purchased from the National Research Council of Canada Biotoxins program (Halifax, Nova Scotia). MC-LA, MC-RR, MC-LF, MC-YR, MC-WR, MC-LY, MC-LW, MC-HtyR, MC-HilR (all >95%) were purchased from Enzo Life Sciences (Farmingdale, NY). Anabaenopeptin A (Apt-A) (> 95%), Apt-B (>95%), and Apt-F (> 95%) were purchased from MARBIONC (Wilmington, NC). To calculate
370   total concentrations, we summed nanomolar measurements of the congeners. Toxin totals are available in a tabular format as Table S2.

Analysis

We defined microbially-relevant seasons (Fig 1A) based on the environmental data. Ice
375   extent, mixing, and clearwater were phenological markers that corresponded well with the microbial community. To divide the summer months we explored multiple seasonal cutoffs based on stratification extent, epilimnion temperature, and anoxia extent. We visualized each season option with non-metric multidimensional scaling (NMDS) plots and manually chose the most relevant definitions based on visual inspection of the NMDS plots. To account for the proportional
380   nature of sequencing data, we performed a centered log ratio (CLR) transformation on our data before creating NMDS plots (Fig 1B, Fig S2). Following the guidance provided by Quinn *et al.* (19), we filtered each 3-season group to contain only OTUs that occurred in at least 10% of the samples, assigned small-values to zeros using the zCompositions R package (49), and performed a CLR-transform and calculated the *phi* distance metric using the propr R package
385   (13). We performed NMDS and confirmed the significance of our season choices by calculating the ANOSIM significance using the vegan R package (12). The season start dates are listed in Table S1.

We observed changes in spring *Cyanobacteria* relative abundance (Fig 2A) by aligning
390   samples by the start of clearwater phase and grouping years into invasion periods. Both invasions were discovered in fall, so we considered the subsequent years the start of the invasion period. The seven 2019 samples were excluded from this analysis because they did not span the entire spring-clearwater-early summer period. We compared the summer diversity before and after the spiny water flea invasion (Fig 2B) by aligning samples by the start of the early summer
395   season and calculating the Shannon diversity index using the vegan R package (12). We

observed no significant difference in diversity between the spiny water flea-only period and the
zebra mussel period, so we grouped these two periods together as a single post-spiny water flea
group. Before calculating *Cyanobacteria* diversity, we excluded samples from 2000-2003 that had
been prefiltered with a 10 μm filter instead of filtered directly onto a 0.22 μm filter. Before

400   calculating zooplankton community diversity (Fig 2B), we excluded spiny water flea and juvenile
copepods from the zooplankton count data and grouped zooplankton based on ecological
relevance and our confidence in the identifications (Table S3). We calculated the mean and
standard deviation of linearly interpolated daily diversity values and performed a t-test to compare
between invasion groups.

         We calculated the concentration of toxins before and after the zebra mussel invasion (Fig
3A) by separating toxin measurement dates by season and splitting the years into invasion
groups. For each season and year group, we calculated the mean and standard deviation of all
toxin measurements within it. We performed a t-test on each seasonal block to identify significant

410   change between invasion groups. We calculated the duration of toxin production (Fig 3B-C) by
normalizing microcystin concentrations in each year by the yearly maximum concentration. We
applied a linear interpolation to the 7-day moving average of measurements (lines in Fig 3B). We
defined days as toxin-producing if their interpolated daily toxin concentration was > 15 % of the
yearly maximum value.

         We identified differentially abundant bacteria (Fig 4) using the ALDEx2 package (50)
following the guidance provided by Quinn *et al.* (19). For each season group, we removed OTUs
present in fewer than 10% of samples and added small numbers to remaining zeros using
zCompositions (49). We identified differentially abundant organisms and their effect sizes using a

420   generalized linear model with the aldex.glm and aldex.glm.effect functions from the ALDEx2
package (50). We manually chose an effect size cutoff of 0.5 by examining plots of OTU
abundance over time for OTUs with varying effect sizes (examples in Fig 4A and Fig 4B). We
identified which seasons were most susceptible to microbial community change (Fig 4C) by
calculating the proportion of the community that was differentially abundant. We defined this

425   proportion as the average abundance of all differentially abundant OTUs divided by the average
abundance of all OTUs. All differentially abundant bacteria are listed in Table S4.

Acknowledgments

430   We thank Jake Vander Zanden for input on the manuscript, as well as Stephen Carpenter and Edward DeLong for friendly reviews. We thank technicians Ame Xiong and Leah Stromberg for assistance measuring cyanotoxins. We thank Jake Walsh for assistance interpreting the zooplankton data. We thank the many field personnel involved in microbial sampling, including sampling leads Angela Kent, Tony Yannarell, Ashley Shade, Stuart Jones, Ryan Newton,

435   Georgia Wolfe, Emily Kara Read, Lucas Beversdorf, and James Muetschler; and initial Microbial Observatory lead Eric W. Triplett.

We thank the funding sources that made this work possible:

U.S. National Science Foundation Postdoctoral Research Fellowship in Biology, award number

440   2011002 (RRR)

E. Michael and Winona Foster WARF Wisconsin Idea Fellowship, 2018 (RRR)

National Institute of Food and Agriculture, U.S. Department of Agriculture, award number 2016-67012-24709 (Joshua J. Hamilton)

National Institute of Food and Agriculture, U.S. Department of Agriculture, Hatch Projects

445   WIS01516, WIS01789, WIS03004 (KDM)

U.S. National Science Foundation North Temperate Lakes Long-Term Ecological Research site, NTL-LTER, award numbers DEB-9632853, DEB-0217533 (Stephen Carpenter) and DEB-0822700, DEB-1440297 (Emily Stanley)

U.S. National Science Foundation Microbial Observatories program, award numbers MCB-

450   9977903 (Eric Triplett) and DEB-0702395 (KDM)

U.S. National Science Foundation INSPIRE award, DEB-1344254 (KDM)

The Laboratory for Aquatic Environmental Microbiology & Chemistry at the University of Wisconsin-Milwaukee, Zilber School of Public Health (TRM)

References

1.   B. E. Lopez, *et al.*, Global environmental changes more frequently offset than intensify detrimental effects of biological invasions. *Proceedings of the National Academy of Sciences* **119**, e2117389119 (2022).

2.   F. Azam, *et al.*, The Ecological Role of Water-Column Microbes in the Sea. *Marine Ecology Progress Series* **10**, 257–263 (1983).

3.   K. Piwosz, I. Mukherjee, M. M. Salcher, V. Grujčić, K. Šimek, CARD-FISH in the Sequencing Era: Opening a New Universe of Protistan Ecology. *Frontiers in Microbiology* **12** (2021).

4. J. R. Walsh, S. R. Carpenter, M. J. V. Zanden, Invasive species triggers a massive loss of ecosystem services through a trophic cascade. *PNAS* **113**, 4081–4085 (2016).

5. S.-I. S. Matsuzaki, *et al.*, Climate and food web effects on the spring clear-water phase in two north-temperate eutrophic lakes. *Limnol. Oceanogr.* **n/a**, 1–17 (2020).

6. M. J. Spear, *et al.*, Early changes in the benthic community of a eutrophic lake following zebra mussel (Dreissena polymorpha) invasion. *Inland Waters* **0**, 1–19 (2022).

7. R. Naddafi, K. Pettersson, P. Eklöv, The effect of seasonal variation in selective feeding by zebra mussels (Dreissena polymorpha) on phytoplankton community composition. *Freshwater Biology* **52**, 823–842 (2007).

8. O. Bykova, A. Laursen, V. Bostan, J. Bautista, L. McCarthy, Do zebra mussels (Dreissena polymorpha) alter lake water chemistry in a way that favours Microcystis growth? *Science of The Total Environment* **371**, 362–372 (2006).

9. C. C. Carey, P. C. Hanson, R. C. Lathrop, S. Amand, A. L, Using wavelet analyses to examine variability in phytoplankton seasonal succession and annual periodicity. *J Plankton Res* **38**, 27–40 (2016).

10. A. Shade, *et al.*, Interannual dynamics and phenology of bacterial communities in a eutrophic lake. *Limnol. Oceanogr.* **52**, 487–494 (2007).

11. J. J. Magnuson, Long-Term Ecological Research and the Invisible Present. *BioScience* **40**, 495–501 (1990).

12. J. Oksanen, *et al.*, *vegan: Community Ecology Package* (2020).

13. T. P. Quinn, M. F. Richardson, D. Lovell, T. M. Crowley, propr: An R-package for Identifying Proportionally Abundant Features Using Compositional Data Analysis. *Scientific Reports* **7**, 16252 (2017).

14. L. J. Beversdorf, S. D. Chaston, T. R. Miller, K. D. McMahon, Microcystin mcyA and mcyE Gene Abundances Are Not Appropriate Indicators of Microcystin Concentrations in Lakes. *PLOS ONE* **10**, e0125353 (2015).

15. Robert E. Zuellig, Jennifer L. Graham, Erin A. Stelzer, Keith A. Loftin, Barry H. Rosen, "Cyanobacteria, Cyanotoxin Synthetase Gene, and Cyanotoxin Occurrence Among Selected Large River Sites of the Conterminous United States, 2017–18" (U.S. Department of the Interior, U.S. Geological Survey, 2021).

16. T. R. Miller, S. L. Bartlett, C. A. Weirich, J. Hernandez, Automated Subdaily Sampling of Cyanobacterial Toxins on a Buoy Reveals New Temporal Patterns in Toxin Dynamics. *Environ. Sci. Technol.* **53**, 5661–5670 (2019).

17. T. R. Miller, L. J. Beversdorf, C. A. Weirich, S. L. Bartlett, Cyanobacterial Toxins of the Laurentian Great Lakes, Their Toxicological Effects, and Numerical Limits in Drinking Water. *Marine Drugs* **15**, 160 (2017).

18. G. Juhel, *et al.*, Pseudodiarrhoea in zebra mussels Dreissena polymorpha (Pallas)exposed to microcystins. *Journal of Experimental Biology* **209**, 810–816 (2006).

19. T. P. Quinn, *et al.*, A field guide for the compositional analysis of any-omics data. *GigaScience* **8**, giz107 (2019).

20. A. D. Fernandes, *et al.*, Unifying the analysis of high-throughput sequencing datasets: characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data analysis. *Microbiome* **2**, 15 (2014).

21. T. H. Oliver, *et al.*, Biodiversity and Resilience of Ecosystem Functions. *Trends in Ecology & Evolution* **30**, 673–684 (2015).

22. R. R. Rohwer, J. J. Hamilton, R. J. Newton, K. D. McMahon, TaxAss: Leveraging a Custom Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *mSphere* **3**, e00327-18 (2018).

23. R. J. Newton, S. E. Jones, A. Eiler, K. D. McMahon, S. Bertilsson, A Guide to the Natural History of Freshwater Lake Bacteria. *Microbiology and Molecular Biology Reviews* **75**, 14–49 (2011).

24. M. W. Hall, R. R. Rohwer, J. Perrie, K. D. McMahon, R. G. Beiko, Ananke: temporal clustering reveals ecological dynamics of microbial communities. *PeerJ* **5**, e3812 (2017).

25. T. W. Ghylin, *et al.*, Comparative single-cell genomics reveals potential ecological niches for the freshwater acI Actinobacteria lineage. *ISME J* **8**, 2503–2516 (2014).

26. M. M. Salcher, T. Posch, J. Pernthaler, In situ substrate preferences of abundant bacterioplankton populations in a prealpine freshwater lake. *ISME J* **7**, 896–907 (2013).

27. R. J. Newton, S. L. McLellan, A unique assemblage of cosmopolitan freshwater bacteria and higher community diversity differentiate an urbanized estuary from oligotrophic Lake Michigan. *Frontiers in Microbiology* **6**, 1028 (2015).

28. M. A. Berry, *et al.*, Cyanobacterial harmful algal blooms are a biological disturbance to Western Lake Erie bacterial communities. *Environ Microbiol* **19**, 1149–1162 (2017).

29. S. Sharma, *et al.*, Loss of Ice Cover, Shifting Phenology, and More Extreme Events in Northern Hemisphere Lakes. *Journal of Geophysical Research: Biogeosciences* **126**, e2021JG006348 (2021).

30. R. I. Woolway, *et al.*, Phenological shifts in lake stratification under climate change. *Nat Commun* **12**, 2318 (2021).

31. T. D. Loecke, *et al.*, Weather whiplash in agricultural regions drives deterioration of water quality. *Biogeochemistry* **133**, 7–15 (2017).

32. M. R. Magee, C. H. Wu, D. M. Robertson, R. C. Lathrop, D. P. Hamilton, Trends and abrupt changes in 104 years of ice cover and water temperature in a dimictic lake in response to air temperature, wind speed, and water clarity drivers. *Hydrology and Earth System Sciences* **20**, 1681–1702 (2016).

33. R. Ladwig, *et al.*, Lake thermal structure drives interannual variability in summer anoxia dynamics in a eutrophic lake over 37 years. *Hydrology and Earth System Sciences* **25**, 1009–1032 (2021).

34. J. R. Walsh, S. E. Munoz, M. J. Vander Zanden, Outbreak of an undetected invasive species triggered by a climate anomaly. *Ecosphere* **7**, n/a-n/a (2016).

35. D. Simberloff, B. Von Holle, Positive Interactions of Nonindigenous Species: Invasional
545 Meltdown? *Biological Invasions* **1**, 21–32 (1999).

36. J. Huisman, *et al.*, Cyanobacterial blooms. *Nat Rev Microbiol* **16**, 471–483 (2018).

37. M. Scheffer, S. Carpenter, J. A. Foley, C. Folke, B. Walker, Catastrophic shifts in ecosystems. *Nature* **413**, 591–596 (2001).

38. R. R. Rohwer, K. D. McMahon, Lake iTAG measurements over nineteen years, introducing
550 the limony dataset. *in prep.* (in prep.).

39. R. R. Rohwer, K. D. McMahon, Lake Mendota Microbial Observatory Temperature, Dissolved Oxygen, pH, and conductivity data, 2006-present. *Environmental Data Initiative* **knb-lter-ntl.415** (2022).

40. R. R. Rohwer, K. D. McMahon, Lake Mendota Microbial Observatory Secchi Disk
555 Measurements 2012-present. *Environmental Data Initiative* **knb-lter-ntl.416** (2022).

41. J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Physical Limnology of Primary Study Lakes 1981 - current. *Environmental Data Initiative* **knb-lter-ntl.29** (2021).

42. J. Magnuson, S. Carpenter, E. Stanley, North Temperate Lakes LTER: High Frequency
560 Water Temperature Data - Lake Mendota Buoy 2006 - current. *Environmental Data Initiative* **knb-lter-ntl.130** (2020).

43. D. M. Robertson, Lake Mendota water temperature secchi depth snow depth ice thickness and meterological conditions 1894 - 2007. *Environmental Data Initiative* **knb-lter-ntl.335** (2016).

565 44. J. J. Magnuson, S. R. Carpenter, E. H. Stanley, Lake Mendota Multiparameter Sonde Profiles: 2017 - current. *Environmental Data Initiative* **knb-lter-ntl.400** (2021).

45. J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Secchi Disk Depth; Other Auxiliary Base Crew Sample Data 1981 - current. *Environmental Data Initiative* **knb-lter-ntl.31** (2021).

570 46. J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Ice Duration - Madison Lakes Area 1853 - current. *Environmental Data Initiative* **knb-lter-ntl.33** (2021).

47. J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Phytoplankton - Madison Lakes Area 1995 - current. *Environmental Data Initiative* **knb-lter-ntl.88** (2022).

575 48. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Zooplankton - Madison Lakes Area 1997 - current. *Environmental Data Initiative* **knb-lter-ntl.90** (2010).

49. J. Palarea-Albaladejo, J. A. Martín-Fernández, zCompositions — R package for multivariate imputation of left-censored data under a compositional approach. *Chemometrics and Intelligent Laboratory Systems* **143**, 85–96 (2015).

580   50.   A. D. Fernandes, J. M. Macklaim, T. G. Linn, G. Reid, G. B. Gloor, ANOVA-Like Differential
      Expression (ALDEx) Analysis for Mixed Population RNA-Seq. *PLOS ONE* **8**, e67019
      (2013).

Figures and Tables

590   **Figure 1. Sample Dates and microbial phenology. (A)** Sample dates are represented by

points, and season durations are represented by shading. **(B)** Non-metric multidimensional

scaling of CLR-transformed bacterial community compositions. Points represent sample dates,

and colors indicate seasons. Three seasons are compared in each NMDS plot to highlight season

boundaries. All NMDS stress values are ≤ 0.1.

Figure 2. Shifts in *Cyanobacteria* phenology. (A) Spring relative abundance of *Cyanobacteria*
aligned as days since clearwater. Average and standard deviation duration of clearwater is
indicated by grey bars. Blue represents the pre-invasion period of 2000-2009, orange the post-
spiny water flea period of 2010-2015, and red the post-zebra mussel period of 2016-2018. **(B)**
Summer diversity of zooplankton and *Cyanobacteria*. Thick lines represent the mean Shannon
diversity and shading represents the standard deviation. Thin lines represent individual years.
Here the orange post-spiny water flea period includes years 2010-2018. Grey starred regions
indicate the period of significant difference in mean diversity (p < 0.05). *Cyanobacteria* diversity is
shown at the order level, and Zooplankton diversity is shown at ecologically relevant groupings
with spiny water flea counts removed.

Figure 3. Shifts in cyanotoxin phenology. Orange represents the pre-zebra mussel invasion

615   period of 2013-2015, red represents the post-zebra mussel invasion period of 2016-2018. Stars

represent significant differences (p < 0.05). **(A)** Concentration of ananbaenopeptin and

microcystin during the seasons with regular toxicity. **(B)** Microcystin toxin concentrations

normalized by the yearly maximum concentrations to compare annual patterns. Shaded regions

represent the duration of toxin production, defined as days when toxins were present at ≥ 15% of

620   maximum toxicity. **(C)** Toxin production duration compared before and after the invasion of zebra

mussels.

Figure 4. Shifts in heterotrophic bacteria phenology. Differential abundance was calculated on CLR-transformed abundances with a GLM model. Pink represents "winners" that differentially increased, and teal represents "losers" that differentially decreased. Points represent OTUs, and point sizes are log-scaled by the OTU's mean abundance. **(A)** Left, an example relative abundance over time curve of an OTU with positive effect size. Right, effect sizes for differential abundance of phylum *Bacteroidota* OTUs around the zebra mussel invasion. **(B)** Middle, effect sizes for differential abundance of order *Frankiales* OTUs around the zebra mussel invasion. Right and left are example winner and loser OTU abundances over time. **(C)** Percent of each season's microbial community that was differentially abundant over time. OTUs were normalized by their mean abundance before calculating the percentage of community change. The impact of the spiny water flea invasion is shown on the left, and the impact of the zebra mussel invasion is shown on the right.

Supplemental Figures and Tables

Table S1. Season start dates. The dates for the start of each season are listed for every year.

Year	Spring	Clearwater	Early Summer	Late Summer	Fall	Ice-on
2000	2000-03-07	2000-04-12	2000-06-01	2000-07-21	2000-10-07	2000-12-13
2001	2001-04-07	2001-04-30	2001-06-23	2001-07-15	2001-09-26	2002-01-02
2002	2002-03-15	2002-04-24	2002-06-25	2002-07-22	2002-10-10	2003-01-04
2003	2003-04-03	2003-04-21	2003-06-17	2003-07-24	2003-10-07	2004-01-07
2004	2004-03-27	2004-04-20	2004-06-24	2004-07-21	2004-10-06	2004-12-25
2005	2005-04-05	2005-04-20	2005-06-22	2005-07-17	2005-10-21	2005-12-19
2006	2006-03-24	2006-03-24	2006-06-18	2006-07-21	2006-10-09	2007-01-20
2007	2007-03-27	2007-05-07	2007-06-06	2007-07-20	2007-10-28	2007-12-26
2008	2008-04-10	2008-04-21	2008-06-17	2008-07-22	2008-10-13	2008-12-16
2009	2009-03-23	2009-05-02	2009-06-25	2009-08-10	2009-10-03	2009-12-29
2010	2010-03-26	2010-05-14	2010-06-05	2010-07-05	2010-10-09	2010-12-15
2011	2011-04-03	2011-05-24	2011-06-21	2011-07-19	2011-10-11	2012-01-14
2012	2012-03-11	2012-04-10	2012-06-10	2012-07-03	2012-10-03	2013-01-14
2013	2013-04-11	2013-05-19	2013-06-18	2013-07-10	2013-10-22	2013-12-16
2014	2014-04-12	2014-05-29	2014-06-14	2014-07-14	2014-10-12	2015-01-02
2015	2015-04-03	2015-05-26	2015-06-10	2015-07-04	2015-10-14	2016-01-11
2016	2016-03-13	2016-05-03	2016-05-30	2016-07-04	2016-10-26	2017-01-01
2017	2017-03-07	2017-05-19	2017-06-13	2017-07-08	2017-10-21	2017-12-27
2018	2018-03-31	2018-05-19	2018-06-02	2018-07-04	2018-10-14	2018-12-15
2019	2019-03-31	2019-05-20	2019-07-03	2019-07-17	2019-10-10	2020-01-12

Table S2. Cyanotoxin concentrations. The measured microcystin and anabaenopeptin concentrations in nM.

650   *Table is larger than a single page*

Table S3. Zooplankton grouping for diversity calculation. Zooplankton were grouped by the finest taxonomic level that they could be confidently identified to before calculating the Shannon diversity. This table lists the taxonomic levels used to group zooplankton.

LTER Species Code	LTER Name	Group
50201	ACROPERUS HARPAE	Chydoridae Acroperus harpae
50300	ALONA	Chydoridae Alona
51000	CHYDORUS	Chydoridae Chydorus
51103	DAPHNIA MENDOTAE	Daphnia mendotae
51105	DAPHNIA PARVULA	Daphnia parvula
51130	DAPHNIA PULICARIA	Daphnia pulicaria
51107	DAPHNIA RETROCURVA	Daphnia retrocurva
51100	DAPHNIA	Daphnia unclassified
53701	BYTHOTREPHES LONGIMANUS	Bythotrephes longimanus
53601	SINOBOSMINA FRYEI	Bosmina sinobosmina fryei
51202	DIAPHANOSOMA BIRGEI	Diaphanosoma
52301	LEPTODORA KINDTI	Leptodora
40000	HARPACTICOID	Harpacticoida
30000	CALANOID	Calanoida unclassified
30901	AGLAODIAPTOMUS CLAVIPES	Calanoida Diaptomidae
30400	DIAPTOMID	Calanoida Diaptomidae
20100	ACANTHOCYCLOPS	Acanthocyclops
20302	DIACYCLOPS THOMASI	Diacyclops thomasi
20601	MESOCYCLOPS EDAX	Mesocyclops edax
20901	TROPOCYCLOPS PRASINUS MEXICANUS	Tropocyclops
20900	TROPOCYCLOPS	Tropocyclops
20000	CYCLOPOID	Cyclopoida uncalssified
10000	COPEPOD NAUPLII	Copepoda juvenile
15000	COPEPODITES	Copepoda juvenile
70100	CHAOBORUS	Chaoborus

Table S4. Differentially abundant OTUs. The effect sizes of all differentially abundant OTUs are

660   listed, along with the OTUs' taxonomic assignments.

Table is larger than a single page

Fig S1. Phytoplankton biovolume. The biovolume of *Cyanobacteria* and eukaryotic
phytoplankton in each season, broken down by invasion status.

Fig S2. NMDS plots of invasion impacts by season. Panels correspond to different seasons,
675   points represent sample dates included in each season, and colors indicate the sample date's
invasion status. NMDS was performed on CLR-transformed data.