Species invasions shift microbial phenology in a two-decade

freshwater time series.

Robin R. Rohwer[1*], Riley J. Hale[2], M. Jake Vander Zanden[3], Todd R. Miller[4], Katherine D.

McMahon[2,5]

[1]Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA.
[2]Department of Civil and Environmental Engineering, University of Wisconsin-Madison;
Madison, WI, USA.
[3]Center for Limnology, University of Wisconsin-Madison; Madison, WI, USA
[4]Zilber School of Public Health, University of Wisconsin-Milwaukee; Milwaukee, WI, USA.
[5]Department of Bacteriology, University of Wisconsin-Madison; Madison, WI, USA.
*Robin R. Rohwer
Email: Robin.Rohwer@gmail.com
13

Author Contributions: Conceptualization: RRR, KDM**;** Funding acquisition: RRR, TRM, KDM**;**

Methodology: RRR, TRM, JVZ**;** Investigation: RRR, RJH**;** Data curation: RRR**;** Formal analysis: RRR**;**

Visualization: RRR**;** Writing – original draft: RRR**;** Writing – review & editing: RRR, RJH, JVZ, TRM,

KDM

Competing Interest Statement: Authors declare that they have no competing interests.

Keywords: Invasive Species, Microbial Ecology, Phenology, Limnology, Time Series

This PDF file includes:
Main Text
Figure Captions
Table Headings
25

Abstract

Invasive species impart abrupt changes on ecosystems, but their impacts on microbial
communities are often overlooked. We paired a 20-year freshwater microbial community time
series with zooplankton and phytoplankton counts, rich environmental data, and a 6-year
cyanotoxin time series. We observed strong microbial phenological patterns that were disrupted
by the invasions of spiny water flea (*Bythotrephes cederströmii*) and zebra mussels (*Dreissena*
polymorpha). First, we detected shifts in *Cyanobacteria* phenology. After the spiny water flea
invasion, *Cyanobacteria* dominance crept earlier into clearwater; and after the zebra mussel
invasion, *Cyanobacteria* abundance crept even earlier into the diatom-dominated spring. During
summer, the spiny water flea invasion sparked a cascade of shifting diversity where zooplankton
diversity decreased and *Cyanobacteria* diversity increased. Second, we detected shifts in
cyanotoxin phenology. After the zebra mussel invasion, microcystin increased in early summer
and the duration of toxin production increased by over a month. Third, we observed shifts in
heterotrophic bacteria phenology. The *Bacteroidota* phylum and members of the acI
Nanopelagicales lineage were differentially more abundant. The proportion of the bacterial
community that changed also differed by season; the spring and clearwater bacterial
communities changed most following the spiny water flea invasion that lessened clearwater
duration and intensity, while the diverse summer bacterial community changed least following
the zebra mussel invasion despite the observed shifts in diversity and toxicity during summer.
These long-term invasion-mediated shifts in microbial phenology demonstrate the
interconnectedness of microbes with the broader food web, and their susceptibility to long-
term environmental change.

Significance Statement

Microbial communities are typically studied as part of the microbial loop, separately from the
broader food web. Using a two-decade freshwater time series, we explored whether two
species invasions that shifted the metazoan food web (spiny water flea and zebra mussels) also
impacted the microbial communities. We looked for seasonal responses because the microbial
communities had strong seasonal patterns. We discovered that *Cyanobacteria* increased early in
the year, and *Cyanobacteria* diversity increased in the summer. Cyanotoxins also increased,
along with the duration of toxin production. In the heterotrophic bacterial community, some
organisms changed consistently within lineages and seasons while others diverged. These
findings illustrate the importance of seasonal context, and highlight the interconnectedness of
bacteria with the broader food web.
61

Main Text

Introduction

Invasive species have wide-ranging impacts on ecosystems, often rewiring food webs and disrupting nutrient cycling, and they can be even more disruptive than global abiotic change (1). However, the impact of species invasions on entire microbial communities are rarely considered. Microbes are usually considered separately from metazoan lake ecology; bacterial communities are thought to interact with the broader food web as part of a microbial loop, where phytoplankton-derived dissolved organic matter is returned to higher trophic levels (2, 3). Bacteria that consume organic matter are in turn consumed by ciliates and nanoflagellates, which are eaten by microzooplankton, thus linking microbial communities with higher trophic levels. Although this microbial food web is often studied separately from the metazoan food web, we hypothesized that metazoan species invasions would change the bacterial community.

We collected a 20-year microbial community time series (2000-2019, 496-samples) from Lake Mendota, a eutrophic temperate lake in WI, USA. This period includes two metazoan species invasions: the predatory zooplankton spiny water flea in 2009 (*Bythotrephes cederströmii*) and zebra mussels in 2015 (*Dreissena polymorpha*). The spiny water flea invasion triggered a food web cascade that resulted in a one meter loss in water clarity (4), as increased zooplankton predation by spiny water flea led to reduced grazing pressure on phytoplankton. This impact was particularly strong during the spring clearwater phase, which decreased in intensity and duration (5). The invasion of zebra mussels triggered a 300% increase in benthic zooplankton and phytoplankton abundance, and they themselves became abundant enough to filter the epilimnion volume every 45 days (6). Zebra mussels increase filtration, but they can also worsen *Cyanobacteria* blooms via selective retention of eukaryotic algae and release of Cyanobacteria back into the lake (7), and by altering nutrient stoichiometry (8). In Lake Mendota, zebra mussels did not impact the pelagic water clarity in either direction (6). These mid-time series disturbances set up a natural experiment with which we explored the impact of metazoan species invasions on the bacterial community.

Lake Mendota is a temperate, dimictic lake and its ecology follows consistent seasonal

patterns (phenology). The lake freezes in the winter, and ice-off is followed by spring mixing and

a diatom bloom. As the water warms, zooplankton grazing increases and results in a short-lived

period of high water clarity, which ends when planktivorous fish begin eating zooplankton (5).

After this clearwater phase, diatoms give way to *Cyanobacteria* which persist throughout the

summer (9). During the summer the water column is thermally stratified, and anoxia develops in

the bottom layer while nutrient draw-down occurs in the upper layer. When temperatures drop

in the fall, Lake Mendota mixes and remains mixed until ice-on. Seasonal patterns have also

been observed in microbial community composition (10), and these seasonal cycles can mask

long-term changes (11). Therefore, we split our time series into seasons and looked for changes

in bacterial phenology.

Here we identify seasons that described the bacterial phenology, and then identify

phenological change in *Cyanobacteria*, cyanotoxins, and heterotrophic bacteria. These observed

phenological shifts are a direct observation of multi-decadal microbial change in response to

invasive species.

Results

We characterized the microbial community composition via 16S rRNA gene amplicon

sequencing performed on duplicate water grabs (961 total samples, 496 sample dates) (Fig 1).

The water grabs integrated the top 12 m of the water column, which approximates the mixed

upper layer. In addition to the sequencing-based relative abundances, we obtained

environmental data available from the North Temperate Lakes Long Term Ecological Research

program (NTL-LTER), including water physical and chemical measurements and microscopy-

based counts of zooplankton and phytoplankton.

Microbial Phenology

To compare across years, we defined lake seasons using environmental variables and

compared samples within the same season. We compared across multiple years by season

instead of by Julian date to discriminate shifts in each season's community composition from

shifts in seasonal timing. We created multiple potential season definitions based on

environmental data, and then selected the definitions that were most representative of the

microbial phenology. We applied non-metric multidimensional scaling (NMDS) (12) to sample

distance matrices calculated with the phi distance metric for centered log-ratio (CLR)-

transformed relative abundances (13). To select the season definitions most relevant to the

microbial community, we chose definitions that resulted in clear separation between season

boundaries in NMDS plots (Fig 1). Samples grouped by our selected season definitions were

significantly less variable within groups than between groups for every season boundary

(ANOSIM < 0.001).

The season definitions that best described the microbial phenology were ice-on, spring,

clearwater, early summer, late summer, and fall (Fig 1, Table S1). The ice-on season was

determined by the onset and break-up of continuous ice-cover across a central lake transect.

Spring began with ice-off and ended with clearwater onset, spanning a period that includes the

spring diatom bloom (9). Clearwater was defined based on water clarity measurements, and is

an annual phase driven by high zooplankton grazing pressure (5). Early summer was a period of

growing stratification, bounded by clearwater and the point at which hypolimnetic anoxia

extended above 12 m. Late summer was a period of strong stratification and anoxia, which

ended with fall mixing. We defined mixing as a difference in average epilimnion and

hypolimnion temperatures < 3°C. Fall began with mixing and ended with ice-on.

Shifts in *Cyanobacteria* phenology

Given spiny water flea's marked disruption of clearwater phenology, *i.e.* shortened

duration and intensity, we first looked for changes in spring phytoplankton phenology.

Historically, the end of clearwater phase marks the transition from a diatom-dominated system

to a *Cyanobacteria*-dominated system (9). We found that after the spiny water flea invasion,

Cyanobacteria began appearing during clearwater phase (Fig 2A). We did not see a similar

increase in chloroplast relative abundance or diatom counts during clearwater phase, indicating

that the transition to *Cyanobacteria* dominance shifted earlier. We further found that after the

zebra mussel invasion, *Cyanobacteria* began appearing even before the start of clearwater

phase (Fig 2A). Count data confirmed that the biovolume of *Cyanobacteria* increased 2-fold,

growing to over 10% of total phytoplankton (Fig S1). This earlier onset of *Cyanobacteria*

describes a kingdom-level regime shift from a eukaryotic algae-dominated period to one that

includes *Cyanobacteria*.

After identifying these changes in the spring phenology, we next looked for spiny water

flea invasion impacts on the summer cyanobacterial phenology. *Cyanobacteria* abundance

(relative to the microbial community) during summer and fall did not change with the spiny

water flea invasion. However, *Cyanobacteria* diversity during the late summer season increased

(Fig 2B). This increase in cyanobacterial diversity was observed at fine and coarse-resolution

taxonomic levels, including levels as coarse as order (Fig 2B). It is possible that the shift in

summer diversity could be explained by the earlier onset of the *Cyanobacteria*-dominated

season resulting in an earlier onset of high *Cyanobacteria* diversity. However, the spring

cyanobacterial incursion was comprised of a single amplicon sequence assigned to

Aphanizomenon, not to a more complex community typical of early summer. Additionally, we

observed a concurrent late summer change in zooplankton diversity; after the spiny water flea

invasion, zooplankton diversity decreased (Fig 2B, Table S2). Therefore, we attribute the shift in

cyanobacterial summer diversity to a cascade of shifting diversity originating in the zooplankton

community. The lower diversity of zooplankton may have resulted in narrower grazing pressure,

allowing a wider range of *Cyanobacteria* to flourish.

Next, we looked for changes in the summer cyanobacterial community stemming from

the zebra mussel invasion. We first compared total *Cyanobacteria* abundance before and after

zebra mussels but did not see strong and consistent changes in relative abundance or biovolume

(Fig S1). We also did not observe strong shifts in cyanobacterial community composition.

Cyanobacteria can produce potent toxins, but *Cyanobacteria* abundance does not always

correlate with cyanotoxin concentrations (14, 15), suggesting that toxin concentrations could

yet have changed as a result of the invasion.

Shifts in cyanotoxin phenology

To examine the impact of zebra mussels on cyanobacterial toxicity, we collected a 6-

year, 246-sample toxin time series spanning the 3 years preceding and following the zebra

mussel invasion. We measured the hepatotoxin microcystin and the bioactive peptide

anabaenopeptin using high performance liquid chromatography coupled with tandem mass

spectrometry (HPLC-MS/MS) (Table S3) (16). We observed a 3-fold decrease in anabaenopeptin

and a concurrent 3-fold increase in microcystin during early summer ($p < 0.05$) (Fig 3A). The

family *Microcystaceae* is considered the main microcystin producer in Lake Mendota, although it

is also capable of producing anabaenopeptin (14). However, the relative abundance of

Microcystaceae did not change. This suggests that the switch from anabaenopeptin to

microcystin is due to more complex ecological drivers than the abundance of *Microcystaceae*,

and illustrates the complexity of invasion impacts. This shift in toxin concentrations has

implications for human health, because microcystin is a more potent toxin than anabaenopeptin

(17).

After observing this increase in early summer microcystin, we next explored the

phenology of microcystin production (Fig 3B). Total toxin concentrations were variable between

years, so for each year we normalized total microcystin by the maximum measured

concentration within that year. We found that the duration of toxin production, defined as the

period during which toxins were at least 15% of maximum annual toxicity, increased after the

zebra mussel invasion by 53 ± 9 days, or 170 ± 10%. ($p < 0.001$) (Fig 3C). This increase was due to

a longer late summer season, as well as toxin production extending further into the early

summer and fall seasons. We observed these increases in toxicity despite the lack of change in

the *Cyanobacteria* abundance and diversity. This might be explained by changes in gene

expression, which can be variable enough that the abundance of microcystin *mcy* genes is a

poor predictor of microcystin concentrations (14). Another possible explanation is preferential

retention of non-toxic strains or individual cells, with toxic cells released back into the water

column. Zebra mussels were previously found to preferentially retain a non-toxic *Microcystis*

strain over a toxic one (18).

In summary, cyanobacterial phenology shifted in response to the two metazoan

invasions, with changes in timing, diversity, and toxicity.

Shifts in heterotrophic bacteria phenology

To explore whether invasion impacts extended further, into the heterotrophic bacterial

community, we compared each season's heterotrophic community before and after invasion

using NMDS analyses of their CLR-transformed phi dissimilarities (13). We observed modest

grouping based on invasion status (Fig S2), and an ANOSIM analysis confirmed that within-group

differences were smaller than between-group differences in all seasons (ANOSIM significance =

0.001).

To identify which bacteria were responsible for the shift in community composition, we

performed a differential abundance analysis after applying a CLR transformation to account for

the relativized nature of sequencing data (19). We used a generalized linear model to test for

organisms that were differentially abundant before and after each invasion (20), and we looked

for changes within each season. We considered an operational taxonomic unit (OTU)

differentially abundant when it had an effect size > 0.5, and we manually confirmed the

ecological significance of this effect size cutoff by inspecting relative abundance over time

(examples in Fig 4).

To assess which seasonal communities were most susceptible to change, we looked at

the proportion of OTUs within the community that were differentially abundant. To better

reflect the ecological impact of the differentially abundant OTUs, we normalized the OTUs by

their mean relative abundance. The invasions impacted seasonal community composition to

different degrees. For example, spiny water flea had the largest impact on the proportion of the

community that changed during spring and clearwater, with 38% of the average community

becoming differentially abundant (Fig 4C, left). This means that in addition to changing the

seasonal timing of clearwater (5), spiny water flea also changed the heterotrophic bacteria

community. Meanwhile, zebra mussels had the smallest impact on the late summer microbial

community, with only 4% of the average community becoming differentially abundant (Fig 4C,

right). The late summer microbial community had the smallest proportion of differentially

abundant organisms, and also the fewest overall differentially abundant OTUs (Fig 4). The late

summer community also has the highest alpha diversity, and high biodiversity has been linked to

ecosystem stability (21). It is possible that this high diversity makes the late summer ecosystem

more robust. The seasonal differences in invasion impacts emphasize the importance of a

phenological context.

Among the organisms that were differentially abundant, OTUs in the phylum

Bacteroidota were zebra mussel invasion "winners" broadly (Fig 4A, Table S4). These winners

included abundant OTUs from the bacI, bacII, bacIII, and bacIV lineages (*Chitinophagales*,

Flavobacteriales, and *Cytophagales* orders, lineage nomenclature from (22, 23)). However, most

changes in differential abundance were not consistent within taxonomic lineages or across

seasons. For example, within the abundant lineages of order *Frankiales*, *Nanopelagicaceae* (acI)

tended to win while acTH1 and acSTL tended to lose (Fig 4B, Table S4).

The *Bacteroidota* phylum is often particle-associated (23, 24), and *Bacteroidota* OTUs

were broadly differentially abundant after both the spiny water flea invasion and the zebra

mussel invasion. In contrast, the *Frankiales* order is comprised of small, free-living bacteria and

contained both winner and loser OTUs. The non-uniform response of the *Frankiales* order

suggests that they occupy divergent niche spaces. Further, it appears that the niche spaces for

acI may be seasonally dependent. The acI lineage are oligotrophs (25), and acI-A and acI-B

tended to win during clearwater and early summer even though they are abundant throughout

the year, while acI-C is abundant during fall and also tended to win during fall. AcTH1 and acSTL

are also small, free-living *Frankiales*, but little is known about their ecology and metabolism.

They comprised many of the abundant invasion losers in all seasons after both invasions, which

suggests that their niche is less seasonally dependent and different from their close relatives'.

In summary, heterotrophic bacteria phenology was shifted by metazoan species

invasions. Some organisms showed consistent responses within phyla and across seasons, and

some organisms showed divergent responses within orders and between seasons, suggesting

that some niche spaces were changing broadly while others experienced seasonally dependent

changes. Overall, the amount of change bacterial communities experienced was seasonal, and

some seasonally-defined communities were more robust to change than others.

Species invasions are drivers of change

To compare the importance of species invasions to other environmental drivers, we

evaluated metrics related to invasion status, landscape change, and climate change in a

modeling framework. We predicted the lag between *Cyanobacteria* onset and clearwater (Fig

2A) and the CLR-transformed abundance of the winner acI-B OTU 1 (Fig 4B) using invasion

status, phosphorus loading, ice duration, and hot summer days. We used Akaike Information

Criterion (AIC) to identify which drivers were included in the best-performing models (Table S5

and S6). Spiny water flea and zebra mussel invasion status were included in all the best models,

which emphasizes the importance of species invasions as drivers of changing microbial

phenology. The number of hot summer days and mean phosphorus loading was also included in

some of the best-performing models. This suggests that even though species invasions were a

strong driver of microbial change during the 19-year time frame of our dataset, climate and

landscape are interacting drivers of microbial change.

Discussion

Over century-long time frames, the interacting driver of climate change is shifting the

phenology of northern hemisphere lakes. Winters are shortening (26), summer stratification and

anoxia is increasing (27), and extreme weather is becoming more common (28). In Lake

Mendota specifically, ice duration (29) and clearwater (5) are shortening, while the stratified

(29) and anoxic (30) periods are lengthening. We posit that lake microbial communities will also

change in response to climate change because the lake microbial community has phenological

patterns that track these shifting seasons. Further, we directly observed that food web-

mediated shifts in seasonal timing can change microbial community timing, composition, and

toxicity. It follows that the interacting driver of climate change shifting seasons over longer time

frames will likely shift microbial phenology too.

Moreover, the spiny water flea invasion itself was driven by climate change. Previous

work on Lake Mendota found that an unusually cool year allowed a sleeper population of spiny

water flea to irrupt (31), and weather extremes like this are expected to increase with climate

change (28). It is possible that the spiny water flea-mediated food web shift also paved the way

for the zebra mussel invasion. Zebra mussels likely had multiple introductions to Lake Mendota

based on the large amount of recreational boat traffic on the lake, so it is plausible that they

were introduced prior to 2015 and existed as a low abundance sleeper population. The

establishment of one invasive species can destabilize an ecosystem in a way that primes it for

additional invasions (32), or that triggers the population irruption of existing sleeper populations

(33). A climate anomaly triggered a population irruption of the invasive spiny water flea sleeper

population, and the resulting food web disruptions may have paved the way for the

establishment of zebra mussels. The impact of these invasions reached all the way into the

bacterial community, illustrating how the interacting drivers of species invasions and climate

change can have complex and far-reaching impacts.

Harmful cyanobacterial blooms are predicted to increase with warmer temperatures,

higher CO_2, and increasingly intense rain events (34). We observed an earlier onset of

Cyanobacteria into the clearwater and spring seasons. In other lakes, regime changes have been

observed as oligotrophic systems switch to a new stable state dominated by *Cyanobacteria* (35).

It is possible that we may be observing the beginning of a transition to a new stable state at a

seasonal level. In the summer seasons already dominated by *Cyanobacteria*, we found the

fewest differentially abundant organisms. This suggests that even if tipping points are reached in

other seasons, the community is less likely to shift when it would be most desired: during the

summer months plagued by toxic cyanobacterial blooms.

Long-term change is difficult to distinguish from short-term variability and seasonal

cycles, a phenomenon termed the "invisible present" (11). Observations of the microbial

community for almost twenty years have allowed us to distinguish consistent phenological

patterns, and to identify separate long-term, invasion-mediated shifts. The establishment and

continuation of time series like ours are the only way to directly observe the long-term impacts

of environmental change. Our study found that microbial communities are sensitive to long-

term change, specifically to the cascading impacts of invasive species. These impacts included

many aspects of phenology, from timing and abundance, to diversity and community

composition, to human health and cyanobacterial toxicity.

Materials and Methods

Bacterial Community Data

Microbial sampling, sequencing, and data processing was performed as described in

detail in Rohwer & McMahon (36). Briefly, integrated epilimnion samples were collected from

the central deep hole of Lake Mendota (WI, USA) using a 12 m tube, and water was well mixed

before filtering onto a 0.22 um filter (Pall Corporation). Filters were stored in a -80°C freezer

until DNA extraction in 2018-2019. DNA was extracted using a FastDNA Spin Kit (MP

Biomedicals) after sample order was randomized. Samples were sent to Argonne National lab

for amplification of the 16S rRNA gene V4-V5 region (515F-Y GTGYCAGCMGCCGCGGTAA, 926R

CCGYCAATTYMTTTRAGTTT) and sequencing on an Illumina MiSeq instrument.

After sequencing, data was processed using custom R scripts available at

https://github.com/rrohwer/limony and described in Rohwer & McMahon (36). Microbial relative

abundances from replicate samples were averaged, and a standard deviation was calculated.

This processed data is available as part of an R package at https://github.com/rrohwer/limony. The

raw sequencing data is available through the NCBI Sequence Read Archive under accession

number PRJNA846788.

Environmental Data

We collected lake physical and chemical data alongside microbial samples, including

water temperature and dissolved oxygen measured with a YSI ProPlus multiparameter sonde

(Yellow Springs Instruments), and water clarity measured with a Secchi disk. This data is

available through the Environmental Data Initiative (EDI) portal under identifiers knb-lter-

ntl.415.2 (37) and knb-lter-ntl.416.1 (38). We combined our measurements with additional

water temperature, dissolved oxygen, water clarity, and ice cover datasets made available by

the North Temperate Lakes Long-Term Ecological Research program (NTL-LTER) under EDI

identifiers knb-lter-ntl.29.29 (39), knb-lter-ntl.130.29 (40), knb-lter-ntl.335.1 (41), knb-lter-

ntl.400.2 (42), knb-lter-ntl.31.30 (43), and knb-lter-ntl.33.35 (44). We obtained mean annual

phosphorus loading at the Yahara River Windsor site from the USGS surface water annual

statistics (45).

We defined lake seasons using metrics calculated from this environmental data. We

identified the epilimnion, metalimnion, and hypolimnion depths by identifying the inflection

points of temperature profile vectors, calculated the average epilimnion and hypolimnion

temperatures for all complete profiles, and smoothed these values with a 7-day moving

average. We defined mixing as the day when the linearly interpolated difference in epilimnion

and hypolimnion temperature fell below 3°C. We considered depths with < 1 mg/L dissolved

oxygen anoxic, and we applied a 7-day moving average to the depth of the anoxic layer. We

defined the start of the late summer season as the day when the linearly interpolated anoxic

layer depth reached 12 m. We defined the clearwater season's start and end for each year by

manually identifying the edges of the abrupt drop in Secchi depth measurements that occurs

after ice-off, and we refined the edge dates based on chloroplast and *Cyanobacteria* abundance

when sequencing data existed without a paired Secchi depth measurement.

We obtained absolute abundance and biovolume of large *Cyanobacteria* and eukaryotic

algae using microscopy-based phytoplankton counts from the NTL-LTER, and we used

microscopy-based zooplankton counts to calculate zooplankton diversity. This data is available

under EDI identifiers knb-lter-ntl.88.30 (46) and knb.lter.ntl.90.31 (47).

Cyanotoxin Data

We measured toxins in 246 whole-water samples spanning 6 years (2013-2018). These

samples were taken from the same water grabs used to characterize the microbial community.

We quantified cyanotoxins using methods described in detail by Miller *et al.* (16). Briefly,

lyophilized samples were resuspended in formic acid, subjected to three freeze-thaw cycles, and

extracted in methanol and formic acid in a sonicating water bath. The extract supernatant was

analyzed by targeted high performance liquid chromatography coupled with tandem mass

spectrometry (HPLC-MS/MS).

Eleven microcystin (MC) congeners and three anabaenopeptin (Apt) congeners were

analyzed. Certified reference standards of MC-LR, [Dha^7]MC-LR and nodularin were purchased

from the National Research Council of Canada Biotoxins program (Halifax, Nova Scotia). MC-LA,

MC-RR, MC-LF, MC-YR, MC-WR, MC-LY, MC-LW, MC-HtyR, MC-HilR (all >95%) were purchased

from Enzo Life Sciences (Farmingdale, NY). Anabaenopeptin A (Apt-A) (> 95%), Apt-B (>95%),

and Apt-F (> 95%) were purchased from MARBIONC (Wilmington, NC). To calculate total

concentrations, we summed nanomolar measurements of the congeners. 99 ± 4% of total

microcystin was comprised of the congeners MC-LA and MC-LR. Toxin totals are available in a

tabular format as Table S2.

Analysis

We defined microbially-relevant seasons (Fig 1A) based on the environmental data. Ice

extent, mixing, and clearwater were phenological markers that corresponded well with the

microbial community. To divide the summer months we explored multiple seasonal cutoffs

based on stratification extent, epilimnion temperature, and anoxia extent. We visualized each

season option with non-metric multidimensional scaling (NMDS) analysis and manually chose

the most relevant definitions based on visual inspection of the NMDS plots. To account for the

proportional nature of sequencing data, we performed a centered log ratio (CLR) transformation

on our data before the NMDS analysis (Fig 1B, Fig S2). Following the guidance provided by Quinn

et al. (19), we filtered each 3-season group to contain only OTUs that occurred in at least 10% of

the samples, assigned small-values to zeros using the zCompositions R package (48), and

performed a CLR-transform and calculated the *phi* distance metric using the propr R package

(13). We performed NMDS and confirmed the significance of our season choices by calculating

the ANOSIM significance using the vegan R package (12). The season start dates are listed in

Table S1. We used season start dates to align samples in our subsequent analyses; instead of

aligning samples in different years by Julian date, we aligned samples by the days since the start

of a given season. Comparisons across Julian dates do not account for shifts in seasonal timing

across years, but aligning samples by season start dates enables comparison of phenologically

similar microbial communities.

We observed changes in spring *Cyanobacteria* relative abundance (Fig 2A) by aligning

samples by the start of clearwater phase and grouping years into invasion periods. Both

invasions were discovered in fall, so we considered the subsequent years the start of the

invasion period. The seven 2019 samples were excluded from this analysis because they did not

span the entire spring-clearwater-early summer period. We compared the summer diversity

before and after the spiny water flea invasion (Fig 2B) by aligning samples by the start of the

early summer season and calculating the Shannon diversity index using the vegan R package

(12). We observed no significant difference in diversity between the spiny water flea-only period

and the zebra mussel period, so we grouped these two periods together as a single post-spiny

water flea group. Before calculating *Cyanobacteria* diversity, we excluded samples from 2000-

2003 that had been prefiltered with a 10 µm filter instead of filtered directly onto a 0.22 µm

filter. Before calculating zooplankton community diversity (Fig 2B), we excluded spiny water flea

and juvenile copepods from the zooplankton count data and grouped zooplankton based on

ecological relevance and our confidence in the identifications (Table S3). We calculated the

mean and standard deviation of linearly interpolated daily diversity values and performed a t-

test to compare between invasion groups.

We calculated the concentration of toxins before and after the zebra mussel invasion

(Fig 3A) by separating toxin measurement dates by season and splitting the years into invasion

groups. For each season and year group, we calculated the mean and standard deviation of all

toxin measurements within it. We performed a t-test on each seasonal block to identify

significant change between invasion groups. We calculated the duration of toxin production (Fig

3B-C) by normalizing microcystin concentrations in each year by the yearly maximum

concentration. We applied a linear interpolation to the 7-day moving average of measurements

(lines in Fig 3B). We defined days as toxin-producing if their interpolated daily toxin

concentration was > 15 % of the yearly maximum value.

We identified differentially abundant bacteria (Fig 4) using the ALDEx2 package (49)

following the guidance provided by Quinn *et al.* (19). For each season group, we removed OTUs

present in fewer than 10% of samples and added small numbers to remaining zeros using

zCompositions (48). We identified differentially abundant organisms and their effect sizes using

a generalized linear model with the aldex.glm and aldex.glm.effect functions from the ALDEx2

package (49). We manually chose an effect size cutoff of 0.5 by examining plots of OTU

abundance over time for OTUs with varying effect sizes (examples in Fig 4A and Fig 4B). We

identified which seasons were most susceptible to microbial community change (Fig 4C) by

calculating the proportion of the community that was differentially abundant. We defined this

proportion as the average abundance of all differentially abundant OTUs divided by the average

abundance of all OTUs. All differentially abundant bacteria are listed in Table S4.

We created generalized linear models to predict microbial phenology from invasion

status, Yahara River phosphorus loading into the lake, ice duration, and hot summer days (Table

S4-5). We used the glm function in R to create models, and the dredge function in the MuMIn R

package to compute the corrected AIC for each combination of predictors. For a *Cyanobacteria*

phenology response variable, we modeled the lag between clearwater season start and the first

day with *Cyanobacteria* abundance ≥ 3%. For a heterotroph phenology response variable, we

modeled the average CLR-transformed clearwater abundance of OTU 1, a *Frankiales* acI-B that

was a spiny water flear and zebra mussel clearwater winner and is also the most abundant OTU

in the dataset. As predictor variables, we used the previous-winter ice duration as a climate

metric (44), the mean annual Yahara River phosphorus loading as a landscape metric (45), and

the number of days with an average epilimnion temperature ≥ 23°C as a climate metric related

to the initial spiny water flea invasion (31).

Acknowledgments

We thank Stephen Carpenter and Edward DeLong for friendly reviews. We thank

technicians Ame Xiong and Leah Stromberg for assistance measuring cyanotoxins. We thank

Jake Walsh for assistance interpreting the zooplankton data. We thank the many field personnel

involved in microbial sampling, including sampling leads Angela Kent, Tony Yannarell, Ashley

Shade, Stuart Jones, Ryan Newton, Georgia Wolfe, Emily Kara Read, Lucas Beversdorf, and

James Mutschler; and initial Microbial Observatory lead Eric W. Triplett.

We thank the funding sources that made this work possible:

U.S. National Science Foundation Postdoctoral Research Fellowship in Biology, award number

2011002 (RRR)

E. Michael and Winona Foster WARF Wisconsin Idea Fellowship, 2018 (RRR)

National Institute of Food and Agriculture, U.S. Department of Agriculture, award number 2016-

67012-24709 (Joshua J. Hamilton)

National Institute of Food and Agriculture, U.S. Department of Agriculture, Hatch Projects

WIS01516, WIS01789, WIS03004 (KDM)

U.S. National Science Foundation North Temperate Lakes Long-Term Ecological Research site,

NTL-LTER, award numbers DEB-9632853, DEB-0217533 (Stephen Carpenter) and DEB-0822700,

DEB-1440297 (Emily Stanley)

U.S. National Science Foundation Microbial Observatories program, award numbers MCB-

9977903 (Eric Triplett) and DEB-0702395 (KDM)

U.S. National Science Foundation INSPIRE award, DEB-1344254 (KDM)

The Laboratory for Aquatic Environmental Microbiology & Chemistry at the University of

Wisconsin-Milwaukee, Zilber School of Public Health (TRM)

References

1.   B. E. Lopez, *et al.*, Global environmental changes more frequently offset than intensify
469   detrimental effects of biological invasions. *Proceedings of the National Academy of*
470   *Sciences* **119**, e2117389119 (2022).

471   2.   F. Azam, *et al.*, The Ecological Role of Water-Column Microbes in the Sea. *Marine Ecology*
472   *Progress Series* **10**, 257–263 (1983).

473   3.   K. Piwosz, I. Mukherjee, M. M. Salcher, V. Grujčić, K. Šimek, CARD-FISH in the Sequencing
474   Era: Opening a New Universe of Protistan Ecology. *Frontiers in Microbiology* **12** (2021).

475   4.   J. R. Walsh, S. R. Carpenter, M. J. V. Zanden, Invasive species triggers a massive loss of
476   ecosystem services through a trophic cascade. *PNAS* **113**, 4081–4085 (2016).

477   5.   S.-I. S. Matsuzaki, *et al.*, Climate and food web effects on the spring clear-water phase in
478   two north-temperate eutrophic lakes. *Limnol. Oceanogr.* **n/a**, 1–17 (2020).

479   6.   M. J. Spear, *et al.*, Early changes in the benthic community of a eutrophic lake following
480   zebra mussel (Dreissena polymorpha) invasion. *Inland Waters* **0**, 1–19 (2022).

481   7.   R. Naddafi, K. Pettersson, P. Eklöv, The effect of seasonal variation in selective feeding by
482   zebra mussels (Dreissena polymorpha) on phytoplankton community composition.
483   *Freshwater Biology* **52**, 823–842 (2007).

484   8.   O. Bykova, A. Laursen, V. Bostan, J. Bautista, L. McCarthy, Do zebra mussels (Dreissena
485   polymorpha) alter lake water chemistry in a way that favours Microcystis growth? *Science*
486   *of The Total Environment* **371**, 362–372 (2006).

487   9.   C. C. Carey, P. C. Hanson, R. C. Lathrop, S. Amand, A. L, Using wavelet analyses to examine
488   variability in phytoplankton seasonal succession and annual periodicity. *J Plankton Res* **38**,
489   27–40 (2016).

10. A. Shade, *et al.*, Interannual dynamics and phenology of bacterial communities in a
eutrophic lake. *Limnol. Oceanogr.* **52**, 487–494 (2007).

11. J. J. Magnuson, Long-Term Ecological Research and the Invisible Present. *BioScience* **40**,
495–501 (1990).

12. J. Oksanen, *et al.*, vegan: Community Ecology Package (2020).

13. T. P. Quinn, M. F. Richardson, D. Lovell, T. M. Crowley, propr: An R-package for Identifying
Proportionally Abundant Features Using Compositional Data Analysis. *Scientific Reports* **7**,
16252 (2017).

14. L. J. Beversdorf, S. D. Chaston, T. R. Miller, K. D. McMahon, Microcystin mcyA and mcyE
Gene Abundances Are Not Appropriate Indicators of Microcystin Concentrations in Lakes.
PLOS ONE **10**, e0125353 (2015).

15. Robert E. Zuellig, Jennifer L. Graham, Erin A. Stelzer, Keith A. Loftin, Barry H. Rosen,
"Cyanobacteria, Cyanotoxin Synthetase Gene, and Cyanotoxin Occurrence Among Selected
Large River Sites of the Conterminous United States, 2017–18" (U.S. Department of the
Interior, U.S. Geological Survey, 2021).

16. T. R. Miller, S. L. Bartlett, C. A. Weirich, J. Hernandez, Automated Subdaily Sampling of
Cyanobacterial Toxins on a Buoy Reveals New Temporal Patterns in Toxin Dynamics.
Environ. Sci. Technol. **53**, 5661–5670 (2019).

17. T. R. Miller, L. J. Beversdorf, C. A. Weirich, S. L. Bartlett, Cyanobacterial Toxins of the
Laurentian Great Lakes, Their Toxicological Effects, and Numerical Limits in Drinking
Water. *Marine Drugs* **15**, 160 (2017).

18. G. Juhel, *et al.*, Pseudodiarrhoea in zebra mussels Dreissena polymorpha (Pallas)exposed
to microcystins. *Journal of Experimental Biology* **209**, 810–816 (2006).

19.   T. P. Quinn, *et al.*, A field guide for the compositional analysis of any-omics data.
GigaScience **8**, giz107 (2019).

20.   A. D. Fernandes, *et al.*, Unifying the analysis of high-throughput sequencing datasets:
characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by
compositional data analysis. *Microbiome* **2**, 15 (2014).

21.   T. H. Oliver, *et al.*, Biodiversity and Resilience of Ecosystem Functions. *Trends in Ecology &*
Evolution **30**, 673–684 (2015).

22.   R. R. Rohwer, J. J. Hamilton, R. J. Newton, K. D. McMahon, TaxAss: Leveraging a Custom
Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *mSphere* **3**, e00327-18
(2018).

23.   R. J. Newton, S. E. Jones, A. Eiler, K. D. McMahon, S. Bertilsson, A Guide to the Natural
History of Freshwater Lake Bacteria. *Microbiology and Molecular Biology Reviews* **75**, 14–
49 (2011).

24.   M. W. Hall, R. R. Rohwer, J. Perrie, K. D. McMahon, R. G. Beiko, Ananke: temporal
clustering reveals ecological dynamics of microbial communities. *PeerJ* **5**, e3812 (2017).

25.   R. J. Newton, S. L. McLellan, A unique assemblage of cosmopolitan freshwater bacteria and
higher community diversity differentiate an urbanized estuary from oligotrophic Lake
Michigan. *Frontiers in Microbiology* **6**, 1028 (2015).

26.   S. Sharma, *et al.*, Loss of Ice Cover, Shifting Phenology, and More Extreme Events in
Northern Hemisphere Lakes. *Journal of Geophysical Research: Biogeosciences* **126**,
e2021JG006348 (2021).

27.   R. I. Woolway, *et al.*, Phenological shifts in lake stratification under climate change. *Nat*
Commun **12**, 2318 (2021).

28.   T. D. Loecke, *et al.*, Weather whiplash in agricultural regions drives deterioration of water
537   quality. *Biogeochemistry* **133**, 7–15 (2017).

538   29.   M. R. Magee, C. H. Wu, D. M. Robertson, R. C. Lathrop, D. P. Hamilton, Trends and abrupt
539   changes in 104 years of ice cover and water temperature in a dimictic lake in response to
540   air temperature, wind speed, and water clarity drivers. *Hydrology and Earth System*
541   *Sciences* **20**, 1681–1702 (2016).

542   30.   R. Ladwig, *et al.*, Lake thermal structure drives interannual variability in summer anoxia
543   dynamics in a eutrophic lake over 37 years. *Hydrology and Earth System Sciences* **25**,
544   1009–1032 (2021).

545   31.   J. R. Walsh, S. E. Munoz, M. J. Vander Zanden, Outbreak of an undetected invasive species
546   triggered by a climate anomaly. *Ecosphere* **7**, n/a-n/a (2016).

547   32.   D. Simberloff, B. Von Holle, Positive Interactions of Nonindigenous Species: Invasional
548   Meltdown? *Biological Invasions* **1**, 21–32 (1999).

549   33.   M. J. Spear, J. R. Walsh, A. Ricciardi, M. J. V. Zanden, The Invasion Ecology of Sleeper
550   Populations: Prevalence, Persistence, and Abrupt Shifts. *BioScience* (2021)
551   https:/doi.org/10.1093/biosci/biaa168 (March 26, 2021).

552   34.   J. Huisman, *et al.*, Cyanobacterial blooms. *Nat Rev Microbiol* **16**, 471–483 (2018).

553   35.   M. Scheffer, S. Carpenter, J. A. Foley, C. Folke, B. Walker, Catastrophic shifts in ecosystems.
554   *Nature* **413**, 591–596 (2001).

555   36.   R. R. Rohwer, K. D. McMahon, Lake iTag measurements over nineteen years, introducing
556   the limony dataset. *bioRxiv*, 2022.08.04.502869 (2022).

557   37.   R. R. Rohwer, K. D. McMahon, Lake Mendota Microbial Observatory Temperature,
558   Dissolved Oxygen, pH, and conductivity data, 2006-present. (2022)

https:/doi.org/10.6073/PASTA/7E533C197ED8EBD27777A89A2C8D7DFE (October 17,
2022).

38.   R. R. Rohwer, K. D. McMahon, Lake Mendota Microbial Observatory Secchi Disk
Measurements 2012-present (2022)
https:/doi.org/10.6073/PASTA/3B650E19D28CBC7B9ED631F0A7878033 (October 17,
2022).

39.   J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Physical
Limnology of Primary Study Lakes 1981 - current (2021)
https:/doi.org/10.6073/PASTA/316203040EA1B8ECE89673985AB431B7 (October 17,
2022).

40.   J. Magnuson, S. Carpenter, E. Stanley, North Temperate Lakes LTER: High Frequency Water
Temperature Data - Lake Mendota Buoy 2006 - current (2020)
https:/doi.org/10.6073/PASTA/8CEFF296AD68FA8DA6787076E0A5D992 (October 17,
2022).

41.   D. Robertson, Lake Mendota water temperature secchi depth snow depth ice thickness
and meterological conditions 1894 - 2007 (2016)
https:/doi.org/10.6073/PASTA/F20F9A644BD12E4B80CB288F1812C935 (October 17,
2022).

42.   J. J. Magnuson, S. R. Carpenter, E. H. Stanley, Lake Mendota Multiparameter Sonde
Profiles: 2017 - current (2021)
https:/doi.org/10.6073/PASTA/5F15BF453851987FC030B2F07A110B21 (October 17,
2022).

43.   J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Secchi Disk
Depth; Other Auxiliary Base Crew Sample Data 1981 - current (2021)
https:/doi.org/10.6073/PASTA/26FA98B39F9758FDA2109021F5B88076 (October 17,
2022).

44.   J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Ice Duration -
Madison Lakes Area 1853 - current (2021)
https:/doi.org/10.6073/PASTA/AB31F2489EE436BEB73FC8F1D0213D97 (October 17,
2022).

45.   U.S. Geological Survey, National Water Information System data available on the World
Wide Web (USGS Water Data for the Nation) (2022)
https:/doi.org/https://waterdata.usgs.gov/nwis/annual?site_no=05427718&agency_cd=U
SGS&por_05427718_220293=1353769,91050,220293,1990,2021&year_type=W&referred
_module=sw&format=rdb (December 12, 2022).

46.   J. J. Magnuson, S. R. Carpenter, E. H. Stanley, North Temperate Lakes LTER: Phytoplankton
- Madison Lakes Area 1995 - current (2022)
https:/doi.org/10.6073/PASTA/43D3D401AF88CC05C6595962BDB1AB5C (October 17,
2022).

47.   J. Magnuson, S. Carpenter, E. Stanley, North Temperate Lakes LTER: Zooplankton -
Madison Lakes Area 1997 - current (2019)
https:/doi.org/10.6073/PASTA/8B265C0300252C87805F26F41E174AA4 (October 17,
2022).

48.   J. Palarea-Albaladejo, J. A. Martín-Fernández, zCompositions — R package for multivariate
imputation of left-censored data under a compositional approach. *Chemometrics and*
Intelligent Laboratory Systems **143**, 85–96 (2015).

49.   A. D. Fernandes, J. M. Macklaim, T. G. Linn, G. Reid, G. B. Gloor, ANOVA-Like Differential
Expression (ALDEx) Analysis for Mixed Population RNA-Seq. *PLOS ONE* **8**, e67019 (2013).

Species invasions shift microbial phenology in a two-decade freshwater time series.

Robin R. Rohwer[1*], Riley J. Hale[2], M. Jake Vander Zanden[3], Todd R. Miller[4], Katherine D. McMahon[2,5]

[1]Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA.
[2]Department of Civil and Environmental Engineering, University of Wisconsin-Madison; Madison, WI, USA.
[3]Center for Limnology, University of Wisconsin-Madison; Madison, WI, USA
[4]Zilber School of Public Health, University of Wisconsin-Milwaukee; Milwaukee, WI, USA.
[5]Department of Bacteriology, University of Wisconsin-Madison; Madison, WI, USA.
*Robin R. Rohwer
Email: Robin.Rohwer@gmail.com

Author Contributions: Conceptualization: RRR, KDM**;** Funding acquisition: RRR, TRM, KDM**;**

Methodology: RRR, TRM, JVZ**;** Investigation: RRR, RJH**;** Data curation: RRR**;** Formal analysis: RRR**;**

Visualization: RRR**;** Writing – original draft: RRR**;** Writing – review & editing: RRR, RJH, JVZ, TRM, KDM

Competing Interest Statement: Authors declare that they have no competing interests.

Keywords: Invasive Species, Microbial Ecology, Phenology, Limnology, Time Series

This PDF file includes:
Figures 1-4
Supplemental Figures 1-2
Supplemental Tables 1, 3, 5-6

Figure 1. Sample Dates and microbial phenology. (A) Sample dates are represented by points, and season durations are represented by shading. **(B)** Non-metric multidimensional scaling of CLR-transformed bacterial community compositions. Points represent sample dates, and colors indicate seasons. Three seasons are compared in each NMDS plot to highlight season boundaries. All NMDS stress values are ≤ 0.1.

Figure 2. Shifts in *Cyanobacteria* phenology. (A) Spring relative abundance of *Cyanobacteria* aligned as days since clearwater. Average and standard deviation duration of clearwater is indicated by grey bars. Blue represents the pre-invasion period of 2000-2009, orange the post-spiny water flea period of 2010-2015, and red the post-zebra mussel period of 2016-2018. **(B)** Summer diversity of zooplankton and *Cyanobacteria*. Thick lines represent the mean Shannon diversity and shading represents the standard deviation. Thin lines represent individual years. Here the orange post-spiny water flea period includes years 2010-2018. Grey starred regions indicate the period of significant difference in mean diversity (p < 0.05). *Cyanobacteria* diversity is shown at the order level, and Zooplankton diversity is shown at ecologically relevant groupings with spiny water flea counts removed.

Figure 3. Shifts in cyanotoxin phenology. Orange represents the pre-zebra mussel invasion period of 2013-2015, red represents the post-zebra mussel invasion period of 2016-2018. Stars represent significant differences (p < 0.05). **(A)** Concentration of ananbaeopeptin and microcystin during the seasons with regular toxicity. **(B)** Microcystin toxin concentrations normalized by the yearly maximum concentrations to compare annual patterns. Shaded regions represent the duration of toxin production, defined as days when toxins were present at ≥ 15% of maximum toxicity. **(C)** Toxin production duration compared before and after the invasion of zebra mussels.

Figure 4. Shifts in heterotrophic bacteria phenology. Differential abundance was calculated on CLR-transformed abundances with a GLM model. Pink represents "winners" that differentially increased, and teal represents "losers" that differentially decreased. Points represent OTUs, and point sizes are log-scaled by the OTU's mean abundance. **(A)** Left, an example relative abundance over time curve of an OTU with positive effect size. Right, effect sizes for differential abundance of phylum *Bacteroidota* OTUs around the zebra mussel invasion. **(B)** Middle, effect sizes for differential abundance of order *Frankiales* OTUs around the zebra mussel invasion. Right and left are example winner and loser OTU abundances over time. **(C)** Percent of each season's microbial community that was differentially abundant over time. OTUs were normalized by their mean abundance before calculating the percentage of community change. The impact of the spiny water flea invasion is shown on the left, and the impact of the zebra mussel invasion is shown on the right.

Figure S1. Phytoplankton biovolume. The biovolume of *Cyanobacteria* and eukaryotic phytoplankton in each season, broken down by invasion status.

Figure S2. NMDS plots of invasion impacts by season. Panels correspond to different seasons, points represent sample dates included in each season, and colors indicate the sample date's invasion status. NMDS analysis was performed on CLR-transformed data.

Table S1. Season start dates.

Year	Spring	Clearwater	Early Summer	Late Summer	Fall	Ice-on
2000	2000-03-07	2000-04-12	2000-06-01	2000-07-21	2000-10-07	2000-12-13
2001	2001-04-07	2001-04-30	2001-06-23	2001-07-15	2001-09-26	2002-01-02
2002	2002-03-15	2002-04-24	2002-06-25	2002-07-22	2002-10-10	2003-01-04
2003	2003-04-03	2003-04-21	2003-06-17	2003-07-24	2003-10-07	2004-01-07
2004	2004-03-27	2004-04-20	2004-06-24	2004-07-21	2004-10-06	2004-12-25
2005	2005-04-05	2005-04-20	2005-06-22	2005-07-17	2005-10-21	2005-12-19
2006	2006-03-24	2006-03-24	2006-06-18	2006-07-21	2006-10-09	2007-01-20
2007	2007-03-27	2007-05-07	2007-06-06	2007-07-20	2007-10-28	2007-12-26
2008	2008-04-10	2008-04-21	2008-06-17	2008-07-22	2008-10-13	2008-12-16
2009	2009-03-23	2009-05-02	2009-06-25	2009-08-10	2009-10-03	2009-12-29
2010	2010-03-26	2010-05-14	2010-06-05	2010-07-05	2010-10-09	2010-12-15
2011	2011-04-03	2011-05-24	2011-06-21	2011-07-19	2011-10-11	2012-01-14
2012	2012-03-11	2012-04-10	2012-06-10	2012-07-03	2012-10-03	2013-01-14
2013	2013-04-11	2013-05-19	2013-06-18	2013-07-10	2013-10-22	2013-12-16
2014	2014-04-12	2014-05-29	2014-06-14	2014-07-14	2014-10-12	2015-01-02
2015	2015-04-03	2015-05-26	2015-06-10	2015-07-04	2015-10-14	2016-01-11
2016	2016-03-13	2016-05-03	2016-05-30	2016-07-04	2016-10-26	2017-01-01
2017	2017-03-07	2017-05-19	2017-06-13	2017-07-08	2017-10-21	2017-12-27
2018	2018-03-31	2018-05-19	2018-06-02	2018-07-04	2018-10-14	2018-12-15
2019	2019-03-31	2019-05-20	2019-07-03	2019-07-17	2019-10-10	2020-01-12

Table S2. Cyanotoxin concentrations.

Larger than a single page.

Table S3. Zooplankton grouping for diversity calculation.

LTER Species Code	LTER Name	Group[†]
50201	ACROPERUS HARPAE	Chydoridae Acroperus harpae
50300	ALONA	Chydoridae Alona
51000	CHYDORUS	Chydoridae Chydorus
51103	DAPHNIA MENDOTAE	Daphnia mendotae
51105	DAPHNIA PARVULA	Daphnia parvula
51130	DAPHNIA PULICARIA	Daphnia pulicaria
51107	DAPHNIA RETROCURVA	Daphnia retrocurva
51100	DAPHNIA	Daphnia unclassified
53701	BYTHOTREPHES LONGIMANUS	Bythotrephes longimanus
53601	SINOBOSMINA FRYEI	Bosmina sinobosmina fryei
51202	DIAPHANOSOMA BIRGEI	Diaphanosoma
52301	LEPTODORA KINDTI	Leptodora
40000	HARPACTICOID	Harpacticoida
30000	CALANOID	Calanoida unclassified
30901	AGLAODIAPTOMUS CLAVIPES	Calanoida Diaptomidae
30400	DIAPTOMID	Calanoida Diaptomidae
20100	ACANTHOCYCLOPS	Acanthocyclops
20302	DIACYCLOPS THOMASI	Diacyclops thomasi
20601	MESOCYCLOPS EDAX	Mesocyclops edax
20901	TROPOCYCLOPS PRASINUS MEXICANUS	Tropocyclops
20900	TROPOCYCLOPS	Tropocyclops
20000	CYCLOPOID	Cyclopoida uncalssified
10000	COPEPOD NAUPLII	Copepoda juvenile
15000	COPEPODITES	Copepoda juvenile
70100	CHAOBORUS	Chaoborus

[†]*Zooplankton were grouped by the finest taxonomic level that they could be confidently identified to before calculating the Shannon diversity.*

Table S4. Differentially abundant OTUs.

This table is larger than a single page.

Table S5. AIC table for models predicting *Cyanobacteria* phenology.

Hot[1]	Ice[2]	Pload[3]	SWF[4]	ZM[5]	AICc	delta	weight	df
0.31			-36.52	-37.19	163.22	0.00	0.30	5
			-34.27	-35.83	163.29	0.07	0.29	4
0.51		0.53	-35.83	-42.24	163.77	0.55	0.23	6
		0.09	-33.91	-36.57	166.92	3.70	0.05	5
	-0.01		-34.15	-36.05	167.03	3.82	0.05	5
0.33	0.06		-37.17	-36.26	167.39	4.18	0.04	6
0.50	-0.02	0.55	-35.52	-42.83	168.90	5.69	0.02	7
				-46.21	170.74	7.52	0.01	3
	-0.04	0.12	-33.43	-37.52	171.23	8.01	0.01	6
0.26			-48.47		172.55	9.33	0.00	4
	0.11		-46.56		173.60	10.38	0.00	4
		-0.12	-46.36		173.87	10.65	0.00	4
0.33	0.18		-49.58		175.23	12.01	0.00	5
				-57.25	175.84	12.62	0.00	3
0.31		0.13	-48.70		176.20	12.98	0.00	5
	0.14	-0.22	-46.95		176.98	13.76	0.00	5
		0.28		-58.78	178.59	15.37	0.00	4
0.16				-58.68	178.68	15.46	0.00	4
	-0.13			-58.76	178.69	15.48	0.00	4
0.34	0.17	0.03	-49.62		179.60	16.39	0.00	6
0.39		0.62		-64.14	180.60	17.38	0.00	5
	-0.21	0.43		-62.18	181.34	18.13	0.00	5
0.13	-0.10			-59.65	182.18	18.96	0.00	5
0.39	-0.21	0.78		-67.46	183.91	20.69	0.00	6
					186.74	23.52	0.00	2
	0.04				189.57	26.35	0.00	3
-0.03					189.58	26.37	0.00	3
		-0.02			189.59	26.37	0.00	3
	0.05	-0.05			192.82	29.60	0.00	4
-0.02	0.04				192.82	29.61	0.00	4
-0.06		-0.07			192.83	29.61	0.00	4
-0.05	0.05	-0.09			196.56	33.35	0.00	5

[1]Days with average epilimnion temperature ≥ 23°C.
[2]Pevious winter ice duration in days.
[3]Mean annual daily phosphorus loading.
[4]Spiny water flea invasion status.
[5]Zebra mussel invasion status.

Table S6. AIC table for models predicting *Frankiales* phenology.

Hot[1]	Ice[2]	Pload[3]	SWF[4]	ZM[5]	AICc	delta	weight	df
			0.45	0.54	12.35	0.00	0.40	4
0.00			0.49	0.56	14.35	2.00	0.15	5
-0.01		-0.01	0.48	0.66	14.81	2.46	0.12	6
		0.00	0.43	0.57	15.68	3.33	0.08	5
	0.00		0.44	0.57	15.86	3.50	0.07	5
			0.63		16.44	4.09	0.05	3
				0.81	17.86	5.51	0.03	3
-0.01	0.00	-0.01	0.44	0.73	18.80	6.44	0.02	7
0.00	0.00		0.48	0.57	18.86	6.51	0.02	6
0.00			0.66		18.98	6.62	0.01	4
	0.00	-0.01	0.40	0.64	19.14	6.78	0.01	6
	0.00		0.63		19.80	7.44	0.01	4
		0.00	0.63		19.80	7.45	0.01	4
		-0.01		0.84	20.18	7.83	0.01	4
	0.00			0.84	20.45	8.10	0.01	4
0.00				0.83	21.01	8.66	0.01	4
	0.00	-0.01		0.92	22.05	9.70	0.00	5
-0.01		-0.01		0.93	22.33	9.98	0.00	5
-0.01		0.00	0.67		22.46	10.11	0.00	5
0.00	0.00		0.67		22.78	10.43	0.00	5
	0.00	0.00	0.63		23.72	11.36	0.00	5
0.00	0.00			0.85	24.28	11.93	0.00	5
-0.01	0.00	-0.02		1.02	24.55	12.20	0.00	6
					26.30	13.95	0.00	2
-0.01	0.00	0.00	0.68		27.06	14.71	0.00	6
		0.00			29.15	16.80	0.00	3
0.00					29.17	16.81	0.00	3
	0.00				29.22	16.86	0.00	3
	0.00	0.00			32.50	20.15	0.00	4
0.00		0.00			32.50	20.15	0.00	4
0.00	0.00				32.52	20.17	0.00	4
0.00	0.00	0.00			36.41	24.06	0.00	5

[1]Days with average epilimnion temperature ≥ 23°C.
[2]Pevious winter ice duration in days.
[3]Mean annual daily phosphorus loading.
[4]Spiny water flea invasion status.
[5]Zebra mussel invasion status.