Title

N-terminal region of *Drosophila melanogaster* Argonaute2 forms amyloid-like aggregates

Authors

Haruka Narita, Tomohiro Shima#, Ryo Iizuka, and Sotaro Uemura#

Affiliation

Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Tokyo, Japan

Corresponding authors

T.S.,

Tel: (+81)-35841-4399

Email: tomohiro.shima@bs.s.u-tokyo.ac.jp

S.U.,

Tel: (+81)-35841-4396

Email: uemura@bs.s.u-tokyo.ac.jp

Running title

Amyloid-like aggregation by the N-terminus of DmAgo2

Keywords

Argonaute, aggregation, amyloid, prion, RNA interference

Abstract

Argonaute proteins play a central role in RNA silencing by forming protein-small RNA complexes responsible for the silencing process. While most Argonaute proteins have a short N-terminal region, Argonaute2 in *Drosophila melanogaster* (DmAgo2) harbors a long and unique N-terminal region. Previous *in vitro* biochemical studies have shown that the loss of this region does not impair the RNA silencing activity of the complex. However, an N-terminal mutant of *Drosophila melanogaster* has demonstrated abnormal RNA silencing activity. To explore the causes of this discrepancy between *in vitro* and *in vivo* studies, we investigated the biophysical properties of the region. Because the N-terminal region is highly rich in glutamine and glycine residues, which is a well-known property for prion-like domains (PrLD), the possibility of the N-terminal region functioning as a PrLD was tested. Our biochemical assays demonstrated that the N-terminal region can form aggregates that are not dissociated even in the presence of SDS. Also, the aggregates enhanced the fluorescence intensity of thioflavin-T, an amyloid detection reagent. The kinetics of the aggregation followed that of typical amyloid formation exhibiting the self-propagating activity. Further, we directly visualized the aggregation process of the N-terminal region under fluorescence microscopy and found that the aggregations took fractal or fibril shapes. Together, the results indicate that the N-terminal region is a PrLD. Many other PrLDs have been reported to modulate the function of proteins through their aggregation. Therefore, our results raise the possibility that aggregation of the N-terminal region regulates the RNA silencing activity of DmAgo2.

Introduction

RNA silencing is involved in various biological processes including antiviral defense, development, and the maintenance of genomic integrity (1). Among the RNA silencing machinery, Argonaute proteins play key roles by directly binding to small RNAs to form the RNA Induced Silencing Complex (RISC) (2). After the small RNA in RISC recognizes its RNA targets, Argonaute proteins display endonucleolytic activity or translational repression (3). Despite the functional variety of Argonaute proteins, their structural organization is highly conserved, sharing four distinct domains: N, PAZ, MID, and PIWI (4). Previous studies have revealed the functions of each domain. The N domain initiates duplex unwinding immediately after small RNA binds to Argonaute (5).

The PAZ and MID domains recognize the 5′ and 3′ ends of the small RNA, respectively

(6–8). The PIWI domain, which shares structural similarities with ribonucleases,

cleaves the target RNA(9, 10).

In contrast to these highly conserved domains, the N-terminal region, which is located

upstream of the N domain, shows extensive diversity among species (11). While most

Argonaute proteins have a short N-terminal region (e.g., 24 residues in human Ago2),

Argonaute2 in *Drosophila melanogaster* (DmAgo2) harbors a long and unique

N-terminal region. The N-terminal region (residues 1–398; henceforth, Nter) comprises

almost one-third of the residues in DmAgo2 (1208 residues). Partial truncation of Nter

(residues 326–371 deletion) has been reported to impair RNA silencing in mutant flies

(12). However, an *in vitro* study demonstrated that another DmAgo2 mutant with Nter

deletion (residues 1–278 deletion) still forms RISC and retains RNA cleavage

activity(13).

Many eukaryotic RNA-binding proteins contain a prion-like domain (PrLD), which

organizes intracellular condensates and regulates biochemical reactions (14–17). PrLDs

consist of intrinsically disordered, low-complexity sequences often enriched in

glutamine and glycine residues (18, 19). The Nter sequence shares this property. The

content of glutamine and glycine residues in DmAgo2 Nter reach approximately 40%

and 20%, respectively (11, 12). Inspired by these findings, here we tested whether

DmAgo2 Nter is a PrLD. Our *in silico* prediction and biochemical assays demonstrated

that Nter is a PrLD that can form amyloid-like aggregates. We also directly visualized

by fluorescence microscopy how Nter aggregates and found that the aggregates are

polymorphic. Since many PrLDs have been shown to regulate biochemical reactions

through their aggregate formation (14–17), our results raise the possibility that Nter

regulates RNA silencing activities in cells by forming aggregates.

Results

PrLD prediction based on the amino acid sequence of DmAgo2

To confirm the characteristics of DmAgo2 based on its amino acid sequence (Figure

1A), we performed intrinsically disordered regions (IDRs) prediction and three kinds of

PrLDs prediction. First, to confirm that DmAgo2 harbors IDRs, we analyzed the

DmAgo2 amino acid sequence using PONDR (20), a neural network-based prediction

algorithm of IDRs. Two parts of Nter (residues 1–87 and 105–412) were identified as

regions with high disorder scores (Figure 1B). Then, the possibility that Nter is not only an IDR but also a PrLD that can form amyloid-like aggregates was predicted by three PrLD prediction algorithms: the prion-like amino acid composition (PLAAC) (21), Prion Aggregation Prediction Algorithm (PAPA) (22), and PrionW (23). PLAAC, a Hidden Markov Model (HMM)-based prion prediction algorithm, identified residues 14–386 in Nter as the region with a high prion-like score (Figure 1B). Because the predictions of Nter being PrLD varied across the algorithms (Supplementary Figure 1), we next tested experimentally whether Nter has the capacity to form amyloid-like aggregates.

Nter can form SDS-resistant amyloid-like aggregates in a typical amyloid formation manner.

We prepared a recombinant protein of Nter fused with mCherry at its N-terminal (mCherry-Nter) to test whether Nter behaves as a PrLD. PrLDs are known to form aggregates even in the presence of SDS (24). To test if Nter forms SDS-resistant aggregates, we first incubated 5 µM of monomeric mCherry-Nter in aggregation buffer (20 mM sodium phosphate, 50 mM sodium chloride, pH 7.4) for 3 days at room temperature. Then, the aggregates in the solution were collected by ultracentrifugation and subjected to semi-denaturing detergent agarose gel electrophoresis (SDD-AGE) (24). As references for the assay, mCherry and Sup35NM-mCherry were also subjected to SDD-AGE. mCherry is a monomeric protein (25), whereas Sup35NM is a well-known PrLD and can form amyloid-like aggregates (26). mCherry did not appear as a distinct band, indicating that mCherry did not form any aggregates (Figure 2). Contrarily, Sup35NM-mCherry and mCherry-Nter predominantly migrated as smears with high molecular weights, suggesting that mCherry-Nter formed SDS-resistant aggregates the same as Sup35NM-mCherry (Figure 2). The aggregates of Sup35NM-mCherry and mCherry-Nter were dissolved into soluble fractions when boiled. Thus, the results showed that Nter forms SDS-resistant aggregates.

To further test whether Nter aggregates take amyloid-like structures, we performed Thioflavin-T (ThT) assays. ThT is a fluorescent probe with specific fluorescence enhancement upon binding with amyloid fibrils (27, 28). First, we measured the fluorescence spectrum of 20 µM ThT with and without 5 µM Nter solution incubated for 12 h beforehand. In the presence of Nter, the fluorescence intensity of ThT was enhanced approximately four times compared to that without Nter, and a peak at 490 nm was clearly visible in the fluorescence spectrum (Figure 3A), indicating the presence of amyloid-like structures in the Nter solution.

To confirm that these aggregates were formed in a typical amyloid formation manner, we tracked the aggregation process over time by monitoring the ThT fluorescence intensity under microscopy. Consistent with the canonical amyloid formation from monomers generally requiring a lag phase for nucleation (29), the ThT fluorescence intensity with Nter showed an initial lag phase of ~300 min and then reached a plateau at ~800 min after mixing monomeric Nter with ThT in aggregation buffer (Figure 3B, Supplementary Figure 2A). Prion-like proteins generally exhibit seeded self-propagation *in vivo* and *in vitro* (26, 30). Therefore, we prepared the seeds of Nter aggregates and examined whether the seeds can accelerate the aggregation process. The addition of the seeds reduced the initial lag phase to less than 100 min (Figure 3C, Supplementary Figure 2B), showing that the Nter aggregation process follows canonical amyloid formation kinetics with a self-propagation property.

Direct fluorescence visualization of Nter aggregates.

Next, we directly visualized the shape of Nter aggregates using fluorescence microscopy to investigate their morphology. The fluorescence of mCherry-Nter that had been flushed into a glass chamber and incubated for 12 h to form aggregates revealed fractal or fibril-shaped aggregates. When ThT was added to the chamber, the fluorescent images of ThT and mCherry matched completely, suggesting that the entire region of the Nter aggregates took an amyloid-like structure (Figure 4A–D).

Since the aggregates showed various shapes (Figure 4A–C and Supplementary Figure 3, we analyzed and classified the shapes using principal component analysis (PCA) with k-means clustering into three groups: linear fractal-shaped, branched fractal-shaped and fibril-shaped aggregates (Figure 4E). The most frequently observed group (56%) was linear fractal-shaped aggregates (Figure 4A and Supplementary Figure 3A). In these aggregates, ~0.5 μm fluorescent puncta were linearly connected to each other. The median length of these aggregates was 1.4 [1.2–2.1] μm (median with quartile range, *n* = 83, Supplementary Figure 4A), corresponding to ~2 puncta. For the second-largest group, 32% of the whole aggregates were classified as branched fractal-shaped aggregates (Figure 4B and Supplementary Figure 3B). In the aggregates of this group, several puncta were bound to branch off from the main chain of the aggregates. The median length of the longest chain of the aggregates was 3.3 [2.2–4.9] μm (median with quartile range, *n* = 47, Supplementary Figure 4B), and the median number of branched points was 4 [3–6] (median with quartile range, *n* = 47, Supplementary Figure 4C). The remaining 12% of the whole aggregates were classified as fibril-shaped aggregates (Figure 4C and Supplementary Figure 3C). This fibril shape is the most common

structure for amyloids (31). The median length of the Nter fibril was 27 [19–50] μm
(median with quartile range, $n = 17$, Supplementary Figure 4D), taking a shape much
longer than the fractal-shaped aggregates. Most Nter fibrils contained a few number of
puncta, but these puncta did not directly bind to each other. Thus, we could classify the
shapes of the Nter aggregates according to the presence of branching, the length of the
aggregates and the alignment of puncta.

Time-lapse imaging of the Nter aggregation process demonstrated how the
fractal-shaped aggregates formed. First, fluorescent puncta docked to each other,
forming linear fractal-shaped aggregates (Figure 5A). Next, linear fractal-shaped
aggregates docked to each other, forming branched fractal-shaped aggregates (Figure
5B). Finally, branched fractal-shaped aggregates docked to each other, forming larger
branched fractal-shaped aggregates (Figure 5C). Therefore, it is most likely that the
linear and branched fractal-shaped aggregates share the same formation process, and the
difference in the stage of the formation process is seen as the difference in shape.

Discussion

Here, we demonstrated that Nter is a PrLD by *in silico* prediction and biochemical
assays. The PLAAC algorithm predicted Nter is a PrLD based on its characteristic
sequence. mCherry-Nter fusion protein formed aggregates that exhibit amyloid-specific
properties of resistance to SDS and binding to ThT. Furthermore, Nter aggregated in a
manner consistent with typical amyloids, including a lag phase for nucleation and
self-propagation activity. These results strongly suggest that Nter can form amyloid-like
aggregates.

From a structural viewpoint, we found that Nter aggregates can take two distinct
conformational states. Most aggregates took fractal shapes, while the remaining had
fibril shapes. This observation is consistent with previous studies showing that
fractal-shaped aggregates tend to form under diffusion limited-conditions (32–35),
because protein diffusion is limited in the 145-μm thick chamber in our assays. Within
cells, protein diffusion should be even more restricted due to molecular crowding and
the size of the cytoplasm. Therefore, Nter aggregates are expected to be more prone to
taking on a fractal shape, although it is yet to be demonstrated that these observed
fractal-shaped aggregates are really formed in cells. In addition, fractal-shaped
aggregates docked to form larger fractal-shaped aggregates. On the other hand, although
the forming process of fibril-shaped aggregates was not observed, fibril-shaped
aggregates frequently contained a few numbers of puncta in their fibrils (Figure 4C and

Supplementary Figure 3C). This finding implies that fibril-shaped aggregates can elongate from the edges of the puncta. Thus, we propose a growth model of Nter aggregates in which the docking of several puncta form fractal-shaped aggregates and polymerization of monomeric Nter from a single punctum forms fibril-shaped aggregates (Figure 5D).

Nter aggregation may indirectly modulate the RNA silencing activity of DmAgo2. It has been shown that Nter is not essential for DmAgo2 to exhibit RNA silencing activity under *in vitro* conditions, but the partial truncation of Nter causes RNA silencing defects in the compound eyes of flies (12). Combining these previous studies with our current results raises the possibility that *in vivo* Nter aggregation modulates RISC formation and/or RNA cleavage activity by other domains of DmAgo2. Protein aggregation can both be inhibitory on protein activity by steric hindrance effects on the substrate binding or facilitative by increasing local protein concentrations (36). Furthermore, it has been reported that protein aggregation is a major driving force for liquid-liquid phase separation in the cytoplasm, allowing for the proper subcellular localization of biomolecules (14). Indeed, DmAgo2 is localized in cytoplasmic D2 bodies, where endogenous siRNAs are loaded onto DmAgo2, and this localization is essential for the formation of RISC with the proper small RNA (37). The ability of Nter to form amyloid-like aggregates may contribute to such modulation of DmAgo2 activity and localization. Although it still needs to be verified whether the aggregation of endogenous DmAgo2 actually occurs in cells, this hypothesis can explain the apparent inconsistency among previous *in vitro* and *in vivo* studies regarding the effects of Nter in RNA silencing.

We focused on DmAgo2 because *Drosophila melanogaster* is one of the most well-studied model organisms in RNA silencing studies. However, the aggregation of the N-terminal region may be a general phenomenon not limited to DmAgo2. Although Argonaute proteins of many organisms have only a short N-terminal region, Ago2 among arthropods share glutamine- and glycine-enriched properties in the N-terminal region (38). Their amino acid composition, which is a major factor in amyloid-formation potential (39), is close to DmAgo2 Nter. Therefore, the N-terminal regions of Ago2 in other arthropods may also form amyloid-like aggregates as PrLDs. Also, some plant Argonaute proteins possess glutamine- and glycine-rich N-terminal regions (Supplementary Figure 5), implying they share the same property. Since many Argonaute proteins are likely to have the similar characteristic sequence at the N-terminus, we propose to call such region N-terminal Argonaute Prion-like (NAP)

domain. The physiological features and function of this aggregation property of NAP
domains await further study, but the property has the potential to function as a
regulatory mechanism for RNA silencing across species.

Experimental procedures

In silico analysis

The amino acid sequence of DmAgo2 (equivalent to Uniprot entry Q9VUQ5 with the
deletion of residues 43–48) was derived from native *Drosophila* DNA. The sequence
was analyzed using the programs PLAAC (http://plaac.wi.mit.edu/) (21), PAPA
(https://combi.cs.colostate.edu/supplements/papa/) (40), and PrionW
(http://bioinf.uab.cat/prionw/) (23) with default settings. Disorder scores were
calculated using the Predictor of Natural Disordered Regions (PONDR) algorithm with
the VLXT predictor (http://www.pondr.com/) (41).

Plasmid construction

The expression plasmid for N-terminal 6×His-tagged mCherry-Nter was constructed by
inserting the sequence encoding DmAgo2 Nter (residues 1–398) amplified from the
native DmAgo2 sequence into pCold I (TAKARA Bio) using the InFusion HD cloning
kit (Clontech). The expression plasmid for C-terminal 6×His-tagged
Sup35NM-mCherry was constructed by inserting the sequence encoding mCherry into
the expression plasmid for Sup35NM based on pET29b (residues 1–253) (a kind gift
from Prof. M. Tanaka (RIKEN CBS, Japan)).

Protein expression and purification

For the protein expression, *E. coli* BL21(DE3) cells were transformed with the
constructed plasmids and selected using 50 mg/L carbenicillin (for pCold I and pET11a
plasmids) or 50 mg/L kanamycin (for pET29b plasmid). These antibiotics were added in
all culture media described below. The transformed *E. coli* cells were pre-cultured in 5
mL of LB medium overnight at 37°C and then inoculated into 2 L of TB medium. The
cells were grown at 37°C until OD_{600} reached 1.2 (6×His-tagged mCherry-Nter and
6×His-tagged Sup35NM-mCherry) or 0.6 (mCherry-His). 6×His-tagged mCherry-Nter
was expressed with 1 mM IPTG at 12°C. mCherry-His and 6×His-tagged
Sup35NM-mCherry were expressed with 1 mM IPTG at 28°C. After 20 h of culture, the
cells were harvested and then lysed in 80 mL of lysis buffer (20 mM Hepes-KOH pH8.0,
100 mM NaCl, and 6 M GdnHCl) with 10-s sonication at an intensity of 2 on ice using
the ultrasonic disruptor (UD211, TOMY) a total of 9 times with 20 s intervals in
between. Following centrifugation of the lysate at 18,000 × g for 30 min, 6 mL of 2 ×
Ni-NTA resin (Ni-NTA Agarose HP, FUJIFILM Wako Pure Chemical) was added to

the supernatant. After a 60-min rotation at room temperature, the solution was transferred to a disposal column (Muromac Mini-column M, Muromachi Chemical Inc). The resin was washed twice with 3 column volumes (CVs) of wash buffer (40 mM imidazole-HCl, 20 mM Hepes-KOH, pH8.0, 100 mM NaCl, and 6 M GdnHCl) and eluted with 1 CV of elution buffer (400 mM imidazole-HCl, 20 mM Hepes-KOH, pH8.0, 100 mM NaCl, and 6 M GdnHCl) three times. The elution fractions were resolved on 12% denaturing polyacrylamide gels and visualized by Coomassie Brilliant blue (Nacalai Tesque) staining. Elution fractions containing proteins of interest were pooled and ultra-filtered to remove large aggregates using Vivaspin 6, 100,000 MWCO (Sartorius Stedim Biotech GmbH). Finally, the flow-through was concentrated using Vivaspin 6, 30,000 MWCO (Sartorius Stedim Biotech GmbH). The purified proteins were snap-frozen in liquid nitrogen and stored at -80°C until use.

Monomers of mCherry-Nter, mCherry, and Sup35NM-mCherry were prepared as follows. First, purified proteins were diluted 50 times by aggregation buffer (20 mM sodium phosphate, pH 7.4, 50 mM NaCl). Next, the protein solution was ultracentrifuged at 418,000 × g for 20 min at 25°C to remove large aggregates. Following the ultracentrifugation, the supernatant, equivalent to a half volume of the solution, was immediately transferred to a new 1.5 mL tube. The protein concentrations were determined using the following molar extinction coefficients at 280 nm: mCherry-Nter, 66,240 $M^{-1}cm^{-1}$; Sup35NM-mCherry, 65,670 $M^{-1}cm^{-1}$; and mCherry, 35,870 $M^{-1}cm^{-1}$. The molar extinction coefficients were calculated using the ProtParam tool of ExPASy (https://web.expasy.org/protparam/) (42). Then, the protein solution was diluted to 5 μM by aggregation buffer. Nter aggregates were prepared by incubating the above solution at room temperature for the given time.

Seeds were prepared as follows. First, purified mCherry-Nter was diluted 10 times by elution buffer. Next, the solution was applied to a gel filtration spin column (Micro Bio-Spin 30 column, Bio-Rad) equilibrated with aggregation buffer to remove GdnHCl and ultracentrifuged at 418,000 × g for 20 min at 25°C to remove large aggregates. Finally, the supernatant, equivalent to 90% solution volume, was immediately transferred to a new 1.5 mL tube as a seed.

SDD-AGE

After monomers were incubated for 3 days at room temperature without agitation, aggregates were pelleted by ultracentrifugation at 418,000 × g for 20 min at 25°C. After completely removing the supernatant, the pellet was resuspended with aggregation

buffer. Prepared samples were subjected to SDD-AGE analysis. First, aggregate
solutions were mixed with $4 \times$ sample buffer ($2 \times$ TAE, 20% (v/v) glycerol, 4% (w/v)
SDS, 0.25% (w/v) bromophenol blue). Next, after incubation for 15 min at room
temperature with or without heat treatment at 95°C for 2 min, the samples were loaded
onto a 1.5% agarose gel containing $1 \times$ TAE and 0.1% SDS. Finally, the aggregates and
monomer protein were resolved on the gel in running buffer ($1 \times$ TAE, and 0.1% SDS)
at 75 V for 90 min at 4°C, followed by capillary blotting onto a PVDF membrane
(FUJIFILM Wako Pure Chemical) for the western blotting analysis, as described
previously (24). Anti-RFP polyclonal antibody (1/5,000 dilution from the product,
PM005, MBL) and HRP-labeled IgG detector (1/5,000 dilution from the product,
Western BLoT Rapid Detect v2.0, Takara Bio) were used as primary and secondary
antibodies, respectively. Proteins were detected with SuperSignal West Femto (Thermo
Scientific) on a gel imager (Amersham Imager 600, Cytiva).

Measurement of ThT spectrum

5 μM mCherry-Nter monomers were incubated for 12 h at 25°C to prepare
mCherry-Nter aggregates. Fluorescence spectra of mCherry-Nter aggregates in solution
with or without 20 μM ThT (FUJIFILM Wako Pure Chemical) were obtained using a
fluorescence spectrometer (RF-6000, Shimadzu) at room temperature. The excitation
wavelength was set at 455 nm (bandwidth: 5 nm), and the emission was recorded from
465 to 560 nm (bandwidth: 5 nm).

Chamber preparation for fluorescence microscopy

Flow chambers were prepared as described previously (Narita et al. 2020 *bioRxiv*) with
some modifications. Briefly, the coverslips (No. 1S 22×22 mm and No.1S 24×32 mm,
Matsunami) were cleaned in 1 N KOH for 15 min with sonication (Bransonic tabletop
cleaner, Emerson). All subsequent preparation procedures were performed in a clean
hood (Matsusada Precision). After 20 times rinsing with Milli-Q water and drying in a
dryer, the coverslips were cleaned using a plasma cleaner (YHS-R,
SAKIGAKE-Semiconductor Co., Ltd.). A 25 μL volume micro-chamber was made by
placing a small coverslip of 22×22 mm over a 24×32 mm glass coverslip using
double-sided adhesive tape (145 μm thickness, TERAOKA SEISAKUSHO CO., LTD)
in a clean hood. First, Lipidure-BL103 (NOF Corporation) was flowed into the chamber
to coat the glass surface. After a 2-min incubation and excess Lipidure-BL103 removal

by three washes with 25 µL aggregation buffer, each sample was flushed into the glass
chamber.
To evaluate the colocalization of ThT and mCherry and the size distribution of the
aggregates, mCherry-Nter aggregates were prepared by a 12 h-incubation of monomer
mCherry-Nter at room temperature. mCherry-Nter aggregates were observed as follows.
mCherry-Nter aggregates solution was flushed into a Lipidure-coated glass chamber and
incubated for 2 min at room temperature. Finally, excess mCherry-Nter unbound on the
glass surface was removed by three washes with 25 µL aggregation buffer with 20 µM
ThT.
To measure the aggregation kinetics with or without the aggregate seeds, 5 µM
mCherry-Nter monomers were flushed into a Lipidure-coated glass chamber and sealed
with VALAP (1:1:1 mixture of vaseline, lanolin, and paraffin). In the presence of seeds,
the final 5 µM of seeds was added to the monomer solution.

Fluorescence Microscopy
Nter aggregates were observed using an inverted microscope (Nikon Ti-E) equipped
with a filter cube for mCherry imaging consisting of an excitation filter (#67-033,
Edmund Optics), a dichroic mirror (#67-083, Edmund Optics) and an emission filter
(#67-036, Edmund Optics) and a filter cube for ThT imaging consisting of an excitation
filter (#67-026, Edmund Optics), a dichroic mirror (#67-078, Edmund Optics) and an
emission filter (#67-028, Edmund Optics). mCherry and ThT were illuminated with a
532 nm-laser (OBIS LX/LS, Coherent) and 445 nm-LED light (SOLIS-445C,
THORLABS), respectively. The images were obtained through a Plan Fluor 10×/0.30 or
Plan Apoλ 40×/0.95 objective (Nikon) and recorded at 10 frames/s using an Orca
Flash4.0 V3 digital CMOS camera (Hamamatsu Photonics). All equipment in the
microscopy system was controlled by Micro-Manager software (43). To measure
aggregates formation kinetics, ThT fluorescence images were recorded at 10-min
intervals.

Image Analysis
Image analysis was performed using either ImageJ or Python. To accurately evaluate
the colocalization of ThT and mCherry, we corrected the chromatic aberration of the
images obtained from both fluorescence channels using ImageJ and the TurboReg
plugin   (http://bigwww.epfl.ch/thevenaz/turboreg/).   The   translational   distortion

parameters were obtained by comparing the averaged image from 10 images of an
objective micrometer (OB-M, 1/100, Olympus) in each channel. Four regions of interest
(3 × 3 pixels) were used for the calculation to minimize artifacts. The images of
mCherry-Nter aggregates corrected using the obtained parameters were used for further
colocalization analysis.

To classify the shapes of the aggregates observed by the ThT fluorescence, we first
extracted features listed below: the longest chain length, the mean width, the mean
width per longest chain length, the total fluorescence intensity, the number of branched
points per longest chain length, the number of fluorescence intensity peaks per longest
chain length, and the percentage of the longest chain length to total length. To calculate
these features, the images were preprocessed as described below. First, the background
of each image was subtracted using the "Subtract Background" algorithm in ImageJ
with a rolling ball radius of 200 pixels. Next, bright particles with more than 9 pixels
with fluorescence intensity beyond the given threshold (60 a.u. out of 65,535 a.u.) were
detected as aggregates. Following the above preprocessing, to calculate the total length
of each aggregate, the "Skeletonize" algorithm was applied to each detected aggregate.
Next, the longest chain length and the number of branched points of each aggregate
were calculated from the skeletonized images. The mean width of each aggregate along
the longest chain was calculated by averaging the widths along the longest chain using
ImageJ. The number of fluorescence intensity peaks per longest chain length was
detected using the Laplacian of Gaussian (LoG) method available in the scikit-image
module in Python (44). This number was evaluated because the smallest unit of a
fractal-shaped aggregate is a round-shaped aggregate detected as a fluorescence
punctum. Finally, we performed a PCA analysis based on the extracted eight features to
obtain the two axes corresponding to the maximum variation (PC1) and second-most
variation (PC2). Using the plot, the k-means clustering algorithm classified these
aggregates into four groups. One of these groups consisted of only one sample, where
both PC1 and PC2 values were outliers because two ends of the aggregate were
connected to form a ring. We re-classified this one sample into "fibril-shaped
aggregate". The PCA result was plotted using a Python plotting library (Matplotlib:
http://matplotlib.org) (45).

To calculate the amount of ThT bound to the aggregates at each time point, bright
particles with more than 9 pixels of high fluorescence intensity (more than 250 a.u. out
of 65,535 a.u.) were detected as the aggregates. ThT fluorescence intensity of the pixels
with the detected aggregates were plotted against time.

Data availability
All image data sets and source codes for image analysis were deposited at Figshare
(10.6084/m9.figshare.20438139)

Acknowledgement
We thank Dr. Motomasa Tanaka (RIKEN Center for Brain Science) for providing the
expression plasmid of Sup35NM. We also thank the members of the Uemura and Siomi
laboratories for valuable discussions.

Author contributions
Conceptualization, H.N. and T.S.; Data curation, H.N.; Methodology, H.N. and T.S.;
Investigation, H.N.; Analysis, H.N.; Supervision, T.S.; Visualization, H.N. and T.S.;
Writing – original draft, H.N. and T.S.; Writing – review & editing, H.N., T.S., R.I. and
S.U.

Funding and additional information
HN is supported by a JSPS Research Fellowship for Young Scientists and WINGS-LST,
the University of Tokyo. This work is supported by Grant-in-Aid for JSPS Fellows,
21J11218 (to HN) and by JSPS KAKENHI (18K06147,19H05379 and 21H00387 to
T.S.).

Conflict of interests
The authors declare no conflicts of interest associated with this manuscript.

References

1. Denli, A. M., and Hannon, G. J. (2003) RNAi: an ever-growing puzzle. *Trends Biochem. Sci.* **28**, 196–201

2. Siomi, H., and Siomi, M. C. (2009) On the road to reading the RNA-interference code. *Nature*. **457**, 396–404

3. Okamura, K. (2004) Distinct roles for Argonaute proteins in small RNA-directed RNA cleavage pathways. *Genes Dev*. **18**, 1655–1666

4. Olina, A. V., Kulbachinskiy, A. V., Aravin, A. A., and Esyunina, D. M. (2018) Argonaute Proteins and Mechanisms of RNA Interference in Eukaryotes and Prokaryotes. *Biochemistry Moscow*. **83**, 483–497

5. Kwak, P. B., and Tomari, Y. (2012) The N domain of Argonaute drives duplex unwinding during RISC assembly. *Nat Struct Mol Biol*. **19**, 145–151

6. Boland, A., Tritschler, F., Heimstädt, S., Izaurralde, E., and Weichenrieder, O. (2010) Crystal structure and ligand binding of the MID domain of a eukaryotic Argonaute protein. *EMBO Rep*. **11**, 522–527

7. Ma, J.-B., Yuan, Y.-R., Meister, G., Pei, Y., Tuschl, T., and Patel, D. J. (2005) Structural basis for 5′-end-specific recognition of guide RNA by the A. fulgidus Piwi protein. *Nature*. **434**, 666–670

8. Lingel, A., Simon, B., Izaurralde, E., and Sattler, M. (2003) Structure and nucleic-acid binding of the Drosophila Argonaute 2 PAZ domain. *Nature*. **426**, 465–469

9. Parker, J. S., Roe, S. M., and Barford, D. (2004) Crystal structure of a PIWI protein suggests mechanisms for siRNA recognition and slicer activity. *EMBO J*. **23**, 4727–4737

10. Song, J.-J. (2004) Crystal Structure of Argonaute and Its Implications for RISC Slicer Activity. *Science*. **305**, 1434–1437

11. Hain, D., Bettencourt, B. R., Okamura, K., Csorba, T., Meyer, W., Jin, Z., Biggerstaff, J., Siomi, H., Hutvagner, G., Lai, E. C., Welte, M., and Müller, H.-A. J. (2010) Natural Variation of the Amino-Terminal Glutamine-Rich Domain in Drosophila Argonaute2 Is Not Associated with Developmental Defects. *PLoS ONE*. **5**, e15264

12. Meyer, W. (2006) Overlapping Functions of Argonaute Proteins in Patterning and Morphogenesis of Drosophila Embryos. *PLoS Genet*. **2**, e134

13. Liu, Y., Ye, X., Jiang, F., Liang, C., Chen, D., Peng, J., Kinch, L. N., Grishin, N. V., and Liu, Q. (2009) C3PO, an Endoribonuclease That Promotes RNAi by Facilitating RISC Activation. *Science*. **325**, 750–753

14.   Banani, S. F., Lee, H. O., Hyman, A. A., and Rosen, M. K. (2017) Biomolecular
condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol*. **18**,
285–298

15.   Gotor, N. L., Armaos, A., Calloni, G., Vabulas, R. M., de Groot, N. S., and
Tartaglia, G. G. (2020) *RNA-Binding and Prion Domains: The Yin and Yang of*
Phase Separation, Molecular Biology, 10.1101/2020.01.14.904383

16.   Jakobson, C. M., and Jarosz, D. F. (2018) Organizing biochemistry in space and
time using prion-like self-assembly. *Curr. Opin. Syst. Biol.* **8**, 16–24

17.   Keleman, K., Krüttner, S., Alenius, M., and Dickson, B. J. (2007) Function of the
Drosophila CPEB protein Orb2 in long-term courtship memory. *Nat Neurosci*. **10**,
1587–1593

18.   Alberti, S., Halfmann, R., King, O., Kapila, A., and Lindquist, S. (2009) A
Systematic Survey Identifies Prions and Illuminates Sequence Features of
Prionogenic Proteins. *Cell*. **137**, 146–158

19.   King, O. D., Gitler, A. D., and Shorter, J. (2012) The tip of the iceberg:
RNA-binding proteins with prion-like domains in neurodegenerative disease.
Brain Research. **1462**, 61–80

20.   Romero, P., Obradovic, Z., Li, X., Garner, E. C., Brown, C. J., and Dunker, A. K.
(2001) Sequence complexity of disordered protein. *Proteins*. **42**, 38–48

21.   Lancaster, A. K., Nutter-Upham, A., Lindquist, S., and King, O. D. (2014)
PLAAC: a web and command-line application to identify proteins with prion-like
amino acid composition. *Bioinformatics*. **30**, 2501–2502

22.   Toombs, J. A., Petri, M., Paul, K. R., Kan, G. Y., Ben-Hur, A., and Ross, E. D.
(2012) De novo design of synthetic prion domains. *PNAS*. **109**, 6519–6524

23.   Zambrano, R., Conchillo-Sole, O., Iglesias, V., Illa, R., Rousseau, F., Schymkowitz,
J., Sabate, R., Daura, X., and Ventura, S. (2015) PrionW: a server to identify
proteins containing glutamine/asparagine rich prion-like domains and their
amyloid cores. *Nucleic Acids Res*. **43**, W331–W337

24.   Halfmann, R., and Lindquist, S. (2008) Screening for Amyloid Aggregation by
Semi-Denaturing Detergent-Agarose Gel Electrophoresis. *J Vis Exp*. **17**, e838

25.   Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N. G., Palmer, A.
E., and Tsien, R. Y. (2004) Improved monomeric red, orange and yellow
fluorescent proteins derived from Discosoma sp. red fluorescent protein. *Nat*
Biotechnol. **22**, 1567–1572

26.   King, C.-Y., Tittmann, P., Gross, H., Gebert, R., Aebi, M., and Wuthrich, K. (1997)
Prion-inducing domain 2–114 of yeast Sup35 protein transforms in vitro into

amyloid-like filaments. *Proc. Natl. Acad. Sci. USA*. **94**, 6618–6622

27. Gade Malmos, K., Blancas-Mejia, L. M., Weber, B., Buchner, J., Ramirez-Alvarado, M., Naiki, H., and Otzen, D. (2017) ThT 101: a primer on the use of thioflavin T to investigate amyloid formation. *Amyloid*. **24**, 1–16

28. Levine, H. (1993) Thioflavine T interaction with synthetic Alzheimer's disease β-amyloid peptides: Detection of amyloid aggregation in solution: Thioflavine T fluorescence with β/A4 peptides. *Protein Sci.* **2**, 404–410

29. Iadanza, M. G., Jackson, M. P., Hewitt, E. W., Ranson, N. A., and Radford, S. E. (2018) A new era for understanding amyloid structures and disease. *Nat Rev Mol Cell Biol*. **19**, 755–773

30. Glover, J. R., Kowal, A. S., Schirmer, E. C., Patino, M. M., Liu, J.-J., and Lindquist, S. (1997) Self-Seeded Fibers Formed by Sup35, the Protein Determinant of [PSI+], a Heritable Prion-like Factor of S. cerevisiae. *Cell*. **89**, 811–819

31. Wu, H., and Fuxreiter, M. (2016) The Structure and Dynamics of Higher-Order Assemblies: Amyloids, Signalosomes, and Granules. *Cell*. **165**, 1055–1066

32. Murr, M. M., and Morse, D. E. (2005) Fractal intermediates in the self-assembly of silicatein filaments. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 11657–11662

33. Herrera, M. G., Benedini, L. A., Lonez, C., Schilardi, P. L., Hellweg, T., Ruysschaert, J.-M., and Dodero, V. I. (2015) Self-assembly of 33-mer gliadin peptide oligomers. *Soft Matter*. **11**, 8648–8660

34. Hernández, N. E., Hansen, W. A., Zhu, D., Shea, M. E., Khalid, M., Manichev, V., Putnins, M., Chen, M., Dodge, A. G., Yang, L., Marrero-Berríos, I., Banal, M., Rechani, P., Gustafsson, T., Feldman, L. C., Lee, S.-H., Wackett, L. P., Dai, W., and Khare, S. D. (2019) Stimulus-responsive self-assembly of protein-based fractals by computational design. *Nat. Chem.* **11**, 605–614

35. Khatun, S., Singh, A., Maji, S., Maiti, T. K., Pawar, N., and Gupta, A. N. (2020) Fractal self-assembly and aggregation of human amylin. *Soft Matter*. **16**, 3143–3153

36. Otzen, D., and Riek, R. (2019) Functional Amyloids. *Cold Spring Harb Perspect Biol*. **11**, a033860

37. Nishida, K. M., Miyoshi, K., Ogino, A., Miyoshi, T., Siomi, H., and Siomi, M. C. (2013) Roles of R2D2, a Cytoplasmic D2 Body Component, in the Endogenous siRNA Pathway in Drosophila. *Mol. Cell*. **49**, 680–691

38. Palmer, W. H., and Obbard, D. J. (2016) Variation and Evolution in the Glutamine-Rich Repeat Region of *Drosophila* Argonaute-2. *G3*. **6**, 2563–2572

39. Toombs, J. A., McCarty, B. R., and Ross, E. D. (2010) Compositional
Determinants of Prion Formation in Yeast. *Mol Cell Biol*. **30**, 319–332
40. Cascarina, S. M., Paul, K. R., and Ross, E. D. (2017) Manipulating the aggregation
activity of human prion-like proteins. *Prion*. **11**, 323–331
41. Xue, B., Dunbrack, R. L., Williams, R. W., Dunker, A. K., and Uversky, V. N.
(2010) PONDR-FIT: A meta-predictor of intrinsically disordered amino acids.
Biochim. Biophys. Acta - Proteins Proteom. **1804**, 996–1010
42. Gasteiger, E. (2003) ExPASy: the proteomics server for in-depth protein
knowledge and analysis. *Nucleic Acids Res.* **31**, 3784–3788
43. Edelstein, A. D., Tsuchida, M. A., Amodaj, N., Pinkard, H., Vale, R. D., and
Stuurman, N. (2014) Advanced methods of microscope control using μManager
software. *J Biol Methods*. **1**, e10
44. van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D.,
Yager, N., Gouillart, E., and Yu, T. (2014) scikit-image: image processing in
Python. *PeerJ*. **2**, e453
45. Hunter, J. D. (2007) Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**,
90–95

Figure legends

Figure 1. DmAgo2 has a long characteristic N-terminal region

(A) A schematic diagram of the full-length DmAgo2 protein and the sequence of N-ter.

The four conserved domains are shown in gray, while the N-terminal region is shown in

orange. In the N-ter sequence, glutamine and glycine residues are shown in orange. **(B)**

Plot of the prion-like probability of the DmAgo2 sequence predicted by PLAAC (top) and

a heatmap of the degree of disorder predicted by PONDR analysis (bottom).

Figure 2. mCherry-Nter formed SDS-resistant aggregates

SDD-AGE analysis of mCherry-Nter aggregates. mCherry and Sup35NM-mCherry were

used as reference controls. mCherry-tagged polypeptides were detected by western

blotting with anti-RFP polyclonal antibodies. NB, non-boiled fraction; B, boiled fraction.

Figure 3 Nter aggregated in the same manner as common amyloids

(**A**) Fluorescence spectra of ThT in the absence (blue) and presence (orange) of Nter

aggregates.

(**B**) Time-course of the ThT fluorescence intensity in the absence (blue) and presence

(orange) of mCherry-Nter monomers. The solid curves and filled area indicate the mean

value and standard deviation of three repetitive experiments, respectively.

(**C**) Time-course of ThT fluorescence intensity in the absence (blue) and presence

(orange) of mCherry-Nter seeds. The solid curves and filled area indicate the mean value

and standard deviation of three repetitive experiments, respectively.

Figure 4 Fluorescence imaging of the Nter aggregate morphology

(**A–C**) Representative fluorescent images of linear fractal-shaped (**A**), branched

fractal-shaped (**B**) and fibril-shaped aggregates (**C**). Fluorescence signals from mCherry

(left panels, magenta) and ThT (middle panels, green) showed almost identical images.

Merged images are shown in the right panels. Scale bars represent 1 μm (**A**, **B**) and 5 μm

(**C**).

(**D**) Fluorescence intensity profiles of mCherry and ThT along the line in the merged

image of Figure 4C. Intensities were normalized.

(**E**) PCA analysis of the shapes of the Nter aggregates. The plots were colored according

to the classified group (blue, linear fractal-shaped; green, branched fractal-shaped; orange, fibril-shaped).

Figure 5 Fractal-shaped Nter aggregates grew by docking between puncta

(**A–C**) Time-lapse images of the Nter aggregates formation. Ten-minute intervals are shown from left to right. Scale bars represent 5 µm (**A, B**) and 10 µm (**C**).

(**D**) Model of the aggregate formation. Assuming both fibril- and fractal-shaped Nter aggregates grew from a single punctum, the polymerization of Nter monomers from the punctum results in fibril-shaped aggregates, while the fractal-shaped aggregates result from puncta binding to each other (i). The binding of several linear fractal-shaped aggregates produces branched fractal-shaped aggregates (ii). The branched fractal-shaped aggregate becomes bigger by docking with other fractal-shaped aggregates (iii).

Figure 1

A

14 S A A A P Q P Q Q Q Q K Q Q Q Q R Q Q Q P Q Q L Q Q P Q Q L Q Q P Q Q Q Q Q Q Q P H Q Q Q Q
Q S S R Q Q P S T S S G G S R A S G F Q Q G G Q Q Q K S Q D A E G W T A Q K K Q G K Q Q V Q
G W T K Q G Q Q G G H Q Q G R Q G Q D G G Y Q Q R P P G Q Q Q G G H Q Q G R Q G Q E G G Y Q
Q R P P G Q Q Q G G H Q Q G R Q G Q E G G Y Q Q R P S G Q Q Q G G H Q Q G R Q G Q E G G Y Q
Q R P P G Q Q Q G G H Q Q G R Q G Q E G G Y Q Q R P S G Q Q Q G G H Q Q G R Q G Q E G G Y Q
Q R P P G Q Q Q G G H Q Q G R Q G Q E G G Y Q Q R P P G Q Q Q G G H E Q G R Q G Q E G G Y Q
Q R P S G Q Q Q G G H Q Q G R Q G Q E G G Y Q Q R P S G Q Q Q G G H Q Q G R Q G Q E G G Y Q
Q R P S G Q Q Q G G H Q Q G R Q G Q E G G Y Q Q R P P G Q Q P N Q T Q S Q G Q Y Q S R G P P
Q Q Q Q A A P 386

DmAgo2 — Gln- and Gly-rich region — N — PAZ — MID — PIWI — 1208 residues

14   386   602   711   879   1180

B

Figure 2

mCherry | Sup35NM-mCherry | mCherry-Nter

NB B   NB B   NB B

Amyloid fraction

Soluble fraction

1   2   3   4   5   6

Figure 3

Figure 4

Figure 5

A 290 min 300 310 320 330 340 350

B 250 min 260 270 280
   290 300 310 320

C 130 min 140 150 160 170
   180 190 200 210 220

D

puncta to fibril-shaped aggregates   puncta to fractal-shaped aggregates

(i)   (ii)   (iii)