Comparative genomics of *Streptococcus oralis* identifies large scale homologous

recombination and a genetic variant associated with infection

Luke R. Joyce[1]*, Madison A. Youngblom[2,3]*, Harshini Cormaty[1], Evelyn Gartstein[1], Katie E.

Barber[4], Ronda L. Akins[5], Caitlin S. Pepperell[3,6#], and Kelli L. Palmer[1#]

*Authors contributed equally, listed in alphabetical order

[1] Department of Biological Sciences, The University of Texas at Dallas, Richardson, TX, 75080

[2] Microbiology Doctoral Training Program, University of Wisconsin-Madison

[3] Department of Medical Microbiology and Immunology, School of Medicine and Public Health,

University of Wisconsin-Madison, Madison, WI, 53706

[4] Department of Pharmacy Practice, University of Mississippi School of Pharmacy, The

University of Mississippi, Jackson, MS, 38677

[5] Methodist Charlton Medical Center, Dallas, TX, 75237

[6] Department of Medicine (Infectious Diseases), School of Medicine and Public Health,

University of Wisconsin-Madison, Madison, WI, 53706

Running title: Genomic comparison of *Streptococcus oralis.*

- Corresponding authors

Kelli L. Palmer: Kelli.Palmer@UTDallas.edu

Caitlin S. Pepperell: cspepper@medicine.wisc.edu

Abstract

The viridans group streptococci (VGS) are a large consortium of commensal streptococci that colonize the human body. Many species within this group are opportunistic pathogens causing bacteremia and infective endocarditis (IE), yet little is known about why some strains cause invasive disease. Identification of virulence determinants is complicated by the difficulty of distinguishing between the closely related species of this group. Here, we analyzed genomic data from VGS isolated from patient blood cultures with invasive infections and from oral swabs of healthy volunteers and determined the best performing methods for species identification. Using whole-genome sequence data, we characterized the population structure of a diverse sample of *Streptococcus oralis* isolates and found evidence of frequent recombination. We used multiple genome-wide association study tools to identify candidate determinants of invasiveness. These tools gave consistent results, leading to the discovery of a single synonymous single nucleotide polymorphism (SNP) that was significantly associated with invasiveness. This SNP is within a previously undescribed gene that is conserved across the majority of VGS species. Using growth in the presence of human serum and a simulated infective endocarditis vegetation model, we were unable to identify a phenotype for the enriched allele in laboratory assays, suggesting a phenotype may be specific to natural infection. These data highlight the power of analyzing natural populations for gaining insight into pathogenicity, particularly for organisms with complex population structures like the VGS.

Importance

The viridians group streptococci (VGS) are a large collection of closely related commensal streptococci, with many being opportunistic pathogens causing invasive diseases such as

bacteremia and infective endocarditis. Little is known about virulence determinants in these species, and there is a distinct lack of genomic information available for the VGS. In this study, we collected VGS isolates from invasive infections and healthy volunteers and performed whole genome sequencing for a suite of downstream analyses. We focused on a diverse sample of *Streptococcus oralis* genomes and identified high rates of recombination in the population as well as a single genome variant highly enriched in invasive isolates. The variant lies within a previously uncharacterized gene, *nrdM*, which shares homology with the anaerobic ribonucleoside triphosphate reductase, *nrdD*, and is highly conserved among VGS. This work increases our knowledge of VGS genomics and indicates that differences in virulence potential among *S. oralis* isolates is, at least in part, genetically determined.

Introduction

The viridans group streptococci (VGS) comprise a diverse collection of alpha and non-hemolytic streptococci that inhabit the oral cavity and gastrointestinal and genitourinary tracts of healthy humans (*1*). VGS are also associated with invasive disease, particularly in immunocompromised hosts, and are estimated to cause ~23% of Gram-positive bacteremia (*2*, *3*) and ~17% of infective endocarditis (IE) cases (*4*). Bacterial determinants of invasiveness among the VGS are not well understood. Research in this area is hampered by the fact that the specific species of VGS causing bacteremia and IE are infrequently determined in a clinical context due to lack of resolution of existing diagnostic microbiological tools (*5*, *6*). VITEK®2 and MicroScan allow for general assignment of isolates to VGS, and assignment of a limited number of VGS species specifically. Even a relatively newer technique in clinical diagnostics, matrix assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF MS), fails to resolve certain

VGS species, including *Streptococcus mitis* and *Streptococcus oralis* (*7*). 16S rRNA gene

sequencing, multilocus sequence analysis and other genotyping schemes (*5–10*), and GyrB

typing (*11*) are commonly used methods for molecular VGS identification but are not generally

employed in clinical laboratories.

Retrospective studies using molecular approaches have determined that in addition to being

present as oral commensals in healthy individuals, *S. mitis* and *S. oralis* stand out among VGS

as major causative agents of VGS bacteremia and IE (*8, 12–15*)**.** *S. mitis* and *S. oralis* are

members of the mitis group streptococci, a subgroup within the VGS that is closely related to the

major human pathogen *Streptococcus pneumoniae* (*16*, *17*). A recent study in oncology patients

demonstrated that *S. mitis* (58%) and *S. oralis* (19%) were the most frequently identified species

in VGS infections over ~1.5 years (*8*). While *S. mitis* and *S. oralis* are a significant burden on

immunocompromised patients, the mechanisms of virulence within these species have not been

fully elucidated. More specifically, it is not known whether all members of these species have

equal pathogenic potential, or whether some strains have a higher propensity for causing

invasive disease than others.

In this study, we aimed to investigate the mechanisms of invasiveness among VGS by

characterizing species diversity of presumptive VGS obtained clinically from bacteremia and

endocarditis patients. Clinical isolate genomes obtained in this study were supplemented by

existing genome sequences with curated metadata in public databases and genome sequences

of oral isolates collected from healthy volunteers. Our results support metagenomic sequence

binning as a high-resolution tool for differentiating species within the VGS grouping. After a

preliminary analysis in which we determined species designations for clinical isolates that were

diagnosed as VGS, we focused our study on *S. oralis* as a prominent cause of invasive infection.

Within a large sample of *S. oralis* isolates from all three described sub-species (*18*) (*subsp.*

dentisani, *subsp. tigurinus*, and *subsp. oralis*) we found high levels of diversity and strikingly high

recombination rates. We used multiple genome-wide association study (GWAS) methods to test

the hypothesis that specific genetic variations are associated with invasive infection (when

compared to commensal isolates) among *S. oralis* isolates. We discovered a SNP in a previously

uncharacterized gene that is significantly enriched in invasive isolates compared to non-invasive

isolates*.* The contribution of this novel locus to growth with human serum and in a simulated

infective endocarditis vegetation model (*19*) was assessed, although we were unable to identify

a phenotype for a gene knockout or for either allele of the significant variant under the conditions

tested. This work has 1) increased the genomic information available for VGS strains, 2)

described population structure and large-scale homologous recombination within the *S. oralis*

species, and 3) provided evidence that the propensity for virulence in *S. oralis* is at least in part

genetically determined.

Results

GyrB typing and Kraken efficiently speciate *S. mitis* and *S. oralis*.

VGS isolates from Dallas, Texas and Jackson, Mississippi areas were collected from clinically

confirmed VGS bacteremia and infective endocarditis patients. Isolates were initially

characterized using either VITEK®2 or MicroScan platforms. A total of 66 clinical isolates were

successfully sub-cultured in the laboratory, however two isolates were incorrectly identified as

Streptococcus spp.: one *Enterococcus faecalis*, and one *Aerococcus urinae* strain, resulting in

64 presumptive VGS strains collected (Table S1). To compare isolates causing invasive disease

against commensal isolates, healthy volunteers were recruited for oral swab collection on the

University of Texas at Dallas campus, for a total of 81 VGS isolates (Table S1).

As has previously been reported, clinical methods of VGS species identification (VITEK®2 and

MicroScan) were not effective for delineating closely related species, particularly within the mitis

group; however, they were almost always correct in identifying an isolate as a part of the VGS

(Table S1). The taxonomy of the clinical and oral isolates was analyzed by 16S rRNA gene

sequencing, GyrB typing (11), and analysis of Illumina whole genome sequencing (WGS) data

using Kraken (20). While we expected that WGS would provide the clearest results, we also

wanted to assess the utility of the GyrB typing scheme for a diverse set of VGS species, as this

method is cheaper, faster and more feasible for researchers without computational experience

and resources. Using these three different methods of species identification, we classified

isolates into VGS groups using the taxonomy described by Facklam (21). All six of the major

VGS groups (1) were represented in our sample of 81 isolates, with two isolates not fitting into

any of these six groups (Figure 1A). 16S rRNA sequencing was frequently unable to provide

resolution to the species level but was often able to identify which group the isolate belonged to.

GyrB typing (11) and Kraken (20) were both effective in distinguishing *S. oralis* and *S. mitis*,

however GyrB typing is not generally accurate for other VGS groups (Figure 1A).

One downside of species identification with Kraken is it seems unable to distinguish between

anginosus group isolates: *S. anginosus* and *S. constellatus* are species within the anginosus

group (also known as the milleri group) of the VGS and are recognized as abscess-causing

bacteria, with more recent data suggesting their emergence as uropathogens (*1*, *22*, *23*). Kraken

identified 19 *S. anginosus* isolates, and via GyrB typing a total of 20 anginosus group isolates

were identified (12 *S. anginosus* and 8 *S. constellatus*). These data suggest that *S. mitis* and *S.*

oralis are accurately speciated with either Kraken or GyrB typing, yet anginosus group species

may be better identified by GyrB typing.

One drawback to species identification with GyrB typing is the current scheme is limited in the

species it can distinguish (Figure 1A). To further assess the relative functionality of these two

methods we inferred a phylogenetic tree from GyrB sequences and mapped species as defined

by Kraken onto the tree (Figure 1B). The added benefit of a phylogeny of GyrB sequences is the

visualization of additional species not included in the typing scheme (e.g. species in the bovis

group; Figure 1B) which when combined with Kraken output allowed us to confidently assign

species to most isolates. There was only a single instance where Kraken may have mis-identified

the species: an isolate identified by Kraken as *S. cristatus* (a member of the mitis group) appears

to cluster with *S. sanguinis* isolates (of the sanguinis group) on the GyrB phylogeny (Figure 1B).

Additionally, combining data from Kraken and GyrB typing showed that 2 out of the 3 isolates

identified as *S. sp. oral taxon 431* by Kraken were identified as *S. infantis* (a member of the mitis

group) by GyrB typing, and all 3 cluster together on the GyrB phylogeny, indicating that *S. infantis*

is likely the correct designation (Figure 1B).

Overall, our data show that closely related *S. mitis* and *S. oralis* can be speciated accurately via

either Kraken or GyrB typing; however, anginosus group isolates are better distinguished by

GyrB typing. For datasets suspected to contain a diverse sample of different VGS species,

species identification using Kraken with whole genome sequence data appears the most robust.

GyrB typing is an accurate method for making distinctions within species subsets, and a

phylogeny of GyrB sequences may also provide additional information. Our results support

previous assertions that 16S rRNA sequencing is not an effective method for distinguishing the

closely related species of the VGS (1).

When we separated our sample into isolates from blood cultures and those from the mouths of

healthy persons, we saw that invasive isolates spanned all 6 groups while isolates from oral

isolates were less diverse and only originated from the mitis and salivarius groups (Figure 1C).

The lack of diversity in the oral isolates is due to the use of Mitis-Salivarius Agar which inhibits

Gram-negative and most Gram-positive bacterial growth due to the presence of inhibitory

nutrients; however, *S. mitis, S. salivarius,* and enterococci will still grow and produce different

colony morphologies to allow preferential selection (24). It has been reported that *S. mitis* is the

predominant species found in healthy oral microbiomes (25–30) with other commonly found VGS

species including *S. oralis* (27, 28), *S. sanguinis* (28, 29), and *S. salivarius* (29).

High diversity among commensal and invasive *S. oralis* strains

S. oralis was the predominant single species (20/81 isolates) in our sample, and so to identify

possible genetic variants associated with invasive infection, we gathered a larger sample of *S.*

oralis. We obtained all isolates labeled as *S. oralis* and *S. mitis* from NCBI (see Methods for

inclusion criteria) and used Kraken, a core genome phylogeny, and accessory gene content to

confirm *S. oralis* isolates and identify mislabeled isolates. We found that a core genome

alignment produced by Roary (31) effectively delineates *S. oralis* from *S. mitis*, as do patterns

of accessory gene content (Figure S1). We were able to identify 11 isolates mis-labeled as *S.*

mitis within NCBI databases, which we identified as *S. oralis* using these methods (Table S2).

We ended up with a total of 108 *S. oralis* isolates: 57 oral commensals and 51 from invasive

infections (including bacteremia and IE) that we refer to as "oral" and "blood" isolates,

respectively (Table S2). After assembly and annotation of all genomes, we performed a

pangenome analysis with Roary and found that our sample has a core genome of only 801

genes – with an average genome size of 1898 genes. This means that a significant proportion

of genes (57%) encoded by an individual isolate are variable accessory genes (Figure 2A). The

accessory gene content in our sample of *S. oralis* isolates is highly diverse, with 88% of all genes

found in our sample at a frequency of 14% or less, and > 19,000 genes identified in the pan

genome (Figure 2A). This high level of gene content diversity is mirrored in the core genome

phylogeny, which has long terminal branch lengths indicating high variability between core

genome sequences (Figure 2B). Isolates from commensal and invasive sources are

interspersed on the phylogeny and do not generally form monophyletic clades, nor are the

commensal or invasive phenotypes associated any sub-clades within the phylogeny (Figure 2B).

We identified three sub-clades within our core genome phylogeny (Figure 2B) that we found to

correspond with the three previously described sub-species of *S. oralis*: *subsp. dentisani, subsp.*

tigurinus, and *subsp. oralis* (*18*) by cross-referencing our tree with the subspecies of some of

the publicly available isolates in our dataset (Table S2). Average nucleotide identity (ANI) values

for core genome sequences within each subspecies show that *subsp. oralis* is the most

conserved, followed by *subsp. tigurinus* and *subsp. dentisani* (Figure 2C). *Subsp. dentisani* is

unique in that it has ANI values resembling those measured between isolates of different

subspecies – this indicates that levels of diversity within this subspecies are similar to those

found between subspecies (Figure 2C). Following the trends we identified in core genome

diversity levels, the pangenome of *subsp. oralis* isolates appears more conserved (more core

genes, fewer accessory genes) than either *subsp. tigurinus* or *subsp. dentisani* (Figure 2D).

Higher core genome and pangenome diversity levels among *subsp. tigurinus* and *subsp.*

dentisani isolates could indicate that these isolates have access to more diverse partners for

horizontal gene transfer (HGT).

High levels of recombination among *S. oralis* isolates

Viridans group streptococci (VGS) are known for being naturally competent (*32*) and for evolving

rapidly via widespread homologous recombination (*33, 34*). We characterized the signatures of

recombination in the core genome of *S. oralis* using Gubbins (*35*) and identified extreme

amounts of recombination: 99.9% of the core genome is within a predicted recombinant fragment

in at least one isolate (Figure 3A). Additionally, we noted that recombinant fragments are not

shared across multiple isolates but are usually present in only a handful of isolates (Figure 3B).

Using ClonalFrameML (*36*) we calculated the *r/m* value - the ratio of SNPs imported via

recombination (r) to those introduced randomly (m) – and found that with an *r/m* of 5.77, genetic

diversity in our sample is ~6x more likely to be introduced via recombination. This is slightly less

than the notoriously recombinogenic *S. pneumoniae* [*r/m* = ~7; (*33*)] but much higher than other

IE causing bacteria such as *S. aureus* [*r/m* = <1; (*37*)].

Genome wide association study reveals variant associated with invasiveness

With a large sample balanced between our phenotypes of interest, we used multiple genome-

wide association study (GWAS) methods to identify genetic variants associated with invasive

disease, which included isolates from both IE and bacteremia patients. We started by looking for

associations between accessory genes and invasiveness using Scoary (38), which yielded no

significant results. This is perhaps not surprising given that accessory gene content is so diverse

in the sample and individual accessory genes are thus unlikely to be shared by a large proportion

of isolates (Figure 1A).

To identify core genome variants associated with invasiveness, we started with a F_{ST} outlier

analysis, which delineates allele frequency differences between sub-populations and identifies

variants with extreme measures of differentiation. We defined sub-populations of our sample as

being of "oral" or "blood" source and used vcflib (https://github.com/vcflib/vcflib) to calculate Weir

and Cockerham's F_{ST} (wcF_{ST}, abbreviated to F_{ST}) for bi-allelic SNPs (n=67,026) in the core

genome alignment (Figure 4A). We identified a single SNP with a significant F_{ST} value, indicating

significant allele frequency differences between oral and blood sub-populations of *S. oralis*

(Figure 4A). The SNP of interest lies within a gene originally annotated as a homologue of *nrdD*,

an anaerobic ribonucleoside triphosphate reductase present in many streptococci, involved in

synthesis of deoxyribonucleotides under anaerobic conditions (39). However, further inspection

of the sequence revealed that the canonical *nrdD* gene was annotated separately in our *S. oralis*

isolates, and that the novel protein was shorter than *nrdD*. The novel protein does share some

sequence features with *nrdD* including an ATP cone domain, and so we will refer to the novel

locus as *nrdM*. Given the amount of recombination present in our sample (Figure 3A) we

validated the results of our F_{ST} outlier analysis using GWAS methods specifically designed for

use with microbial genomes. We used two programs: treeWAS (*40*), which corrects for the

presence of recombination, and BugWAS (*41*), which identifies lineage effects and controls for

population structure. Results from both of these tools replicated the results of our F_{ST} outlier

analysis with both methods returning a single variant that was significantly associated with the

phenotype: the same SNP in *nrdM* (Figure S2).

The SNP of interest in *nrdM* is a synonymous mutation (I78I), and mapping the nucleotide alleles

of the mutation (C233T) onto the core genome phylogeny shows interspersal of alleles on the

tree, with some structure in *subsp. dentisani* and *subsp. tigurinus* (Figure S3). Most interesting

is the striking association between isolates from invasive infections and the *nrdM* allele: out of

51 invasive isolates in our sample, 39 encode the *nrdM* SNP associated with invasiveness

(Figure S3; Table S2). A multiple sequence alignment of NrdM from our sample of *S. oralis* strains

reveals high protein sequence conservation (Figure S4), and so to determine whether *nrdM* is

more or less conserved than other core genes we calculated the average πN/πS (piN/piS) values

for each core gene (n=801). The distribution of gene-wise values of piN/piS indicate the majority

of genes are evolving under relatively strong purifying selection, including *nrdM*, which had a

piN/piS value slightly below the mean (Figure 4B). This is not surprising given that recombination

strengthens the efficiency of selection by enabling rapid removal of deleterious mutations (*42*).

Selection on *nrdM* variant

Given the strong association between the *nrdM* variant and invasiveness, we wondered if this

SNP is under positive selection in our sample. One method for identifying positive selection is

the identification of homoplastic mutations, *i.e*. mutations that arose more than once on the

phylogeny, which we have used previously to screen for drug resistance loci in *Mycobacterium*

tuberculosis (*43*). We used TreeTime (*44*) to identify homoplastic mutations in our sample and

found numerous homoplastic mutations – this is expected since intergenomic recombination can

produce homoplasies by lateral transfer of sequence variants. The variant in *nrdM* arose 20

times on the phylogeny, which was in the 85th percentile of mutation multiplicity (Figure S5). This

finding, along with gene-wise piN/piS values (Figure 4B), suggests that selection pressures are

similar at this locus to others in the genome; however, further analysis with a larger sample size

could elucidate more subtle signs of selection.

nrdM is conserved among VGS species

In order to investigate the presence of *nrdM* homologues in other *Streptococcus* spp. we used

Blast to search annotated genes in all newly sequenced isolates and found *nrdM* to be present

in all VGS species in our original sample, except for two bovis group species: *S. lutetiensis* and

S. pasteuranius (Figure 5A). Additionally, the position of the variant associated with invasiveness

in *S. oralis* is conserved, and we were able to identify variation in *nrdM* alleles between different

species. As in *S. oralis*, NrdM is highly conserved at the protein level across the VGS species in

our sample, indicating the gene may have a conserved function across multiple species (Figure

5B).

Deletion of *nrdM* does not affect *in vitro* growth

To investigate the impact the *nrdM* orphan allele has on growth in the two infective endocarditis

strains *S. oralis* 1648 (SO48) and mitis group *Streptococcus* 1643 (SM43), we deleted *nrdM*,

locus tags MP387_03665 and FD735_06230, respectively, from each strain, via homologous

recombination (Table 1). SO48 possesses the *nrdM*-C ("C" = commensal) allele, and SM43

possesses the *nrdM*-I ("I" = invasive) allele. Knockout mutants were confirmed via Sanger

sequencing. No difference in growth was observed for knockout mutants compared to wild-type

in Todd-Hewitt broth (THB) (Figure S6). Further, *nrdM* allele swaps from SO48 and SM43 were

placed back into the genome of SO48Δ*nrdM* and SM43Δ*nrdM* to generate Δ*nrdM*::*nrdM*-I or

Δ*nrdM*::*nrdM*-C strains in each strain background (Table 1). Since the presence of human serum

has been shown to impact mitis group streptococci physiology (*45*, *46*), we assessed the impact

of *nrdM* during growth with human serum by performing growth curves of allele replacement

strains in chemically defined medium supplemented with 5% v/v human serum (Figure S6). No

difference in growth profile was observed between alleles in either strain background. Thus, the

nrdM locus does not impact growth in rich laboratory medium or chemically defined medium

supplemented with 5% v/v human serum *in vitro*.

***nrdM* has no impact on growth in a simulated infective endocarditis vegetation model**

Finally, the allele swapped strains were investigated using the pharmacological *in vitro* simulated

infective endocarditis vegetation model (SIEVM). We reasoned that if *nrdM*-I confers enhanced

fitness in simulated endocardial vegetations, we would observe significantly different vegetation

CFUs for SO48Δ*nrdM*::*nrdM*-I versus SO48Δ*nrdM*::*nrdM*-C. The SM43 strain was similarly

tested, to assess the effect of *nrdM* allele swapping in a closely related but different (i.e. non-

oralis) genetic background. Strains were inoculated into vegetation clots at ~ 1 x10^8 CFU,

incubated in the chemostat model with Mueller Hinton Broth (MHB) in parallel for 48hrs (see

Material and Methods). At 4, 24, and 48 h post inoculation, 4 clots from each model were

removed for CFU enumeration (Figure 6). No significant difference was observed during growth

in the SIEVM between *nrdM* allele swap in either strain background. Ultimately, these data and

the growth curve data together show that under these *in vitro* conditions, *nrdM* is not essential,

and no phenotype could be assigned to either of the *nrdM* alleles.

Discussion

The viridans group streptococci (VGS) are a large collection of closely related *Streptococcus*

spp. that inhabit the oral cavity and gastrointestinal and genitourinary tracts of humans as

commensals, but can invade other tissues to cause severe diseases such as bacteremia and

infective endocarditis (*1*). Our understanding of the pathogenesis of VGS disease is limited by

lack of knowledge surrounding the genetic and environmental conditions that facilitate a switch

from commensalism to pathogen. Additionally, clinical detection and identification of bacterial

infections is critical for patient care and recovery, especially in neutropenic and

immunocompromised patients. A major etiological agent of disease in this patient demographic

are the VGS, especially *S. mitis* and *S. oralis.* Yet, clinical methods of speciation are still largely

inaccurate, and even newer methods like MALDI-TOF MS struggle to differentiate between

closely related species such as *S. oralis* and *S. mitis* (*7*). In this study, we collected and whole

genome-sequenced a variety of invasive and non-invasive VGS isolates, compared

bioinformatic methods for delineating closely related species, and identified a SNP that may

predispose certain isolates of *S. oralis* to invasive disease.

Complex population structure of *S. oralis*

We characterized the population structure of 81 *S. oralis* isolated from healthy oral microbiota

and from blood, and found high levels of diversity, both in the core (Figure 2B-C) and accessory

genomes (Figure 2A,D). We found that using our methods, previously described sub-species of

S. oralis corresponded to sub-clades we identified in a phylogeny inferred from core genome

sequences (Table S2). We also identified *subsp. oralis* as being the least diverse of the three

subspecies, with respect to both gene content and sequence variation in the core genome

(Figure 2). We assert that recombination is likely the primary mechanism for generating and

maintaining diversity in this species, as 99.9% of the *S. oralis* core genome has been affected

by recombination (Figure 3). The rarity of a given recombinant fragment in our sample also

indicates that *S. oralis* isolates are likely participating in HGT with diverse species – made

possible by the fact that they inhabit complex communities. We observed a mean piN/piS value

across *S. oralis* core genes of 0.039 which is an order of magnitude lower than comparator

species such as *S. aureus* for which values of 0.32 (*47*) and 0.55 (*48*) have been reported, and

lung-colonizing *P. aeruginosa* from cystic-fibrosis patients that have a mean of 0.14. (*49*). This

indicates that *S. oralis* core genes are under strong purifying selection. We recently identified a

similar phenomenon among environmental isolates of *Mycobacterium abscessus* that are also

highly recombinogenic and manifest high amounts of synonymous diversity in their core

genomes (*50*). This suggests that like *M. abscessus*, *S. oralis* inhabits environments alongside

diverse microbial species where high rates of recombination enable ready acquisition of novel

genetic material and rapid removal of deleterious mutations.

Switching from commensal to pathogen

Our identification of extremely high recombination rates within *S. oralis* points to the unique

genomic features characterizing this VGS species. High rates of recombination allow the species

to adapt to fluctuating environments encountered within the human body, and potentially enable

invasion of new niches, including pathogenic niches. We used three different GWAS methods

that replicably identified a strong association between a synonymous SNP within an undescribed

protein, NrdM, and invasive disease (Figure 4, Figure S2). The association was robust across

diverse isolates of all three subspecies (Figure S3). We also found that *nrdM* was conserved

among *S. oralis* isolates (Figure S4) and is one of only ~440 genes conserved across all VGS

species in our sample indicating that it is likely to play an important, and perhaps a similar, role

across species. Utilizing *in vitro* growth in the presence of human serum and a simulated

infective endocarditis vegetation model, we compared *nrdM* knockout strains and allele-

swapped invasive and commensal isolate alleles in two strain backgrounds. No phenotype was

observed under the tested conditions, suggesting that if NrdM is contributing to invasive disease,

it requires different conditions (possibly specific to the *in vivo* environment) for a potential

phenotype to be observed. NrdD, which shares the same ATPase cone domain as NrdM, is

required for anaerobic growth in *S. sanguinis* (*39*), and, notably, disruption of this gene results

in attenuated virulence (*51*). Long thought to have neutral fitness effects, synonymous mutations

are increasingly recognized as having significant effects on bacterial fitness, for example by

impacting gene expression and protein folding (*52–56*). We believe that further study of this

synonymous variant could reveal fitness effects and that adaptation within *nrdM* may affect a

multitude of phenotypes that aid in transition from the oral cavity to the bloodstream, although

further investigation is necessary.

A major hurdle to successful genomic analyses in VGS is the correct classification of the different

species. It is possible that incorrect species identification has contributed to the lack of clear

results from studies looking at the genetic determinants of virulence in VGS species. One such

study by Rasmussen *et. al.* used a sample of both *S. mitis* and *S. oralis* to search for known

virulence factors, which they found at varying frequencies, indicating that genetic differences

may be responsible for variability in virulence among strains (*57*). Similar to our sample of *S.*

oralis isolates (Figure 2B), studies of the closely related VGS *S. sanguinis* and *S. gordonii* were

unable to identify phylogenetic patterns based on invasive disease (*58, 59*). These studies,

however, were unable to identify specific genomic variants associated with invasiveness. What

our work and the work of others indicates, is that virulence properties differ between VGS

species, despite being so closely related and often causing similar disease. This is further

illustrated by the fact that although *nrdM* is conserved in *S. mitis,* the association between the

invasive allele identified in *S. oralis* and pathogenicity in *S. mitis* is not significant (data not

shown). Going forward, analyses using larger samples of well-defined, individual species will

provide better resolution for identifying the genetic determinants of virulence in the VGS.

Materials and methods

Collection of clinical Isolates

Clinical strains obtained from routine blood cultures and identified as viridans group streptococci

were stored in the microbiology laboratory at each hospital (Methodist Health System [MHS] and

University of Mississippi Medical Center [UMMC]). Respective site investigators reviewed patient

clinical information to confirm invasive infection (i.e., bacteremia or endocarditis), then de-

identified specimens and provided blinded isolates to the University of Texas at Dallas (UTD)

laboratory for study. Specimens were not included for further analysis if the isolate was deemed

a contaminant and did not require antibiotic therapy as determined by the treating physician. Site

investigators obtained approval from respective Institutional Review Boards (MHS UTD IRB 18-

121 and UMMC 2018-0068).

Processing of clinical isolates

Clinical isolates were struck on Mitis-Salivarius Agar (MSA) (BD Bacto and incubated overnight

in 37˚C and 5% CO_2. MSA plates were observed for homogenous colony morphology, and a

single colony was inoculated into 10 mL THB for overnight growth at 37˚C and 5% CO_2. If more

than one colony morphology was identified on MSA plates, broth cultures were made from each

colony morphology. Overnight cultures were stored at -80°C in 25% glycerol. The remaining

culture volume was pelleted at 4,280 x g in a Sorvall RC6+ floor centrifuge and genomic DNA

was extracted using Qiagen DNeasy Blood and Tissue Kit per manufacturer protocols, with

minor modifications as described in (*6*).

Collection of oral swab samples

Oral swabs were obtained from healthy adult volunteers at the UTD campus (UTD IRB 17-170).

The following exclusion criteria were applied: previous history of bacteremia or endocarditis;

recent antibiotic exposure (prior 30 days); history of periodontal disease; personal or family

history of immunocompromise. No participants were excluded based on these criteria. The

volunteer was asked to rinse their mouth with sterile saline, and then self-swab their teeth and

tongue with a sterile swab (Puritan). Swabs were stored at 4˚C until processing.

Processing of oral swabs

Oral swabs were processed as described above. Briefly, swabs were struck onto MSA plates

and grown overnight. After incubation, the MSA plates were observed and colony morphologies

consistent with *Streptococcus mitis* and *S. oralis* were selected for overnight growth in THB. In

addition, approximately 3 random colonies of different morphologies were also selected for

overnight growth. Cultures were processed as described above.

16S rRNA sequence analysis and GyrB typing

PCR reactions were performed using Taq polymerase (New England Biolabs) with primer

sequences in Table S3. 16S rRNA genes were amplified using universal primers 8F and 1492R

(*60*). The DNA Gyrase B gene (*gyrB*) was amplified using previously reported primers (Table

S3). PCR reactions were analyzed by agarose gel electrophoresis and purified using the

GeneJET PCR purification kit (Thermo Fisher) per the manufacturer protocols. Products were

sequenced at the Massachusetts General Hospital DNA Core. 16S rRNA sequences were

trimmed using Geneious R11 (https://www.geneious.com) allowing a maximum of 10 low quality

bases and 6 ambiguities. Trimmed sequences were used as queries for NCBI Blastn against the

16S ribosomal RNA sequences (Bacteria and Archaea) database and species were assigned

only when the forward and reverse sequencing reactions had the same top Blastn result (Table

S1). GyrB nucleotide sequences were translated, and amino acid sequences pairwise aligned

to *Streptococcus mitis* ATCC 49456 GyrB (locus tag SM12261_0755). Amino acid variations

were identified using the method of Galloway-Peña et al. (*11*).

Ilumina sequencing

Sequencing was performed at The University of Texas at Dallas Genome Core using Illumina

Nextseq 500 platform with a mid-output 300 cycle of 2 x 75 bp paired-end reads for clinical

isolates S1-S31, or 2 x 150 bp paired-end reads for all other isolates.

Genome assembly & annotation

Using the raw sequencing data from newly sequenced clinical and oral isolates (Table S1),

species identification was additionally performed using Kraken2 (*20*) Raw data was quality

checked and trimmed using FastQC v0.11.8 (*61*) and TrimGalore v0.6.4,

(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore), respectively. Contigs were

assembled using SPAdes v3.13.0 with default parameters (*62*). Assemblies were checked for

quality using Quast v5.0.2 (*63*) filtering out contigs shorter than 500bp or with coverage lower

than 5x, as well as confirming all assemblies had a N50 > 50,000bp. Contigs were annotated

using Prokka v1.13.3 (*64*) before a pangenome analysis was performed with Roary v3.12.0

using a blastp identity threshold of 75% (*31*). Using a nucleotide sequence alignment of GyrB

(as clustered by Roary) a phylogenetic tree was made using FastTree v2.1.9 (*65*) and visualized

in R with ggtree (*66*). GyrB typing was confirmed using these sequences with a custom script

(code available at https://github.com/myoungblom/VGS_GWAS.git) assigning species based on

the scheme outlined by Galloway-Peña et al. (*11*). *nrdM* sequences from all newly sequenced

VGS isolates were identified with Blastp using the *S. oralis* NrdM amino acid sequence as the

query sequence.

S. oralis **genome collection for GWAS analyses**

From our sample of clinical and oral isolates, *S. oralis* made up the largest part of our sample

so we decided to proceed with analyses of just this species. To create a dataset large enough

for a powerful genome-wide association study (GWAS) we identified all *S. oralis* and *S. mitis*

isolates from NCBI (NCBI SRA & Assembly databases accessed June 2019) with the proper

metadata indicating they were isolated from the mouth (e.g oral cavity, dental plaque, dental

biofilm, etc) or from blood (e.g. infective endocarditis, bloodstream infection, blood, etc). We

assumed isolates uploaded to NCBI with various "oral" sources were all commensal and all those

from "blood" were from an invasive infection (Table S2). We chose to start with both *S. oralis*

and *S. mitis* because these species are so closely related, they are often mistaken for each other

and uploaded to NCBI under the wrong species (Table S2) as has previously been reported (*10*).

We pulled out the true *S. oralis* isolates using Kraken and GyrB typing as described above.

Samples for which raw sequence data was available were assembled as described above, and

then annotated with the remainder of assemblies downloaded from NCBI (Table S2). Note that

since the time of data collection, three of the assemblies used in this dataset have been

suppressed (Table S2). We then performed a pangenome analysis on the *S. oralis* sample as

described above (using a Blastp identity threshold of 95% and Prank to align core genes) and a

phylogenetic tree was inferred from the resulting core genome alignment using RAxML v8.2.3

(*67*) and visualized in R with ggtree (*66*).

Recombination analyses

We identified recombinant fragments in the *S. oralis* core genome using Gubbins v2.4.1 with

default parameters (*35*). Recombinant fragments were visualized alongside the core genome

phylogeny using Phandango (*68*). We used ClonalFrameML v1.11 (*36*) with default parameters

to estimate *r/m*.

Population genetics statistics

Pairwise average nucleotide identity (ANI) values of *S. oralis* core genome sequences were

calculated with OrthoANI (*69*). piN/piS values were calculated for all pairwise combinations for

each *S. oralis* core gene were calculated using Egglib (*70*), then the average piN/piS value for

each gene was calculated.

Rarefaction & accumulation plots

Rarefaction and accumulation curves for the *S. oralis* subspecies were calculated from Roary

gene presence absence files. Briefly, separate pangenome analyses were performed as

described above for each individual subspecies and then each dataset was iteratively

subsampled to the size of the smallest dataset (n=25) and the median number of core and total

genes was plotted from all iterations.

GWAS

We first queried for genetic associations with the 'invasive' phenotype in our dataset by

identifying accessory gene content significantly associated with invasiveness using Scoary

v1.16.6 (*38*). We then performed a preliminary GWAS of core genome variants using an F_{ST}

outlier analysis. Briefly, a VCF file containing all core genome variants was made using SnpSites

v2.0.3 (*71*) and reformatted using a custom script (code available

https://github.com/myoungblom/VGS_GWAS.git). Then we calculated Weir and Cockerham's

F_{ST} for bi-allelic SNPs using vcflib (https://github.com/vcflib/vcflib). Using a custom script (code

available https://github.com/myoungblom/VGS_GWAS.git) we permuted the phenotypes in this

analysis 100x and used the maximum F_{ST} value observed in the null distribution as a cutoff to

identify significant F_{ST} outliers. To validate the results of our F_{ST} outlier analysis we also used

two GWAS programs designed specifically for use with microbial genomes: treeWAS (*40*) that

corrects for the presence of recombination, and BugWAS (*41*) that identifies lineage effects and

controls for population structure. TreeWAS was run using the recombination-adjusted

phylogenetic tree made with Gubbins (see above) using 10x the number of SNPs in the core

genome for the parameter "n.snps.sim". BugWAS was run using default parameters.

Homoplasy analysis

Homoplasy analysis was performed using TreeTime v0.9.0-b.2 (*44*) with default parameters.

Deletion of *nrdM*

Knockout constructs of *nrdM* in SM43 (Locus ID FD735_06230) and SO48 (Locus ID

MP387_03665) were generated as previously described (*45*). Briefly, linear constructs were

generated by amplifying ~2 kb regions upstream and downstream of *nrdM* using Phusion

polymerase (Thermo Fisher) using primers in Table S3. SOEing PCR was used to stitch

fragments together and the amplified product was assessed via agarose gel electrophoresis.

Gel extraction was performed using the QIAQuick Gel Extraction kit (Qiagen). Linear constructs

were transformed by natural transformation as described in (*45*). Transformation plates were

incubated overnight, and putative transformant colonies were screened via PCR for the *nrdM*

deletion.

nrdM allele swaps

Allele swap strains were generated using the same strategy as the deletion, except the linear

construct contained either the _nrdM_-I or _nrdM_-C allele coupled with the flanking regions for the

respective strain. The linear construct was transformed into SM43Δ_nrdM_ and SO48Δ_nrdM._ The

allele swap region in transformants was amplified using primers in Table S3. Products were

sequenced for validation of the allele swap (Massachusetts General Hospital DNA Core).

Growth curves

Growth curves in THB were performed in biological duplicate. Wild-type and Δ_nrdM_ strains were

grown overnight as described, then diluted to OD_{600nm} 0.05 in approximately 12 mL THB. OD_{600nm}

was monitored every hour using a Thermo Scientific Genesys 30 spectrophotometer. For growth

curves in the presence of human serum, biological triplicate overnight cultures were grown in

streptococcal defined medium (_45, 46, 72_) and diluted to OD_{600nm} 0.1 in defined medium

supplemented with 5% v/v human serum (Sigma-Aldrich). OD_{600nm} was monitored at 3, 6 and 24

hrs as described above.

Simulated Infective Endocarditis Vegetation Model (SIEVM)

Strains were inoculated from freezer stocks into 5 mL Mueller Hinton Broth (MHB) (BD Bacto)

and incubated overnight at 37˚C and 5% CO_2. 1 mL was expanded into 100 mL pre-warmed

MHB and incubated overnight as described above. The SIEVM was set up and performed as

previously described (_19_) using human blood products from the American Red Cross (UTD IRB

19MR0160). Briefly, 500 µL of pooled human cryoprecipitate (American Red Cross), 50 µL ~2

TIU/mL aprotinin (Sigma-Aldrich), ~100,000 human platelets (American Red Cross), and 10^8

CFU/g bacteria were combined in a sterile Eppendorf tube and vortexed. Sterile monofilament

line was positioned before 100 µL of ~2 KU/mL high activity bovine thrombin (Sigma-Aldrich)

was added to congeal the vegetation. Vegetations were placed into the glass apparatus in a

37˚C water bath, and MHB was pumped through the model at a pre-calibrated rate of 0.4 mL/min.

Four vegetations were removed at designated time points for each strain, weighed, removed

from the monofilament line and placed in 1.25% Trypsin solution (Sigma-Aldrich) in sterile screw

cap microcentrifuge tubes (Fisher Scientific) with ~5-8 2.7 mm glass beads (BioSpec). Clots

were homogenized for 10-15 min horizontally on a vortex before serial dilution and plating on

THB agar plates for enumeration. CFU/g was calculated by multiplying the observed CFU/mL

by the net weight of the vegetation. SIEVM was performed in biological triplicate for each strain,

with four vegetations per time point per strain.

***S. oralis* 1648 hybrid genome assembly**

Pacific Biosciences single molecule real time (SMRT) sequencing was performed by the Johns

Hopkins Genome Core. The SO48 whole genome was assembled using the Unicyler assembly

pipeline (*73*) combining SMRT long reads generated in this study and Illumina reads previously

generated for SO48 (accession PRJNA354070; (*74*)).

Accession numbers

The SO48 whole genome sequence generated in this study has been deposited in Genbank

under the accession number CP094226. Genome constructs and Illumina and SMRT sequence

reads generated in this study have been deposited in the Sequence Read Archive under the

BioProject accession PRJNA817585, see Table S1.

Acknowledgements

This work was supported by R21 AI130666 from the National Institutes of Health and the Cecil

H. and Ida Green Chair in Systems Biology Science to K.L.P., and by the National Science

Foundation Graduate Research Fellowship to M.A.Y under grant number DGE-1747503. The

content of this work is solely the responsibility of the authors and does not necessarily represent

the official views of the National Institutes of Health.

References

1. C. D. Doern, C.-A. D. Burnham, It's not easy being green: the viridans group streptococci, with a
focus on pediatric clinical manifestations. *J. Clin. Microbiol.* **48**, 3829–3835 (2010).

2. C. Gudiol, M. Bodro, A. Simonetti, F. Tubau, E. González-Barca, M. Cisnal, E. Domingo-Domenech, L.
Jiménez, J. Carratalà, Changing aetiology, clinical features, antimicrobial resistance, and outcomes
of bloodstream infection in neutropenic cancer patients. *Clin Microbiol Infect*. **19**, 474–479 (2013).

3. M. Marín, C. Gudiol, C. Garcia-Vidal, C. Ardanuy, J. Carratalà, Bloodstream infections in patients
with solid tumors: epidemiology, antibiotic therapy, and outcomes in 528 episodes in a single
cancer center. *Medicine (Baltimore)*. **93**, 143–149 (2014).

4. D. R. Murdoch, G. R. Corey, B. Hoen, J. M. Miró, V. G. Fowler, A. S. Bayer, A. W. Karchmer, L.
Olaison, P. A. Pappas, P. Moreillon, S. T. Chambers, V. H. Chu, V. Falcó, D. J. Holland, P. Jones, J. L.
Klein, N. J. Raymond, K. M. Read, M. F. Tripodi, R. Utili, A. Wang, C. W. Woods, C. H. Cabell,
International Collaboration on Endocarditis-Prospective Cohort Study (ICE-PCS) Investigators,
Clinical presentation, etiology, and outcome of infective endocarditis in the 21st century: the
International Collaboration on Endocarditis-Prospective Cohort Study. *Arch Intern Med*. **169**, 463–
473 (2009).

5. C. Teles, A. Smith, G. Ramage, S. Lang, Identification of clinically relevant viridans group
streptococci by phenotypic and genotypic analysis. *Eur J Clin Microbiol Infect Dis*. **30**, 243–250
(2011).

6.   T. Hoshino, T. Fujiwara, M. Kilian, Use of phylogenetic and phenotypic analyses to identify
nonhemolytic streptococci isolated from bacteremic patients. *J Clin Microbiol*. **43**, 6073–6085
(2005).

7.   J. Isaksson, M. Rasmussen, B. Nilson, L. S. Stadler, S. Kurland, L. Olaison, E. Ek, B. Herrmann,
Comparison of species identification of endocarditis associated viridans streptococci using rnpB
genotyping and 2 MALDI-TOF systems. *Diagn Microbiol Infect Dis*. **81**, 240–245 (2015).

8.   S. A. Shelburne, P. Sahasrabhojane, M. Saldana, H. Yao, X. Su, N. Horstmann, E. Thompson, A. R.
Flores, Streptococcus mitis Strains Causing Severe Clinical Disease in Cancer Patients. *Emerg Infect*
Dis. **20**, 762–771 (2014).

9.   C. J. Bishop, D. M. Aanensen, G. E. Jordan, M. Kilian, W. P. Hanage, B. G. Spratt, Assigning strains to
bacterial species via the internet. *BMC Biol*. **7**, 3 (2009).

10. I. M. Velsko, M. S. Perez, V. P. Richards, Resolving Phylogenetic Relationships for Streptococcus
mitis and Streptococcus oralis through Core- and Pan-Genome Analyses. *Genome Biology and*
Evolution. **11**, 1077–1087 (2019).

11. J. Galloway-Peña, P. Sahasrabhojane, J. Tarrand, X. Y. Han, S. A. Shelburne, GyrB Polymorphisms
Accurately Assign Invasive Viridans Group Streptococcal Species. *J Clin Microbiol*. **52**, 2905–2912
(2014).

12. P. Sahasrabhojane, J. Galloway-Peña, L. Velazquez, M. Saldaña, N. Horstmann, J. Tarrand, S. A.
Shelburne, Species-level assessment of the molecular basis of fluoroquinolone resistance among
viridans group streptococci causing bacteraemia in cancer patients. *Int J Antimicrob Agents*. **43**,
558–562 (2014).

13. T. Kitten, C. L. Munro, N. Q. Zollar, S. P. Lee, R. D. Patel, Oral streptococcal bacteremia in
hospitalized patients: taxonomic identification and clinical characterization. *J Clin Microbiol*. **50**,
1039–1042 (2012).

14. S. Chun, H. J. Huh, N. Y. Lee, Species-specific difference in antimicrobial susceptibility among
viridans group streptococci. *Ann Lab Med*. **35**, 205–211 (2015).

15. S. L. Kim, S. M. Gordon, N. K. Shrestha, Distribution of streptococcal groups causing infective
endocarditis: a descriptive study. *Diagnostic Microbiology and Infectious Disease*. **91**, 269–272
(2018).

16. M. Kilian, D. R. Riley, A. Jensen, H. Brüggemann, H. Tettelin, Parallel evolution of Streptococcus
pneumoniae and Streptococcus mitis to pathogenic and mutualistic lifestyles. *mBio*. **5**, e01490-
01414 (2014).

17. M. Kilian, K. Poulsen, T. Blomqvist, L. S. Håvarstein, M. Bek-Thomsen, H. Tettelin, U. B. S. Sørensen,
Evolution of Streptococcus pneumoniae and Its Close Commensal Relatives. *PLoS One*. **3**, e2683
(2008).

18. A. Jensen, C. F. P. Scholz, M. 2016 Kilian, Re-evaluation of the taxonomy of the Mitis group of the
genus Streptococcus based on whole genome phylogenetic analyses, and proposed
reclassification of Streptococcus dentisani as Streptococcus oralis subsp. dentisani comb. nov.,
Streptococcus tigurinus as Streptococcus oralis subsp. tigurinus comb. nov., and Streptococcus
oligofermentans as a later synonym of Streptococcus cristatus. *International Journal of Systematic*
and Evolutionary Microbiology. **66**, 4803–4820.

19. R. L. Akins, B. D. Katz, C. Monahan, D. Alexander, Characterization of High-Level Daptomycin
Resistance in Viridans Group Streptococci Developed upon In Vitro Exposure to Daptomycin.
Antimicrob Agents Chemother. **59**, 2102–2112 (2015).

20. D. E. Wood, J. Lu, B. Langmead, Improved metagenomic analysis with Kraken 2. *Genome Biology*.
20, 257 (2019).

21. R. Facklam, What happened to the streptococci: overview of taxonomic and nomenclature
changes. *Clin Microbiol Rev*. **15**, 613–630 (2002).

22. M. Furuichi, Y. Horikoshi, Sites of infection associated with Streptococcus anginosus group among
children. *J Infect Chemother*. **24**, 99–102 (2018).

23. M. M. Pearce, E. E. Hilt, A. B. Rosenfeld, M. J. Zilliox, K. Thomas-White, C. Fok, S. Kliethermes, P. C.
Schreckenberger, L. Brubaker, X. Gai, A. J. Wolfe, The female urinary microbiome: a comparison of
women with and without urgency urinary incontinence. *mBio*. **5**, e01283-01214 (2014).

24. G. H. Chapman, The isolation and testing of fecal streptococci. *Am J Dig Dis*. **13**, 105–107 (1946).

25. C. Pearce, G. H. Bowden, M. Evans, S. P. Fitzsimmons, J. Johnson, M. J. Sheridan, R. Wientzen, M. F.
Y. 1995 Cole, Identification of pioneer viridans streptococci in the oral cavity of human neonates.
Journal of Medical Microbiology. **42**, 67–72.

26. J. A. Aas, B. J. Paster, L. N. Stokes, I. Olsen, F. E. Dewhirst, Defining the Normal Bacterial Flora of the
Oral Cavity. *Journal of Clinical Microbiology*. **43**, 5721–5732 (2005).

27. D. L. Mager, L. A. Ximenez-Fyvie, A. D. Haffajee, S. S. Socransky, Distribution of selected bacterial
species on intraoral surfaces. *Journal of Clinical Periodontology*. **30**, 644–654 (2003).

28. S. N. Peterson, E. Snesrud, J. Liu, A. C. Ong, M. Kilian, N. J. Schork, W. Bretz, The Dental Plaque
Microbiome in Health and Disease. *PLOS ONE*. **8**, e58487 (2013).

29. E. Caselli, C. Fabbri, M. D'Accolti, I. Soffritti, C. Bassi, S. Mazzacane, M. Franchi, Defining the oral
microbiome by whole-genome sequencing and resistome analysis: the complexity of the healthy
picture. *BMC Microbiology*. **20**, 120 (2020).

30. N. N. Al-hebshi, A. T. Nasher, M. Y. Maryoud, H. E. Homeida, T. Chen, A. M. Idris, N. W. Johnson,
Inflammatory bacteriome featuring Fusobacterium nucleatum and Pseudomonas aeruginosa
identified in association with oral squamous cell carcinoma. *Sci Rep*. **7**, 1834 (2017).

31. A. J. Page, C. A. Cummins, M. Hunt, V. K. Wong, S. Reuter, M. T. G. Holden, M. Fookes, D. Falush, J.
691    A. Keane, J. Parkhill, Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*. **31**,
692    3691–3693 (2015).

693 32. J. Mitchell, Streptococcus mitis: walking the line between commensalism and pathogenesis. *Mol*
694    *Oral Microbiol*. **26**, 89–98 (2011).

695 33. C. Chaguza, J. E. Cornick, D. B. Everett, Mechanisms and impact of genetic recombination in the
696    evolution of Streptococcus pneumoniae. *Computational and Structural Biotechnology Journal*. **13**,
697    241–247 (2015).

698 34. C. P. Andam, W. P. Hanage, Mechanisms of genome evolution of Streptococcus. *Infection, Genetics*
699    *and Evolution*. **33**, 334–342 (2015).

700 35. N. J. Croucher, A. J. Page, T. R. Connor, A. J. Delaney, J. A. Keane, S. D. Bentley, J. Parkhill, S. R.
701    Harris, Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome
702    sequences using Gubbins. *Nucleic Acids Res.* **43**, e15 (2015).

703 36. X. Didelot, D. J. Wilson, ClonalFrameML: efficient inference of recombination in whole bacterial
704    genomes. *PLoS Comput. Biol.* **11**, e1004041 (2015).

705 37. R. G. Everitt, X. Didelot, E. M. Batty, R. R. Miller, K. Knox, B. C. Young, R. Bowden, A. Auton, A.
706    Votintseva, H. Larner-Svensson, J. Charlesworth, T. Golubchik, C. L. C. Ip, H. Godwin, R. Fung, T. E.
707    A. Peto, A. S. Walker, D. W. Crook, D. J. Wilson, Mobile elements drive recombination hotspots in
708    the core genome of Staphylococcus aureus. *Nat Commun*. **5**, 3956 (2014).

709 38. O. Brynildsrud, J. Bohlin, L. Scheffer, V. Eldholm, Rapid scoring of genes in microbial pan-genome-
710    wide association studies with Scoary. *Genome Biology*. **17**, 238 (2016).

711 39. D. V. Rhodes, K. E. Crump, O. Makhlynets, M. Snyder, X. Ge, P. Xu, J. Stubbe, T. Kitten, Genetic
712    Characterization and Role in Virulence of the Ribonucleotide Reductases of Streptococcus
713    sanguinis. *J. Biol. Chem.* **289**, 6273–6287 (2014).

714 40. C. Collins, X. Didelot, A phylogenetic method to perform genome-wide association studies in
715    microbes that accounts for population structure and recombination. *PLoS Comput. Biol.* **14**,
716    e1005958 (2018).

717 41. S. G. Earle, C.-H. Wu, J. Charlesworth, N. Stoesser, N. C. Gordon, T. M. Walker, C. C. A. Spencer, Z.
718    Iqbal, D. A. Clifton, K. L. Hopkins, N. Woodford, E. G. Smith, N. Ismail, M. J. Llewelyn, T. E. Peto, D.
719    W. Crook, G. McVean, A. S. Walker, D. J. Wilson, Identifying lineage effects when controlling for
720    population structure improves power in bacterial association studies. *Nature Microbiology*. **1**,
721    16041 (2016).

722 42. M. Nordborg, B. Charlesworth, D. Charlesworth, The effect of recombination on background
723    selection. *Genet Res*. **67**, 159–174 (1996).

43. T. D. Mortimer, A. M. Weber, C. S. Pepperell, Signatures of Selection at Drug Resistance Loci in
Mycobacterium tuberculosis. *mSystems*. **3**, e00108-17 (2018).

44. P. Sagulenko, V. Puller, R. A. Neher, TreeTime: Maximum-likelihood phylodynamic analysis. *Virus*
Evol. **4** (2018), doi:10.1093/ve/vex042.

45. L. R. Joyce, Z. Guan, K. L. Palmer, Phosphatidylcholine Biosynthesis in Mitis Group Streptococci via
Host Metabolite Scavenging. *J Bacteriol*. **201**, e00495-19 (2019).

46. L. R. Joyce, Z. Guan, K. L. Palmer, Streptococcus pneumoniae, S. pyogenes and S. agalactiae
membrane phospholipid remodelling in response to human serum. *Microbiology (Reading)*. **167**,
001048 (2021).

47. T. Azarian, P. P. Martinez, B. J. Arnold, X. Qiu, L. R. Grant, J. Corander, C. Fraser, N. J. Croucher, L. L.
Hammitt, R. Reid, M. Santosham, R. C. Weatherholtz, S. D. Bentley, K. L. O'Brien, M. Lipsitch, W. P.
Hanage, Frequency-dependent selection can forecast evolution in Streptococcus pneumoniae.
PLOS Biology. **18**, e3000878 (2020).

48. T. Golubchik, E. M. Batty, R. R. Miller, H. Farr, B. C. Young, H. Larner-Svensson, R. Fung, H. Godwin,
K. Knox, A. Votintseva, R. G. Everitt, T. Street, M. Cule, C. L. C. Ip, X. Didelot, T. E. A. Peto, R. M.
Harding, D. J. Wilson, D. W. Crook, R. Bowden, Within-Host Evolution of Staphylococcus aureus
during Asymptomatic Carriage. *PLOS ONE*. **8**, e61319 (2013).

49. U. Muthukumarasamy, M. Preusse, A. Kordes, M. Koska, M. Schniederjans, A. Khaledi, S. Häussler,
Single-Nucleotide Polymorphism-Based Genetic Diversity Analysis of Clinical Pseudomonas
aeruginosa Isolates. *Genome Biology and Evolution*. **12**, 396–406 (2020).

50. L. L. Bohr, M. A. Youngblom, V. Eldholm, C. S. Pepperell, Genome reorganization during emergence
of host-associated Mycobacterium abscessus. *Microbial Genomics*. **7** (2021),
doi:10.1099/mgen.0.000706.

51. S. Paik, L. Senty, S. Das, J. C. Noe, C. L. Munro, T. Kitten, Identification of Virulence Determinants for
Endocarditis in Streptococcus sanguinis by Signature-Tagged Mutagenesis. *Infection and*
Immunity. **73**, 6064–6074 (2005).

52. S. F. Bailey, A. Hinz, R. Kassen, Adaptive synonymous mutations in an experimentally evolved
Pseudomonas fluorescens population. *Nat Commun*. **5**, 4076 (2014).

53. S. F. Bailey, L. A. Alonso Morales, R. Kassen, Effects of Synonymous Mutations beyond Codon Bias:
The Evidence for Adaptive Synonymous Substitutions from Microbial Evolution Experiments.
Genome Biology and Evolution. **13**, evab141 (2021).

54. A. Ballard, S. Bieniek, D. B. Carlini, The fitness consequences of synonymous mutations in
Escherichia coli: Experimental evidence for a pleiotropic effect of translational selection. *Gene*.
694, 111–120 (2019).

55. J. Kristofich, A. B. Morgenthaler, W. R. Kinney, C. C. Ebmeier, D. J. Snyder, W. M. Old, V. S. Cooper,
S. D. Copley, Synonymous mutations make dramatic contributions to fitness when growth is
limited by a weak-link enzyme. *PLOS Genetics*. **14**, e1007615 (2018).

56. E. Lundin, P.-C. Tang, L. Guy, J. Näsvall, D. I. Andersson, Experimental Determination and Prediction
of the Fitness Effects of Random Point Mutations in the Biosynthetic Enzyme HisA. *Molecular*
Biology and Evolution. **35**, 704–718 (2018).

57. L. H. Rasmussen, K. Højholt, R. Dargis, J. J. Christensen, O. Skovgaard, U. S. Justesen, F. S.
Rosenvinge, C. Moser, O. Lukjancenko, S. Rasmussen, X. C. Y. 2017 Nielsen, In silico assessment of
virulence factors in strains of Streptococcus oralis and Streptococcus mitis isolated from patients
with Infective Endocarditis. *Journal of Medical Microbiology*. **66**, 1316–1323.

58. S. P. Baker, T. J. Nulton, T. Kitten, Genomic, Phenotypic, and Virulence Analysis of Streptococcus
sanguinis Oral and Infective-Endocarditis Isolates. *Infection and Immunity*. **87**, e00703-18 (2019).

59. K. H. Iversen, L. H. Rasmussen, K. Al-Nakeeb, J. J. A. Armenteros, C. S. Jensen, R. Dargis, O.
Lukjancenko, U. S. Justesen, C. Moser, F. S. Rosenvinge, X. C. Nielsen, J. J. Christensen, S.
Rasmussen, Similar genomic patterns of clinical infective endocarditis and oral isolates of
Streptococcus sanguinis and Streptococcus gordonii. *Sci Rep*. **10**, 2728 (2020).

60. J. P. Galkiewicz, C. A. Kellogg, Cross-kingdom amplification using bacteria-specific primers:
complications for studies of coral microbial ecology. *Appl Environ Microbiol*. **74**, 7828–7831
(2008).

61. S. Andrews, FastQC: A Quality Control tool for High Throughput Sequence Data (2010).

62. A. Bankevich, S. Nurk, D. Antipov, A. A. Gurevich, M. Dvorkin, A. S. Kulikov, V. M. Lesin, S. I.
Nikolenko, S. Pham, A. D. Prjibelski, A. V. Pyshkin, A. V. Sirotkin, N. Vyahhi, G. Tesler, M. A.
Alekseyev, P. A. Pevzner, SPAdes: A New Genome Assembly Algorithm and Its Applications to
Single-Cell Sequencing. *J Comput Biol*. **19**, 455–477 (2012).

63. A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for genome
assemblies. *Bioinformatics*. **29**, 1072–1075 (2013).

64. T. Seemann, Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, btu153 (2014).

65. M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2 – Approximately Maximum-Likelihood Trees for
Large Alignments. *PLOS ONE*. **5**, e9490 (2010).

66. G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol*
Evol. **8**, 28–36 (2017).

67. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics*. **30**, 1312–1313 (2014).

68. J. Hadfield, N. J. Croucher, R. J. Goater, K. Abudahab, D. M. Aanensen, S. R. Harris, Phandango: an
interactive viewer for bacterial population genomics. *Bioinformatics*. **34**, 292–293 (2018).

69. I. Lee, Y. Ouk Kim, S.-C. Park, J. Chun, OrthoANI: An improved algorithm and software for
calculating average nucleotide identity. *Int J Syst Evol Microbiol*. **66**, 1100–1103 (2016).

70. S. De Mita, M. Siol, EggLib: processing, analysis and simulation tools for population genetics and
genomics. *BMC Genetics*. **13**, 27 (2012).

71. A. J. Page, B. Taylor, A. J. Delaney, J. Soares, T. Seemann, J. A. Keane, S. R. Harris, SNP-sites: rapid
efficient extraction of SNPs from multi-FASTA alignments. *Microbial Genomics*. **2** (2016),
doi:10.1099/mgen.0.000056.

72. I. van de Rijn, R. E. Kessler, Growth characteristics of group A streptococci in a new chemically
defined medium. *Infect Immun*. **27**, 444–448 (1980).

73. R. R. Wick, L. M. Judd, C. L. Gorrie, K. E. Holt, Unicycler: Resolving bacterial genome assemblies
from short and long sequencing reads. *PLoS Comput Biol*. **13**, e1005595 (2017).

74. H. M. Adams, L. R. Joyce, Z. Guan, R. L. Akins, K. L. Palmer, Streptococcus mitis and S. oralis Lack a
Requirement for CdsA, the Enzyme Required for Synthesis of Major Membrane Phospholipids in
Bacteria. *Antimicrob Agents Chemother*. **61**, e02552-16 (2017).

Table and Figures
811
812 **Table 1:** Strains used in this study.
813

Organism	Strain	Description	Ref
Mitis group streptococci	1643 (SM43)	Wild-type infective endocarditis isolate	(19)
	SM43ΔNrdM	SM43 with clean deletion of *nrdM*, FD735_06230	This work
	SM43:NrdM-I	SM43 with SM43 *nrdM* allelic replacement into SM43Δ*nrdM*	This work
	SM43:NrdM-C	SM43 with SO48 *nrdM* allelic replacement into SM43Δ*nrdM*	This work
S. oralis	1648 (SO48)	Wild-type infective endocarditis isolate	(19)
	SO48ΔNrdM	SO48 with clean deletion of *nrdM*, MP387_03665	This work
	SO48:NrdM-I	SM43 *nrdM* allelic replacement into SO48Δ*nrdM*	This work
	SO48:NrdM-C	SO48 *nrdM* allelic replacement into SO48Δ*nrdM*	This work

814
815

817
818
Figure 1: Species identification of blood and oral commensal viridans group streptococci
(VGS). A) Three different methods for species identification were used (Kraken, GyrB typing &
16S rRNA sequencing) providing differing results. "na" values in the gyrB column indicate
species not included in this typing scheme. "No species identified" indicates that the isolate could
only be identified to the group level. **B)** Phylogenetic tree of all 81 isolates inferred from an
alignment of GyrB sequences and identified by VGS group and species as assigned by Kraken.
The isolate which appears to have been mis-identified by Kraken is outlined in red. **C)** Invasive
(left) and oral (right) isolates by VGS group according to species identification performed by
Kraken.
828

Figure 2: Pangenome analysis of *S. oralis* strains. A) Summary statistics from pangenome analysis of all *S. oralis* GWAS isolates (n=108). **B)** Core genome phylogeny of *S. oralis* isolates inferred using RAxML and midpoint rooted. Tips colored by isolation source (oral-commensal in blue, blood-invasive in red). 108 isolates total, 57 isolated from the oral cavity and 51 isolated from invasive infections. Three subspecies are labeled and scale bar given in SNPs per site. Nodes colored in purple represent bootstrap values < 50. **C)** Average nucleotide identity (ANI) calculated within and between the three subspecies of *S. oralis* phylogeny show relatively low sequence conservation (~94-96% ANI) between even strains of the same subspecies. ANI values within *subsp. oralis* and *subsp. tigurinus* show significantly higher sequence conservation within compared to between subspecies (Mann-Whitney U test with Benjamini-Hochberg correction). **D)** Accumulation and rarefaction curves for three subspecies. All samples were

841 repeatedly sub-sampled to the size of the smallest sample (*subsp. dentisani,* n=25) and the
842 median core and total gene values plotted for 100 iterations.
843

844
Figure 3: *S. oralis* strains are exceptionally recombinogenic. A) Recombination tracts
predicted by Gubbins in the core genome of *S. oralis* isolates plotted alongside the core genome
phylogeny. Red tracts represent recombination within the sample and blue tracts represent
recombination with isolates outside this sample. 99.9% of the core genome has been affected
by recombination. Visualization performed with Phandango. **B)** Histogram showing the number
of strains in our sample that share a given recombinant fragment. The x-axis is the number of
strains that share a recombinant fragment (with a maximum of 108 isolates, the size of our
sample) and the y-axis is the number of fragments identified by Gubbins that are shared by that
number of isolates.
854

Figure 4: A Single core genome variant is associated with invasiveness in *S. oralis* isolates. A) Weir and Cockerham's F_{ST} (wcF_{ST}) values calculated for each core genome variant (n=67,026) and plotted against the core genome position. Significance threshold (red dotted line) estimated by taking the maximum wcFst value from 100 random permutations of phenotypes. Variants with non-significant F_{ST} values shown in light blue, a single variant with a significant F_{ST} value shown in dark blue. **B)** Pairwise piN/piS values calculated and averaged for each core gene and plotted as a histogram. Average piN/piS value across all genes is 0.039, red dotted line represents the average piN/piS value for *nrdM* (0.024).

Figure 5: A) Homologous *nrdM* genes identified in all newly sequenced VGS isolates with alleles
from our variant of interest plotted on a phylogeny made from GyrB sequences. Only 2 species
in our sample lacked *nrdM* (*S. lutetiensis* and *S. pasteuranius*) both of which are in the bovis
group and are shown in grey. All other species contained *nrdM* and were conserved at the
position of interest. **B)** Multiple sequence alignment (MSA) of NrdM plotted next to the *gyrB*
phylogeny (same tree as in A) where tips have been colored by *nrdM* allele. Scale on top of the
MSA represents length in amino acids, the position of the synonymous mutation of interest (I78I)
indicated with an arrow. Clade labels shown for the most common species: *S. oralis, S. mitis, S.*
anginosus and *S. constellatus*.

876
Figure 6: Simulated infective endocarditis vegetation model of *nrdM* allele swapped
strains. Survival of either **A)** SM43 or **B)** SO48 harboring either the infectious *nrdM*-I allele or
the commensal *nrdM*-C allele during growth in vegetations. Biological triplicates were performed
for each strain with four technical replicates per time point. Mean and SEM indicated. No
significant difference was observed under these conditions. Two-way ANOVA with Sidak's
multiple comparisons test.
883