MetaTiME: Meta-components of the Tumor Immune Microenvironment

Yi Zhang[1,2], Guanjue Xiang[1,2], Alva Yijia Jiang[1], Allen Lynch[1,2], Zexian Zeng[1,2], Chenfei Wang[1,2],

Wubing Zhang[1,2], Jingyu Fan[1,2], Jiajinlong Kang[1], Shengqing Stan Gu[4], Changxin Wan[1,2],

Boning Zhang[1,2], X. Shirley Liu[1,2,3]*, Myles Brown[3,4]*, Clifford A Meyer[1,2,3]*

1. Department of Data Science, Dana-Farber Cancer Institute, Boston, MA 02215, USA

2. Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA 02215 USA.

3. Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, MA, USA

4. Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA 02215, USA.

*Corresponding author. Email: cliff_meyer@ds.dfci.harvard.edu

*Co-corresponding author. Email: myles_brown@dfci.harvard.edu

*Co-corresponding author. Email: xsliu.res@gmail.com

Abstract

Recent advances in single-cell RNA sequencing have revealed heterogeneous cell types and

gene expression states in the non-cancerous cells in tumors. The integration of multiple scRNA-

seq datasets across tumors can reveal common cell types and states in the tumor

microenvironment (TME). We developed a data driven framework, MetaTiME, to overcome the

limitations in resolution and consistency that result from manual labelling using known gene

markers. Using millions of TME single cells, MetaTiME learns meta-components that encode

independent components of gene expression observed across cancer types. The meta-

components are biologically interpretable as cell types, cell states, and signaling activities. By

projecting onto the MetaTiME space, we provide a tool to annotate cell states and signature

continuums for TME scRNA-seq data. Leveraging epigenetics data, MetaTiME reveals critical

transcriptional regulators for the cell states. Overall, MetaTiME learns data-driven meta-

components that depict cellular states and gene regulators for tumor immunity and cancer

immunotherapy.

Keywords

Tumor microenvironment, cancer immunity, single-cell RNA sequencing, transcriptional

regulators, gene expression cell state, cell state representation.

Introduction

Recent advances in cancer research have revealed the integral role of the tumor

microenvironment (TME) in tumor progression and therapy responses[1–6]. Understanding

interactions between cancer cells and the non-cancer compartments, including immune cells,

fibroblasts, and endothelial cells, has revealed potential targets for cancer immunotherapy.

Specifically, single-cell RNA-sequencing (scRNA-Seq) applied on multiple patient tumors has

enabled the high-resolution identification of TME constituents that interfere with the elimination

of cancer cells. For example, exhausted tumor-infiltrating lymphocytes (TILs)[4,7,8], and certain

tumor-associated macrophages subtypes[9–11], have been associated with tumor development.

However, the definition of cell types and cell states in tumor scRNA analyses still relies on

manual labeling by experts using known exclusive biomarkers following unsupervised

clustering[12,13], which lacks consistency and varies between different cohorts.

As single-cell data accumulate, integrating a large collection of cells from multiple cohorts can

help unify the definition of cell types and states to facilitate the automatic annotation of new

scRNA-seq data[14,15]. One approach to cell annotation is to use predefined biomarker lists.

However, these biomarkers might not cover domain-specific cellular states, for example, well-

defined immune cell markers derived from blood immune cells may not fully cover the TME

disease context[14]. Moreover, although cell type definitions in reference databases such as

CIBERSORT, Azimuth, and Human Primary Cell Atlas[16–18] can be useful, the granularity of

these definitions varies between databases. Several efforts integrating pan-cancer scRNA data

have revealed subtypes in the TME through the manual annotation of clusters using a shortlist

of exclusive gene markers[13,19,20].

Another approach is to map a dataset containing unannotated cell states onto an annotated

reference. Methods to obtain such representations include canonical correlation analysis

(CCA)[21], adjusted principal components (Harmony)[22], or generative deep learning models using

variational autoencoders (scVI)[23]. These methods use dimension reduction onto a common

latent space to align cells with similar states between datasets, without ascribing meaning to the

latent space representations. An alternative data driven approach is to identify low dimensional

latent space representations in which a biological meaning can be ascribed to each latent

dimension. Several matrix factorization algorithms have been developed to represent high

dimensional data in a low-dimensional space with interpretable components, including non-

negative matrix factorization (NMF)[24] and independent component analysis (ICA)[25,26].

In this study, we developed a computational framework for mapping millions of single cells from

multiple cohorts onto a comprehensive and interpretable latent space, learnt from the data. The

framework, MetaTiME (Meta-components of the Tumor immune MicroEnvironment), identifies

reproducible low-dimensional meta-components that reflect independent components of gene

expression variation across cohorts and cancer types. MetaTiME adopted ICA for dimensional

reduction to maximize the mutual independence among components. We used MetaTiME to

obtain meta-components (MeCs) from 1.7 million single cells across 79 tumor datasets. These

MeCs represent the TME landscape along 86 data-driven transcriptional directions mirroring

lineage-specific cell states and signaling activities. Furthermore, we developed a MetaTiME

toolkit for using the MeCs to annotate cellular states and signature continuums in tumor scRNA

datasets, and to reveal differential signatures across immunotherapy responses. Finally, by

incorporating transcription factor binding data, MetaTiME revealed and prioritized putative

transcriptional regulators that may modulate tumor immunity.

Results

MetaTiME as a general framework to discover consensus transcriptomic programs

The MetaTiME framework consists of three stages: meta-component (MeC) discovery,

interpretation of MeCs, and application of cell state annotations (**Fig.1a**). The MeC discovery

stage detects repeatable sources of variation from multiple single-cell measurements sharing

similar cellular properties. The MeC interpretation step involves a one-time curation effort using

biomarker databases, pathway information and Cistrome DB chromatin profiling data[27]. In the

third step, users map MeCs onto their new tumor scRNA-seq datasets using MetaTiME

application tools, to obtain annotated cell states and signature continuums.

To train MeCs for the TME context, we collected and curated 2,157,387 cells from 76 studies

ranging across 27 cancer types, using publicly available tumor scRNA-Seq data mostly from

TISCH[28]. After removing the TISCH annotated malignant cells using MAESTRO[17], 102,703

stromal cells and 1,617,110 immune cells were retained for downstream training (**Fig.S1**). The

76 studies were further partitioned according to cancer type, resulting in 93 datasets, including 7

datasets with immune checkpoint blockade (ICB) treatment and 3 10x Genomics provided

datasets representing peripheral blood mononuclear cells (PBMC) sampled from healthy

donors.

In the MeC discovery stage, MetaTiME first decomposes the log-transformed expression matrix

of each single dataset using Independent Component Analysis (ICA)[25]. We adopted ICA to

maximize mutual independence among gene expression components. In simulations ICA

performed slightly better than Non-negative Matrix Factorization (NMF) in simulated single-cell

data with pre-embedded transcriptional signatures (**Fig.S2**). The feature weight distribution of

each Independent Component (IC) also enables normalization of the gene contribution scores

for measuring similarity among components. MetaTiME then applies two transformations to the

IC vectors, z-weight normalization and skewness alignment, to ensure the scales of gene

representation scores are comparable among components (**Methods, Fig.S3a**). Next,

MetaTiME filters ICs to retain ones that are reproducible across multiple cohorts (the minimum

Pearson correlation with any other IC >=0.3). These are passed to the Louvain graph clustering

algorithm to merge IC groups into MeCs (**Methods**). Lastly, MetaTiME computes averaged

profiles of gene z-weights within each IC cluster, yielding 86 MeCs trained for the TME (**Fig.1a,**

Fig.2a). The number of MeCs was automatically determined by simultaneously optimizing

granularity and independence in IC clustering (**Fig.S3b**). Importantly, the MeC clustering does

not depend on cohort source (**Fig.S4a,** cohort source of component), and is robust during leave-

one-out testing in which the MeCs are learnt with a single cohort left out of the analysis

(**Fig.S5**). The leave-one-out MeCs are highly correlated with the MeCs learnt on the complete

data (mean maximum correlation=0.988), and the numbers of components are similar (mean

leave-one-out MeC number=84.64, full-set MeC number=86). This integration after

decomposition approach overcomes batch effects, which are often a challenge in single cell

RNA-seq data analysis. Moreover, we implemented the standard approach in which cluster-wise

signatures are obtained by batch effect removal, clustering, and differential expression analysis

(**Fig.S6a**). Limited by server computing memory, the maximum number of datasets we could

integrate in this way was 21, and the cluster-wise signatures display a lower level of specificity

in the test data (**Fig.S6b,c**). Therefore, the MetaTiME approach allows for the effective

integration of large numbers of single cell datasets, which will become increasingly important as

data accumulates.

MetaTiME defines interpretable meta-components

In principle, each MeC represents one independent source of transcriptional variation commonly

present in the TME. We investigated top ranked genes in MeCs and found MeCs are highly

interpretable, reflecting common biological processes in the TME. For instance, the MeC

derived from the largest IC cluster is highly enriched in interferon response genes, such as

ISG15, *IFI6*, *LY6E*, and *MX1*, indicating that the underlying interferon response is among the

most common source of transcriptional variation shared across tumor samples and cohorts

(**Fig.2a, b**). Intriguingly, top genes of each MeC are enriched in known biomarkers or regulators.

For example, several T cell-related MeCs identify different gene modules co-expressed in T

cells reflecting activation of different T-cell related processes (**Fig.2c, Fig.S7**). The "T cell co-

signaling" MeC-65 features T cell receptors in co-stimulatory and co-inhibitory pathways[29,30],

such as *TNFRSF4* (OX40), *TNFRSF18* (GITR) *TNFRSF9* (4-1BB), and *ICOS* (**Fig.2c**, left). The

"CXCL13+ exhausted CD8 T cell" MeC-40 features receptors characterizing the exhausted CD8

T cell state[8], including *HAVCR2* (TIM3), *LAG3*, *TIGIT*, and *PDCD1* (PD1) (**Fig.S7**), each being

potential ICB targets[31]. In addition, this MeC is characterized by a high level of *CXCL13*

(**Fig.2c**, second panel), a cytokine mediating immune cell trafficking to tertiary lymphoid

structures[32]. In contrast, a related MeC representing T cell co-signaling receptors in regulatory

CD4T cells (Treg) has a different ranking, including *TNFRSF18*, *TNFRSF4*, *TIGIT, TNFRSF1B*,

CTLA4, *CD27* among the top 20 genes, along with the regulatory T cell-specific marker *FOXP3*

(**Fig.2c**, right). Though ICB has been an extremely successful therapy for some patients, it has

not yet had an impact on the majority of patients[33]. Investigating the top members in the MeCs

involving T cell receptor pathways may help identify new ICB targets.

MetaTiME depicts the functional landscape of transcriptomic variation and cell states in

the tumor microenvironment

We provided functional annotations of all MeCs by examining top z-weight genes and compared

these with functional gene sets, such as immune cell type markers[15,18] and gene ontology

databases[34]. We found that 86 MeCs clearly mirror gene expression patterns corresponding to

cell types, cell states and signaling pathway activities, depicting a landscape of non-cancer cell

states in the TME (**Fig.2, Table S1**: MeC annotation). The top genes of the cell type MeCs

match well-known lineage-specific markers[15,18]. Examples include *CD74*, *CD79A*, *MS4A1* for B

cells ("B cell" MeC-18), *CD3D*, *CD8A*, *CD8B* for T cells ("CD3 - CD8 T cell" MeC-32), and *LYZ*,

VCAN, *S100A9* for CD14+ Monocytes ("CD14 monocyte" MeC-17) (**Fig.2b, Fig.2d**). The

majority of MeCs define high resolution lineage-specific cell states (**Fig.2b, Fig.S4b**). Taking the

B cell lineage as an example, multiple MeCs harbor genes specific to B cell developmental

stages[35], ranging from a progenitor B cell state (*CD69* and *PAX5* in the "PAX5 B cell" in MeC-

50), to a mature B cell state (*CD79A* in the "B cell" MeC-18), an antibody-secreting plasma cell

state (*XBP1* in the "B_Plasma" MeC-4, and *JCHAIN* in the "JCHAIN+ plasma B cell" MeC-77),

and immunoglobulin secretion states (IGK and IGH in the "immunoglobin kappa B cell" MeC-30,

IGL and IGH in the "immunoglobin lambda B cell" MeC-50) (**Fig.2b**). Lastly, like the interferon

responsive MeC mentioned above, we found a subset of MeCs that are more accurately

interpreted as signaling pathways because their top genes are more related to pathways or

molecular functions than to cell identities.

We organized the 86 annotated MeCs into six cell lineage-focused categories and one signaling

pathway-focused category (**Fig.2a,b** and **Table S1**: MeC annotation, MeC enrichment). Among

these, there are 6 B lineage-related MeCs for B cells; 20 T cell lineage MeCs covering CD8 T

cells, CD4 T cells, and natural killer (NK) cells; 4 dendritic cell (DC) lineage MeCs; 12 monocyte

and macrophage-related MeCs; 3 platelet, erythrocyte, and mast cell MeCs; 6 stromal cell-

related MeCs for fibroblasts, myofibroblasts and endothelial cells; and 35 MeCs in the signaling

category (**Fig.2b**). We demonstrated that the MeCs are of high specificity, visualizing the z-

weights of known cell subtype markers and pathway biomarkers (**Fig.2d**). Correlating MeCs

with the comprehensive immune cell type database Azimuth[15] validated the lineage-specificity of

several MeCs, while most MeCs reflect cell states that appear specific to the tumor context

(**Fig.S4b**).

MetaTiME annotates cell states and signature continuums when applied to the tumor

microenvironment single-cell data

As MetaTiME MeCs provides a highly interpretable basis for the TME in single cells, we

provided a toolkit to reveal MeC signature continuums and enriched cell states in scRNA-seq

TME data (code deposited in https://github.com/yi-zhang/MetaTiME). The MetaTiME annotation

toolkit takes as input the scRNA-seq expression matrix after depth normalization and log

transformation, maps each single cell onto the pre-trained MeC space, and annotates the most

highly enriched cell states for pre-defined cell clusters. The cell clusters are by default

calculated using graph clustering with high resolution after an optional batch effect correction

with Harmony[22]. We demonstrate the application of MetaTiME on basal cell carcinoma (BCC)

single-cells from *Yost et al.*[8]. These test cells were excluded from the MetaTiME training stage.

MetaTiME annotates the enriched cell states (**Fig.3a**) highlighting gradients of exhausted CD8 T

cells and follicular helper T cells (Tfh) (**Fig.3c**). The most enriched cell states consistently match

the manual labelling from the original study with improved resolution (**Fig.3b**) and highly

express corresponding markers (**Fig.3d**). In addition, compared to Seurat's[14] automated

CIBERSORT marker-based annotations (14 cell types, **Fig.S6a**), MetaTiME provides higher

resolution (38 cell states, **Fig.3b**). Other automatic annotation gene panels were also tested,

including the human primary cell atlas (HPCA) panel and Blueprint-ENCODE panel used in

SingleR[16], where macrophages and plasma cells appear to be mislabeled as subclusters within

T cell clusters (**Fig.S8b, c**). Interestingly, the MetaTiME annotation not only indicates the CD8 T

cell and CD4 T cell subtypes, but also splits cells further into cell states with polarized

expression in proliferation, cytotoxicity, exhaustion level, heat stress, co-signaling pathways,

and so on. (**Fig.3a, Fig.S8a**). The B cell group is further partitioned into distinct B cell

developmental states including a B cell cluster with cell cycle and MYC activities (**Fig.3a,b,**

Fig.S8), which possibly represent germinal center (GC) B cells undergoing active expansion

and maturation[36].

We thus re-annotated all tumor scRNA cohorts using MetaTiME and investigated the distribution

of cell state compositions across cancer cohorts. As shown in **Fig. 3e** and **Fig.S9**, tumors are

highly heterogenous and the TME cellular composition is only partially determined by cancer

type. For example, Cholangiocarcinoma (CHOL) is highly enriched in stromal cells including

collagen-secreting fibroblast, as expected[37], while other samples including ovarian cancer (OV),

pancreatic adenocarcinoma (PAAD), and multiple myeloma (MM) are also stromal-rich.

Furthermore, tumors with high infiltration of the "GZMK+ CCL5+ CD8 T cell" state include

multiple tumor types including colon cancer, breast cancer, and skin cancer, suggesting that

immune infiltration is sample-dependent and that cancer treatments should be personalized[38].

Differential MetaTiME analysis detects alterations of transcriptional programs in

immunotherapy

Single-cell data derived from ICB trials is invaluable for identifying cell types associated with ICB

treatment or response[8]. However, the detection of differential cell type abundances in ICB

cohorts has been challenging due to the heterogeneity of cell type proportions and to the limited

numbers of patients in each cohort[39]. We compared differences in MeC signatures instead of

cell count proportions, to understand immune responses during ICB. We analyzed two ICB

cohorts, a basal cell carcinoma (BCC) cohort with samples from pre- or post-ICB treatment[8],

and a bladder cancer (BLCA) cohort with samples from ICB responders and non-responders[40].

We applied MetaTiME for per-cluster cell state annotation and per-cell MeC signature

evaluation. For each cell state cluster, we tested all MeC signatures passing significance

(average z-weight>2) between conditions using the Wilcoxon rank-sum test. We plotted cluster-

wise signatures in the significance – effect size scatterplot to highlight the most significant

differential MeCs (**Methods**). In a comparison of pre- and post-ICB treatment, we observed

higher expression of cytotoxic T cell and B cell MeCs in the post-ICB samples. Moreover,

several monocyte and macrophage states are also suppressed after ICB treatment (**Fig.4a**).

Notably, the IL1B-positive macrophage signature is also found to be elevated in non-responders

compared to responders in the BLCA ICB cohort (**Fig.4b**). Since activation of the IL1B pathway

is a known regulator of inflammatory processes[41], we sought to investigate whether the IL1B-

positive macrophage signature is associated with tumor survival prognosis in bulk RNA-seq

data from The Cancer Genome Atlas Program (TCGA). We evaluated TCGA tumors using the

averaged expression of the top 20 genes from the "Macrophage IL1-NFkB" MeC (MeC-42)

which ranks first in elevated MeCs in non-responders (**Fig.4a**). We found that higher expression

of the *IL1B* signature is associated with lower survival rate in multiple cancer types, especially in

Low Grade Glioma (LGG) and in Kidney renal cell carcinoma (KIRC) (**Fig.S10**). This suggests

the macrophage state with *IL1B* pathway activation is associated with poor prognosis and lower

ICB efficacy.

MetaTiME delineates myeloid cells in different metabolic states

As specific myeloid cell states have been associated with cancer survival and treatment

response, we sought to systematically characterize MeCs related to monocytes and

macrophages. Although the canonical definition of M1 and M2 macrophages is derived from

cytokine polarized macrophages *in vitro*[42], MetaTiME's myeloid-related MeCs represent a more

complex framework for understanding tumor-infiltrating macrophages. MetaTiME's 12 monocyte

and macrophage related MeCs can be summarized into six central monocyte or macrophage

states for the TME, after merging similar states such as "Macrophage IL1-NFkB" and

"Macrophage IL1- JUN" due to similarity among top genes (**Fig.4c**). Monocytes are classified as

two categories, CD14+ and CD16+. For macrophages, four MeCs define common states of

intra-tumor macrophages: C1Q+, SPP1+, lipid-rich, and IL1B+ macrophages, and two MeCs,

representing interferon and MHC-II signaling pathways, are less frequently observed among

macrophages (**Fig.2b, Fig.4c**). In comparison, previous studies defined different tumor

associated macrophages in terms of manually selected representative genes after clustering

myeloid cells. For example, *Cheng et al.*[10] defined several tumor associated macrophage types

including *ISG15+*, *SPP1+*, *INHBA+*, *VCAN+*, *NLRP3+*, and *FN1+* macrophages, while *Bi et al.*[43]

defined *CXCL10*-high, *GPNMB*-high, *FOLR2*-high, *VSIR*-high, and cycling macrophages for

advanced renal cell carcinoma (ccRCC). We find that the MetaTiME-defined macrophage MeCs

reflect co-expression relationships with the selected marker genes. For example, macrophage

markers from *Bi et al*. rank high in several macrophage-related MeCs (**Fig.S11**), and the

expression pattern of the marker genes picked by *Bi et al.* (*CXCL10*, *GPNMB*, *VSIR*, *FOLR2*,

Cycling marker *MKI67*) correspond to several MeCs ("interferon response" MeC-0, "PPARG+

lipid-rich" MeC-37, "MHCII-high" MeC-58, and "RNASE1+,C1Q+" MeC-49) (**Fig.S11**). However,

MetaTiME reveals additional distinct components such as the "SPP1+" and "C1Q+" MeCs,

which were detected as separate myeloid types in the *Cheng et al.* multi-cohort study (**Fig.S12**).

While the manual reconciliation of cell types from multi-cohort scRNA data shows many marker

genes to be consistent with the top genes in the MetaTiME MeCs, the myeloid cell population is

not neatly partitioned into cell clusters and might be better represented in terms of expression

signature continuums. For example, when mapping myeloid MeCs onto the kidney myeloid

cells, the IL1B+ MeC signature (MeC-42, MeC-45) is distributed across the "Macro_IL1B"

cluster as well as the CD14 monocyte cluster (**Fig.S12**).

To investigate functional differences among the different macrophage states, we applied gene

set enrichment (GSEA)[34] analysis using the top MeC genes (**Methods**). Interestingly, the

different macrophage states have different metabolic preferences (**Fig.4d**). Glucose metabolism

and the glycosylation pathway are highly active in SPP1+ macrophages, while lysosome and

phagosome activity are the most highly enriched in C1Q+ macrophages. Lysosome and

cholesterol metabolism, including *PPARG* signaling, are enriched in the lipid-rich state. The

inflammatory IL1B and NFkB pathways are highly active in IL1B+ macrophages. Several

macrophage states are related to cell signaling. *SPP1* for example, encodes Osteopontin, which

has been found to foster an environment that promotes cancer metastasis[44]. The C1Q+ MeC

features *C1QA*, *C1QB*, and *C1QC*, members of the family of complement molecules that could

play dual roles in chronic inflammation[45]. The IL1B+ meta-components features cytokines co-

expressed with *IL1B*, including *CXCL8*, *CXCL2*, and *CXCL3*, all of which can interact with other

cells in the TME by binding to cytokine receptors[46] (**Fig.2b**).

Incorporation of epigenetic data prioritizes transcriptional regulators of tumor immunity

We next investigated the transcription factors (TFs) that regulate the MeCs, hypothesizing that

the co-expression of genes in a subset of MeCs is determined through TF regulatory events.

Our group previously developed the Cistrome Data Browser and Lisa to predict transcriptional

regulators of gene sets based on chromatin immunoprecipitation with sequencing (ChIP-seq)

data[27,47]. Thus, we used Lisa to predict the TFs that regulate the top genes of each MeC, and

compared these Lisa regulatory prediction scores with the MeC z-weights across TFs. We found

that, for many MeCs, the same TFs were predicted to be both regulators of the MeC and were

highly expressed in the MeC itself, indicating an autoregulatory control scheme. Often, however,

TFs that were predicted by Lisa to be MeC regulators were not represented by high MeC z-

weights, and TFs with high MeC z-weight were not always found to have high Lisa scores

(**Fig.5, Table S1**: MeC regulators). TFs predicted by Lisa but not represented by high MeC z-

weight could be the result of TF activities being regulated through non-transcriptional

mechanisms[48] or multiple TFs in a family having similar binding patterns but only a subset being

the regulators[49]. TFs that have high MeC z-weights but low Lisa scores are most likely not well

represented in the relevant cell types in available ChIP-seq data. In the "interferon response"

MeC-0, *STAT1* is highly represented in the MeC z-weight and Lisa ranks *STAT1* as the top

regulator, consistent with *STAT1* being known as the master regulator of the interferon

response (**Fig.5a**). Several lineage-defining TFs display the autoregulatory pattern, including

TCF4 in plasmacytoid dendritic cells (pDC) (**Fig.5b**) and *XBP1* in B plasma cells (**Table S1:**

MeC regulators). The macrophage related MeCs are regulated by myeloid lineage TFs like

CEBPB, and TFs related to immune stimulus responses, including NFkB complex TFs. In the

"lipid-rich macrophage" MeC-37, although PPARG ranks among the top Lisa-predicted

regulators (**Fig.5c**), *PPARG* expression is not highly represented as a MeC z-weight. In this

MeC, the top co-expressed genes are indeed enriched in the PPARG signaling pathway

(**Fig.4d**); this result can be accounted for by PPARG being regulated through its ligands, which

include a variety of lipophilic acids[48].

We found glucocorticoid receptor (GR) signaling to be implicated in the regulation of the

"CXCL13+ Tfh" MeC-20, with GR being most highly ranked TF in both MeC and Lisa scores

(**Fig.5d**). Top genes in the "CXCL13+ Tfh" MeC include several direct target genes of GR,

including *SRGN* and *FKBP5* (**Fig.S13b**). We investigated whether the *CXCL13* cytokine itself

could be a direct target of GR in *CXCL13* secreting Tfh cells. Since GR ChIP-seq data is not

available for the exact Tfh cell state, we collected GR ChIP-seq data from several other cell

types. Direct binding of GR is observed at the *CXCL13* gene promoter and nearby the gene

locus at putative enhancers, which are conserved across multiple cell lines (**Fig.S13a**), including

the B cell line Nalm6, the monocyte cell line THP1, and cancer cell lines. Moreover, in another

CXCL13 secreting cell state, the "CXCL13+ exhausted CD8 T cell" MeC-40, GR is also highly

ranked in both Lisa and MeC scores (**Fig.5d**). Thus, we hypothesize that GR is likely to be a

transcriptional driver of the CXCL13-secreting cell states in exhausted CD8 T cells[50] as well as

in CD4 T follicular helper T cells. Thus, the GR pathway could be a candidate target in tumor

immunity modulation.

Discussion

We developed the MetaTiME (Meta-components of the tumor immune Microenvironment)

framework and performed a large-scale and pan-cancer integration of tumor single cell datasets

using ICA to optimize information independence among components[51]. We identified 86

interpretable meta-components (MeCs) that describe common aspects of TME gene expression

variation across multiple tumors.

The MetaTiME MeCs serve as comprehensive transcriptional signatures that depict a functional

landscape of TME transcriptional programs and cell states. For monocytes and macrophages,

the related MeCs revealed heterogeneity and plasticity of tumor-associated macrophages

(TAMs). We thus propose that TAMs, especially for solid tumors, should be classified based on

the major states with different metabolic preferences instead of the canonical M1 and M2

classification[42]. Similar states that do not fit well into the M1 versus M2 classification scheme

were also observed in previous studies analyzing myeloid cells in the TME, where single cells

were clustered and labeled using differential markers[10,12,43]. *Cheng et al.*[10] defined several TAM

types by clustering myeloid cells separately for each study and naming the TAMs with manually

selected top marker, chosen based on consistency across cohorts. Similarly, *Bi et al.*[43] defined

TAM types by harmonizing patients and naming the TAMs with top genes in each cluster for an

advanced renal cell carcinoma (ccRCC) cohort. Cell type definitions in the previous studies

were based on representative genes, which were chosen differently in the respective studies.

We propose that the MetaTiME derived monocyte and macrophage MeCs could be used to

define macrophage states and functional co-expressed gene modules more consistently for the

TME. Reexamining the marker genes from the previous studies: *NLRP3* is highly ranked in the

"IL1B+" MeC; in fact, the NLRP3 inflammasome mediates interleukin-1β production. *GPNMB* is

weighted among the top 20 genes in both the "C1Q+ macrophage" and "SPP1+ macrophage"

MeC-19 and MeC-26; it encodes a membrane glycoprotein which is typically highly expressed in

macrophages. *FOLR2* is ranked 29th in the "C1Q+ macrophage" MeC, indicating this

macrophage state also encodes a high folate-activated pathway. Finally, the "SPP1/C1Q

macrophage" MeC features an intermediate state with both SPP1 and C1QA, indicating the

plasticity and mixed nature of pathways activated in TAMs that could not be defined using

exclusive markers. Thus, the myeloid MeCs may provide a consistent definition of TAM states

corresponding to different metabolic processes.

MetaTiME provides a toolkit for analyzing independent TME scRNA-seq datasets by mapping

gene expression onto the MeC space. The outputs include signature continuums and the most

highly enriched cell states. Recent useful single-cell dataset integration algorithms such as

Harmony[22] and scArches[52] infer a joint low-dimensional representation among data. In these

approaches the shared space is re-computed every time a new dataset is incorporated. The

MetaTiME strategy builds upon previous approaches that transfer latent representations from

large datasets, but provides a stable and interpretable representation specialized for the TME.

By leveraging ChIP-seq data, MetaTiME reveals critical transcriptional regulators in tumor

immunity. In many cases, we found the joint consideration of MeC-specific co-expression

patterns and TF binding enrichments reveals the roles of TFs in defining cellular states and

gene expression programs. MetaTiME captured multiple known TFs critical to tumor immunity

and could serve as immune modulation targets; this includes *TOX* in the "CXCL13-secreting

exhausted CD8 T cell" MeC-40, a recently discovered regulator of T cell exhaustion[53] (**Table

S1**). The MeCs further implicated the glucocorticoid receptor pathway in the regulation of

several T cell states. Glucocorticoids are a class of steroid hormones essential to the

modulation of multiple biological processes, including immune related ones[54], although the role

of the GR pathway in different immune cell types is not fully understood. Since GR is broadly

expressed in many cell types, and is regulated through ligand binding, differential analysis of

GR expression is unlikely to fully capture GR regulation in single-cell data analysis. Though GR

ChIP-seq is not available in the contexts of the relevant T cell states, GR ChIP-seq in other cell

lines demonstrate robust binding nearby the top gene *CXCL13*. *CXCL13* is crucial to T follicular

helper cell communication with germinal center B cells, through interaction with its receptor

CXCR5[55,56].

Overall, MetaTiME depicts the functional landscape of transcriptomic variation and cell states in

the tumor microenvironment. It provides a computational framework to facilitate the elucidation

of the identity and function of cells in the TME in future studies and will facilitate the

identification of potential new therapeutic targets for immune modulation.

Methods

Tumor single-cell RNA-seq data collection and processing

For an extensive collection of single cells from tumor microenvironment, we utilized the public

tumor scRNA-seq collection, TISCH[28]. The TISCH collection uniformly processed each dataset

with MAESTRO[17] and isolated non-malignant environmental cells from malignant cells. Overall,

we collected 2,157,387 cells from 76 studies ranging 27 cancer types. The MAESTRO

annotation was labeled using CIBERSORT gene panels[18] followed by curation, enabling

selection of 1,719,813 tumor micro-environment cells, including 1,617,110 immune cells and

102,703 stromal cells, were retained for integrative analysis in this study. For studies with data

measured from multiple cancer types, cells different cancer types were split into independent

datasets, resulting in 93 datasets; it includes 3 PBMC datasets from healthy donors from 10X

Genomics as baseline and 7 datasets with ICB treatment.

For an unsupervised component analysis, each dataset was re-analyzed. For datasets with raw

count matrices available, gene expression was normalized towards per-cell read depth 10,000

followed by log transformation. For datasets with only TPM or FPKM values available, including

Smart-seq data or studies with only normalized matrix available, gene expression underwent log

transformation. Cells were filtered based on minimum library size 1000, gene number 500, and

maximum mitochondrial read proportion 5%.

Decomposing individual studies and denoising low-dimensional components.

We then decomposed the expression matrix of each scRNA-seq dataset using fastICA[51] into an

independent component (IC) vector matrix and a projection weight matrix. We tested different

values for the number of components (k) and chose k to be 100 uniformly for each dataset,

given it could cover more variations than the number of cell types in the TME, which is around

twenty. We applied two denoising approaches to handle sparsity and potentially noisy ICs. First,

we performed a z-score transformation of the gene loadings in the component, scaling all gene

loading values by the standard deviation of each IC. The gene loadings indicate the degree of

contribution to the component as a "metagene" from each gene, and we observed that most

genes contribute neutrally to the metagene. Thus, genes with significant contributions are

selected using the two-standard deviation threshold from either the positive or the negative side.

Second, we aligned the positive skewness of components (using scipy.stats.skewtest[57]) since

the sign of an independent component is randomly assigned in fastICA optimization. We

observed that asymmetrically extreme gene loading values highlight genes representative of the

component's function; thus, we computed each component's skewness statistics and flipped the

sign of component loadings if the skewness is negative. We excluded genes with a low

contribution (gene weight not passing two standard deviations) to any reproducible components

and kept 6623 genes with potentials in driving the reproducible components. The post-

decomposition steps ensured the attitude and sign of gene weights are comparable across

cohorts, depicting degree of contribution from each gene in the genome-wide background.

Meta-component calling and functional annotation

We then aim to discover reproducible patterns from all components from each dataset. We

evaluated similarity between pairs of components using cosine distance and retained a set of

1043 candidate reproducible components from 69 datasets, each with a minimum Pearson

correlation coefficient 0.3 with at least one different IC. We then clustered ICs using Louvain

clustering, a graph-based community detection algorithm where the resolution parameter

controls segmentation granularity. Clusters with at least five ICs were retained as reproducible

IC clusters. The number of clusters is determined by optimizing both Silhouette's score for

optimal within-cluster similarity compared to inter-cluster similarity and number of reproducible

clusters. The final resolution parameter was chosen to be 1.25 resulting in 86 clusters for meta-

component (MeC) calling. The consensus gene z-weights in each MeC were then calculated by

averaging ICs in each cluster. Genes of outlier z-weights passing two standard deviations were

highlighted as significant, and the ones with positive largest z-weights were considered

representative of the MeC.

MetaTiME MeCs were assigned curated annotation by matching top z-weighted genes to

functional biological information including cell type markers, pathway databases from GSEA,

cell types expressing top MeC genes, and high-rank transcription factors. In detail, GSEA

enrichment analyses utilized the top 100 highest z-weighted genes and TF database was

obtained from AnimalTFDB[49]. GSEA was also performed using the z-weight rank of all genes[58].

The 86 MeCs were first ordered by MeC cluster size, then organized into seven functional

categories with six lineage-related categories and one signaling category.

Simulating multi-cohort single-cell RNA data with expression programs

To benchmark dimensional reduction methods, we built upon previous effort from *Kotliar et al.*[59]

to use the scsim package to simulate multiple count matrices with built-in transcriptional

programs. In principle, the built-in gene expression programs (GEP) were sampled as random

scaling factors on a subset of genes mimicking overexpression or suppression of a pathway.

For testing whether a higher number of cohorts facilitate GEP recovery, we simulated 20 single-

cell datasets and tested usage of 5 cohorts, 10 cohorts, and 20 cohorts. Each dataset was

embedded with a subset of 14 pre-defined GEPs, since the real tumor scRNA data may not

cover every possible cell type or gene program in every dataset. The 14 GEPs contain 13 cell

type-specific programs with distinct cell type-specific genes, and one signaling gene expression

program that is randomly active in multiple cell types. Two low-dimensional reduction method

are benchmarked using simulated scRNA data: independent component analysis (ICA) and

non-negative matrix factorization (NMF). Decomposition was performed on each single cohort

separately, and meta-component calling was done as similar in MetaTiME: components are

filtered, clustered into meta-components, followed by averaging gene z-weights per cluster as

predicted gene expression programs (GEP). The predicted GEPs were compared with pre-

defined True GEPs using Pearson correlation. Overall, both ICA and NMF can recover GEPs,

while the ICA-based GEPs are more mutually independent and performs slightly better. Since

the GEP recovered in the 20 cohorts case matched true GEP better than 5 cohorts and 10

cohorts, the increased number of cohorts also improves GEP recovery. Thus, we chose ICA for

component integration and use all available datasets for GEP discovery for tumor

microenvironmental cells.

MeC calling robustness with leave-one-out testing.

MetaTiME's robustness was examined by re-calling MeCs after leaving one dataset out and

calculate the Pearson correlation between the left out dataset MeCs and the full-set MeCs. The

leave-one-out analysis was first performed by removing the SCC cells from the Yost et al.

dataset, which was shown as an example of MetaTiME annotation. Each dataset in the MeC

pool was left out to call new MeCs using the same parameters, resolution=1.25 and

min_cluster_size=5. We used two metrics to check the similarity of each leave-one-out

experiments compared to the full set. The first metric is the mean maximum correlation between

leave-one-out and the full set. This is calculated by taking the maximum correlation, row-wise

and column-wise in the correlation heatmap, and then taking the average. The second metric is

the number of MeCs.

Comparing MeCs, signatures by post-embedding integration, to signatures by cluster-

wise differential expression.

We compared MetaTiME MeCs to the standard approach that extracts signatures after

integrating cells across datasets. The standard approach to obtain cluster-wise signatures is to

perform dataset harmonization, cell clustering, and cluster-wise differential expression analysis.

When performing dataset integration, we first used CCA to include as many datasets as

possible given our maximum memory available in our computing resource (150GB). The

maximum number of datasets integrated by CCA is 10 when using datasets ranked by the

number of TME cells. We then used scanpy to achieve the successful integration of 21

datasets. The integrated cells are harmonized with Harmony[22] and clustered, followed by

differential expression analysis using the Wilcoxon test to extract cluster-wise signatures. The

signatures are in the format of gene vector of log fold change, and named as "Cluster DE

signatures". To control the confounding effect of signature numbers, the resolution of clustering

is chosen to reach 90 clusters, a similar number with the number of full-set MeCs which is 86.

Then, we compared MetaTiME MeCs to the Cluster DE signature projected onto a test dataset.

We used same BCC cells as the test cells that are never seen in the training step from either

method. To reflect how variable the projected scores are across cells, we grouped cells from the

test BCC data into 25 clusters with Leiden clustering. The heatmap of the cluster-wise signature

mapping is plotted to observe how specific each type of score is across cell groups. We further

used cross-cluster entropy to quantify the information content of scores relative to clusters in the

test dataset.

The MetaTiME annotator for analyzing new tumor scRNA-seq data.

MetaTiME provides an analytical toolkit for annotating cell states and signature activities for

tumor scRNA-seq data (https://github.com/yi-zhang/MetaTiME). The scRNA-seq data is first

processed following standard procedures, which includes cell depth normalization, log-

transformation, batch effect removal using Harmony[22], neighboring graph construction, graph

clustering, and UMAP embedding for visualization. Specifically, the clustering step uses an

over-clustering strategy, which sets a high-resolution parameter (default 8) that generates a

larger number of clusters and help reveal fine structures among the cells. Then, the MetaTiME

annotator tool takes as input a single log-transformed expression matrix for TME cells from the

dataset. The outputs include both per-cell MetaTiME MeC signature scores and per-cluster

enriched MeC state. For the per-cell score, MetaTiME projects each cell onto the MeC space by

calculating the dot product between the expression vector and the z-weight vector of each MeC,

using genes passing the significant z-weight criterion (z-weight >2). The projection matrix is

then scaled across all cells to ensure normally distributed scores within each MeC, outputting

the cell-by-MeC score matrix. Meanwhile, the UMAP view of the projection score shows the

signature gradient across the cells positioned by similarity. Lastly, the cluster-wise MeC

enrichment results are also generated. The per-cluster MeC enrichment score is calculated by

averaging profile of cells along each MeC; MeCs with mean score passing the significant cutoff

(2 in the z-weight scale) are called as the set of enriched MeCs. Each cluster may enrich

multiple number of MeCs, and the top enriched MeC with highest score is used in UMAP

visualization.

Differential MeC signature analysis

For tumor scRNA-seq data with different conditions, a differential signature analysis can be

carried out following MetaTiME annotation, which provides enriched MeCs for each cluster and

names each cluster with the top enriched MeC. Thus, for each cell cluster, the MeC signature

strength can be compared across conditions, for all enriched MeC in the current cluster. In

details, a simple Wilcoxon rank-sum test is adopted to compare MeC scores of cells in one

condition with another. The log-fold change of MeC scores were calculated by the ratio between

cell means from the two conditions in comparison. To plot the cluster-specific differential

signature plot, the signatures are marked using "EnrichedMeC@ClusterName", where the

"ClusterName" is the top first enriched MeC used as cell state as current cluster. When

"EnrichedMeC" is the same as the "ClusterName", only "ClusterName" is marked on the

Significance-Effect size plot.

Incorporation of epigenetic data using Lisa

Our group previously developed Lisa that predicts the influence of TFs on a set of genes. Lisa

models public chromatin accessibility and TF binding profiles to score TFs in gene regulation

from an epigenetic perspective. We developed Lisa2 that improves on running speed and

pipeline integration, which is applied on each MeC to score TFs in regulation potential on top

100 high z-weighted genes. The impact scores of TFs are thus from two sources: MeC z-

weights for expression representation, and Lisa scores for binding potential. The TFs are

grouped into three classes, TFs highly ranked based on both MeC gene weights and Lisa

significance, TFs representative only in MeC, and TFs based on binding information only. In

TableS1, we marked TFs from different classes in different colored columns. Significant TFs

based on both MeC and Lisa (MeC z-weight >=2, Lisa score -log p-value >=2) are marked as

orange color; furthermore, the TFs ranking among top 40 (aggregated rank of MeC and Lisa)

compared to all genes are further marked red. TFs ranking among top 10 only in MeC z-weight

are colored green, and TFs ranking among top 10 only in Lisa score are colored blue.

Grant information

NIH grants P01CA163222, U24CA237617, and R01HG011139.

Acknowledgement

We acknowledge all authors of the original tumor single-cell studies that generated publicly

available data. The results appearing here are in part based upon the data generated by the

TCGA Research Network (http://cancergenome.nih.gov/).

Conflict of Interest

MB is a consultant to and receives sponsored research support from Novartis. MB serves on the

SAB of H3 Biomedicine, Kronos Bio, and GV20 Oncotherapy. X.S.L conducted the work while

being on the faculty at DFCI, and is currently a board member and CEO of GV20 Therapeutics.

The remaining authors declare no competing interests.

Figures

Fig.1.

Fig.1. Overview of MetaTiME. MetaTiME integrates 1.7 million single cells to learn common

transcriptional programs in the tumor microenvironment (TME). **(a)** Steps for Meta-components

(MeCs) discovery. For each scRNA dataset, the expression matrix of TME cells is decomposed

into a loading matrix (red) and an independent component (IC) matrix through independent

component analysis (ICA). The ICs represent mutually independent sources of transcriptional

variation. ICs from each dataset are concatenated and clustered into groups of ICs with high

similarity, representing transcriptional programs shared across TME. MeCs are then calculated

as averaged profiles of ICs from each cluster. Each MeC is interpretable, representing gene

signatures of cell type, cell states, or signaling pathway activities. **(b)** Left: MetaTiME provides

86 functionally annotated MeCs that depict the TME transcriptional landscape. They are

grouped into six lineage-related categories and one category reflecting signaling activities.

Middle: the MetaTiME annotation tool facilitates automatic annotation of cell states for new

tumor scRNA data. Right: candidate regulators of each MeC are prioritized by combining MeC

gene weights with epigenetics data. MeC: meta-components, TME: tumor microenvironment,

ICA: independent component analysis, MeC: meta-component, TF: transcription factor.

Fig. 2.

Fig.2. MetaTiME meta-components are biologically interpretable with top genes. (a)

Heatmap of top ten most recurrent clusters of MeCs showing normalized gene weights. **(b)**

Biological characterization of each MeC with top genes. To facilitate biological interpretation,

MeCs are categorized into six lineage-associated classes (B cells, T cells for CD4T, CD8T and

NK cells, dendritic cells, monocyte and macrophages, other myeloid cell types, and stroma

cells) and one signaling pathway-associated class. **(c)** Examples of T cell related MeCs with top

20 genes with largest weights. **(d)** Gene contribution of known lineage-related biomarkers for

each MeC, and correlation with known immune markers from Azimuth. In the top dot plot, size

and color represents MeC z-weights of each gene in each MeC. In the bottom dot plot, size and

color represents the maximum correlation coefficient between MeC and Azimuth defined marker

genes per cell type. MeC: meta-component, DC: dendritic cell, Mono/Mac: monocytes and

macrophages.

Fig.3.

Fig.3. MetaTiME annotates cell states with high resolution on tumor microenvironment

single-cell data. (a) MetaTiME cell state annotation of cell clusters in a basal cell carcinoma

scRNA dataset based on top enriched MeCs. **(b)** Manual annotation labels by experts from the

original study shown on the same UMAP space. **(c)** Signature continuums of four MeCs

representing the mature dendritic cell state, the CXCL13-secreting exhausted T cell state, the

CXCL13-secreting T follicular helper cell state, and the IL1B pathway-activated macrophage

state. **(d)** Marker gene expression for each annotated cell cluster as in (a)**. (e)** Bar plot showing

cell state composition of tumor microenvironment for tumor scRNA dataset cell states. The

proportion of cell states from the same MeC category are aggregated.

Fig.4

Fig.4. Differential signature analysis and delineated macrophage states in TME. (a)

Differential MeC signature testing for enriched cell states comparing pre- and post-

immunotherapy conditions in basal cell carcinoma (BCC). X-axis: Log odds ratio of mean

signature scores between post- and pre-immunotherapy conditions. Y-axis: minus log p-value

from Wilcoxon test. The significant cluster-wise differential signature is marked as

"EnrichedMeC@ClusterName"; when the enriched MeC is the same as cluster name, the

signature is marked "ClusterName". **(b)** Differential signature testing for enriched cell states

comparing non-responders and responders from pre-treatment condition in bladder carcinoma

(BLCA). **(c)** Model of monocytes and macrophage states in tumor and their metabolic

differences. **(d)** Top pathways enriched in different macrophage MeCs. BCC: basal cell

carcinoma, BLCA: bladder carcinoma.

Fig.5.

Fig.5. MetaTiME prioritizes tumor immunity transcriptional regulators. For selected MeCs,

TFs are prioritized by their MeC expression representation and Lisa, ChIP-seq based,

regulatory potentials. X-axis: gene z-weight of the TF for the current MeC. Y-axis: Lisa-based

regulatory potential significance for top genes in the current MeC. Red and orange factors: MeC

regulators prioritized based on both MeC gene weights and Lisa regulatory potential

significance. Green factors: TFs highly weighted in MeCs and not in Lisa analysis. Blue factors:

TFs with high Lisa regulatory potential and not highly weighted in MeCs. **(a)** TFs prioritized for

three MeCs in the signaling category. **(b)** TFs prioritized for three MeCs in the dendritic cell

category. **(c)** TFs prioritized for three MeCs representing different macrophage states. **(d)** TFs

prioritized for three MeCs representing different T cell states. TF: transcription factor, MeC:

meta-component.

Reference

1. Liu, X. S. & Mardis, E. R. Applications of Immunogenomics to Cancer. *Cell* **168**, 600–612

(2017).

2. Jin, M. Z. & Jin, W. L. The updated landscape of tumor microenvironment and drug

repurposing. *Signal Transduct. Target. Ther.* **5**, (2020).

3. Penter, L. *et al.* Local and Systemic Effects of Immune Checkpoint Blockade on Relapsed

Myeloid Malignancies Following Allogeneic Hematopoietic Stem Cell Transplantation.

Blood **136**, 34–35 (2020).

4. Zhang, Y. *et al.* The cancer-associated genetic variant rs3903072 modulates immune

cells in the tumor microenvironment. *Front. Genet.* (2019). doi:10.3389/fgene.2019.00754

5. Gu, S. S. *et al.* Therapeutically Increasing MHC-I Expression Potentiates Immune

Checkpoint Blockade. *Cancer Discov.* **11**, 1524–1541 (2021).

6. Zeng, Z. *et al.* TISMO: syngeneic mouse tumor database to model tumor immunity and

immunotherapy response. *Nucleic Acids Res.* **50**, D1391–D1397 (2022).

7. Jiang, P. *et al.* Signatures of T cell dysfunction and exclusion predict cancer

immunotherapy response. *Nat. Med.* (2018). doi:10.1038/s41591-018-0136-1

8. Yost, K. E. *et al.* Clonal replacement of tumor-specific T cells following PD-1 blockade.

Nat. Med. (2019). doi:10.1038/s41591-019-0522-3

9. Zhang, L. *et al.* Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted Therapies

in Colon Cancer. *Cell* **181**, 442-459.e29 (2020).

10. Cheng, S. *et al.* A pan-cancer single-cell transcriptional atlas of tumor infiltrating myeloid

cells. *Cell* **184**, 792-809.e23 (2021).

11. Wang, X. *et al.* In vivo CRISPR screens identify the E3 ligase Cop1 as a modulator of

macrophage infiltration and cancer immunotherapy target. *Cell* **184**, 5357-5374.e22

(2021).

12.   Wu, S. Z. *et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nat.*

Genet. **53**, 1334–1347 (2021).

13.   Nieto, P. *et al.* A single-cell tumor immune atlas for precision oncology. *Genome Res.* **31**,

1913–1926 (2021).

14.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

15.   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

16.   Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a

transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).

17.   Wang, C. *et al.* Integrative analyses of single-cell transcriptome and regulome using

MAESTRO. *Genome Biol.* **21**, (2020).

18.   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.

Nat. Methods 2015 125 **12**, 453–457 (2015).

19.   Andreatta, M. *et al.* Interpretation of T cell states from single-cell transcriptomics data

using reference atlases. *Nat. Commun.* **12**, 1–19 (2021).

20.   Zheng, A. L., Qin, S., Wang, A., Xing, B. & Gao, R. Pan-Cancer Single Cell Landscape of

Tumor-Infiltrating T Cells. **6474**, (2021).

21.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat.*

Biotechnol. 2018 365 **36**, 411–420 (2018).

22.   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

23.   Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for

single-cell transcriptomics. *Nat. Methods* (2018). doi:10.1038/s41592-018-0229-2

24.   Pelka, K. *et al.* Spatially organized multicellular immune hubs in human colorectal cancer.

Cell **184**, 4734-4752.e20 (2021).

25.   Hyvärinen, A. Fast and robust fixed-point algorithms for independent component analysis.

IEEE Trans. Neural Networks (1999). doi:10.1109/72.761722

26.   Wang, W. *et al.* Independent component analysis based gene co-expression network

inference (ICAnet) to decipher functional modules for better single-cell clustering and

batch integration. *Nucleic Acids Res.* **49**, E54 (2021).

27.   Zheng, R. *et al.* Cistrome Data Browser: expanded datasets and new tools for gene

regulatory analysis. *Nucleic Acids Res.* **47**, D729–D735 (2019).

28.   Sun, D. *et al.* TISCH: A comprehensive web resource enabling interactive single-cell

transcriptome visualization of tumor microenvironment. *Nucleic Acids Res.* **49**, D1420–

D1430 (2021).

29.   Wykes, M. N. & Lewin, S. R. Immune checkpoint blockade in infectious diseases. *Nat.*

Rev. Immunol. 2017 182 **18**, 91–104 (2017).

30.   Zhang, Q. & Vignali, D. A. A. Co-stimulatory and co-inhibitory pathways in autoimmunity.

Immunity **44**, 1034 (2016).

31.   Qin, S. *et al.* Novel immune checkpoint targets: moving beyond PD-1 and CTLA-4. *Mol.*

Cancer 2019 181 **18**, 1–14 (2019).

32.   Thommen, D. S. *et al.* A transcriptionally and functionally distinct PD-1+ CD8+ T cell pool

with predictive potential in non-small-cell lung cancer treated with PD-1 blockade. *Nat.*

Med. 2018 247 **24**, 994–1004 (2018).

33.   Chowell, D. *et al.* Improved prediction of immune checkpoint blockade efficacy across

multiple cancer types. *Nat. Biotechnol. 2021 404* **40**, 499–506 (2021).

34.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–

15550 (2005).

35.   Morgan, D. & Tergaonkar, V. Unraveling B cell trajectories at single cell resolution.

747   *Trends Immunol.* **43**, 210–229 (2022).

748 36. Finkin, S., Hartweger, H., Oliveira, T. Y., Kara, E. E. & Nussenzweig, M. C. Protein

749   Amounts of the MYC Transcription Factor Determine Germinal Center B Cell Division

750   Capacity. *Immunity* **51**, 324-336.e5 (2019).

751 37. Vaquero, J., Aoudjehane, L. & Fouassier, L. Cancer-associated fibroblasts in

752   cholangiocarcinoma. *Curr. Opin. Gastroenterol.* **36**, 63–69 (2020).

753 38. Luca, B. A. *et al.* Atlas of clinically distinct cell states and ecosystems across human solid

754   tumors. *Cell* **184**, 5482-5496.e28 (2021).

755 39. Zhao, J. *et al.* Detection of differentially abundant cell subpopulations in scrna-seq data.

756   *Proc. Natl. Acad. Sci. U. S. A.* **118**, (2021).

757 40. Yuen, K. C. *et al.* High systemic and tumor-associated IL-8 correlates with reduced

758   clinical benefit of PD-L1 blockade. *Nat. Med.* **26**, 693–698 (2020).

759 41. Kaplanov, I. *et al.* Blocking IL-1β reverses the immunosuppression in mouse breast

760   cancer and synergizes with anti-PD-1 for tumor abrogation. *Proc. Natl. Acad. Sci. U. S. A.*

761   **116**, 1361–1369 (2019).

762 42. Duan, Z. & Luo, Y. Targeting macrophages in cancer immunotherapy. *Signal Transduct.*

763   *Target. Ther. 2021 61* **6**, 1–21 (2021).

764 43. Bi, K. *et al.* Tumor and immune reprogramming during immunotherapy in advanced renal

765   cell carcinoma. *Cancer Cell* **39**, 649-661.e5 (2021).

766 44. Zhao, H. *et al.* The role of osteopontin in the progression of solid organ tumour. *Cell*

767   *Death Dis.* **9**, (2018).

768 45. Spivia, W., Magno, P. S., Le, P. & Fraser, D. A. Complement protein C1q promotes

769   macrophage anti-inflammatory M2-like polarization during the clearance of atherogenic

770   lipoproteins. *Inflamm. Res.* **63**, 885–893 (2014).

771 46. Liu, Q. *et al.* The CXCL8-CXCR1/2 pathways in cancer. *Cytokine Growth Factor Rev.* **31**,

772   61–71 (2016).

47.   Qin, Q. *et al.* Lisa: inferring transcriptional regulators through integrative modeling of

public chromatin accessibility and ChIP-seq data. *Genome Biol.* (2020).

doi:10.1186/s13059-020-1934-6

48.   Peters, J. M., Shah, Y. M. & Gonzalez, F. J. The role of peroxisome proliferator-activated

receptors in carcinogenesis and chemoprevention. *Nat. Rev. Cancer 2012 123* **12**, 181–

195 (2012).

49.   Hu, H. *et al.* AnimalTFDB 3.0: a comprehensive resource for annotation and prediction of

animal transcription factors. *Nucleic Acids Res.* **47**, D33–D38 (2019).

50.   Li, H. *et al.* Dysfunctional CD8 T Cells Form a Proliferative, Dynamically Regulated

Compartment within Human Melanoma. *Cell* **176**, 775-789.e18 (2019).

51.   Hyvärinen, A. & Oja, E. Independent component analysis: algorithms and applications.

Neural Networks **13**, 411–430 (2000).

52.   Lotfollahi, M. *et al.* Mapping single-cell data to reference atlases by transfer learning. *Nat.*

Biotechnol. (2021). doi:10.1038/s41587-021-01001-7

53.   Khan, O. *et al.* TOX transcriptionally and epigenetically programs CD8+ T cell

exhaustion. *Nat. 2019 5717764* **571**, 211–218 (2019).

54.   Tang, Q. & Brown, M. A. Abstract A218: Modulating glucocorticoid receptor-mediated

signaling for enhancement of cancer immunotherapy. *Cancer Immunol. Res.* **7**, A218–

A218 (2019).

55.   Ohandjo, A. Q. *et al.* Transcriptome Network Analysis Identifies CXCL13-CXCR5

Signaling Modules in the Prostate Tumor Immune Microenvironment. *Sci. Reports 2019*

91 **9**, 1–13 (2019).

56.   Havenar-Daughton, C. *et al.* CXCL13 is a plasma biomarker of germinal center activity.

Proc. Natl. Acad. Sci. U. S. A. **113**, 2702–2707 (2016).

57.   D'Agostino, R. B., Belanger, A. & D'Agostino, R. B. A Suggestion for Using Powerful and

Informative Tests of Normality. *Am. Stat.* **44**, 316 (1990).

58.   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: An R package for comparing

800   biological themes among gene clusters. *Omi. A J. Integr. Biol.* **16**, 284–287 (2012).

801   59.   Kotliar, D. *et al.* Identifying gene expression programs of cell-type identity and cellular

802   activity with single-cell RNA-Seq. *Elife* **8**, 1–26 (2019).

a

Tumor scRNA
- 1.7 million TME cells

→ ICA

Independent Components
- Signal sources e.g. cell states

→ Clustering & Integration

Meta-Components
- Cell type, cell state, signaling

Dataset 1

Dataset 2

...

Dataset n

b

Functional MeC Landscape for TME

- 86 Functional Annotated MeCs

- Categories

MeC	Top Genes
B_B	CD74,MS4A1,…
T_CD8T-CD3	CD3D,CD8A,…
DC_pDC	GZMB,TCF4,…
M_Macrophage-IL1-NFKB	CXCL8,IL1B,…
Pan_Inteferon_response	STAT1, ISG15,…
…	…

Categories:
- B
- T/NK
- DC
- Mono/Mac
- OtherMyeloid
- Stroma
- Signaling

Cell State Annotation for Tumor scRNA

- MeC Signature Continuum
- Top Enriched State

Tumor Immunity Regulators

- Incorporated Epigenetic Data

- Prioritized TF regulators

MeC	TF
DC_pDC	TCF4,IRF8
M_Macrophage-IL1-NFKB	NFKB
Pan_Inteferon_response	STAT1
…	…