Intracytoplasmic membranes develop in *Geobacter sulfurreducens* under

thermodynamically limiting conditions

Ethan Howley[a], Anna Mangus[b], Dewight Williams[c], César I. Torres[b]

[a] School for Sustainable Engineering and the Built Environment, Arizona State University

[b] School for Engineering Mass Transport and Energy, Arizona State University

[c] Eyring Materials Center, Arizona State University

Abstract

Geobacter sulfurreducens is an electroactive bacterium capable of reducing metal oxides in the

environment and electrodes in engineered systems[1,2]. To respire extracellular electron acceptors

with a wide range of redox potentials, *G. sulfurreducens* has a complex network of respiratory

proteins, many of which are membrane-bound[3–5]. We have identified intracytoplasmic membrane

(ICM) structures in *G. sulfurreducens*. This ICM is an invagination of the inner membrane that

has folded and organized by an unknown mechanism, often but not always located near the tip of

a cell. Using confocal microscopy, we can identify that at least half of the cells contain an ICM

when grown at low potentials, whereas cells grown at higher potentials or using fumarate as

electron acceptor had significantly lower ICM frequency. 3D models developed from cryo-

electron tomograms show the ICM to be a continuous extension of the inner membrane in

contact with the cytoplasmic and periplasmic space. The differential abundance of ICM in cells

grown under different thermodynamic conditions supports the hypothesis that it is an adaptation

to limited energy availability, as an increase in membrane-bound respiratory proteins could

increase electron flux. Thus, the ICM provides extra inner-membrane surface to increase the

abundance of these proteins. *G. sulfurreducens* is the first Thermodesulfobacterium or metal-

oxide reducer found to produce ICMs.

Introduction

While we classically differentiate prokaryotes from eukaryotes by a difference in organelle

compartmentalization of the cytoplasm, reality is more complicated. Prokaryotes with a diverse

variety of metabolisms and ecological niches express various well-defined intracellular

organelles [6–8]. Most of the organelles that have been characterized in prokaryotes fall into one of

two categories. The first are isolated compartments where specialized conditions are maintained

to perform chemical processes not possible in the cytoplasmic space e.g. the anammoxosome[9],

the carboxysome [10], and the acidocalcisome [11]. The second category of prokaryotic organelles

consists of densely packed membrane structures that facilitate higher throughput for membrane-

dependent metabolic processes by increasing the available surface area in a cell e.g. the

thylakoid, the chlorosome[12], and membranous structures in methane, nitrite, and ammonia

oxidizers[13–16]. We use the general term 'intracytoplasmic membrane' (ICM) to describe all these

lipidic structures in prokaryotes as it includes organelles with membranous structures with

unknown functions. For organisms operating with thin thermodynamic margins or performing

slow chemical reactions, the rate of enzyme activity, e.g., ATP production, should be

proportional to the membrane surface area available for those enzymes. In methane-oxidizing

bacteria, for example, two essential membrane-associated metabolic enzymes – methane

monooxygenase and methanol dehydrogenase – are localized in the ICM, providing higher

throughput for a potentially rate-limiting reaction[14,17], and the same has been observed with

ammonia monooxygenase in ammonia-oxidizing bacteria[15]. Interestingly, the relationship

between membrane proteins and ICMs goes both ways, as modifying a bacterium to overexpress

a membrane-bound enzyme can spur ICM-like structures in a bacterium that normally lacks any

organelles [18,19].

Geobacter sulfurreducens is a gram-negative Thermodesulfobacterium (previously classified as

a δ-proteobacterium) that reduces iron and other metals in anaerobic environments[1]. As an

organism adapted to respire insoluble metal oxides in nature, *G. sulfurreducens* is capable of

respiring man-made solid electron acceptors as well[2]. In an engineered system, we can take

advantage of this extracellular electron transfer (EET) to produce a measurable electrical current.

G. sulfurreducens reduces electron acceptors with a wide range of estimated redox potentials[20,21]

(–0.17 [goethite] to +0.98 V vs. SHE [palladium]), produces a relatively high current density in

engineered systems (as high as 10 $A \cdot m^{-2}$)[22], and has a complex network of electron

carriers[5,21,23,24]. In order to adapt as the redox potential of its electron acceptor changes, *G.*

sulfurreducens expresses at least three different electron transfer pathways that each have an

optimal growth condition and distinct electrochemical signal[3,5,21,24]. For these reasons, *G.*

sulfurreducens is considered a model electroactive organism[25].

Metal-reducing bacteria like *G. sulfurreducens* can operate in a relatively energy-limited niche.

G. sulfurreducens oxidizing acetate and reducing natural iron (III) minerals can have as little as

0.12 Volts of redox differential in the case of goethite[26], versus ~1.1 V for aerobic acetate

oxidation. *G. sulfurreducens'* ability to adapt to varying redox conditions depends on a complex

network of electron carriers. This flexible metabolism is strictly dependent on membrane

processes; the inner membrane electron transport chain in *G. sulfurreducens* requires different

electron carrier proteins dependent on the amount of energy available to the cell i.e., the redox

potential of the terminal electron acceptor [5,21,27,28]. When energy-limited by electron acceptors

with low redox potentials, *G. sulfurreducens*' growth could be limited by both a lower rate of

respiration according to Nernstian kinetics[29,30] and by a lower yield of ATP generation per

electron respired. The effect is a membrane-limited respiratory metabolism. This limitation

results in a decreased growth and respiratory rate for *G. sulfurreducens* growing on low redox

potential electron acceptors[27,31].

In this work, we have discovered ICM structures in *G. sulfurreducens*. We used confocal

microscopy, plastic embedded thin-section transmission electron microscopy (TEM), and

cryogenic electron tomography (CryoET), to identify ICM structures in *G. sulfurreducens* that

are localized in specific subcellular regions. By observing cells in different redox conditions, we

can test the hypothesis that ICM in *G. sulfurreducens* is associated with thermodynamic

conditions where cells must respire at low potential. These observations have significant

implications for the holistic understanding of respiration under energy-limited conditions.

<u>Results and Discussion</u>

Morphology of ICM in *G. sulfurreducens* through TEM imaging- In thin plastic sections

prepared via freeze substitution of plunge-frozen cells, we observed ICM structures in *G.*

sulfurreducens that were grown on an anode at -0.07 V vs. SHE. The ICMs mostly appear as

parallel bands of membrane in the cytoplasm and are localized to a fraction of the cell's entire

volume (Figure 1). In some cases, the ICMs were also observed as curved or circular structures.

In most cases, when the TEM section showed the full length of the cell, ICMs mostly appeared

towards one of the tips of the cell (Figure 1B, D). We did not find cells with ICM in more than

one area of the cytoplasm, and not every cell in our plastic sections had evidence of the structure.

We repeated the plastic sectioning procedure with cells grown using fumarate, a soluble electron

acceptor with a higher redox potential ($E'_0 = 0.03$ V vs SHE), and we did not find evidence of

ICM in the resulting micrographs (Figure S1-S4).

Figure 1: Plunge frozen, freeze substituted plastic embedded TEM micrographs of *G.*

sulfurreducens collected from an anode poised at -0.07 V vs. SHE. ICM structures present as

parallel bands in one area of a cell. 3A, 3C, and 3E show ICM in cells sliced perpendicular to the

major axis, while 3B, 3D, and 3F show ICM in cells sliced parallel to the major axis where the

ICM is located near the tip of the cell. Micrographs were collected on an FEI TF20. Scale bars:

100 nm.

Conventional plastic embedded TEM has been used to characterize bacterial ICMs for over 50

years[32]. The ICM of *G. sulfurreducens* shares some morphological characteristics with

previously described structures in other bacteria. Ammonia-oxidizing bacteria (AOB) can

produce ICMs with one continuous membrane folded tightly into parallel bands localized to one

area of the cell, although the ICM in AOB appears to occupy a larger fraction of the cell volume

compared to what we observed in *G. sulfurreducens*[32]. The ICM in AOB forms from

invagination of the inner membrane [32]. In the AOB *Nitrosomonas eutropha*, ICM development is

stimulated by ammonia-oxidizing conditions, but ICM is not developed during anoxic

denitrification[33]. Similarly, we observed the production of ICM in *G. sulfurreducens* in response

to certain environmental redox conditions. In *G. sulfurreducens*, we have not yet determined the

signal mechanism that activates ICM expression or if there are cytoskeletal-like proteins

involved in their organization.

3D structure of ICM- Using CryoET of whole cells, we observed ICM without the artifacts and

membrane damage commonly introduced by dehydration[34]. We grew *G. sulfurreducens* directly

on TEM grids with the grid itself serving as the anode in an electrochemical cell followed by

immediate vitrification. Reconstructions of *G. sulfurreducens* cells display ICM that are not as

tightly packed and regular as what we observed with plastic section TEM (Figure 2, Figure 3).

The difference in ICM appearance between cryoET and plastic embedded thin-section TEM

could be related to artifacts from the dehydration process prior to embedding, or due to a

difference in the growth stage of the biofilm. CryoET revealed that the ICM in *G. sulfurreducens*

has significant variance in morphology. In some cells, the ICM is a loosely organized mass of

membrane structures (Figure 3), but in others it is more regular and composed of smaller units

(Figure 2). This variance may represent different stages of development of the structure, since

cryotomography samples were taken only after 24 hours of introducing an EM grid into the

electrochemical cell, while plastic-section TEM images capture cells that were collected from a

fully developed biofilm. ICMs as observed in cryotomograms were closely associated to the

inner membrane in most cases, and typically near the tip of the cell (Figure 2).

Figure 2: (A) Tomogram slices illustrating how 3D models were created via tomogram

segmentation of ICM located in the tip of a *G. sulfurreducens* cell with the inner membrane,

outer membrane, ICM, and several nanowires modeled from the tomogram. The scale bar is 100

nm. (B,C,D) 3D models of three separate cells displaying ICM near the tip of each cell. These

cells were grown at -0.07 V vs. SHE directly on a grid.

Protein nanowires are important to the respiration of *G. sulfurreducens* [35,36]. We observed

extracellular protein nanowires in most of the cryotomograms that we collected, but it is unclear

if there is a direct relationship between nanowires and ICMs (Figure 2, SI Videos of tomograms).

Some nanowires intersect the outer membrane near the ICM locations, but others do not. As the

ICM is expected to be an area of high metabolic activity, the location of nanowires for

extracellular electron transfer might be preferentially localized near this area.

In a sample grown with a lower potential anode (−0.17 vs. SHE), we can observe a putative

earlier ICM development stage (Figure 3). The ICM is clearly shown to be formed by

invagination of the inner membrane, which is consistent with ICM formation in other bacteria

(e.g, *Rhodobacter sphaeroides*)[37,38]. While a complex ICM network is evident, close inspection

shows all sections to be interconnected with cytoplasmic space within them. Thus, it is a

continuous inner membrane invagination. Most importantly, the periplasmic space is

continuous, providing a path to the outer membrane from all regions of the ICM. This

continuous periplasmic space is most crucial for *G. sulfurreducens'* metabolism of extracellular

respiration, where electrons from the inner membrane must be transported extracellularly,

passing through periplasmic cytochromes along the way[39,40].

Figure 3: Cryotomograph slices and 3D model of a *G. sulfurreducens* cell grown at –0.17 V vs

SHE on a holey-carbon/gold cryo-EM grid as anode for 24 hrs. Cryotomogram shows how ICM

seems to form by invagination of the inner membrane. Top – selected tomogram slices in the z

axis displaying sections of the invagination. Bottom – 3D model of the inner membrane and

outer membrane through (bottom left) the entire thickness of the modeled volume and (bottom

right) the model sliced approximately in half in the z axis to show the ICM profile at a different

depth. Scale bar: 100 nm.

ICM abundance differences- We can also identify the presence of ICM in *G. sulfurreducens*

with confocal microscopy. A similar technique has been used to identify ICM in

methanotrophs[41]. We took confocal images of fixed *G. sulfurreducens* biofilm cells grown at

different anode potentials, or from cell suspensions grown with fumarate as electron acceptor,

and each image had numerous individual cells (Table S1). The ICM is a localized bright area

within a cell when it is stained with Nile red, a lipid-selective fluorescent dye[42] . Some cells have

a single ICM, while others have multiple distinct ICM regions (Figure 4b). Generally, the ICM

are located near the tips of cells, but we also observed ICM in locations throughout the length of

a cell. By using the ImageJ plugin MicrobeJ[43], we can detect cell boundaries and the local

maxima within them that we have identified as ICM. The ICM area as a fraction of total cell area

had a median value of 5.7% in the electrode biofilm grown at -0.07 V and 4.2% in the fumarate

cells, but there is a high variance in fractional area with some cells having over 30% of the area

occupied by ICM (Figure S5 and Figure S6). This is lower than the cell area occupied by ICMs

in methanotrophs[41]. By applying identical image analysis (ICM counting) to cells collected from

different conditions, we found that a change in electron acceptor significantly affects the

frequency of *G. sulfurreducens* cells displaying ICM (Figure 4a). In biofilms grown on anodes at

higher potentials we observed a relatively lower fraction of cells with ICM (41 $\pm$ 4% at -0.03 V

vs. SHE vs. 58 $\pm$ 8% at -0.17 V vs. SHE). Cells grown with fumarate had the lowest incidence of

ICM (14 $\pm$ 10%) despite the fumarate/succinate redox couple having a potential around +0.03 vs.

SHE, similar in redox potential to our highest anode potential studied. In a biofilm, however,

there will be a redox potential gradient because of ohmic losses and diffusion limitations,[22,44,45]

so we anticipate that cells grown with fumarate will experience a higher redox potential on

average than anode biofilm cells on an electrode at a similar potential. Since lower anode

potentials provide less potential energy for growth[24], the higher abundance of ICM may be an

adaptation to energy limitation. Our image analysis used conservative parameters for identifying

ICM within cells, so the absolute frequency of ICM is likely higher.

Figure 4: **a:** Fraction of cells with an ICM in *G. sulfurreducens* under different growth

conditions. Each dot represents a unique image with an average of 113 cells, see Table S1 for

raw counts. The three potentials – -0.17, -0.07, and +0.07 – represent cells collected from anodes

poised at the respective potentials vs. SHE. The 'fumarate' condition cells were grown

planktonically with fumarate as the electron acceptor. Statistical testing consisted of a two-tailed

t-test with Hochberg multiple comparison correction. $*(\rho<0.05)$, $***(\rho\leq0.001)$, $****($

$\rho\leq0.0001)$. **b:** Sum projection of confocal z-stacks of *G. sulfurreducens* cells grown on an

electrode at -0.07 V vs. SHE with examples of ICM annotated. **c:** Sum projection of cells grown

with fumarate as an electron acceptor exemplifying typical cell morphology when ICM is not

present. Both cell images were cropped from larger images taken at 100X magnification using

Nile red as a phospholipid-selective fluorophore.

Significance

G. sulfurreducens has a complex and efficient respiratory metabolism. At the inner membrane,

electrons are trifurcated into different respiration pathways depending on the redox potential of

the terminal electron acceptor[3,5,21]. Being an invagination of the inner membrane, the ICM must

contain the critical inner membrane cytochromes for respiration, but whether respiration is

occurring in the ICM remains to be proven. In ammonia- and methane-oxidizing bacteria, the

rate-limiting respiratory enzymes – ammonia monooxygenase and methane monooxygenase,

respectively – are present within the ICM[15,17]. For organisms with slim thermodynamic margins,

an ICM could allow a higher respiration rate by increasing membrane surface area and the total

number of respiratory proteins. Producing an ICM must be a significant energy investment for a

cell, and the structural organization of the ICM in *G. sulfurreducens* suggests a specialized

function outside of lipid storage. The production of ICMs explains why *G. sulfurreducens* is

higher in lipid content than other gram negative bacteria[46], as the production of ICM in other

bacteria causes elevated lipid fractions[41,47].

In our study we found higher incidence of ICM in cells growing with less thermodynamically

favorable conditions (Figure 4a), consistent with the use of an ICM to increase respiratory rates

in limiting conditions. A respiratory ICM in *G. sulfurreducens* would require a mechanism to

pass electrons to the rest of the extracellular electron transfer network, and if there is an open

connection with the periplasm, as shown in Figure 3, electrons could transfer by diffusion of

periplasmic cytochromes (e.g., PpcA, GSU1996)[39,48]. Yet, transport of these cytochromes from

ICMs to the outer membrane would be a much larger distance than the typical periplasmic space,

since ICMs seem to span the whole cell thickness and have regions that are up to 200 nm from

the outer membrane.

While *G. sulfurreducens* is not the only bacterium with an ICM, nothing like the ICM has been

found in any closely related organisms. To our knowledge, it is the first organism from the

phylum Thermodesulfobacteriota identified to produce an ICM, as well as the first metal oxide

reducer that does so. The ICM in *G. sulfurreducens* provides a good opportunity to study ICM

formation in general, because we can easily control its expression by changing the redox

potential of the electron acceptor, and the polar localization of the ICM in the tip presents an

opportunity to study intracellular organization in bacteria. The varying production of ICMs in *G.*

sulfurreducens suggests that this structure is naturally formed to increase respiratory rates in

thermodynamically and kinetically limiting conditions, a hypothesis that has been proposed

before for other microorganisms[8]. Unlike other bacteria that produce ICMs, *G. sulfurreducens*

cannot complete its entire respiratory pathway within the ICM. Electrons from the ICM in *G.*

sulfurreducens must travel to the outer membrane for extracellular respiration.

G. sulfurreducens has various approaches to optimize energy conservation under limiting and

varying redox conditions. Three characterized pathways seem to be expressed concomitantly

within an electrogenic biofilm to maximize energy conservation[49]. The generation of ICMs at

energy-limiting conditions seem to be an additional tool to maximize energy production within

the cell. Nernstian models that are used to predict rates of respiration in *G. sulfurreducens* would

predict a slow respiration rate at lower potentials[29,50,51] The slower respiration rate is the

consequence of a rate-limiting electron transfer protein, proposed to be at the inner

membrane[29,52] . If *G. sulfurreducens* can increase the amount of this rate-limiting protein by

increasing the amount of inner membrane present, the apparent limitation is alleviated, and

higher respiratory rates can be achieved. Thus, knowledge of how ICMs are used and under

which conditions they are produced will be important to predict rates of electrical current

generation by *G. sulfurreducens.* Future *G. sulfurreducens* ICM studies will likely take

advantage of the genetic tools that have been developed for the organism as well as creative

imaging techniques.

<u>Methods</u>

***G. sulfurreducens* growth**

G. sulfurreducens PCA (ATCC, Virginia USA) was grown from glycerol freezer stocks using

fumarate or an electrode as the electron acceptor as described previously[24]. Fumarate cells were

grown in sealed anaerobic culture tubes with ATCC 1957 medium. Electrode cells were grown

in 100 mL single chamber microbial electrochemical cells on 6-8 cm^2 graphite electrodes

(Graphitestore, Illinois USA) poised at either -0.17, -0.07, or +0.07 V vs. SHE and electrical

current was monitored with a VMP3 potentiostat (BioLogic, Tennessee USA). Each reactor had

an Ag/AgCl reference electrode (BASi, Indiana USA).

TEM

G. sulfurreducens cells from a mature biofilm (~30 days of growth) were fixed in phosphate

buffer containing 4% paraformaldehyde, 2% glutaraldehyde (v/v) and then plunge frozen in

liquid propane before dehydration via freeze substitution in 2% osmium tetroxide dissolved in

acetone on dry ice at -80° C. Cells were slowly returned to room temperature for embedding in

Araldite 502 resin (Ted Pella, Inc.). These sections were cut at 70 nm thickness and secondary

stained with lead acetate to improve contrast All thin section imaging was performed on a Tecnai

TF-20 (FEI) or a Phillips CM12.

CryoET

To grow electrode biofilm cells for cryoET, we designed a holder for fenestrated carbon TEM

grids to insert into bioelectrochemical cells with actively growing cultures. The grids in the

holder are held against a titanium plate that functions as an anode and electron acceptor for the

bacteria. 6 nm BSA conjugated gold nanoparticle fiducials were dried then baked at 60° C to fix

the gold fiducials onto the grids before introduction to the bioreactor. The TEM grids in the

holder were removed after 24 hours and immediately plunge frozen into liquid nitrogen cooled

ethane using an in house designed manual plunge freezer to capture the cells in their active state

(Figure S7 and S8). For the fumarate condition, a suspension of cells grown for 7-10 days was

pipetted onto each grid, blotted to remove excess liquid, and frozen with the FEI Vitrobot Mark

IV.

The frozen hydrated cells were imaged on a Krios G2 (FEI, Oregon USA) at tilts from $-65°$ to

$+65°$ in $2.0°$ angular steps using a dose symmetric collection scheme[53] in regions where

individual cells could be observed over fenestrations in the carbon. Images were collected at a

nominal magnification of 6500x giving a 1.8 Å pixel size in super-resolution mode on the K2

summit camera with a dose rate of 0.5 electron per $Å^2$ * second for three seconds with a frame

rate of 0.2 frames per second (total dose of 100 electrons per $Å^2$). Individual movie frames were

gain corrected, aligned with MotionCor2[54], and sum images were binned by 2 resulting in an

image with a 3.6 Å/pixel scale. Summed images were restacked, then tomographic

reconstruction in eTOMO[55] was performed. Slices from each tomogram were output from IMOD

using the ZAP window saving feature which saves at the native resolution of the monitor. 3D

modeling was performed in IMOD and visualized in ChimeraX[56].

Data Availability

The tomograms used to create figures for this manuscript can be accessed in the EMDB-EBI

repository under accession numbers EMD-27710, EMD-27729, EMD-27748, and EMD-27747.

Confocal microscopy sample preparation and acquisition properties

G. sulfurreducens anode biofilm cells were resuspended, fixed, and imaged between 12-20 days

after current began growing exponentially and was at least 2 A/m^2. *G. sulfurreducens* biofilm

cells were removed from the electrode by gently vortexing in phosphate buffer, pelleted at 4000

RPM for 5 minutes, resuspended in 4% paraformaldehyde for 1 hour, then rinsed and stored in

phosphate buffer at 7 ˚C. Live *G. sulfurreducens* cells grown under fumarate conditions were

imaged 5-7 days after inoculation. Cells under both anode and fumarate conditions were diluted

with 50 mM phosphate buffer and Nile red (ThermoFisher/Invitrogen, red, excitation: 561 nm)

for lipid staining to a final concentration of 2 µg/mL. Samples were allowed to incubate for at

least 15 minutes at room temperature in the dark. Stained cells were imaged on a standard glass

slide, or 2% poly-L-lysine coated glass slide with a 1 ½ cover slip sealed with nail polish.

Fluorescence images were acquired with a Nikon C2+ confocal microscope equipped with a

100X Plan Apo λ (NA 1.45) oil objective using the adjacent NIS Elements software, operated

inside an anaerobic glovebox. Nile red was excited with a 561 nm laser and filtered with a

525/50 561 LP filter cube.

Fluorescence image analysis

Z stacks were processed into sum projections and a Gaussian blur ($\sigma=1$) filter applied for

smoothing in ImageJ. The ImageJ MicrobeJ plug-in[43] was used for bacterial and ICM detection

of *G. sulfurreducens* cells under both anode and fumarate conditions with settings recorded in

Figure S8 and summary output in Table S1. Statistical analysis was performed using the

Student's t-Test corrected for multiple comparisons using the Benjamini-Hochberg method.

Acknowledgements

The funding for this work was provided by the Office of Naval Research (ONR award N0014-

20-1-2269). Confocal microscopy was performed in a Nikon C2+ obtained through an ONR

DURIP grant #N00014-19-1-2531. All electron microscopy was performed at ASU's electron

microscopy core facilities.

3D model manipulation was partially performed with UCSF ChimeraX, developed by the

Resource for Biocomputing, Visualization, and Informatics at the University of California, San

Francisco, with support from National Institutes of Health R01-GM129325 and the Office of

Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious

Diseases.

References

1.   Caccavo, F. *et al.* Geobacter sulfurreducens sp. nov., a Hydrogen- and Acetate-Oxidizing

Dissimilatory Metal-Reducing Microorganism. *Appl. Environ. Microbiol.* **60**, 3752–3759

(1994).

2.   Bond, D. R. & Lovley, D. R. Electricity Production by Geobacter sulfurreducens Attached

to Electrodes. *AEM* **69**, 1548–1555 (2003).

3.   Levar, C. E., Chan, C. H., Mehta-Kolte, M. G. & Bond, D. R. An inner membrane

cytochrome required only for reduction of high redox potential extracellular electron

acceptors. *MBio* **5**, 1–9 (2014).

4.   Zacharoff, L., Chan, C. H. & Bond, D. R. Reduction of low potential electron acceptors

requires the CbcL inner membrane cytochrome of Geobacter sulfurreducens.

Bioelectrochemistry **107**, 7–13 (2016).

5.   Joshi, K., Chan, C. H. & Bond, D. R. Geobacter sulfurreducens inner membrane

cytochrome CbcBA controls electron transfer and growth yield near the energetic limit of

respiration. *Mol. Microbiol.* **116**, 1124–1139 (2021).

6.   Shively, J. M. *Complex Intracellular Structures in Prokaryotes*. (Springer Berlin

Heidelberg, 2006). doi:https://doi.org/10.1007/11497158.

7.   Chowdhury, C., Sinha, S., Chun, S., Yeates, T. O. & Bobik, T. A. Diverse Bacterial

Microcompartment Organelles. *Microbiol. Mol. Biol. Rev.* **78**, 438–468 (2014).

8.   Greening, C. & Lithgow, T. Formation and function of bacterial organelles. *Nat. Rev.*

Microbiol. **18**, 677–689 (2020).

9.   Strous, M. *et al.* Missing lithotroph identified as new planctomycete. *Nature* **400**, 446

(1999).

10.   Shively, J. M., Ball, F. L. & Kline, B. W. Electron microscopy of the carboxysomes

360   (polyhedral bodies) of Thiobacillus neapolitanus. *J. Bacteriol.* **116**, 1405–1411 (1973).

361   11.   Seufferheld, M. *et al.* Identification of organelles in bacteria similar to acidocalcisomes of

362   unicellular eukaryotes. *J. Biol. Chem.* **278**, 29971–29978 (2003).

363   12.   Staehelin, L. A., Golecki, J. R. & Drews, G. SUPRAMOLECULAR ORGANIZATION

364   OF CHLOROSOMES (CHLOROBIUM VESICLES) AND OF THEIR MEMBRANE

365   ATTACHMENT SITES IN CHLOROBIUM LIMICOLA. *Biochim. Biophys. Acta* **589**,

366   30–45 (1980).

367   13.   Davies, S. L. & Whittenbury, R. Fine structure of methane and other hydrocarbon-

368   utilizing bacteria. *J. Gen. Microbiol.* **61**, 227–232 (1970).

369   14.   Fassel, T. A., Buchholz, L. A., Collins, M. L. P. & Remsen, C. C. Localization of

370   methanol dehydrogenase in two strains of methylotrophic bacteria detected by

371   immunogold labeling. *Appl. Environ. Microbiol.* **58**, 2302–2307 (1992).

372   15.   Fiencke, C. & Bock, E. Immunocytochemical localization of membrane-bound ammonia

373   monooxygenase in cells of ammonia oxidizing bacteria. *Arch. Microbiol.* **185**, 99–106

374   (2006).

375   16.   Watson, S. Taxonomic Considerations of the Family Nitrobacteraceae Buchanan. *Int. J.*

376   *Syst. Bacteriol.* **21**, 254–270 (1971).

377   17.   Brantner, C. A., Remsen, C. C., Owen, H. A., Buchholz, L. A. & Collins, M. Intracellular

378   localization of the particulate methane monooxygenase and methanol dehydrogenase in

379   Methylomicrobium album BG8. *Arch. Microbiol.* **178**, 59–64 (2002).

18.   von Meyenburg, K., Jørgensen, B. B. & van Deurs, B. Physiological and morphological

effects of overproduction of membrane-bound ATP synthase in Escherichia coli K-12.

EMBO J. **3**, 1791–1797 (1984).

19.   Weiner, J. H. & Lemire, B. D. Overproduction of Fumarate Reductase in Escherichia.

Microbiology **158**, 590–596 (1984).

20.   Pat-Espadas, A. M., Razo-Flores, E., Rangel-Mendez, J. R. & Cervantes, F. J. Reduction

of palladium and production of nano-catalyst by Geobacter sulfurreducens. *Appl.*

Microbiol. Biotechnol. **97**, 9553–9560 (2013).

21.   Levar, C. E., Hoffman, C. L., Dunshee, A. J., Toner, B. M. & Bond, D. R. Redox potential

as a master variable controlling pathways of metal reduction by Geobacter sulfurreducens.

11, 741–752 (2017).

22.   Torres, C. I., Marcus, A. K. & Rittmann, B. E. Proton transport inside the biofilm limits

electrical current generation by anode-respiring bacteria. *Biotechnol. Bioeng.* **100**, 872–

881 (2008).

23.   Bonanni, P. S., Schrott, G. D. & Busalmen, J. P. A long way to the electrode: how do

Geobacter cells transport their electrons? *Biochem. Soc. Trans.* **40**, 1274–9 (2012).

24.   Howley, E., Krajmalnik-brown, R. & Torres, C. I. Cytochrome expression shifts in

Geobacter sulfurreducens to maximize energy conservation in response to changes in

redox conditions. 1–34 (2022).

25.   Lovley, D. R. *et al. Geobacter. The Microbe Electric's Physiology, Ecology, and*

Practical Applications. Advances in Microbial Physiology vol. 59 (Elsevier Ltd., 2011).

26.   Orsetti, S., Laskov, C. & Haderlein, S. B. Electron transfer between iron minerals and

quinones: Estimating the reduction potential of the Fe(II)-goethite surface from AQDS

speciation. *Environ. Sci. Technol.* **47**, 14161–14168 (2013).

27.   Levar, C. E., Chan, C. H., Mehta-kolte, M. G. & Bond, D. R. An Inner Membrane

Cytochrome Required Only for Reduction of. *MBio* **5**, 02034–14 (2014).

28.   Yoho, R. A., Popat, S. C. & Torres, C. I. Dynamic potential-dependent electron transport

pathway shifts in anode biofilms of Geobacter sulfurreducens. *ChemSusChem* **7**, 3413–

3419 (2014).

29.   Torres, C. I. *et al.* A kinetic perspective on extracellular electron transfer by anode-

respiring bacteria. *FEMS Microbiol. Rev.* **34**, 3–17 (2010).

30.   Andrew Kato Marcus, Cesar I. Torres, B. E. R. Conduction-Based Modeling of the

Biofilm Anode of a Microbial Fuel Cell. *Biotechnol. Bio* (2007) doi:10.1002/bit.

31.   Wei, J., Liang, P., Cao, X. & Huang, X. A new insight into potential regulation on growth

and power generation of geobacter sulfurreducens in microbial fuel cells based on energy

viewpoint. *Environ. Sci. Technol.* **44**, 3187–3191 (2010).

32.   Murray, R. G. & Watson, S. W. Structure of Nitrosocystis Oceanus and Comparison With

Nitrosomonas and Nitrobacter. *J. Bacteriol.* **89**, 1594–1609 (1965).

33.   Schmidt, I., Zart, D. & Bock, E. Effects of gaseous NO2 on cells of Nitrosomonas

eutropha previously incapable of using ammonia as an energy source. *Antonie van*

Leeuwenhoek, Int. J. Gen. Mol. Microbiol. **79**, 39–47 (2001).

34.   Bouchet-Marquis, C. & Hoenger, A. Cryo-electron tomography on vitrified sections: A

critical analysis of benefits and limitations for structural cell biology. *Micron* **42**, 152–162

(2011).

35.　Peng, L. & Zhang, Y. Cytochrome OmcZ is essential for the current generation by

Geobacter sulfurreducens under low electrode potential. *Electrochim. Acta* **228**, 447–452

(2017).

36.　Wang, F. *et al.* Structure of Microbial Nanowires Reveals Stacked Hemes that Transport

Electrons over Micrometers. *Cell* **177**, 361-369.e10 (2019).

37.　Tucker, J. D. *et al.* Membrane invagination in Rhodobacter sphaeroides is initiated at

curved regions of the cytoplasmic membrane, then forms both budded and fully detached

spherical vesicles. *Mol. Microbiol.* **76**, 833–847 (2010).

38.　Woronowicz, K., Harrold, J. W., Kay, J. M. & Niederman, R. A. Structural and functional

proteomics of intracytoplasmic membrane assembly in Rhodobacter sphaeroides. *J. Mol.*

Microbiol. Biotechnol. **23**, 48–62 (2013).

39.　Lloyd, J. R. *et al.* Biochemical and genetic characterization of PpcA, a periplasmic c-type

cytochrome in Geobacter sulfurreducens. *Biochem. J.* **369**, 153–161 (2003).

40.　Aklujkar, M. *et al.* Proteins involved in electron transfer to Fe(III) and Mn(IV) oxides by

Geobacter sulfurreducens and Geobacter uraniireducens. *Microbiol. (United Kingdom)*

159, 515–535 (2013).

41.　Whiddon, K. T., Gudneppanavar, R., Hammer, T. J., West, D. A. & Konopka, M. C.

Fluorescence-based analysis of the intracytoplasmic membranes of type I methanotrophs.

Microb. Biotechnol. **12**, 1024–1033 (2019).

42.   Greenspan, P., Mayer, E. P. & Fowler, S. D. Nile red: A selective fluorescent stain for

intracellular lipid droplets. *J. Cell Biol.* **100**, 965–973 (1985).

43.   Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput

bacterial cell detection and quantitative analysis. *Nat. Microbiol.* **1**, 1–7 (2016).

44.   Popat, S. C. & Torres, C. I. Critical transport rates that limit the performance of microbial

electrochemistry technologies. *Bioresour. Technol.* **215**, 265–273 (2016).

45.   Bond, D. R., Strycharz-Glaven, S. M., Tender, L. M. & Torres, I. C. I. On Electron

Transport through Geobacter Biofilms. *ChemSusChem* **5**, 1099–1105 (2012).

46.   Howley, E., Ki, D., Krajmalnik-brown, R. & Torres, C. I. Geobacter sulfurreducens '

unique metabolism results in cells with a high iron and lipid content. **2**, 1–28 (2022).

47.   Patt, T. E. & Hanson, R. S. Intracytoplasmic membrane, phospholipid, and sterol content

of Methylobacterium organophilum cells grown under different conditions. *J. Bacteriol.*

134, 636–644 (1978).

48.   Pokkuluri, P. R. *et al.* Structure of a novel dodecaheme cytochrome c from Geobacter

sulfurreducens reveals an extended 12nm protein with interacting hemes. *J. Struct. Biol.*

174, 223–233 (2011).

49.   He, X., Chadwick, G., Jiménez Otero, F., Orphan, V. & Meile, C. Spatially Resolved

Electron Transport through Anode-Respiring Geobacter sulfurreducens Biofilms: Controls

and Constraints. *ChemElectroChem* 1–13 (2021) doi:10.1002/celc.202100111.

50.   Kato Marcus, A., Torres, C. I. & Rittmann, B. E. Conduction-based modeling of the

biofilm anode of a microbial fuel cell. *Biotechnol. Bioeng.* **98**, 1171–1182 (2007).

51.   Richter, H. *et al.* Cyclic voltammetry of biofilms of wild type and mutant Geobacter

sulfurreducens on fuel cell anodes indicates possible roles of OmcB, OmcZ, type IV pili,

and protons in extracellular electron transfer. *Energy Environ. Sci.* **2**, 506–516 (2009).

52.   Lusk, B. G. *et al.* PH Dependency in Anode Biofilms of Thermincola ferriacetica

Suggests a Proton-Dependent Electrochemical Response. *J. Am. Chem. Soc.* **140**, 5527–

5534 (2018).

53.   Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron

tomography tilt-scheme optimized for high resolution subtomogram averaging. *J. Struct.*

Biol. **197**, 191–198 (2017).

54.   Zheng, S. Q. *et al.* MotionCor2: anisotropic correction of beam-induced motion for

improved cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).

55.   Mastronarde, D. N. & Held, S. R. Automated tilt series alignment and tomographic

reconstruction in IMOD. *J. Struct. Biol.* **197**, 102–113 (2017).

56.   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,

and developers. *Protein Sci.* **30**, 70–82 (2021).