**NF-κB-Inducing Kinase (NIK) Governs the Mitochondrial Respiratory Capacity, Differentiation, and Inflammatory Status of Innate Immune Cells**

Justin N. Keeney[1], Ashley Winters[2], Raquel Sitcheran[1]*, A. Phillip West[2]*

[1]Department of Cell Biology and Genetics & [2]Department of Microbial Pathogenesis and Immunology, School of Medicine, Texas A&M University, Bryan, TX, USA

**Running Title**: NIK controls metabolism and function of innate immune cells.

**Funding**: This work was supported by NIH grants R01HL148153 to A.P.W. and R01NS082554 to R.S. Additional support was provided by a pilot award from the Texas A&M School of Medicine and an X-Grant from the Texas A&M University Office of the Vice President for Research.

***Correspondence**: Address correspondence and reprint requests to Dr. Raquel Sitcheran (sitcheran@tamu.edu) or Dr. A. Phillip West (awest@tamu.edu).

**Keywords**: NIK, macrophage, myeloid cells, innate immunity, mitochondria, oxidative metabolism, inflammation

## Abstract

NF-κB-Inducing Kinase (NIK), which is essential for the activation of the noncanonical NF-κB pathway, regulates diverse processes in immunity, development, and disease. While recent studies have elucidated important functions of NIK in adaptive immune cells and cancer cell metabolism, the role of NIK in metabolic-driven inflammatory responses in innate immune cells remains unclear. Here, we demonstrate that NIK-deficient bone marrow-derived macrophages exhibit defects in mitochondrial-dependent metabolism and oxidative phosphorylation (OXPHOS), which impairs the acquisition of a pro-repair, anti-inflammatory phenotype. Subsequently, NIK-deficient mice exhibit skewing of myeloid cells characterized by aberrant eosinophil, monocyte, and macrophage cell populations in the blood, bone marrow, and adipose tissue. Furthermore, NIK-deficient blood monocytes display hyperresponsiveness to bacterial lipopolysaccharide and elevated TNFα production ex vivo. These findings suggest that NIK governs metabolic rewiring, which is critical for balancing pro- and anti-inflammatory myeloid immune cell function. Overall, our work highlights a previously unrecognized role for NIK as a molecular rheostat that fine-tunes immunometabolism in innate immunity and suggests that metabolic dysfunction may be an important driver of inflammatory diseases caused by aberrant NIK expression or activity.

**Key Points:**

- NIK-deficient macrophages exhibit impaired mitochondrial oxidative phosphorylation
- NIK-deficient mice have more inflammatory myeloid cells in blood and bone marrow

43 • NIK-dependent metabolic rewiring shapes pro- and anti-inflammatory innate immunity

## Introduction

Macrophages are important components of the innate immune system that play key roles in immunity, tissue homeostasis, and cancer (1). Macrophages exist across a spectrum of polarization states in vivo and can adopt pro-inflammatory or anti-inflammatory properties depending on local cytokine gradients and tissue residency (2). Murine bone marrow-derived macrophages (BMDMs) can be polarized toward the extreme end of the pro-inflammatory spectrum by culture in bacterial lipopolysaccharide (LPS) or interferon gamma. These so-called M1 or M(LPS) macrophages produce pro-inflammatory cytokines such as TNFα, have bactericidal and phagocytic functionality, and can produce reactive oxygen species (ROS) and nitric oxide (NO) (3). Conversely, BMDMs cultured in interleukin-4 (IL-4) are skewed toward the anti-inflammatory end of the spectrum. These so-called M2 or M(IL-4) macrophages can dampen inflammatory responses by secreting anti-inflammatory cytokines such as interleukin-10 (IL-10), promote extracellular matrix remodeling, and enhance stromal cell proliferation (3). It is well appreciated that metabolic switches promote the acquisition of distinct macrophage polarization states and play pivotal roles in the final effector cell functionality (4). M1 macrophages upregulate glycolysis, have a broken tricarboxylic acid (TCA) cycle, and display impaired oxidative phosphorylation (OXPHOS), leading to enhanced generation of TCA intermediates such as succinate and other metabolites necessary for ROS and NO generation (5–7). On the other hand, M2 macrophages have an intact TCA cycle and upregulate OXPHOS and fatty acid oxidation to generate anti-inflammatory products such as glucocorticoids, IL-10, IL-13, etc. (8, 9). These metabolic shifts are critical for macrophage function, as small molecules that inhibit the major metabolic enzymes in glycolysis, TCA, and OXPHOS can dramatically alter polarization states in vitro and in vivo (10–12).

Nuclear factor-kappaB (NF-κB) signaling has a well-established role in inflammatory processes with the canonical NF-κB pathway rapidly responding to inflammatory stimuli (13). Alternatively the non-canonical NF-κB pathway, which requires NF-κB Inducing Kinase (NIK; also known as MAP3K14), regulates immune cell differentiation, development, and tissue homeostasis (13–16). Over the last decade, NIK has been found to play many novel roles in a diverse set of immune cells and their progenitors (17, 18). Within the bone marrow, NIK promotes mitochondrial biogenesis during osteoclastogenesis (14). In microglia, which are tissue-resident macrophages of the brain, NIK is important for the induction of cytokine expression, which aids in T-cell recruitment in an animal model of experimental autoimmune encephalomyelitis (19, 20).

Recently, we and others have demonstrated roles for NIK in regulating mitochondrial function and metabolism in cancer cells (19, 21–23). For example, we previously reported that NIK regulates mitochondrial dynamics (21) and promotes oxidative metabolism (22) in glioblastoma, an aggressive form of brain cancer. Moreover, NIK-deficient glioblastoma cells exhibit impaired mitochondrial fission and increased glycolysis to compensate for diminished oxidative metabolism (21, 22) and are less invasive. Notably, we have shown that these novel metabolic roles of NIK are largely independent of non-canonical NF-κB signaling, suggesting that NIK plays a direct role in regulating mitochondrial function. Another recent study showed that NIK shapes the metabolism of CD8 T cells independently of non-canonical NF-κB by stabilizing hexokinase 2 and promoting NADPH generation (23). Similar results were found in B cells containing a deletion of TRAF3, an upstream inhibitor of NIK (24). Although NIK has emerged as a key regulator of tumor cell and lymphocyte metabolism, whether NIK regulates immunometabolic switches in myeloid innate immune cells remains unclear.

Studies using NIK knockout (NIK$^{KO}$) mice have revealed that NIK deficiency promotes a

dramatic increase in peripheral blood eosinophil abundance (25). NIK$^{KO}$ mice develop a

hypereosinophilic inflammatory disease driven by the infiltration of pathogenic eosinophilic

granulocytes into diverse organs, including skin, liver, lung, and esophagus, leading to early

death (25, 26). Recent studies have shown that eosinophils upregulate OXPHOS upon activation

(27); however, it remains unclear whether immunometabolic switches contribute to the

inflammatory myeloid phenotypes observed in NIK$^{KO}$ mice. To begin to address this knowledge

gap, we measured mitochondrial and metabolic parameters in BMDMs from NIK$^{KO}$ mice. Here,

we show that NIK promotes oxidative metabolism in macrophages and that loss of NIK results in

hyperinflammatory myeloid skewing of the bone marrow, blood, and other tissues. Our findings

demonstrate that NIK is an important molecular rheostat that maintains mitochondrial function to

control innate immune cell function, and they suggest that metabolic alterations may contribute

to diseases where NIK activity is dysregulated.

**Materials and methods**

**Mice**

All animal experiments were performed in accordance with animal use protocols with approved IACUC guidelines at Texas A&M University. *Map3k14* null mice were purchased from The Jackson Laboratory (strain 025557) and maintained by heterozygous breeding. ==*Map3k14* floxed mice were obtained from Genentech and bred with LysMCre mice (Jackson Lab strain 004781).== Equivalent numbers of male and female mice were used for all experiments. Animal numbers used are described in figure legends.

**Reagents**

A comprehensive list of antibodies, reagents, and primers is provided in Tables 1-3, respectively.

**Flow Cytometry**

*Blood and spleen immune phenotyping:* Whole mouse blood was collected in sodium heparin tubes. Spleens were collected and filtered through a 100uM cell strainer. RBCs were lysed twice with ACK lysis buffer, and leukocytes were subjected to LPS stimulation (1 μg/ml) in the presence of brefeldin A and monensin for 4 hours. Fc receptors were blocked with anti-mouse CD16/CD32 (2.4G2) antibody, and cells were stained with antibodies against surface proteins, permeabilized with Foxp3/Transcription Factor Staining Buffer Kit (TNB-0607-KIT, Tonbo Biosciences, San Diego, CA, USA), and stained with antibodies against intracellular proteins. Cells were analyzed with a Cytek Aurora 5L Spectral Analyzer. The results were plotted and analyzed using FlowJo software (BD Bioscience; Franklin Lakes, NJ, USA).

*Bone marrow phenotyping:* Cells were isolated from the tibia and femurs of mice as described

below (see Cell Culture). After an ACK lysis step, 500,000 cells per mouse were taken for

immune cell phenotyping and 1 million cells were stained for the stem cell analysis. Cells were

stained in a 96 well plate with Live/Dead Fix Violet (eBioscience, Waltham, MA, USA) at 4°C

for 20 minutes. After being washed thoroughly, cells were blocked with FC blocker

(eBioscience, Waltham, MA, USA) for 10 minutes at 4°C before the addition of the primary

antibody cocktail which incubated with the cells for 30 minutes. After, cells were washed and

fixed with 4% paraformaldehyde for 10 minutes at room temperature in the dark. Flow

cytometric analysis was performed on a BD Fortessa X-20 (BD Bioscience; Franklin Lakes, NJ,

USA). The results were plotted and analyzed using FlowJo software (BD Bioscience; Franklin

Lakes, NJ, USA).

*Fat and small intestine immune phenotyping:* Visceral fat deposits and a 5cm section of the

ileum were harvested from mice and washed with PBS before being placed in a 0.2% (2mg/mL)

collagenase 2 solution, (Worthington Biochemical; Lakewood, NJ, USA) diluted in RIPA buffer

(Gibco, ThermoFisher Scientific; Waltham, MA, USA), under contestant stirring for 60 minutes

at 37°C. The isolated cells were filtered through a 70 μM cell strainer and red blood cells were

lysed with ACK lysis buffer (VWR; Radnor, PA, USA). Cells were then stained as described

above for the bone marrow.

*Mitochondrial flow cytometry assays:* This assay was previously described (28), but in brief:

NIK$^{WT}$ and NIK$^{KO}$ BMDMs were treated with 20ng/mL IL-4 for 5 hours before cells were

detached using PBS containing 1mM EDTA (Gibco, ThermoFisher Scientific; Waltham, MA,

USA). Cells were then stained with a combination of Mitotracker Green FM, Mitotracker Red

FM, Mitosox, Anti-MO CD11b, and Violet live dead dye for 15 minutes in a 37°C tissue culture

incubator. Cells were then washed twice before analysis on a BD Fortessa X-20 flow cytometer.

The results were plotted and analyzed using FlowJo software (BD Bioscience; Franklin Lakes,

NJ, USA).

**Cell Culture**

L929 and HEK293T cells were obtained from the American Type Culture Collection (ATCC)

and maintained in Dulbecco's Modified Eagle's Medium (DMEM; Gibco, ThermoFisher

Scientific; Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS; VWR;

Radnor, PA, USA). BMDMs were generated by crushing the tibia and femurs of unpooled mice.

After lysis of red blood cells with ACK lysis buffer (VWR; Radnor, PA, USA) cells were filtered

through a 40 μM cell strainer and grown on petri plates in DMEM containing 10% FBS and 30%

L929 conditioned media for 7 days with fresh L929 containing media media being added on day

4 post plating. IL-4 (PeproTech; Rocky Hill, NJ, USA) was used at a concentration of 20ng/mL

for *in vitro* experiments unless otherwise stated. Human U937 monocytic cells were a gift from

Dr. Kristin Patrick and maintained in Roswell Park Memorial Institute 1640 Medium (RPMI

1640; Gibco, ThermoFisher Scientific, Waltham, MA, USA) supplemented with 10% FBS. To

differentiate U937 monocytes into macrophages, cells were treated with 100ng/mL phorbol 12-

myristate 13-acetate (PMA; AmBeed, Arlington Hts, IL, USA) for 48 hours. After that time

period the media was changed, and cells were treated similarly as BMBMs. All cells were

cultured at 37°C with 95% humidity and 5% $CO_2$.

**Lentiviral production**

Twenty-four micrograms of each Lenti-CrispR-V2 plasmids containing a guide RNA against

NIK and 72□μg of polyethyleneimine were used to transfect HEK293T cells. After 3 days of

transfection, viral supernatant was harvested and filtered through a 0.45-μM syringe filter. After

filtration, viral particles were concentrated 20-fold to 400□μl using Lenti-X Concentrator

(Clontech, #631231, Mountain View, CA), and 100□μl of concentrated virus was used to infect

cells. 1 μg/mL Puromycin (InvivoGen; San Diego, CA, USA) was used for selection until no

non-transduced cells were alive.

**CRISPR-Cas9 gene knockout**

U937 cells were transduced with a mixture of Lenti-CrispR-v2 virus' carrying three gRNAs for

each target. The gRNA sequences for human NIK were previously described (20, 21). Loss of

NIK was confirmed by immunoblot of puromycin-resistant cells. Single colony cells were

isolated by serial dilution. All experiments were repeated with at least three knockdown clones

**Seahorse Extracellular Flux Analysis**

The Seahorse XFe96 Analyzer (Agilent; Santa Clara, CA, USA) was used to measure

mitochondrial respiration. 5 x $10^4$ cells were plated per well in at least triplicate in 100 μL of

DMEM media containing 10%FBS in an Agilent Seahorse XF96 Cell Culture Microplate. After

allowing the cells to settle at room temperature for one hour, cells were moved to a 37°C

incubator for 3 hours until stimulated overnight with 20 ng/mL IL-4. After 16 hours the cells

were washed and replaced with XF assay medium (Base Medium Minimal DMEM

supplemented with 200mM L-glutamine pH 7.4) and incubated with a 37°C incubator without

CO2 for an hour before analysis. The Oxygen Consumption Rate (OCR) was measured after

sequential addition of 25 mM glucose, 1.5 μM Oligomycin A (Sigma Aldrich; Saint Louis, MO,

USA), 1.5 μM FCCP (Sigma Aldrich; Saint Louis, MO, USA) plus 1 mM Sodium Pyruvate

(Gibco, ThermoFisher Scientific; Waltham, MA, USA), and 2.5 μM Antimycin A (Sigma

Aldrich; Saint Louis, MO, USA) plus 1.25 μM Rotenone (Enzo Life Sciences, Farmingdale, NY,

USA) following a previously published protocol (29).

*NIK Inhibition:* Same workflow as above with the exception of after allowing the cells to settle,

DMSO or 5μM B022 (MedChem Express, Monmouth Junction, NJ, USA) were added to the

cells and allowed to incubate for up to 5 hours before treating the cells overnight with 20ng/mL

IL-4.

**RNA Sequencing**

$1 \times 10^6$ $NIK^{WT}$ and $NIK^{KO}$ BMDMs were stimulated with 20ng/mL IL-4 or PBS for 6 hours

before being lifted with PBS containing 1mM EDTA (Gibco, ThermoFisher Scientific; Waltham,

MA, USA). Cells were spun down and washed once with PBS before being flash frozen in liquid

nitrogen. Cell pellets were shipped on dry ice to Genewiz (South Plainfield, NJ, USA) where

they isolated the RNA, constructed the sequencing library, and conducted the sequencing. RNA-

sequencing data was analyzed using the Salmon pipeline (version 1.4.0-gompi-2020b) on the

Texas A&M High Performance Research Computing Cluster. Salmon was used to align the

results to the reference genome *Mus muculus* GRCm39. The data was then processed through

DESeq2 (version 1.32.0) on R Studio where the annotated gene expression files were used to

determine statistically significant changes in gene expression in stimulated $NIK^{KO}$ BMDMs

relative to stimulated NIK$^{WT}$ BMDMs. Ingenuity Pathway Analysis software (IPA; QIAGEN,

Germantown, MD, USA) was used to identify gene families and potential upstream regulators

within the datasets. GraphPad Prism (San Diego, CA, USA) was used to make heatmaps. Raw

and analyzed RNASeq datasets can be accessed at the NCBI GEO, accession number XXXXXX.

(the GEO submission is pending and will be updated as soon as approved)

**Gene Expression**

5x10$^5$ NIK$^{WT}$ and NIK$^{KO}$ BMDMs were stimulated with 20ng/mL IL-4 for 6 hours before being

lifted with PBS containing 1mM EDTA (Gibco, ThermoFisher Scientific; Waltham, MA, USA).

Cells were spun down and washed once with PBS before total RNA was isolated from cells by

Purelink™ RNA Mini Kit (Life Technologies). cDNA was synthesized from 1 µg total RNA

using iScript reverse transcription supermix (Bio-Rad, Hercules, CA) following the

manufacturer's protocol. Quantitative RT-PCR was performed using Universal SYBR Green

Supermix with StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA).

Data was graphed as a 2$^{\Delta\Delta CT}$ fold change with the gene expression data being normalized to the

house keeping gene RPS18.

**Immunofluorescence**

BMDMs were grown on 12mm coverslips at $6 \times 10^5$ cells/well. Cells were treated with 200ng/ml

LPS for 24 hours. After washing in PBS, cells were fixed with 4% paraformaldehyde for 20 min,

permeabilized with 0.1% Triton X-100 in PBS for 5 min, blocked with PBS containing 5% FBS

for 30 min, stained with primary antibodies for 1 hour, and stained with secondary antibodies for

1 hour. Cells were washed with PBS containing 5% FBS between each step Coverslips were

mounted with Prolong Gold anti-fade reagent containing DAPI (Molecular Probes). Cells were

imaged on an Olympus FV3000 confocal laser scanning microscope using a 100X oil-immersed

objective.

**Mitochondrial isolation**

$2 \times 10^6$ NIK$^{WT}$ and NIK$^{KO}$ BMDMs were plated in a 6 well plate and treated as described

previously (22). At the time of harvest, cells were washed with PBS and scraped into a 1.5mL

tube before spinning down. PBS was replaced with 0.5mL isolation buffer (320mM Sucrose,

1mM EDTA, 10mM HEPES, pH 7.2 with KOH and stored at 4°C) containing protease inhibitors

(ThermoFisher Scientific; Waltham, MA, USA). While on ice, cells in isolation buffer were

passed through a 27-gauge needle 10 times before spinning at 700xg for 8 minutes at 4°C.

Supernatant was removed and placed into a new tube and kept on ice while the pellet was

resuspended in 0.5mL isolation buffer with protease inhibitors before spinning at 700xg for 5

minutes at 4°C. Supernatant was removed and added with the previous supernatant, pellet

contains nucleus, and the 1mL of supernatant was spun down at 17,000xg for 11 minutes at 4°C.

After spinning the supernatant was placed into a new tube labeled cytoplasm and the pellet was

resuspended in isolation buffer and spun down at 17,000xg for 11 minutes at 4°C. Following this

last spin, the supernatant was discarded, and the mitochondrial pellet was resuspended in 40uL

isolation buffer and protein concentration was measured by BCA assay using the Victor X3

microplate reader.

**Electron transport chain activity assays**

Electron transport chain activity assays were obtained from Cayman Chemical Company (Ann

Arbor, MI, USA) and manufacturer's directions were followed with the following changes. 50uL

of tube A was added to each well in the 96 well plate but activity buffer was used in place of the

supplied mitochondria. We then added 5ug of mitochondrial protein (isolation described above)

to a total volume of 20uL activity buffer per well. At this point each well should have 70uL per

well before adding 30uL of tube B to start the reaction. Reactions were conducted at 25°C using

an Epoch microplate reader with readings taken every 45 seconds for 20 minutes.

**ELISA**

$2 \times 10^6$ NIK$^{WT}$ or NIK$^{KO}$ BMDMs were plated in a 6 well plate and allowed to adhere overnight.

The following day the cells were treated with PBS or 20ng/mL IL-4 for 24 hours. Media was

then removed from the cells, spun down, and froze at -20°C until ready for analysis (cells were

used for other experiments). IL-10 ELISA matched antibodies were obtained from Tonbo

Biosciences (San Diego, CA, USA) and the ELISA was carried out according to the

manufacturer's instructions. TMB Substrate was used to detect the peroxidase-labeled secondary

antibody in the ELSIA kit (KPL, Gaithersburg, MD, USA). 100uL of undiluted media was used

and ran in duplicate for each biological replicate.

**Macrophage Heptotaxis Assay**

This assay was described previously (30). In brief, NIK$^{WT}$ and NIK$^{KO}$ BMDMs were stimulated

with IL-4 (20ng/mL) for 24 hours before lifted with PBS containing 1mM EDTA (Gibco,

ThermoFisher Scientific; Waltham, MA, USA). 75,000 DiO (ThermoFisher Scientific; Waltham,

MA, USA) labeled BMDMs were plated on the insert of an 8.0μm pore, polyester membrane

285 Transwell Permeable Support (Corning Incorporated; Corning, NY, USA) which was precoated

286 for 3 hours at 37°C with 4ug/mL fibrinogen (Sigma-Aldrich; Saint Louis, MO, USA). After

287 letting the cells settle for 3 hours, the media was removed and a Matrigel Matrix (50% Matrigel,

288 49% Culture media, 1% FBS) was added on top of the cell layer (Corning Incorporated; Corning,

289 NY, USA). Once the Matrigel matrix solidified, a solution of 100nM

290 N-Formyl-Met-Leu-Phe [FMLP] (Sigma-Aldrich; Saint Louis, MO, USA) was added to the top

291 of the Matrigel to induce a chemokine gradient. Cells were allowed to migrate for 24 hours

292 before z-stack images were acquired on a Nikon A1R Confocal microscope (Nikon Instruments;

293 Melville, NY, USA). Images were then processed in Imaris 9.5.0 (Oxford Instruments;

294 Abingdon, Oxfordshire, UK) before counting the number of cells invaded. Data was graphed as a

295 fold change to $NIK^{WT}$ of the number of total cells that invaded.

**Macrophage Adhesion Assay**

298 This assay was described previously (30). In brief, $NIK^{WT}$ and $NIK^{KO}$ BMDMs were stimulated

299 with IL-4 (20ng/mL) for 24 hours before being lifted with PBS containing 1mM EDTA (Gibco,

300 ThermoFisher Scientific; Waltham, MA, USA). Cells were diluted to a concentration of 1

301 million cells/mL and 50uL of cell solution was plated on a precoated 4ug/mL fibrinogen and

302 0.5% polyvinyl alcohol (Sigma-Aldrich; Saint Louis, MO, USA) 96-well plate. Cells were plated

303 with technical replicates and allowed to sit for 30 minutes in a 37°C tissue culture incubator.

304 Cells were then washed once with PBS before staining the adhered cells with the CyQUANT

305 DNA stain according to the manufacturer's instructions (Invitrogen, ThermoFisher Scientific;

306 Waltham, MA, USA). Fluorescence signal was measured on the Victor X3 microplate reader and

307 cell number was determined through a standard curve of cells.

**Macrophage Phagocytosis Assay**

NIK$^{WT}$ and NIK$^{KO}$ BMDMs were plated subconfluently in a 6-well plate before being stimulated

with IL-4 (20ng/mL) for 24 hours. $10^8$ 1um carboxylate microspheres (Polysciences Inc;

Warrington, PA, USA) were coated in a 50% mouse serum (MP Biomedicals; Irvine, CA, USA)

solution and incubated for 30 minutes at 37˚C. The solution was then diluted to 5% serum before

being added to cells and allowed to incubate for 2 hours. After, cells were washed twice with

PBS before fixing with 4%PFA. Once the PFA was washed, cells were stained with Phalloidin

and Hoechst (Invitrogen, ThermoFisher Scientific; Waltham, MA, USA) and imaged on a Nikon

Eclipse Ti microscope.

**Statistical Analysis**

320  Statistical analyses were carried out using GraphPad Prism (San Diego, CA, USA), and details

321  can be found in the figure legends where applicable. The data presented herein were considered

322  statistically significant at p□<□0.05.

**Results**

**NIK regulates macrophage metabolism by shaping mitochondrial respiratory capacity and OXPHOS complex activity.**

To investigate how NIK regulates innate immune cell function, we examined metabolic shifts during the polarization of bone marrow-derived macrophages (BMDMs) obtained from wild-type (NIK$^{WT}$) and NIK knockout (NIK$^{KO}$) mice. After differentiation in L929-conditioned media for 7 days, we noted that M0 macrophages from NIK$^{KO}$ mice displayed reduced basal and maximal respiration, as well as impaired mitochondrial spare respiratory capacity (SRC) (**Figure 1A**). Upon treatment with interleukin-4 (IL-4) to stimulate an M2-like phenotype (M(IL-4)), NIK$^{KO}$ BMDMs were unable to increase basal and maximal respiration, abrogating SRC and indicating a pronounced and persistent defect in oxidative metabolism (**Figure 1B**). Decreased maximal respiration and SRC were also observed in M0 and M(IL-4) NIK CRISPR knockout (NIK$^{-/-}$) human monocytic U937 cells after their differentiation into macrophages (**Figure 1C-E**). Similar decreases in basal and maximal respiration were observed in NIK conditional knockout BMDMs (LysMCre; NIK$^{fl/fl}$) and in NIK$^{WT}$ BMDMs treated with the NIK inhibitor B022 (31) (**Supplemental Figure 1**). Taken together, these data indicated that NIK regulates oxidative metabolism by controlling OXPHOS and maximal respiration.

To gain insight into the underlying OXPHOS deficit in NIK$^{KO}$ BMDMs, we performed RNA sequencing on M0 and M(IL-4) skewed BMDMs. We did not observe significant changes in the expression of genes involved in glycolysis, TCA cycle, pentose phosphate pathway, glucose transporters, or electron transport chain subunits (**Supplemental Figure 2**), indicating that loss of respiratory capacity in NIK$^{KO}$ BMDMs occurs at a post-transcriptional level. However, we did note a significant increase in mitochondrial superoxide production in both M0

and M(IL-4) NIK$^{KO}$ BMDMs compared to NIK$^{WT}$ (**Figure 2A,B**), indicating enhanced electron

leak from inefficient respiratory complexes. Mitochondrial reactive oxygen species (mROS) can

be produced from OXPHOS complexes I, II, and III, so we next performed enzymatic assays

with isolated mitochondria to assess respiratory complex efficiency. Consistent with our mROS

data, we observed that NIK$^{KO}$ BMDMs exhibited impaired activity of Complexes I and II

compared to NIK$^{WT}$ BMDMs (**Figure 2C,D**). This apparent electron transport chain defect

correlated with defects in other markers of mitochondrial fitness, including increased numbers of

mitochondria in the NIK$^{KO}$ BMDMs with reduced membrane potential (**Supplemental Figure**

**3A-C**) and a trend toward increased mitochondrial size (**Supplemental Figure 3E-G**).

Complexes III-V exhibited no changes in activity between NIK$^{WT}$ and NIK$^{KO}$, regardless of

treatment with IL-4 (**Figure 2E-G**). Similar to our previous findings in cancer cells (21, 22),

NIK localized to the mitochondria after treatment with IL-4 (**Supplemental Figure 3D**). These

data suggest that mitochondria-associated NIK controls oxidative metabolism and electron

transport chain function through OXPHOS Complexes I and II.

**NIK$^{KO}$ Macrophages Fail to Polarize to an M(IL-4) Phenotype *in vitro* and *in vivo*.**

362   To investigate the downstream pathways impacted by the dysregulated metabolism

observed in the absence of NIK, we performed transcriptome analysis of NIK$^{WT}$ and NIK$^{KO}$

BMDMs, and evaluated transcriptional changes in macrophage phenotype signatures. We

observed that there were significant defects in M2-related wound healing gene signatures in the

NIK$^{KO}$ BMDMs compared to NIK$^{WT}$ in response to IL-4 treatment (**Figure 3A**). The RNA-

sequencing data were verified via qPCR for changes in expression of typical IL-4 response

genes, such as Arginase 1 (Arg1) and CD163, and a significant decrease in expression was

observed in Arg1 in the NIK$^{KO}$ BMDMs compared to NIK$^{WT}$ (**Figure 3B**). Consistent with this

finding, we observed a significant decrease in IL-10 secretion to the cell culture media in

response to IL-4 treatment (**Figure 3C**). Additionally, after IL-4 treatment, we observed that

NIK$^{KO}$ BMDMs harbored functional defects in migration and adhesion (**Figure 3D,E**), although

their phagocytic capabilities remained intact (**Figure 3F**). Taken together with the metabolic

findings discussed above, these data suggest that NIK$^{KO}$ BMDMs fail to fully polarize into an

M(IL-4) phenotype due to a deficiency in OXPHOS activity.

Eosinophils have been shown to secrete IL-4 and promote an M(IL-4) macrophage

phenotype *in vivo* (32–35). Increased peripheral and tissue eosinophil numbers have been

previously described in NIK$^{KO}$ mice and are associated with dermatitis and esophageal

eosinophilia (25, 26). Because little is known about the role of NIK in macrophages, we

investigated the relationship between the accumulated eosinophils and their effect on

macrophage polarization in visceral white adipose tissue (WAT) deposits, which are known to

have low eosinophil numbers and high levels of macrophages, along with the small intestine,

which has high eosinophil numbers and low levels of macrophages (33, 35, 36). It was

previously reported that NIK$^{KO}$ mice exhibit decreased body weight compared to NIK$^{WT}$

controls, which also correlates with decreased fat mass and smaller adipocytes (37). We found

that NIK$^{KO}$ mice had a significant accumulation of lineage positive immune cells within the

WAT compared to NIK$^{WT}$ littermates and further analysis of the immune populations revealed a

significant increase in macrophages and a similar trend in eosinophils (**Figure 3G-J**). In contrast,

total immune cells, macrophages, and eosinophils were either unchanged or decreased in the

NIK$^{KO}$ ileum compared to NIK$^{WT}$ (**Figure 3K-N**). Taken together, the changes in immune

populations within WAT suggest persistent inflammation due to high levels of macrophages and
eosinophils.

**NIK<sup>KO</sup> Mice Have Increased Levels of Pro-Inflammatory Myeloid Cells in both the Blood**
**and Bone Marrow.**

Given the changes in immune populations observed in the WAT and small intestine in
NIK$^{KO}$ mice, we performed high dimensional multiparameter flow cytometry and standard
gating to phenotype circulating immune cell populations. Consistent with prior findings and roles
for NIK in B cell development and function (15, 16, 24, 38), we observed markedly reduced
CD19+ B lymphocytes, but not CD3+ T cells, in the blood of NIK$^{KO}$ mice (**Figure 4A-B**). We
also observed a large increase in the percentage of CD11b+ myeloid cells, which was
characterized by a shift from neutrophils to eosinophils and Ly6C$^{hi}$ inflammatory monocytes
(**Figure 4C-F**). Next, using t-Distributed Stochastic Neighbor Embedding (tSNE) analysis to
investigate myeloid immune marker expression more broadly, we confirmed that NIK$^{KO}$ mice
contain dramatically more inflammatory eosinophils (**Figure H, green dashed line**) (25, 26).
These myeloid cells were characterized as side scatter high (SSC-A), CD11b+, Ly6C+, Ly6G$^{lo}$,
F4-80+, and high expression of TNFα at rest (**Figure 4G-I**). We also observed that NIK$^{KO}$ mice
possess elevated TNFα+, Ly6C$^{hi}$ inflammatory monocytes in the blood at rest (**Figure 4G-I,**
**black dashed line**). Ex vivo LPS stimulation of blood leukocytes revealed a greater
enhancement of CD11b+ Ly6C$^{hi}$ inflammatory monocytes in NIK$^{KO}$ mice compared to NIK$^{WT}$
(**Figure 4J**). Four hours after LPS stimulation, CD11b+ Ly6C$^{hi}$ monocytes from NIK$^{KO}$ mice
produced very high levels of intracellular TNFα relative to NIK$^{WT}$ monocytes **(Figure 4J,K,L).**
Overall, these results reveal that NIK$^{KO}$ mice have a significant increase in blood myeloid cells

that are both basally producing pro-inflammatory cytokines and hyper-responsive to innate

immune stimuli, suggesting unresolved systemic inflammation.

Given the role that NIK plays a role in hemopoietic stem cell differentiation (17, 18), we

next investigated the bone marrow of NIK$^{KO}$ mice for defects in immune populations. We found

that NIK$^{KO}$ mice have a significant increase in the total number of bone marrow cells (**Figure**

**5A**), which are primarily derived from increased CD45+ lineage positive cells (**Figure 5B,C**).

Analysis of CD45+ cells in NIK$^{KO}$ mice showed a significant increase in granulocytes compared

to blast cells, monocytes, and lymphocytes (**Figure 5D-G**). Moreover, we found that the increase

in granulocytes is due to an increase in eosinophils as opposed to neutrophils (**Figure 5H,I**).

Consistent with our blood findings, we observed that monocytic cells were slightly elevated

(**Figure 5J**), total T cells were unchanged (**Figure 5K**), and B cells were significantly decreased

(**Figure 5L**). These data suggest that dysregulated terminal differentiation of cell populations in

circulation and the bone marrow stem from an enrichment of low proliferative cells within the

stem cell niche of the bone marrow in NIK$^{KO}$ mice (**Supplemental Figure 4**). Taken together,

our findings demonstrate that NIK$^{KO}$ mice exhibit a systemic skewing toward myeloid and pro-

inflammatory phenotypes in the bone marrow, circulation, and peripheral tissues, driven at least

in part by dysregulated mitochondrial homeostasis and oxidative metabolism (**Figure 5M,N**).

**Discussion**

Although prior studies have established a critical role for NIK in controlling adaptive immunity and development (15, 16, 23, 24, 38), much less is known about whether and how NIK regulates innate immune functions. While we have demonstrated that NIK acts as a stress sensor to regulate both basal and stress-induced mitochondrial metabolism to promote glioblastoma pathogenesis (22), it was not clear whether NIK played similar roles in non-cancerous immune cells. In this study, we identify a previously unreported requirement for NIK in regulating immunometabolic switches in myeloid cells to promote their acquisition of a pro-repair, anti-inflammatory immune phenotype. Specifically, we observed that BMDMs isolated from NIK$^{KO}$ mice exhibited impaired OXPHOS, with a significant reduction in the maximal respiratory and spare respiratory capacities, and the inability to fully acquire an M(IL-4) polarization state, as seen by a failure to induce M2 markers (see **Figure 3**). Similarly, decreased mitochondrial respiration was also observed in BMDMs isolated from mice with a conditional deletion of NIK (LysMCre NIK$^{fl/fl}$, **see Supplemental Figure 1**), as well as NIK knockout human monocytic U937 cells differentiated into macrophages, indicating cell intrinsic mitochondrial functions for NIK. Furthermore, we observed that pharmacological inhibition of NIK in BMDMs also impaired mitochondrial respiration, indicating a requirement for the NIK kinase functionality in innate immune cell metabolic regulation. On a systemic level, we observed that NIK$^{KO}$ mice exhibit a skewing toward pro-inflammatory myeloid phenotypes in the bone marrow, blood, and adipose tissue, characterized by increased numbers of macrophages, eosinophils, and monocytes expressing high levels of TNFα. Since both cell-intrinsic and cell-extrinsic NIK deficiency affects systemic inflammation (25), future studies are required to address whether the phenotypic changes observed in myeloid cell populations in NIK$^{KO}$ mice are controlled cell-autonomously.

Our findings are the first to demonstrate a requirement for NIK in myeloid cell

mitochondrial metabolism and macrophage polarization through regulation of electron transport

chain function. Moreover, while we noted a decrease in bone marrow and blood CD45+ B

lineage cells in NIK[KO] mice as previously reported (38), we demonstrate a significant increase in

eosinophils and CD11b+, Ly6C[hi] monocytes expressing elevated TNFα at baseline and after LPS

challenge. These findings suggest that previously reported hyper-eosinophilia and dermatitis in

NIK[KO] mice are associated with an inability of eosinophils to induce OXPHOS in macrophages

to resolve inflammation (25, 26). Interestingly, development of eosinophilia in NIK[KO] mice was

shown to be independent of IKK/NF-κB transcriptional activity (25). This is consistent with our

RNA-seq data confirming no significant transcriptional alterations in metabolic or macrophage

polarization profiles in NIK[KO] BMDMs, and suggests that the regulation of macrophage

mitochondrial metabolism and TCA complex activity does not occur at a transcriptional level.

Moreover, our findings are consistent with studies demonstrating that control of oxidative

metabolism and mitochondrial dynamics in GBM cells (21, 22), glycolytic metabolism in T cells

(23), and ATP production and respiratory capacity in B cells (24) are NIK dependent and

independent of IKK/NF-κB dependent transcriptional regulation.

The most prominent reported phenotypes of NIK[KO] mice, as well as mice harboring a

spontaneous *NIK* mutation (NIK[aly/aly]), are their lack of lymph nodes and Peyer's patches,

aberrant splenic and thymic architecture, leading to profound defects in adaptive immunity and

increased susceptibility to infection (15, 39, 40). Primary Immunodeficiency Disease (PID)

patients who harbor loss-of-function mutations in *NIK* have remarkably similar clinical

manifestations to NIK[KO] mice, including lymphopenia, abnormal splenic architecture, and

markedly reduced immunoglobulin levels leading to recurrent infections (15, 39–42). PID

resulting from NIK mutation has been largely attributed to aberrant NF-κB transcriptional regulatory activity leading to adaptive immunodeficiency. How loss of NIK function alters innate immune cell development and function in patients with PID is unclear, and the impact of NIK dysregulation on immunometabolic homeostasis in PID has not been previously considered. We have recently observed that NIK$^{KO}$ mice exhibit altered systemic metabolism characterized by an increase in aerobic glycolysis and impaired respiration (37), which is similar to patients with primary mitochondrial diseases (MD) caused by mutations in OXPHOS genes. Patients with MD often experience recurrent infections and display adaptive immunodeficiency (43, 44), paralleling those with NIK loss-of-function PID. We and others have found that mouse models of human mitochondrial disease display pronounced myeloid skewing in the bone marrow and blood, characterized by high levels of inflammatory neutrophils and monocytes (45, 46). Moreover, we observed that the Polymerase gamma (POLG) 'mutator' model of mitochondrial DNA disease exhibits OXPHOS deficits and a switch to glycolytic metabolism, which drive enhanced pro-inflammatory phenotypes, macrophage polarization, and tissue damage (46). Myeloid phenotypes in these mitochondrial mutant mice resemble findings here with NIK$^{KO}$ mice. Taken together, these findings suggest that mitochondrial and metabolic dysfunction leading to innate immune rewiring could be a common feature of patients with NIK-deficient PID, Leigh Syndrome or other primary mitochondrial diseases. Future research to more thoroughly investigate these emerging links is warranted.

In conclusion, our findings elucidate a previously unrecognized role for NIK as a molecular rheostat that shapes the metabolic status, differentiation, and inflammatory phenotypes of myeloid cells in vitro and in vivo. Moreover, our work identifies NIK as a key factor governing innate immune cell mitochondrial function and suggests that metabolic dysfunction

may be an important contributor to inflammatory and/or immunodeficiency diseases caused by

aberrant NIK expression or activity. Our findings have important implications for human disease,

as the bioenergetic pathways underlying immune cell phenotype and function are increasingly

recognized as important nodes for therapeutic intervention in immune and inflammatory

disorders. Moreover, our results provide a strong rationale for future analysis of overlapping

phenotypes in inborn errors of immunity and inborn errors of metabolism and suggest that

strategies aimed at boosting mitochondrial function may help rebalance the immune system in

primary immunodeficiency caused by NIK loss of function.

**Acknowledgements**

We thank Dr. Kristin Patrick for providing U937 cells.

**Disclosures**

The authors declare no financial conflicts of interest.

**References**

1. Shapouri-Moghaddam, A., S. Mohammadian, H. Vazini, M. Taghadosi, S.-A. Esmaeili, F.

Mardani, B. Seifi, A. Mohammadi, J. T. Afshari, and A. Sahebkar. 2018. Macrophage plasticity,

polarization, and function in health and disease. *J. Cell. Physiol.* 233: 6425–6440.

2. Murray, P. J., J. E. Allen, S. K. Biswas, E. A. Fisher, D. W. Gilroy, S. Goerdt, S. Gordon, J. A.

Hamilton, L. B. Ivashkiv, T. Lawrence, M. Locati, A. Mantovani, F. O. Martinez, J.-L. Mege, D.

M. Mosser, G. Natoli, J. P. Saeij, J. L. Schultze, K. A. Shirey, A. Sica, J. Suttles, I. Udalova, J.

A. van Ginderachter, S. N. Vogel, and T. A. Wynn. 2014. Macrophage Activation and

Polarization: Nomenclature and Experimental Guidelines. *Immunity* 41: 14–20.

3. Orecchioni, M., Y. Ghosheh, A. B. Pramod, and K. Ley. 2019. Macrophage Polarization:

Different Gene Signatures in M1(LPS+) vs. Classically and M2(LPS−) vs. Alternatively

Activated Macrophages. *Front. Immunol.* 10: 1–14.

4. Viola, A., F. Munari, R. Sánchez-Rodríguez, T. Scolaro, and A. Castegna. 2019. The

Metabolic Signature of Macrophage Responses. *Front. Immunol.* 10: 1–16.

5. Ahmed, D., D. Roy, A. Jaworski, A. Edwards, A. Abizaid, A. Kumar, A. Golshani, and E.

Cassol. 2019. Differential remodeling of the electron transport chain is required to support TLR3

and TLR4 signaling and cytokine production in macrophages. *Sci. Rep.* 9: 1-13.

6. Scialò, F., D. J. Fernández-Ayala, and A. Sanz. 2017. Role of Mitochondrial Reverse Electron

Transport in ROS Signaling: Potential Roles in Health and Disease. *Front. Physiol.* 8: 1–7.

7. Palmieri, E. M., M. Gonzalez-Cotto, W. A. Baseler, L. C. Davies, B. Ghesquière, N. Maio, C.

M. Rice, T. A. Rouault, T. Cassel, R. M. Higashi, A. N. Lane, T. W.-M. Fan, D. A. Wink, and D.

W. McVicar. 2020. Nitric oxide orchestrates metabolic rewiring in M1 macrophages by targeting

aconitase 2 and pyruvate dehydrogenase. *Nat. Commun.* 11: 1–17.

8. Zhang, Q., H. Wang, C. Mao, M. Sun, G. Dominah, L. Chen, and Z. Zhuang. 2018. Fatty acid

oxidation contributes to IL-1β secretion in M2 macrophages and promotes macrophage-mediated

tumor cell migration. *Mol. Immunol.* 94: 27–35.

9. Mills, E. L., and L. A. O'Neill. 2016. Reprogramming mitochondrial metabolism in

macrophages as an anti-inflammatory signal. *Eur. J. Immunol.* 46: 13–21.

10. Fan, J., K. A. Krautkramer, J. L. Feldman, and J. M. Denu. 2015. Metabolic regulation of

histone post-translational modifications. *ACS Chem. Biol.* 10: 95–108.

11. Rath, M., I. Müller, P. Kropf, E. I. Closs, and M. Munder. 2014. Metabolism via Arginase or

Nitric Oxide Synthase: Two Competing Arginine Pathways in Macrophages. *Front. Immunol.* 5:

1–10.

12. Makowski, L., M. Chaib, and J. C. Rathmell. 2020. Immunometabolism: From basic

mechanisms to translation. *Immunol. Rev.* 295: 5–14.

13. Pflug, K. M., and R. Sitcheran. 2020. Targeting NF-κB-Inducing Kinase (NIK) in Immunity,

Inflammation, and Cancer. *Int. J. Mol. Sci.* 21: 1–19.

14. Zeng, R., R. Faccio, and D. V. Novack. 2015. Alternative NF-κB Regulates RANKL-

Induced Osteoclast Differentiation and Mitochondrial Biogenesis via Independent Mechanisms.

*J. Bone Miner. Res. Off. J. Am. Soc. Bone Miner. Res.* 30: 2287–2299.

15. Yin, L., L. Wu, H. Wesche, C. D. Arthur, J. M. White, D. V. Goeddel, and R. D. Schreiber.

2001. Defective lymphotoxin-beta receptor-induced NF-kappaB transcriptional activity in NIK-

deficient mice. *Science* 291: 2162–2165.

16. Hahn, M., A. Macht, A. Waisman, and N. Hövelmeyer. 2016. NF-κB-inducing kinase is

essential for B-cell maintenance in mice. *Eur. J. Immunol.* 46: 732–741.

17. Zhao, C., Y. Xiu, J. Ashton, L. Xing, Y. Morita, C. T. Jordan, and B. F. Boyce. 2012.

Noncanonical NF-κB signaling regulates hematopoietic stem cell self-renewal and

microenvironment interactions. *Stem Cells Dayt. Ohio* 30: 709–718.

18. Xiu, Y., W. Y. Xue, A. Lambertz, M. Leidinger, K. Gibson-Corley, and C. Zhao. 2017.

Constitutive Activation of NIK Impairs the Self-Renewal of Hematopoietic Stem/Progenitor

Cells and Induces Bone Marrow Failure. *STEM CELLS* 35: 777–786.

19. Jie, Z., C.-J. Ko, H. Wang, X. Xie, Y. Li, M. Gu, L. Zhu, J.-Y. Yang, T. Gao, W. Ru, S.-J.

Tang, X. Cheng, and S.-C. Sun. 2021. Microglia promote autoimmune inflammation via the

noncanonical NF-κB pathway. *Sci. Adv.* 7: 1–13.

20. Duran, C. L., D. W. Lee, J.-U. Jung, S. Ravi, C. B. Pogue, L. G. Toussaint, K. J. Bayless, and

R. Sitcheran. 2016. NIK regulates MT1-MMP activity and promotes glioma cell invasion

independently of the canonical NF-κB pathway. *Oncogenesis* 5: 1–12.

21. Jung, J.-U., S. Ravi, D. W. Lee, K. McFadden, M. L. Kamradt, L. G. Toussaint, and R.

Sitcheran. 2016. NIK/MAP3K14 Regulates Mitochondrial Dynamics and Trafficking to Promote

Cell Invasion. *Curr. Biol.* 26: 3288–3302.

22. Kamradt, M. L., J.-U. Jung, K. M. Pflug, D. W. Lee, V. Fanniel, and R. Sitcheran. 2021. NIK

promotes metabolic adaptation of glioblastoma cells to bioenergetic stress. *Cell Death Dis.* 12:

1–18.

23. Gu, M., X. Zhou, J. H. Sohn, L. Zhu, Z. Jie, J.-Y. Yang, X. Zheng, X. Xie, J. Yang, Y. Shi,

H. D. Brightbill, J. B. Kim, J. Wang, X. Cheng, and S.-C. Sun. 2021. NF-κB-inducing kinase

maintains T cell metabolic fitness in antitumor immunity. *Nat. Immunol.* 22: 193–204.

24. Mambetsariev, N., W. W. Lin, A. M. Wallis, L. L. Stunz, and G. A. Bishop. 2016. TRAF3

deficiency promotes metabolic reprogramming in B cells. *Sci. Rep.* 6: 1–9.

25. Häcker, H., L. Chi, J. E. Rehg, and V. Redecke. 2012. NIK Prevents the Development of

Hypereosinophilic Syndrome-like Disease in Mice Independent of IKKα Activation. *J. Immunol.*

188: 4602–4610.

26. Eden, K., D. E. Rothschild, D. K. McDaniel, B. Heid, and I. C. Allen. 2017. Noncanonical

NF-κB signaling and the essential kinase NIK modulate crucial features associated with

eosinophilic esophagitis pathogenesis. *Dis. Model. Mech.* 10: 1517–1527.

27. Porter, L., N. Toepfner, K. R. Bashant, J. Guck, M. Ashcroft, N. Farahi, and E. R. Chilvers.

2018. Metabolic Profiling of Human Eosinophils. *Front. Immunol.* 9: 1–11.

28. Monteiro, L. de B., G. G. Davanzo, C. F. de Aguiar, and P. M. M. Moraes-Vieira. 2020.

Using flow cytometry for mitochondrial assays. *MethodsX* 7: 1–9.

29. Van den Bossche, J., J. Baardman, and M. P. J. de Winther. 2015. Metabolic Characterization

of Polarized M1 and M2 Bone Marrow-derived Macrophages Using Real-time Extracellular Flux

Analysis. *J. Vis. Exp. JoVE* 1–7.

30. Cui, K., C. L. Ardell, N. P. Podolnikova, and V. P. Yakubenko. 2018. Distinct Migratory

Properties of M1, M2, and Resident Macrophages Are Regulated by αDβ2 and αMβ2 Integrin-

Mediated Adhesion. *Front. Immunol.* 9: 1–14.

31. Ren, X., X. Li, L. Jia, D. Chen, H. Hou, L. Rui, Y. Zhao, and Z. Chen. 2017. A small-

molecule inhibitor of NF-κB-inducing kinase (NIK) protects liver from toxin-induced

inflammation, oxidative stress, and injury. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 31:

711–718.

32. Rose, C. E., J. A. Lannigan, P. Kim, J. J. Lee, S. M. Fu, and S. J. Sung. 2010. Murine lung

eosinophil activation and chemokine production in allergic airway inflammation. *Cell. Mol.*

*Immunol.* 7: 361–374.

33. Yoon, J., H.-N. Um, J. Jang, Y.-A. Bae, W.-J. Park, H. J. Kim, M.-S. Yoon, I. Y. Chung, and

Y. Jung. 2019. Eosinophil Activation by Toll-Like Receptor 4 Ligands Regulates Macrophage

Polarization. *Front. Cell Dev. Biol.* 7: 1–14.

34. Qiu, Y., K. D. Nguyen, J. I. Odegaard, X. Cui, X. Tian, R. M. Locksley, R. D. Palmiter, and

A. Chawla. 2014. Eosinophils and type 2 cytokine signaling in macrophages orchestrate

development of functional beige fat. *Cell* 157: 1292–1308.

35. Wu, D., A. B. Molofsky, H.-E. Liang, R. R. Ricardo-Gonzalez, H. A. Jouihan, J. K. Bando,

A. Chawla, and R. M. Locksley. 2011. Eosinophils sustain adipose alternatively activated

macrophages associated with glucose homeostasis. *Science* 332: 243–247.

36. Jung, Y., and M. E. Rothenberg. 2014. Roles and Regulation of Gastrointestinal Eosinophils

in Immunity and Disease. *J. Immunol. Baltim. Md 1950* 193: 999–1005.

37. Pflug, K. M., D. W. Lee, and R. Sitcheran. 2021. NF-κB-Inducing Kinase Rewires Metabolic

Homeostasis and Promotes Diet-Induced Obesity. 1-51.

38. Brightbill, H. D., J. K. Jackman, E. Suto, H. Kennedy, C. Jones, S. Chalasani, Z. Lin, L.

Tam, M. Roose-Girma, M. Balazs, C. D. Austin, W. P. Lee, and L. C. Wu. 2015. Conditional

Deletion of NF-κB-Inducing Kinase (NIK) in Adult Mice Disrupts Mature B Cell Survival and

Activation. *J. Immunol. Baltim. Md 1950* 195: 953–964.

39. Dejardin, E., N. M. Droin, M. Delhase, E. Haas, Y. Cao, C. Makris, Z.-W. Li, M. Karin, C.

F. Ware, and D. R. Green. 2002. The lymphotoxin-beta receptor induces different patterns of

gene expression via two NF-kappaB pathways. *Immunity* 17: 525–535.

40. Weih, F., and J. Caamaño. 2003. Regulation of secondary lymphoid organ development by

the nuclear factor-kappaB signal transduction pathway. *Immunol. Rev.* 195: 91–105.

41. Schlechter, N., B. Glanzmann, E. G. Hoal, M. Schoeman, B.-S. Petersen, A. Franke, Y.-L.

Lau, M. Urban, P. D. van Helden, M. M. Esser, M. Möller, and C. Kinnear. 2017. Exome

Sequencing Identifies a Novel MAP3K14 Mutation in Recessive Atypical Combined

Immunodeficiency. *Front. Immunol.* 8: 1–15.

42. Willmann, K. L., S. Klaver, F. Doğu, E. Santos-Valente, W. Garncarz, I. Bilic, E. Mace, E.

Salzer, C. Domínguez Conde, H. Sic, P. Májek, P. P. Banerjee, G. I. Vladimer, Ş. Haskoloğlu,

M. Gökalp Bolkent, A. Küpesiz, A. Condino-Neto, J. Colinge, G. Superti-Furga, W. F. Pickl, M.

C. van Zelm, H. Eibel, J. S. Orange, A. Ikincioğulları, and K. Boztuğ. 2014. Biallelic loss-of-

function mutation in NIK causes a primary immunodeficiency with multifaceted aberrant

lymphoid immunity. *Nat. Commun.* 5: 1–13.

43. McGuire, P. J. 2019. Mitochondrial Dysfunction and the Aging Immune System. *Biology* 8:

1–10.

44. Kapnick, S. M., S. E. Pacheco, and P. J. McGuire. 2018. The emerging role of immune

dysfunction in mitochondrial diseases as a paradigm for understanding immunometabolism.

*Metabolism.* 81: 97–112.

45. Jin, Z., W. Wei, M. Yang, Y. Du, and Y. Wan. 2014. Mitochondrial Complex I Activity

Suppresses Inflammation and Enhances Bone Resorption by Tipping the Balance of

Macrophage-Osteoclast Polarization. *Cell Metab.* 20: 483–498.

46. Lei, Y., C. Guerra Martinez, S. Torres-Odio, S. L. Bell, C. E. Birdwell, J. D. Bryant, C. W.

Tong, R. O. Watson, L. C. West, and A. P. West. 2021. Elevated type I interferon responses

potentiate metabolic dysfunction, inflammation, and accelerated aging in mtDNA mutator mice.

*Sci. Adv.* 7: 1–16.

**Figure 1: NIK regulates oxidative metabolism in macrophages.** Seahorse extracellular flux

analysis investigating changes in the oxygen consumption rate (OCR) of basal (M0) NIK$^{WT}$ and

NIK$^{KO}$ BMDMs **(A)** or after an overnight treatment with IL-4 M(IL-4) **(B)**. Line graphs are

combined technical (n > 23) and biological replicates with bar graphs representing the averaged

technical replicate value for each animal (n > 6 biological replicates). Data are mean +/- SEM

and statistics are a Student's T-test *p < 0.05, **p < 0.01, ***p < 0.001. Seahorse extracellular

flux analysis showing **(C)** OCR readouts of basal (M0) and **(D)** IL-4 treated (M(IL-4)) control

and NIK$^{-/-}$ U937 cells. Graphs represent 3 independent experiments with treatment conditions ran

in at least technical triplicates (n = 3 single cell isolated clones). Data are mean +/- SEM and

statistics are a Student's T-test. *p < 0.05, **p < 0.01. **(E)** Representative western blot verifying

loss of NIK in U937 cells using the empty CRISPR vector, LCV2, as a control. Samples were

treated with 200ng/mL LPS and 10μM MG132 for 4 hours before extracting protein with RIPA

buffer.

**Figure 2: NIK deficient macrophages display elevated mROS and impaired OXPHOS**

**Complex I and II activity. (A,B)** Representative histograms and quantification of Mitosox

levels in NIK$^{WT}$ and NIK$^{KO}$ BMDMs post 6-hour treatment with IL-4 (n = 5 biological

replicates). Statistics are a Student's T-test. *p < 0.05, **p < 0.01. Electron transport chain

activity assays for Complex I **(C)**, Complex II **(D)**, Complex III **(E)**, Complex IV **(F)**, and

Complex V **(G)** using isolated mitochondria from basal or 24 hour IL-4 treated NIK$^{WT}$ and

NIK$^{KO}$ BMDMs. Graphs represent biological replicates (n > 3 biological replicates ran in at least

duplicate). Statistics are a Student's T-test. *p < 0.05.

**Figure 3: NIK$^{KO}$ macrophages fail to polarize to an M(IL-4) phenotype *in vitro* and *in vivo*.**

**(A)** RNA-sequencing data of classical pro-repair pathways in NIK$^{KO}$ macrophages after 6 hours

of IL-4 treatment. Volcano plots represent 2 biological replicates with a Log2Fold change of

NIK$^{KO}$ BMDMs compared back to NIK$^{WT}$ of the same treatment. Statistics were calculated by

DESeq2 and genes above the dotted line are deemed significant for having a -log(p-value) of

greater than 1.3. **(B-C)** The trends in RNA-sequencing were confirmed by assessing IL-4

response genes in by RT-qPCR **(B)** and validating IL-10 levels by ELISA post 24hr IL-4

treatment in NIK$^{WT}$ and NIK$^{KO}$ macrophages **(C)** (n > 3 biological replicates ran in triplicates).

Statistics are a Student's T-test *p < 0.05, ***p < 0.001. NIK$^{WT}$ and NIK$^{KO}$ BMDMs were

stimulated for 24 hours with IL-4 before undergoing the following assays: **(D)** Cell invasion

assay was performed with media containing the chemoattractant FMLP. After 24 hours invaded

cells were calculated percent migrated compared to NIK$^{WT}$ (n > 4 biological replicates).

Statistics are a Student's T-test *p < 0.05. **(E)** Cells were plated on a PVA coated 96 well plate

and allowed to settle for 30 minutes before washing with PBS and quantifying percent attached

cells via CyQUANT DNA staining and comparing the NIK$^{KO}$ value back to NIK$^{WT}$ (n = 3

biological replicates). Statistics are a Student's T-test *p < 0.05. **(F)** Representative fluorescent

images of phagocytosis assay. Graph represents % BMDMs containing beads (n = 3 biological

replicates). Statistics are a Student's T-test. *In vivo* analysis of NIK$^{WT}$ and NIK$^{KO}$ mice

investigating immune cells in the WAT **(G-J)** and ileum **(K-N)** gated through single and live

cells. Quantification of biological replicates analyzing total immune cells, macrophages, and

eosinophils in WAT (top) and ileum (bottom) is graphed as a percent of CD45+ parent gate (n =

4). Statistics are a Student's T-test. *p < 0.05.

**Figure 4: NIK$^{KO}$ mice exhibit increased pro-inflammatory myeloid cells in the blood at**

**baseline and after innate immune stimulation.** Blood was harvested from NIK$^{WT}$ and NIK$^{KO}$

mice and subjected to red blood cell lysis with ACK buffer. Total blood leucocytes were

quantified by flow cytometry using markers of B cells **(A)**, T cells **(B)**, CD11b+ myeloid cells **(C)**, neutrophils **(D)**, eosinophils **(E)**, and monocytes **(F)**. (n = 4 biological replicates). Statistics are a Student's T-test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$. **(G-I)** T-SNE clustering of NIK$^{WT}$ and NIK$^{KO}$ blood leukocytes representing 4 biological replicates. **(J-L)**. Blood leukocytes were stimulated with LPS ex vivo and TNFα mean fluorescence intensity (MFI) levels were quantified. Statistics are a Student's T-test. $**p < 0.01$.

**Figure 5: NIK$^{KO}$ mice exhibit myeloid skewing in the bone marrow. (A)** Total bone marrow cell count after ACK red blood cell lysis (n = 23 biological replicates). Statistics are a Student's T-test $**p < 0.01$. **(B,C)** Quantification and flow cytometry plot of lineage positive (CD45+) cells in NIK$^{WT}$ and NIK$^{KO}$ mouse bone marrow (n = 7). Statistics are a Student's T-test $**p < 0.01$. **(D-G)** Quantification of different bone marrow cell populations in NIK$^{WT}$ and NIK$^{KO}$ bone marrow based off of Side Scatter Area (SSC-A) and CD45 (n = 7). **(H-L)** Quantification of specific immune populations in the CD45+ parent gate utilizing the indicated surface markers. Statistics are a Student's T-test. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. Model showing the role of NIK in WT myeloid cells promoting oxidative metabolism and having a homeostatic balance of innate immune cells **(M)** and the consequences of NIK loss resulting in decreased oxidative metabolism and systemic inflammation **(N)**. Model Created with BioRender.com.

**Supplemental Figure 1: Conditional knockout and chemical inhibition of NIK reduce maximal respiration.** Seahorse extracellular flux analysis investigating changes in the oxygen consumption rate (OCR) of basal (M0) LysMCre;NIK$^{+/+}$ and LysMCre;NIK$^{fl/fl}$ BMDMs **(A)** or after an overnight treatment with IL-4 M(IL-4) **(B)**. Line graphs are combined technical (n > 19) and biological replicates with bar graphs representing each animal (n = 5 biological replicates). Data are mean +/- SEM and statistics are a Student's T-test $*p < 0.05$, $**p < 0.01$. **(C)**

Representative western blot verifying loss of NIK in BMDMs from LysMCre mice. Samples were treated with 20ng/mL IL-4 and 10μM MG132 for 4 hours before extracting protein with RIPA buffer. Seahorse extracellular flux analysis showing **(D)** OCR readouts of IL-4 treated (M(IL-4)) NIK$^{WT}$ BMDMs pretreated for 5 hours with 5μM B022 or vehicle (DMSO). Line graphs are combined technical (n > 20) and biological replicates with bar graphs representing each animal (n > 5 biological replicates). Data are mean +/- SEM and statistics are a Student's T-test with pvalues shown on samples close to significant **$p < 0.01$. **(E)** Representative western blot showing inhibition of NIK post 5μM B022 treatment. Samples were pretreated with B022 for 5 hours before treatment with 20ng/mL IL-4 for 4 hours and extracting protein with RIPA buffer.

**Supplemental Figure 2: NIK$^{KO}$ macrophages have no significant transcriptional changes in metabolic genes.** RNA-sequencing heatmaps represent 2 biological replicates with a Log2Fold change post 6 hour IL-4 treatment over baseline in NIK$^{WT}$ **(A)** and NIK$^{KO}$ BMDMs **(B)** probing for changes in genes involved in glycolysis, TCA cycle, pentose phosphate pathway, and glucose transporters. RNA-sequencing heatmaps of NIK$^{KO}$ macrophages compared to NIK$^{WT}$ at baseline **(C)** or after 6 hour treatment of IL-4 **(D)** representing changes in gene expression of electron transport chain complex subunits. Statistics were calculated by DESeq2 and -log10 (p value) of 1.3 represents a p value of 0.05.

**Supplemental Figure 3: Mitochondrial homeostasis is disrupted due to a loss of NIK. (A)** 24 hour IL-4 stimulated BMDMs were costained with Mitotracker green and Mitotracker red to assess mitochondrial membrane potential via flow cytometry as a marker of mitochondrial functionality. Cells with low mitochondrial membrane potential were termed "dysfunctional" **(B)** and those with high mitochondrial membrane potential "functional" (**C**). Statistics are a

741 Student's T-test. \*p < 0.05 (n = 5 biological replicates). **(D)** Representative western blots from

742 human U937 cells showing NIK localization to a mitochondrial enriched fraction post 0-4 hours

743 of IL-4 treatment along with mitochondrial fractionation controls. **(E)** Isolated mitochondria

744 were analyzed by flow cytometry against standard sized beads to calculate mitochondrial size

745 with the quantification of mitochondrial size based on the standard curve of the beads **(F)** (n = 4

746 biological replicates). Statistics are a Student's T-test and pvalues are shown. **(G)** Representative

747 immunofluorescent images characterizing the mitochondrial network in NIK$^{WT}$ and NIK$^{KO}$

748 macrophages at baseline using antibodies against HSP60 (green), mitochondrial DNA (red),

749 TFAM (purple), and DAPI (blue).

750 **Supplemental Figure 4: Lineage negative bone marrow cells are enriched for low**

751 **proliferative cells due to a loss of NIK.** Gating scheme for stem cell analysis in NIK$^{WT}$ **(A)** and

752 NIK$^{KO}$ **(B)** bone marrow. Quantification of the stem cell compartment probing for differences in

753 the LSK populations (early lymphoid progenitor) **(C)**, MPP (multi potent progenitor) **(D)**, LT-

754 HSC (long term hemopoietic stem cell) **(E)**, and ST-HSC (short term hemopoietic stem cell) **(F)**.

755 Cell populations (n ≥ 4 biological replicates). Statistics are a Student's T-test. \*p < 0.05, \*\*p <

756 0.01.

757 **Table 1: Primary antibodies and antibody-based reagents used in the study.**

| Catalog Number | Company | Reagent | Conjugate |
|---|---|---|---|
| **Stemcell Panel** | | | |
| A15423 | Invitrogen | Anti-MO CD117/C-Kit | APC-Cy7 |
| 15-5981-81 | Invitrogen | Anti-MO Ly-6A/E (SCA1) | PE-Cyn5 |
| 25-0481-80 | Invitrogen | Anti-MO CD48 | PE-Cyn7 |
| 12-1351-81 | Invitrogen | Anti-MO CD135 | PE |
| 17-1502-80 | Invitrogen | Anti-MO CD150 | APC |
| 11-0341-82 | Invitrogen | Anti-MO CD34 | FITC |
| **Immune Panel** | | | |
| 25-0032-80 | Invitrogen | Anti-MO CD3 | PE-Cyn7 |
| 12-0452-81 | Invitrogen | Anti-MO CD45R(B220) | PE |
| 11-5931-81 | Invitrogen | Anti-MO Ly-6G | FITC |
| 35-0451-80 | Invitrogen | Anti-MO CD45 | PE-Cyn5.5 |
| 17-0112-81 | Invitrogen | Anti-MO CD11b | APC |
| 155505 | Biolegend | Anti-MO Siglec F | PE |
| **In vivo Polarization** | | | |
| 117351 | Biolegend | Anti-MO CD11c | APC |
| 144509 | Biolegend | Anti-MO CD193 (CCR3) | FITC |
| 123111 | Biolegend | Anti-MO F4/80 | PE-Cyn5 |
| **Blood and Spleen Immune Phenotyping** | | | |
| 25-0112-U100 | Tonbo Biosciences | Anti-HU/MO M1/70 (CD11b) | APC-Cy7 |
| 115539 | Biolegend | Anti-MO CD19 | BV605 |
| 100267 | Biolegend | Anti-MO CD3 | APC-Fire810 |
| 65-4801-U100 | Tonbo Biosciences | Anti-MO F4/80 | PerCP-C5.5 |
| 128049 | Biolegend | Anti-MO Ly6C | BV650 |
| 50-1276-U100 | Tonbo Biosciences | Anti-MO Ly-6G | PE |
| 107643 | Biolegend | Anti-MO MHCII | BV711 |
| 506324 | Biolegend | Anti-MO TNFα | PE-Cy7 |
| 13-0871-T100 | Tonbo Biosciences | Viability | Ghost Dye 710 |
| 70-0161-U100 | Tonbo Biosciences | Anti-MO CD16 / CD32 (Fc Shield) (2.4G2) | |
| **ELISA Kit** | | | |
| 41-9234-P002 | Tonbo Biosciences | Mouse IL-10 ELISA Matched Antibody Pair Kit | |

| Immunofluorescence | | | |
|---|---|---|---|
| MABT166 | Millipore | Anti-TFAM | |
| CBL186 | Millipore | Anti-DNA | |
| N-20 | Santa Cruz Biotechnology | Anti-HSP60 | |
| **Western Blotting** | | | |
| 4994 | Cell Signaling | Anti-NIK | |
| 2697 | Cell Signaling | Anti-pIKKα/IKKβ S176/180 | |
| OP133 | EMD Millipore | Anti-IKKα | |
| 4882 | Cell Signaling | Anti-p100/p52 | |
| SC-365062 | Santa Cruz Biotechnology | Anti-GAPDH | |
| SC-69879 | Santa Cruz Biotechnology | Anti-β-Actin | |
| SC-5274 | Santa Cruz Biotechnology | Anti-β-Tubulin | |
| 4661T | Cell Signaling | Anti-VDAC | |

760 **Table 2: Commercial reagents used in the study.**

| Catalog Number | Supplier | Reagent |
|---|---|---|
| L31963 | Invitrogen | Violet Live/Dead dye |
| 14-9161-71 | Invitrogen | HU FCR Binding Inhibitor |
| M36008 | Invitrogen | MitoSox |
| M22425 | Invitrogen | Mitotracker Red FM |
| M7514 | Invitrogen | Mitotracker Green FM |
| 8778s | Cell Signaling | Mitotracker Deep Red FM |
| TLRL-PB5LPS | Invivogen | Ultrapure LPS-B5 |
| 214-14 | PeproTech | Recombinant Murine IL-4 |
| LS004174 | Worthington Biochemical | Collagenase, Type 2 |
| LS004186 | Worthington Biochemical | Collagenase, Type 4 |
| 354234 | Corning | Matrigel Basement Membrane Matrix |
| F3879 | Sigma Aldrich | Fibrinogen |
| 341584 | Sigma Aldrich | Polyvinyl Alcohol |
| C35007 | Invitrogen | CyQUANT |
| D275 | Invitrogen | DiO |
| F3506 | Sigma Aldrich | FMLP |
| TNB-0607-KIT | Tonbo Biosciences | Foxp3/Transcription Factor Staining Buffer Kit |
| 75351 | Sigma Aldrich | Oliogmycin A |
| 15218 | Cayman Chemical | FCCP |
| A8674 | Sigma Aldrich | Antimycin A |
| ALX-350-360-G001 | Enzo Life Sciences | Rotenone |
| 1708891 | BioRad | iScript |
| A25742 | Universal SYBR Green | Invitrogen |
| 700930 | Cayman Chemical | MitoCheck Complex I Activity Assay Kit |
| 700940 | Cayman Chemical | MitoCheck Complex II Activity Assay Kit |
| 700950 | Cayman Chemical | MitoCheck Complex II/III Activity Assay Kit |
| 700990 | Cayman Chemical | MitoCheck Complex IV Activity Assay Kit |
| 701000 | Cayman Chemical | MitoCheck Complex V Activity Assay Kit |
| 5120-0075 | KPL | TMB Substrate |
| 15702-10 | Polysciences Inc | 1um carboxylate microspheres - FITC |
| HY-120501 | MedChem Express | B022 NIK inhibitor |
| P7170-1 | Sigma Aldrich | Ponceau S |
| 24592 | Thermo Scientific | GelCode Blue |

763    **Table 3: Oligonucleotides used in quantitative real-time PCR.**

| Name | Sequence |
|---|---|
| mCD163 F | CATCCGCCTTTGAATCCATCTCTTG |
| mCD163 R | CCTCCTCATTGTCTTCCTCCTGTG |
|  |  |
| mArg1 F | ACC TGG CCT TTG ATG TCC CTA |
| mArg1 R | AGA GAT GCT TCC AAC TGC CAG ACT |
|  |  |
| mRps 18 F | ACTTTTGGGGCCTTCGTGTC |
| mRps 18 R | GCCCAGAGACTCATTTCTTCTTG |

# Figure 1

# Figure 2

**Figure 3**

# Figure 4

**A.** CD19+ B cells

% of Blood Leukocytes

NIK^WT ****
NIK^KO

**B.** CD3+ T cells

% of Blood Leukocytes

**C.** CD11b+ cells

% of Blood Leukocytes

**

**D.** Ly6G+ Ly6C+ Neutrophils

% of CD11b+ Blood Leukocytes

*

**E.** F4-80+ Eosinophils

% of CD11b+ Blood Leukocytes

**

**F.** Ly6C^hi Monocytes

% of CD11b+ Blood Leukocytes

***

○ NIK^WT
● NIK^KO

**G.**
Eosinophils   Neutrophils
Myeloid cells   Ly6C^hi Monocytes
T cells   B cells

**H.** NIK^WT   NIK^KO   NIK^WT   NIK^KO

**I.** TNFα   CD11b   Ly6C   Ly6G

Expression Intensity
Low   High

SSC-A   F4/80   CD19   MHCII

**Blood leukocytes - LPS ex vivo stimulation**

**J.** NIK^WT   NIK^KO

Ly6C

CD11b+Ly6C^hi   CD11b+Ly6C^hi

TNFα Low   High   TNFα Low   High

CD11b

**K.**

Normalized to Mode

%TNFα^hi

12.8%

46.2%

TNFα

**L.** CD11b+ Ly6C^hi Monocytes

TNFα MFI

**

○ NIK^WT
● NIK^KO

# Figure 5