Optogenetic activation of dorsal raphe serotonin neurons induces brain-wide activation

Hiro Taiyo Hamada[1,3,*], Yoshifumi Abe[2], Norio Takata[2], Masakazu Taira[1], Kenji F. Tanaka[2], Kenji Doya[1,*]

[1] Neural Computation Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan
[2] Division of Brain Sciences, Institute for Advanced Medical Research, Keio University School of Medicine, Tokyo, Japan
[3] Research & Development Department, Araya Inc., Tokyo, Japan
* Correspondence to hamada_h@araya.org, doya@oist.jp

Abstract

Serotonin is a neuromodulator that affects multiple behavioral and cognitive functions. Nonetheless, how serotonin causes such a variety of effects via brain-wide projections and various receptors remains unclear. Here we measured brain-wide responses to optogenetic stimulation of serotonin neurons in the dorsal raphe nucleus (DRN) of the male mouse brain using functional MRI with an 11.7T scanner and a cryoprobe. Transient activation of DRN serotonin neurons caused brain-wide activation, including the medial prefrontal cortex, the striatum, and the ventral tegmental area. The same stimulation under anesthesia with isoflurane decreased brain-wide activation, including the hippocampal complex. These brain-wide response patterns can be explained by DRN serotonergic projection topography and serotonin receptor expression profiles, with enhanced weights on 5-HT1 receptors. Together, these results provide insight into the DR serotonergic system, which is consistent with recent discoveries of its functions in adaptive behaviors.

Introduction

Serotonin, 5-hydroxytryptamine (5-HT), serves crucial behavioral and cognitive functions, including regulation of the awake-sleep cycle, mood, reward, and punishment predictions [1, 2], waiting behaviors [3, 4], and learning [5]. Serotonin is also a major target of medications for psychiatric disorders, notably major depression, by selective serotonin re-uptake inhibitors (SSRIs). The dorsal raphe nucleus (DRN) is a major source of serotonergic projections to brain-wide targets, including the frontal cortex, the basal ganglia, the ventral tegmental area (VTA), and other mid-brain regions [6]. The DRN contains glutamatergic, GABAergic, and dopaminergic neurons, in addition to serotonergic neurons [7]. Optogenetic manipulation of activities of serotonin neurons has revealed their functions in reward [8, 9] , waiting for delayed rewards [3, 4], anxiety [10, 11], and learning [5, 12]. Nonetheless, associations between brain-wide serotonergic projections and brain responses remain unclear [13, 14].

In order to clarify the global picture of serotonergic control of brain activities, we performed optogenetic functional MRI experiments using Tph2-ChR2(C128S) transgenic mice and an 11.7T MRI scanner with a cryoprobe for high S/N measurements. A previous optogenetic fMRI experiment activated DRN serotonin neurons using the Pet1 promoter under anesthesia and showed widespread suppression of cortical activities [15]. We introduced an acclimation protocol to allow stable fMRI recording in awake mice [16-18] and found that DRN serotonin activation in an awake state induced increased brain-wide activity in both cortical and subcortical areas. In contrast, under anesthesia with isoflurane, DRN serotonin activation decreased activities in those areas, consistent with a previous study [15].

We further found that regional BOLD responses can be approximated by a linear model with projection densities of DRN serotonin neurons and expression profiles of different types of serotonin receptors, with negative coefficients for 5-HT1 receptors and positive coefficients for 5-HT2 receptors. These results suggest that serotonergic neurons activate brain-wide regions, and they demand reconsideration of implications of results under anesthesia.

Results

In order to clarify effects of brain-wide serotonergic projections, we employed optogenetic fMRI, combining optogenetic stimulation of serotonergic neurons in the dorsal raphe nucleus (DRN) and measurements of whole-brain activities using functional MRI. We used Tph2-ChR2(C128S) transgenic mice [11, 19] which selectively express step-type channel rhodopsin in serotonin neurons [4, 11]. We used an 11.7T MRI scanner (Bruker BioSpec 117/11) with a cryoprobe for high S/N recording of BOLD signals from the whole brain, except the olfactory bulb and the posterior cerebellum.

Prior to experiments, the transgenic and wild-type (WT) groups (Transgenic group: n = 8; Wild-type: n = 6) underwent surgery to attach a plastic fixation bar and implant an optical cannula in the DRN. Acclimation training in a mock chamber followed (Figure 1. See Methods for details). After fMRI experiments, animals were tested in a reward

85   waiting task [3, 4] to confirm the effectiveness of optogenetic stimulation
86   (Supplementary Figure S1).
87
DRN serotonin activation induces a brain-wide increase in BOLD signals
We measured BOLD responses under optogenetic stimulation of DRN serotonin
neurons in an ON/OFF block design protocol (Figure 1e). Photo-stimulation increased
BOLD signals in the DRN (Figure 1f), with a tendency for reduced responses after
repeated stimulation. On average, over five stimulations, blue light stimulation induced
a significantly higher BOLD response than did yellow light stimulation ($p_{FDR-corrected} <$
0.05, correction of multiple comparisons with a false discovery rate (FDR), see 'General
Linear Model (GLM) for BOLD signals' in Methods; Figure 1g and Supplementary
Figure S5).
97
A group analysis with a general linear model (GLM) of photo-activated BOLD
responses between blue and yellow stimulation from the transgenic group(n = 8)
revealed a brain-wide increase in BOLD responses, significantly in the frontal cortex,
the caudate putamen (CPu), temporal cortical regions, the thalamic complex, and the
VTA ($p_{FWE-corrected} < 0.05$, cluster-wise correction controlling family wise error (FWE)
with probabilistic threshold-free cluster enhancement (pTFCE); Figure 2a). Trial-wise
BOLD responses also showed consistent brain-wide activation by blue, but not yellow
stimulation ($p_{uncorrected} < 0.05$, Supplementary Figure S2a-b).
106
We next selected 28 regions of interest (ROIs) from cortical and subcortical areas that
are involved in valence processing, according to the Allen mouse brain atlas (Table 1,
see Regions of interest in Methods). We selected components of basal ganglia such as
the caudate putamen (CPu), nucleus accumbens (NAc), and globus pallidus (GP) [20].
Furthermore, other subcortical regions, including the bed nucleus of the stria terminalis
(BST) [21], the hippocampus [22], and the lateral habenula (LHb) [23] were also
selected. Then, we extracted regional average beta values (regression coefficients) from
the GLM. Comparison of regional beta values under blue and yellow stimulation from
the transgenic group revealed that BOLD responses were significantly higher in blue
illumination sessions in 12 of 14 cortical regions, including the OFC, mPFC and ACC,
and 12 of 14 subcortical regions, including the CPu, NAc, GP, VTA, LHb, and DRN
(Figure 2b) although no statistical differences were found in the WT group
(Supplementary Figure S4). Moreover, there were no significantly higher BOLD
responses in vRSC, V1, DG, or CA1.
121
The week after Session 1, we performed another session under the same stimulation
parameters as in Session 1. Patterns of BOLD responses were similar to those in
Session 1 (transgenic: n = 7, WT: n = 6, Supplementary Figure S3). Pearson's
correlation of beta values in Sessions 1 and 2 was r = 0.875, which was statistically
significant ($p_{uncorrected} < 1e^{-8}$; Supplementary Figure S7.a). We further checked regional
differences of responses in peak amplitude and peak timing by extracting regional
BOLD signals (Supplementary Figure S5-9). Peak amplitude and peak timing of BOLD
responses varied across the 28 ROIs (Supplementary Figure S8.c-d), with the largest
responses in the VTA, SNr, and FP. Both amplitude and timing of BOLD responses in
Sessions 1 and 2 were significantly correlated under blue light stimulation (amplitude: r

= 0.886, $p_{uncorrected} < 1e^{-8}$; timing: r = 0.679, $p_{uncorrected} < 0.005$), but not yellow
stimulation (amplitude: r = -0.06, $p_{uncorrected} = 0.766$; r = 0.037, $p_{uncorrected} = 0.855$;
Supplementary Figure S8.a, b). These results indicate that our photo-stimulation of
DRN induced consistent activation of DRN serotonergic neurons, which caused distinct
spatial and temporal responses in multiple cortical and sub-cortical regions.
137
DRN serotonin activation induces a brain-wide negative response in BOLD signals
under anesthesia
Contrary to our above results, a previous optogenetic fMRI study reported that DR
serotonin activation evoked negative cortical cerebral blood volume (CBV) responses,
an enhanced proxy of BOLD signals, from the entire cortical region under anesthesia
[15]. There are two critical differences between that study and that one, optogenetic
probes and anesthesia. The previous study used the ePET-Cre mouse line and AAV
harboring Cre-dependent ChR2 and applied 20-Hz photostimulation. In contrast, the
current study used Tph2-ChR2(C128S) transgenic mice, which express step-type opsin,
to allow continuous activation by a pulse of blue light (1.0 s, 473 nm) at the beginning
and yellow light (1.0 s, 593 nm) at the end (Figure 1e). Another difference is that the
previous study employed isoflurane, a general gas anesthetic which is also a muscle
relaxant, while this experiment employed no anesthetics.
151
In order to test the possibility that the difference was due to anesthesia, we compared
BOLD responses to DRN serotonin stimulation between anesthetized and awake states
in the same animals. We employed the same ON/OFF stimulation protocol under
anesthesia with 1% isoflurane with the same groups (Transgenic: n = 7, WT: n = 5,
Figure 3a). We also performed another ON/OFF stimulation session the next day in an
awake state in order to eliminate possible confounding by diminished responses after
repetitive photo-stimulation (Figure 3b).
159
Under anesthesia, we observed mostly negative BOLD responses from the entire
cortical region, while BOLD signals in the DRN remained positive (statistical t > 1,
uncorrected; Figure 3a & Supplementary S10a). On the other hand, Session 4 in an
awake state replicated positive BOLD signals from mid-line cortical regions and DRN
(statistical t > 1, uncorrected; Figure 3b & Supplementary Figure S10b). We then
contrasted regional beta values in anesthetized and awake states (Figure 3c). In the
anesthetized state, multiple regions, such as the ACC, dRSC, vRSC, DG and CA1
showed negative beta values, significantly different from those in an awake state (p_{FDR-}
$_{corrected} < 0.05$, correction of multiple comparisons with FDR). We further found a
statistically significant difference in peaks of % BOLDs in these regions between
anesthetized and awake states ($p_{FDR-corrected} < 0.05$, Figure 3d; Supplementary Figure
S11). Our results support the hypothesis that general anesthesia induces negative BOLD
responses by DR serotonin photo-activation.
173
Brain-wide BOLD responses are explained by serotonin projection density and
receptor expression profiles
Our results revealed a brain-wide profile of evoked BOLD signals by DR serotonin
activation (Figure 2a). One plausible mechanism underlying such a spatial profile is the
density of the serotonergic projections from the DRN. In order to test such a possibility,

we checked the correlation between intensity of projections of DRN serotonin neurons
and peak levels of BOLD responses. We extracted serotonin-specific structural
projection density from the DRN using the Allen Brain Atlas (Figure 4a and
Supplementary Figure S12).

Serotonin projection density was statistically correlated with beta values under blue
stimulation ($r = 0.462$, $p_{FDR\text{-}corrected} < 0.05$), but not under yellow stimulation ($r = 0.132$,
$p = 0.529$; Figure 4b). We also found that projection density was significantly correlated
with photo-stimulated BOLD responses in awake and anesthetized states (Anesthetized
(Session 3): $r = 0.546$, $p_{FDR\text{-}corrected} < 0.05$; Awake (Session 4): $r = 0.758$, $p_{FDR\text{-}corrected} <$
0.001; Figure 4c).

The previous study revealed that serotonergic structural density was not correlated with
the cortical blood volume (CBV) response of DRN serotonin stimulation under 0.5%
isoflurane + 0.2mg/kg/h s.c. medetomidine [15]. We checked whether the difference
originated from our ROI difference. After preprocessing with the current study, time
series of CBV signals were extracted from 28 ROIs (see "Supplementary Methods").
We first applied GLM to the previous dataset [15]. Although we consistently found
negative beta values across multiple brain regions, stimulation of DRN serotonin
neurons under anesthesia showed a negative BOLD response, contrary to the previous
result [15] ($p_{FDR\text{-}corrected} < 0.05$; Supplementary Figure S13). We then compared peak
intensity of CBV and BOLD signals considering the methodological difference of GLM
between these two studies. Pearson correlation between peak intensity of time series
from ΔCBV signals and %BOLD responses under anesthesia (Session 3) showed
statistical significance ($r = 0.53$, $p_{uncorrected} < 0.01$; Supplementary Figure S14) although
neither was significantly correlated with serotonergic structural projection density (peak
intensity of CBV signals (Grandjean2019): $r = 0.35$, $p_{uncorrected} = 0.08$; peak intensity of
BOLD signals (Session 3): $r = 0.34$, $p_{uncorrected} = 0.09$; Supplementary Figure S14).

We further investigated whether photo-stimulated BOLD responses are associated with
serotonin synaptic receptors. Gene expression maps for 5-HT1A, 5-HT1B, 5-HT1F, 5-
HT2A, and 5-HT2C receptors were obtained from the Allen Brain Atlas
(Supplementary Figure S12&S15). We then constructed a multiple linear regression
model (MLRM; Figure 5a) composed of one dependent variable of the BOLD response
and six independent variables of the projection density and five gene expression
patterns (5-HT1a, 5-HT1b, 5-HT1f, 5-HT2a, and 5-HT2c receptors) after normalization.
The model reproduced the BOLD response with $R^2=0.66$ ($p_{uncorrected} < 0.01$, Table 2).
We also checked the consistency of the model using BOLD responses in different
sessions. Using the fit beta values with Session 1, we found a significant correlation
with the beta value in Session 4 (Awake: $r = 0.615$, $p_{FDR\text{-}corrected} < 0.01$, Figure 5b), but
not in Session 3 (Anesthetized (Session 3): $r = 0.365$, $p_{FDR\text{-}corrected} = 0.079$; Figure 5c).

We investigated the contribution of each receptor by the estimated regression
coefficients in anesthetized and awake states (Figure 5d, Table 3, and Supplementary
Table 3). For each subject, we estimated regression coefficients from each session. In
the awake state (Sessions 1, 2, and 4), gene expression of 5-HT1 receptors (5-HT1a, 5-
HT1b, and 5-HT1f) showed average negative coefficients, indicating negative

contributions to BOLD responses. On the other hand, gene expression of 5-HT2
receptors showed average positive coefficients, indicating positive contributions to
BOLD responses. We also checked whether expression of each receptor type differed
between awake and anesthetized states using one-way analysis of variance (ANOVA).
These results revealed that in the anesthetized state (Session 3), weights for 5-HT1f and
5-HT2c receptors were significantly different ($p_{Bonferroni-corrected} < 0.05$, Table 3).

These results suggest that photo-activated BOLD responses reflect projection density of
serotonin neurons and serotonin receptor expression profiles and that general anesthesia
affects responses of 5-HT1 and 5-HT2 receptors differently to induce altered BOLD
responses.

Discussion

Our opto-fMRI investigation of DRN serotonergic modulations of brain-wide dynamics
yielded three major discoveries. First, DRN serotonergic activation causes brain-wide
positive BOLD responses in an awake state. Second, DRN serotonergic activation under
general anesthesia causes brain-wide negative BOLD responses. Third, the whole-brain
profile of DRN serotonin-activated BOLD signals is associated with the density of DRN
serotonergic projection and expression profiles of 5-HT receptors, with opposite
weights for 5-HT1 and 5-HT2 receptors.

The present results, which demonstrate brain-wide activation by DRN serotonergic
activation, provide a means of interpreting recordings and manipulation studies in
humans and animals [3-5, 14, 24-28]. Previous studies have demonstrated serotonin's
involvement in valence processing [2, 26] through modulation of the ACC and
ventromedial PFC (vmPFC) on learning signals [27], the CPu on short/long-term
prediction of rewards [24], the mPFC and OFC on waiting for delayed rewards [14],
and the BST on anxiety-like behaviors [10, 21]. A reduction of reward sensitivity is a
symptom of major depression. In healthy subjects, 2-week selective SSRI
administration potentiated reward signals in the ACC and vmPFC [27]. VTA-projecting
DR serotonin neurons induce self-stimulating behaviors, an indication of positive
reinforcement [29]. Brain response from brain-wide cortical regions and the VTA may
reflect neural mechanisms underlying recovery of reward sensitivity. High-delay
discounting of future rewards is another indication of psychiatric disorders, including
major depression [30]. Various studies have shown that optogenetic activations of DR
serotonergic projections to the orbitofrontal cortex and the mPFC, but not the NAcc,
enhance waiting for delayed rewards [3, 14]. Our brain response results demonstrate
that activations of the OFC and mPFC are induced by optogenetic stimulation of DR
serotonin neurons. Anxiety is a general symptom in psychiatric disorders. By increasing
extra-synaptic serotonin levels, SSRI contributes to a reduction of anxiety. However,
simple activation of DR serotonin neurons did not induce antidepressant effects [11].
Antidepressant roles of DR serotonin neurons on anxiety can be region-dependent [21,
31] and work in a time-locked manner [32]. The frontal-cortex- and BST-projecting DR
serotonin subsystem promotes antidepressant effects, whereas the amygdala-projecting
DR serotonin subsystem enhances anxiogenic effects [21, 31]. Nishitani et al. (2018)
showed that optogenetic serotonin neurons increase active coping with inescapable
stress [32]. Our opto-fMRI results support those studies, as those areas were

consistently activated by DRN serotonin stimulation, although such resting-state brain-
wide activation also may blunt effects due to co-activations of specific subsystems,
without time-locked stimulation.
276
In addition, average amplitudes of BOLD responses and peak timing are region-
dependent. This implies that serotonin-activated BOLD responses are coupled with
different regional distributions of receptor densities and serotonin projections, but not
with the distance of the activation site. In spite of indirect evidence, our linear
approximation of serotonin-activated BOLD responses supports such a possibility. A
subpopulation of DR serotonin projections to the cortex is also expressed with the
vesicular glutamate transporter Vglut3 (Slc17a8). Co-release of glutamate has recently
been reported [6, 31]. Our results may reflect the influence of serotonin as well as
glutamate co-release on downstream circuitry.
286
Serotonergic modulation in short-term windows using optogenetics can avoid
limitations of long-term windows, such as pharmacological manipulation [33, 34] and
chemogenetic manipulation with DREADD (designed receptors exclusively activated
by designed drugs) [35]. Multiple pharmacological studies have applied SSRIs to
manipulate the serotonin system. Therapeutic benefits of SSRI, including alleviation of
depressed mood and psychic anxiety, arise from chronic SSRI administration [36].
Chronic SSRI administration normalizes aberrant activity in the amygdala and ACC
during major depressive episodes [37], as well as in brain networks associated with
obsessive-compulsive disorder [38]. In healthy individuals, Scholl et al. (2017) found
that chronic SSRI administration amplifies both reward and effort signals, fostering
learning processes in the ACC, amygdala, mPFC, and striatum [27]. The brain-wide
positive response from the ACC, mPFC, and striatum with optogenetic activations in an
awake state may partly imitate chronic SSRI administration. On the other hand, acute
systemic SSRI administration causes brain-wide inhibitory modulation [33, 34].
However, the influence of acute SSRI administration on brain activity can be
confounded by the administration period, due to receptor sensitivity [34, 39, 40]. A
previous study stimulated the serotonin system with specifically designed drugs under
fMRI [35] and revealed brain-wide positive responses in the somatosensory cortex and
subcortical regions, including the hippocampus, the hypothalamus, and the cerebellum.
That study did not reveal BOLD responses of mid-line cortical regions, such as the
OFC, mPFC and ACC. A possible explanation for the difference between their results
and ours is that BOLD responses cause different time scales, depending on receptor
types. Our optogenetic fMRI approach revealed an influence on motivation-related
regions at different timings in a transient time window (Supplementary Figure S5, S6
and S8). Further studies using different methodologies, including optogenetics and
chemogenetics, will uncover neural mechanisms underlying different time scales of
serotonin modulation.
314
315 The mechanism underlying different serotonin modulation between awake and
316 anesthetized states remains unknown. One possible mechanism is a reduction of
317 glutamate transmission via serotonin in target regions [41]. Previous studies have shown
318 that isoflurane suppresses neural activity via inhibition of glutamatergic transmission
319 [41, 42]. We found that isoflurane reduced photo-stimulated BOLD signals where they

were potentiated in an awake state. Our results also showed that some regions show potentiated negative BOLD responses. In addition to reduced glutamatergic transmission, isoflurane also enhances sensitivity of GABA neurons [43, 44]. Such elevated responses of GABA neurons and reduced responses of glutamatergic neurons may result in diminished positive BOLD signals and potentiated negative BOLD signals. It is interesting to ask whether different anesthetics, like NMDA antagonists, induce different effects on receptor responses while neuron-specific distributions of serotonin receptors are unclear. Serotonin receptors in the prefrontal cortex are differentially distributed [45]. 5-HT1A and 2A receptors are enriched on glutamatergic pyramidal neurons in the PFC [46], whereas 5-HT3A receptors are located on GABAergic interneurons [47]. Since different anesthetics influence different neuron types in various ways, reactivity of serotonin receptors may induce different responses. In addition, anesthetics may change effects of serotonin receptors. Some studies showed that isoflurane changes binding affinities of 5-HT$_{2B}$ receptors [48] *in vitro* and 5-HT$_{1A}$ receptors [49] *in vivo*. Anesthetics can influence ligand binding to specific subtypes of serotonin receptors.

It is important to consider whether the negative brain-wide response is caused by the neuronal response. Granjean et al. (2019) revealed that DRN serotonin activation reduces the amplitude of local field potentials (LFPs) and burst frequency with multiunit activity (MUA) from a variety of cortical regions including M1, M2, S1, and FA under 0.5% isoflurane + 0.2mg/kg/h s.c. medetomidine [15]. That study also examined the association between the brain response to DRN serotonin activation and delta power/burst frequency. This study revealed that a negative brain-wide response is caused by a reduction of cortical neuronal activity under isoflurane & medetomidine. Another study also showed that the negative BOLD response by activation of medial spiny neurons (MSNs) is associated with decreased neuronal activity under isoflurane 0.3-0.7% [50]. There is also the possibility that the negative brain-wide response is due to an increase in cerebral blood flow (CBF), which subsequently reduces neuronal activity under isoflurane. It is well established that isoflurane affects blood vessels and amplifies CBF. Abe et al. (2021) showed that an increase in CBF in the striatum doesn't alter neuronal activity [51]. Furthermore, increasing isoflurane dosages elevates BOLD signal levels [52]. Thus, these studies suggest that the decrease in BOLD signals isn't due to increased CBF. Overall, a brain-wide negative response can be induced by decreased neuronal activity rather than increased CBF.

Furthermore, our results showed a negative BOLD response in dorsal hippocampus under anesthesia, despite a lack of projections to the hippocampus from the DRN. One potential cause is an indirect influence of downstream circuitry of DR serotonin neurons. For example, DR serotonin neurons have dense projections to the PFC, which interact with the hippocampus [53, 54]. Reduction of the BOLD response in the hippocampus under anesthesia can be indirectly caused by activation of DR serotonergic neurons through their downstream circuitry, although this influence is not observable in awake states. The brain-wide response by activation of DR serotonin neurons under anesthesia may have implications for the sleep state. The similarity of brain states between general anesthesia and non-rapid-eye-movement (NREM) sleep is implied [55]. Previous studies also revealed that DR serotonin neurons mediate the

sleep/wake cycle [56, 57]. Optogenetic activation of DR serotonin neurons prolonged
wakefulness and decreased NREM sleep, whereas their inactivation prevented arousal
from CO_2-induced sleep [56]. Similarly, optogenetic and chemogenetic activation
induced arousal from isoflurane [57]. The negative response under anesthesia may
reflect a brain response associated with arousal. Future studies are needed to clarify the
neural mechanism underlying BOLD signals under anesthesia.

We also demonstrated that expression of different 5-HT genes are associated with
photo-activated BOLD responses. Discrete contributions of serotonin receptor types
influence neural activities, mostly at metabotropic receptors [58]. 5-HT1 receptors are
inhibitory metabotropic receptors that suppress neural activity by modulating the
cellular signal cascade, whereas 5-HT2 receptors are excitatory metabotropic receptors
that enhance neural activity [59, 60]. Our MLRM analysis revealed that regression
coefficients of gene expression of 5-HT1 and 5-HT2 receptors are negative and positive,
respectively. The correspondence between functions of receptor types and signs of
coefficients implies that BOLD responses by DRN serotonin photo-activation can be
influenced by different receptor distributions. Localization of 5-HT receptors is
subtype-dependent. 5-HT_{1A} and 5-HT_{1B} receptors are pre-synaptically expressed, either
on somato-dendrites or within axon terminals of DR serotonin neurons [61, 62],
whereas 5-HT_{2A} and 5-HT_{2C} receptors are predominantly expressed on postsynaptic
neurons [63, 64]. However, whether 5-HT_{1F} receptors are pre- or post-synaptically
expressed is unclear, although they are enriched in cortical layers (layer IV and V) and
the caudate putamen [65]. Nonetheless, specifically how 5-HT receptors react to
serotonin transmission under general anesthetics is unclear. It is also plausible that
general anesthetics change binding affinities of 5-HT receptors [48, 49]. As another
potential mechanism, serotonin transmission can be influenced by general anesthetics.
Some studies revealed that isoflurane lessens spontaneous serotonin levels and
spontaneous firing [66, 67]. These changes of serotonin transmission and receptor
binding affinity may influence responses of 5-HT receptors under anesthesia.

The substantial reduction of magnitude of brain-wide responses between the first,
second, and fourth sessions is noteworthy, despite consistent activation patterns.
Repeated optogenetic stimulation of DR serotonin neurons and repeated stress
potentially account for this reduction. Persistent changes in spontaneous behaviors by
repeated optogenetic activation of DR serotonin neurons have been reported [68].
Serotonin induces synaptic plasticity, and repeated stimulation may cause plasticity
changes. In addition, repeated restraint stress also may change responses to optogenetic
stimulation of DR serotonin neurons, although the habituation protocol to awake fMRI
is introduced to prevent restraint stress. Restraint stress increases serotonin release in
the amygdala [69]. Molecular and neuronal mechanisms underlying such diminutions
are needed to understand long-term effects of serotonin stimulation.

It is also worth mentioning limitations of the relationship between the BOLD response
and behavioral variability in the present study. Although the present study showed
variability of the evoked BOLD response by serotonin stimulation, its behavioral
relationship remains unclear. The serotonin system has complementary distinct
subsystems [10, 14, 21, 31]. Ren et al. showed that amygdala-projecting DRN serotonin

neurons enhance anxiety-like behaviors, whereas frontal-cortex-projecting DRN
serotonin neurons enhance anxiolytic behaviors [31]. Miyazaki et al. demonstrated that
OFC-projecting serotonin neurons, but not mPFC-projecting serotonin neurons
contribute to waiting for delayed rewards [14]. BST-projecting serotonin neurons
promote anxiety-related behaviors [10, 21, 31]. Neural mechanisms underlying diverse
ways in which serotonin modulates behavioral functions remain to be clarified.
420
While our results replicated negative cortical responses under anesthesia and found the
association with the CBV response identified in a previous study [15], there are several
important differences in methodology and results between previous studies and this one.
First, one previous study utilized cerebral blood volume (CBV) signals with a contrast
agent to enhance influence of serotonin modulation, although the current study
employed BOLD signals. Although CBV signals are an enhanced proxy of BOLD
signals, the methodological difference in imaging may show different responses from
various brain regions. Second, we employed a transgenic group targeting the tryptophan
hydroxylase 2(Tph2) promoter [11, 19] although the previous study employed the
ePET-Cre line, targeting transcription factor Pet1 [15]. While expression of ChR2 by
Tph2 is selective for serotonergic neurons [29], Pet1 promoters are also expressed in
non-serotonergic neurons [9, 70]. While CA1 and DG were significantly deactivated by
serotonin photo-activation in our study, responses in temporal regions were not found in
the previous study [15]. The difference in GLM tools between the two studies should be
noted. We used SPM for fitting the BOLD response, whereas Grandjean et al. used a
different analytic tool, BROCCOLI [71] (https://github.com/wanderine/BROCCOLI),
for fitting the CVB response in SPM12. Beta values of the previous dataset using SPM
showed negative beta values in the DRN, which did not match ΔCVB signals, although
consistently negative brain responses across multiple brain regions were consonant with
ΔCVB signals (Supplementary Figure S13 & S14) [20]. GLM using SPM12 overlooks
the negative CBV response. The methodological difference in GLM tools may cause
different patterns of functional maps.
443
Additionally, our results reveal an association between 5HT projection density and
BOLD responses from GLM analysis using SPM12, but the previous study did not find
this association. One critical methodological difference is that we employed different
ROIs from those of the previous study. In the current study, 28 ROIs from the Allen
brain atlas included other sources of serotonin and dopamine, the MRN and VTA. In
contrast, the previous study used 38 ROIs from the Australian Mouse Brain Mapping
Consortium (AMBMC, https://www.imaging.org.au/AMBMC) with multiple ROI
hippocampal formations and midbrain regions. Our analysis revealed an association
between BOLD and CBV responses with our selected 28 ROIs. The ROI difference
may yield different results in the association between structural connectivity and
responses to optogenetic serotonin activation. Moreover, our results also indicated that
neither peak intensity of BOLD signals nor CBV signals are associated with 5-HT
projection density (Supplementary Figure S14c). Stimulation of the serotonin system
and its modulation through metabolic 5-HT receptors promote slow and sustained
activities [72, 73]. An advantage of GLM analysis is to capture not only acutely
activated responses, but also sustained responses by comparing target blocks with other
blocks. The correlation of 5-HT projection with peak intensity of brain response may

overlook the influence of sustained brain activities. Further studies are required to
reveal the association between serotonin projection density in brain regions and the
underlying variety of neural responses.
464
Beside the aforementioned methodological difference, different anesthetics (isoflurane
1% vs. 0.5% isoflurane + 0.2mg/kg/h s.c. medetomidine), canula locations (horizontal
vs. vertical insertion), stimulation power (473nm laser with 4 mW vs. 4-40 mW), and
scanner differences (an 11.7T MRI scanner with a cryoprobe vs. a 7□T Pharmascan
scanner) may also cause distinct brain responses. Further research is required to dissect
differences between the two sets of experimental results.
471
Testing acute and chronic SSRI treatments on serotonin modulation is important for
clinical modulation of the serotonin system. For example, one study observed an effect
of fluoxetine, an SSRI with acutely stressed mice under anesthesia [15]. That study
showed enhanced negative responses from brain regions such as the CPu and mPFC. It
is unclear whether such effects remain under awake, long-term SSRI administration. It
is also important to examine whether reactions of serotonin levels and 5-HT receptors
show different dynamics under acute and chronic SSRI administration. A meta-analysis
showed that serotonin levels in each region change during the course of SSRI
administration [40]. Salvan et al. demonstrated that acute SSRI administration may
change regulation of receptor responses [74]. Further studies on physiological and
molecular mechanisms under SSRI administrations are required.
483
In conclusion, we reported brain-wide responses of DRN serotonin activation via opto-
fMRI for the first time in awake animals. We found serotonergic modulation of
widespread cortical and sub-cortical targets at different times as a key controller of
brain-wide information transmission. We also showed different modulation by the
serotonin system in awake and anesthetized states. We further showed that serotonergic
structural projections and gene expression profiles of 5-HT receptors are associated with
BOLD responses. Further research is required to understand complex serotonin
modulation of brain-wide dynamics, given its neural and molecular mechanisms, such
as interactions with neural populations and receptors. Our findings reveal a functional
basis for understanding serotonergic control of brain-wide dynamics and functions.
494

Methods

496 All experimental procedures were performed under protocols approved by the Okinawa
497 Institute of Science and Technology Experimental Animal Committee.
498

Animals

Serotonin neuron-specific channelrhodopsin 2 variant (C128S) mice were generated by
crossing Tph2-tTA mice with tetO-ChR(C128S)-EYFP knock-in mice under control of
the tryptophan hydroxylase 2 (Tph2) promoter [11, 19]. Eight bi-genic and six wild-
type (WT) C57BL/6J male mice (age > 12 w.o.) were used in this study. The male mice
were housed at 24 ◦C on an inverted dark:light cycle (lights on 10:00-22:00, GMT+9).
After stereotaxic surgery, male mice underwent habituation training [16], MRI
experiments, and a reward delay task in turn.
507

Stereotaxic surgery

Male mice were initially sedated with 1-3% isoflurane, and were given three types of mixed anesthetic agents (1 (mg/mL) of medetomidine, 5.0 (mg/mL) of midaloram, and 5.0 (mg/mL) of butorphanol, i.p.) [75]. Cranial skins were removed with a sterilized surgical knife, and a plastic bar (3x3x27mm) for head-fixation in the MRI environment and an optic fiber (480 µm diameter, 4 mm length, Doric Lenses) were fixed to their skulls with dental acrylic cement. The optical fiber was inserted horizontally toward the DRN from the cerebellum in each male mouse. Male Mice were housed separately after surgery and underwent at least a 1-week recovery period. Mice were habituated to the MRI environment for at least another week. They subsequently underwent imaging sessions.

Acclimation

In order to habituate awake mice to the MRI imaging protocol, an acclimation protocol was employed after the surgeries for head-fixation [16-18] . Male mice were fixed in a mock MRI environment and exposed to MRI imaging sounds and blue illumination for 2 h to habituate them to a leak of illumination during scanning (2.0 Hz). Acclimation to sounds and illumination was repeated 7 days prior to the first imaging session. Following imaging sessions, at least 3 days of acclimation were permitted.

Functional magnetic resonance imaging (fMRI)

Each animal was placed in the MRI animal bed. T2-weighted structural imaging and three T2* functional imaging sessions were performed. MRI images were acquired with an 11.7-tesla MRI scanner for small animals (Biospec 117/11 system, Bruker Biospin, EmbH, Ettlingen, Germany) with a cryogenic quadrature RF surface probe (Cryoprobe, Bruker BioSpin AG, Fällanden, Switzerland). We performed T2-weighted structural imaging with a RARE sequence with the following parameters: 140 x 140 matrix, 13.5 x 13.5 mm^2 field-of-view, repetition time (TR)/effective echo time (TE) 4000/18.75 (ms), 31 coronal slices, slice thickness: 300 µm.

After T2-weighted structural imaging was performed, a functional MRI session was then executed. A functional MRI session was composed of two imaging runs, a blue-yellow-light stimulation run and a yellow-light stimulation run. All functional imaging was performed using a GE-EPI sequence with the following parameters: 67 x 67 matrix, 13.5 x 13.5 mm^2 field-of-view, TR/TE 1000/10.7 (ms), flip angle: 50◦, bandwidth: 333k(Hz), 31 coronal slices and slice thickness: 300 µm, and 690 repetitions.

ON/OFF stimulation paradigm

A stimulation protocol was composed of ON/OFF stimulation patterns (Figure 1d). In each activation session, stimulation was initiated 90 volumes after the onset of an imaging session. DRN serotonin neurons were stimulated with blue illumination (490 nm, 3.0 mW, 1.0 sec), and stimulation subsequently ceased with yellow illumination (630 nm, 3.0 mW, 1.0 sec) 20 seconds after the onset of blue illumination. This stimulation was repeated 5 times for 120 sec after the onset of blue illumination. For the control session, instead of blue illumination, only yellow illumination was applied in stimulation sessions.

Preprocessing and de-noising

The first 10 volume images of each session were removed from subsequent analysis due to initial imaging instability caused by motion and non-steady-state image quality. 10-fold magnification of images was done to process images with Statistical Parametric Mapping 12 (SPM12), which is designed to process human brain images. Therefore, the voxel size was 2 x 2 x 3 mm in analytical steps. Pre-processing, including motion correction, realignment, co-registration, normalization to a C57BL6/J template [76], and spatial smoothing (kernel with 4 x 4 x 6 mm) was executed with SPM12. Since each slice of a 3D brain image was taken at a different time, slice timing correction was applied to correct slice-timing differences by temporal interpolation. Next, realignment of 3D brain images was conducted to correct motion-related changes of brain position. Then, co-registration was executed to overlay functional T2* images onto structural T2 images and to save coordination changes of T2* images for normalization. During normalization, T2 structural images were first warped to fit the average C57BL6/J template [76]. Then T2* functional images were further warped to the template using the co-registered coordination change of the images. In spatial smoothing, voxel signals were spatially smoothed using a gaussian kernel (4 x 4 x 6 mm). After pre-processing steps, the general linear model in SPM12 was executed to map spatial activation patterns.

Regions of Interest

The Allen brain institute provides parcellation of the C57BL6 brain. In order to extract BOLD signals from anatomically defined brain regions, this parcellation was utilized for subsequent analyses. Parcellation was first warped to the MRI template using Advanced Normalization Tools (ANTs), and 28 regions of interest (ROIs) were selected from the warped parcellation. 28 ROIs were selected with the following three criteria. First, primary cortical regions, especially mid-line regions, were selected. Second, primary reward- and punishment-related sub-cortical regions, including basal ganglia, VTA, and DRN should were included. Third, ventro-lateral cortical and sub-cortical regions, such as the amygdala complex were not selected due to its sensitivity to artifacts near air cavities by the ear canals [77].

General Linear Model (GLM) for BOLD signals

A general linear model (GLM) was performed to map beta values of parameter estimation of optogenetic stimulation onto the mouse brain template with SPM12. For statistical functional mapping of optogenetic stimulation, each voxel under an uncorrected P value < 0.001 was thresholded, and cluster-wise multiple comparisons were subsequently executed with family-wise error (FWE) by thresholding pFWE < 0.05 with probabilistic threshold-free cluster enhancement (pTFCE) [78]. Statistical second-level, voxel-wise analysis was finally executed to compare functional maps of blue and yellow illumination from transgenic mice. For Session 1, all procedures were executed accordingly. For Sessions 2, 3, and 4, none of the voxels under uncorrected P values < 0.001 survived the thresholding.

Beta values from BOLD responses

600   Furthermore, regional beta values were extracted based on ROIs. Statistical significance
601   of regional beta values was subsequently tested with Student's t tests with multiple
602   comparisons with the false discovery rate (FDR; p value (FDR) < 0.05).
603
604   **Extraction of regional BOLD signal changes**
605   Each regional BOLD signal change (%) is calculated by averaging regional time series
606   on voxels, which are scaled with the mean of time series during the no-stimulation
607   period and followed with multiplication by 100.
608
Structural density and RNA expression maps
The structural density map of serotonin connections was obtained from the mouse brain
database of the Allen Brain Institute (Experimental IDs: 480074702) [79]. As with the
structural density map, gene expression of serotonin receptors was obtained from the
mouse brain database of Allen Brain Institute (Experimental IDs: 79556616 (5HT1a),
583 (5HT1b), 69859867 (5HT1f), 81671344 (5HT2a), and 73636098 (5HT2c)).
Intensity of the structural density and gene expression was normalized ranging from 0 to
100. Normalized energy intensity was extracted from the normalized atlas. Multiple
linear model analysis of ROIs between betas from GLM analyses and gene expressions
was performed.
619
Multivariable Linear Regression Model (MLRM)
Beta values from each session were obtained based on 28 ROIs. Structural density and
RNA expression of 5HT1a, 1b, 1f, 2a, and 2c were also extracted based on 28 ROIs.
Multivariable linear models were used to estimate weights of structural and gene
expression of 5HT receptors; where beta values from each session were the dependent
variables, and intensity of structural and gene expression was the explanatory variable.
Results of statistical values for fitting are described in Table 2.
627
One-way Analysis of Variance (ANOVA) on Receptor Contribution
Regression coefficients of structural density and RNA expression of 5-HT receptors
were calculated for each subject for all sessions. One-way ANOVA was applied to
regression coefficients of each receptor type in order to test whether receptor expression
is statistically different between awake (Sessions 1, 2, &4) and anesthetized states
(Session 3). For analysis, data from all sessions were concatenated (total data: n=29,
session 1 (awake): n=8; Session 2(awake): n=7, Session 3 (anesthetized): n=7, Session 4
(awake): n=7). p values from one-way ANOVA were further corrected with the
Bonferroni method. Results of ANOVA are described in Supplementary Table ST3.
637
A reward delay task
After imaging sessions, some mice from both groups (Transgenic: n=3, WT: n=2)
subsequently performed a waiting experiment for a delayed reward. A sequential tone-
food waiting task was utilized as a free operant task [3, 4] (Supplementary Figure S1a).
After MRI sessions, individual subjects were trained and tested in an operant-
conditioning box (Med-Associates, 12.7x12.7 cm). In short, subjects were required to
initiate trials by nose-poking into a tone site after 500 (ms). The reward-delay task was
composed of reward trials (75%) and omission trials (25%). We adopted two types of
stimulation protocols (Supplementary Figure S1b): fixed-delay and random-delay

protocols. At the onset of reward and omission trials, either blue or yellow illumination
was delivered. In reward trials, blue/yellow light stimulation was delivered at the onset
of each trial until nose-poking was performed, and yellow light was delivered when
subjects waited for 6.0 (s) for the fixed protocol and 2, 6, or 10 (s) for the random delay
protocol. However, yellow illumination was also given when subjects failed to wait for
the delay. On the other hand, during omission trials, yellow stimulation was given when
mice gave up waiting for delayed rewards whereas light stimulation was delivered in the
same manner as in the reward trials.
655

Data availability

657 Preprocessed data, including functional and structural images, are available from
658 https://openneuro.org (Project_ID: optofMRI_DRN).
659

Code availability

661 In-house codes are also available from https://github.com/Taiyou/OptofMRI_5HT
662 (DOI:10.5281/zenodo.10956557). All software used in this study is also available
663 (Supplementary Table ST1).
664

Acknowledgments

We thank Dr. Koji Toda for his comments on awake optogenetic stimulation, Dr.
Katsuhiko Miyazaki for his comments on protocol of optogenetic stimulation, Dr.
Keigo Hikishima for his comments on fMRI, and Mr. Yoshiaki Ohba for creation of
acclimation tools. We thank Dr. Jiafu Zeng for animal support for MRI acquisition. We
also acknowledge Prof. Grandjean for his support in replicating results from a CBV
fMRI dataset. This work was supported by Moonshot R&D from the Japan Science and
Technology Agency (JST), Japan, grant PMJMS2295 (to H.T.H.). WE also
acknowledge a grant-in-aid from the Japan Society for the Promotion of Science (JSPS),
Japan, grant JP19KK0387 (to N.T.) and JP16H06563 (to K.D.), Transformative
Research Areas (A) "Glia decoding" grant 20H05896 (to K.F.T.) and "Unified Theory"
grant 23H04975 (to K.D.); Brain Mapping by Integrated Neurotechnologies for Disease
Studies (Brain/MINDS) by the Japan Agency for Medical Research and Development
(AMED), Japan, grant JP22dm0207069 (to K.F.T.), JP18dm0207030 and
22dm0207001 (to K.D.); a grant from; the Naito Foundation, Japan (to H.T.H.); and
research support of Okinawa Institute of Science and Technology Graduate University
for the Neural Computation Unit.

Author contributions

684 H.T.H., K.F.T. and K.D. conceived the experiments. H.T.H., Y.A., and N.T. designed
685 and prepared experiments. H.T.H. performed MRI experiments. M.T. conducted
686 behavioral experiments. H.T.H. analyzed results, and H.T.H. wrote the manuscript.
687 H.T.H., Y.A., N.T., M.T., K.F.T., and K.D. reviewed the manuscript.
688

Competing interests

690 The authors declare no competing interests.

691
Figure 1. Experimental setup for opto-fMRI. (a) Surgical setup. A fixation bar was mounted on a
mouse's head with dental cement and an optical cannula was inserted into the dorsal raphe nucleus
(DRN) (b) Acclimation protocol. A mouse was exposed daily to recorded MRI scanning sounds for 2 h
for 7 days prior to fMRI sessions. (c) Location of an optical cannula. An optical cannula was inserted
horizontally into the DRN from the cerebellum. A cyan window indicates an imaging window for T2*
sessions. (d) Experimental schedule. After surgery and 7 days acclimation, the first fMRI recording
session was performed. A replicate session was also performed a few weeks later. Finally, experiments
for anesthesia effects were performed. On the 1st day, an fMRI recording with general anesthesia using
isoflurane 1% was performed, and an fMRI recording without anesthesia was conducted thereafter (e)
Stimulation protocol. In an fMRI session, optogenetic stimulation was initiated 90 s after an imaging
session began. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow
illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after blue illumination. (f)

Average time course of % BOLD signals in DRN. Time series of % BOLD signals were individually
extracted from the DRN in Sessions 1 and 2, and time series were averaged across transgenic mice (n =
8). Each cyan window indicates the stimulation period during a session for 20 s. (g) Alignment of
average % BOLD responses in the DRN. % BOLD responses were aligned at the time of blue
illumination (1.0 s) in session 1. Blue and yellow lines indicate % BOLD responses in the DRN from
blue and yellow stimulation sessions from the transgenic group (n = 8), respectively. Blue and yellow
highlights indicate the range of mean values +/- standard deviation (SD). A cyan window indicates
stimulation duration during a session for 20 s. ** indicates $p_{FDR-corrected}$<0.01 with two-tailed t tests, FDR
corrected.

713
Figure 2. BOLD response to optogenetic stimulation of serotonin neurons in the DRN. (a) Functional
map of optogenetic illumination. A functional map was calculated by contrasting BOLD responses
between blue and yellow illumination in the transgenic (n = 8) group ($p_{FWE\text{-}corrected}$<0.05, two-tailed paired
t tests and cluster-corrected controlling family-wise error (FWE) with probabilistic threshold-free cluster
enhancement (pTFCE, see 'General Linear Model (GLM) for BOLD signals' in the Method section)).
Magenta color indicates the t value. (b) Comparison of beta values in multiple ROIs. Beta values were
obtained from GLM analysis in Session 1, and were extracted and averaged based on ROIs from the
Allen Brain Atlas (Table 1). Differences of beta values between blue and yellow illumination sessions
were contrasted in the transgenic group. * indicates $p_{FDR\text{-}corrected}$<0.05 (Two-tailed paired t tests were
performed, and multiple comparison was corrected with false discovery rate (FDR)). Cyan and magenta
colors indicate blue and yellow illumination, respectively. (c) Alignment of average % BOLD responses
in the mPFc, OFC, CPu, and VTA. % BOLD responses were aligned at the time of blue illumination (1.0
s) in Session 1. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow
illumination (1.0 second; 593 nm) followed 20 s the offset of the stimulation after blue illumination. A

cyan window indicates stimulation duration during a session for 20 s. Blue and yellow lines indicate %
BOLD responses in the mPFc, OFC, CPu, and VTA from blue and yellow illumination from transgenic
groups (n = 8), respectively. Blue and yellow highlights indicate the range of mean values +/- SD. A
cyan window indicates stimulation duration during a session for 20 s. ** indicates $p_{FDR-corrected}<0.01$ with
two-tailed t tests, FDR corrected.
733

735 Figure 3. Comparison of BOLD responses between anesthetized and awake states. (a) Optogenetic fMRI
736 session with general anesthesia in Session 3. General anesthesia, 1% isoflurane, was applied to the
737 transgenic group (n=7) during optogenetic fMRI sessions. GLM analysis following optogenetic

stimulation was executed and visualized (T > 1, uncorrected). Magenta and cyan colors indicate positive and negative t values. (b) An optogenetic fMRI session in the awake state in Session 4. As in Session 1, no anesthesia was applied to the transgenic group during optogenetic fMRI sessions. As on day 1, GLM analysis was executed and visualized (T > 1, uncorrected). (c) Comparison between beta values of Sessions 3 and 4 in multiple ROIs. Beta values of blue illumination sessions in Session 3 and Session 4 were statistically contrasted using two-tailed paired Student's t test. * indicates $p_{FDR-corrected}<0.05$ with two-tailed t tests, FDR-corrected. (d) Alignment of average % BOLD responses in vRSC, CA1, DG, and DRN in Session 3 from transgenic mice. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after the blue illumination. Blue and magenta highlights indicate the range of mean values +/- SD. A cyan window indicates stimulation duration for 20 seconds. * indicates $p_{FDR-corrected}<0.05$ with two-tailed t tests, FDR corrected.

751
Figure 4. Correlation between the BOLD response and serotonergic projections. (a) Structural density
map of serotonergic projections. A structural density map of serotonergic projections was obtained from
the Allen Brain Atlas. The left color bar indicates beta values of BOLD responses corresponding to
Figure 2a. The right color bar indicates normalized intensity of structural density ranging from 0-50. (b)
Correlation between projection density and regional beta values. The correlation between regional
structural density and average regional beta values across transgenic groups was calculated from Session
1 (blue stimulation: $r = 0.462$, $p_{FDR} = 0.020$ ($p_{uncorrected} = 0.030$); yellow stimulation: $r = 0.132$, $p_{FDR} =$
0.529, two-tailed t tests and FDR corrected). Blue and yellow colors indicate data from blue and yellow
illumination sessions, respectively. Cyan and yellow highlights indicate 95% confidence intervals (CI)
for correlation coefficients. (c) Correlation between projection density and regional beta values. The
correlation between regional structural density and average regional beta values across transgenic groups
was calculated (Anesthetized (Session3): $r = 0.546$, $p_{FDR-corrected} = 0.012$ ($p_{uncorrected} = 0.006$); Awake
(Session4): $r = 0.746$, $p_{FDR-corrected} = 0.0002$ ($p_{uncorrected} = 0.00003$), two-tailed t tests and FDR corrected).
Magenta and blue colors indicate data from blue illumination sessions in Sessions 3 and 4, respectively.
Cyan and magenta highlight indicates 95% CI for correlation coefficients.
767

Figure 5. Regression analysis with 5-HT receptors and serotonergic projections. (a) Procedure to make a
model of predicted beta values in Session 1 of blue illumination. Independent variables, including
regional structural densities and regional gene expression of 5ht1a, 5ht1b, 5ht1f, 5ht2a, and 5ht2c were
fit with a linear regression model to beta values of multiple ROIs from blue illumination sessions. The
right color bar indicates normalized intensity of mRNA expression of 5-HT receptors and intensity of
structural density for projections. The color bar ranges from 0-80. (b) Correlation between modeled beta
values from Session 1 and beta values from Session 3 under anesthesia (r = 0.365, $p_{FDR-corrected}$ = 0.079,

two-tailed t test and FDR corrected). A magenta highlight indicates 95% CI. (c) Correlation between modeled beta values from Session 1 and beta values from Session 4 (Awake: $r = 0.615$, $p_{FDR-corrected} = 0.0019$ ($p_{uncorrected} = 0.0014$), two-tailed t test and FDR corrected). A cyan highlight indicates 95% CI. (d) Weights of fitted models from Session 1, Session 2, and Session 3 under anesthesia, and Session 4 without anesthesia. Each box indicates the range between the 25th percentile (bottom) and the 75th percentile (upper), and a horizontal line within each box is a median value. Whiskers of each box cover 99.3% (+/–2.7σ), and each red dot indicates a data point beyond 99.3% coverage. Cyan, black, magenta, and green colors indicate weights of expression profiles from Session 1 (n=8), Session 2 (n=7), Session 3 (n=7) and Session 4 (n=7), respectively.

ROI id	Area	ROI name	Abbreviations
1	Cortical regions	Frontal polar area	FP
2		Orbital area	OFC
3		Medial prefrontal cortex	mPFC
4		Secondary motor area	M2
5		Anterior cingulate cortex	ACC
6		Primary motor area	M1
7		Supplementary somatosensory area	SS
8		Ventral retrosplenial cortex	vRSC
9		Dorsal retrosplenial cortex	dRSC
10		Temporal association area	TAA
11		Primary auditory area	A1
12		Primary somatosensory area	S1
13		Posterior parietal association area	PPA
14		Primary visual area	V1
15	Sub-cortical regions	Nucleus accumbens	NAc
16		Caudate putamen	CPu
17		Bed nucleus of steria terminalis	BST
18		Globus pallidus external part	GPe
19		Globus pallidus internal part	GPi
20		Thalamic reticular nucleus	RT
21		Ventral pallidus	VP
22		Dentate gyrus of hippocampus	DG
23		CA1 of hippocampus	CA1
24		Lateral habenula	LHb
25		Substantia nigra pars reticulata	SNr
26		Ventral tegmental area	VTA
27		Medial raphe nucleus	MRN
28		Dorsal raphe nucleus	DRN

786
787 Table1. Regions of interest (ROIs) and their abbreviations. ROIs are ordered from anterior to posterior for
788 cortical and subcortical areas.
789

Model Statistics (Session 1)				
	R^2 statistic	F-statistic	P value (uncorrected)	Estimate of the error variance
Session 1	0.6603	5.5063	0.0025	0.0643

Table 2. Model with Parameter Estimates of Session 1. Multiple linear regression was performed to fit parameter estimates from Session 1 (blue stimulation) with the density of serotonergic projection and expression of 5-HT receptors.

5-ht1a		SS	df	MS	F-statistic	p value
	Group	0.00111	1	0.00111	0	0.945
	Error	6.20697	27	0.22989		
	Total	6.20808	28			
5-ht1b						
	Group	1.6679	1	1.66795	5.07	0.0327
	Error	8.8794	27	0.32886		
	Total	10.5473	28			
5-ht1f						
	Group	4.6674	1	4.66744	12.95	**0.0013***
	Error	9.7293	27	0.36034		
	Total	14.4967	28			
5-ht2a						
	Group	0.40786	1	0.40786	3.43	0.075
	Error	3.21128	27	0.11894		
	Total	3.61914	28			
5-ht2c						
	Group	10.103	1	10.103	31.08	**0.0019***
	Error	8.779	27	0.3251		
	Total	18.8809	28			
Structural density						
	Group	0.18596	1	0.18596	1.73	0.1999
	Error	2.90782	27	0.1077		
	Total	3.0938	28			
Intercept						
	Group	0.1135	1	0.11348	0.26	0.6171
	Error	11.9762	27	0.44356		
	Total	12.0896	28			

Table 3. Results of one-way ANOVA for 5-HT receptor type and structural intensity. One-way ANOVA was applied to regression coefficients from each factor to test whether each factor showed statistically different expressions between awake state and anesthetized states. SS, df, and MS indicate Sum of Squares, Degrees of freedom, and Mean Squares, respectively. * indicates statistical significance after multiple correction with the Bonferroni method ($p_{\text{Bonferroni-corrected}} < 0.05$).

References

1. Matias, S., et al., *Activity patterns of serotonin neurons underlying cognitive*
flexibility. Elife, 2017. **6**.
2. Boureau, Y.L. and P. Dayan, *Opponency revisited: competition and cooperation*
between dopamine and serotonin. Neuropsychopharmacology, 2011. **36**(1): p.
74-97.
3. Miyazaki, K., et al., *Reward probability and timing uncertainty alter the effect*
of dorsal raphe serotonin neurons on patience. Nat Commun, 2018. **9**(1): p.
2048.
4. Miyazaki, K.W., et al., *Optogenetic activation of dorsal raphe serotonin*
neurons enhances patience for future rewards. Curr Biol, 2014. **24**(17): p. 2033-
40.
5. Iigaya, K., et al., *An effect of serotonergic stimulation on learning rates for*
rewards apparent after long intertrial intervals. Nat Commun, 2018. **9**(1): p.
2477.
6. Ren, J., et al., *Single-cell transcriptomes and whole-brain projections of*
serotonin neurons in the mouse dorsal and median raphe nuclei. Elife, 2019. **8**.
7. Fu, W., et al., *Chemical neuroanatomy of the dorsal raphe nucleus and adjacent*
structures of the mouse brain. J Comp Neurol, 2010. **518**(17): p. 3464-94.
8. Li, Y., et al., *Serotonin neurons in the dorsal raphe nucleus encode reward*
signals. Nat Commun, 2016. **7**: p. 10503.
9. Liu, Z., et al., *Dorsal raphe neurons signal reward through 5-HT and glutamate.*
Neuron, 2014. **81**(6): p. 1360-1374.
10. Marcinkiewcz, C.A., et al., *Serotonin engages an anxiety and fear-promoting*
circuit in the extended amygdala. Nature, 2016. **537**(7618): p. 97-101.
11. Ohmura, Y., et al., *Optogenetic activation of serotonergic neurons enhances*
anxiety-like behaviour in mice. Int J Neuropsychopharmacol, 2014. **17**(11): p.
1777-83.
12. Ohmura, Y., et al., *Disruption of model-based decision making by silencing of*
serotonin neurons in the dorsal raphe nucleus. Curr Biol, 2021. **31**(11): p. 2446-
2454 e5.
13. Xu, S., et al., *Dorsal Raphe Serotonergic Neurons Control Intertemporal Choice*
under Trade-off. Curr Biol, 2017. **27**(20): p. 3111-3119 e3.
14. Miyazaki, K., et al., *Serotonergic projections to the orbitofrontal and medial*
prefrontal cortices differentially modulate waiting for future rewards. Sci Adv,
2020. **6**(48).
15. Grandjean, J., et al., *A brain-wide functional map of the serotonergic responses*
to acute stress and fluoxetine. Nat Commun, 2019. **10**(1): p. 350.
16. Yoshida, K., et al., *Physiological effects of a habituation procedure for*
functional MRI in awake mice using a cryogenic radiofrequency probe. J
Neurosci Methods, 2016. **274**: p. 38-48.
17. Abe, Y., et al., *Diffusion functional MRI reveals global brain network functional*
abnormalities driven by targeted local activity in a neuropsychiatric disease
mouse model. Neuroimage, 2020. **223**: p. 117318.
18. Youssef, M.M.M., et al., *TOB is an effector of the hippocampus-mediated acute*
stress response. Transl Psychiatry, 2022. **12**(1): p. 302.

19.   Tanaka, K.F., et al., *Expanding the repertoire of optogenetically targeted cells*
with an enhanced gene expression system. Cell Rep, 2012. **2**(2): p. 397-406.

20.   Sesack, S.R. and A.A. Grace, *Cortico-Basal Ganglia reward network:*
microcircuitry. Neuropsychopharmacology, 2010. **35**(1): p. 27-47.

21.   Garcia-Garcia, A.L., et al., *Serotonin inputs to the dorsal BNST modulate*
anxiety in a 5-HT(1A) receptor-dependent manner. Mol Psychiatry, 2018.
23(10): p. 1990-1997.

22.   Izquierdo, I., C.R. Furini, and J.C. Myskiw, *Fear Memory.* Physiol Rev, 2016.
96(2): p. 695-750.

23.   Trusel, M., et al., *Punishment-Predictive Cues Guide Avoidance through*
Potentiation of Hypothalamus-to-Habenula Synapses. Neuron, 2019. **102**(1): p.
120-127 e4.

24.   Tanaka, S.C., et al., *Serotonin differentially regulates short- and long-term*
prediction of rewards in the ventral and dorsal striatum. PLoS One, 2007.
2(12): p. e1333.

25.   Fonseca, M.S., M. Murakami, and Z.F. Mainen, *Activation of dorsal raphe*
serotonergic neurons promotes waiting but is not reinforcing. Curr Biol, 2015.
25(3): p. 306-315.

26.   Meyniel, F., et al., *A specific role for serotonin in overcoming effort cost.* Elife,
2016. **5**.

27.   Scholl, J., et al., *Beyond negative valence: 2-week administration of a*
serotonergic antidepressant enhances both reward and effort learning signals.
PLoS Biol, 2017. **15**(2): p. e2000756.

28.   Lottem, E., et al., *Activation of serotonin neurons promotes active persistence in*
a probabilistic foraging task. Nat Commun, 2018. **9**(1): p. 1000.

29.   Nagai, Y., et al., *The Role of Dorsal Raphe Serotonin Neurons in the Balance*
between Reward and Aversion. Int J Mol Sci, 2020. **21**(6).

30.   Pulcu, E., et al., *Temporal discounting in major depressive disorder.* Psychol
Med, 2014. **44**(9): p. 1825-34.

31.   Ren, J., et al., *Anatomically Defined and Functionally Distinct Dorsal Raphe*
Serotonin Sub-systems. Cell, 2018. **175**(2): p. 472-487 e20.

32.   Nishitani, N., et al., *Manipulation of dorsal raphe serotonergic neurons*
modulates active coping to inescapable stress and anxiety-related behaviors in
mice and rats. Neuropsychopharmacology, 2019. **44**(4): p. 721-732.

33.   Anderson, I.M., et al., *Assessing human 5-HT function in vivo with*
pharmacoMRI. Neuropharmacology, 2008. **55**(6): p. 1029-37.

34.   Schaefer, A., et al., *Serotonergic modulation of intrinsic functional connectivity.*
Curr Biol, 2014. **24**(19): p. 2314-8.

35.   Giorgi, A., et al., *Brain-wide Mapping of Endogenous Serotonergic*
Transmission via Chemogenetic fMRI. Cell Rep, 2017. **21**(4): p. 910-918.

36.   Boschloo, L., et al., *The complex clinical response to selective serotonin*
reuptake inhibitors in depression: a network perspective. Transl Psychiatry,
2023. **13**(1): p. 19.

37.   Arnone, D., *Functional MRI findings, pharmacological treatment in major*
depression and clinical response. Prog Neuropsychopharmacol Biol Psychiatry,
2019. **91**: p. 28-37.

38. Kim, M., et al., *The effects of selective serotonin reuptake inhibitors on brain functional networks during goal-directed planning in obsessive-compulsive disorder.* Sci Rep, 2020. **10**(1): p. 20619.

39. Klomp, A., et al., *Lasting effects of chronic fluoxetine treatment on the late developing rat brain: age-dependent changes in the serotonergic neurotransmitter system assessed by pharmacological MRI.* Neuroimage, 2012. **59**(1): p. 218-26.

40. Fritze, S., R. Spanagel, and H.R. Noori, *Adaptive dynamics of the 5-HT systems following chronic administration of selective serotonin reuptake inhibitors: a meta-analysis.* J Neurochem, 2017. **142**(5): p. 747-755.

41. Wang, H.Y., et al., *Frequency-Dependent Block of Excitatory Neurotransmission by Isoflurane via Dual Presynaptic Mechanisms.* J Neurosci, 2020. **40**(21): p. 4103-4115.

42. Winegar, B.D. and M.B. MacIver, *Isoflurane depresses hippocampal CA1 glutamate nerve terminals without inhibiting fiber volleys.* BMC Neurosci, 2006. **7**: p. 5.

43. Caraiscos, V.B., et al., *Selective enhancement of tonic GABAergic inhibition in murine hippocampal neurons by low concentrations of the volatile anesthetic isoflurane.* J Neurosci, 2004. **24**(39): p. 8454-8.

44. Ying, S.W., et al., *Isoflurane modulates excitability in the mouse thalamus via GABA-dependent and GABA-independent mechanisms.* Neuropharmacology, 2009. **56**(2): p. 438-47.

45. Puig, M.V. and A.T. Gulledge, *Serotonin and prefrontal cortex function: neurons, networks, and circuits.* Mol Neurobiol, 2011. **44**(3): p. 449-64.

46. Amargos-Bosch, M., et al., *Co-expression and in vivo interaction of serotonin1A and serotonin2A receptors in pyramidal neurons of prefrontal cortex.* Cereb Cortex, 2004. **14**(3): p. 281-99.

47. Puig, M.V., et al., *In vivo excitation of GABA interneurons in the medial prefrontal cortex through 5-HT3 receptors.* Cereb Cortex, 2004. **14**(12): p. 1365-75.

48. Matsunaga, F., et al., *Molecular interactions between general anesthetics and the 5HT2B receptor.* J Biomol Struct Dyn, 2015. **33**(1): p. 211-8.

49. Yokoyama, C., et al., *Marmoset Serotonin 5-HT1A Receptor Mapping with a Biased Agonist PET Probe 18F-F13714: Comparison with an Antagonist Tracer 18F-MPPF in Awake and Anesthetized States.* Int J Neuropsychopharmacol, 2016. **19**(12).

50. Lee, H.J., et al., *Activation of Direct and Indirect Pathway Medium Spiny Neurons Drives Distinct Brain-wide Responses.* Neuron, 2016. **91**(2): p. 412-24.

51. Abe, Y., et al., *Optical manipulation of local cerebral blood flow in the deep brain of freely moving mice.* Cell Rep, 2021. **36**(4): p. 109427.

52. Abe, Y., T. Tsurugizawa, and D. Le Bihan, *Water diffusion closely reveals neural activity status in rat brain loci affected by anesthesia.* PLoS Biol, 2017. **15**(4): p. e2001494.

53. Eichenbaum, H., *Prefrontal-hippocampal interactions in episodic memory.* Nat Rev Neurosci, 2017. **18**(9): p. 547-558.

54. Malik, R., et al., *Top-down control of hippocampal signal-to-noise by prefrontal long-range inhibition.* Cell, 2022. **185**(9): p. 1602-1617 e17.

55.   Oikonomou, G., et al., *The Serotonergic Raphe Promote Sleep in Zebrafish and*
949   *Mice.* Neuron, 2019. **103**(4): p. 686-701 e8.
950   56.   Smith, H.R., et al., *Dorsal Raphe Serotonin Neurons Mediate CO(2)-Induced*
951   *Arousal from Sleep.* J Neurosci, 2018. **38**(8): p. 1915-1925.
952   57.   Li, A., et al., *Dorsal raphe serotonergic neurons promote arousal from*
953   *isoflurane anesthesia.* CNS Neurosci Ther, 2021. **27**(8): p. 941-950.
954   58.   Mengod, G., et al., *5-HT receptors in mammalian brain: receptor*
955   *autoradiography and in situ hybridization studies of new ligands and newly*
956   *identified receptors.* Histochem J, 1996. **28**(11): p. 747-58.
957   59.   Nichols, D.E. and C.D. Nichols, *Serotonin receptors.* Chem Rev, 2008. **108**(5):
958   p. 1614-41.
959   60.   Pytliak, M., et al., *Serotonin receptors - from molecular biology to clinical*
960   *applications.* Physiol Res, 2011. **60**(1): p. 15-25.
961   61.   Boschert, U., et al., *The mouse 5-hydroxytryptamine1B receptor is localized*
962   *predominantly on axon terminals.* Neuroscience, 1994. **58**(1): p. 167-82.
963   62.   Riad, M., et al., *Somatodendritic localization of 5-HT1A and preterminal axonal*
964   *localization of 5-HT1B serotonin receptors in adult rat brain.* J Comp Neurol,
965   2000. **417**(2): p. 181-94.
966   63.   Cornea-Hebert, V., et al., *Cellular and subcellular distribution of the serotonin*
967   *5-HT2A receptor in the central nervous system of adult rat.* J Comp Neurol,
968   1999. **409**(2): p. 187-209.
969   64.   Giorgetti, M. and L.H. Tecott, *Contributions of 5-HT(2C) receptors to multiple*
970   *actions of central serotonin systems.* Eur J Pharmacol, 2004. **488**(1-3): p. 1-9.
971   65.   Lucaites, V.L., et al., *[3H]LY334370, a novel radioligand for the 5-HT1F*
972   *receptor. II. Autoradiographic localization in rat, guinea pig, monkey and*
973   *human brain.* Naunyn Schmiedebergs Arch Pharmacol, 2005. **371**(3): p. 178-84.
974   66.   Whittington, R.A. and L. Virag, *Isoflurane decreases extracellular serotonin in*
975   *the mouse hippocampus.* Anesth Analg, 2006. **103**(1): p. 92-8, table of contents.
976   67.   Massey, C.A., et al., *Isoflurane abolishes spontaneous firing of serotonin*
977   *neurons and masks their pH/CO(2) chemosensitivity.* J Neurophysiol, 2015.
978   **113**(7): p. 2879-88.
979   68.   Correia, P.A., et al., *Transient inhibition and long-term facilitation of*
980   *locomotion by phasic optogenetic activation of serotonin neurons.* Elife, 2017.
981   **6**.
982   69.   Mo, B., et al., *Restraint stress increases serotonin release in the central nucleus*
983   *of the amygdala via activation of corticotropin-releasing factor receptors.* Brain
984   Res Bull, 2008. **76**(5): p. 493-8.
985   70.   McDevitt, R.A., et al., *Serotonergic versus nonserotonergic dorsal raphe*
986   *projection neurons: differential participation in reward circuitry.* Cell Rep,
987   2014. **8**(6): p. 1857-1869.
988   71.   Eklund, A., et al., *BROCCOLI: Software for fast fMRI analysis on many-core*
989   *CPUs and GPUs.* Front Neuroinform, 2014. **8**: p. 24.
990   72.   Miyazaki, K., K.W. Miyazaki, and K. Doya, *Activation of dorsal raphe*
991   *serotonin neurons underlies waiting for delayed rewards.* J Neurosci, 2011.
992   **31**(2): p. 469-79.

993   73.   Puig, M.V., et al., *Serotonin modulates fast-spiking interneuron and*
994   *synchronous activity in the rat prefrontal cortex through 5-HT1A and 5-HT2A*
995   *receptors.* J Neurosci, 2010. **30**(6): p. 2211-22.
996   74.   Salvan, P., et al., *Serotonin regulation of behavior via large-scale*
997   *neuromodulation of serotonin receptor networks.* Nat Neurosci, 2023. **26**(1): p.
998   53-63.
75.   Kawai, S., et al., *Effect of three types of mixed anesthetic agents alternate to*
ketamine in mice. Exp Anim, 2011. **60**(5): p. 481-7.
76.   Hikishima, K., et al., *In vivo microscopic voxel-based morphometry with a brain*
template to characterize strain-specific structures in the mouse brain. Sci Rep,
2017. **7**(1): p. 85.
77.   Li, R., et al., *Restoring susceptibility induced MRI signal loss in rat brain at 9.4*
T: A step towards whole brain functional connectivity imaging. PLoS One,
2015. **10**(4): p. e0119450.
78.   Spisak, T., et al., *Probabilistic TFCE: A generalized combination of cluster size*
and voxel intensity to increase statistical power. Neuroimage, 2019. **185**: p. 12-
26.
79.   Harris, J.A., et al., *Anatomical characterization of Cre driver mice for neural*
circuit mapping and manipulation. Front Neural Circuits, 2014. **8**: p. 76.
1012