Optogenetic activation of dorsal raphe serotonin neurons induces brain-wide activation, including reward-related circuits

Hiro Taiyo Hamada[1,3,*], Yoshifumi Abe[2], Norio Takata[2], Masakazu Taira[1], Kenji F. Tanaka[2], Kenji Doya[1,*]

[1] Neural Computation Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

[2] Department of Neuropsychiatry, Keio University School of Medicine, Tokyo, Japan

[3] Research & Development Department, Araya Inc., Tokyo, Japan

* Correspondence to hamada_h@araya.org, doya@oist.jp

Abstract

Serotonin is a neuromodulator that affects multiple behavioral and cognitive functions. Nonetheless, how serotonin causes such a variety of effects through brain-wide projections and various receptors remains unclear. Here we measured the brain-wide response to optogenetic stimulation of serotonin neurons in the dorsal raphe nucleus (DRN) with functional MRI using an 11.7T scanner with a cryoprobe. Transient activation of DRN serotonin neurons caused brain-wide activation, including reward-related brain areas. The same stimulation under anesthesia with isoflurane decreased activation of reward-related areas and the hippocampal complex. These brain-wide response patterns can be explained by DRN serotonergic projection topography and serotonin receptor expression profiles, with enhanced weights on 5-HT1 receptors. Together, these results provide foundational understanding of the primarily facilitatory effect on brain areas by the DR serotonergic system, which is consistent with recent discoveries of its roles in adaptive behaviors.

Introduction

Serotonin, 5-hydroxytryptamine (5-HT), serves crucial behavioral and cognitive functions, including awake-sleep cycle, mood, reward and punishment predictions [5, 32], patience [37, 39], and learning [19]. Serotonin is also a major target of medications for psychiatric disorders, notably major depression, by selective serotonin re-uptake inhibitors (SSRIs). The dorsal raphe nucleus (DRN) is a major source of serotonergic projections to brain-wide targets, including the frontal cortex, the basal ganglia, the ventral tegmental area (VTA), and other mid-brain regions [50]. The DRN contains glutamatergic, GABAergic, and dopaminergic neurons, in addition to serotonergic neurons [12]. Optogenetic manipulation of serotonin neuron activities has revealed their roles in reward [25-26] , patience [37, 39], anxiety [30, 43], and learning [19, 44]. Nonetheless, how brain-wide serotonergic projections cause such effects and by which target areas and receptors, remain poorly understood [38, 63].

In order to clarify the global picture of serotonergic control of brain activities, we performed optogenetic functional MRI experiments using Tph2-ChR2(C128S) transgenic mice and an 11.7T MRI scanner with a cryoprobe for high S/N measurement. A previous optogenetic fMRI experiment activated DRN serotonin neurons using the Pet1 promoter under anesthesia and showed widespread suppression of cortical activities [16]. We introduced an acclimation protocol to allow stable fMRI recording from awake mice [1, 66] and found that DRN serotonin activation in an awake state induced a brain-wide activity increase both in the cortical and subcortical areas, including reward-related circuits. In contrast, under anesthesia with isoflurane, DRN serotonin activation decreased activities in those areas, consistent with a previous study [16].

We further found that regional BOLD responses can be approximated by a linear model with projection densities of DRN serotonin neurons and expression profiles of different types of serotonin receptors, with negative coefficients for 5-HT1 receptors and positive coefficients for 5-HT2 receptors. These results suggest that serotonergic neurons activate brain-wide regions, including reward-related circuits, and they demand reconsideration of implications of results under anesthesia.

Results

In order to clarify effects of brain-wide serotonergic projections, we employed optogenetic fMRI combining optogenetic stimulation of serotonergic neurons in the dorsal raphe nucleus (DRN) and measurements of whole-brain activities by functional MRI. We used Tph2-ChR2(C128S) transgenic mice [43, 57] which express step-type channel rhodopsin selectively in serotonin neurons [39, 43]. We used an 11.7T MRI scanner (Bruker BioSpec 117/11) with a cryoprobe for high S/N recording of BOLD signals from the whole brain, except the olfactory bulb and the posterior cerebellum.

Prior to experiments, the transgenic and wild-type (WT) groups (Transgenic group: n = 8; Wild-type: n = 6) underwent surgery to attach a plastic fixation bar and implant an optical cannula in the DRN. Acclimation training in a mock chamber followed (Figure 1. See Methods for details). After fMRI experiments, animals were tested in a reward

84   waiting task [37, 39] to confirm the effectiveness of optogenetic stimulation
85   (Supplementary Figure S1).
86
DRN Serotonin activation induces a brain-wide increase in BOLD signals
We measured BOLD responses under optogenetic stimulation of DRN serotonin
neurons in an ON/OFF block design protocol (Figure 1e). Photo-stimulation increased
BOLD signals in the DRN (Figure 1f), with a tendency for reduced response after
repeated stimulation. On average, over five stimulations, blue light stimulation induced
a significantly higher BOLD response than did yellow light stimulation ($p < 0.05$,
correction of multiple comparisons with the false discovery rate (FDR), see 'General
Linear Model (GLM) for BOLD signals' in Methods; Figure 1g and Supplementary
Figure S5).
96
A group analysis with a general linear model (GLM) of photo-activated BOLD
responses between blue and yellow stimulation from the transgenic group($n = 8$)
revealed a brain-wide increase in BOLD responses, significantly in the frontal cortex,
the caudate putamen (CPu), the temporal cortical regions, the thalamic complex, and the
VTA ($p < 0.05$, cluster-wise correction controlling family wise error (FWE) with
probabilistic threshold-free cluster enhancement (pTFCE); Figure 2a). Trial-wise BOLD
responses also showed consistent brain-wide activation by blue, but not yellow
stimulation ($p < 0.05$, uncorrected, Supplementary Figure S2a-b).
105
We next selected 28 regions of interest (ROIs) from the cortex and subcortical areas that
are involved in reward and punishment, according to the Allen mouse brain atlas (Table
1, see Regions of interest in Methods). For the reward system, we selected components
of basal ganglia such as caudate putamen (CPu), nucleus accumbens (NAc), and globus
pallidus (GP) [54]. For the punishment system, the bed nucleus of the stria terminalis
(BST) [13], the hippocampus [20], and the lateral habenula (LHb) [59] were selected.
Then, we extracted regional average beta values (regression coefficients) from the
GLM. Comparison of regional beta values between blue and yellow stimulation from
the transgenic group revealed that the BOLD response was significantly higher in the
blue illumination session in 12 of 14 cortical regions, including the OFC, mPFC and
ACC, and 12 of 14 subcortical regions, including the CPu, NAc, GP, VTA, LHb, and
DRN (Figure 2b) although no statistical differences were found in the WT group
(Supplementary Figure S4). Moreover, there was no significantly higher BOLD
response in vRSC, V1, DG, or CA1.
120
The week after session 1, we performed another session under the same stimulation
parameters as in session 1. Patterns of BOLD responses were similar to those in session
1(transgenic: $n = 7$, WT: $n = 6$, Supplementary Figure S3). Pearson's correlation of the
beta values in sessions 1 and 2 was $r = 0.875$, which was statistically significant ($p < 1e-$
8; Supplementary Figure S7.a). We further checked regional differences of responses in
peak amplitude and peak timing by extracting regional BOLD signals (Supplementary
Figure S5-9). Peak amplitude and peak timing of BOLD responses varied across the 28
ROIs (Supplementary Figure S8.c-d), with the largest responses in the VTA, SNr, and
FP. Both amplitude and timing of BOLD responses in sessions 1 and 2 were
significantly correlated for blue light stimulation (amplitude: $r = 0.886$, $p < 1e-8$;

timing: r = 0.679, p < 0.005), but not for yellow stimulation (amplitude: r = -0.06, p =
0.766; r = 0.037, p = 0.855; Supplementary Figure S8.a, b). These results indicate that
our photo-stimulation of DRN induced consistent activation of DRN serotonergic
neurons, which caused distinct spatial and temporal responses in multiple cortical and
sub-cortical regions.
136
DRN Serotonin activation induces a brain-wide negative response in BOLD signals
under anesthesia
Contrary to our above results, a previous optogenetic fMRI study reported that DR
serotonin activation evoked negative cortical cerebral blood volume (CBV) responses,
an enhanced proxy of BOLD signals, from the entire cortical region under anesthesia
[16]. There are two critical differences between that study and this one: optogenetic
probes and anesthesia. The previous study used the ePET-Cre mouse line and AAV
harboring Cre-dependent ChR2 and applied 20-Hz photostimulation. On the other hand,
the current study used Tph2-ChR2(C128S) transgenic mice, which express step-type
opsin, to allow continuous activation by a pulse of blue light (1.0 s, 473 nm) at the
beginning and yellow light (1.0 s, 593 nm) at the end (Figure 1e). Another difference is
that the previous study employed isoflurane, a general gas anesthetic with a muscle
relaxant, while our experiment employed no anesthetics.
150
In order to test the possibility that the difference is due to anesthesia, we compared
BOLD responses to DRN serotonin stimulation between anesthetized and awake states
in the same animals. We performed the same ON/OFF stimulation protocol under
anesthesia with 1% isoflurane with the same groups (Transgenic: n = 7, WT: n = 5,
Figure 3a). We also performed another ON/OFF stimulation session the next day in an
awake state in order to eliminate possible confounding by diminished responses after
repetitive photo-stimulation (Figure 3b).
158
Under anesthesia, we observed mostly negative BOLD responses from the entire
cortical region, while BOLD signals in the DRN remained positive (statistical t > 1,
uncorrected; Figure 3a & Supplementary S10a). On the other hand, session 4 under an
awake state replicated positive BOLD signals from mid-line cortical regions and DRN
(statistical t > 1, uncorrected; Figure 3b & Supplementary Figure S10b). We then
contrasted regional beta values in anesthetized and awake states (Figure 3c). In the
anesthetized state, multiple regions, such as ACC, dRSC, vRSC, DG and CA1 showed
negative beta values, significantly different from those in the awake state (p < 0.05,
correction of multiple comparisons with FDR). We further found a statistically
significant difference in peaks of % BOLDs of these regions between anesthetized and
awake states (p < 0.05, Figure 3d; Supplementary Figure S11). Our results support the
hypothesis that general anesthesia induced negative BOLD responses by DR serotonin
photo-activation.
172
Brain-wide BOLD responses are explained by serotonin projection density and receptor
expression profiles
Our results revealed a brain-wide profile of evoked BOLD signals by DR serotonin
activation (Figure 2a). One plausible mechanism underlying such a spatial profile is the
density of the serotonergic projection pattern from the DRN. In order to test such a

possibility, we checked the correlation between intensity of projections of DRN
serotonin neurons and peak levels of BOLD responses. We extracted serotonin-specific
structural projection density from DRN from the Allen Brain Atlas (Figure 4a).
181
Serotonin projection density was statistically correlated with beta values under blue
stimulation ($r = 0.462$, $p < 0.05$, FDR corrected) but not under yellow stimulation ($r =$
0.132, $p = 0.529$; Figure 4b). We also found that projection density was significantly
correlated with photo-stimulated BOLD responses in awake and anesthetized states
(Anesthetized (Session3): $r = 0.546$, $p < 0.05$; Awake (Session4): $r = 0.758$, $p < 0.001$,
FDR corrected; Figure 4c).
188
We further investigated whether photo-stimulated BOLD responses are associated
with serotonin synaptic receptors. Gene expression maps for 5-HT1A, 5-HT1B, 5-
HT1F, 5-HT2A, and 5-HT2C receptors were obtained from the database of the Allen
Brain Atlas (Supplementary Figure S12&S13). We then constructed a multiple linear
regression model (MLRM; Figure 5a) composed of one dependent variable of the
BOLD response and six independent variables of the projection density and five gene
expression patterns (5-HT1a, 5-HT1b, 5-HT1f, 5-HT2a, and 5-HT2c receptors) after
normalization. The model reproduced the BOLD response with $R^2=0.66$ ($p < 0.01$,
Table 2). We also checked the consistency of the model using BOLD responses in
different sessions. Using the fit beta values with session 1, we found a significant
correlation with the beta value in session 4 (Awake: $r = 0.615$, $p < 0.01$, FDR corrected,
Figure 5b), but not in session 3 (Anesthetized (Session3): $r = 0.365$, $p = 0.079$, FDR
corrected; Figure 5c).
202
We investigated the contribution of each receptor by the estimated regression
coefficients in anesthetized and awake states (Figure 5d, Table 3, and Supplementary
Table 3). For each subject, we estimated regression coefficients from each session. In
the awake state (Session 1,2, and 4), gene expression of 5-HT1 receptors (5-HT1a, 5-
HT1b, and 5-HT1f) showed average negative coefficients, indicating negative
contributions to BOLD responses. On the other hand, gene expression of 5-HT2
receptors showed average positive coefficients, indicating positive contributions to
BOLD responses. We also checked whether expression of each receptor type differed
between awake and anesthetized states with one-way analysis of variance (ANOVA).
The results revealed that in the anesthetized state (Session 3), weights for 5-HT1f and 5-
HT2c receptors were significantly different ($p < 0.05$, Bonferroni corrected, Table 3).
214
215 These results suggest that photo-activated BOLD responses reflect projection density of
216 serotonin neurons and serotonin receptor expression profiles and that general anesthesia
217 affects responses of 5-HT1 and 5-HT2 receptors differently to induce altered BOLD
218 responses.
219

Discussion

Our opto-fMRI investigation of the DRN serotonergic modulations of brain-wide
dynamics brought three major discoveries. First, DRN serotonergic activation causes
brain-wide positive BOLD responses in an awake state. Second, DRN serotonergic
activation under general anesthesia causes brain-wide negative BOLD responses. Third,

the whole-brain profile of DRN serotonin-activated BOLD signals is associated with the
density of DRN serotonergic projection and the expression profiles of 5-HT receptors,
with opposite weights for 5-HT1 and 5-HT2 receptors.
228
The present result demonstrating brain-wide activation by DRN serotonergic activation
provides a means of interpreting recording and manipulation studies in humans and
animals [10, 19, 27, 36-39, 53, 58]. Previous studies have demonstrated serotonin's
involvement in motivation and reward-related learning[5, 36] through modulation of the
ACC and ventromedial PFC on learning signals [53], CPu on short/long-term prediction
of rewards [58], and mPFC and OFC on waiting for delayed rewards [38]. Our opto-
fMRI results support those studies, as those areas were consistently activated by DRN
serotonin stimulation. In addition, average amplitudes of BOLD responses and peak
timing are region-dependent. This finding implies that serotonin-activated BOLD
responses are coupled with different regional distributions of receptor densities and
serotonin projection, but not with the distance of the activation site. In spite of indirect
evidence, our linear approximation of serotonin-activated BOLD responses support
such a possibility. A subpopulation of DR serotonin projections to the cortex is also
expressed with the vesicular glutamate transporter Vglut3 (Slc17a8). Co-release of
glutamate has recently been reported [49-50]. Our results may reflect the influence of
serotonin as well as glutamate co-release on downstream circuitry.
245
Serotonergic modulation in short-term windows using optogenetics can avoid
limitations of long-term windows, such as pharmacological manipulation [3, 52] and
chemogenetic manipulation with DREADD (designed receptors exclusively activated
by designed drugs) [15]. Multiple pharmacological studies have applied SSRIs to
manipulate the serotonin system. Systemic SSRI administration causes brain-wide
inhibitory modulation [3, 52]. However, the influence of SSRI administration on brain
activity can be confounded by the administration period due to receptor sensitivity [11,
22, 52].
254
A previous study stimulated the serotonin system with specifically designed drugs
under fMRI [15] and revealed brain-wide positive responses in the somatosensory
cortex and subcortical regions, including the hippocampus, the hypothalamus, and the
cerebellum. That study did not reveal BOLD responses of mid-line cortical regions,
such as the OFC, mPFC and ACC. A possible explanation for the difference between
their results and ours is that BOLD responses cause different time scales, depending on
receptor types. Our optogenetic fMRI approach revealed an influence on motivation-
related regions at different timings in a transient time window (Supplementary Figure
S5, S6 and S8). Further studies using different methodologies, including optogenetics
and chemogenetics, will uncover neural mechanisms underlying different time scales of
serotonin modulation.
266
The mechanism underlying different serotonin modulation between awake and
anesthetized states remains unknown. One possible mechanism is a reduction of
glutamate transmission via serotonin in target regions [60]. Previous studies revealed
that isoflurane suppresses neural activity via inhibition of glutamatergic transmission
[60, 62]. We found that isoflurane reduced photo-stimulated BOLD signals where they

were potentiated in an awake state. Our results also showed that some regions show
potentiated negative BOLD responses. In addition to reduced glutamatergic
transmission, isoflurane also enhances sensitivity of GABA neurons [6, 64]. Such
elevated responses of GABA neurons and reduced responses of glutamatergic neurons
may result in diminished positive BOLD signals and potentiated negative BOLD
signals. It is also interesting to ask whether different anesthetics like NMDA antagonists
induce different effects on receptor responses while neuron-specific distributions of
serotonin receptors are unclear. Serotonin receptors in the prefrontal cortex are
differentially distributed [47]. 5-HT1A and 2A receptors are enriched on glutamatergic
pyramidal neurons in the PFC [2] . Meanwhile, 5-HT3A receptors are located on
GABAergic interneurons [46]. Since different anesthetics influence different neuron
types in various ways, reactivity of serotonin receptors may induce different responses.
In addition, anesthetics may change effects of serotonin receptors. Some studies showed
that isoflurane changes binding affinities of 5-HT$_{2B}$ receptors [33] in vitro and 5-HT$_{1A}$
receptors [65] in vivo. Anesthetics can influence ligand binding to specific subtypes of
serotonin receptors.
288
Furthermore, our results showed negative BOLD response in dorsal hippocampus under
anesthesia despite no projections to hippocampus from the DRN. One potential cause is
indirect influence by downstream circuitry of DR serotonin neurons. For example, DR
serotonin neurons have dense projections to the PFC, which interacts with the
hippocampus [9, 29]. Reduction of the BOLD response in the hippocampus under
anesthesia can be indirectly caused by activation of DR serotonergic neurons through
their downstream circuitry, although the influence is not observable in awake states.
The brain-wide response by activation of DR serotonin neurons under anesthesia may
have implications for sleep state. Similarity of brain states between general anesthesia
and non-rapid-eye-movement (NREM) sleep is implicated [45]. Previous studies also
revealed that DR serotonin neurons mediate the sleep/wake cycle [23, 55]. Optogenetic
activation of DR serotonin neurons prolonged wakefulness and decreased NREM sleep
while their inactivation prevented arousal from CO_2-induced sleep [55]. Similarly,
optogenetic and chemogenetic activation induced arousal from isoflurane [23]. The
negative response under anesthesia may reflect a brain response associated with arousal.
Future studies are needed to clarify the neural mechanism underlying BOLD signals
under anesthesia.
306
We have also demonstrated that different types of 5-HT gene expression are associated
with photo-activated BOLD responses. Distinctive contributions of serotonin receptor
types influence neural activities mostly at metabotropic receptors [35]. 5-HT1 receptors
are inhibitory metabotropic receptors that suppress neural activity by modulating the
cellular signal cascade, whereas 5-HT2 receptors are excitatory metabotropic receptors
that activate neural activity [42, 48]. Our MLRM analysis revealed that regression
coefficients of gene expression of 5-HT1 receptors and 5-HT2 receptors are negative
and positive, respectively. The correspondence between functions of receptor types and
signs of coefficients implies that BOLD responses by DRN serotonin photo-activation
can be influenced by different receptor distributions. Localization of 5-HT receptors is
subtype-dependent. 5-HT$_{1A}$ and 5-HT$_{1B}$ receptors are pre-synaptically expressed either
on somato-dendrites or within axon terminals of DR serotonin neurons [4, 51], whereas

5-HT$_{2A}$ and 5-HT$_{2C}$ receptors are predominantly expressed on postsynaptic neurons [7,
14]. However, whether 5-HT$_{1F}$ receptors are pre- or post-synaptically expressed is
unclear, although they are enriched in cortical layers (layer IV and V) and the caudate
putamen [28]. Nonetheless, specifically how 5-HT receptors react to serotonin
transmission under general anesthetics is unclear. It is also plausible that general
anesthetics change binding affinity of 5-HT receptors [33, 65]. As another potential
mechanism, serotonin transmission can be influenced by general anesthetics. Some
studies revealed that isoflurane lessens spontaneous serotonin levels and spontaneous
firing [31, 61]. These changes of serotonin transmission and receptor binding affinity
may influence responses of 5-HT receptors under anesthesia.
329
The substantial reduction of magnitude of brain-wide responses between the first,
second, and third sessions is noteworthy, despite consistent activation patterns.
Repeated optogenetic stimulation of DR serotonin neurons and repeated stress
potentially account for this reduction. Persistent changes in spontaneous behaviors by
repeated optogenetic activation of DR serotonin neurons have been reported [8].
Serotonin is an inducing factor of synaptic plasticity, and repeated stimulation may
cause plasticity changes. In addition, repeated restraint stress also may change responses
to optogenetic stimulation of DR serotonin neurons, although the habituation protocol to
awake fMRI is introduced to prevent restraint stress. Restraint stress increases serotonin
release in the amygdala [40]. Molecular and neuronal mechanisms underlying such
diminitions are needed to understand long-term effects of serotonin stimulation.
341
While our results replicated negative cortical responses under anesthesia [16], there are
several important differences in methodology and results between previous studies and
this one. First, one previous study utilized cerebral blood volume (CBV) signals with a
contrast agent to enhance influence of serotonin modulation, although the current study
employed BOLD signals. Although CBV signals are an enhanced proxy of BOLD
signals, the methodological difference in imaging may show different responses from
various brain regions. Second, we employed a transgenic group targeting the tryptophan
hydroxylase 2(Tph2) promoter [43, 57] although the previous study employed the
ePET-Cre line, targeting transcription factor Pet1 [16]. While expression of ChR2 by
Tph2 is selective for serotonergic neurons [41], Pet1 promoters are also expressed in
non-serotonergic neurons [26, 34]. While CA1 and DG were significantly deactivated
by serotonin photo-activation in our study, responses in temporal regions were not
found in the previous study [16]. Additionally, our results reveal an association between
5HT projection density and BOLD responses, but the previous study did not find this
association. One critical methodological difference is that we employed different ROIs
from those of the previous study. In the current study, 28 ROIs from the Allen brain
atlas, included other sources of serotonin and dopamine, the MRN and VTA. In
contrast, the previous study used 38 ROIs from the Australian Mouse Brain Mapping
Consortium (AMBMC, https://www.imaging.org.au/AMBMC) with multiple ROI
hippocampal formations and midbrain regions. The ROI difference may yield different
results in the association between structural connectivity and responses to optogenetic
serotonin activation. Further research is required to dissect differences between the two
sets of experimental results.
365

In conclusion, we reported brain-wide responses of DRN serotonin activation via opto-
fMRI for the first time in awake animals. We found serotonergic modulation of
widespread cortical and sub-cortical targets at different timing as a key controller of
brain-wide information transmission. We also showed different modulation by the
serotonin system in awake and anesthetized states. We further showed that serotonergic
structural projections and gene expression profiles of 5-HT receptors are associated with
BOLD responses. Further research is required to understand complex serotonin
modulation of brain-wide dynamics, given its neural and molecular mechanisms, such
as interactions with neural populations and receptors. Our findings reveal a new
physiological basis for understanding serotonergic control of brain-wide dynamics and
functions.
377
Materials and Methods
379 All experimental procedures were performed under protocols approved by the Okinawa
380 Institute of Science and Technology Experimental Animal Committee.
381
Animals
Serotonin neuron-specific channelrhodopsin 2 variant(C128S) mice were generated by
crossing Tph2-tTA mice with tetO-ChR(C128S)-EYFP knock-in mice under control of
the tryptophan hydroxylase 2(Tph2) promoter [43, 57]. Eight bi-genic and six wild-type
(WT) C57BL/6J male mice (age > 12 w.o.) were used in this study. Mice were housed
at 24 ∘C. on a inverted dark:light cycle (lights on 10:00-22:00, GMT+9). After
stereotaxic surgery, mice underwent habituation training [66], MRI experiments, and a
reward delay task in turn.
390
Stereotaxic surgery
Mice were initially sedated with 1-3% isoflurane, and were administered three types of
mixed anesthetic agents (1 (mg/mL) of medetomidine, 5.0 (mg/mL) of midaloram, and
5.0 (mg/mL) of butorphanol, i.p.) [21]. Cranial skins were removed with a sterilized
surgical knife, and a plastic bar (3x3x27mm) for head-fixation in the MRI environment
and an optic fiber (480 µm diameter, 4 mm length, Doric Lenses) was fixed to their
skulls with dental acrylic cement. The optical fiber was inserted horizontally toward the
DRN from the cerebellum in each mouse. Mice were housed separately after surgery
and underwent at least a 1-week recovery period. Mice were habituated to the MRI
environment for at least another week. They subsequently underwent imaging sessions.
401
Acclimation
In order to habituate awake mice to the MRI imaging protocol, an acclimation protocol
was applied to the mice after the surgeries for head-fixation [1, 66] . Mice were fixed in
a mock MRI environment and exposed to MRI imaging sounds and blue illumination
for 2 h to habituate them to a leak of illumination during scanning (2.0 Hz). Acclimation
to sounds and illumination was repeated 7 days prior to the first imaging session.
Following imaging sessions, at least 3 days acclimation were permitted.
409
Functional magnetic resonance imaging (fMRI)
Each animal was placed in the MRI animal bed. T2-weighted structural imaging and
three T2* functional imaging sessions were performed. MRI images were acquired with

an 11.7-tesla MRI scanner for small animals (Biospec 117/11 system, Bruker Biospin,
EmbH, Ettlingen, Germany) with a cryogenic quadrature RF surface probe (Cryoprobe,
Bruker BioSpin AG, Fällanden, Switzerland). We performed T2-weighted structural
imaging with a RARE sequence with subsequent parameters: 140 x 140 matrix, 13.5 x
13.5 mm2 field-of-view, repetition time(TR)/effective echo time(TE) 4000/18.75 (ms),
31 coronal slices, slice thickness: 300 µm.
419
After T2-weighted structural imaging was performed, a functional MRI session was
further executed. A functional MRI session was composed of two imaging runs, a blue-
yellow light stimulation run and a yellow-light stimulation run. All functional imaging
was performed using a GE-EPI sequence with following parameters: 67 x 67 matrix,
13.5 x 13.5 mm2 field-of-view, TR/TE 1000/10.7 (ms), flip angle: 50◦, bandwidth:
333k(Hz), 31 coronal slices and slice thickness: 300 µm, and 690 repetitions.
426
ON/OFF stimulation paradigm
A stimulation protocol was composed of ON/OFF stimulation patterns (Figure 1d). In
each activation session, stimulation was initiated 90 volumes after the onset of an
imaging session. DRN serotonin neurons were stimulated with blue illumination (490
nm, 3.0 mW, 1.0 sec), and stimulation subsequently ceased with yellow illumination
(630 nm, 3.0 mW, 1.0 sec) 20 seconds after the onset of blue illumination. This
stimulation was repeated 5 times for 120 seconds after the onset of blue illumination.
For the control session, instead of blue illumination, only yellow illumination was
applied in stimulation sessions.
436
Preprocessing and de-noising
The first 10 volume images of each session were removed from subsequent analysis due
to initial imaging instability caused by motion and non-steady state image quality. 10-
fold magnification of images was done to process images with Statistical Parametric
Mapping 12 (SPM12), which is designed to process human brain images. Therefore, the
voxel size was 2 x 2 x 3 mm in analytical steps. Pre-processing, including motion
correction, realignment, co-registration, normalization to a C57BL6/J template [18], and
spatial smoothing (kernel with 3 x 3 x 6 mm) was executed with Statistical Parametric
Mapping (SPM12). Since each slice of a 3D brain image was taken at a different time,
slice timing correction was applied to correct slice-timing differences by temporal
interpolation. Next, realignment of 3D brain images was conducted to correct motion-
related changes of brain position. Then, co-registration was executed to overlay
functional T2* images onto structural T2 images and to save coordination changes of
T2* images for normalization. During normalization, T2 structural images were first
warped to fit the average C57BL6/J template [18]. Then T2* functional images were
further warped to the template using the co-registered coordination change of the
images. In spatial smoothing, voxel signals were spatially smoothed using a gaussian
kernel (4 x 4 x 6 mm). After pre-processing steps, the general linear model in SPM12
was executed to map spatial activation patterns.
456
Regions of Interest
The Allen brain institute provides a parcellation of the C57BL6 mouse brain. In order to
extract BOLD signals from anatomically defined brain regions, this parcellation was

utilized for subsequent analyses. Parcellation was first warped to the MRI template
using Advanced Normalization Tools (ANTs), and 28 regions of interest (ROIs) were
selected from the warped parcellation. 28 ROIs were selected with the following three
criteria. First, primary cortical regions, especially mid-line regions, were selected.
Second, primary reward- and punishment-related sub-cortical regions, including basal
ganglia, VTA, and DRN should were included. Third, ventro-lateral cortical and sub-
cortical regions, such as the amygdala complex were not selected due to its sensitivity to
artifacts near air cavities by the ear canals [24].
468
General Linear Model (GLM) for BOLD signals
A general linear model (GLM) was performed to map beta values of parameter
estimation of optogenetic stimulation onto the mouse brain template with SPM12. For
statistical functional mapping of optogenetic stimulation, each voxel under an
uncorrected P value < 0.001 was thresholded, and cluster-wise multiple comparisons
were subsequently executed with family-wise error (©) by thresholding pFWE < 0.05
with probabilistic threshold-free cluster enhancement (pTFCE) [56]. Statistical second-
level voxel-wise analysis was finally executed to compare functional maps of blue and
yellow illumination from transgenic mice. For session 1, all procedures were executed
accordingly. For sessions 2, 3, and 4, none of the voxels under uncorrected P values <
0.001 survived the thresholding.
480
Beta values from BOLD responses
Furthermore, regional beta values were extracted based on ROIs. Statistical significance
of regional beta values was subsequently tested with Student's t tests with multiple
comparisons with the false discovery rate (FDR; p value (FDR) < 0.05). Each regional
BOLD signal change (%) is calculated by averaging regional time series on voxels,
which are scaled with the mean of time series during the no-stimulation period and
followed with multiplication by 100.
488
Structural density and RNA expression maps
The structural density map of serotonin connections was obtained from the mouse brain
database of the Allen Brain Institute (Experimental IDs: 480074702) [17]. As with the
structural density map, gene expression of serotonin receptors was obtained from the
mouse brain database of Allen Brain Institute (Experimental IDs: 79556616 (5HT1a),
583 (5HT1b), 69859867 (5HT1f), 81671344 (5HT2a), and 73636098 (5HT2c)).
Normalized energy intensity was extracted from the normalized atlas. Multiple linear
model analysis of ROIs between betas from GLM analyses and gene expressions was
performed.
498
Multivariable Linear Regression Model (MLRM)
Beta values from each session were obtained based on 28 ROIs. Structural density and
RNA expression of 5HT1a, 1b, 1f, 2a, and 2c was also extracted based on 28 ROIs.
Multivariable linear models were used to estimate weights of structural and gene
expression of 5HT receptors; where beta values from each session were the dependent
variables, and intensity of structural and gene expression was the explanatory variable.
The results of statistical values for fitting are described in Table 2.
506

One-way Analysis of Variance (ANOVA) on Receptor Contribution
Regression coefficients of structural density and RNA expression of 5-HT receptors
were calculated for each subject for all sessions. One-way ANOVA was applied to
regression coefficients of each receptor type in order to test whether receptor expression
is statistically different between awake (sessions 1, 2, &4) and anesthetized states
(session 3). For the analysis, data from all sessions were concatenated (total data: n=29,
session 1 (awake): n=8; session 2(awake): n=7, session 3 (anesthetized): n=7, session 4
(awake): n=7). p values from one-way ANOVA were further corrected with the
Bonferroni method. Results of ANOVA are described in Supplementary Table ST3.
516
A reward delay task
After imaging sessions, some mice from both groups (Transgenic: n=3, WT: n=2)
subsequently performed a waiting experiment for a delayed reward. A sequential tone-
food waiting task was utilized for a free operant task [37, 39] (Supplementary Figure
S1a). After MRI sessions, individual subjects were trained and tested in an operant-
conditioning box (Med-Associates, 12.7x12.7 cm). In short, subjects were required to
initiate trials by nose-poking into a tone site 500 (ms). The reward-delay task was
composed of reward trials (75%) and omission trials (25%). We adopted two types of
stimulation protocols (Supplementary Figure S1b): fixed-delay and random-delay
protocols. At the onset of reward and omission trials, either blue or yellow illumination
was delivered. In the reward trials, blue/yellow light stimulation was delivered at the
onset of each trial until nose-poking was performed, and yellow light was delivered
when subjects waited for 6.0 (s) for the fixed protocol and 2, 6, or 10 (s) for the random
delay protocol. However, yellow illumination was also given when subjects failed to
wait for the delay. On the other hand, during omission trials, yellow stimulation was
given when mice gave up waiting for delayed rewards while light stimulation was
delivered in the same manner as in the reward trials.
534
535   Data and code availability
536   Raw data including functional and structural images will be available soon. Codes and
537   pre-processed data are also available from https://github.com/Taiyou/OptofMRI_5HT.
538   All software used in this study is also available (Supplementary Table ST1)
539

540   Acknowledgments
541   We thank Dr. Koji Toda (Keio University) for his comments on awake optogenetic
542   stimulation, Dr. Keigo Hikishima for his comments on fMRI, Mr. Yoshiaki Ohba for
543   creation of acclimation tools, and DMD. Jiafu Zeng for animal support for MRI
544   acquisition.
545
546   Author contributions
547   HTH and KD conceived the experiments. HTH, AY, NT designed and prepared
548   experiments, HTH performed MRI experiments. MT conducted behavioral experiments.
549   HTH analysed results, and HTH wrote the manuscript. All authors reviewed the
550   manuscript.

Figure 1. (a) Surgical setup. A fixation bar was mounted on the mouse head with dental cement and an optical cannula was inserted into the dorsal raphe nucleus (DRN) (b) Acclimation protocol. A mouse was exposed daily to recorded MRI scanning sounds for 2 h for 7 days prior to fMRI sessions. (c) Location of an optical cannula. An optical cannula was inserted horizontally into the DRN from the cerebellum. A cyan window indicates the imaging window for T2* sessions. (d) Experimental schedule. After surgery and 7 days acclimation, the first fMRI recording session was performed. A replication session was also performed a few weeks later. Finally, experiments for anesthesia effects were performed. On the 1st day, an fMRI recording with general anesthesia using isoflurane 1% was performed, and an fMRI recording without anesthesia was conducted thereafte©(e) Stimulation protocol. In an fMRI session, the onset of optogenetic stimulation was applied 90 s after an imaging session began. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after blue illumination. (f) Average time course of % BOLD signals in DRN. Time

series of % BOLD signals were individually extracted from the DRN in session 1 and session 2, and the time series were averaged across transgenic mice (n = 8). Each cyan window indicates the stimulation period during a session for 20 s. (g) Alignment of average % BOLD responses in the DRN. % BOLD responses were aligned at the time of blue illumination (1.0 s) in session 1. Blue and yellow lines indicate % BOLD responses in the DRN from blue and yellow stimulation sessions from the transgenic group (n = 8), respectively. A cyan window indicates stimulation duration during a session for 20 s.

595
Figure 2. (a) Functional map of optogenetic illumination. A functional map was calculated by contrasting
BOLD responses between blue and yellow illumination in the transgenic (n = 8) group (p < 0.05, cluster-
corrected controlling family-wise e©r (FWE) with probabilistic threshold-free cluster enhancement
(pTFCE, see 'General Linear Model (GLM) for BOLD signals' in the Method section)). Magenta color
indicates the t value. (b) Comparison of beta values in multiple ROIs. Beta values were obtained from
GLM analysis in session 1, and were extracted and averaged based on ROIs from the Allen Brain Atlas
(Table 1). Differences of beta values between blue and yellow illumination sessions were contrasted in the
transgenic group. * indicates p<0.05 (multiple comparison was corrected with false discovery rate (FDR)).
Cyan and magenta colors indicate blue and yellow illumination, respectively. (c) Alignment of average %
BOLD responses in mPFc, OFC, CPu, and VTA. % BOLD responses were aligned at the time of blue
illumination (1.0 s) in session 1. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation,
and yellow illumination (1.0 second; 593 nm) followed 20 s the offset of the stimulation after blue
illumination. A cyan window indicates stimulation duration during a session for 20 s. Blue and yellow
lines indicate % BOLD responses in mPFc, OFC, CPu, and VTA from blue and yellow illumination from

transgenic groups (n = 8), respectively. A cyan window indicates stimulation duration during a session for
20 s.

641 Figure 3. (a) Optogenetic fMRI session with general anesthesia in session 3. General anesthesia, 1%
642 isoflurane, was applied to the transgenic group (n=7) during optogenetic fMRI sessions. GLM analysis
643 following optogenetic stimulation was executed and visualized (T > 1, uncorrected). Magenta and cyan

colors indicate positive and negative statistical t values. (b) An optogenetic fMRI session in the awake state in session 4. As in session 1, no anesthesia was applied to the transgenic group during optogenetic fMRI sessions. As on day 1, GLM analysis was executed and visualized (T > 1, uncorrected). (c) Comparison between beta values of sessions 3 and 4 in multiple ROIs. Beta values of blue illumination sessions in session 3 and session 4 were statistically contrasted using Student's t test. * indicates $p<0.05$, FDR-corrected. (d) Alignment of average % BOLD responses in vRSC, CA1, DG, and DRN in session 3 from transgenic mice. Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after the blue illumination. A cyan window indicates stimulation duration for 20 seconds. * indicates $p<0.05$, FDR corrected.

Figure 4. (a) Structural density map of serotonergic projection. A structural density map of serotonergic projections was obtained from the Allen Brain Atlas. (b) Correlation between projection density and regional beta values. The correlation between regional structural density and average regional beta values across transgenic groups was calculated from session 1 (blue stimulation: r = 0.462, p < 0.05; yellow stimulation: r = 0.132, p = 0.529, FDR corrected). Blue and yellow colors indicate data from blue and yellow illumination sessions, respectively. (c) Correlation between projection density and regional beta values. The correlation between regional structural density and average regional beta values across transgenic groups was calculated (Anesthetized (Session3): r = 0.546, p < 0.05; Awake (Session4): r = 0.758, p < 0.001, FDR corrected). Magenta and blue colors indicate data from blue illumination sessions in sessions 3 and 4, respectively.

Figure 5. (a) Procedure to make a model of predicted beta values in session 1 of blue illumination. Independent variables, including regional structural densities and regional gene expression of 5ht1a, 5ht1b, 5ht1f, 5ht2a, and 5ht2c were fit with a linear regression model to beta values of multiple ROIs from blue illumination sessions. (b) Correlation between modeled beta values from session 1 and beta values from session 3 under anesthesia (r = 0.365, p = 0.079, FDR corrected). (c) Correlation between modeled beta values from session 1 and beta values from session 4 (Awake: r = 0.615, p < 0.01, FDR corrected). (d) Weights of fitted models from session 1, session 2, and session 3 under anesthesia, and session 4 without

697 anesthesia. Cyan, black, magenta, and green colors indicate weights of expression profiles from session 1
698 (n=8), session2 (n=7), session 3 (n=7) and session 4 (n=7), respectively.

ROI id	Area	ROI name	Abbreviations
1	Cortical regions	Frontal polar area	FP
2		Orbital area	OFC
3		Medial prefrontal cortex	mPFC
4		Secondary motor area	M2
5		Anterior cingulate cortex	ACC
6		Primary motor area	M1
7		Supplementary somatosensory area	SS
8		Ventral retrosplenial cortex	vRSC
9		Dorsal retrosplenial cortex	dRSC
10		Temporal association area	TAA
11		Primary auditory area	A1
12		Primary somatosensory area	S1
13		Posterior parietal association area	PPA
14		Primary visual area	V1
15	Sub-cortical regions	Nucleus accumbens	NAc
16		Caudate putamen	CPu
17		Bed nucleus of steria terminalis	BST
18		Globus pallidus external part	GPe
19		Globus pallidus internal part	GPi
20		Thalamic reticular nucleus	RT
21		Ventral pallidus	VP
22		Dentate gyrus of hippocampus	DG
23		CA1 of hippocampus	CA1
24		Lateral habenula	LHb
25		Substantia nigra pars reticulata	SNr
26		Ventral tegmental area	VTA
27		Medial raphe nucleus	MRN
28		Dorsal raphe nucleus	DRN

744
745 Table1. Regions of interest (ROIs) and their abbreviations. ROIs are ordered from anterior to posterior for
746 cortical and subcortical areas.
747
748

Model Statistics (Session 1)				
	R^2 statistic	F-statistic	P value (uncorrected)	Estimate of the error variance
Session 1	0.6603	5.5063	0.0025	0.0643

749
750 Table 2. Model with Parameter Estimate of Session 1. Multiple linear regression was performed to fit
751 parameter estimates from session 1 (blue stimulation) with density of serotonergic projection and
752 expression of 5-HT receptors.
753
754

5-ht1a		SS	df	MS	F-statistic	p value
	Group	0.00111	1	0.00111	0	0.945
	Error	6.20697	27	0.22989		
	Total	6.20808	28			
5-ht1b						
	Group	1.6679	1	1.66795	5.07	0.0327
	Error	8.8794	27	0.32886		
	Total	10.5473	28			
5-ht1f						
	Group	4.6674	1	4.66744	12.95	**0.0013***
	Error	9.7293	27	0.36034		
	Total	14.4967	28			
5-ht2a						
	Group	0.40786	1	0.40786	3.43	0.075
	Error	3.21128	27	0.11894		
	Total	3.61914	28			
5-ht2c						
	Group	10.103	1	10.103	31.08	**0.0019***
	Error	8.779	27	0.3251		
	Total	18.8809	28			
Structural density						
	Group	0.18596	1	0.18596	1.73	0.1999
	Error	2.90782	27	0.1077		
	Total	3.0938	28			
Intercept						
	Group	0.1135	1	0.11348	0.26	0.6171
	Error	11.9762	27	0.44356		
	Total	12.0896	28			

755
756 Table 3. Results of one-way ANOVA for 5-HT receptor type and structural intensity. One-way ANOVA
757 was applied to regression coefficients from each factor to test whether each factor showed statistically
758 different expressions between awake state and anesthetized states. SS, df, and MS indicate Sum of
759 Squares, Degrees of freedom, and Mean Squares, respectively. * indicates statistical significance after
760 multiple correction with the Bonferroni method ($p < 0.05$).
761

References

1. Abe, Y., et al., Diffusion functional MRI reveals global brain network functional
abnormalities driven by targeted local activity in a neuropsychiatric disease
mouse model. Neuroimage, 2020. 223: p. 117318.
2. Amargos-Bosch, M., et al., Co-expression and in vivo interaction of
serotonin1A and serotonin2A receptors in pyramidal neurons of prefrontal
cortex. Cereb Cortex, 2004. 14(3): p. 281-99.
3. Anderson, I.M., et al., Assessing human 5-HT function in vivo with
pharmacoMRI. Neuropharmacology, 2008. 55(6): p. 1029-37.
4. Boschert, U., et al., The mouse 5-hydroxytryptamine1B receptor is localized
predominantly on axon terminals. Neuroscience, 1994. 58(1): p. 167-82.
5. Boureau, Y.L. and P. Dayan, Opponency revisited: competition and cooperation
between dopamine and serotonin. Neuropsychopharmacology, 2011. 36(1): p.
74-97.
6. Caraiscos, V.B., et al., Selective enhancement of tonic GABAergic inhibition in
murine hippocampal neurons by low concentrations of the volatile anesthetic
isoflurane. J Neurosci, 2004. 24(39): p. 8454-8.
7. Cornea-Hebert, V., et al., Cellular and subcellular distribution of the serotonin 5-
HT2A receptor in the central nervous system of adult rat. J Comp Neurol, 1999.
409(2): p. 187-209.
8. Correia, P.A., et al., Transient inhibition and long-term facilitation of
locomotion by phasic optogenetic activation of serotonin neurons. Elife, 2017. 6.
9. Eichenbaum, H., Prefrontal-hippocampal interactions in episodic memory. Nat
Rev Neurosci, 2017. 18(9): p. 547-558.
10. Fonseca, M.S., M. Murakami, and Z.F. Mainen, Activation of dorsal raphe
serotonergic neurons promotes waiting but is not reinforcing. Curr Biol, 2015.
25(3): p. 306-315.
11. Fritze, S., R. Spanagel, and H.R. Noori, Adaptive dynamics of the 5-HT systems
following chronic administration of selective serotonin reuptake inhibitors: a
meta-analysis. J Neurochem, 2017. 142(5): p. 747-755.
12. Fu, W., et al., Chemical neuroanatomy of the dorsal raphe nucleus and adjacent
structures of the mouse brain. J Comp Neurol, 2010. 518(17): p. 3464-94.
13. Garcia-Garcia, A.L., et al., Serotonin inputs to the dorsal BNST modulate
anxiety in a 5-HT(1A) receptor-dependent manner. Mol Psychiatry, 2018.
23(10): p. 1990-1997.
14. Giorgetti, M. and L.H. Tecott, Contributions of 5-HT(2C) receptors to multiple
actions of central serotonin systems. Eur J Pharmacol, 2004. 488(1-3): p. 1-9.
15. Giorgi, A., et al., Brain-wide Mapping of Endogenous Serotonergic
Transmission via Chemogenetic fMRI. Cell Rep, 2017. 21(4): p. 910-918.
16. Grandjean, J., et al., A brain-wide functional map of the serotonergic responses
to acute stress and fluoxetine. Nat Commun, 2019. 10(1): p. 350.
17. Harris, J.A., et al., Anatomical characterization of Cre driver mice for neural
circuit mapping and manipulation. Front Neural Circuits, 2014. 8: p. 76.

18.   Hikishima, K., et al., In vivo microscopic voxel-based morphometry with a brain
807   template to characterize strain-specific structures in the mouse brain. Sci Rep,
808   2017. 7(1): p. 85.
809   19.   Iigaya, K., et al., An effect of serotonergic stimulation on learning rates for
810   rewards apparent after long intertrial intervals. Nat Commun, 2018. 9(1): p.
811   2477.
812   20.   Izquierdo, I., C.R. Furini, and J.C. Myskiw, Fear Memory. Physiol Rev, 2016.
813   96(2): p. 695-750.
814   21.   Kawai, S., et al., Effect of three types of mixed anesthetic agents alternate to
815   ketamine in mice. Exp Anim, 2011. 60(5): p. 481-7.
816   22.   Klomp, A., et al., Lasting effects of chronic fluoxetine treatment on the late
817   developing rat brain: age-dependent changes in the serotonergic
818   neurotransmitter system assessed by pharmacological MRI. Neuroimage, 2012.
819   59(1): p. 218-26.
820   23.   Li, A., et al., Dorsal raphe serotonergic neurons promote arousal from isoflurane
821   anesthesia. CNS Neurosci Ther, 2021. 27(8): p. 941-950.
822   24.   Li, R., et al., Restoring susceptibility induced MRI signal loss in rat brain at 9.4
823   T: A step towards whole brain functional connectivity imaging. PLoS One,
824   2015. 10(4): p. e0119450.
825   25.   Li, Y., et al., Serotonin neurons in the dorsal raphe nucleus encode reward
826   signals. Nat Commun, 2016. 7: p. 10503.
827   26.   Liu, Z., et al., Dorsal raphe neurons signal reward through 5-HT and glutamate.
828   Neuron, 2014. 81(6): p. 1360-1374.
829   27.   Lottem, E., et al., Activation of serotonin neurons promotes active persistence in
830   a probabilistic foraging task. Nat Commun, 2018. 9(1): p. 1000.
831   28.   Lucaites, V.L., et al., [3H]LY334370, a novel radioligand for the 5-HT1F
832   receptor. II. Autoradiographic localization in rat, guinea pig, monkey and human
833   brain. Naunyn Schmiedebergs Arch Pharmacol, 2005. 371(3): p. 178-84.
834   29.   Malik, R., et al., Top-down control of hippocampal signal-to-noise by prefrontal
835   long-range inhibition. Cell, 2022. 185(9): p. 1602-1617 e17.
836   30.   Marcinkiewcz, C.A., et al., Serotonin engages an anxiety and fear-promoting
837   circuit in the extended amygdala. Nature, 2016. 537(7618): p. 97-101.
838   31.   Massey, C.A., et al., Isoflurane abolishes spontaneous firing of serotonin
839   neurons and masks their pH/CO(2) chemosensitivity. J Neurophysiol, 2015.
840   113(7): p. 2879-88.
841   32.   Matias, S., et al., Activity patterns of serotonin neurons underlying cognitive
842   flexibility. Elife, 2017. 6.
843   33.   Matsunaga, F., et al., Molecular interactions between general anesthetics and the
844   5HT2B receptor. J Biomol Struct Dyn, 2015. 33(1): p. 211-8.
845   34.   McDevitt, R.A., et al., Serotonergic versus nonserotonergic dorsal raphe
846   projection neurons: differential participation in reward circuitry. Cell Rep, 2014.
847   8(6): p. 1857-1869.
848   35.   Mengod, G., et al., 5-HT receptors in mammalian brain: receptor
849   autoradiography and in situ hybridization studies of new ligands and newly
850   identified receptors. Histochem J, 1996. 28(11): p. 747-58.
851   36.   Meyniel, F., et al., A specific role for serotonin in overcoming effort cost. Elife,
852   2016. 5.

853   37.   Miyazaki, K., et al., Reward probability and timing uncertainty alter the effect of
854   dorsal raphe serotonin neurons on patience. Nat Commun, 2018. 9(1): p. 2048.
855   38.   Miyazaki, K., et al., Serotonergic projections to the orbitofrontal and medial
856   prefrontal cortices differentially modulate waiting for future rewards. Sci Adv,
857   2020. 6(48).
858   39.   Miyazaki, K.W., et al., Optogenetic activation of dorsal raphe serotonin neurons
859   enhances patience for future rewards. Curr Biol, 2014. 24(17): p. 2033-40.
860   40.   Mo, B., et al., Restraint stress increases serotonin release in the central nucleus
861   of the amygdala via activation of corticotropin-releasing factor receptors. Brain
862   Res Bull, 2008. 76(5): p. 493-8.
863   41.   Nagai, Y., et al., The Role of Dorsal Raphe Serotonin Neurons in the Balance
864   between Reward and Aversion. Int J Mol Sci, 2020. 21(6).
865   42.   Nichols, D.E. and C.D. Nichols, Serotonin receptors. Chem Rev, 2008. 108(5):
866   p. 1614-41.
867   43.   Ohmura, Y., et al., Optogenetic activation of serotonergic neurons enhances
868   anxiety-like behaviour in mice. Int J Neuropsychopharmacol, 2014. 17(11): p.
869   1777-83.
870   44.   Ohmura, Y., et al., Disruption of model-based decision making by silencing of
871   serotonin neurons in the dorsal raphe nucleus. Curr Biol, 2021. 31(11): p. 2446-
872   2454 e5.
873   45.   Oikonomou, G., et al., The Serotonergic Raphe Promote Sleep in Zebrafish and
874   Mice. Neuron, 2019. 103(4): p. 686-701 e8.
875   46.   Puig, M.V., et al., In vivo excitation of GABA interneurons in the medial
876   prefrontal cortex through 5-HT3 receptors. Cereb Cortex, 2004. 14(12): p. 1365-
877   75.
878   47.   Puig, M.V. and A.T. Gulledge, Serotonin and prefrontal cortex function:
879   neurons, networks, and circuits. Mol Neurobiol, 2011. 44(3): p. 449-64.
880   48.   Pytliak, M., et al., Serotonin receptors - from molecular biology to clinical
881   applications. Physiol Res, 2011. 60(1): p. 15-25.
882   49.   Ren, J., et al., Anatomically Defined and Functionally Distinct Dorsal Raphe
883   Serotonin Sub-systems. Cell, 2018. 175(2): p. 472-487 e20.
884   50.   Ren, J., et al., Single-cell transcriptomes and whole-brain projections of
885   serotonin neurons in the mouse dorsal and median raphe nuclei. Elife, 2019. 8.
886   51.   Riad, M., et al., Somatodendritic localization of 5-HT1A and preterminal axonal
887   localization of 5-HT1B serotonin receptors in adult rat brain. J Comp Neurol,
888   2000. 417(2): p. 181-94.
889   52.   Schaefer, A., et al., Serotonergic modulation of intrinsic functional connectivity.
890   Curr Biol, 2014. 24(19): p. 2314-8.
891   53.   Scholl, J., et al., Beyond negative valence: 2-week administration of a
892   serotonergic antidepressant enhances both reward and effort learning signals.
893   PLoS Biol, 2017. 15(2): p. e2000756.
894   54.   Sesack, S.R. and A.A. Grace, Cortico-Basal Ganglia reward network:
895   microcircuitry. Neuropsychopharmacology, 2010. 35(1): p. 27-47.
896   55.   Smith, H.R., et al., Dorsal Raphe Serotonin Neurons Mediate CO_2-Induced
897   Arousal from Sleep. J Neurosci, 2018. 38(8): p. 1915-1925.

56. Spisak, T., et al., Probabilistic TFCE: A generalized combination of cluster size
and voxel intensity to increase statistical power. Neuroimage, 2019. 185: p. 12-
26.
57. Tanaka, K.F., et al., Expanding the repertoire of optogenetically targeted cells
with an enhanced gene expression system. Cell Rep, 2012. 2(2): p. 397-406.
58. Tanaka, S.C., et al., Serotonin differentially regulates short- and long-term
prediction of rewards in the ventral and dorsal striatum. PLoS One, 2007. 2(12):
p. e1333.
59. Trusel, M., et al., Punishment-Predictive Cues Guide Avoidance through
Potentiation of Hypothalamus-to-Habenula Synapses. Neuron, 2019. 102(1): p.
120-127 e4.
60. Wang, H.Y., et al., Frequency-Dependent Block of Excitatory
Neurotransmission by Isoflurane via Dual Presynaptic Mechanisms. J Neurosci,
2020. 40(21): p. 4103-4115.
61. Whittington, R.A. and L. Virag, Isoflurane decreases extracellular serotonin in
the mouse hippocampus. Anesth Analg, 2006. 103(1): p. 92-8, table of contents.
62. Winegar, B.D. and M.B. MacIver, Isoflurane depresses hippocampal CA1
glutamate nerve terminals without inhibiting fiber volleys. BMC Neurosci,
2006. 7: p. 5.
63. Xu, S., et al., Dorsal Raphe Serotonergic Neurons Control Intertemporal Choice
under Trade-off. Curr Biol, 2017. 27(20): p. 3111-3119 e3.
64. Ying, S.W., et al., Isoflurane modulates excitability in the mouse thalamus via
GABA-dependent and GABA-independent mechanisms. Neuropharmacology,
2009. 56(2): p. 438-47.
65. Yokoyama, C., et al., Marmoset Serotonin 5-HT1A Receptor Mapping with a
Biased Agonist PET Probe 18F-F13714: Comparison with an Antagonist Tracer
18F-MPPF in Awake and Anesthetized States. Int J Neuropsychopharmacol,
2016. 19(12).
66. Yoshida, K., et al., Physiological effects of a habituation procedure for
functional MRI in awake mice using a cryogenic radiofrequency probe. J
Neurosci Methods, 2016. 274: p. 38-48.

Supplementary materials

Optogenetic activation of dorsal raphe serotonin neurons induces brain-wide activation, including reward-related circuits

Hiro Taiyo Hamada[1,3,*], Yoshifumi Abe[2], Norio Takata[2], Maaskazu Taira[1], Kenji F. Tanaka[2], Kenji Doya[1,*]

[1] Neural Computation Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

[2] Department of Neuropsychiatry, Keio University School of Medicine, Tokyo, Japan

[3] Research & Development Department, Araya Inc., Tokyo, Japan

* Correspondence to hamada_h@araya.org, doya@oist.jp

Software List				
IDs	Name	Contributor	Purpose	URLs
1	Statistical Parametric Mapping 12(SPM12)	The Wellcome Centre for Human Neuroimaging, UCL	MRI data analysis	https://www.fil.ion.ucl.ac.uk/spm/
2	MarsBaR region of interest toolbox for SPM		Brain Reference Registration	http://marsbar.sourceforge.net/
3	MRIcroGL	Chris Rorden, Neuropsychology Lab, University of South Carolina	3D visualization of MRI data	https://www.nitrc.org/ projects/mricrogl/
4	Probabilistic Threshold-free Cluster Enhancement (pTFCE) [3]	Tamás Spisák	A cluster-enhancement method	https://spisakt.github.io/pTFCE/
5	Allen Brain Atlas	Allen Brain Institutes	Reference atlas, 5HT projection map, 5HT receptor gene expressions	https://portal.brain-map.org/
6	Templates for In Vivo Mouse Brain [1]	Keigo Hikishima (k-hikishima@ aist.go.jp)	Mouse brain templates for registration and normalization	https:// www.nitrc.org/ projects/tpm_mouse
7	Gramm [2]	Pierre Morel	Visualization of Violin Plotting	https://github.com/piermorel/gramm
8	in-house analysis programs	Neural Computational Unit	Analysis for time series, Behavioral analysis	https://github.com/Taiyou/OptofMRI_5HT
9	MATLAB	MathWorks	Main analyses and data visualization	https://mathworks.com/

Supplementary Table S1. Software employed in this study

ROI id	ROI name	Allen Brain Atlas Name
1	Frontal polar area (FP)	Forntaol pole, layer 1, 2/3, 5, 6a, 6b
2	Orbital area (OFC)	Orbital area, lateral/medial/ventral/ventrolateral part, layer 1, 2/3, 5, 6a, 6b
3	Medial prefrontal cortex (mPFC)	Prelimbic area, layer 1, 2, 2/3, 5, 6a, 6b & Infralimbic area, layer 1, 2, 2/3, 5, 6a, 6b
4	Secondary motor area (M2)	Secondary Motor area, layer 1, 2/3, 5, 6a, 6b
5	Anterior cingulate cortex (ACC)	Anterior cingulate area, layer 1, 2/3, 5, 6a, 6b & Anterior cingulate area, dorsal/ventral part, layer 1, 2/3, 5, 6a, 6b
6	Primary motor area (M1)	Primary motor area, layer 1, 2/3, 5, 6a, 6b
7	Supplementary somatosensory area (SS)	Supplemental somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b
8	Ventral retrosplenial cortex (vRSC)	Retrosplenial area, ventral part, layer 1, 2/3, 4, 5, 6a, 6b
9	Dorsal retrosplenial cortex (dRSC)	Retrosplenial area, dorsal part, layer 1, 2/3, 4, 5, 6a, 6b
10	Temporal association area (TAA)	Temporal association areas, layer 1, 2/3, 4, 5, 6a, 6b
11	Primary auditory area (A1)	Primary auditory are, layer 1, 2/3, 4, 5, 6a, 6b
12	Primary somatosensory area (S1)	Somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b, Primary somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b, Primary somatosensory area, nose/barrel field, layer 1, 2/3, 4, 5, 6a, 6b
13	Posterior parietal association area (PPA)	Posteior parietal association areas, layer 1, 2/3, 4, 5, 6a, 6b
14	Primary visual area (V1)	Primary visual area, layer 1, 2/3, 4, 5, 6a, 6b
15	Nucleus accumbens (NAc)	Nucleus accumbens
16	Caudate putamen (CPu)	Caudoputamen
17	Bed nucleus of steria terminalis (BST)	Bed nuclei of the stria terminalis, anterior/posterior division
18	Globus pallidus external part (GPe)	Globus pallidus, external segment
19	Globus pallidus internal part (GPi)	Globus pallidus, internal segment
20	Thalamic reticular nucleus (RT)	Reticular nucleus of the thalamus
21	Ventral pallidus (VP)	Pallidum, ventral region
22	Dentate gyrus of hippocampus (DG)	Dentate gyrus, molecular/polymorph/granule cell layer/subgranular zone, Dentate gyrus crest, molecular/polymorph/granule cell layer, Dentate gyrus lateral/medial blade, molecular/polymorph/granucle cell layer
23	CA1 of hippocampus (CA1)	Field CA1, stratum lacunosum-moleculare/stratum oriens/pyramidal layer/stratum radiatum
24	Lateral habenula (LHb)	Lateral habenula
25	Substantia nigra pars reticulata (SNr)	Substantia nigra, reticular part
26	Ventral tegmental area (VTA)	Ventral tegmental area
27	Medial raphe nucleus (MRN)	Superior central nucleus raphe, lateral/medial part
28	Dorsal raphe nucleus (DRN)	Dorsal nucleus raphe

1033
1034 Supplementary Table ST2. A list of correspondence between ROIs and names in Allen Brain Atlas
1035
1036
1037

	Session 1	Session 2	Session 3	Session 4
R^2 statistics	0.5851 ± 0.1574	0.5505 ± 0.0863	0.5240 ± 0.1166	0.5437 ± 0.1100
F statistics	5.1849 ± 3.9732	3.6719 ± 1.2474	3.4689 ± 1.6842	3.7058 ± 1.6093
P value	0.0506 ± 0.0774	0.0352 ± 0.0464	0.0599 ± 0.0609	0.0468 ± 0.0570
EEV	0.1252 ± 0.0644	0.1009 ± 0.0224	0.1316 ± 0.0994	0.0630 ± 0.0259

Supplementary Table ST 3. Summary of multiple linear regression with beta values from session1-4. Multiple linear regression was applied to beta values from each subject for all sessions. Each value indicates mean ± standard error of mean.

1043
Supplementary Figure S1. (a) Sequence of the tone-food waiting task. (b) Time sequence of optogenetic
illumination. In serotonin activation trials, blue light stimulation was delivered until a trial ended with
delivery of a food pellet or quitting. In serotonin no—activation trials, yellow light stimulation was
delivered until the trial ended. After the trial end, 1 s of yellow light was delivered. Blue and yellow color
indicate blue and yellow light stimulation. Red regions indicate reward delay periods. (c) Experiment 1:
Fixed time waiting (6.0 s). In this experiment, the reward delay duration for a success trial was fixed at 6
s. For an omission trial, no food reward was given. ** $p < 0.01$ by Wilcoxon rank sum test with
Bonferroni correction. Blue and orange regions indicate distributions of blue light and yellow light
stimulation. (d) Experiment 2: Random time waiting (2.0, 6.0, or 10.0 s). In this experiment, the reward
delay duration for a successful trial was randomly delivered from 2, 6, and 10 s. For an omission trial, no
food reward was given to mice. ** $p < 0.01$ by Wilcoxon rank sum test with Bonferroni correction. Blue
and orange regions indicate distributions of blue light and yellow light stimulation.

1062
1063

1064
1065 Supplementary Figure S2. (a) Trial-wise functional maps of transgenic mice with blue illumination in
1066 session 1 ($p < 0.05$, uncorrected, Transgenic: n = 8). The numbers 1-5 correspond to timing of each
1067 stimulus. (b) Trial-wise functional maps of transgenic mice with yellow illumination in session 1 ($p <$
1068 0.05, uncorrected, Transgenic: n = 8).
1069
1070

Supplementary Figure S3. (a) Functional map of transgenic mice with blue illumination in session 1 ($p <$ 0.05, uncorrected, Transgenic: n = 8). (b) Functional map of transgenic mice with blue illumination in replication session ($p < 0.05$, uncorrected, Transgenic: n = 7).

Supplementary Figure S4. Comparison of beta values in multiple ROIs with the WT group (n = 6). Beta values were obtained from GLM analysis in session 1, and were extracted and averaged based on ROIs from the Allen Brain Atlas (Table1.). The difference of beta values between blue and yellow illumination sessions were contrasted in the WT group. Cyan and yellow colors indicate blue and yellow illumination trials, respectively.

Supplementary Figure S5. Average % BOLD responses to blue illumination in the session 1. Time series
of % BOLD responses to yellow illumination in each mouse were averaged at the onset of illumination.
Average % BOLD responses were contrasted between the transgenic (n = 8) and WT groups (n = 6). Blue
illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow illumination (1.0 second;
593 nm) followed the offset of stimulation 20 s after blue illumination. Cyan highlighting indicates the
stimulation duration for 20 s. * and ** indicate statistical p values under 0.05 and 0.01, respectively
(corrected with FDR).

cortex

subcortical regions

Yellow stimulation
- Transgenic (awake, n=8)   *: p < 0.05
- WT (awake, n=6)

Supplementary Figure S6. Average % BOLD responses of yellow illumination in session 1. Time series of % BOLD responses to yellow illumination in each mouse were averaged at the onset of illumination. Average % BOLD responses were contrasted between the transgenic (n = 8) and WT groups (n = 6). Yellow highlighting indicates the stimulus duration for 20 s. * indicate statistical p values under 0.05 (corrected with FDR).

Supplementary Figure S7. (a) Replication of average beta values from GLM to blue stimulation between sessions 1 and 2 in transgenic mice. Statistically significant correlation was found in blue photo-stimulation (r = 0.875, p <1e-8), but not in yellow photo-stimulation (r = 0.449, p = 0.0688). (b) Relationship of average regional beta values from GLM to blue photo-stimulation between session 2 and sessions 3 and 4 in transgenic mice. Statistically significant correlations were found in session 3 under general anesthetics (r = 0.552, p < 0.05) and in session 4 without general anesthesia (r = 0.900, p < 1e^{-8}).

1146
Supplementary Figure S8. (a) Replication of peaks of average % BOLD signals between sessions 1 and 2
in transgenic mice. Statistically significant correlation between peaks of % BOLD responses to blue
photo-stimulation in sessions 1 and 2 was found, but not to yellow photo-stimulation (blue illumination: r
= 0.8863, p < 1e-8; yellow photo stimulation: r = -0.06, p = 0.766). (b) Replication of peaks of average %
BOLD signals between sessions 1 and 2 in transgenic mice. Statistically significant correlation between
peak timings of % BOLD responses to blue photo-stimulation in sessions 1 and 2 was found but not to
yellow photo-stimulation (blue illumination: r = 0.6785, p < 0.005; yellow photo stimulation: r = 0.0369,
p = 0.8551). (c) Alignment of peaks of regional % BOLD signals in sessions 1 and 2. Peaks of % BOLD

signals are sorted in ascending order. (d) Alignment of peak timing of regional % BOLD signals in sessions 1 and 2. Peak timings of regional % BOLD signals are sorted in ascending order.

Supplementary Figure S9. Association between peak timing and peaks of average %BOLD signals in sessions 1 and 2. No statistically significant correlations between peak timing and peaks of photo-stimulated %BOLD signals were found in either session (session 1: r = 0.092, p = 0.639; session 2: r = - 0.047, p = 0.811).

Supplementary Figure S10. (a) Functional map of transgenic mice (n = 7) with blue illumination in session 3 under anesthesia (t > 1, uncorrected). (b) Functional map of transgenic mice (n = 7) with blue illumination in session 4 without anesthesia (t > 1, uncorrected).

1215
1216
Supplementary Figure S11. Average % BOLD responses to blue illumination in sessions 3 and 4. Time
series of % BOLD responses by blue illumination in each mouse were averaged at the onset of
illumination. Average % BOLD responses were contrasted between sessions 3 and 4 in the transgenic
group (n = 7). Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, yellow
illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after blue illumination. Cyan
highlighting indicates the stimulus duration for 20 s. * indicate statistical p values under 0.05 (corrected
with FDR).

Supplementary Figure S12. Comparison of functional map, structural density, and gene expression maps. The functional map from session 1 under blue illumination was aligned with structural density and gene expression maps. The structural density map and gene expression of serotonin receptors are obtained from the mouse brain database of the Allen Brain Institute (Experimental IDs: 480074702 (5HT projection), 79556616 (5HT1a), 583 (5HT1b), 69859867 (5HT1f), 81671344 (5HT2a), and 73636098 (5HT2c)).

1231
1232
1233 Supplementary Figure S13. Correlation between 5HT projection density and expression profiles of
1234 serotonin receptors. (a) Pearson's correlation 5HT projection density and expression profiles of serotonin
1235 receptors. (b) Corresponding statistical p values of the correlation between 5HT projection density and
1236 expression profiles of serotonin receptors ($p < 0.05$, uncorrected).

References

1. Hikishima, K., et al., In vivo microscopic voxel-based morphometry with a brain template to characterize strain-specific structures in the mouse brain. Sci Rep, 2017. 7(1): p. 85.
2. Morel, P., Gramm: grammar of graphics plotting in Matlab. Journal of Open Source Software, 2018. 3(23): p. 568.
3. Spisak, T., et al., Probabilistic TFCE: A generalized combination of cluster size and voxel intensity to increase statistical power. Neuroimage, 2019. 185: p. 12-26.

Optogenetic activation of dorsal raphe serotonin neurons induces brain-wide activation, including reward-related circuits

Hiro Taiyo Hamada[1,3,*], Yoshifumi Abe[2], Norio Takata[2], Maaskazu Taira[1], Kenji F. Tanaka[2], Kenji Doya[1,*]

[1] Neural Computation Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan
[2] Department of Neuropsychiatry, Keio University School of Medicine, Tokyo, Japan
[3] Research & Development Department, Araya Inc., Tokyo, Japan

* Correspondence to hamada_h@araya.org, doya@oist.jp

Software List				
IDs	Name	Contributor	Purpose	URLs
1	Statistical Parametric Mapping 12(SPM12)	The Wellcome Centre for Human Neuroimaging, UCL	MRI data analysis	https://www.fil.ion.ucl.ac.uk/spm/
2	MarsBaR region of interest toolbox for SPM		Brain Reference Registration	http://marsbar.sourceforge.net/
3	MRIcroGL	Chris Rorden, Neuropsychology Lab, University of South Carolina	3D visualization of MRI data	https://www.nitrc.org/projects/mricrogl/
4	Probabilistic Threshold-free Cluster Enhancement (pTFCE) [3]	Tamás Spisák	A cluster-enhancement method	https://spisakt.github.io/pTFCE/
5	Allen Brain Atlas	Allen Brain Institutes	Reference atlas, 5HT projection map, 5HT receptor gene expressions	https://portal.brain-map.org/
6	Templates for In Vivo Mouse Brain [1]	Keigo Hikishima (k-hikishima@aist.go.jp)	Mouse brain templates for registration and normalization	https:// www.nitrc.org/projects/tpm_mouse
7	Gramm [2]	Pierre Morel	Visualization of Violin Plotting	https://github.com/piermorel/gramm
8	in-house analysis programs	Neural Computational Unit	Analysis for time series, Behavioral analysis	https://github.com/Taiyou/OptofMRI_5HT
9	MATLAB	MathWorks	Main analyses and data visualization	https://mathworks.com/

Supplementary Table S1. Software employed in this study

ROI id	ROI name	Allen Brain Atlas Name
1	Frontal polar area (FP)	Forntaol pole, layer 1, 2/3, 5, 6a, 6b
2	Orbital area (OFC)	Orbital area, lateral/medial/ventral/ventrolateral part, layer 1, 2/3, 5, 6a, 6b
3	Medial prefrontal cortex (mPFC)	Prelimbic area, layer 1, 2, 2/3, 5, 6a, 6b & Infralimbic area, layer 1, 2, 2/3, 5, 6a, 6b
4	Secondary motor area (M2)	Secondary Motor area, layer 1, 2/3, 5, 6a, 6b
5	Anterior cingulate cortex (ACC)	Anterior cingulate area, layer 1, 2/3, 5, 6a, 6b & Anterior cingulate area, dorsal/ventral part, layer 1, 2/3, 5, 6a, 6b
6	Primary motor area (M1)	Primary motor area, layer 1, 2/3, 5, 6a, 6b
7	Supplementary somatosensory area (SS)	Supplemental somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b
8	Ventral retrosplenial cortex (vRSC)	Retrosplenial area, ventral part, layer 1, 2/3, 4, 5, 6a, 6b
9	Dorsal retrosplenial cortex (dRSC)	Retrosplenial area, dorsal part, layer 1, 2/3, 4, 5, 6a, 6b
10	Temporal association area (TAA)	Temporal association areas, layer 1, 2/3, 4, 5, 6a, 6b
11	Primary auditory area (A1)	Primary auditory are, layer 1, 2/3, 4, 5, 6a, 6b
12	Primary somatosensory area (S1)	Somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b, Primary somatosensory area, layer 1, 2/3, 4, 5, 6a, 6b, Primary somatosensory area, nose/barrel field, layer 1, 2/3, 4, 5, 6a, 6b
13	Posterior parietal association area (PPA)	Posteior parietal association areas, layer 1, 2/3, 4, 5, 6a, 6b
14	Primary visual area (V1)	Primary visual area, layer 1, 2/3, 4, 5, 6a, 6b
15	Nucleus accumbens (NAc)	Nucleus accumbens
16	Caudate putamen (CPu)	Caudoputamen
17	Bed nucleus of steria terminalis (BST)	Bed nuclei of the stria terminalis, anterior/posterior division
18	Globus pallidus external part (GPe)	Globus pallidus, external segment
19	Globus pallidus internal part (GPi)	Globus pallidus, internal segment
20	Thalamic reticular nucleus (RT)	Reticular nucleus of the thalamus
21	Ventral pallidus (VP)	Pallidum, ventral region
22	Dentate gyrus of hippocampus (DG)	Dentate gyrus, molecular/polymorph/granule cell layer/subgranular zone, Dentate gyrus crest, molecular/polymorph/granule cell layer, Dentate gyrus lateral/medial blade, molecular/polymorph/granucle cell layer
23	CA1 of hippocampus (CA1)	Field CA1, stratum lacunosum-moleculare/stratum oriens/pyramidal layer/stratum radiatum
24	Lateral habenula (LHb)	Lateral habenula
25	Substantia nigra pars reticulata (SNr)	Substantia nigra, reticular part
26	Ventral tegmental area (VTA)	Ventral tegmental area
27	Medial raphe nucleus (MRN)	Superior central nucleus raphe, lateral/medial part
28	Dorsal raphe nucleus (DRN)	Dorsal nucleus raphe

Supplementary Table ST2. A list of correspondence between ROIs and names in Allen Brain Atlas

	Session 1	Session 2	Session 3	Session 4
R^2 statistics	0.5851 ± 0.1574	0.5505 ± 0.0863	0.5240 ± 0.1166	0.5437 ± 0.1100
F statistics	5.1849 ± 3.9732	3.6719 ± 1.2474	3.4689 ± 1.6842	3.7058 ± 1.6093
P value	0.0506 ± 0.0774	0.0352 ± 0.0464	0.0599 ± 0.0609	0.0468 ± 0.0570
EEV	0.1252 ± 0.0644	0.1009 ± 0.0224	0.1316 ± 0.0994	0.0630 ± 0.0259

Supplementary Table ST 3. Summary of multiple linear regression with beta values from session1-4. Multiple linear regression was applied to beta values from each subject for all sessions. Each value indicates mean ± standard error of mean.

Supplementary Figure S1. (a) Sequence of the tone-food waiting task. (b) Time sequence of optogenetic illumination. In serotonin activation trials, blue light stimulation was delivered until a trial ended with delivery of a food pellet or quitting. In serotonin no—activation trials, yellow light stimulation was delivered until the trial ended. After the trial end, 1 s of yellow light was delivered. Blue and yellow color indicate blue and yellow light stimulation. Red regions indicate reward delay periods. (c) Experiment 1: Fixed time waiting (6.0 s). In this experiment, the reward delay duration for a success trial was fixed at 6 s. For an omission trial, no food reward was given. ** p < 0.01 by Wilcoxon rank sum test with Bonferroni correction. Blue and orange regions indicate distributions of blue light and yellow light stimulation. (d) Experiment 2: Random time waiting (2.0, 6.0, or 10.0 s). In this experiment, the reward delay duration for a successful trial was randomly delivered from 2, 6, and 10 s. For an omission trial, no food reward was given to mice. ** p < 0.01 by Wilcoxon rank sum test with Bonferroni correction. Blue and orange regions indicate distributions of blue light and yellow light stimulation.

Supplementary Figure S2. (a) Trial-wise functional maps of transgenic mice with blue illumination in session 1 ($p < 0.05$, uncorrected, Transgenic: n = 8). The numbers 1-5 correspond to timing of each stimulus. (b) Trial-wise functional maps of transgenic mice with yellow illumination in session 1 ($p < 0.05$, uncorrected, Transgenic: n = 8).

127
128 Supplementary Figure S3. (a) Functional map of transgenic mice with blue illumination in session 1 ($p <$
129 0.05, uncorrected, Transgenic: n = 8). (b) Functional map of transgenic mice with blue illumination in
130 replication session ($p < 0.05$, uncorrected, Transgenic: n = 7).

141
142 Supplementary Figure S4. Comparison of beta values in multiple ROIs with the WT group (n = 6). Beta
143 values were obtained from GLM analysis in session 1, and were extracted and averaged based on ROIs
144 from the Allen Brain Atlas (Table1.). The difference of beta values between blue and yellow illumination
145 sessions were contrasted in the WT group. Cyan and yellow colors indicate blue and yellow illumination
146 trials, respectively.

cortex

subcortical regions

Blue illumination
- Transgenic (awake, n=8)   *: p < 0.05
- WT (awake, n=6)   **: p < 0.01

Supplementary Figure S5. Average % BOLD responses to blue illumination in the session 1. Time series
of % BOLD responses to yellow illumination in each mouse were averaged at the onset of illumination.
Average % BOLD responses were contrasted between the transgenic (n = 8) and WT groups (n = 6). Blue
illumination (1.0 s; 473 nm) was applied at the onset of stimulation, and yellow illumination (1.0 second;
593 nm) followed the offset of stimulation 20 s after blue illumination. Cyan highlighting indicates the
stimulation duration for 20 s. * and ** indicate statistical p values under 0.05 and 0.01, respectively
(corrected with FDR).

168
169
Supplementary Figure S6. Average % BOLD responses of yellow illumination in session 1. Time series
of % BOLD responses to yellow illumination in each mouse were averaged at the onset of illumination.
Average % BOLD responses were contrasted between the transgenic (n = 8) and WT groups (n = 6).
Yellow highlighting indicates the stimulus duration for 20 s. * indicate statistical p values under 0.05
(corrected with FDR).

Supplementary Figure S7. (a) Replication of average beta values from GLM to blue stimulation between
sessions 1 and 2 in transgenic mice. Statistically significant correlation was found in blue photo-
stimulation (r = 0.875, p <1e-8), but not in yellow photo-stimulation (r = 0.449, p = 0.0688). (b)
Relationship of average regional beta values from GLM to blue photo-stimulation between session 2 and
sessions 3 and 4 in transgenic mice. Statistically significant correlations were found in session 3 under
general anesthetics (r = 0.552, p < 0.05) and in session 4 without general anesthesia (r = 0.900, p < 1e^{-8}).

202
Supplementary Figure S8. (a) Replication of peaks of average % BOLD signals between sessions 1 and 2
in transgenic mice. Statistically significant correlation between peaks of % BOLD responses to blue
photo-stimulation in sessions 1 and 2 was found, but not to yellow photo-stimulation (blue illumination: r
= 0.8863, p < 1e-8; yellow photo stimulation: r = -0.06, p = 0.766). (b) Replication of peaks of average %
BOLD signals between sessions 1 and 2 in transgenic mice. Statistically significant correlation between
peak timings of % BOLD responses to blue photo-stimulation in sessions 1 and 2 was found but not to
yellow photo-stimulation (blue illumination: r = 0.6785, p < 0.005; yellow photo stimulation: r = 0.0369,
p = 0.8551). (c) Alignment of peaks of regional % BOLD signals in sessions 1 and 2. Peaks of % BOLD

signals are sorted in ascending order. (d) Alignment of peak timing of regional % BOLD signals in
sessions 1 and 2. Peak timings of regional % BOLD signals are sorted in ascending order.

Supplementary Figure S9. Association between peak timing and peaks of average %BOLD signals in sessions 1 and 2. No statistically significant correlations between peak timing and peaks of photo-stimulated %BOLD signals were found in either session (session 1: r = 0.092, p = 0.639; session 2: r = -0.047, p = 0.811).

263
264 Supplementary Figure S10. (a) Functional map of transgenic mice (n = 7) with blue illumination in
265 session 3 under anesthesia (t > 1, uncorrected). (b) Functional map of transgenic mice (n = 7) with blue
266 illumination in session 4 without anesthesia (t > 1, uncorrected).
267
268
269
270

Supplementary Figure S11. Average % BOLD responses to blue illumination in sessions 3 and 4. Time
series of % BOLD responses by blue illumination in each mouse were averaged at the onset of
illumination. Average % BOLD responses were contrasted between sessions 3 and 4 in the transgenic
group (n = 7). Blue illumination (1.0 s; 473 nm) was applied at the onset of stimulation, yellow
illumination (1.0 second; 593 nm) followed the offset of stimulation 20 s after blue illumination. Cyan
highlighting indicates the stimulus duration for 20 s. * indicate statistical p values under 0.05 (corrected
with FDR).

Supplementary Figure S12. Comparison of functional map, structural density, and gene expression maps. The functional map from session 1 under blue illumination was aligned with structural density and gene expression maps. The structural density map and gene expression of serotonin receptors are obtained from the mouse brain database of the Allen Brain Institute (Experimental IDs: 480074702 (5HT projection), 79556616 (5HT1a), 583 (5HT1b), 69859867 (5HT1f), 81671344 (5HT2a), and 73636098 (5HT2c)).

287
288
289 Supplementary Figure S13. Correlation between 5HT projection density and expression profiles of
290 serotonin receptors. (a) Pearson's correlation 5HT projection density and expression profiles of serotonin
291 receptors. (b) Corresponding statistical p values of the correlation between 5HT projection density and
292 expression profiles of serotonin receptors ($p < 0.05$, uncorrected).

References

1. Hikishima, K., et al., In vivo microscopic voxel-based morphometry with a brain template to characterize strain-specific structures in the mouse brain. Sci Rep, 2017. 7(1): p. 85.
2. Morel, P., Gramm: grammar of graphics plotting in Matlab. Journal of Open Source Software, 2018. 3(23): p. 568.
3. Spisak, T., et al., Probabilistic TFCE: A generalized combination of cluster size and voxel intensity to increase statistical power. Neuroimage, 2019. 185: p. 12-26.