Macrophage differentiation is marked by increased abundance of the mRNA 3' end processing machinery, altered poly(A) site usage, and sensitivity to the level of CstF64

Srimoyee Mukherjee[1], Joel H. Graber[2], Claire L. Moore[1*]

[1]Department of Developmental, Molecular, and Chemical Biology, Tufts University School of Medicine, Boston, Massachusetts 02111, USA

[2]Mount Desert Island Biological Laboratory, Bar Harbor, Maine, 04609, USA

*Corresponding author and lead contact: Claire L. Moore

Correspondence: claire.moore@tufts.edu

Running title: Role of alternative polyadenylation in monocyte: macrophage differentiation

Key words: alternative polyadenylation, monocyte, macrophage, CstF64, CSTF2, differentiation.

ABBREVIATIONS

CstF: Cleavage Stimulation Factor; CPSF: Cleavage and Polyadenylation Specificity Factor; C/P: Cleavage and Polyadenylation; KD: Knockdown; OE: Overexpression; UTR: Untranslated Region.

ABSTRACT

Alternative polyadenylation (APA) results in mRNA isoforms that vary in the amount of
coding sequence or 3' UTR regulatory elements. APA is important for response to
external signals and differentiation in several cell types, but its role in differentiation of
monocytes to macrophages has not been investigated. Macrophages are key effectors
of the innate immune system that help control infection and promote tissue-repair.
However, overactivity of macrophages contributes to pathogenesis of many diseases. In
this study, we show that macrophage differentiation is characterized by shortening and
lengthening of mRNAs in relevant cellular pathways. The cleavage/polyadenylation
(C/P) proteins increase during differentiation, suggesting a possible mechanism for APA
changes. This was surprising since higher C/P protein levels correlate with higher
proliferation rates in other systems, but monocytes stop dividing after induction of
differentiation. Depletion of CstF64, a C/P protein and known APA regulator, delayed
macrophage marker expression, cell cycle exit, attachment, and acquisition of structural
complexity, and impeded shortening of mRNAs with functions relevant to macrophage
biology. Conversely, CstF64 overexpression increased use of promoter-proximal
poly(A) sites and caused the appearance of differentiated phenotypes in the absence of
induction. Our findings indicate that regulation of polyadenylation plays an important
role in macrophage differentiation.
20

INTRODUCTION

Macrophages are a critical component of the innate immune system that have a protective and beneficial role through the phagocytosis and killing of microbes, removal of dying cells and cellular debris, production of immune-regulatory molecules, and presentation of antigens to T-cells (1). Macrophages are derived from precursor cells called monocytes, which, in turn, are derived from myeloid progenitors. Monocytes undergo phenotypical and functional differentiation when they are recruited to tissues, where they differentiate into macrophages with altered morphological features and then engage in inflammatory or tissue remodeling processes (2). Robust macrophage function is critical for tissue repair and combating infection. However, inappropriate or overly robust activity contributes to the pathogenesis of diseases such as atherosclerosis, arthritis, asthma, lung fibrosis, type 2 diabetes, insulin resistance, inflammatory bowel disease, cancer and most recently, COVID-19 and the formation of intra-abdominal scar tissue (3-9).

Abnormal proliferation of macrophage progenitor cells in lieu of differentiation not only impairs normal macrophage responsibilities but can also lead to blood cancers such as acute myeloid leukemias (AML). A promising approach to treat AML is differentiation therapy, in which malignant cells are encouraged to differentiate into more mature forms using pharmacological agents. This approach has been remarkably successful for a small subset of AML cases involving a PML-retinoic acid receptor fusion (10,11), and is a valuable alternative approach to killing cancer cells through cytotoxicity. To improve treatment of leukemias by forcing differentiation as well as to modulate macrophage function to help abate diseases influenced by macrophages, we must understand the critical molecular pathways that govern macrophage differentiation.

In many cell types, differentiation requires not only changes in transcription, but also pervasive changes in mRNA splicing and 3' end cleavage and polyadenylation (C/P). In the C/P step, pre-mRNA is cleaved at the poly(A) site and a poly(A) tail is added to the new end (12). Alternative polyadenylation (APA) plays an increasingly appreciated role in regulation of gene expression by affecting protein output at many levels. Genes with multiple poly(A) sites can produce distinct mRNA isoforms with different lengths of 3' untranslated regions (3' UTRs), which in turn affects nuclear export, translation, stability, association with stress granules, and localization in the cytoplasm (13-16). Alternate sequences in the 3' UTR can even facilitate co-translational protein complex formation (17) or act as a sink to bind miRNAs that down-regulate tumor suppressor mRNAs (18). Polyadenylation that occurs in an intron or less frequently an exon upstream of the 3' most exon affects the coding region, leading to expression of different protein isoforms (19,20).

Genome-wide high-throughput studies have demonstrated that the APA changes accompanying transitions in cell state are complex and cannot always be predicted based simply on proliferation status. Proliferative and tumor cells have been reported to produce mRNAs with shorter 3' UTRs, thus escaping regulation by miRNAs and RNA binding proteins (21,22). With terminally differentiated cells, the opposite can occur, with a preference for 3' UTR lengthening, as seen during the differentiation of myoblast cells

to myotubes (23) or of embryonic stem cells into neurons (24). Conversely, if differentiation is reversed, and induced pluripotent stem cells are generated from fully differentiated somatic cells, a general 3' UTR shortening has been observed (25). However, in contrast to this generalization, spermatogenesis and differentiation of secretory cells leads to a prevalence of shorter mRNA isoforms (26,27), and when B cells are transformed with Epstein-Barr virus, there is only a slight preference for shortening (28).

Many examples of APA have been documented in the immune system. For example, stimulation of human monocytes with lipopolysaccharide (LPS) and interferon-γ or activation of B cells and T cells leads to proliferation and a switch to upstream poly(A) sites (21,26). A specific example of the importance of APA in immune cell differentiation is the switch to the upstream poly(A) site of the transcription factor NFATC1, which leads to NFATC1 accumulation and maturation of naïve T cells into effector T cells (29). Preferential utilization of the upstream poly(A) site of the immunoglobulin heavy chain gene in activated B cells is needed to switch to producing the secreted form of the protein instead of the membrane-attached form and is mediated by the CstF64 C/P protein (30-32). Increased levels of CstF64 and changes in APA have also been observed after treatment of RAW 264.7 mouse monocyte/macrophage-like cells with LPS (33), which activates these cells to produce an inflammatory response (34).

While the above studies have illuminated key aspects of mRNA 3' end processing in immune cell regulation, the potential role of C/P proteins in differentiation of monocytes to macrophages, which results in a dramatic change of cellular characteristics including attachment to surfaces, increased size with extended filopodia, and cell cycle arrest, has not been explored. We hypothesized that the changes in cell morphology that occur during macrophage differentiation will be accompanied by changes in mRNA 3' end processing efficiency as well as choice of poly(A) site, and that perturbation of the C/P machinery will affect the course of this differentiation. In the current study, we show that this is indeed the case. There are significant changes in APA when monocytes are differentiated to form macrophages, and an increase in the components of the C/P machinery during the differentiation correlates with an increase in processing efficiency. Genetic manipulation of the C/P protein CstF64 led to altered APA and differentiation properties of monocytes. Overall, our findings indicate an important role for regulation of polyadenylation in macrophage differentiation.

RESULTS

Establishment of the monocyte-macrophage differentiation systems.

To study the impact of alternative polyadenylation in monocyte to macrophage differentiation, we used two well-established systems – monocytic cell lines and human primary monocytes (35-37). The monocytic cell lines U937 and THP-1 can be differentiated to macrophages by treatment with phorbol 12-myristate 13-acetate (PMA) (38). While both are commonly used to generate macrophages for functional studies, the two cell lines have very different origins, with THP-1 being derived from a patient with Acute Myeloid Leukemia and U937 from a patient with histiocytic lymphoma (39).

After 6 hours of treatment with 30 nM of PMA, U937 cells are in a transition phase in which they stop dividing and are loosely adhered to the dish, with little change in morphology (Fig. 1A). By 18-24 hours, almost all cells showed macrophage-like phenotypes including increased cell size, strong adherence to the dish, and appearance of filopodia. Monocyte-macrophage differentiation is associated with decreased expression of monocyte surface markers and increased expression of macrophage markers (40). The classic macrophage surface markers such as CD16 (41), CD68 (42), HLA-DRA (43), ICAM1 (44) and CD38 (45) were considerably increased over the 24-hour time course and the monocyte marker CD14 (46) had reduced expression in differentiated U937 cells (Fig. 1B). As observed previously (47,48), these cells also displayed increased attachment and no longer proliferated after PMA treatment (Fig. 1C and 1D). THP-1 cells behaved similarly to the U937 cells in terms of morphological changes, marker expression, attachment, and proliferation (Supplementary Fig. S1A-D). Upon differentiation of human blood-derived primary monocytes with M-CSF, characteristic morphological changes were also observed (Supplementary Fig. S1F) and protein expression of the macrophage markers such as CD16, CD68, HLA-DRA, ICAM1 and CD38 were increased, while the monocyte marker CD14 decreased (Supplementary Fig. S1G).

To further validate that differentiation was successful in our systems, we measured the mRNA levels of genes important for macrophage differentiation and function at two critical time points in the transition from monocytes to macrophages - 6 hours after induction when U937 and THP-1 cells show some signs of differentiation and 24 hours, when differentiation is complete. Genes encoding proteins such as the surface marker CD86, the adhesion molecule ICAM1, transcription factors including JUN, FOS, NFKB1, and IKZF1, cytokines such as Oncostatin-M (OSM) and IL12A, all increased in expression at 6 hours and some also at 24 hours after treating U937 and THP-1 cells with PMA (Fig. 1E and Supplementary Fig. S1E). The expression of these genes also increased during differentiation of human primary monocytes (Supplementary Fig. S1H). These differentiation protocols all create naïve macrophages that are poised for different types of activity depending on what additional stimuli they receive.

Integrative analysis of poly(A) site mapping data identifies expression changes and 3′ UTR shortening and lengthening events during macrophage differentiation.

To determine the effects of PMA treatment on global poly(A) site usage and gene expression in U937 cells, we used Poly(A)-Click sequencing, which returns sequences adjacent to the poly(A) site and has been used in recent studies to assess APA and expression changes (49-53). The differential expression for a particular gene can be determined by collapsing all mapped poly(A) sites within each gene into one count. As might be expected for monocytes undergoing differentiation, genes with increased expression at 6h are strongly enriched in pathways involved in the immune system and macrophage function (Supplementary Fig. S2A). Pathways associated with DNA repair were predominant in the decreased expression set. At 24 hours and similar to what was seen at 6 hours, the categories with significant increases in expression were heavily skewed towards the immune system and pathways related to macrophage function (Supplementary Fig. S2B). The increased expression of macrophage-related genes

Fig 1. Morphological changes and altered expression of cell-surface markers and mRNAs during monocyte differentiation.

(A) The effect of differentiation on cellular morphology. U937 cells exposed to PMA (30 nM) for 6 and 24 hours and observed by phase contrast microscopy (40X). (B) Western blots of total cell lysates from U937 cells differentiated with PMA (30 nM) for 0, 1, 6, 18, and 24 hours. Blots were probed with antibodies against CD16, CD68, HLA-DRA, ICAM1, CD38 and CD14 with β-actin as the loading control. (C) Attachment assay of U937 cells. Cells were treated with PMA for 0, 6, and 24 hours and live cells visualized and counted by the Trypan blue exclusion assay. The graph presents the percentage of cells that are suspended or attached. (D) Proliferation assay of U937 cells. The number of cells before and after differentiation depicts the absolute cell number in millions, compared to normal cell proliferation in the absence of PMA. The figure represents mean ± SE from three independent experiments. (E) Real-time quantitative PCR (RT-qPCR)–based analysis of the expression of genes important for macrophage differentiation and function *(CD86, ICAM1, JUN, FOS, NFKB1, IKZF1, OSM and IL12A)*. The RT-qPCR analysis used primers designed in the first exon to represent the total transcript level and the figure shows $\log_2$-fold changes in expression of these genes at 6 hours and 24 hours post PMA treatment of U937 cells, where values are normalized to ACTB mRNA and the 0 hour (no PMA) time point. The figure represents mean ± SE from three independent experiments. P value <0.05 was considered significant, where * = $P \leq 0.05$; ** = $P \leq 0.01$; *** = $P \leq 0.001$; **** = $P \leq 0.0001$.

Supplementary Figure S1. Morphological changes and altered expression of cell-surface markers and mRNAs during differentiation of THP-1 cells and human primary monocytes
(A) The effect of differentiation on THP-1 cellular morphology. THP-1 cells exposed to PMA (30 nM) for 6 and 24 hours and observed by phase contrast microscopy (40X). (B) Western blots of total cell lysates from THP-1 cells differentiated with PMA (30 nM) for 0, 1, 6, 18 and 24 hours. Blots were probed with antibodies against CD16, CD68, HLA-DRA, ICAM1, CD38 and CD14 with β-actin as the loading control. (C) Attachment assay for THP-1 cells, performed as described for Fig. 1C. (D) Proliferation assay for THP-1 cells performed as described for Fig. 1D. (E) Effect of THP-1 differentiation on the expression level of mRNAs from genes involved in macrophage differentiation and function. RT-qPCR assays performed for genes as described in Fig. 1E. (F) The effect of M-CSF on cellular characteristics of human primary monocytes. Human primary cells were exposed to 50 ng/ml human M-CSF for 5 and 7 days and morphological features with respect to control cells observed by phase contrast microscopy (40X). (G) Western blots of total cell lysates from human primary monocytes differentiated with M-CSF for 0, 5 and 7 days probed with antibodies against CD16, CD68, HLA-DRA, ICAM1, CD14 and CD38, with GAPDH as the loading control. (H) RT-qPCR analysis of mRNAs from human primary monocytes (undifferentiated and differentiated for 5 and 7 days) for similar genes as Fig. 1E.

1
2

Supplementary Figure S2. Gene set enrichment analysis of mRNAs with expression changes in U937 cells after PMA treatment.

Functional annotation clustering of mRNAs whose expression is regulated after 6h (A) and 24h (B) of differentiation. The 10 most significant GO-process enriched gene groups according to GSEA analysis are ranked based on negative $\log_{10}$ (P-values) for upregulated (left panel) and downregulated (right panel) transcripts. Green indicates biological process; red, canonical process. Pathways classified according to specific cellular processes are grouped together as indicated by the individual symbols at the bottom of the graphs.

further confirms that our differentiation protocol is effective. The down-regulated set at 24 hours showed marked enrichment of genes involved in cell cycle and chromosome organization. Decreased expression of these genes is consistent with the exit from the cell cycle upon PMA treatment.

We also asked if there were changes in poly(A) site usage during differentiation. The degree of APA usage was calculated by changes in Percentage of Distal poly(A) site Usage Index (delta-PDUI), which identifies lengthening (positive index) or shortening (negative index) of 3′ UTRs having statistical significance between differentiated and control samples. For U937 cells differentiated for 6 hours, sequencing identified polyadenylated mRNAs from a total of 12,208 genes of which 331 genes had mRNAs with significant changes in length in the terminal 3' UTR of the 3' most exon, designated as terminal 3' UTR. Of these, 139 exhibited an increase in shorter isoforms, and 113 showed an increase in longer isoforms, while 79 showed a complex pattern involving multiple poly(A) sites, where use of some went up and others went down (Fig. 2A). At 24 hours, signals from 12,145 genes were obtained, and mRNAs from a greater number of genes (864) had changes in poly(A) site usage compared to 6 hours. Of these, 376 exhibited shortening, 305 exhibited lengthening and 183 showed a complex pattern (Fig. 2A). At both time points, the number of significantly shortened APA events was slightly higher compared to the significantly lengthened APA events. Detailed outputs for all the APA events are supplied in Supplementary Tables S3 and S4.

Shortening of transcripts in the terminal 3' UTR has sometimes, but not always been associated with increased mRNA expression, with the converse for mRNA lengthening (21,28,54-58). To assess how gene expression correlated with changes in terminal 3' UTR length during macrophage differentiation, the 3' UTR shortened and lengthened gene sets were categorized into those with up-regulated or down-regulated expression and delta-PDUI versus fold change in expression was plotted for the 6 and 24 hour time points (Fig. 2B and Fig. 2C). A comparable number of genes in the lengthened or shortened groups at both time points exhibited increases or decreases in mRNA expression, and many genes with significant changes in isoform length showed no or only small changes in expression, indicating a poor correlation between 3' UTR length and steady-state mRNA level.

Fig 2. Evidence of APA in U937 cells after 6 and 24 hours of PMA treatment.
(A) Bar graph representing the numbers of genes with APA changes in the terminal 3' UTR after 6h and 24h of PMA treatment with respect to control undifferentiated U937 cells. Red, green and blue represents shortening, lengthening and complex (where some went up and others went down) APA patterns in the dataset. The criteria for a significant APA change is that at least one PAC in an exon must change greater than 1.5 fold with p_{Adj} <0.1, resulting in a difference in fractional usage of that PAC of at least 10%. (B & C) Log$_2$-fold change in gene expression is plotted against the change in Percentage of Distal poly(A) site Usage Index (delta-PDUI) for 3'-UTR-altered genes in 6h PMA-treated U937 cells with respect to control undifferentiated ones (B) or in 24h PMA-treated U937 samples (C). Plots for fold change vs poly(A) site usage (delta PDUI) where genes on the left half of each graph represent 3′-UTR-shortened hits, whereas those on the right half represent 3′-UTR lengthened hits.

Previous work has shown that genes with significant changes in their 3' UTR isoform expression ratios are cell type- and pathway-specific (28). To determine the biological pathways impacted by APA during macrophage differentiation, we performed Gene Set Enrichment Analysis (GSEA) (59) on genes which exhibited significant shortening or lengthening of transcripts. Analysis of the 0 vs 6h shortened set showed an enrichment of categories related to immune system or signal pathways related to macrophage function, as well as metabolism and cellular transport (Fig. 3A). Intracellular transport is important for both cytokine secretion and phagocytosis, and modulation of metabolism is recognized as a central player in macrophage activation (60). In contrast, the lengthened set was primarily related to the immune system.

The affected pathways for the 0 vs 24h APA changes were quite different from those affected at 6 hours. Unlike what was observed for 6h, APA changes at 24h that involved shortening affected pathways not specific to the immune system but nevertheless relevant to macrophage differentiation such as cytoskeletal organization, cell cycle, and chromosome organization, while the lengthened set was more mixed and included categories related to post-translational modification, cell cycle and signaling pathways involved in macrophage-mediated inflammatory responses (Fig. 3B).

Examples of genome browser plots of the mapped poly(A) sites for genes which showed mRNA shortening or lengthening in U937 cells are shown in Supplementary Fig. S3A for the 6 hour time point (CIAPIN1 and EIF1) and in Fig. 4 for the 24 hour time point (shortened: SERPINB1 and PTCH1; lengthened: PSAT1 and PSMD10). One way to confirm the presence of isoforms differing in their 3' ends and shifts in use of these isoforms is 3' rapid amplification of cDNA ends (RACE) assays (61-64). Thus, 3' RACE assays were used to detect the different mRNA isoforms, and these confirmed the APA changes depicted by the genome browser plots from the poly(A)-sequencing data of U937 cells (Fig. 5 and Supplementary Fig. S3B). SERPINB1 has 3 major poly (A) sites (PAC-1, 3 and 5 as shown in Fig. 4A), but the 3' RACE was not able to detect the longest isoform, perhaps because of the strong bias towards amplifying the short isoform limited the sensitivity for longer products. Nevertheless, in accordance with the genome browser plot, we did observe a reduction in use of the middle p(A) site (marked as D in Fig. 4A). The differentiation of the other monocytic cell line, THP-1 also produced shortening or lengthening of the same genes after 24 hours of PMA treatment (Supplementary Fig. S4A and B). Three of the four APA events were also detected after differentiation of human primary monocytes (Supplementary Fig. S4C and D).

Some genes have poly(A) sites located upstream of the gene's last exon. Use of these intronic sites creates a new terminal exon and changes the amount of coding sequence in the mRNA. We also observed changes in intronic poly(A) site use, but at a lower frequency than changes in the terminal 3' UTR. At 6 hours, 9 genes showed increased use of these poly(A) sites and one showed decreased use, and at 24 hours, 45 genes showed an increase and 30 showed a decrease (Supplementary Fig. S5A and B).

Fig 3. Gene set enrichment analysis (GSEA) of APA events in U937 cells after PMA treatment.
Functional annotation clustering of mRNAs whose APA is regulated and after 6h (A) and 24h (B) of
differentiation. The 10 most significant GO-process enriched gene groups for biological process (green) or
canonical process (red) are ranked based on negative $\log_{10}$ (P-values) for shortened (left panel) and
lengthened (right panel) transcripts. Pathways classified according to specific cellular processes are
grouped together as indicated by the individual symbols at the bottom of the graphs.

**Supplementary Figure S3. Changes in poly(A) site use in U937 cells after 6 hours of PMA
treatment and validation by 3'RACE.**
(A) UCSC genome browser plots of RNA sequencing tracks highlighting the 3′-UTR profile differences for
shortened and lengthened genes after 6h PMA treatment with respect to control (0h). The colors of the
tracks represent 0h (red) and 6h (blue). Proximal (P) and distal (D) poly(A) sites are indicated with red
stars. The green arrow defines the direction of the coding strand, blue arrow defines the direction of
chromosome co-ordinates and tag counts are indicated on the y axis. Additionally, positions and
chromosome co-ordinates of the annotated PACs are indicated at the top of each browser plot. (B)
Representative 1% agarose gel images for the 3′ RACE RT-PCR from U937 cells (untreated and 6h
PMA-treated) for shortened gene CIAPIN1 and lengthened gene EIF1 (upper panel) using gene specific
primer at the last exon-exon junction and reverse anchor primer. Normalized intensities of long and short
bands for each PCR were quantified and a long/short ratio determined and normalized to the No PMA
control. The graph (lower panel) represents mean ± SE from at least two independent experiments. P
value <0.05 was considered significant, where * = P ≤ 0.05; ** = P ≤ 0.01; *** = P ≤ 0.001.

Fig 4. Changes in poly(A) site use in U937 cells after 24 hours of PMA treatment.
UCSC genome browser plots of RNA sequencing tracks highlighting the 3′-UTR profile differences for
shortened genes (A) or lengthened genes (B) after 24 hours of PMA treatment with respect to control
(0h). The colors of the tracks represent 0h (red) and 24h (green). Proximal (P) and distal (D) poly(A) sites
are indicated with red stars. The green arrow defines the direction of the coding strand, and tag counts
are indicated on the y axis. Additionally, positions and chromosome co-ordinates of the annotated PACs
are indicated at the top of each plot.

Fig 5. Validation of APA events in U937 by 3' RACE.
Representative 1% agarose gel images for the 3′ RACE RT-PCR from U937 cells (untreated and 24h
PMA-treated) for (A) shortened genes SERPINB1, PTCH1 and (B) lengthened genes PSAT1, PSMD10
using gene specific primers at the last exon-exon junction and a common reverse anchor primer.
Normalized intensities of long and short bands for each PCR were quantified and a long/short ratio
determined and normalized to the No PMA control. The corresponding graphs represent mean ± SE from
at least two independent experiments. P value <0.05 was considered significant, where * = P ≤ 0.05; ** =
P ≤ 0.01; *** = P ≤ 0.001.

**Supplementary Figure S4. Validation of APA events in THP-1 and human primary monocytes by 3'
RACE.**
Representative 1% agarose gel images for the 3′ RACE RT-PCR from control and differentiated THP-1
cells and human primary monocytes for shortened genes SERPINB1 and PTCH1 (A and C), lengthened
genes PSAT1 and PSMD10 (B and D) with gene specific primer at the last exon-exon junction and
reverse anchor primer. Normalized intensities of long and short bands for each PCR were quantified and
a long/short ratio determined and normalized to the No PMA or undifferentiated controls. The
corresponding graphs (right) represents mean ± SE from at least two independent experiments. P value
<0.05 was considered significant, where * = P ≤ 0.05; ** = P ≤ 0.01; *** = P ≤ 0.001.

Supplementary Figure S5. Evidence of intronic polyadenylation events. (A) and (B) mRNAs undergoing intronic polyadenylation in U937 cells treated with PMA for 6h and 24h, classified according to their delta PDUI changes in both directions. Every exon with multiple PACs having "splicing APA" event (Supplementary Tables S3 and S4) is plotted here, and the PDU for 0h is on the x-axis and 24h on the y-axis. Depending upon the significance levels reported by the DPAC pipeline, blue dots indicate lengthening or decreased usage of the poly(A) site whereas pink dots indicate shortening or increased usage of the poly(A) site. Dotted grey lines indicate a change in PDU of 0.1 in either direction and the events (black dots) within the lines are considered not significantly altered.

Levels of cleavage/polyadenylation (C/P) complex proteins are upregulated at different times during monocyte-to-macrophage differentiation.

The analysis of the poly(A)-Click seq data described above shows that the APA occurring during monocyte-macrophage differentiation affected relevant functional categories of genes. We next wanted to explore possible mechanisms for eliciting this APA and examined the C/P complex proteins, as numerous studies have shown that changing the levels of these proteins can affect poly(A) site choice (65,66). The core C/P complex is comprised of six factors: CPSF, CstF, CFI$_m$, CFII$_m$, symplekin and poly(A) polymerase (PAP) (Fig. 6B) [reviewed in (67)]. The C/P proteins, with the exception of PAP, increased in level during differentiation of U937 cells, but with different kinetics and not necessarily in the same manner for the individual subunits of each factor (Fig. 6A). For example, CstF50, CstF64 and CstF77 increased early at 1 hour after PMA treatment, before differentiated phenotypes are observable. CPSF100, CPSF73 and CFI$_m$59 increased at 6 hours when cells begin to attach, whereas CFI$_m$68, FIP1, CPSF160, Clp1, Pcf11, symplekin and CstF64т, a conserved paralog of CstF64 (68) in mammals, increased later (18-24 hours), when the cells are mostly differentiated and attached. Some subunits, such as WDR33 and CstF77, decreased when differentiation was completed. The C/P proteins increased in similar ways during differentiation of the THP-1 monocytic cell line, with comparable kinetics (Supplementary Fig. S6A). We also examined the level of one subunit from each of the four multi-subunit C/P factors in human primary monocytes and found that CFI$_m$25, FIP1, CstF64 and Pcf-11 showed a striking increase upon differentiation (Supplementary Fig. S6B). Overall, our findings show that C/P proteins increase during macrophage differentiation but that some may be limiting in concentration at different points in the differentiation process.

To determine whether the increase in levels of C/P proteins during differentiation was reflected in changes in their mRNA levels, we analyzed the expression from poly(A)-Click mapping data of U937 cells before and after differentiation. Interestingly, the increase in C/P proteins did not correlate with an increase in mRNA expression (Fig. 6C). On the contrary, the mRNA levels remained either unchanged or decreased over the course of differentiation. RNA seq data from studies on differentiation of U937 cells (69), THP-1 and human primary monocyte cells (70) revealed the same outcome for C/P factor mRNAs (Supplementary Fig. S6C). These results indicate that steady-state mRNA levels are not a good predictor of the polyadenylation capacity of a cell.

Fig 6. Effect of macrophage differentiation on the expression of C/P proteins and mRNAs in U937 cells.
(A) Whole cell lysates from U937 cells treated with PMA for 0, 1, 6, 18 and 24 hours were separated by 10% SDS-PAGE and western blotting was performed for the indicated subunits of the C/P complex, with β-actin as the loading control. Each western blot (upper panel) was performed in three biological replicates of the differentiation process and the quantified data is shown in the lower panel. (B) Schematic of the mammalian core C/P complex. (C) Normalized RNA expression levels of C/P factors. The normalized gene counts for each replicate was obtained from the DEseq2 pipeline of the poly(A)-Click seq data for control vs treatment with PMA for 6 and 24 hours, and log2 fold change were plotted as bar graphs for the indicated C/P proteins. Dotted grey lines indicate a change in normalized expression of 1 in either direction and the events within the lines are considered not significantly altered. It was not possible to include all factors because of insufficient reads. (D) Processing efficiency of MYC and ACTB transcripts for U937 cells. The bar graph shows the ratio of unspliced (US) RNA transcripts (detected by RT-qPCR with a primer pair upstream of the poly(A) site) to unprocessed RNA (detected by RT-qPCR with a primer pair that spans the poly(A) site) as shown in the schematic. The cDNA preparation was done by using random hexamer.

Supplementary Figure S6. Effect of macrophage differentiation on the expression of C/P proteins and mRNAs in THP-1 cells and human primary monocytes.
(A) Whole cell lysates from THP-1 cells treated with PMA for 0, 1, 6, 18 and 24 hours were separated by 10% SDS-PAGE and western blotting was performed for the indicated subunits of the C/P complex, with β-actin as the loading control. Each western blot (upper panel) was performed in three biological replicates of the differentiation process and the quantified data is shown in the lower panel.
(B) Western blot analysis of CF1$_m$25, hFip1, CstF64 and Pcf-11 in human primary monocytes treated with MCSF for 0, 5, or 7 days. (C) Differential expression of C/P factors between differentiated and undifferentiated samples. The normalized gene counts for each replicate was obtained from the expression changes of C/P protein encoding genes using published datasets of undifferentiated and differentiated U937 (69), THP-1 and human primary monocytes (70). Details of dataset parameters are given in the Supplementary Table S2 (sheets 3 and 4). Mean values were plotted as bar graphs for the indicated C/P proteins. Dotted grey lines indicate a change in normalized expression of 1 in either direction and the events within the lines are considered not significantly altered. (D) Processing efficiency of MYC and ACTB transcripts for THP-1 cells. The bar graph shows the ratio of unspliced (US) RNA transcripts (detected by RT-qPCR with a primer pair upstream of the poly(A) site) to unprocessed RNA (detected by RT-qPCR with a primer pair that spans the poly(A) site) as shown in the schematic. The cDNA preparation was done by using random hexamers.

Differentiation of monocytic cells increases processing efficiency at single poly(A) sites.

To test whether PMA-mediated differentiation alters the efficiency of 3'-end processing *in vivo*, we determined the level of uncleaved transcripts that will contain sequence upstream and downstream of a poly(A) site. We used previously described primers to monitor cleavage of ACTB or MYC transcripts in vivo by RT-qPCR of total RNA reverse transcribed with random hexamers (71,72). ACTB has a single poly(A) site and MYC has a major poly(A) site used 93% of the time (73). One primer pair detects RNA containing sequence that spans the poly(A) sites (Span) of the ACTB or MYC genes. Because we expect uncleaved RNA to be nuclear, we compared this value to that obtained using a primer pair which detects unspliced transcripts (US), which would also be nuclear. The processing efficiency at each gene's poly(A) site can be quantified by the ratio of "US" PCR product to that of "Span" product. If processing efficiency increases, fewer "Span" products are expected relative to the "US" control Fig. 6D (72). The processing efficiency at each gene's poly(A) site can be quantified by the ratio of US PCR product to that of Span product. Compared to undifferentiated cells, more mRNA from both genes was processed at 24 hours, the time when most of the U937 or THP-1 cells have differentiated (Fig. 6D and Supplementary Fig. S6D). Respective "minus RT" controls gave insignificant signals (Supplementary Fig. S7A) indicating the effect to be entirely coming from the transcripts. This data indicates that consistent with the increase in C/P proteins, differentiation positively affected processing efficiency at these ACTB and MYC poly(A) sites.

Supplementary Figure S7. Processing efficiency minus RT control and cellular complexity of U937 cells by FACS.
(A) Minus RT control for processing efficiency of MYC and ACTB transcripts for U937 and THP-1 cells.
PCR was done by adding equivalent amount of RNA that was used to prepare cDNA for Fig. 6D and S6D
(1µg). The bar graph shows the ratio of unspliced (US) RNA transcripts (detected by RT-qPCR with a
primer pair upstream of the poly(A) site) to unprocessed RNA (detected by RT-qPCR with a primer pair
that spans the poly(A) site) as shown in Fig. 6D.
(B) Changes in cellular complexity determined by FACs analysis of side (SSC) and forward scattering
(FSC) values for untreated and 24h PMA-treated U937 cells.

Genetic manipulation of CstF64 alters monocyte-macrophage differentiation and switches poly(A) site choice.

As described above, monocyte differentiation led to the increased expression of C/P proteins and altered APA of mRNAs from groups of genes known to be important for macrophage differentiation and function. We thus wanted to test whether altering the level of one of these proteins could impact the APA and macrophage differentiation. We focused on CstF64, because it has been shown to influence APA during several types of transitions in cell state, in particular some of which involve the immune system (21,26,29,33,74,75). Based on these previous studies, we hypothesized that while CstF64 is likely not to be the only C/P protein affecting APA in macrophage differentiation, alteration of its levels would affect both APA and the progression of differentiation.

To test our hypothesis, we generated stable cell lines in which the expression of CstF64 in U937 cells could be knocked down (KD) by doxycycline-inducible expression of shRNA against CstF64 mRNAs. Depletion of CstF64 was done by doxycycline treatment for 3 days and then PMA was added for one day before harvesting the cells. Assessment of the extent of protein KD by two different shRNA sequences against CstF64 showed considerable decrease in expression, indicating efficient KD, while levels of the homolog CstF64τ remained unaffected (Fig. 7A). The macrophage surface markers CD16, CD68, HLA-DRA and CD38 were substantially reduced compared to the control shRNA when either CstF64 shRNA was used for KD, while ICAM1 was not affected. On the other hand, the levels of monocyte marker CD14 increased on CstF64 KD (Fig. 7A).

The changes in expression of monocyte-macrophage markers suggested that CstF64 KD might also affect the cellular changes that occur during macrophage differentiation. Indeed, counting of suspended and attached cells after 24 hours of PMA treatment showed that CstF64 KD in U937 cells significantly reduced cell attachment with both shRNAs (Fig. 7B, left panel). Normally by 24 hours, the cells have stopped dividing but are viable. However, the total cell number at 24 hours was greater in CstF64 KD cells compared to the control, indicating a slowing of the PMA-induced cell cycle arrest (Fig. 7B, right panel). Macrophage differentiation is also characterized by the formation of large, structurally complex cells (Fig. 1) (2), and this change can be detected using flow cytometry to measure the side (SSC) and forward scattering (FSC) of live cells (76). An increase in intensities of both FSC and SSC indicate the formation of cells with complex morphological features (also called increased granularity or complexity) upon PMA-mediated differentiation (Supplementary Fig. S7B). U937 cells depleted of CstF64 demonstrated reduced granularity (Fig. 7C).

Next, to determine if overexpression (OE) of CstF64 would promote macrophage differentiation, we generated stably transfected U937 cells inducibly overexpressing this C/P protein. CstF64 was efficiently overexpressed after 72 hours of doxycycline exposure, while CstF64τ levels did not change (Fig. 7D). The cell proliferation rate was not affected by CstF64 OE (Fig. 7E, right panel). However, overexpression increased expression of macrophage markers such as CD16, CD68, HLA-DRA, ICAM1 and CD38.

and decreased CD14 expression (Fig. 7D). In addition, there was an increase in the
number of attached cells from 0 to 16% (Fig. 7E, left panel) and by FACS analysis,
U937 cells overexpressing CstF64 demonstrated an almost two-fold increase in
granularity (Fig. 7F). It is important to note that these cells were not treated with the
differentiation inducer PMA.

Fig 7. Knockdown or overexpression of CstF64 in U937 cells alters macrophage differentiation.
(A) Western blot analysis of C/P factors CstF64 and CstF64τ, macrophage-specific markers CD16, CD68,
HLA-DRA, ICAM1, CD38 and the monocyte marker CD14 in U937 cells stably transfected with
doxycycline-inducible control shRNA and two different CstF64 shRNAs (shRNA 2 and shRNA 3). Addition
of 3 nM doxycycline for 3 days supplemented with 30 nM PMA one day prior to harvesting was done for
all three cell lines. (B) Attachment assay (left panel) and changes in total number of PMA-treated cells
(right panel) after knockdown of CstF64 compared to the shRNA control. Cells were treated with
doxycycline, after which equal number of cells were treated with PMA for 24 h and then counted. The
figures are representative of at least three independent experiments. (C) Changes in cellular complexity
determined by FACs analysis of side (SSC) and forward scattering (FSC) values for 24h PMA-treated
U937 cells stably expressing control shRNA and shRNA against CstF64. (D) Western blot analysis of C/P
factors CstF64, CstF64τ and macrophage-specific markers CD68, HLA-DR, ICAM1, CD38 and the
monocyte marker CD14 in U937 cells stably transfected with control overexpression vector (OE-control)
and those overexpressing CstF64 (OE-CstF64) and treated with 3 nM doxycycline for 3 days prior to
harvesting. No PMA treatment has been done. (E) Attachment assay (left panel) and changes in total
number of cells (right panel) after overexpression of CstF64 compared to the OE-control. (F) Changes in
cellular complexity or granularity determined by FACs analysis of side (SSC) and forward scattering
(FSC) values for doxycycline-induced U937 cells transfected with OE-control or OE-CstF64. Percentage
indicates a quantitation of cells that have altered granularity.

In addition to these changes in cellular properties, 3' RACE analysis showed that
CstF64 KD resulted in increased usage of the distal poly(A) sites of the mRNAs
SERPINB1 and *PTCH1,* which would otherwise be shortened after 24 hours of PMA
treatment, whereas CstF64 OE without PMA treatment caused decreased usage of the
distal poly(A) sites of these mRNAs (Fig. 8). This finding is consistent with studies in
other systems showing that CstF64 promotes use of proximal poly(A) sites (64,77,78).

In summary, depletion of CstF64 caused changes in cell properties indicative of poorer
macrophage differentiation and shifts to longer mRNA isoforms of genes which are
normally shortened during differentiation. In contrast, CstF64 overexpression caused
the appearance of differentiated phenotypes (cell surface markers, attachment,
granularity, and transcript shortening) in the absence of the differentiation inducer.

Fig 8. Knockdown or overexpression of CstF64 in U937 cells alters APA.
Representative 1% agarose gel images for the 3′ RACE RT-PCR in U937 cells for shortened genes
SERPINB1 (A) and *PTCH1* (B), using gene specific primers at the last exon-exon junction and a common
reverse anchor primer. In each subfigure, the effect of depletion (knockdown) of CstF64 and the effect of
overexpression of CstF64 are indicated, using the same treatment protocols as described in Fig. 7.
Normalized intensities of long and short bands for each PCR were quantified and a long/short ratio
determined and normalized to the shRNA or OE control. The corresponding graphs (right) represents
mean ± SE from at least two independent experiments. P value <0.05 was considered significant, where *
= P ≤ 0.05; ** = P ≤ 0.01; *** = P ≤ 0.001.

DISCUSSION

Macrophage activity is needed for a robust response to infection and tissue damage but
when deregulated, it also contributes to many human diseases (3-7). As such, it is
critical that monocyte to macrophage differentiation is appropriately regulated to
preserve the balance of various monocyte and macrophage populations and ensure the
appropriate functional consequences. The findings of our study indicate that regulation
of polyadenylation is one of the ways in which gene expression is controlled during
macrophage differentiation.

Macrophage differentiation is characterized by both 3' UTR shortening and
lengthening events.

APA profiles are known to vary according to the differentiation and growth state of cells,
with a general shift to distal poly(A) sites (3' UTR lengthening) often observed with
terminal differentiation (12,23,25,66). We found that upon macrophage differentiation,
the distribution of mRNA isoforms shifts as the monocyte precursor stops dividing and
changes morphology. There were similar numbers of 3' UTR shortening and
lengthening events, indicating that exit from the cell cycle during differentiation is not
always associated with predominant lengthening of mRNA isoforms. GSEA analysis
showed that lengthened and shortened gene sets are both enriched in groups of genes
important for macrophage differentiation and function. We have validated several APA
events by 3'RACE in both the U937 and THP-1 monocytic cell lines and in human
primary monocytes. Previous studies have shown that many of these genes examined
by 3'RACE have functions relevant to macrophage biology. For example, *SERPINB1*
has been reported to influence the innate immune system during influenza infection
(79), *PTCH1* is important for macrophage chemotaxis during gastric inflammation (80).
PSAT1 regulates the inflammatory response in macrophages (81), and PSMD10 can
regulate the nuclear localization of NF-κB (82). Our analysis shows that the APA profiles
are highly dynamic during macrophage differentiation. Moreover, the APA changes are
not globally affecting genes with multiple poly(A) sites but target sets of genes whose
functions are consistent with the need to exit the cell cycle and develop properties of a
mature macrophage.

For many genes, 3' UTR shortening has been shown to increase mRNA and protein
expression (54,83,84). However, other studies have demonstrated that global 3′-UTR
shortening in variety of tissues and cell types has limited influence on mRNA or protein
levels (28,57,58). Similar to these latter studies, our global analysis showed that the
direction of APA change is not a good general predictor of mRNA expression levels
during macrophage differentiation. It is likely that in this transition, the final levels of
different mRNAs are determined by APA-dependent and APA-independent changes in
mRNA stability as well as changes in transcription. Moreover, the function of some APA
changes during macrophage differentiation may be to affect other mRNA properties
such as translatability and localization. In summary, our findings support the idea that
APA is used as an additional mechanism to insure the timely expression and protein
output of critical groups of genes during the transition of monocytes to macrophages.

The increase in C/P proteins may facilitate APA during macrophage differentiation.

Our study of the U937 and THP-1 monocytic cell lines shows that differentiating monocytic cells display an overall trend of increased expression of proteins of the core C/P machinery. While there were some differences which could be expected given the different origins of the two lines, the general features of shifts in poly(A) site usage and increase in C/P proteins are consistent between the two. This finding was unexpected, as many studies have shown that the expression of C/P mRNA and protein increases with increased proliferation (65,77), but monocyte cell division ceases upon induction. The boosted expression of C/P proteins in mature macrophages was accompanied by an increase in the processing efficiency of genes with single poly(A) sites. The specific set of cellular changes accompanying macrophage differentiation, including increased mobility and metabolic activity, may necessitate an increased polyadenylation capacity. Interestingly, in spite of the increase in protein, the mRNA levels for the C/P proteins did not increase upon differentiation and most showed no change in 3' UTR length. The mRNAs for CFI_m25 and CFI_m68 were shortened and those of $CstF64\tau$ were lengthened (Supplementary Fig. S8), but it is not known whether such shifts can lead to increased translation. Previous studies have shown that PCF11 and CstF77 mRNA and protein levels can be increased by suppression of intronic poly(A) sites (85-87), but we did not see evidence of altered use of these sites in our transcriptome-wide poly(A) site mapping (data not shown). Instead, it is likely that mechanisms other than APA, such as enhanced translation or protein stability, leads to the accumulation of these and other C/P proteins.

Many studies have demonstrated that the core components of the mRNA 3′ processing machinery are also key regulators of global APA profiles. The sensitivity of transcripts to changes in levels of C/P factors depends on the strength of binding sites for these factors in the vicinity of poly(A) sites. In macrophages compared to monocytes, we found increases in proteins such as Fip1 and CstF64, known to promote use of upstream sites and production of mRNAs with shorter 3' UTRs (65,74,75,77,78), and as described below, showed that manipulation of CstF64 levels indeed altered APA in these cells. The increases in subunits of CFI_m, which is documented to promote use of downstream sites (83,88-92) and Pcf11, whose effect can vary depending on gene length and cell type (85,93), could explain why we observe APA changes in both directions. The increase in C/P proteins may not only affect APA, but could also support increased processing of genes which have single poly(A) sites with weak polyadenylation signals.

**Supplementary Figure S8. UCSC genome browser plots of RNA sequencing tracks highlighting
the 3′-UTR profile differences for genes encoding some C/P proteins**.
UCSC genome browser plots of RNA sequencing tracks highlighting the 3′-UTR profile differences for
CFIm25 (A) CFIm68 (B) and CstF64τ (C) after 24 hours of PMA treatment with respect to control (0h). The
color codes represent 0h (red) and 24h (green). Proximal (P) and distal (D) poly(A) sites are indicated
above. The green arrow defines the direction of the sense strand and tag counts are indicated by the y
axis. Additionally, positions of all the annotated PACs are indicated at the top of each browser plots.

CstF64 levels alter monocyte-macrophage differentiation and switch poly(A) site choice.

In this study, we examined the role of CstF64 in macrophage differentiation. CstF64 is the component of the core C/P complex that recognizes GU/U-rich signal sequences located downstream of the poly(A) site, and it also facilitates APA in numerous cellular contexts. In a well-studied example, an increase in CstF64 protein during terminal B-cell differentiation increases use of an intronic poly(A) site in immunoglobulin heavy chain pre-mRNAs that changes the protein product from a membrane-bound form to a secreted form (32,74). CstF64 also promotes use of weaker proximal poly(A) sites in proliferating fibroblasts (77), HeLa cells (78) and activated T cells (64), and non-canonical poly(A) site usage (94) as well as processing of specific acetylcholinesterase isoforms (95). Depletion of CstF64 suppressed proliferation of lung cancer cells and exogenous expression promoted proliferation of HEK293 cells (96). CstF64 has also been shown to regulate the cell cycle by controlling replication-dependent histone mRNA 3' end processing (97,98), CstF64 loss blocked the conversion of mouse embryonic stem cells into the endodermal lineage, but not into ectoderm or mesoderm (99). The same study also showed that CstF64 was needed for cardiomyocyte differentiation but not for neuronal differentiation (99), suggesting cell-specific requirements for CstF64. Activation of RAW 264.7 macrophage-like cells with LPS blocked proliferation and caused an increase in CstF64 but not other C/P subunits and an increase in proximal poly(A) site selection (33). As these examples clearly demonstrate, CstF64 is a major player in APA that can promote proliferation, differentiation or specific responses depending on the cell context. However, its role in the differentiation of monocytes to macrophage, a process that involves cell cycle exit and striking morphological changes, has not been previously studied.

In the current work, we found that CstF64 expression increased 6-8 fold during differentiation of monocytic cell lines. Our subsequent analysis showed that CstF64 depletion not only caused lengthening of transcripts that were otherwise shortened during differentiation but also had functional consequences by slowing the acquisition of macrophage properties such as exit from the cell cycle, attachment, marker expression and morphological changes. In contrast, overexpression of CstF64 stimulated macrophage-like changes and enhanced use of promoter proximal sites. Importantly, these effects of extra CstF64 happened in the absence of the powerful inducer used to initiate differentiation. Thus, our findings show that CstF64 must affect pathways important for macrophage differentiation. They are also in agreement with studies in other systems showing that CstF64 promotes use of promoter proximal sites. However, the increase in other C/P proteins and the observation that some transcripts lengthen during differentiation suggests that CstF64 is not the only driver and that other C/P factors and RNA-binding proteins with roles in APA regulation (12) will also influence isoform expression in macrophages. While our study with CstF64 provides evidence that regulation of polyadenylation is one of the mechanisms that promotes efficient macrophage differentiation, further research is needed to determine the consortium of proteins that regulate APA during this transition in cell state.

In summary, our study has provided new insights into the regulation of gene expression
in the differentiation of macrophages. The increase in levels of C/P proteins during
differentiation and the fact that manipulation of the C/P protein CstF64 alters the
progression of this differentiation supports a model in which changes in 3' end
processing efficiency are coordinated with changes in transcription and mRNA stability,
and this regulatory network helps create mature macrophages poised to respond
appropriately to a variety of extracellular stimuli through specific activities such as
migration, phagocytosis, antigen presentation, and immunomodulation. Given the
critical role of macrophages in regulating the inflammatory response, a greater
understanding of how APA promotes beneficial and harmful macrophage activity could
open new therapeutic treatment avenues.

MATERIAL & METHODS

Bacterial strains, DNA constructs and antibodies. NEB® Stable Competent *E. coli*
(Cat. #C3040) were used for transformation of plasmids. All siRNAs and primers were
purchased from Eton Biosciences (San Diego, CA). Inducible Lentiviral shRNA
constructs targeting CstF64 were purchased from GE Healthcare Dharmacon, Inc.
(sequence included in Supplementary Table S1). For overexpression (OE) of CstF64,
we cloned CstF64 coding sequence into pLIX_402 vector backbone (a gift from David
Root, Broad Institute, MA) and pLIX_402 vector (Addgene plasmid #41394) was used
as control. Antibodies were purchased from Santa Cruz, Bethyl and Abcam
(Supplementary Table S2).

Supplementary Table S1. Primers and shRNA sequence used in this study.

Purpose	Primer name	Forward primer 5'-3'	Reverse primer 5'-3'
RT-qPCR for gene expression	CD86 total mRNA	GACTGAGTAACATTCTCTTTGTGATGGCC	CTACTAGCTCACTCAGGCTTTGG
	ICAMI total mRNA	CCAATGTGCTATTCAAACTGC	CAGCGTAGGGTAAGGTTCTTG
	JUN total mRNA	AAGTTGCGACGGAGAGAAAA	ACGCAACCCAGTCCAACTT
	FOS total mRNA	GACCGAGCCCTTTGATGAC	GCCACTGTGCAGAGGCTC
	NFKB1 total mRNA	CAAGCAGCTCTGCAGCAG	ACTGTCATAGATGGCGTCTG
	IKZFI total mRNA	CACGTGATGGACCAAGC	CGGCTTGTGCAGCTGGTAC
	OSM total mRNA	CTCCTGGACCCCTATATACG	GTGTGGCATTGAGGGTCTG
	IL12A total mRNA	GAAAGTCCTGCCGCGCC	GTTTCTGGCCAAACTGAGGTG
	GAPDH mRNA	TGATGACATCAAGAAGGTGGTGAAG	TCCTTGGAGGCCATGTGGGCCAT
	ACTB mRNA	CATGTACGTTGCTATCCAGGC	CTCCTTAATGTCACGCACGAT
Processing activity	MYC Total	ATTACAGGTGTGAGCCAGGG	AGCCTGCCTCTTTTCCACA
	MYC Span	ATCATTGAGCCAAATCTTAAGTTGTG	CTCTGAAGGGGCAATTGATGA
	ACTB Total	TCAAGGTGGGTGTCTTTCCT	CCTGCTTGCTGATCCACATC
	ACTB Span	GCTTTTGGTCTCCCTGGGA	CTGCACTCTGGGTAAGGACA
CstF64 overexpression (For PCR cloning)		CTTACGGCTAGCGCCACCATGGCGGGTT TGAC (Nhe I enzyme site)	CATTAGACCGGTAGGTGCTCCAG TGG (Ase I enzyme site)

Purpose	Sequence
CstF64 knockdown shRNA #1	CGTAGAGAACGATCCACCG
CstF64 knockdown shRNA #2	TGGCATTCCAACCTGAGCC

CstF64 knockdown shRNA #3	CGTGCATCTATTCCTCGGG
Oligo d(T) adapter for 3'RACE	GCGAGCACAGAATTAATACGACTCACTATAGGTTTTTTTTTTTTTTTTVN
Reverse anchor-primer for 3'RACE	GCGAGCACAGAATTAATACGACT
SERPINB1 3'RACE exon Fwd	GGCCTGAAGAAGATTGAGGAAC
PTCH1 3'RACE exon Fwd	CAGCACCTTGTTACAGACAAGAAG
PSAT1 3'RACE exon Fwd	CATAGGTCTGTGGGAGGC
PSMD10 3'RACE exon Fwd	GCATGAGATCGCTGTCATGTTAC
CIAPIN1 3'RACE exon Fwd	GTGGAAACTGCTACCTGGGC
EIF1 3'RACE exon Fwd	GCGTTTAAGAAAAAGTTTGCCTG

1 **Supplementary Table S2. Antibodies used in this study.**

Catalogue no.	Name	Host	MW
SC-20052	CD16 Or FCGR3B	Mouse monoclonal	50 kDa
A6554	CD68	Rabbit polyclonal	110-140 kDa
SC-55593	HLA-DRα	Mouse monoclonal	36 kDa
SC-18853	ICAM1 (6.5B5)	Mouse monoclonal	85-110 kDa
SC-374650	CD38 (H-11)	Mouse monoclonal	45 kDa
SC -58951	CD14 (5A3B11B5)	Mouse monoclonal	53-55 kDa
ab8227 (Abcam)	Beta Actin	Rabbit polyclonal	43 kDa
SC-365062	GAPDH (G-9)	Mouse monoclonal	37 kDa
Bethyl A301-357A	CFI$_m$68 or CPSF6	Rabbit	68 kDa
Bethyl	CFI$_m$25 or NUDT21	Rabbit	25 kDa
SC-393880	CFI$_m$59 Or CPSF7	Mouse monoclonal	52 kDa
SC-398392	FIP1L1 (C-10)	Mouse monoclonal	67 kDa
SC-166281	CPSF160 or CPSF1 (G-10)	Mouse monoclonal	160 kDa
SC-165983	CPSF100 or CPSF2 (A-11)	Mouse monoclonal	100 kDa
SC-374466	WDR33 (D-1)	Mouse monoclonal	146 kDa
SC-393001	CPSF73 or CPSF3 (C-3)	Mouse monoclonal	73 kDa
SC-393316	CPSF30 or CPSF4 (D-1)	Mouse monoclonal	30 kDa
Bethyl A301-250A	CSTF50 or CSTF1	Rabbit polyclonal	50 kDa
SC-376575	CstF-77 (G-5) or CSTF3	Mouse monoclonal	77 kDa
SC-398862	CstF64 (H-1) or CSTF2	Mouse monoclonal	64 kDa
Bethyl A301-487A	CSTF2T/TauCSTF64/CstF64τ	Rabbit polyclonal	70 kDa
Bethyl AAS02642C	hClp1	Rabbit polyclonal	41 kDa
Bethyl A303-706A	Pcf11	Rabbit Polyclonal	173 kDa
SC-135410	Symplekin (H308)	Rabbit polyclonal	150 kDa
SC-32915	PAP (H-300)	Rabbit polyclonal	64 kDa (may be 90 kDa)

2
Cell culture, treatment and microscopy. All cells were maintained in RPMI 1640
medium supplemented with 2 mM L-glutamine and 10% heat-inactivated fetal bovine
serum (FBS) at 37°C in a humidified atmosphere of 5% CO_2. In order to culture human
primary monocytes (PBMCs), human whole blood in K3-EDTA was purchased from
Vitrologic (Charleston, SC). Collection and use of blood were conducted according to
the guidelines of IRB (Institutional Review Board)-approved informed consent and
protocols. Human PBMCs were isolated from whole blood by the Ficoll-Paque density

centrifugation method (100). Then, monocytes were enriched by negative selection
using EasySep™ Human Monocyte Isolation Kit (Stem Cell Technologies) according to
manufacturer's protocol. Differentiation of the U937 (ATCC #CRL-1593.2) and THP-1
(ATCC #TIB-202) human monocytic cell-lines into macrophages was performed by
treatment with 30 nM phorbol-12-myristate-13-acetate (PMA; Sigma-Aldrich) for up to
24 hours. This PMA concentration has been shown in previous studies to efficiently
differentiate U937 cells (38,101), and as described in the Results section, we found it to
be effective with both U937 and THP-1 cells. The human primary monocytes (isolated
from normal individual whole blood K3-EDTA purchased from Vitrologic #HWB-K3U)
were differentiated with 50 ng/ml recombinant human M-CSF (Peprotech) for 5–7 days
to obtain macrophages. For monitoring morphological changes, cells were cultured in 6-
well plates at a density of 1×10^6 cells/well, after which they were examined and
photographed using an EVOS FL fluorescence phase contrast microscope (Thermo
Fisher Scientific). All results were obtained from three independent experiments.

Western blot analysis. Western blot was performed with total cell extracts of untreated
and differentiated cells. For preparation of total cell extract, 1X RIPA buffer (20 mM Tris-
HCl pH 7.5, 150 mM NaCl, 1 mM Na_2EDTA, 1 mM EGTA, 1% NP-40, 1% sodium
deoxycholate, 2.5 mM sodium pyrophosphate, 1 mM β-glycerophosphate, 1 mM
Na_3VO_4 and 1 µg/ml leupeptin) was added, followed by incubation on ice for 15 min,
centrifugation at 25,000 rcf for 10 minutes at 4°C and supernatant collection. Protein
concentration was measured by BCA reagent (Pierce, Thermo Fisher Scientific,
Waltham, MA), and 50-80 µg protein was separated on a 10% polyacrylamide-SDS or
bis-tris gel and transferred to PVDF membrane. The membrane was blocked for 1 h in
5% non-fat dried milk in TBS-T (Tris Buffer Saline with Tween-20, *i.e.*, 0.05% Tween 20
in 1X TBS) buffer followed by rocking overnight at 4˚C with primary antibodies (antibody
details provided in Supplementary Table S2). Next day, unbound antibodies were
removed by 4 X 5 min washes with TBS-T buffer. The membrane was then incubated
with an HRP-conjugated secondary antibody at a dilution of 1:5000 for 1 h at room
temperature followed by 3 X 5 minute washes with TBS-T and a final wash with TBS.
The blot was developed with SuperSignal™ West Pico PLUS or Femto
Chemiluminescent Substrate (Thermo Fisher Scientific) and visualized with a ChemiDoc
XRS+ System (Bio-Rad) and quantitated using Image J (102). ACTB or GAPDH was
used as the control for individual western blots because their levels did not change upon
differentiation. Pre-stained protein markers were used as internal molecular mass
standards, and each western blot was performed in three biological replicates of each
time point of the differentiation process.

RT-qPCR analysis for expression analysis. Total RNA extraction was carried out on
2×10^6 cells using Trizol according to manufacturer's protocol. 1.5 µg of RNA was
subjected to reverse transcription using oligo dT primer and Superscript III reverse
transcriptase. The cDNA was amplified by qPCR with specific primers (Supplementary
Table S1) using the C1000™ thermal cycler with CFX96 Touch Real-Time PCR
Detection System (Bio-Rad Laboratories). The relative expression of genes was
analyzed quantitatively by the $\Delta\Delta C_t$ method. ACTB RNA was used as the normalization

control for RT-qPCR-based RNA expression analyses because its level did not change upon differentiation according to the poly(A)-seq datasets. Primers for RNA expression were designed at the first exon according to the annotation featured in Ensembl (103).

RT-qPCR analysis for processing activity. We used a previously published assay to measure the processing efficiency at the single poly(A) sites of the MYC and ACTB genes (71,72). This assay uses random hexamers for reverse transcription of 1 μg of total RNA. RT-qPCR is then performed with a primer pair upstream of the poly(A) site, called "US" which measures the level of unspliced transcripts and another primer pair that spans the poly(A) site, called "span". The sequences of these primers are provided in Supplementary Table S1.

Poly(A)-Click sequencing and analysis. This was a paid service from Click-Seq Technologies, in which total RNAs isolated from control and differentiated U937 cells were subjected to Poly(A) Click-sequencing as described (49,50,53). Briefly, reverse-transcription with oligo-dT primer (unanchored) supplemented with small amounts of spiked-in 3'-azido-nucleotides (AzVTPs) was performed to initiate cDNA synthesis from within the poly(A) tail, ensuring stochastic termination upstream of the poly(A) tail. The resulting 3'-azido-blocked cDNA molecules were purified, 'click-ligated' and amplified to generate a cDNA library. These libraries were pooled and sequenced using the manufacturer's standard operating procedures on a NextSeq 550, obtaining 1×150 bp (130M) paired-end reads. Raw data from the sequencing reactions for four biological replicates for 0h vs 24h and two biological replicates for 0h vs 6h was analyzed by Click-Seq Technologies using Differential Poly(A)-Clustering (DPAC), a pipeline to pre-process raw poly(A)-tail targeted RNAseq data, map to a reference genome, identify and annotate the location of poly(A) sites, generate poly(A)-clusters (PACs) and determine the differential abundance of PACs and differential gene expression between two conditions (50). The raw reads and mapped reads for each library is provided in Supplementary Table S2 (sheets 1 and 2). Each of these libraries were trimmed and quality filtered as described (53). Gene counts collapse all mapped poly(A) sites within that one gene into one count, this provides differential gene expression data. On an average 87% of all the mapped reads were found within or 200 nucleotides downstream of annotated mRNA poly(A) sites. Multiple poly(A) sites occurring within 25 nucleotides of one another were grouped into a poly(A) cluster (PAC) and treated as a unit to determine alternative polyadenylation events within exons and introns. The cutoff to be considered a PAC is 10 counts. In the final stage of DPAC, the mapped reads from each individual samples are used to determine the frequency of PACs in each dataset by determining the overlaps of the 3' ends of the mapped reads with the provided poly(A) cluster database using bedtools (104). Data normalization and statistical tests are applied using the canonical DESeq2 (105) pipeline using local dispersion estimation and Independent Hypothesis Weighting (IHW) (106) to estimate false discovery rates and power maximization. The final output of DPAC for the 6h and 24h PMA-treated U937 samples are provided in Supplemental Tables S3 and S4.

Principal component analysis (PCA) comparing control and PMA-treated U937 cells (0h vs 6h and 0h vs 24h) was done with the help of the DESeq2 pipeline (105) to determine and identify outlying samples within datasets. Variables (i.e. the times of differentiation)

are transformed into orthogonal principal components where each principal component indicates the direction with the greatest variability within the data. PCA plots (Supplementary Fig. S9A, B) revealed that treatment time was the most critical variable. Sequencing data sets are uploaded to NIH-GEO with the accession number #GSE169140. Volcano plots illustrating changes in gene counts and PAC counts (Supplementary Fig. S9C, D) were generated for both 6h and 24h PMA treated U937 cells by plotting the $\log_2$ fold change values vs p_{Adj} value. Differential Gene or PAC usage is defined as >1.5 fold change with p_{Adj} <0.1. The volcano plots indicate that the number of differentially expressed genes and the number of PACs showing change in usage increases as differentiation progresses. Gene Set Enrichment Analysis (GSEA) was carried out using online resources from the Broad Institute (59). Poly(A) site locations for genes were extracted into BED files and visualized with the help of UCSC genome browser (107).

Analysis of existing RNA-sequencing datasets. We determined the expression changes of C/P protein encoding genes using published datasets of undifferentiated and differentiated U937 (69), THP-1 and human primary monocytes (70). Details of datasets parameters are given in the Supplementary Table S2 (sheets 3 and 4). FastQ files were downloaded from the NCBI Sequence Read Archive along with sample metadata. Sequences were aligned to the human genome version GRC38, with annotations from Ensembl release (108), using the STAR aligner version 2.7.5.c (109). The resulting gene count matrixes were assessed for differential expression between differentiated and undifferentiated samples using DESeq2 version 1.32.0 (105), in R version 4.1 (110).

3' RACE. We modified the previously published 3'RACE protocols for this study (111,112). Briefly, total RNA extraction was done using Trizol according to manufacturer's protocol. 1 μg of RNA was subjected to reverse transcription for 1 hour using 100 units of Superscript IV reverse transcriptase and an oligo(dT)-containing adapter primer to poly(A) mRNA. After cDNA synthesis, the RNA template is removed from the cDNA:RNA hybrid molecule by digestion with 2 units of RNase H after cDNA synthesis to increase the sensitivity of PCR. For purification, the prepared cDNA was then chloroform extracted and precipitated in 0.3M sodium acetate and 3 volumes of ethanol. Subsequently, cDNA was subjected to PCR reaction with HotStarTaq Master Mix Kit (Qiagen) utilizing gene-specific forward primer (designed at the junction of the last two exons) and a unique anchor primer complimentary to the oligo(dT)-containing adapter primer (sequences provided in Supplementary Table S1). The first primer extension was accomplished by incubation at annealing temperature for 1 min, followed by incubation at 72°C for 40 min. Exponential amplification was accomplished by 30-35 cycles of PCR at 94°C for 1 min, 55°C /60°C (3-5 degrees lower than the lowest primer T_m) for 30 sec and 72°C for 5 min, followed by an additional 10-min extension at 72°C. The products were resolved in 1% agarose gels along with 1 kb Plus DNA Ladder (New England Biolabs) and visualized using SYBR™ Safe DNA Gel Stain (Thermo Fisher) with a ChemiDoc XRS+ System (Bio-Rad) and quantitated using Image J (102).

Supplementary Figure S9. Analysis of p(A) site mapping data.
(A and B) Principal component analysis (PCA) plot of U937 Poly(A)-Click-sequencing data from samples of the control (0h) and cells treated with PMA for 6 hours (A) or 24 hours (B). Each dot represents a sample and each color represents the time of PMA treatment. (C and D) Volcano plots illustrating changes in gene expression counts and PAC counts. They were generated by plotting the $\log_2$ fold change values vs p_{Adj} value for 6 hour (left panel) and 24 hour (right panel) PMA-treated U937 cells with respect to control undifferentiated ones for differential gene expression counts (C) and poly(A)-cluster counts (D). The gene counts for 6h and 24h collapse all mapped p(A) sites within one gene into one number and provides differential gene expression data, and PACs are defined as groupings of p(A) sites that are found within 25 nucleotides of one another. The gene counts in each PAC are used to determine relative changes in site selection between samples. Differential Gene or PAC count is defined as >1.5 fold change with p_{Adj} <0.1 (independent hypothesis weighting).

Lentivirus construction and cell treatment. For shRNA-based KD, three different lentiviral SMARTvector constructs, designed for doxycycline-inducible expression of shRNA for human CstF64 gene silencing, were purchased from GE Healthcare Dharmacon, Inc. These vectors incorporate the Tet-On® 3G bipartite system to inducibly control of shRNA with the Tet 3G promoter (PTRE3G), while the expression of the Tet-On® 3G transactivator protein is under the control of a constitutive human EF-1 promoter for optimal expression in the monocyte-macrophage system. A SMART vector inducible lentiviral control shRNA was used to control for potential non-specific effects caused by expression of shRNA. The sequences of the shRNAs are in Supplementary Table S1. As shown in the Results, shRNA #2 & 3 gave efficient KD but shRNA #1 was not effective. To generate stable cells inducibly overexpressing CstF64, its coding sequence was amplified by PCR from U937 cDNA and cloned into the pLIX_402 inducible lentiviral expression vector by using NheI and AgeI restriction enzyme sites to generate the OE construct. The primers used for cloning are mentioned in Supplementary Table S1.

These recombinant plasmids were co-transfected with the components of Dharmacon™ Trans-Lentiviral packaging kit into HEK293FT cells using FuGENE® HD (Promega Corp.) according to manufacturer's protocol. Transfection of HEK293FT cells was done in 6-well plates when the cells were 80-85% confluent, and transfection media changed after 16 hours. The recombinant lentiviruses were harvested at 48 h post-transfection, spun at 1250 rpm for 5 minutes and filtered by a 0.45 µm filter to remove cells and debris. Purified virus was used for infecting monocytic cells. $1.5X10^6$ target cells were seeded in 4 ml of media per well in a 6-well plate and cultured overnight. Lentiviral particles (1-2 ml per well) were added the next day to the cells in culture media containing 10 µg/ml polybrene for efficient infection. Selection of cells stably expressing CstF64-shRNAs or control-shRNA and pLIX_402 overexpressing CstF64 or control pLIX_402 started 72 h post-transfection or a time determined by the expression of GFP from the cells. Growth medium was replaced with fresh selection medium containing 1 µg/mL of puromycin. Puromycin-containing medium was refreshed every 2–3 days, and selection was completed after approximately 1 week, after which clones were expanded for 2 more weeks and frozen for later use. The Tet-On 3G induction system was activated by 3 nM doxycycline for 3 days for both KD and OE transfections. Additionally for KD clones, 30 nM PMA was added one day prior to harvesting for RNA and protein extraction.

Attachment assay. U937 cells were either untreated or treated with PMA and left for 24 hours to attach. After that, unbound cells were collected from the plate by washing gently with PBS and collected for counting (denoted as "suspended"). The attached cells were collected by trypsinizing to release them from the plate. Cells were then stained with trypan blue and counted in a hemocytometer and represented as percentage of suspended or attached cells over total (suspended + attached). The figures are average of three independent observations.

Flow cytometry. Phenotypic analysis of monocytes and macrophages was performed using flow cytometry. After differentiation, cells were recovered from culture plates and washed with phosphate-buffered saline (PBS) containing 1% paraformaldehyde for

fixation. In order to avoid non-specific binding and background fluorescence of cells expressing high levels of Fc receptors, the cells were incubated with Fc Receptor Blocking Solution (Human TruStain FcX, Biolegend) according to the manufacturer's instructions for 10 minutes, followed by washing with cold PBS. Zombie Aqua™ Fixable Viability kit (BioLegend) at a final dilution of 1:2000 was used for the exclusion of dead cells. Thereafter, cells were analyzed with a LSR II flow cytometer (BD Biosciences) using FACSDiva software, and data were analyzed with the FlowJo software. Primary gating was done on the basis of forward versus side scatter (FSC vs SSC) to compare size and granularity (complexity) of live cells across samples (40).

Statistical analysis. All experiments were performed in at least three independent sets. Data are presented as mean ± SE. Statistical analysis was performed using GraphPad Prism 6.01 (GraphPad Software Inc., La Jolla, CA, USA). Two-way analysis of variance (ANOVA) was performed to determine the significance between the groups. Considerations were * = $P ≤ 0.05$; ** = $P ≤ 0.01$; *** = $P ≤ 0.001$; **** = $P ≤ 0.0001$. A P value <0.05 was considered significant.

Materials availability. All plasmids and cell lines will be available upon request.

DATA AVAILABILITY

Poly(A) sequencing data sets are available in the NCBI Gene Expression Omnibus (GEO), accession number #GSE169140. Supplementary materials are attached with this manuscript.

FUNDING

This work was supported by the National Institutes of Health [grant numbers 1R01GM101010 and 1R01AI152337] to Claire Moore and partially supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health [under grant numbers P20GM103423 and P20GM104318] to Mount Desert Island Biological Laboratory.

ACKNOWLEDGMENTS

We thank Tufts University faculty Alexei Degterev and Karl Munger for providing cell lines and Marta Rodriguez-Garcia for consulting with us on FACS analysis. We thank members of Karl Munger's lab for use of the cell culture facility and want to acknowledge the critical inputs of the members of FACS core, Tufts University, and other members of the Moore lab. We would also like to thank Andrew Routh and Elizabeth Jaworski of ClickSeq Technologies for their valuable input on the bioinformatic analysis, and Nathaniel Maki and Chris Wilson of Mount Desert Island Biological Laboratory for helping perform the alignments for RNA-seq analysis.

COMPETING INTERESTS

No conflict of interest.

AUTHOR CONTRIBUTION

CM and SM conceived the study and CM provided general oversight. SM developed the strategy and methodology, acquired the data, reported and organized the findings. CM and SM designed the experiments, interpreted the results, and wrote and revised the manuscript. JG contributed to the computational analyses.

REFERENCES

1.   Shi, C. and Pamer, E.G. (2011) Monocyte recruitment during infection and inflammation. *Nat Rev Immunol*, **11**, 762-774.
2.   Italiani, P. and Boraschi, D. (2014) From Monocytes to M1/M2 Macrophages: Phenotypical vs. Functional Differentiation. *Front Immunol*, **5**, 514.
3.   Mammana, S., Fagone, P., Cavalli, E., Basile, M.S., Petralia, M.C., Nicoletti, F., Bramanti, P. and Mazzon, E. (2018) The Role of Macrophages in Neuroinflammatory and Neurodegenerative Pathways of Alzheimer's Disease, Amyotrophic Lateral Sclerosis, and Multiple Sclerosis: Pathogenetic Cellular Effectors and Potential Therapeutic Targets. *Int J Mol Sci*, **19**.
4.   Salmaninejad, A., Valilou, S.F., Soltani, A., Ahmadi, S., Abarghan, Y.J., Rosengren, R.J. and Sahebkar, A. (2019) Tumor-associated macrophages: role in cancer development and therapeutic implications. *Cell Oncol (Dordr)*, **42**, 591-608.
5.   Gomez-Rial, J., Rivero-Calle, I., Salas, A. and Martinon-Torres, F. (2020) Role of Monocytes/Macrophages in Covid-19 Pathogenesis: Implications for Therapy. *Infect Drug Resist*, **13**, 2485-2493.
6.   Punzo, F., Bellini, G., Tortora, C., Pinto, D.D., Argenziano, M., Pota, E., Paola, A.D., Martino, M.D. and Rossi, F. (2020) Mifamurtide and TAM-like macrophages: effect on proliferation, migration and differentiation of osteosarcoma cells. *Oncotarget*, **11**, 687-698.
7.   Zindel, J., Peiseler, M., Hossain, M., Deppermann, C., Lee, W.Y., Haenni, B., Zuber, B., Deniset, J.F., Surewaard, B.G.J., Candinas, D. *et al.* (2021) Primordial GATA6 macrophages function as extravascular platelets in sterile injury. *Science*, **371**.
8.   Zhou, X., Li, W., Wang, S., Zhang, P., Wang, Q., Xiao, J., Zhang, C., Zheng, X., Xu, X., Xue, S. *et al.* (2019) YAP Aggravates Inflammatory Bowel Disease by Regulating M1/M2 Macrophage Polarization and Gut Microbial Homeostasis. *Cell Rep*, **27**, 1176-1189 e1175.
9.   Labonte, A.C., Tosello-Trampont, A.C. and Hahn, Y.S. (2014) The role of macrophage polarization in infectious and inflammatory diseases. *Mol Cells*, **37**, 275-285.
10.   de The, H. (2018) Differentiation therapy revisited. *Nat Rev Cancer*, **18**, 117-127.
11.   Sykes, D.B., Kfoury, Y.S., Mercier, F.E., Wawer, M.J., Law, J.M., Haynes, M.K., Lewis, T.A., Schajnovitz, A., Jain, E., Lee, D. *et al.* (2016) Inhibition of Dihydroorotate Dehydrogenase Overcomes Differentiation Blockade in Acute Myeloid Leukemia. *Cell*, **167**, 171-186 e115.
12.   Neve, J., Patel, R., Wang, Z., Louey, A. and Furger, A.M. (2017) Cleavage and polyadenylation: Ending the message expands gene regulation. *RNA Biol*, **14**, 865-890.
13.   Cheng, L.C., Zheng, D., Zhang, Q., Guvenek, A., Cheng, H. and Tian, B. (2021) Alternative 3' UTRs play a widespread role in translation-independent mRNA association with the endoplasmic reticulum. *Cell Rep*, **36**, 109407.
14.   Muller-McNicoll, M., Botti, V., de Jesus Domingues, A.M., Brandl, H., Schwich, O.D., Steiner, M.C., Curk, T., Poser, I., Zarnack, K. and Neugebauer, K.M. (2016) SR proteins are NXF1 adaptors that link alternative RNA processing to mRNA export. *Genes Dev*, **30**, 553-566.

15. Neve, J., Burger, K., Li, W., Hoque, M., Patel, R., Tian, B., Gullerova, M. and Furger, A. (2016) Subcellular RNA profiling links splicing and nuclear DICER1 to alternative cleavage and polyadenylation. *Genome Res*, **26**, 24-35.

16. Zheng, D., Wang, R., Ding, Q., Wang, T., Xie, B., Wei, L., Zhong, Z. and Tian, B. (2018) Cellular stress alters 3'UTR landscape through alternative polyadenylation and isoform-specific degradation. *Nat Commun*, **9**, 2268.

17. Berkovits, B.D. and Mayr, C. (2015) Alternative 3' UTRs act as scaffolds to regulate membrane protein localization. *Nature*, **522**, 363-367.

18. Park, H.J., Ji, P., Kim, S., Xia, Z., Rodriguez, B., Li, L., Su, J., Chen, K., Masamha, C.P., Baillat, D. *et al.* (2018) 3' UTR shortening represses tumor-suppressor genes in trans by disrupting ceRNA crosstalk. *Nat Genet*, **50**, 783-789.

19. Wang, R., Zheng, D., Yehia, G. and Tian, B. (2018) A compendium of conserved cleavage and polyadenylation events in mammalian genes. *Genome Res*, **28**, 1427-1441.

20. MacDonald, C.C. (2019) Tissue-specific mechanisms of alternative polyadenylation: Testis, brain, and beyond (2018 update). *Wiley Interdiscip Rev RNA*, **10**, e1526.

21. Sandberg, R., Neilson, J.R., Sarma, A., Sharp, P.A. and Burge, C.B. (2008) Proliferating cells express mRNAs with shortened 3' untranslated regions and fewer microRNA target sites. *Science*, **320**, 1643-1647.

22. Elkon, R., Drost, J., van Haaften, G., Jenal, M., Schrier, M., Oude Vrielink, J.A. and Agami, R. (2012) E2F mediates enhanced alternative polyadenylation in proliferation. *Genome Biol*, **13**, R59.

23. Ji, Z., Lee, J.Y., Pan, Z., Jiang, B. and Tian, B. (2009) Progressive lengthening of 3' untranslated regions of mRNAs by alternative polyadenylation during mouse embryonic development. *Proc Natl Acad Sci U S A*, **106**, 7028-7033.

24. Guvenek, A. and Tian, B. (2018) Analysis of alternative cleavage and polyadenylation in mature and differentiating neurons using RNA-seq data. *Quant Biol*, **6**, 253-266.

25. Ji, Z. and Tian, B. (2009) Reprogramming of 3' untranslated regions of mRNAs by alternative polyadenylation in generation of pluripotent stem cells from different cell types. *PLoS One*, **4**, e8419.

26. Cheng, L.C., Zheng, D., Baljinnyam, E., Sun, F., Ogami, K., Yeung, P.L., Hoque, M., Lu, C.W., Manley, J.L. and Tian, B. (2020) Widespread transcript shortening through alternative polyadenylation in secretory cell differentiation. *Nat Commun*, **11**, 3182.

27. Li, W., Park, J.Y., Zheng, D., Hoque, M., Yehia, G. and Tian, B. (2016) Alternative cleavage and polyadenylation in spermatogenesis connects chromatin regulation with post-transcriptional control. *BMC Biol*, **14**, 6.

28. Lianoglou, S., Garg, V., Yang, J.L., Leslie, C.S. and Mayr, C. (2013) Ubiquitously transcribed genes use alternative polyadenylation to achieve tissue-specific expression. *Genes Dev*, **27**, 2380-2396.

29. Chuvpilo, S., Zimmer, M., Kerstan, A., Glockner, J., Avots, A., Escher, C., Fischer, C., Inashkina, I., Jankevics, E., Berberich-Siebelt, F. *et al.* (1999) Alternative polyadenylation events contribute to the induction of NF-ATc in effector T cells. *Immunity*, **10**, 261-269.

30. Takagaki, Y. and Manley, J.L. (1998) Levels of polyadenylation factor CstF-64 control IgM heavy chain mRNA accumulation and other events associated with B cell differentiation. *Mol Cell*, **2**, 761-771.

31. Edwalds-Gilbert, G. and Milcarek, C. (1995) Regulation of poly(A) site use during mouse B-cell development involves a change in the binding of a general polyadenylation factor in a B-cell stage-specific manner. *Mol Cell Biol*, **15**, 6420-6429.

32. Peterson, M.L. (2011) Immunoglobulin heavy chain gene regulation through polyadenylation and splicing competition. *Wiley Interdiscip Rev RNA*, **2**, 92-105.

33. Shell, S.A., Hesse, C., Morris, S.M., Jr. and Milcarek, C. (2005) Elevated levels of the 64-kDa cleavage stimulatory factor (CstF-64) in lipopolysaccharide-stimulated macrophages influence gene expression and induce alternative poly(A) site selection. *The Journal of biological chemistry*, **280**, 39950-39961.

34. Schutte, R.J., Parisi-Amon, A. and Reichert, W.M. (2009) Cytokine profiling using monocytes/macrophages cultured on common biomaterials with a range of surface chemistries. *J Biomed Mater Res A*, **88**, 128-139.

35. Harris, P. and Ralph, P. (1985) Human leukemic models of myelomonocytic development: a review of the HL-60 and U937 cell lines. *J Leukoc Biol*, **37**, 407-422.

36. Bekkering, S., Blok, B.A., Joosten, L.A., Riksen, N.P., van Crevel, R. and Netea, M.G. (2016) In Vitro Experimental Model of Trained Innate Immunity in Human Primary Monocytes. *Clin Vaccine Immunol*, **23**, 926-933.

37. Auwerx, J. (1991) The human leukemia cell line, THP-1: a multifacetted model for the study of monocyte-macrophage differentiation. *Experientia*, **47**, 22-31.

38. Mukherjee, S. and Sengupta Bandyopadhyay, S. (2016) Phorbol-12-myristate-13-acetate (PMA) mediated transcriptional regulation of Oncostatin-M. *Cytokine*, **88**, 209-213.

39. Chanput, W., Peters, V. and Wichers, H. (2015) In Verhoeckx, K., Cotter, P., Lopez-Exposito, I., Kleiveland, C., Lea, T., Mackie, A., Requena, T., Swiatecka, D. and Wichers, H. (eds.), *The Impact of Food Bioactives on Health: in vitro and ex vivo models*, Cham (CH), pp. 147-159.

40. Daigneault, M., Preston, J.A., Marriott, H.M., Whyte, M.K. and Dockrell, D.H. (2010) The identification of markers of macrophage differentiation in PMA-stimulated THP-1 cells and monocyte-derived macrophages. *PloS one*, **5**, e8668.

41. Wang, Z.Q., Bapat, A.S., Rayanade, R.J., Dagtas, A.S. and Hoffmann, M.K. (2001) Interleukin-10 induces macrophage apoptosis and expression of CD16 (FcgammaRIII) whose engagement blocks the cell death programme and facilitates differentiation. *Immunology*, **102**, 331-337.

42. Holness, C.L. and Simmons, D.L. (1993) Molecular cloning of CD68, a human macrophage marker related to lysosomal glycoproteins. *Blood*, **81**, 1607-1613.

43. Costabel, U., Bross, K.J., Andreesen, R. and Matthys, H. (1986) HLA-DR antigens on human macrophages from bronchoalveolar lavage fluid. *Thorax*, **41**, 261-265.

44. Novosad, J., Holicka, M., Novosadova, M., Krejsek, J. and Krcmova, I. (2011) Rapid onset of ICAM-1 expression is a marker of effective macrophages activation during infection of Francisella tularensis LVS in vitro. *Folia Microbiol (Praha)*, **56**, 149-154.

45. Amici, S.A., Young, N.A., Narvaez-Miranda, J., Jablonski, K.A., Arcos, J., Rosas, L., Papenfuss, T.L., Torrelles, J.B., Jarjour, W.N. and Guerau-de-Arellano, M. (2018) CD38 Is Robustly Induced in Human Macrophages and Monocytes in Inflammatory Conditions. *Front Immunol*, **9**, 1593.

46. Ziegler-Heitbrock, H.W. and Ulevitch, R.J. (1993) CD14: cell surface receptor and differentiation marker. *Immunol Today*, **14**, 121-125.

47. Chang, M.Y., Huang, D.Y., Ho, F.M., Huang, K.C. and Lin, W.W. (2012) PKC-dependent human monocyte adhesion requires AMPK and Syk activation. *PLoS One*, **7**, e40999.

48. Prasad, A., Sedlarova, M., Balukova, A., Ovsii, A., Rac, M., Krupka, M., Kasai, S. and Pospisil, P. (2020) Reactive Oxygen Species Imaging in U937 Cells. *Front Physiol*, **11**, 552569.

49. Routh, A., Ji, P., Jaworski, E., Xia, Z., Li, W. and Wagner, E.J. (2017) Poly(A)-ClickSeq: click-chemistry for next-generation 3-end sequencing without RNA enrichment or fragmentation. *Nucleic Acids Res*, **45**, e112.

50. Routh, A. (2019) DPAC: A Tool for Differential Poly(A)-Cluster Usage from Poly(A)-Targeted RNAseq Data. *G3 (Bethesda)*, **9**, 1825-1830.

51. Cao, J., Verma, S.K., Jaworski, E., Mohan, S., Nagasawa, C.K., Rayavara, K., Sooter, A., Miller, S.N., Holcomb, R.J., Powell, M.J. *et al.* (2021) RBFOX2 is critical for maintaining alternative polyadenylation patterns and mitochondrial health in rat myoblasts. *Cell Rep*, **37**, 109910.

52. Chu, Y., Elrod, N., Wang, C., Li, L., Chen, T., Routh, A., Xia, Z., Li, W., Wagner, E.J. and Ji, P. (2019) Nudt21 regulates the alternative polyadenylation of Pak1 and is predictive in the prognosis of glioblastoma patients. *Oncogene*, **38**, 4154-4168.

53. Elrod, N.D., Jaworski, E.A., Ji, P., Wagner, E.J. and Routh, A. (2019) Development of Poly(A)-ClickSeq as a tool enabling simultaneous genome-wide poly(A)-site identification and differential expression analysis. *Methods*, **155**, 20-29.

54. Mayr, C. and Bartel, D.P. (2009) Widespread shortening of 3'UTRs by alternative cleavage and polyadenylation activates oncogenes in cancer cells. *Cell*, **138**, 673-684.

55. Ji, Z., Luo, W., Li, W., Hoque, M., Pan, Z., Zhao, Y. and Tian, B. (2011) Transcriptional activity regulates alternative cleavage and polyadenylation. *Mol Syst Biol*, **7**, 534.

56. Chang, J.W., Zhang, W., Yeh, H.S., de Jong, E.P., Jun, S., Kim, K.H., Bae, S.S., Beckman, K., Hwang, T.H., Kim, K.S. *et al.* (2015) mRNA 3'-UTR shortening is a molecular signature of mTORC1 activation. *Nat Commun*, **6**, 7218.

57. Gruber, A.R., Martin, G., Muller, P., Schmidt, A., Gruber, A.J., Gumienny, R., Mittal, N., Jayachandran, R., Pieters, J., Keller, W. *et al.* (2014) Global 3' UTR shortening has a limited effect on protein abundance in proliferating T cells. *Nat Commun*, **5**, 5465.

58. Spies, N., Burge, C.B. and Bartel, D.P. (2013) 3' UTR-isoform choice has limited influence on the stability and translational efficiency of most mRNAs in mouse fibroblasts. *Genome Res*, **23**, 2078-2090.

59. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S. *et al.* (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*, **102**, 15545-15550.

60. Stunault, M.I., Bories, G., Guinamard, R.R. and Ivanov, S. (2018) Metabolism Plays a Key Role during Macrophage Activation. *Mediators Inflamm*, **2018**, 2426138.

61. Graber, J.H., Nazeer, F.I., Yeh, P.C., Kuehner, J.N., Borikar, S., Hoskinson, D. and Moore, C.L. (2013) DNA damage induces targeted, genome-wide variation of poly(A) sites in budding yeast. *Genome Res*, **23**, 1690-1703.

62. Masamha, C.P., Xia, Z., Peart, N., Collum, S., Li, W., Wagner, E.J. and Shyu, A.B. (2016) CFIm25 regulates glutaminase alternative terminal exon definition to modulate miR-23 function. *RNA*, **22**, 830-838.

63. Venkat, S., Tisdale, A.A., Schwarz, J.R., Alahmari, A.A., Maurer, H.C., Olive, K.P., Eng, K.H. and Feigin, M.E. (2020) Alternative polyadenylation drives oncogenic gene expression in pancreatic ductal adenocarcinoma. *Genome Res*, **30**, 347-360.

64. Chatrikhi, R., Mallory, M.J., Gazzara, M.R., Agosto, L.M., Zhu, W.S., Litterman, A.J., Ansel, K.M. and Lynch, K.W. (2019) RNA Binding Protein CELF2 Regulates Signal-Induced Alternative Polyadenylation by Competing with Enhancers of the Polyadenylation Machinery. *Cell Rep*, **28**, 2795-2806 e2793.

65. Di Giammartino, D.C., Nishida, K. and Manley, J.L. (2011) Mechanisms and consequences of alternative polyadenylation. *Mol Cell*, **43**, 853-866.

66. Ren, F., Zhang, N., Zhang, L., Miller, E. and Pu, J.J. (2020) Alternative Polyadenylation: a new frontier in post transcriptional regulation. *Biomark Res*, **8**, 67.

67. Shi, Y. and Manley, J.L. (2015) The end of the message: multiple protein-RNA interactions define the mRNA polyadenylation site. *Genes Dev*, **29**, 889-897.

68. Dass, B., Tardif, S., Park, J.Y., Tian, B., Weitlauf, H.M., Hess, R.A., Carnes, K., Griswold, M.D., Small, C.L. and Macdonald, C.C. (2007) Loss of polyadenylation protein tauCstF-64 causes spermatogenic defects and male infertility. *Proc Natl Acad Sci U S A*, **104**, 20374-20379.

69. Haney, M.S., Bohlen, C.J., Morgens, D.W., Ousey, J.A., Barkal, A.A., Tsui, C.K., Ego, B.K., Levin, R., Kamber, R.A., Collins, H. *et al.* (2018) Identification of phagocytosis regulators using magnetic genome-wide CRISPR screens. *Nat Genet*, **50**, 1716-1727.

70. Liu, H., Lorenzini, P.A., Zhang, F., Xu, S., Wong, M.S.M., Zheng, J. and Roca, X. (2018) Alternative splicing analysis in human monocytes and macrophages reveals MBNL1 as major regulator. *Nucleic Acids Res*, **46**, 6069-6086.

71. Davidson, L., Muniz, L. and West, S. (2014) 3' end formation of pre-mRNA and phosphorylation of Ser2 on the RNA polymerase II CTD are reciprocally coupled in human cells. *Genes Dev*, **28**, 342-356.

72. Eaton, J.D., Davidson, L., Bauer, D.L.V., Natsume, T., Kanemaki, M.T. and West, S. (2018) Xrn2 accelerates termination by RNA polymerase II, which is underpinned by CPSF73 activity. *Genes Dev*, **32**, 127-139.

73. Herrmann, C.J., Schmidt, R., Kanitz, A., Artimo, P., Gruber, A.J. and Zavolan, M. (2020) PolyASite 2.0: a consolidated atlas of polyadenylation sites from 3' end sequencing. *Nucleic Acids Res*, **48**, D174-D179.

74. Takagaki, Y., Seipelt, R.L., Peterson, M.L. and Manley, J.L. (1996) The polyadenylation factor CstF-64 regulates alternative processing of IgM heavy chain pre-mRNA during B cell differentiation. *Cell*, **87**, 941-952.

75. Lackford, B., Yao, C., Charles, G.M., Weng, L., Zheng, X., Choi, E.A., Xie, X., Wan, J., Xing, Y., Freudenberg, J.M. *et al.* (2014) Fip1 regulates mRNA alternative polyadenylation to promote stem cell self-renewal. *EMBO J*, **33**, 878-889.

76. Druszczynska, M., Wlodarczyk, M., Janiszewska-Drobinska, B., Kielnierowski, G., Zawadzka, J., Kowalewicz-Kulbat, M., Fol, M., Szpakowski, P., Rudnicka, K., Chmiela, M. *et al.* (2013) Monocyte signal transduction receptors in active and latent tuberculosis. *Clin Dev Immunol*, **2013**, 851452.

77. Mitra, M., Johnson, E.L., Swamy, V.S., Nersesian, L.E., Corney, D.C., Robinson, D.G., Taylor, D.G., Ambrus, A.M., Jelinek, D., Wang, W. *et al.* (2018) Alternative polyadenylation factors link cell cycle to migration. *Genome Biol*, **19**, 176.

78. Yao, C., Biesinger, J., Wan, J., Weng, L., Xing, Y., Xie, X. and Shi, Y. (2012) Transcriptome-wide analyses of CstF64-RNA interactions in global regulation of mRNA alternative polyadenylation. *Proc Natl Acad Sci U S A*, **109**, 18773-18778.

79. Gong, D., Farley, K., White, M., Hartshorn, K.L., Benarafa, C. and Remold-O'Donnell, E. (2011) Critical role of serpinB1 in regulating inflammatory responses in pulmonary influenza infection. *J Infect Dis*, **204**, 592-600.

80. Chakrabarti, J., Dua-Awereh, M., Schumacher, M., Engevik, A., Hawkins, J., Helmrath, M.A. and Zavros, Y. (2022) Sonic Hedgehog acts as a macrophage chemoattractant during regeneration of the gastric epithelium. *NPJ Regen Med*, **7**, 3.

81. Raines, L.N., Zhao, H., Wang, Y., Chen, H.Y., Gallart-Ayala, H., Hsueh, P.C., Cao, W., Koh, Y., Alamonte-Loya, A., Liu, P.S. *et al.* (2022) PERK is a critical metabolic hub for immunosuppressive function in macrophages. *Nat Immunol*, **23**, 431-445.

82. Chen, Y., Li, H.H., Fu, J., Wang, X.F., Ren, Y.B., Dong, L.W., Tang, S.H., Liu, S.Q., Wu, M.C. and Wang, H.Y. (2007) Oncoprotein p28 GANK binds to RelA and retains NF-kappaB in the cytoplasm through nuclear export. *Cell Res*, **17**, 1020-1029.

83. Masamha, C.P., Xia, Z., Yang, J., Albrecht, T.R., Li, M., Shyu, A.B., Li, W. and Wagner, E.J. (2014) CFIm25 links alternative polyadenylation to glioblastoma tumour suppression. *Nature*, **510**, 412-416.

84. Jenal, M., Elkon, R., Loayza-Puch, F., van Haaften, G., Kuhn, U., Menzies, F.M., Oude Vrielink, J.A., Bos, A.J., Drost, J., Rooijers, K. *et al.* (2012) The poly(A)-binding protein nuclear 1 suppresses alternative cleavage and polyadenylation sites. *Cell*, **149**, 538-553.

85. Wang, R., Zheng, D., Wei, L., Ding, Q. and Tian, B. (2019) Regulation of Intronic Polyadenylation by PCF11 Impacts mRNA Expression of Long Genes. *Cell Rep*, **26**, 2766-2778 e2766.

86. Kamieniarz-Gdula, K., Gdula, M.R., Panser, K., Nojima, T., Monks, J., Wisniewski, J.R., Riepsaame, J., Brockdorff, N., Pauli, A. and Proudfoot, N.J. (2019) Selective roles of vertebrate PCF11 in premature and full-length transcript termination. *Mol Cell*, **74**, 158-172.

87. Luo, W., Ji, Z., Pan, Z., You, B., Hoque, M., Li, W., Gunderson, S.I. and Tian, B. (2013) The conserved intronic cleavage and polyadenylation site of CstF-77 gene imparts control of 3' end processing activity through feedback autoregulation and by U1 snRNP. *PLoS Genet*, **9**, e1003613.

88. Martin, G., Gruber, A.R., Keller, W. and Zavolan, M. (2012) Genome-wide analysis of pre-mRNA 3' end processing reveals a decisive role of human cleavage factor I in the regulation of 3' UTR length. *Cell Rep*, **1**, 753-763.

89. Li, W., You, B., Hoque, M., Zheng, D., Luo, W., Ji, Z., Park, J.Y., Gunderson, S.I., Kalsotra, A., Manley, J.L. *et al.* (2015) Systematic profiling of poly(A)+ transcripts modulated by core 3' end processing and splicing factors reveals regulatory rules of alternative cleavage and polyadenylation. *PLoS Genet*, **11**, e1005166.

90. Brumbaugh, J., Di Stefano, B., Wang, X., Borkent, M., Forouzmand, E., Clowers, K.J., Ji, F., Schwarz, B.A., Kalocsay, M., Elledge, S.J. *et al.* (2018) Nudt21 Controls Cell Fate by Connecting Alternative Polyadenylation to Chromatin Signaling. *Cell*, **172**, 106-120 e121.

91. Zhu, Z.J., Huang, P., Chong, Y.X., Kang, L.X., Huang, X., Zhu, Z.X. and Nie, L. (2016) MicroRNA-181a promotes proliferation and inhibits apoptosis by suppressing CFIm25 in osteosarcoma. *Mol Med Rep*, **14**, 4271-4278.

92. Yang, S.W., Li, L., Connelly, J.P., Porter, S.N., Kodali, K., Gan, H., Park, J.M., Tacer, K.F., Tillman, H., Peng, J. *et al.* (2020) A Cancer-Specific Ubiquitin Ligase Drives mRNA Alternative Polyadenylation by Ubiquitinating the mRNA 3' End Processing Complex. *Mol Cell*, **77**, 1206-1221 e1207.

93. Ogorodnikov, A., Levin, M., Tattikota, S., Tokalov, S., Hoque, M., Scherzinger, D., Marini, F., Poetsch, A., Binder, H., Macher-Goppinger, S. *et al.* (2018) Transcriptome 3'end organization by PCF11 links alternative polyadenylation to formation and neuronal differentiation of neuroblastoma. *Nat Commun*, **9**, 5331.

94. Hwang, H.W., Park, C.Y., Goodarzi, H., Fak, J.J., Mele, A., Moore, M.J., Saito, Y. and Darnell, R.B. (2016) PAPERCLIP Identifies MicroRNA Targets and a Role of CstF64/64tau in Promoting Non-canonical poly(A) Site Usage. *Cell Rep*, **15**, 423-435.

95. Nazim, M., Masuda, A., Rahman, M.A., Nasrin, F., Takeda, J.I., Ohe, K., Ohkawara, B., Ito, M. and Ohno, K. (2017) Competitive regulation of alternative splicing and alternative polyadenylation by hnRNP H and CstF64 determines acetylcholinesterase isoforms. *Nucleic Acids Res*, **45**, 1455-1468.

96. Aragaki, M., Takahashi, K., Akiyama, H., Tsuchiya, E., Kondo, S., Nakamura, Y. and Daigo, Y. (2011) Characterization of a cleavage stimulation factor, 3' pre-RNA, subunit 2, 64 kDa (CSTF2) as a therapeutic target for lung cancer. *Clin Cancer Res*, **17**, 5889-5900.

97. Youngblood, B.A., Grozdanov, P.N. and MacDonald, C.C. (2014) CstF-64 supports pluripotency and regulates cell cycle progression in embryonic stem cells through histone 3' end processing. *Nucleic Acids Res*, **42**, 8330-8342.

98. Romeo, V., Griesbach, E. and Schumperli, D. (2014) CstF64: cell cycle regulation and functional role in 3' end processing of replication-dependent histone mRNAs. *Mol Cell Biol*, **34**, 4272-4284.

99. Youngblood, B.A. and MacDonald, C.C. (2014) CstF-64 is necessary for endoderm differentiation resulting in cardiomyocyte defects. *Stem Cell Res*, **13**, 413-421.

100. Jaatinen, T. and Laine, J. (2007) Isolation of mononuclear cells from human cord blood by Ficoll-Paque density gradient. *Curr Protoc Stem Cell Biol*, **Chapter 2**, Unit 2A 1.

101. Baek, J.Y., Jun, D.Y., Taub, D. and Kim, Y.H. (2003) Characterization of human phosphoserine aminotransferase involved in the phosphorylated pathway of L-serine biosynthesis. *Biochem J*, **373**, 191-200.

102. Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*, **9**, 671-675.

103. Yates, A.D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M., Azov, A.G., Bennett, R. *et al.* (2020) Ensembl 2020. *Nucleic Acids Res*, **48**, D682-D688.

104. Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, **26**, 841-842.

105. Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, **15**, 550.

106. Ignatiadis, N., Klaus, B., Zaugg, J.B. and Huber, W. (2016) Data-driven hypothesis weighting increases detection power in genome-scale multiple testing. *Nat Methods*, **13**, 577-580.

107. Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M. and Haussler, D. (2002) The human genome browser at UCSC. *Genome Res*, **12**, 996-1006.

108. Howe, K.L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M., Azov, A.G., Bennett, R., Bhai, J. *et al.* (2021) Ensembl 2021. *Nucleic Acids Res*, **49**, D884-D891.

109. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M. and Gingeras, T.R. (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**, 15-21.

110. Team, R.C.R. (2018), *R: A language and environment for statistical computing*.

111. Green, M.R. and Sambrook, J. (2019) Rapid Amplification of Sequences from the 3' Ends of mRNAs: 3'-RACE. *Cold Spring Harb Protoc*, **2019**.

112. Scotto-Lavino, E., Du, G. and Frohman, M.A. (2006) 3' end cDNA amplification using classic RACE. *Nat Protoc*, **1**, 2742-2745.

A

Alternative polyadenylation events

- Shortening APA
- Lengthening APA
- Complex pattern

B

C

A Pathway analysis of genes undergoing <u>APA changes</u> after 6h of PMA treatment:

Shortened genes (0 vs 6):

Pathway	Value
GOBP_REGULATION_OF_CATABOLIC_PROCESS	20
GOBP_REGULATION_OF_CELLULAR_CATABOLIC_PROCESS	18
GOBP_POSITIVE_REGULATION_OF_INTRACELLULAR_TRANSPORT	10
GOBP_MACROMOLECULE_CATABOLIC_PROCESS	21
GOBP_INTRACELLULAR_TRANSPORT	23
GOBP_POSITIVE_REGULATION_OF_TRANSPORT	16
GOBP_CELLULAR_MACROMOLECULE_CATABOLIC_PROCESS	18
GOBP_REGULATION_OF_IMMUNE_SYSTEM_PROCESS	21
GOBP_POSITIVE_REGULATION_OF_INTRACELLULAR_PROTEIN_TRANSPORT	8
GOBP_REGULATION_OF_TRANSPORT	22
REACTOME_DAP12_SIGNALING	4
REACTOME_MTOR_SIGNALLING	4
BIOCARTA_TCR_PATHWAY	4
REACTOME_DAP12_INTERACTIONS	4
WP_B_CELL_RECEPTOR_SIGNALING_PATHWAY	5
REACTOME_PLATELET_ACTIVATION_SIGNALING_AND_AGGREGATION	7
REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION	16

x-axis: -log p value (0 2 4 6 8 10)

Lengthened genes (0 vs 6):

Pathway	Value
GOBP_APOPTOTIC_PROCESS	18
GOBP_IMMUNE_SYSTEM_DEVELOPMENT	13
GOBP_RESPONSE_TO_ABIOTIC_STIMULUS	14
GOBP_REGULATION_OF_DNA_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY	9
GOBP_REGULATION_OF_CELL_DEATH	16
GOBP_MACROMOLECULE_CATABOLIC_PROCESS	14
GOBP_NEGATIVE_REGULATION_OF_NF_KAPPAB_TRANSCRIPTION_FACTOR_ACTIVITY	5
GOBP_REGULATION_OF_CELL_DIFFERENTIATION	16
GOBP_NEGATIVE_REGULATION_OF_DNA_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY	6
GOBP_INTERLEUKIN_1_MEDIATED_SIGNALING_PATHWAY	5
KEGG_NEUROTROPHIN_SIGNALING_PATHWAY	6
BIOCARTA_IL1R_PATHWAY	4
WP_SIGNAL_TRANSDUCTION_THROUGH_IL1R	4
WP_ATM_SIGNALING_PATHWAY	4
REACTOME_TOLL_LIKE_RECEPTOR_TLR1_TLR2_CASCADE	5
WP_VEGFAVEGFR2_SIGNALING_PATHWAY	8
WP_IL1_SIGNALING_PATHWAY	4
REACTOME_TOLL_LIKE_RECEPTOR_4_TLR4_CASCADE	5
BIOCARTA_ATM_PATHWAY	3
REACTOME_TOLL_LIKE_RECEPTOR_CASCADES	5

x-axis: -log p value (0 2 4 6 8)

Legend:
- ● Metabolic process
- ■ Immune system or signal pathways related to macrophage function
- ▲ Cellular transport
- ○ DNA damage repair
- → Regulation of cell death
- □ Change in state or activity
- △ Protein modification such as small protein conjugation or phosphorylation
- 🟩 Biological process
- 🟥 Canonical process

B Pathway analysis of genes undergoing <u>APA changes</u> after 24h of PMA treatment:

Shortened genes (0 vs 24):

Pathway	Value
GOBP_CELL_CYCLE	63
GOBP_CHROMOSOME_ORGANIZATION	50
GOBP_MICROTUBULE_BASED_PROCESS	36
GOBP_CELL_CYCLE_PROCESS	48
GOBP_MICROTUBULE_CYTOSKELETON_ORGANIZATION	30
GOBP_CYTOSKELETON_ORGANIZATION	46
GOBP_ORGANELLE_ASSEMBLY	35
GOBP_REGULATION_OF_ORGANELLE_ORGANIZATION	42
GOBP_MICROTUBULE_ORGANIZING_CENTER_ORGANIZATION	15
GOBP_REGULATION_OF_HYDROLASE_ACTIVITY	42
REACTOME_CELL_CYCLE	31
REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION	42
REACTOME_CELL_CYCLE_MITOTIC	24
REACTOME_MITOTIC_PROMETAPHASE	15
REACTOME_M_PHASE	19
REACTOME_SUMOYLATION	13
REACTOME_RNA_POLYMERASE_II_TRANSCRIPTION	36
REACTOME_DNA_REPAIR	16
PID_TELOMERASE_PATHWAY	8
REACTOME_DISEASE	36

x-axis: -log p value (0 5 10 15 20)

Lengthened genes (0 vs 24):

Pathway	Value
GOBP_PROTEIN_MODIFICATION_BY_SMALL_PROTEIN_CONJUGATION_OR_REMOVAL	31
GOBP_REGULATION_OF_RESPONSE_TO_STRESS	36
GOBP_CHROMOSOME_ORGANIZATION	31
GOBP_REGULATION_OF_TRANSFERASE_ACTIVITY	26
GOBP_POSITIVE_REGULATION_OF_TRANSFERASE_ACTIVITY	21
GOBP_CELLULAR_RESPONSE_TO_DNA_DAMAGE_STIMULUS	24
GOBP_POSITIVE_REGULATION_OF_CATALYTIC_ACTIVITY	31
GOBP_POSITIVE_REGULATION_OF_KINASE_ACTIVITY	19
GOBP_PROTEIN_MODIFICATION_BY_SMALL_PROTEIN_CONJUGATION	24
GOBP_CELL_CYCLE	36
REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION	29
REACTOME_CELL_CYCLE	19
REACTOME_CELL_CYCLE_MITOTIC	16
REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM	19
REACTOME_SUMOYLATION_OF_DNA_DAMAGE_RESPONSE_AND_REPAIR_PROTEINS	6
REACTOME_MAPK_FAMILY_SIGNALING_CASCADES	10
PID_P38_ALPHA_BETA_PATHWAY	4
REACTOME_SUMOYLATION	8
REACTOME_MITOTIC_G1_PHASE_AND_G1_S_TRANSITION	7
REACTOME_DISEASE	24

x-axis: -log p value (0 5 10 15 20)

Legend:
- 🟩 Biological process
- 🟥 Canonical process
- ← Cell cycle and chromosome organization
- ✳ Cytoskeletal organization
- △ Protein modification such as small protein conjugation or phosphorylation
- ○ DNA damage repair
- ■ Immune system or signal pathways related to macrophage function

A U937 shortened genes (0 vs 24 hours):

B U937 lengthened genes (0 vs 24 hours):

A <u>U937 shortened genes</u>

SERPINB1   PTCH1

B <u>U937 lengthened genes</u>

PSAT1   PSMD10

A

U937 western blot

CFIm	Time after PMA treatment	0h	1h	6h	18h	24h
	CF Im68					
	CF Im25					
	CFIm59					

CPSF						
	FIP1					
	CPSF160					
	CPSF100					
	Wdr33					
	CPSF73					
	CPSF30					

CstF	Time after PMA treatment	0h	1h	6h	18h	24h
	CstF50					
	CstF77					
	CstF64					
	CstF64т					

CFIIm						
	Clp1					
	Pcf11					

Others						
	Symplekin					
	PAP					
	β-actin					

U937 western blot quantification

B

Mammalian core cleavage/polyadenylation complex

C

Cleavage polyadenylation factors — log2 fold change over 0h (6h PMA / 24h PMA)

D

Primers for unspliced (US)   Primers spanning p(A) site

$$\frac{US}{Span} \propto \text{Processing efficiency}$$

Processing efficiency — PMA-0h / PMA-6h / PMA-24h — MYC, ACTB — U937

A <u>SERPINB1:</u>

B <u>PTCH1:</u>

A Pathway analysis of genes undergoing <u>expression changes</u> after 6h of PMA treatment:

Upregulated genes (0 vs 6):

- ☐ GOBP_REGULATION_OF_INTRACELLULAR_SIGNAL_TRANSDUCTION — 232
- ◼ GOBP_LOCOMOTION — 241
- ◼ GOBP_CELL_MIGRATION — 207
- ☐ GOBP_CELL_ACTIVATION — 194
- ☐ GOBP_NEGATIVE_REGULATION_OF_RESPONSE_TO_STIMULUS — 208
- ◼ GOBP_REGULATION_OF_IMMUNE_SYSTEM_PROCESS — 201
- GOBP_POSITIVE_REGULATION_OF_MULTICELLULAR_ORGANISMAL_PROCESS — 186
- ◼ GOBP_RESPONSE_TO_CYTOKINE — 170
- ◼ GOBP_BIOLOGICAL_ADHESION — 191
- ➡ GOBP_APOPTOTIC_PROCESS — 220
- ◼ REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM — 108
- ◼ REACTOME_SIGNALING_BY_INTERLEUKINS — 76
- ● REACTOME_SIGNALING_BY_RECEPTOR_TYROSINE_KINASES — 74
- ◼ WP_IL18_SIGNALING_PATHWAY — 54
- ◼ KEGG_CYTOKINE_CYTOKINE_RECEPTOR_INTERACTION — 52
- ◼ REACTOME_INNATE_IMMUNE_SYSTEM — 111
- ◼ REACTOME_SIGNALING_BY_GPCR — 82
- ☐ REACTOME_SIGNALING_BY_NTRKS — 35
- ◼ REACTOME_HEMOSTASIS — 80
- WP_NUCLEAR_RECEPTORS_METAPATHWAY — 52

-log p value (0 20 40 60 80)

Downregulated genes (0 vs 6):

- ○ GOBP_DNA_REPAIR — 59
- ○ GOBP_DNA_METABOLIC_PROCESS — 76
- ○ GOBP_CELLULAR_RESPONSE_TO_DNA_DAMAGE_STIMULUS — 71
- ● GOBP_SMALL_MOLECULE_METABOLIC_PROCESS — 115
- ● GOBP_-VE_REGULATION_OF_NUCLEOBASE_COMPOUND_METABOLIC_PROCESS — 94
- ☐ GOBP_NEUROGENESIS — 97
- ○ GOBP_DOUBLE_STRAND_BREAK_REPAIR — 30
- ● GOBP_NEGATIVE_REGULATION_OF_SIGNALING — 86
- ○ GOBP_DNA_REPLICATION — 30
- ● GOBP_ORGANOPHOSPHATE_METABOLIC_PROCESS — 67
- REACTOME_RNA_POLYMERASE_II_TRANSCRIPTION — 86
- ○ REACTOME_HOMOLOGOUS_DNA_PAIRING_AND_STRAND_EXCHANGE — 10
- ○ REACTOME_HDR_THROUGH_HOMOLOGOUS_RECOMBINATION_HRR — 12
- ○ REACTOME_HOMOLOGY_DIRECTED_REPAIR — 17
- ○ REACTOME_DNA_REPAIR — 28
- ▲ REACTOME_RAB_GERANYLGERANYLATION — 11
- ○ REACTOME_RESOLUTION_OF_D_LOOP_STRUCTURES — 8
- ○ REACTOME_RESOLUTION_OF_D_LOOP_STRUCTURES — 7
- △ REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION — 74
- ○ REACTOME_DNA_DOUBLE_STRAND_BREAK_REPAIR — 17

-log p value (0 5 10 15 20)

Legend:
- ◼ Immune system or signal pathways related to macrophage function
- ➡ Regulation of cell death
- ○ DNA damage repair
- ● Metabolic process
- ☐ Change in state or activity
- ▲ Cellular transport
- △ Protein modification such as small protein conjugation or phosphorylation
- 🟩 Biological process
- 🟥 Canonical process

B Pathway analysis of genes undergoing <u>expression changes</u> after 24h of PMA treatment:

Upregulated genes (0 vs 24):

- ✳ GOBP_LOCOMOTION — 334
- ☐ GOBP_CELL_ACTIVATION — 282
- GOBP_REGULATION_OF_INTRACELLULAR_SIGNAL_TRANSDUCTION — 310
- ◼ GOBP_IMMUNE_EFFECTOR_PROCESS — 257
- ✳ GOBP_CELL_MIGRATION — 288
- GOBP_DEFENSE_RESPONSE — 305
- ◼ GOBP_RESPONSE_TO_CYTOKINE — 238
- ◼ GOBP_REGULATION_OF_IMMUNE_SYSTEM_PROCESS — 273
- GOBP_POSITIVE_REGULATION_OF_MOLECULAR_FUNCTION — 286
- GOBP_POSITIVE_REGULATION_OF_SIGNALING — 278
- ◼ REACTOME_INNATE_IMMUNE_SYSTEM — 199
- ◼ REACTOME_NEUTROPHIL_DEGRANULATION — 118
- ◼ REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM — 144
- ◼ REACTOME_SIGNALING_BY_INTERLEUKINS — 94
- REACTOME_HEMOSTASIS — 112
- KEGG_LYSOSOME — 44
- ◼ REACTOME_RHO_GTPASE_CYCLE — 84
- WP_TYROBP_CAUSAL_NETWORK — 32
- ◼ REACTOME_ADAPTIVE_IMMUNE_SYSTEM — 117
- NABA_MATRISOME — 134

-log p value (0 20 40 60 80 100)

Downregulated genes (0 vs 24):

- ⬅ GOBP_CELL_CYCLE — 377
- ⬅ GOBP_CHROMOSOME_ORGANIZATION — 291
- ⬅ GOBP_CELL_CYCLE_PROCESS — 301
- ⬅ GOBP_MITOTIC_CELL_CYCLE — 244
- ○ GOBP_DNA_METABOLIC_PROCESS — 232
- ⬅ GOBP_CELL_DIVISION — 162
- ○ GOBP_DNA_REPLICATION — 114
- ⬅ GOBP_REGULATION_OF_CELL_CYCLE — 221
- ○ GOBP_CELLULAR_RESPONSE_TO_DNA_DAMAGE_STIMULUS — 185
- ⬅ GOBP_CHROMOSOME_SEGREGATION — 113
- ⬅ REACTOME_CELL_CYCLE — 238
- ⬅ REACTOME_CELL_CYCLE_MITOTIC — 195
- ⬅ REACTOME_CELL_CYCLE_CHECKPOINTS — 112
- ⬅ REACTOME_M_PHASE — 120
- ⬅ REACTOME_MITOTIC_PROMETAPHASE — 82
- WP_RETINOBLASTOMA_GENE_IN_CANCER — 56
- ○ REACTOME_DNA_REPAIR — 95
- ⬅ REACTOME_CHROMOSOME_MAINTENANCE — 64
- ○ REACTOME_HOMOLOGY_DIRECTED_REPAIR — 60
- ⬅ REACTOME_RESOLUTION_OF_SISTER_CHROMATID_COHESION — 56

-log p value (0 50 100 150)

Legend:
- ⬅ Cell cycle and chromosome organization
- ✳ Cytoskeletal organization
- ○ DNA damage repair
- ☐ Change in state or activity
- ◼ Immune system or signal pathways related to macrophage function
- 🟩 Biological process
- 🟥 Canonical process

A

Intronic polyadenylation events (6h)

1 decreased usage

9 increased usage

PDU per Exon 6h

PDU per Exon 0h

B

Intronic polyadenylation events (24h)

30 decreased usage

45 increased usage

PDU per Exon 24h

PDU per Exon 0h

A

THP1 western blot

Time after PMA treatment: 0h, 1h, 6h, 18h, 24h

CFIm: CF Im68, CF Im25, CFIm59

FIP1

CPSF: CPSF160, CPSF100, Wdr33, CPSF73, CPSF30

Time after PMA treatment: 0h, 1h, 6h, 18h, 24h

CstF: CstF50, CstF77, CstF64, CstF64τ

CFIIm: Clp1, Pcf11

Others: Symplekin, PAP, β-actin

THP1 western blot quantification

B

Human primary monocytes

MCSF: 0 day, 5 days, 7 days

CFIm25 — CFIm complex
Fip1 — CPSF complex
CstF64 — CstF complex
Pcf-11 — CFIIm complex
β-actin

C

Cleavage polyadenylation factors — log₂ fold change over control

U937, THP-1, Human primary monocytes

Other: SYMPK
CFIIm: Pcf11, Clp1
CstF: CstF64, CstF77, CstF50
CPSF: CPSF30, CPSF73, WDR33, CPSF100, CPSF160, Fip1L1
CFIm: CFIm59, CFIm25, CFIm68

D

Primers for unspliced (US)   Primers spanning p(A) site

$\frac{US}{Span} \propto$ Processing efficiency

Processing efficiency

PMA-0h, PMA-6h, PMA-24h

MYC, ACTB — THP-1

A

Minus RT controls
for processing activity

B

U937 cells

A   **Genome browser map for <u>CFIm25</u>**

B   **Genome browser map for <u>CFIm68</u>**

C   **Genome browser map for <u>CstF64 tau</u>**