Macrophage differentiation is marked by increased abundance of the mRNA 3' end processing machinery, altered poly(A) site usage, and sensitivity to the level of CstF64

Srimoyee Mukherjee[1], Joel H. Graber[2], Claire L. Moore[1*]

[1]Department of Developmental, Molecular, and Chemical Biology, Tufts University School of
Medicine, Boston, Massachusetts 02111, USA

[2]Mount Desert Island Biological Laboratory, Bar Harbor, Maine, 04609, USA

* Correspondence:
Claire L. Moore
claire.moore@tufts.edu

Running title: Polyadenylation regulation in monocyte: macrophage differentiation

Keywords: polyadenylation, monocyte, macrophage, CstF64, CSTF2, differentiation

Abbreviations: CstF: Cleavage Stimulation Factor; CPSF: Cleavage and Polyadenylation Specificity
Factor; C/P: Cleavage and Polyadenylation; KD: Knockdown; OE: Overexpression; UTR:
Untranslated Region

Abstract

Regulation of mRNA polyadenylation is important for response to external signals and differentiation
in several cell types, and results in mRNA isoforms that vary in the amount of coding sequence or 3'
UTR regulatory elements. However, its role in differentiation of monocytes to macrophages has not
been investigated. Macrophages are key effectors of the innate immune system that help control
infection and promote tissue-repair. However, overactivity of macrophages contributes to
pathogenesis of many diseases. In this study, we show that macrophage differentiation is
characterized by shortening and lengthening of mRNAs in relevant cellular pathways. The
cleavage/polyadenylation (C/P) proteins increase during differentiation, suggesting a possible
mechanism for the observed changes in poly(A) site usage. This was surprising since higher C/P
protein levels correlate with higher proliferation rates in other systems, but monocytes stop dividing
after induction of differentiation. Depletion of CstF64, a C/P protein and known regulator of
polyadenylation efficiency, delayed macrophage marker expression, cell cycle exit, attachment, and
acquisition of structural complexity, and impeded shortening of mRNAs with functions relevant to
macrophage biology. Conversely, CstF64 overexpression increased use of promoter-proximal
poly(A) sites and caused the appearance of differentiated phenotypes in the absence of induction. Our
findings indicate that regulation of polyadenylation plays an important role in macrophage
differentiation.

1 Introduction

Macrophages are a critical component of the innate immune system that have a protective and beneficial role through the phagocytosis and killing of microbes, removal of dying cells and cellular debris, production of immune-regulatory molecules, and presentation of antigens to T-cells (1). Macrophages are derived from precursor cells called monocytes, which, in turn, are derived from myeloid progenitors. Monocytes undergo phenotypical and functional differentiation when they are recruited to tissues, where they differentiate into macrophages with altered morphological features and then engage in inflammatory or tissue remodeling processes (2). Robust macrophage function is critical for tissue repair and combating infection. However, inappropriate or overly robust activity contributes to the pathogenesis of diseases such as atherosclerosis, arthritis, asthma, lung fibrosis, type 2 diabetes, insulin resistance, inflammatory bowel disease, cancer and most recently, COVID-19 and the formation of intra-abdominal scar tissue (3-9).

Abnormal proliferation of macrophage progenitor cells in lieu of differentiation not only impairs normal macrophage responsibilities but can also lead to blood cancers such as acute myeloid leukemias (AML). A promising approach to treat AML is differentiation therapy, in which malignant cells are encouraged to differentiate into more mature forms using pharmacological agents. This approach has been remarkably successful for a small subset of AML cases involving a PML-retinoic acid receptor fusion (10, 11), and is a valuable alternative approach to killing cancer cells through cytotoxicity. To improve treatment of leukemias by forcing differentiation as well as to modulate macrophage function to help abate diseases influenced by macrophages, we must understand the critical molecular pathways that govern macrophage differentiation.

In many cell types, differentiation requires not only changes in transcription, but also pervasive changes in mRNA splicing and 3' end cleavage and polyadenylation (C/P). In the C/P step, pre-mRNA is cleaved at the poly(A) site and a poly(A) tail is added to the new end (12). Changes in polyadenylation play an increasingly appreciated role in regulation of gene expression by affecting protein output at many levels. For example, the efficiency at which a particular mRNA precursor is polyadenylated will affect the amount of mRNA available for translation. Through alternative polyadenylation (APA), genes with multiple poly(A) sites can produce distinct mRNA isoforms with different lengths of 3' untranslated regions (3' UTRs), which in turn affects nuclear export, translation, stability, association with stress granules, and localization in the cytoplasm (13-16). Alternate sequences in the 3' UTR can even facilitate co-translational protein complex formation (17) or act as a sink to bind miRNAs that down-regulate tumor suppressor mRNAs (18). Polyadenylation that occurs in an intron or less frequently an exon upstream of the 3' most exon affects the coding region, leading to expression of different protein isoforms (19, 20).

Many examples of APA have been documented in the immune system. For example, stimulation of human monocytes with lipopolysaccharide (LPS) and interferon-γ or activation of B cells and T cells leads to proliferation and a switch to upstream poly(A) sites (21, 22). A specific example of the importance of APA in immune cell differentiation is the switch to the upstream poly(A) site of the transcription factor NFATC1, which leads to NFATC1 accumulation and maturation of naïve T cells into effector T cells (23). Preferential utilization of the upstream poly(A) site of the immunoglobulin heavy chain gene in activated B cells is needed to switch to producing the secreted form of the protein instead of the membrane-attached form and is mediated by the CstF64 C/P protein (24-26). Increased levels of CstF64 and changes in APA have also been observed after treatment of RAW 264.7 mouse monocyte/macrophage-like cells with LPS (27), which activates these cells to produce an inflammatory response (28).

While the above studies have illuminated key aspects of mRNA 3' end processing in immune cell
regulation, the potential role of C/P proteins in differentiation of monocytes to macrophages, which
results in a dramatic change of cellular characteristics including attachment to surfaces, increased
size with extended filopodia, and cell cycle arrest, has not been explored. We hypothesized that the
changes in cell phenotypes that occur during macrophage differentiation will be accompanied by
changes in choice of poly(A) site, and that perturbation of the C/P machinery will affect the course of
this differentiation. In the current study, we show that this is indeed the case. There are significant
changes in APA when the monocytic U937 cell line was differentiated to form macrophages in vitro,
as well as an increase in the components of the C/P machinery. Genetic manipulation of the C/P
protein CstF64 led to altered APA and differentiation properties of monocytes. Overall, our findings
indicate an important role for regulation of polyadenylation in macrophage differentiation.

2   Results

2.1   Establishment of the monocyte-macrophage differentiation systems.

To study alternative polyadenylation in monocyte to macrophage differentiation, we used the well-
established monocytic cell lines U937 and THP-1 (29) as models. These cells can be differentiated to
macrophages by treatment with phorbol 12-myristate 13-acetate (PMA) (30) and are commonly used
to study macrophage differentiation and function. While the two cell lines have different origins,
with THP-1 being derived from a patient with Acute Myeloid Leukemia and U937 from a patient
with histiocytic lymphoma (29), they allow investigations using cells with a homogeneous genetic
background. After 6 hours of treatment with 30 nM of PMA, U937 cells are in a transition phase in
which they stop dividing and loosely adhere to the dish, with little change in morphology (Fig. 1A).
By 18-24 hours, almost all cells showed macrophage-like phenotypes including increased cell size,
strong adherence to the dish, and appearance of filopodia. Monocyte-macrophage differentiation is
associated with decreased expression of monocyte surface markers and increased expression of
macrophage markers (31). The classic macrophage surface markers such as CD16 (32), CD68 (33),
HLA-DRA (34) and CD38 (35) were considerably increased over the 24-hour time course and the
monocyte marker CD14 (36) had reduced expression in differentiated U937 cells (Fig. 1B). As
observed previously (37, 38), these cells also displayed increased attachment and no longer
proliferated after PMA treatment (Fig. 1C and 1D). THP-1 cells behaved similarly to the U937 cells
in terms of morphological changes, marker expression, attachment, and proliferation (Supplementary
Fig. S1A-D).

To further validate that differentiation was successful in our systems, we measured the mRNA levels
of genes important for macrophage differentiation and function at two critical time points in the
transition from monocytes to macrophages - 6 hours after induction when U937 and THP-1 cells
show some signs of differentiation and 24 hours, when differentiation is complete. Genes encoding
proteins such as the surface marker CD86, the adhesion molecule ICAM1, transcription factors
including JUN, FOS, NFKB1, and IKZF1, cytokines such as Oncostatin-M (OSM) and IL12A, all
increased in expression at 6 hours and some also at 24 hours after treating U937 and THP-1 cells
with PMA (Fig. 1E and Supplementary Fig. S1E).

2.2   Integrative analysis of poly(A) site mapping data identifies expression changes and 3′
UTR shortening and lengthening events during macrophage differentiation.

To determine the effects of PMA treatment on global poly(A) site usage and gene expression in U937
cells, we used Poly(A)-Click sequencing, which returns sequences adjacent to the poly(A) site and

has been used in recent studies to assess APA and expression changes (39-43). The differential
expression for a particular gene can be determined by collapsing all mapped poly(A) sites within
each gene into one count. As might be expected for monocytes undergoing differentiation, genes with
increased expression at 6h are strongly enriched in pathways involved in the immune system and
macrophage function (Supplementary Fig. S2A). Pathways associated with DNA repair were
predominant in the decreased expression set. At 24 hours and similar to what was seen at 6 hours, the
categories with significant increases in expression were heavily skewed towards the immune system
and pathways related to macrophage function (Supplementary Fig. S2B). The increased expression of
macrophage-related genes further confirms that our differentiation protocol is effective. The down-
regulated set at 24 hours also showed marked enrichment of genes involved in cell cycle and
chromosome organization. Decreased expression of these genes is consistent with the exit from the
cell cycle upon PMA treatment.

We also asked if there were changes in poly(A) site usage during differentiation. The degree of APA
usage was calculated by changes in the Percentage of Distal poly(A) site Usage Index (delta-PDUI),
which identifies lengthening (positive index) or shortening (negative index) of 3′ UTRs having
statistical significant difference between differentiated and control samples. For U937 cells
differentiated for 6 hours, sequencing identified polyadenylated mRNAs from a total of 12,208 genes
of which 331 genes had mRNAs with significant changes in length in the terminal 3' UTR of the 3'
most exon, designated as terminal 3' UTR. Of these, 139 exhibited an increase in shorter isoforms,
and 113 showed an increase in longer isoforms, while 79 showed a complex pattern involving
multiple poly(A) sites, where use of some went up and others went down (Fig. 2A). At 24 hours,
signals from 12,145 genes were obtained, and mRNAs from a greater number of genes (864) had
changes in poly(A) site usage compared to 6 hours. Of these, 376 exhibited shortening, 305 exhibited
lengthening and 183 showed a complex pattern (Fig. 2A). At both time points, the number of
significantly shortened 3' UTR APA events was slightly higher compared to the significantly
lengthened APA events. Some genes have poly(A) sites located upstream of the gene's last exon. Use
of these intronic sites creates a new terminal exon and changes the amount of coding sequence in the
mRNA. We also observed changes in intronic poly(A) site use, but at a lower frequency than changes
in the terminal 3' UTR. At 6 hours, 9 genes showed increased use of these poly(A) sites and one
showed decreased use, and at 24 hours, 45 genes showed an increase and 30 showed a decrease
(Supplementary Fig. S5A and B). Detailed outputs for all the APA events are supplied in
Supplementary Tables S3-S5.

Shortening of transcripts in the terminal 3' UTR has sometimes, but not always been associated with
increased mRNA expression, with the converse for mRNA lengthening (22, 44-49). To assess how
gene expression correlated with changes in terminal 3' UTR length during macrophage
differentiation, the 3' UTR shortened and lengthened gene sets were categorized into those with up-
regulated or down-regulated expression and delta-PDUI versus fold change in expression was plotted
for the 6 and 24 hour time points (Fig. 2B and Fig. 2C). A comparable number of genes in the
lengthened or shortened groups at both time points exhibited increases or decreases in mRNA
expression, and many genes with significant changes in isoform length showed no or only small
changes in expression, indicating a poor correlation between 3' UTR length and steady-state mRNA
level.

Previous work has shown that genes with significant changes in their 3' UTR isoform expression
ratios are cell type- and pathway-specific (48). To determine the biological pathways impacted by
APA during macrophage differentiation, we performed Gene Set Enrichment Analysis (GSEA) (50)
on genes which exhibited significant shortening or lengthening of transcripts. Analysis of the 0 vs 6h

shortened set showed an enrichment of categories related to immune system or signal pathways
related to macrophage function, as well as metabolism and cellular transport (Fig. 3A). Intracellular
transport is important for both cytokine secretion and phagocytosis, and modulation of metabolism is
recognized as a central player in macrophage activation (51). In contrast, the lengthened set was
primarily related to the immune system.

The affected pathways for the 0 vs 24h APA changes were quite different from those affected at 6
hours. Unlike what was observed for 6h, APA changes at 24h that involved shortening affected
pathways not specific to the immune system but nevertheless relevant to macrophage differentiation
such as cytoskeletal organization, cell cycle, and chromosome organization, while the lengthened set
was more mixed and included categories related to post-translational modification, cell cycle and
signaling pathways involved in macrophage-mediated inflammatory responses (Fig. 3B).

Examples of genome browser plots of the mapped poly(A) sites for genes which showed mRNA
shortening or lengthening in U937 cells are shown in Supplementary Fig. S3A for the 6 hour time
point (CIAPIN1 and EIF1) and in Fig. 4 for the 24 hour time point (shortened: SERPINB1 and
PTCH1; lengthened: PSAT1 and MRPL44). One way to confirm the presence of isoforms differing
in their 3' ends and shifts in use of these isoforms is 3' rapid amplification of cDNA ends (RACE)
assays (52-55). Thus, 3' RACE assays were used to detect the different mRNA isoforms, and these
confirmed the APA changes depicted by the genome browser plots from the poly(A)-sequencing data
of U937 cells (Fig. 5 and Supplementary Fig. S3B). SERPINB1 has 3 major poly (A) sites (PAC-1, 3
and 5 as shown in Fig. 4A), but the 3' RACE was not able to detect the longest isoform, perhaps
because of the strong bias towards amplifying the short isoform limited the sensitivity for longer
products. Nevertheless, in accordance with the genome browser plot, we did observe a reduction in
use of the middle p(A) site (marked as D in Fig. 4A). The differentiation of the other monocytic cell
line, THP-1 also produced shortening or lengthening of the same genes after 24 hours of PMA
treatment (Supplementary Fig. S4A and B).

2.3   Levels of cleavage/polyadenylation (C/P) complex proteins are upregulated at different
times during monocyte-to-macrophage differentiation.

The analysis of the poly(A)-Click seq data described above shows that the APA occurring during
monocyte-macrophage differentiation affected relevant functional categories of genes, and that
similar numbers of genes showed shortening or lengthening in the 3' UTR, with a slight trend
towards shortening. We next wanted to explore possible mechanisms for eliciting this APA and
examined the C/P complex proteins, as numerous studies have shown that changing the levels of
these proteins can affect poly(A) site choice (56, 57). The core C/P complex is comprised of six
factors: CPSF, CstF, CFI_m, $CFII_m$, symplekin and poly(A) polymerase (PAP) (Fig. 6B) [reviewed in
(58)]. The C/P proteins, with the exception of PAP, increased in level during differentiation of U937
cells, but with different kinetics and not necessarily in the same manner for the individual subunits of
each factor (Fig. 6A). For example, CstF50, CstF64 and CstF77 increased early at 1 hour after PMA
treatment, before differentiated phenotypes are observable. CPSF100, CPSF73 and CFI_m59 increased
at 6 hours when cells begin to attach, whereas CFI_m68, FIP1, CPSF160, Clp1, Pcf11, symplekin and
CstF64τ, a conserved paralog of CstF64 (59) in mammals, increased later (18-24 hours), when the
cells are mostly differentiated and attached. Some subunits, such as WDR33 and CstF77, decreased
when differentiation was completed. The C/P proteins increased in similar ways during
differentiation of the THP-1 monocytic cell line, with comparable kinetics (Supplementary Fig.
S6A). Overall, our findings show that C/P proteins increase during macrophage differentiation but
that some may be limiting in concentration at different points in the differentiation process.

2.4   Differentiation of monocytic cells increases processing efficiency at single poly(A) sites.

To test whether PMA-mediated differentiation alters the efficiency of 3'-end processing *in vivo*, we
determined the level of uncleaved transcripts that will contain sequence upstream and downstream of
a poly(A) site. We used previously described primers to monitor cleavage of ACTB or MYC
transcripts in vivo by RT-qPCR of total RNA reverse transcribed with random hexamers (60, 61).
ACTB has a single poly(A) site and MYC has a major poly(A) site used 93% of the time (62). One
primer pair detects RNA containing sequence that spans the poly(A) sites (Span) of the ACTB or
MYC genes. Because we expect uncleaved RNA to be nuclear, we compared this value to that
obtained using a primer pair which detects unspliced transcripts (US), which would also be nuclear.
The processing efficiency at each gene's poly(A) site can be quantified by the ratio of "US" PCR
product to that of "Span" product. If processing efficiency increases, fewer "Span" products are
expected relative to the "US" control Fig. 6C (61). The processing efficiency at each gene's poly(A)
site can be quantified by the ratio of US PCR product to that of Span product. Compared to
undifferentiated cells, more mRNA from both genes was processed at 24 hours, the time when most
of the U937 or THP-1 cells have differentiated (Fig. 6C and Supplementary Fig. S6B). Respective
"minus RT" controls gave insignificant signals (Supplementary Fig. S7A) indicating the effect to be
entirely coming from the transcripts and not DNA contamination. This data indicates that consistent
with the increase in C/P proteins, differentiation positively affected processing efficiency at these
ACTB and MYC poly(A) sites.

**2.5   Genetic manipulation of CstF64 alters monocyte-macrophage differentiation and switches
poly(A) site choice.**

As described above, monocyte differentiation led to the increased expression of C/P proteins and
altered APA of mRNAs from groups of genes known to be important for macrophage differentiation
and function. We thus wanted to test whether altering the level of one of these proteins could impact
the APA and macrophage differentiation. We focused on CstF64, because it has been shown to
influence APA during several types of transitions in cell state, in particular some of which involve
the immune system (21-23, 27, 63, 64). Based on these previous studies, we hypothesized that while
CstF64 is likely not to be the only C/P protein affecting APA in macrophage differentiation,
alteration of its levels would affect both APA and the progression of differentiation.

To test our hypothesis, we generated stable cell lines in which the expression of CstF64 in U937 cells
could be knocked down (KD) by doxycycline-inducible expression of shRNA against CstF64
mRNAs. Depletion of CstF64 was done by doxycycline treatment for 3 days and then PMA was
added for one day before harvesting the cells. Assessment of the extent of protein KD by two
different shRNA sequences against CstF64 showed considerable decrease in expression, indicating
efficient KD, while levels of the homolog CstF64τ remained unaffected (Fig. 7A). The macrophage
surface markers CD16, CD68, HLA-DRA and CD38 were substantially reduced compared to the
control shRNA when either CstF64 shRNA was used for KD. On the other hand, the levels of
monocyte marker CD14 increased on CstF64 KD (Fig. 7A).

The changes in expression of monocyte-macrophage markers suggested that CstF64 KD might also
affect the cellular changes that occur during macrophage differentiation. Indeed, counting of
suspended and attached cells after 24 hours of PMA treatment showed that CstF64 KD in U937 cells
significantly reduced cell attachment with both shRNAs (Fig. 7B, left panel). Normally by 24 hours,
the cells have stopped dividing but are viable. However, the total cell number at 24 hours was greater
in CstF64 KD cells compared to the control, indicating a slowing of the PMA-induced cell cycle

arrest (Fig. 7B, right panel). Macrophage differentiation is also characterized by the formation of
large, structurally complex cells (Fig. 1) (2), and this change can be detected using flow cytometry to
measure the side (SSC) and forward scattering (FSC) of live cells (65). An increase in intensities of
both FSC and SSC indicate the formation of cells with complex morphological features (also called
increased granularity or complexity) upon PMA-mediated differentiation (Supplementary Fig. S7B).
U937 cells depleted of CstF64 demonstrated reduced granularity (Fig. 7C).

Next, to determine if overexpression (OE) of CstF64 would promote macrophage differentiation, we
generated stably transfected U937 cells inducibly overexpressing this C/P protein. CstF64 was
efficiently overexpressed after 72 hours of doxycycline exposure, while CstF64τ levels did not
change (Fig. 7D). The cell proliferation rate was not affected by CstF64 OE (Fig. 7E, right panel).
However, overexpression increased expression of macrophage markers such as CD68, HLA-DRA
and CD38. and decreased CD14 expression (Fig. 7D). In addition, there was an increase in the
number of attached cells from 0 to 16% (Fig. 7E, left panel) and by FACS analysis, U937 cells
overexpressing CstF64 demonstrated an almost two-fold increase in granularity (Fig. 7F). It is
important to note that these cells were not treated with the differentiation inducer PMA.

In addition to these changes in cellular properties, 3' RACE analysis showed that CstF64 KD resulted
in increased usage of the distal poly(A) sites of the mRNAs *SERPINB1* and *PTCH1,* which would
otherwise be shortened after 24 hours of PMA treatment, whereas CstF64 OE without PMA
treatment caused decreased usage of the distal poly(A) sites of these mRNAs (Fig. 8). This finding is
consistent with studies in other systems showing that CstF64 promotes use of proximal poly(A) sites
(55, 66, 67).

In summary, depletion of CstF64 caused changes in cell properties indicative of poorer macrophage
differentiation and shifts to longer mRNA isoforms of genes which are normally shortened during
differentiation. In contrast, CstF64 overexpression caused the appearance of differentiated
phenotypes (cell surface markers, attachment, granularity, and transcript shortening) in the absence of
the differentiation inducer.

3   **Discussion**

Macrophage activity is needed for a robust response to infection and tissue damage but when
deregulated, it also contributes to many human diseases (3-7). As such, it is critical that monocyte to
macrophage differentiation is appropriately regulated to preserve the balance of various monocyte
and macrophage populations and ensure the appropriate functional consequences. The findings of our
study indicate that regulation of polyadenylation is one of the ways in which gene expression is
controlled during macrophage differentiation.

3.1   **Macrophage differentiation is characterized by both 3′ UTR shortening and lengthening**
events.

APA profiles are known to vary according to the differentiation and growth state of cells, with a
general shift to distal poly(A) sites (3' UTR lengthening) often observed with terminal differentiation
and decreased proliferation (12, 57, 68, 69). However, there are counter-examples to this
generalization: spermatogenesis and differentiation of secretory cells leads to a prevalence of shorter
mRNA isoforms (21, 70), and when B cells are transformed with Epstein-Barr virus, there is only a
slight preference for shortening (48). We found that upon macrophage differentiation, the distribution
of mRNA isoforms of many genes shifts as the monocyte precursor stops dividing and changes

morphology. There were similar numbers of 3' UTR shortening and lengthening events, indicating that exit from the cell cycle during macrophage differentiation is not associated with predominant lengthening of mRNA isoforms. GSEA analysis showed that lengthened and shortened gene sets are both enriched in groups of genes important for macrophage differentiation and function. We have validated several APA events by 3'RACE in both the U937 and THP-1 monocytic cell lines, and some of the genes examined by 3'RACE have functions relevant to macrophage biology. For example, *SERPINB1* has been reported to influence the innate immune system during influenza infection (71), *PTCH1* is important for macrophage chemotaxis during gastric inflammation (72), and *PSAT1* regulates the inflammatory response in macrophages (73). Our analysis shows that the APA profiles are highly dynamic during macrophage differentiation. Moreover, the APA changes are not globally affecting all genes with multiple poly(A) sites but target sets of genes whose functions are consistent with the need to exit the cell cycle and develop properties of a mature macrophage.

For many genes, 3' UTR shortening has been shown to increase mRNA and protein expression (44, 74, 75). However, other studies have demonstrated that global 3′-UTR shortening in variety of tissues and cell types has limited influence on mRNA or protein levels (47-49). Similar to these latter studies, our global analysis showed that the direction of APA change is not a good general predictor of mRNA expression levels during macrophage differentiation. It is likely that in this transition, the final levels of different mRNAs are determined by APA-dependent and APA-independent changes in mRNA stability as well as changes in transcription. Moreover, the function of some APA changes during macrophage differentiation may be to affect other mRNA properties such as translatability and localization rather than stability. In summary, our findings support the idea that APA is used as an additional mechanism to insure the timely expression and protein output of critical groups of genes during the transition of monocytes to macrophages.

3.2   The increase in C/P proteins may facilitate APA during macrophage differentiation.

Our study of the U937 and THP-1 monocytic cell lines shows that differentiating monocytic cells display an overall trend of increased expression of proteins of the core C/P machinery. While there were some differences which could be expected given the different origins of the two lines, the general features of shifts in poly(A) site usage and increase in C/P proteins are consistent between the two. This finding was unexpected, as many studies have shown that the expression of C/P mRNA and protein increases with increased proliferation (56, 66), but monocyte cell division ceases upon induction. The boosted expression of C/P proteins in mature macrophages was accompanied by an increase in the processing efficiency of genes with single poly(A) sites. The specific set of cellular changes accompanying macrophage differentiation, including increased mobility and metabolic activity, may necessitate an increased polyadenylation capacity.

Many studies have demonstrated that the core components of the mRNA 3′ processing machinery are also key regulators of global APA profiles. In macrophages compared to monocytes, we found increases in C/P proteins such as Fip1 and CstF64, known to promote use of upstream sites and production of mRNAs with shorter 3' UTRs (56, 63, 64, 66, 67). As described below, we showed that manipulation of CstF64 levels indeed altered APA in these cells. The increases in subunits of CFI_m, which is documented to promote use of downstream sites (74, 76-80) and Pcf11, whose effect can vary depending on gene length and cell type (81-83), could explain why we observe APA changes in both directions.

3.3 CstF64 levels alter monocyte-macrophage differentiation and switch poly(A) site choice.

In this study, we examined the role of CstF64 in macrophage differentiation. CstF64 is the
component of the core C/P complex that recognizes GU/U-rich signal sequences located downstream
of the poly(A) site, and it also facilitates APA in numerous cellular contexts. In a well-studied
example, an increase in CstF64 protein during terminal B-cell differentiation increases use of an
intronic poly(A) site in immunoglobulin heavy chain pre-mRNAs that changes the protein product
from a membrane-bound form to a secreted form (26, 63). CstF64 also promotes use of weaker
proximal poly(A) sites in proliferating fibroblasts (66), HeLa cells (67) and activated T cells (55),
and non-canonical poly(A) site usage (84) as well as processing of specific acetylcholinesterase
isoforms (85). Depletion of CstF64 suppressed proliferation of lung cancer cells and exogenous
expression promoted proliferation of HEK293 cells (86). CstF64 has also been shown to regulate the
cell cycle by controlling replication-dependent histone mRNA 3' end processing (87, 88), and CstF64
loss blocked the conversion of mouse embryonic stem cells into the endodermal lineage, but not into
ectoderm or mesoderm (89). The same study also showed that CstF64 was needed for cardiomyocyte
differentiation but not for neuronal differentiation (89), suggesting cell-specific requirements for
CstF64. Activation of RAW 264.7 macrophage-like cells with LPS blocked proliferation and caused
an increase in CstF64 but not other C/P subunits and an increase in proximal poly(A) site selection
(27). As these examples clearly demonstrate, CstF64 is a major player in APA that can promote
proliferation, differentiation or specific responses depending on the cell context. However, its role in
the differentiation of monocytes to macrophage, a process that involves cell cycle exit and striking
morphological changes, has not been previously studied.

In the current work, we found that CstF64 expression increased 6-8 fold during differentiation of
monocytic cell lines. Our subsequent analysis showed that CstF64 depletion not only caused
lengthening of transcripts that were otherwise shortened during differentiation but also had functional
consequences by slowing the acquisition of macrophage properties such as exit from the cell cycle,
attachment, marker expression and morphological changes. In contrast, overexpression of CstF64
stimulated macrophage-like changes and enhanced use of promoter proximal sites. Importantly, these
effects of extra CstF64 happened in the absence of the powerful inducer used to initiate
differentiation. Thus, our findings show that CstF64 is affecting pathways important for macrophage
differentiation. They are also in agreement with studies in other systems showing that CstF64
promotes use of promoter proximal sites. However, the increase in other C/P proteins and the
observation that some transcripts lengthen during differentiation suggests that CstF64 is not the only
driver and that other C/P factors and RNA-binding proteins with roles in APA regulation (12) will
also influence isoform expression in macrophages. While our study with CstF64 provides evidence
that regulation of polyadenylation is one of the mechanisms that promotes efficient macrophage
differentiation, further research is needed to determine the consortium of proteins that regulate APA
during this transition in cell state.

In summary, our study has provided new insights into the regulation of gene expression in the
differentiation of macrophages. The increase in levels of C/P proteins during differentiation and the
fact that manipulation of the C/P protein CstF64 alters the progression of this differentiation supports
a model in which changes in 3' end processing efficiency and alternative polyadenylation are
coordinated with changes in transcription and mRNA stability, and this regulatory network helps
create mature macrophages poised to respond appropriately to a variety of extracellular stimuli
through specific activities such as migration, phagocytosis, antigen presentation, and
immunomodulation. Given the critical role of macrophages in regulating the inflammatory response,
a greater understanding of how APA promotes beneficial and harmful macrophage activity could
open new therapeutic treatment avenues.

4   Materials and methods

4.1   Bacterial strains, DNA constructs and antibodies.

NEB® Stable Competent *E. coli* (Cat. #C3040) were used for transformation of plasmids. All
siRNAs and primers were purchased from Eton Biosciences (San Diego, CA). Inducible Lentiviral
shRNA constructs targeting CstF64 were purchased from GE Healthcare Dharmacon, Inc. (sequence
included in Supplementary Table S1). For overexpression (OE) of CstF64, we cloned CstF64 coding
sequence into pLIX_402 vector backbone (a gift from David Root, Broad Institute, MA) and
pLIX_402 vector (Addgene plasmid #41394) was used as control. Antibodies were purchased from
Santa Cruz, Bethyl and Abcam (Supplementary Table S2).

4.2   Cell culture, treatment and microscopy.

All cells were maintained in RPMI 1640 medium supplemented with 2 mM L-glutamine and 10%
heat-inactivated fetal bovine serum (FBS) at 37°C in a humidified atmosphere of 5% CO_2.
Differentiation of the U937 (ATCC #CRL-1593.2) and THP-1 (ATCC #TIB-202) human monocytic
cell-lines into macrophages was performed by treatment with 30 nM phorbol-12-myristate-13-acetate
(PMA; Sigma-Aldrich) for up to 24 hours. This PMA concentration has been shown in previous
studies to efficiently differentiate U937 cells (30, 90), and as described in the Results section, we
found it to be effective with both U937 and THP-1 cells. For monitoring morphological changes,
cells were cultured in 6-well plates at a density of 1×10^6 cells/well, after which they were examined
and photographed using an EVOS FL fluorescence phase contrast microscope (Thermo Fisher
Scientific). All results were obtained from three independent experiments.

4.3   Western blot analysis.

Western blot was performed with total cell extracts of untreated and differentiated cells. For
preparation of total cell extract, 1X RIPA buffer (20 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM
Na_2EDTA, 1 mM EGTA, 1% NP-40, 1% sodium deoxycholate, 2.5 mM sodium pyrophosphate, 1
mM β-glycerophosphate, 1 mM Na_3VO_4 and 1 µg/ml leupeptin) was added, followed by incubation
on ice for 15 min, centrifugation at 25,000 rcf for 10 minutes at 4°C and supernatant collection.
Protein concentration was measured by BCA reagent (Pierce, Thermo Fisher Scientific, Waltham,
MA), and 50-80 µg protein was separated on a 10% polyacrylamide-SDS or bis-tris gel and
transferred to PVDF membrane. The membrane was blocked for 1 h in 5% non-fat dried milk in
TBS-T (Tris Buffer Saline with Tween-20, *i.e.*, 0.05% Tween 20 in 1X TBS) buffer followed by
rocking overnight at 4°C with primary antibodies (antibody details provided in Supplementary Table
S2). Next day, unbound antibodies were removed by 4 X 5 min washes with TBS-T buffer. The
membrane was then incubated with an HRP-conjugated secondary antibody at a dilution of 1:5000
for 1 h at room temperature followed by 3 X 5 minute washes with TBS-T and a final wash with
TBS. The blot was developed with SuperSignal™ West Pico PLUS or Femto Chemiluminescent
Substrate (Thermo Fisher Scientific) and visualized with a ChemiDoc XRS+ System (Bio-Rad) and
quantitated using Image J (91). ACTB or GAPDH was used as the control for individual western
blots because their levels did not change upon differentiation. Pre-stained protein markers were used
as internal molecular mass standards, and each western blot was performed in three biological
replicates of each time point of the differentiation process.

4.4   RT-qPCR analysis to quantitate mRNA expression.

Total RNA extraction was carried out on $2x10^6$ cells using Trizol according to manufacturer's
protocol. 1.5 µg of RNA was subjected to reverse transcription using oligo dT primer and Superscript
III reverse transcriptase. The cDNA was amplified by qPCR with specific primers (Supplementary
Table S1) using the C1000™ thermal cycler with CFX96 Touch Real-Time PCR Detection System
(Bio-Rad Laboratories). The relative expression of genes was analyzed quantitatively by the $\Delta\Delta C_t$
method. ACTB RNA was used as the normalization control for RT-qPCR-based RNA expression
analyses because its level did not change upon differentiation according to the poly(A)-seq datasets.
Primers for RNA expression were designed at the first exon according to the annotation featured in
Ensembl (92).

4.5   RT-qPCR analysis for quantitating processing activity.

437 We used a previously published assay to measure the processing efficiency at the single poly(A) sites
438 of the MYC and ACTB genes (60, 61). This assay uses random hexamers for reverse transcription of
439 1 µg of total RNA. RT-qPCR is then performed with a primer pair upstream of the poly(A) site,
440 called "US" which measures the level of unspliced transcripts and another primer pair that spans the
441 poly(A) site, called "span". The sequences of these primers are provided in Supplementary Table S1.

4.6   Poly(A)-Click sequencing and analysis.

This was a paid service from Click-Seq Technologies, in which total RNAs isolated from control and
differentiated U937 cells were subjected to Poly(A) Click-sequencing as described (39, 40, 43).
Briefly, reverse-transcription with oligo-dT primer (unanchored) supplemented with small amounts
of spiked-in 3'-azido-nucleotides (AzVTPs) was performed to initiate cDNA synthesis from within
the poly(A) tail, ensuring stochastic termination upstream of the poly(A) tail. The resulting 3'-azido-
blocked cDNA molecules were purified, 'click-ligated' and amplified to generate a cDNA library.
These libraries were pooled and sequenced using the manufacturer's standard operating procedures
on a NextSeq 550, obtaining 1×150 bp (130M) paired-end reads. Raw data from the sequencing
reactions for four biological replicates for 0h vs 24h and two biological replicates for 0h vs 6h was
analyzed by Click-Seq Technologies using Differential Poly(A)-Clustering (DPAC), a pipeline to
pre-process raw poly(A)-tail targeted RNAseq data, map to a reference genome, identify and annotate
the location of poly(A) sites, generate poly(A)-clusters (PACs) and determine the differential
abundance of PACs and differential gene expression between two conditions (40). The raw reads and
mapped reads for each library are provided in Supplementary Table S2 (sheets 1 and 2). Each of
these libraries were trimmed and quality filtered as described (43). Gene counts collapse all mapped
poly(A) sites within that one gene into one count, this provides differential gene expression data. On
average, 87% of all the mapped reads were found within or 200 nucleotides downstream of annotated
mRNAs. Multiple poly(A) sites occurring within 25 nucleotides of one another were grouped into a
poly(A) cluster (PAC) and treated as a unit to determine alternative polyadenylation events within
exons and introns. The cutoff to be considered a PAC is 10 counts. In the final stage of DPAC, the
mapped reads from each individual samples are used to determine the frequency of PACs in each
dataset by determining the overlaps of the 3' ends of the mapped reads with the provided poly(A)
cluster database using bedtools (93). Data normalization and statistical tests are applied using the
canonical DESeq2 (94) pipeline using local dispersion estimation and Independent Hypothesis
Weighting (IHW) (95) to estimate false discovery rates and power maximization. The final output of
DPAC for the 6h and 24h PMA-treated U937 samples are provided in Supplemental Tables S4 and
S5.

Principal component analysis (PCA) comparing control and PMA-treated U937 cells (0h vs 6h and
0h vs 24h) was done with the help of the DESeq2 pipeline (94) to determine and identify outlying
samples within datasets. Variables (i.e. the times of differentiation) are transformed into orthogonal
principal components where each principal component indicates the direction with the greatest
variability within the data. PCA plots (Supplementary Fig. S8A, B) revealed that treatment time was
the most critical variable. Sequencing data sets are uploaded to NIH-GEO with the accession number
#GSE169140. Volcano plots illustrating changes in gene counts and PAC counts (Supplementary
Fig. S8C, D) were generated for both 6h and 24h PMA treated U937 cells by plotting the $\log_2$ fold
change values vs p_{Adj} value. Differential Gene or PAC usage is defined as >1.5 fold change with p_{Adj}
<0.1. The volcano plots indicate that the number of differentially expressed genes and the number of
PACs showing change in usage increases as differentiation progresses. Gene Set Enrichment
Analysis (GSEA) was carried out using online resources from the Broad Institute (50). Poly(A) site
locations for genes were extracted into BED files and visualized with the help of UCSC genome
browser (96).

4.7   3' RACE.

We modified the previously published 3'RACE protocols for this study (97, 98). Briefly, total RNA
extraction was done using Trizol according to manufacturer's protocol. 1 µg of RNA was subjected
to reverse transcription for 1 hour using 100 units of Superscript IV reverse transcriptase and an
oligo(dT)-containing adapter primer to poly(A) mRNA. After cDNA synthesis, the RNA template is
removed from the cDNA:RNA hybrid molecule by digestion with 2 units of RNase H after cDNA
synthesis to increase the sensitivity of PCR. For purification, the prepared cDNA was then
chloroform extracted and precipitated in 0.3M sodium acetate and 3 volumes of ethanol.
Subsequently, cDNA was subjected to PCR reaction with HotStarTaq Master Mix Kit (Qiagen)
utilizing gene-specific forward primer (designed at the junction of the last two exons) and a unique
anchor primer complimentary to the oligo(dT)-containing adapter primer (sequences provided in
Supplementary Table S1). The first primer extension was accomplished by incubation at annealing
temperature for 1 min, followed by incubation at 72°C for 40 min. Exponential amplification was
accomplished by 30-35 cycles of PCR at 94°C for 1 min, 55°C /60°C (3-5 degrees lower than the
lowest primer T_m) for 30 sec and 72°C for 5 min, followed by an additional 10-min extension at
72°C. The products were resolved in 1% agarose gels along with 1 kb Plus DNA Ladder (New
England Biolabs) and visualized using SYBR™ Safe DNA Gel Stain (Thermo Fisher) with a
ChemiDoc XRS+ System (Bio-Rad) and quantitated using Image J (91).

4.8   Lentivirus construction and cell treatment.

For shRNA-based KD, three different lentiviral SMARTvector constructs, designed for doxycycline-
inducible expression of shRNA for human CstF64 gene silencing, were purchased from GE
Healthcare Dharmacon, Inc. These vectors incorporate the Tet-On® 3G bipartite system to inducibly
control of shRNA with the Tet 3G promoter (PTRE3G), while the expression of the Tet-On® 3G
transactivator protein is under the control of a constitutive human EF-1 promoter for optimal
expression in the monocyte-macrophage system. A SMART vector inducible lentiviral control
shRNA was used to control for potential non-specific effects caused by expression of shRNA. The
sequences of the shRNAs are in Supplementary Table S1. As shown in the Results, shRNA #2 & 3
gave efficient KD but shRNA #1 was not effective. To generate stable cells inducibly overexpressing
CstF64, its coding sequence was amplified by PCR from U937 cDNA and cloned into the pLIX_402
inducible lentiviral expression vector by using NheI and AgeI restriction enzyme sites to generate the
OE construct. The primers used for cloning are mentioned in Supplementary Table S1.

These recombinant plasmids were co-transfected with the components of Dharmacon™ Trans-
Lentiviral packaging kit into HEK293FT cells using FuGENE® HD (Promega Corp.) according to
manufacturer's protocol. Transfection of HEK293FT cells was done in 6-well plates when the cells
were 80-85% confluent, and transfection media changed after 16 hours. The recombinant lentiviruses
were harvested at 48 h post-transfection, spun at 1250 rpm for 5 minutes and filtered by a 0.45 µm
filter to remove cells and debris. Purified virus was used for infecting monocytic cells. $1.5X10^6$ target
cells were seeded in 4 ml of media per well in a 6-well plate and cultured overnight. Lentiviral
particles (1-2 ml per well) were added the next day to the cells in culture media containing 10 µg/ml
polybrene for efficient infection. Selection of cells stably expressing CstF64-shRNAs or control-
shRNA and pLIX_402 overexpressing CstF64 or control pLIX_402 started 72 h post-transfection or
a time determined by the expression of GFP from the cells. Growth medium was replaced with fresh
selection medium containing 1 µg/mL of puromycin. Puromycin-containing medium was refreshed
every 2–3 days, and selection was completed after approximately 1 week, after which clones were
expanded for 2 more weeks and frozen for later use. The Tet-On 3G induction system was activated
by 3 nM doxycycline for 3 days for both KD and OE transfections. Additionally for KD clones, 30
nM PMA was added one day prior to harvesting for RNA and protein extraction.

4.9   Attachment assay.

U937 and THP-1 cells were either untreated or treated with PMA and left for 24 hours to attach.
After that, unbound cells were collected from the plate by washing gently with PBS and collected for
counting (denoted as "suspended"). The attached cells were collected by trypsinizing to release them
from the plate. Cells were then stained with trypan blue and counted in a hemocytometer and
represented as percentage of suspended or attached cells over total (suspended + attached). The
figures are average of three independent observations.

4.10 Flow cytometry.

Phenotypic analysis of monocytes and macrophages was performed using flow cytometry. After
differentiation, cells were recovered from culture plates and washed with phosphate-buffered saline
(PBS) containing 1% paraformaldehyde for fixation. In order to avoid non-specific binding and
background fluorescence of cells expressing high levels of Fc receptors, the cells were incubated with
Fc Receptor Blocking Solution (Human TruStain FcX, Biolegend) according to the manufacturer's
instructions for 10 minutes, followed by washing with cold PBS. Zombie Aqua™ Fixable Viability
kit (BioLegend) at a final dilution of 1:2000 was used for the exclusion of dead cells. Thereafter,
cells were analyzed with a LSR II flow cytometer (BD Biosciences) using FACSDiva software, and
data were analyzed with the FlowJo software. Primary gating was done on the basis of forward
versus side scatter (FSC vs SSC) to compare size and granularity (complexity) of live cells across
samples (31).

4.11 Statistical analysis.

All experiments were performed in at least three independent sets. Data are presented as mean $\pm$ SE.
Statistical analysis was performed using GraphPad Prism 6.01 (GraphPad Software Inc., La Jolla,
CA, USA). Two-way analysis of variance (ANOVA) was performed to determine the significance
between the groups. Considerations were $* = P \leq 0.05$; $** = P \leq 0.01$; $*** = P \leq 0.001$; $**** = P \leq$
0.0001. A P value <0.05 was considered significant.

4.12 Materials availability.

All plasmids and cell lines will be available upon request.

5   Conflict of Interest

559 The authors declare that the research was conducted in the absence of any commercial or financial
560 relationships that could be construed as a potential conflict of interest.

6   Author Contributions

562 CM and SM conceived the study and CM provided general oversight. SM developed the strategy and
563 methodology, acquired the data, reported and organized the findings. CM and SM designed the
564 experiments, interpreted the results, and wrote and revised the manuscript. JG contributed to the
565 computational analyses.

7   Funding

567 This work was supported by the National Institutes of Health [grant numbers 1R01GM101010 and
568 1R01AI152337] to Claire Moore and partially supported by an Institutional Development Award
569 (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health
570 [under grant numbers P20GM103423 and P20GM104318] to Mount Desert Island Biological
571 Laboratory.

8   Acknowledgments

We thank Tufts University faculty Alexei Degterev and Karl Munger for providing cell lines and
Marta Rodriguez-Garcia for consulting with us on FACS analysis. We thank members of Karl
Munger's lab for use of the cell culture facility and want to acknowledge the critical inputs of the
members of FACS core, Tufts University, and other members of the Moore lab. We would also like
to thank Andrew Routh and Elizabeth Jaworski of ClickSeq Technologies for their valuable input on
the bioinformatic analysis, and Nathaniel Maki and Chris Wilson of Mount Desert Island Biological
Laboratory for helping perform the alignments for RNA-seq analysis.

9   References

1.   Shi C, Pamer EG. Monocyte recruitment during infection and inflammation. Nat Rev
Immunol. 2011;11(11):762-74.

2.   Italiani P, Boraschi D. From Monocytes to M1/M2 Macrophages: Phenotypical vs. Functional
Differentiation. Front Immunol. 2014;5:514.

3.   Mammana S, Fagone P, Cavalli E, Basile MS, Petralia MC, Nicoletti F, et al. The Role of
Macrophages in Neuroinflammatory and Neurodegenerative Pathways of Alzheimer's Disease,
Amyotrophic Lateral Sclerosis, and Multiple Sclerosis: Pathogenetic Cellular Effectors and Potential
Therapeutic Targets. Int J Mol Sci. 2018;19(3).

4.   Salmaninejad A, Valilou SF, Soltani A, Ahmadi S, Abarghan YJ, Rosengren RJ, et al.
Tumor-associated macrophages: role in cancer development and therapeutic implications. Cell Oncol
(Dordr). 2019;42(5):591-608.

5.   Gomez-Rial J, Rivero-Calle I, Salas A, Martinon-Torres F. Role of Monocytes/Macrophages
in Covid-19 Pathogenesis: Implications for Therapy. Infect Drug Resist. 2020;13:2485-93.

6.   Punzo F, Bellini G, Tortora C, Pinto DD, Argenziano M, Pota E, et al. Mifamurtide and
TAM-like macrophages: effect on proliferation, migration and differentiation of osteosarcoma cells.
Oncotarget. 2020;11(7):687-98.

7.   Zindel J, Peiseler M, Hossain M, Deppermann C, Lee WY, Haenni B, et al. Primordial
GATA6 macrophages function as extravascular platelets in sterile injury. Science. 2021;371(6533).

8.   Zhou X, Li W, Wang S, Zhang P, Wang Q, Xiao J, et al. YAP Aggravates Inflammatory
Bowel Disease by Regulating M1/M2 Macrophage Polarization and Gut Microbial Homeostasis. Cell
Rep. 2019;27(4):1176-89 e5.

9.   Labonte AC, Tosello-Trampont AC, Hahn YS. The role of macrophage polarization in
infectious and inflammatory diseases. Mol Cells. 2014;37(4):275-85.

10.   de The H. Differentiation therapy revisited. Nat Rev Cancer. 2018;18(2):117-27.

11.   Sykes DB, Kfoury YS, Mercier FE, Wawer MJ, Law JM, Haynes MK, et al. Inhibition of
Dihydroorotate Dehydrogenase Overcomes Differentiation Blockade in Acute Myeloid Leukemia.
Cell. 2016;167(1):171-86 e15.

12.   Neve J, Patel R, Wang Z, Louey A, Furger AM. Cleavage and polyadenylation: Ending the
message expands gene regulation. RNA Biol. 2017;14(7):865-90.

13.   Cheng LC, Zheng D, Zhang Q, Guvenek A, Cheng H, Tian B. Alternative 3' UTRs play a
widespread role in translation-independent mRNA association with the endoplasmic reticulum. Cell
Rep. 2021;36(3):109407.

14.   Muller-McNicoll M, Botti V, de Jesus Domingues AM, Brandl H, Schwich OD, Steiner MC,
et al. SR proteins are NXF1 adaptors that link alternative RNA processing to mRNA export. Genes
Dev. 2016;30(5):553-66.

15.   Neve J, Burger K, Li W, Hoque M, Patel R, Tian B, et al. Subcellular RNA profiling links
splicing and nuclear DICER1 to alternative cleavage and polyadenylation. Genome Res.
2016;26(1):24-35.

16.   Zheng D, Wang R, Ding Q, Wang T, Xie B, Wei L, et al. Cellular stress alters 3'UTR
landscape through alternative polyadenylation and isoform-specific degradation. Nat Commun.
2018;9(1):2268.

17.   Berkovits BD, Mayr C. Alternative 3' UTRs act as scaffolds to regulate membrane protein
localization. Nature. 2015;522(7556):363-7.

18.   Park HJ, Ji P, Kim S, Xia Z, Rodriguez B, Li L, et al. 3' UTR shortening represses tumor-
suppressor genes in trans by disrupting ceRNA crosstalk. Nat Genet. 2018;50(6):783-9.

19.   Wang R, Zheng D, Yehia G, Tian B. A compendium of conserved cleavage and
polyadenylation events in mammalian genes. Genome Res. 2018;28(10):1427-41.

20.   MacDonald CC. Tissue-specific mechanisms of alternative polyadenylation: Testis, brain, and
beyond (2018 update). Wiley Interdiscip Rev RNA. 2019;10(4):e1526.

21.   Cheng LC, Zheng D, Baljinnyam E, Sun F, Ogami K, Yeung PL, et al. Widespread transcript
shortening through alternative polyadenylation in secretory cell differentiation. Nat Commun.
2020;11(1):3182.

22.   Sandberg R, Neilson JR, Sarma A, Sharp PA, Burge CB. Proliferating cells express mRNAs
with shortened 3' untranslated regions and fewer microRNA target sites. Science.
2008;320(5883):1643-7.

23.   Chuvpilo S, Zimmer M, Kerstan A, Glockner J, Avots A, Escher C, et al. Alternative
polyadenylation events contribute to the induction of NF-ATc in effector T cells. Immunity.
1999;10(2):261-9.

24.   Takagaki Y, Manley JL. Levels of polyadenylation factor CstF-64 control IgM heavy chain
mRNA accumulation and other events associated with B cell differentiation. Mol Cell.
1998;2(6):761-71.

25.   Edwalds-Gilbert G, Milcarek C. Regulation of poly(A) site use during mouse B-cell
development involves a change in the binding of a general polyadenylation factor in a B-cell stage-
specific manner. Mol Cell Biol. 1995;15(11):6420-9.

26.   Peterson ML. Immunoglobulin heavy chain gene regulation through polyadenylation and
splicing competition. Wiley Interdiscip Rev RNA. 2011;2(1):92-105.

27.   Shell SA, Hesse C, Morris SM, Jr., Milcarek C. Elevated levels of the 64-kDa cleavage
stimulatory factor (CstF-64) in lipopolysaccharide-stimulated macrophages influence gene
expression and induce alternative poly(A) site selection. The Journal of biological chemistry.
2005;280(48):39950-61.

28.   Schutte RJ, Parisi-Amon A, Reichert WM. Cytokine profiling using monocytes/macrophages
cultured on common biomaterials with a range of surface chemistries. J Biomed Mater Res A.
2009;88(1):128-39.

29.   Chanput W, Peters V, Wichers H. THP-1 and U937 Cells. In: Verhoeckx K, Cotter P, Lopez-
Exposito I, Kleiveland C, Lea T, Mackie A, et al., editors. The Impact of Food Bioactives on Health:
in vitro and ex vivo models. Cham (CH)2015. p. 147-59.

30.   Mukherjee S, Sengupta Bandyopadhyay S. Phorbol-12-myristate-13-acetate (PMA) mediated
transcriptional regulation of Oncostatin-M. Cytokine. 2016;88:209-13.

31.   Daigneault M, Preston JA, Marriott HM, Whyte MK, Dockrell DH. The identification of
markers of macrophage differentiation in PMA-stimulated THP-1 cells and monocyte-derived
macrophages. PloS one. 2010;5(1):e8668.

32.   Wang ZQ, Bapat AS, Rayanade RJ, Dagtas AS, Hoffmann MK. Interleukin-10 induces
macrophage apoptosis and expression of CD16 (FcgammaRIII) whose engagement blocks the cell
death programme and facilitates differentiation. Immunology. 2001;102(3):331-7.

33.   Holness CL, Simmons DL. Molecular cloning of CD68, a human macrophage marker related
to lysosomal glycoproteins. Blood. 1993;81(6):1607-13.

34.   Costabel U, Bross KJ, Andreesen R, Matthys H. HLA-DR antigens on human macrophages
from bronchoalveolar lavage fluid. Thorax. 1986;41(4):261-5.

35.   Amici SA, Young NA, Narvaez-Miranda J, Jablonski KA, Arcos J, Rosas L, et al. CD38 Is
Robustly Induced in Human Macrophages and Monocytes in Inflammatory Conditions. Front
Immunol. 2018;9:1593.

36.   Ziegler-Heitbrock HW, Ulevitch RJ. CD14: cell surface receptor and differentiation marker.
Immunol Today. 1993;14(3):121-5.

37.   Chang MY, Huang DY, Ho FM, Huang KC, Lin WW. PKC-dependent human monocyte
adhesion requires AMPK and Syk activation. PLoS One. 2012;7(7):e40999.

38.   Prasad A, Sedlarova M, Balukova A, Ovsii A, Rac M, Krupka M, et al. Reactive Oxygen
Species Imaging in U937 Cells. Front Physiol. 2020;11:552569.

39.   Routh A, Ji P, Jaworski E, Xia Z, Li W, Wagner EJ. Poly(A)-ClickSeq: click-chemistry for
next-generation 3-end sequencing without RNA enrichment or fragmentation. Nucleic Acids Res.
2017;45(12):e112.

40.   Routh A. DPAC: A Tool for Differential Poly(A)-Cluster Usage from Poly(A)-Targeted
RNAseq Data. G3 (Bethesda). 2019;9(6):1825-30.

41.   Cao J, Verma SK, Jaworski E, Mohan S, Nagasawa CK, Rayavara K, et al. RBFOX2 is
critical for maintaining alternative polyadenylation patterns and mitochondrial health in rat
myoblasts. Cell Rep. 2021;37(5):109910.

42.   Chu Y, Elrod N, Wang C, Li L, Chen T, Routh A, et al. Nudt21 regulates the alternative
polyadenylation of Pak1 and is predictive in the prognosis of glioblastoma patients. Oncogene.
2019;38(21):4154-68.

43.   Elrod ND, Jaworski EA, Ji P, Wagner EJ, Routh A. Development of Poly(A)-ClickSeq as a
tool enabling simultaneous genome-wide poly(A)-site identification and differential expression
analysis. Methods. 2019;155:20-9.

44.   Mayr C, Bartel DP. Widespread shortening of 3'UTRs by alternative cleavage and
polyadenylation activates oncogenes in cancer cells. Cell. 2009;138(4):673-84.

45.   Ji Z, Luo W, Li W, Hoque M, Pan Z, Zhao Y, et al. Transcriptional activity regulates
alternative cleavage and polyadenylation. Mol Syst Biol. 2011;7:534.

46.   Chang JW, Zhang W, Yeh HS, de Jong EP, Jun S, Kim KH, et al. mRNA 3'-UTR shortening
is a molecular signature of mTORC1 activation. Nat Commun. 2015;6:7218.

47.   Gruber AR, Martin G, Muller P, Schmidt A, Gruber AJ, Gumienny R, et al. Global 3' UTR
shortening has a limited effect on protein abundance in proliferating T cells. Nat Commun.
2014;5:5465.

48.   Lianoglou S, Garg V, Yang JL, Leslie CS, Mayr C. Ubiquitously transcribed genes use
alternative polyadenylation to achieve tissue-specific expression. Genes Dev. 2013;27(21):2380-96.

49.   Spies N, Burge CB, Bartel DP. 3' UTR-isoform choice has limited influence on the stability
and translational efficiency of most mRNAs in mouse fibroblasts. Genome Res. 2013;23(12):2078-
90.

50.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
Proc Natl Acad Sci U S A. 2005;102(43):15545-50.

51.   Stunault MI, Bories G, Guinamard RR, Ivanov S. Metabolism Plays a Key Role during
Macrophage Activation. Mediators Inflamm. 2018;2018:2426138.

52.   Graber JH, Nazeer FI, Yeh PC, Kuehner JN, Borikar S, Hoskinson D, et al. DNA damage
induces targeted, genome-wide variation of poly(A) sites in budding yeast. Genome Res.
2013;23(10):1690-703.

53.   Masamha CP, Xia Z, Peart N, Collum S, Li W, Wagner EJ, et al. CFIm25 regulates
glutaminase alternative terminal exon definition to modulate miR-23 function. RNA.
2016;22(6):830-8.

54.   Venkat S, Tisdale AA, Schwarz JR, Alahmari AA, Maurer HC, Olive KP, et al. Alternative
polyadenylation drives oncogenic gene expression in pancreatic ductal adenocarcinoma. Genome
Res. 2020;30(3):347-60.

55.   Chatrikhi R, Mallory MJ, Gazzara MR, Agosto LM, Zhu WS, Litterman AJ, et al. RNA
Binding Protein CELF2 Regulates Signal-Induced Alternative Polyadenylation by Competing with
Enhancers of the Polyadenylation Machinery. Cell Rep. 2019;28(11):2795-806 e3.

56.   Di Giammartino DC, Nishida K, Manley JL. Mechanisms and consequences of alternative
polyadenylation. Mol Cell. 2011;43(6):853-66.

57.   Ren F, Zhang N, Zhang L, Miller E, Pu JJ. Alternative Polyadenylation: a new frontier in post
transcriptional regulation. Biomark Res. 2020;8(1):67.

58.   Shi Y, Manley JL. The end of the message: multiple protein-RNA interactions define the
mRNA polyadenylation site. Genes Dev. 2015;29(9):889-97.

59.   Dass B, Tardif S, Park JY, Tian B, Weitlauf HM, Hess RA, et al. Loss of polyadenylation
protein tauCstF-64 causes spermatogenic defects and male infertility. Proc Natl Acad Sci U S A.
2007;104(51):20374-9.

60.   Davidson L, Muniz L, West S. 3' end formation of pre-mRNA and phosphorylation of Ser2
on the RNA polymerase II CTD are reciprocally coupled in human cells. Genes Dev.
2014;28(4):342-56.

61.   Eaton JD, Davidson L, Bauer DLV, Natsume T, Kanemaki MT, West S. Xrn2 accelerates
termination by RNA polymerase II, which is underpinned by CPSF73 activity. Genes Dev.
2018;32(2):127-39.

62.   Herrmann CJ, Schmidt R, Kanitz A, Artimo P, Gruber AJ, Zavolan M. PolyASite 2.0: a
consolidated atlas of polyadenylation sites from 3' end sequencing. Nucleic Acids Res.
2020;48(D1):D174-D9.

63.   Takagaki Y, Seipelt RL, Peterson ML, Manley JL. The polyadenylation factor CstF-64
regulates alternative processing of IgM heavy chain pre-mRNA during B cell differentiation. Cell.
1996;87(5):941-52.

64.   Lackford B, Yao C, Charles GM, Weng L, Zheng X, Choi EA, et al. Fip1 regulates mRNA
alternative polyadenylation to promote stem cell self-renewal. EMBO J. 2014;33(8):878-89.

65.   Druszczynska M, Wlodarczyk M, Janiszewska-Drobinska B, Kielnierowski G, Zawadzka J,
Kowalewicz-Kulbat M, et al. Monocyte signal transduction receptors in active and latent
tuberculosis. Clin Dev Immunol. 2013;2013:851452.

66.   Mitra M, Johnson EL, Swamy VS, Nersesian LE, Corney DC, Robinson DG, et al.
Alternative polyadenylation factors link cell cycle to migration. Genome Biol. 2018;19(1):176.

67.   Yao C, Biesinger J, Wan J, Weng L, Xing Y, Xie X, et al. Transcriptome-wide analyses of
CstF64-RNA interactions in global regulation of mRNA alternative polyadenylation. Proc Natl Acad
Sci U S A. 2012;109(46):18773-8.

68.   Ji Z, Lee JY, Pan Z, Jiang B, Tian B. Progressive lengthening of 3' untranslated regions of
mRNAs by alternative polyadenylation during mouse embryonic development. Proc Natl Acad Sci U
S A. 2009;106(17):7028-33.

69.   Ji Z, Tian B. Reprogramming of 3' untranslated regions of mRNAs by alternative
polyadenylation in generation of pluripotent stem cells from different cell types. PLoS One.
2009;4(12):e8419.

70. Li W, Park JY, Zheng D, Hoque M, Yehia G, Tian B. Alternative cleavage and polyadenylation in spermatogenesis connects chromatin regulation with post-transcriptional control. BMC Biol. 2016;14:6.

71. Gong D, Farley K, White M, Hartshorn KL, Benarafa C, Remold-O'Donnell E. Critical role of serpinB1 in regulating inflammatory responses in pulmonary influenza infection. J Infect Dis. 2011;204(4):592-600.

72. Chakrabarti J, Dua-Awereh M, Schumacher M, Engevik A, Hawkins J, Helmrath MA, et al. Sonic Hedgehog acts as a macrophage chemoattractant during regeneration of the gastric epithelium. NPJ Regen Med. 2022;7(1):3.

73. Raines LN, Zhao H, Wang Y, Chen HY, Gallart-Ayala H, Hsueh PC, et al. PERK is a critical metabolic hub for immunosuppressive function in macrophages. Nat Immunol. 2022;23(3):431-45.

74. Masamha CP, Xia Z, Yang J, Albrecht TR, Li M, Shyu AB, et al. CFIm25 links alternative polyadenylation to glioblastoma tumour suppression. Nature. 2014;510(7505):412-6.

75. Jenal M, Elkon R, Loayza-Puch F, van Haaften G, Kuhn U, Menzies FM, et al. The poly(A)-binding protein nuclear 1 suppresses alternative cleavage and polyadenylation sites. Cell. 2012;149(3):538-53.

76. Martin G, Gruber AR, Keller W, Zavolan M. Genome-wide analysis of pre-mRNA 3' end processing reveals a decisive role of human cleavage factor I in the regulation of 3' UTR length. Cell Rep. 2012;1(6):753-63.

77. Li W, You B, Hoque M, Zheng D, Luo W, Ji Z, et al. Systematic profiling of poly(A)+ transcripts modulated by core 3' end processing and splicing factors reveals regulatory rules of alternative cleavage and polyadenylation. PLoS Genet. 2015;11(4):e1005166.

78. Brumbaugh J, Di Stefano B, Wang X, Borkent M, Forouzmand E, Clowers KJ, et al. Nudt21 Controls Cell Fate by Connecting Alternative Polyadenylation to Chromatin Signaling. Cell. 2018;172(1-2):106-20 e21.

79. Zhu ZJ, Huang P, Chong YX, Kang LX, Huang X, Zhu ZX, et al. MicroRNA-181a promotes proliferation and inhibits apoptosis by suppressing CFIm25 in osteosarcoma. Mol Med Rep. 2016;14(5):4271-8.

80. Yang SW, Li L, Connelly JP, Porter SN, Kodali K, Gan H, et al. A Cancer-Specific Ubiquitin Ligase Drives mRNA Alternative Polyadenylation by Ubiquitinating the mRNA 3' End Processing Complex. Mol Cell. 2020;77(6):1206-21 e7.

81. Wang R, Zheng D, Wei L, Ding Q, Tian B. Regulation of Intronic Polyadenylation by PCF11 Impacts mRNA Expression of Long Genes. Cell Rep. 2019;26(10):2766-78 e6.

82. Ogorodnikov A, Levin M, Tattikota S, Tokalov S, Hoque M, Scherzinger D, et al. Transcriptome 3'end organization by PCF11 links alternative polyadenylation to formation and neuronal differentiation of neuroblastoma. Nat Commun. 2018;9(1):5331.

83. Turner RE, Henneken LM, Liem-Weits M, Harrison PF, Swaminathan A, Vary R, et al. Requirement for cleavage factor IIm in the control of alternative polyadenylation in breast cancer cells. RNA. 2020;26(8):969-81.

84. Hwang HW, Park CY, Goodarzi H, Fak JJ, Mele A, Moore MJ, et al. PAPERCLIP Identifies MicroRNA Targets and a Role of CstF64/64tau in Promoting Non-canonical poly(A) Site Usage. Cell Rep. 2016;15(2):423-35.

85.   Nazim M, Masuda A, Rahman MA, Nasrin F, Takeda JI, Ohe K, et al. Competitive regulation
of alternative splicing and alternative polyadenylation by hnRNP H and CstF64 determines
acetylcholinesterase isoforms. Nucleic Acids Res. 2017;45(3):1455-68.

86.   Aragaki M, Takahashi K, Akiyama H, Tsuchiya E, Kondo S, Nakamura Y, et al.
Characterization of a cleavage stimulation factor, 3' pre-RNA, subunit 2, 64 kDa (CSTF2) as a
therapeutic target for lung cancer. Clin Cancer Res. 2011;17(18):5889-900.

87.   Youngblood BA, Grozdanov PN, MacDonald CC. CstF-64 supports pluripotency and
regulates cell cycle progression in embryonic stem cells through histone 3' end processing. Nucleic
Acids Res. 2014;42(13):8330-42.

88.   Romeo V, Griesbach E, Schumperli D. CstF64: cell cycle regulation and functional role in 3'
end processing of replication-dependent histone mRNAs. Mol Cell Biol. 2014;34(23):4272-84.

89.   Youngblood BA, MacDonald CC. CstF-64 is necessary for endoderm differentiation resulting
in cardiomyocyte defects. Stem Cell Res. 2014;13(3 Pt A):413-21.

90.   Baek JY, Jun DY, Taub D, Kim YH. Characterization of human phosphoserine
aminotransferase involved in the phosphorylated pathway of L-serine biosynthesis. Biochem J.
2003;373(Pt 1):191-200.

91.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods. 2012;9(7):671-5.

92.   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al. Ensembl 2020.
Nucleic Acids Res. 2020;48(D1):D682-D8.

93.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 2010;26(6):841-2.

94.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15(12):550.

95.   Ignatiadis N, Klaus B, Zaugg JB, Huber W. Data-driven hypothesis weighting increases
detection power in genome-scale multiple testing. Nat Methods. 2016;13(7):577-80.

96.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The human
genome browser at UCSC. Genome Res. 2002;12(6):996-1006.

97.   Green MR, Sambrook J. Rapid Amplification of Sequences from the 3' Ends of mRNAs: 3'-
RACE. Cold Spring Harb Protoc. 2019;2019(5).

98.   Scotto-Lavino E, Du G, Frohman MA. 3' end cDNA amplification using classic RACE. Nat
Protoc. 2006;1(6):2742-5.

10   Data Availability Statement

Poly(A) sequencing data sets are available in the NCBI Gene Expression Omnibus (GEO), accession
number #GSE169140. Supplementary materials are attached with this manuscript.

11   Figure captions

Figure 1. Morphological changes and altered expression of cell-surface markers and mRNAs
during monocyte differentiation.

(A) The effect of differentiation on cellular morphology. U937 cells exposed to PMA (30 nM) for 6
and 24 hours and observed by phase contrast microscopy (40X). (B) Western blots of total cell
lysates from U937 cells differentiated with PMA (30 nM) for 0, 1, 6, 18, and 24 hours. Blots were
probed with antibodies against CD16, CD68, HLA-DRA, CD38 and CD14 with β-actin as the
loading control. (C) Attachment assay of U937 cells. Cells were treated with PMA for 0, 6, and 24
hours and live cells visualized and counted by the Trypan blue exclusion assay. The graph presents
the percentage of cells that are suspended or attached. (D) Proliferation assay of U937 cells. The
number of cells before and after differentiation depicts the absolute cell number in millions,
compared to normal cell proliferation in the absence of PMA. The figure represents mean ± SE from
three independent experiments. (E) Real-time quantitative PCR (RT-qPCR)–based analysis of the
expression of genes important for macrophage differentiation and function *(CD86, ICAM1, JUN,*
FOS, NFKB1, IKZF1, OSM and *IL12A)*. The RT-qPCR analysis used primers designed in the first
exon to represent the total transcript level and the figure shows $\log_2$-fold changes in expression of
these genes at 6 hours and 24 hours post PMA treatment of U937 cells, where values are normalized
to ACTB mRNA and the 0 hour (no PMA) time point. The figure represents mean ± SE from three
independent experiments. P value <0.05 was considered significant, where * = $P \leq 0.05$; ** = $P \leq$
0.01; *** = $P \leq 0.001$; **** = $P \leq 0.0001$.

Figure 2. Evidence of APA in U937 cells after 6 and 24 hours of PMA treatment.

(A) Bar graph representing the numbers of genes with APA changes in the terminal 3' UTR after 6h
and 24h of PMA treatment with respect to control undifferentiated U937 cells. Red, green and blue
represents shortening, lengthening and complex (where some went up and others went down) APA
patterns in the dataset. The criteria for a significant APA change is that at least one PAC in an exon
must change greater than 1.5 fold with $p_{Adj} <0.1$, resulting in a difference in fractional usage of that
PAC of at least 10%. (B & C) $\log_2$-fold change in gene expression is plotted against the change in
Percentage of Distal poly(A) site Usage Index (delta-PDUI) for 3′-UTR-altered genes in 6h PMA-
treated U937 cells with respect to control undifferentiated ones (B) or in 24h PMA-treated U937
samples (C). Plots for fold change vs poly(A) site usage (delta PDUI) where genes on the left half of
each graph represent 3′-UTR-shortened hits, whereas those on the right half represent 3′-UTR
lengthened hits.

Figure 3. Gene set enrichment analysis (GSEA) of APA events in U937 cells after PMA
treatment.

Functional annotation clustering of mRNAs whose APA is regulated and after 6h (A) and 24h (B) of
differentiation. The 10 most significant GO-process enriched gene groups for biological process
(green) or canonical process (red) are ranked based on negative $\log_{10}$ (P-values) for shortened (left
panel) and lengthened (right panel) transcripts. Pathways classified according to specific cellular
processes are grouped together as indicated by the individual symbols at the bottom of the graphs.

Figure 4. Changes in poly(A) site use in U937 cells after 24 hours of PMA treatment.

UCSC genome browser plots of RNA sequencing tracks highlighting the 3′-UTR profile differences
for shortened genes (A) or lengthened genes (B) after 24 hours of PMA treatment with respect to
control (0h). The colors of the tracks represent 0h (red) and 24h (green). Proximal (P) and distal (D)
poly(A) sites are indicated with red stars. The green arrow defines the direction of the coding strand,
and tag counts are indicated on the y axis. Additionally, positions and chromosome co-ordinates of
the annotated PACs are indicated at the top of each plot.

Figure 5. Validation of APA events in U937 by 3' RACE.

Representative 1% agarose gel images for the 3′ RACE RT-PCR from U937 cells (untreated and 24h
PMA-treated) for (A) shortened genes SERPINB1, PTCH1 and (B) lengthened genes PSAT1 and
MRPL44 using gene specific primers at the last exon-exon junction and a common reverse anchor
primer. To confirm the direction of the shift in poly(A) site use, normalized intensities of long and
short bands for each PCR were quantified and a long/short ratio determined and normalized to the No
PMA control. The corresponding graphs represent mean ± SE from at least two independent
experiments. P value <0.05 was considered significant, where * = P ≤ 0.05; ** = P ≤ 0.01; *** = P ≤
0.001.

Figure 6. Effect of macrophage differentiation on the expression of C/P proteins in U937 cells.

(A) Whole cell lysates from U937 cells treated with PMA for 0, 1, 6, 18 and 24 hours were separated
by 10% SDS-PAGE and western blotting was performed for the indicated subunits of the C/P
complex, with β-actin as the loading control. Each western blot (left panel) was performed in three
biological replicates of the differentiation process and the quantified data is shown in the right panel.
(B) Schematic of the mammalian core C/P complex. (C) Processing efficiency of MYC and ACTB
transcripts for U937 cells. The bar graph shows the ratio of unspliced (US) RNA transcripts (detected
by RT-qPCR with a primer pair upstream of the poly(A) site) to unprocessed RNA (detected by RT-
qPCR with a primer pair that spans the poly(A) site) as shown in the schematic. The cDNA
preparation was done using random hexamers.

Figure 7. Knockdown or overexpression of CstF64 in U937 cells alters macrophage
differentiation.

(A) Western blot analysis of C/P factors CstF64 and CstF64τ, macrophage-specific markers CD16,
CD68, HLA-DRA, CD38 and the monocyte marker CD14 in U937 cells stably transfected with
doxycycline-inducible control shRNA and two different CstF64 shRNAs (shRNA 2 and shRNA 3).
Addition of 3 nM doxycycline for 3 days supplemented with 30 nM PMA one day prior to harvesting
was done for all three cell lines. (B) Attachment assay (left panel) and changes in total number of
PMA-treated cells (right panel) after knockdown of CstF64 compared to the shRNA control. Cells
were treated with doxycycline, after which equal number of cells were treated with PMA for 24 h and
then counted. The figures are representative of at least three independent experiments. (C) Changes
in cellular complexity determined by FACs analysis of side (SSC) and forward scattering (FSC)
values for 24h PMA-treated U937 cells stably expressing control shRNA and shRNA against CstF64.
(D) Western blot analysis of C/P factors CstF64, CstF64τ and macrophage-specific markers CD68,
HLA-DR, CD38 and the monocyte marker CD14 in U937 cells stably transfected with control
overexpression vector (OE-control) and those overexpressing CstF64 (OE-CstF64) and treated with 3
nM doxycycline for 3 days prior to harvesting. No PMA treatment has been done. (E) Attachment
assay (left panel) and changes in total number of cells (right panel) after overexpression of CstF64
compared to the OE-control. (F) Changes in cellular complexity or granularity determined by FACs
analysis of side (SSC) and forward scattering (FSC) values for doxycycline-induced U937 cells
transfected with OE-control or OE-CstF64. Percentage indicates a quantitation of cells that have
altered granularity.

Figure 8. Knockdown or overexpression of CstF64 in U937 cells alters APA.

Representative 1% agarose gel images for the 3′ RACE RT-PCR in U937 cells for shortened genes
SERPINB1 (A) and *PTCH1* (B), using gene specific primers at the last exon-exon junction and a

common reverse anchor primer. In each subfigure, the effect of depletion (knockdown) of CstF64
and the effect of overexpression of CstF64 are indicated, using the same treatment protocols as
described in Fig. 7. Normalized intensities of long and short bands for each PCR were quantified and
a long/short ratio determined and normalized to the shRNA or OE control. The corresponding graphs
(right) represents mean ± SE from at least two independent experiments. P value <0.05 was
considered significant, where * = P ≤ 0.05; ** = P ≤ 0.01; *** = P ≤ 0.001.

Fig 1

Fig 2

A

B

C

Fig 3

A Pathway analysis of genes undergoing <u>APA changes</u> after 6h of PMA treatment:

Shortened genes (0 vs 6):

- ● GOBP_REGULATION_OF_CATABOLIC_PROCESS — 20
- ● GOBP_REGULATION_OF_CELLULAR_CATABOLIC_PROCESS — 18
- ▲ GOBP_POSITIVE_REGULATION_OF_INTRACELLULAR_TRANSPORT — 10
- ● GOBP_MACROMOLECULE_CATABOLIC_PROCESS — 21
- ▲ GOBP_INTRACELLULAR_TRANSPORT — 23
- ▲ GOBP_POSITIVE_REGULATION_OF_TRANSPORT — 16
- ● GOBP_CELLULAR_MACROMOLECULE_CATABOLIC_PROCESS — 18
- ■ GOBP_REGULATION_OF_IMMUNE_SYSTEM_PROCESS — 21
- ▲ GOBP_POSITIVE_REGULATION_OF_INTRACELLULAR_PROTEIN_TRANSPORT — 8
- ▲ GOBP_REGULATION_OF_TRANSPORT — 22

- ■ REACTOME_DAP12_SIGNALING — 4
- ● REACTOME_MTOR_SIGNALLING — 4
- ○ BIOCARTA_TCR_PATHWAY — 4
- ■ REACTOME_DAP12_INTERACTIONS — 4
- ■ WP_B_CELL_RECEPTOR_SIGNALING_PATHWAY — 5
- ■ REACTOME_PLATELET_ACTIVATION_SIGNALING_AND_AGGREGATION — 7
- △ REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION — 16

0 2 4 6 8 10
-log p value

Lengthened genes (0 vs 6):

- → GOBP_APOPTOTIC_PROCESS — 18
- ■ GOBP_IMMUNE_SYSTEM_DEVELOPMENT — 13
- □ GOBP_RESPONSE_TO_ABIOTIC_STIMULUS — 14
- GOBP_REGULATION_OF_DNA_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY — 9
- → GOBP_REGULATION_OF_CELL_DEATH — 16
- ● GOBP_MACROMOLECULE_CATABOLIC_PROCESS — 14
- ■ GOBP_NEGATIVE_REGULATION_OF_NF_KAPPAB_TRANSCRIPTION_FACTOR_ACTIVITY — 5
- □ GOBP_REGULATION_OF_CELL_DIFFERENTIATION — 16
- GOBP_NEGATIVE_REGULATION_OF_DNA_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY — 6
- ■ GOBP_INTERLEUKIN_1_MEDIATED_SIGNALING_PATHWAY — 5

- ■ KEGG_NEUROTROPHIN_SIGNALING_PATHWAY — 6
- ■ BIOCARTA_IL1R_PATHWAY — 4
- ■ WP_SIGNAL_TRANSDUCTION_THROUGH_IL1R — 4
- ○ WP_ATM_SIGNALING_PATHWAY — 4
- ■ REACTOME_TOLL_LIKE_RECEPTOR_TLR1_TLR2_CASCADE — 5
- ■ WP_VEGFAVEGFR2_SIGNALING_PATHWAY — 8
- ■ WP_IL1_SIGNALING_PATHWAY — 4
- ■ REACTOME_TOLL_LIKE_RECEPTOR_4_TLR4_CASCADE — 5
- ○ BIOCARTA_ATM_PATHWAY — 3
- ■ REACTOME_TOLL_LIKE_RECEPTOR_CASCADES — 5

0 2 4 6 8
-log p value

Legend:
- ● Metabolic process
- ■ Immune system or signal pathways related to macrophage function
- ▲ Cellular transport
- ○ DNA damage repair
- → Regulation of cell death
- □ Change in state or activity
- △ Protein modification such as small protein conjugation or phosphorylation

- 🟩 Biological process
- 🟥 Canonical process

B Pathway analysis of genes undergoing <u>APA changes</u> after 24h of PMA treatment:

Shortened genes (0 vs 24):

- ← GOBP_CELL_CYCLE — 63
- ← GOBP_CHROMOSOME_ORGANIZATION — 50
- ✹ GOBP_MICROTUBULE_BASED_PROCESS — 36
- ← GOBP_CELL_CYCLE_PROCESS — 48
- ✹ GOBP_MICROTUBULE_CYTOSKELETON_ORGANIZATION — 30
- ✹ GOBP_CYTOSKELETON_ORGANIZATION — 46
- GOBP_ORGANELLE_ASSEMBLY — 35
- GOBP_REGULATION_OF_ORGANELLE_ORGANIZATION — 42
- ✹ GOBP_MICROTUBULE_ORGANIZING_CENTER_ORGANIZATION — 15
- GOBP_REGULATION_OF_HYDROLASE_ACTIVITY — 42

- ← REACTOME_CELL_CYCLE — 31
- △ REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION — 42
- ← REACTOME_CELL_CYCLE_MITOTIC — 24
- ← REACTOME_MITOTIC_PROMETAPHASE — 15
- ← REACTOME_M_PHASE — 19
- △ REACTOME_SUMOYLATION — 13
- REACTOME_RNA_POLYMERASE_II_TRANSCRIPTION — 36
- ○ REACTOME_DNA_REPAIR — 16
- PID_TELOMERASE_PATHWAY — 8
- REACTOME_DISEASE — 36

0 5 10 15 20
-log p value

Lengthened genes (0 vs 24):

- △ GOBP_PROTEIN_MODIFICATION_BY_SMALL_PROTEIN_CONJUGATION_OR_REMOVAL — 31
- GOBP_REGULATION_OF_RESPONSE_TO_STRESS — 36
- ← GOBP_CHROMOSOME_ORGANIZATION — 31
- GOBP_REGULATION_OF_TRANSFERASE_ACTIVITY — 26
- GOBP_POSITIVE_REGULATION_OF_TRANSFERASE_ACTIVITY — 21
- ○ GOBP_CELLULAR_RESPONSE_TO_DNA_DAMAGE_STIMULUS — 24
- GOBP_POSITIVE_REGULATION_OF_CATALYTIC_ACTIVITY — 31
- △ GOBP_POSITIVE_REGULATION_OF_KINASE_ACTIVITY — 19
- △ GOBP_PROTEIN_MODIFICATION_BY_SMALL_PROTEIN_CONJUGATION — 24
- ← GOBP_CELL_CYCLE — 36

- △ REACTOME_POST_TRANSLATIONAL_PROTEIN_MODIFICATION — 29
- ← REACTOME_CELL_CYCLE — 19
- ← REACTOME_CELL_CYCLE_MITOTIC — 16
- ■ REACTOME_CYTOKINE_SIGNALING_IN_IMMUNE_SYSTEM — 19
- △○ REACTOME_SUMOYLATION_OF_DNA_DAMAGE_RESPONSE_AND_REPAIR_PROTEINS — 6
- ■ REACTOME_MAPK_FAMILY_SIGNALING_CASCADES — 10
- ■ PID_P38_ALPHA_BETA_PATHWAY — 4
- △ REACTOME_SUMOYLATION — 8
- ← REACTOME_MITOTIC_G1_PHASE_AND_G1_S_TRANSITION — 7
- REACTOME_DISEASE — 24

0 5 10 15 20
-log p value

- 🟩 Biological process
- 🟥 Canonical process

Legend:
- ← Cell cycle and chromosome organization
- ✹ Cytoskeletal organization
- △ Protein modification such as small protein conjugation or phosphorylation
- ○ DNA damage repair
- ■ Immune system or signal pathways related to macrophage function

Fig 4

A U937 shortened genes (0 vs 24 hours):

B U937 lengthened genes (0 vs 24 hours):

Fig 5

A   U937 shortened genes

SERPINB1   **PTCH1**

B   U937 lengthened genes

PSAT1   **MRPL44**

Fig 6

A

U937 western blot

U937 western blot quantification

B

C

Fig 7

Fig 8

A <u>SERPINB1:</u>

B <u>PTCH1:</u>