Isotope effects on radical pair performance in cryptochrome: a new hypothesis for the evolution of animal migration

Ismael Galván[1], Abbas Hassasfar[2], Betony Adams[2,3] and Francesco Petruccione[2,4,5]

[1]Department of Evolutionary Ecology, National Museum of Natural Sciences, CSIC, 28006 Madrid, Spain.

[2]Quantum Research Group, School of Chemistry and Physics, University of KwaZulu-Natal, Durban 4001, South Africa.

[3]The Guy Foundation, Beaminster, DT8 3HY Dorset, UK.

[4]School for Data Science and Computational Thinking, Stellenbosch University, Stellenbosch 7600, South Africa.

[5]National Institute for Theoretical and Computational Sciences (NITheCS), Stellenbosch 7600, South Africa.

ORCID:

Ismael Galván: 0000-0002-6523-8592; Abbas Hassasfar: 0000-0003-1079-8848; Betony Adams: 0000-0001-9746-5442; Francesco Petruccione: 0000-0002-8604-0913.

Authors for correspondence:

Ismael Galván

e-mail: galvan@mncn.csic.es

Abbas Hassasfar

e-mail: omid.hassasfar@gmail.com

Abstract

Mechanisms occurring at the atomic level are now known to drive processes essential for life, as revealed by quantum effects on biochemical reactions. Some macroscopic characteristics of organisms may thus show an atomic imprint, which may be transferred across organisms and affect their evolution. This possibility is considered here for the first time, with the aim of elucidating the appearance of an animal innovation with an unclear evolutionary origin: migratory behaviour. This trait may be mediated by a radical pair (RP) mechanism in the retinal flavoprotein cryptochrome, providing essential magnetic orientation for migration. Isotopes may affect the performance of quantum processes through their nuclear spin. Here, we consider a simple model and then apply the standard open quantum system approach to the spin dynamics of cryptochrome RP. We changed the spin quantum number (I) and g-factor of hydrogen and nitrogen isotopes to investigate their effect on RP's yield and magnetic sensitivity. Strong differences arose between isotopes with $I=1$ and $I=1/2$ in their contribution to cryptochrome magnetic sensitivity, particularly regarding Earth's magnetic field strengths (25-65 µT). In most cases, isotopic substitution improved RP's magnetic sensitivity. Migratory behaviour may thus have been favoured in animals with certain isotopic compositions of cryptochrome.

Keywords: animal migration, birds, Isotopic Resonance Hypothesis, nuclear spin effects, quantum biology, radical pairs, stable isotopes.

1. Introduction

Life depends upon interactions between the constituent molecules of organisms. Isolated molecules are inanimate. But the physicochemical characteristics of biomolecules, which determine their interactions, are given by their constituent atoms [1]. Life therefore occurs, to some extent, at an atomic scale. Not all atoms of a given element are equal: elements occur as isotopes, i.e., atoms of the same element that differ in the number of neutrons in their nuclei, thus having different mass numbers and nuclear spins. The isotopic composition of molecules might be transferred across molecules, and across organisms, during molecular biosynthesis independently of the genetic code, as nucleic acids cannot determine their own isotopic composition [2]. Atomic properties like this may be responsible for some macroscopic characteristics of organisms, but this possibility has been only envisaged by some authors [3-5]. However, if the isotopic composition of biomolecules is responsible for macroscopic characteristics of organisms, this may have profound consequences for organic evolution.

Different isotopic compositions of the same molecule (i.e., isotopologues) do not behave equally in some chemical reactions [6]. Stable isotopes, i.e., those not undergoing radioactive decay, are particularly relevant for organisms, because the nuclei of the most common biological elements (C, H, O, N and S) exist as 13 stable isotopes, making those biomolecules appear as mixed isotopologues. The different mass numbers of these stable isotopes give them slightly different properties, leading to their partial separation during chemical, biological and physical transformations in what is termed isotope fractionation. Fractionation occurs because of the influence in chemical reactions of the different masses (nuclei with more neutrons are heavier), volumes (nuclei with more neutrons are bigger) and spins (nuclei with certain spin values have a magnetic moment) of isotopes [7]. This explains the distribution of isotopes in the biosphere, as fractionation occurs first in the environment, then during plant photosynthesis, and lastly during animal metabolic reactions [8].

Different experiments have proved that organisms grown under certain isotopic

media or given diets with certain isotopes experience oxidative stress and resistance to

change the isotopic composition of their tissues [9]. This has been observed at all levels of

complexity, from microorganisms to vertebrates such as birds and humans, and in isotopes

of elements such as H, C and N [10-12]. These studies indicate that organisms have a

preferred isotopic environment, because there is an optimum isotopic composition of

molecules under which biochemical reactions accelerate [9,13,14]. Concomitantly, an

evolutionary optimization explains the negative physiological effects that organisms

experience under deviations of their isotopic composition, even if adaptation can be

achieved [9]. Chemical and evolutionary optimizations thus make likely that every organism

has a specific isotopic profile that is not randomly determined by environmental influences.

Yet, a small proportion of variation in the isotopic composition of animals is due to the

incorporation of isotopes with the food [15,16]. Ecological studies even assume that the

whole-body composition of an animal closely resembles that of its diet when tracking the

trophic behavior of animal groups within a given species [8]. This is used to infer the trophic

niche (isotopic niche or "isospace") of species of animals by plotting the composition of

certain tissues in C isotopes against that of N isotopes. Isospaces show significantly

differentiated, characteristic isotopic compositions of many species of animals, including

hominids [17,18]. Importantly, isospaces do not overlap even between coexisting species

that are morphologically and ecologically similar [19]. This supports the idea that, even

considering the dietary source of variability in the isotopic composition of animals, every

species may have a characteristic isotopic composition. This actually agrees with the

existence of niche conservatism, i.e. the retention of ecological traits in species across

space and time, as revealed by non-overlapping isospaces in ecologically similar species of

animals [19,20]. These differentiated isospaces of animals have never been considered an

intrinsic trait of species, but here it is hypothesized that the isotopic composition of molecules

might be more similar within animals belonging to the same species than between animals

of different species (i.e., interspecific variability > intraspecific variability). Indeed, even if

part of variation in the isotopic composition of animals is due to dietary incorporations,

individuals of the same species share biochemical reactions of metabolism, thus

experiencing isotope fractionation processes that must be more similar than in individuals

of other species, which possess different metabolic profiles. This is because metabolic

pathways are shared by all individuals of the same species, an observation that has allowed

the development of metabolomics, in a similar way as genomics, as a useful tool to elucidate

the evolution of species-specific traits in organisms [21,22]. Certain isotopic compositions in

certain species may therefore favour some biochemical reactions and thus promote the

evolution of important phenotypic traits.

The development of the field of quantum biology suggests that biochemical reactions

involved in physiological processes are mediated by quantum mechanisms [23,24]. One

such mechanism is represented by radical pairs (RP), which are strongly influenced by

isotopes [25] and are potentially responsible for animal magnetoreception: the ability to

perceive the Earth's magnetic field and use it to orientate during migratory movements. This

ability is most notably observed in birds, but has evolved in all major vertebrate groups and

some invertebrates [26,27]. The mechanism of magnetoreception and the ability for

magnetic orientation was proposed to be mediated by a light-initiated RP in the eyes of birds,

whose recombination disruption by the magnetic field is recognized by the bird's nervous

system [28]. This is currently the leading theory of avian magnetoreception. Cryptochrome

4 (CRY4), a pigment belonging to the only vertebrate protein class that forms radical pairs

upon photo-excitation (flavoproteins), has been suggested to be the likely magnetoreceptive

protein, localized in double-cones and long-wavelength single cones in the retina of birds

[29]. Recent research has demonstrated that the photochemistry of CRY4 in night-migratory

birds is more magnetically sensitive than that of some non-migratory species [30]. However,

the exact type of cryptochrome responsible for magnetoreception remains a subject of

debate. There are five different types of cryptochrome found in birds' retinas. In addition to

CRY4, cryptochrome 1a (CRY1a) has been put forward as the likely site for a light-sensitive

magnetic compass, due to its specific location in the outer segments of the ultraviolet cone

cells of birds [31]. The majority of research into the role of RPs in magnetoreception has

focused on birds. The RP mechanism could, however, be the same in other animals with

magnetic orientation capabilities, as the expression of magneto-sensitive cryptochrome has

been shown to occur in the retina of insects [32], and even of humans [33]. The capacity for

magnetic orientation has also been shown to reside in the eyes of other mammals [34].

In its simplest iteration, an RP consists of two radicals (molecules with an odd number

of electrons) that have been created simultaneously by chemical bond breaking or light-

induced electron transfer, forming a spatially separated but spin-correlated and possibly

entangled electron pair. This entanglement is due to the fact that the RP is often assumed

to be created in a singlet, or maximally entangled state, although other options have been

discussed [35]. Due to external and local (hyperfine interaction) magnetic fields, the spin

character of the radical pair is coherently interconverted between singlet (anti-parallel

electron spins) and triplet (parallel electron spins) states [36]. The spin state of the electron

(as a purely quantum mechanical concept) then influences subsequent chemical reactions,

converting information about the magnetic environment into a chemical signature. As

outlined previously, this mechanism has been hypothesized to be responsible for

magnetoreception in birds: cryptochrome in the retina is photoexcited by light, forming an

RP that is initially in the singlet state and is interconverted between singlet and triplet states

by hyperfine coupling. The rate of recombination of singlet states also depends on the

orientation of the Earth's magnetic field, and the interconversion is oscillatory coherent,

generating a quantum beat that is used by birds for orientation [37]. See Figure 1 for details

of the RP formation.

Magnetoreception in birds is strongly believed to be an RP mechanism based on

retinal cryptochrome [38]. However, the strong influence that isotopes exert on RP

mechanisms has been largely overlooked, with the exception of Player and Hore [39]. This

influence is due to the fact that one of the dominant interactions involved in RP dynamics is

the internal magnetic interactions, known as hyperfine interactions, which are caused by the

surrounding nuclear spins that can interact with an unpaired electron. Nuclei can have no

spin (e.g., ^{12}C and ^{16}O), integer spin (e.g., ^{14}N and ^{2}H) and also half-integer spin (e.g., ^{13}C

and ^{1}H) [40]. Thus, different isotopes of an element can have different magnetic effects on

the dynamics of the system. An RP with isotopes with certain spin undergoes fast triplet-

singlet conversion, while this is strongly delayed in the case of isotopes with other spins [25].

Cryptochrome molecules with isotopes with certain spin numbers might thus impair the RP

mechanism, which might be favoured in cryptochrome isotopologues with other spin

numbers (Figure 1). The ability for magnetic orientation and long-distance migration may

therefore be more likely to arise in animals that have inherited cryptochrome molecules with

higher proportions of isotopes with certain spin numbers.

Here we hypothesize that the ability to conduct long-distance migration has evolved

only in animals with a prevailing contribution of isotopes with certain spin to the

cryptochrome molecule. To evaluate the potential of this hypothesis to explain interspecific

variability in migratory ability, we modeled spin dynamics of the broadly accepted RP model

of the cryptochrome-based avian compass.

2. Methods

We considered two different theoretical scenarios to investigate our hypothesis. Both involve

the conventional radical pair approach, though there is some suggestion that adding a third

radical improves the performance of the radical-based compass [41]. In the first scenario

(**Scenario 1**), we explored a simple model with one hyperfine-coupled nucleus for the

calculation of the magnetic sensitivity (MS) of cryptochrome depending on different quantum

spin numbers associated to different nuclei. We explored MS in a different range of

anisotropic hyperfine coupling constants in the presence of an external magnetic field,

similar to the Earth's geomagnetic field. The second scenario (**Scenario 2**) looks more

closely at isotope substitution of the specific model by changing the spin quantum number

and also the g-factor for scaling the strength of hyperfine interaction for different

geomagnetic field intensities. In this second scenario, we used the DFT calculations of

nuclear hyperfine coupling tensors by Hiscock et al. [42] for cryptochrome, in addition to a

simpler version with up to three nuclei that has previously been used by Jain et al. [43] and

that assumes that all hyperfine tensors are simultaneously diagonalized. In this second

scenario, we show that both singlet yield and MS change with isotope substitution.

The spin dynamics of RPs are described by the following Hamiltonian:

$$H = H_{\text{Zeeman}} + H_{\text{Hyperfine}} + H_{\text{Exchange}} + H_{\text{Dipolar}} + H_{\text{Nuclear}} \tag{1}$$

Among these five terms, the exchange, dipolar and nuclear Zeeman interactions usually are

negligible as compared to the other two terms [44].

Two dominant interactions are conventionally taken to be involved in RP dynamics:

one of them is the Zeeman interaction (H_{Zeeman}), due to the magnetic field of an external

source (e.g., the geomagnetic field). The other is due to internal magnetic interactions,

known as hyperfine interactions ($H_{\text{Hyperfine}}$) which are caused by the local surrounding

nuclear spins that can interact with an unpaired electron. The RP Hamiltonian with one

nucleus can thus be written as:

$$H = \gamma\mathbf{B}.(\hat{S}_1 + \hat{S}_2) + \hat{I}.\mathbf{A}.\hat{S}_1 \tag{2}$$

where 'I' is the nuclear spin operator, S_1, $S_2 \in \frac{1}{2}(\sigma_x, \sigma_y, \sigma_z)$ are electron spin operators (σ_x, σ_y,

σ_z are the Pauli matrices), $\gamma = \mu_0 g$ is the gyromagnetic ratio, μ_0 is the Bohr magneton, and g

is the electron g factor (= 2.0023). We will consider the simplest case in which the hyperfine

interaction is anisotropic, A = diag(0, 0, a) as well as the more complex case of A = diag(a_x,

a_y, a_z). The external field (geomagnetic field) is characterized by $B = B_0(\sin\theta\cos\varphi, \sin\theta\sin\varphi,$

$\cos\theta)$; $B_0 = 47$ µT is the local geomagnetic field, and θ is the magnetic field orientation with

respect to the molecular axis [45]. These interactions dictate the dynamics of the electrons'

spin state and can cause coherent conversion between singlet and triplet states.

Different approaches have previously been used to tackle RP dynamics and

recombination. These include Haberkorn's [46] master equation, the Jones-Hore

measurement master equation [47], Kominis' [48] measurement master equation approach,

which commonly includes the Liouville-von Neumann equation, as well as other open

quantum systems approaches [49]. There remains some debate as to which is the best

approach [44]. We use the Markovian quantum master equation approach to simulate the

dynamics of the RP system of cryptochrome, similar to Gauger et al. [50]. In this quantum

master equation approach, which conserves probability unlike the Liouville equations, we

use the 'shelving states' $|S\rangle$ and $|T\rangle$ to represent the singlet and triplet products that are

spin-selected from the initial electronic singlet or triplet states of the RP. Mathematically, we

use the direct sum to extend the Hilbert space of the radical pair to include the shelving

states $|S\rangle$ and $|T\rangle$ as extra basis vectors. The recombination of the RP into singlet and

triplet channels is modeled through decay operators. For example, for one nucleus model

with I=1/2, there are eight projections, four projections for the 'up' ($|\uparrow\rangle$) state of nucleus

$P_{S,\uparrow} = |S\rangle\langle s,\uparrow|, P_{T_0,\uparrow} = |T\rangle\langle t_0,\uparrow|, P_{T_+,\uparrow} = |T\rangle\langle t_+,\uparrow|, P_{T_-,\uparrow} = |T\rangle\langle t_-,\uparrow|$ and, similarly, four projections

for   the   'down'   ($|\downarrow\rangle$)   nuclear   state $P_{S,\downarrow} = |S\rangle\langle s,\downarrow|, P_{T_0,\downarrow} = |T\rangle\langle t_0,\downarrow|,$

$P_{T_+,\downarrow} = |T\rangle\langle t_+,\downarrow|, P_{T_-,\downarrow} = |T\rangle\langle t_-,\downarrow|$ [49]. The singlet yield is defined as the proportion of

recombined chemical products originating from the singlet precursor. One of the two events

will occur, and the final populations of $|S\rangle$ and $|T\rangle$ give the singlet and triplet yield.

The standard, GKSL (Gorini–Kossakowski–Sudarshan–Lindblad equation) master

equation can be defined as [51]:

$$\dot{\rho} = -\frac{i}{\hbar}[H,\rho] + k_s \sum_{i=1}^{N} P_i \rho P_i^\dagger - \frac{1}{2}(P_i^\dagger P_i \rho + \rho P_i^\dagger P_i) + k_t \sum_{i=1}^{M} P_i \rho P_i^\dagger - \frac{1}{2}(P_i^\dagger P_i \rho + \rho P_i^\dagger P_i) + \mathrm{L}(\rho)$$

$$(3)$$

where P_i are the projections on singlet and triplet, and N and M are the number of projection operators into singlet and triplet that depend on the Hilbert space of the nucleus. In the case of one nucleus with I=1/2, then N=2 and M=6, but in the case of one nucleus with I=1, then N=3 and M=9. L(ρ) is the standard Lindblad dissipator, used here to describe noise in the system. Several different noise models have previously been proposed, with different noise rates [50,52]. Here, we do not aim to explore all of the different noise models. As an important example we consider a special type of noise, perfectly correlated pure local dephasing noise, given by:

$$\mathrm{L}(\rho) = \sum_{i=1,2} \Gamma\left(L_i \rho L_i^\dagger - \frac{1}{2}(L_i^\dagger L_i \rho + \rho L_i^\dagger L_i) \right), \quad L_i = \frac{\sigma_z^{(1)}}{2} + \frac{\sigma_z^{(2)}}{2}, \qquad (4)$$

where Γ is the dephasing rate. Results suggest that with this noise model MS is quite robust and is even enhanced by the presence of correlated dephasing Cai et al. [52].

The master equation is given in terms of the rates of recombination to singlet and triplet products, k_s and k_t, where H is given in Eq. (3). The dynamics of the master equation of the RP system were calculated using the QuTiP quantum toolbox in the Python module [53], which is developed for simulating quantum systems, particularly open quantum systems.

Here, we aim at showing that isotopes react preferentially (in terms of singlet or triplet yield and MS) depending on their nuclear spin quantum number I and strength of hyperfine interaction, which is embedded in their g-factor. We thus explored the quantum spin dynamics of the RP to investigate isotope-dependent magnetic field effects in the specific system of cryptochrome.

We only consider the spin dynamics, neglecting any effects related to nuclear mass

and volume. This is possible due to the fact that, within the Born-Oppenheimer

approximation, electronic properties (such as spin density) do not depend on the nuclear

mass and nuclei are considered point charges in quantum chemistry, meaning that nuclear

volume is also irrelevant [54]. Therefore, we need only to change the spin quantum number

and the g-factor to explore the effect of isotope substitution (we explain in section 2.1 how

g-factor affects hyperfine coupling constant).

We made the following assumptions for the sake of simplification:

1. All projectors have the same recombination rates. We have used two different

recombination rates: $k_s = k_t = k = 0.1$ µs^{-1} and 0.5 µs^{-1} ($\tau = 1/k = 10$ µs and 2 µs, the lifetime of

the radical pair, τ, is defined as the reciprocal of k) [50]. It is clear from the results that the

sensitivity of the avian compass relies on the recombination rate, where a longer lifetime

allows for greater singlet-triplet interconversion and resultant greater sensitivity.

2. One electron of the pair has no hyperfine interaction and the other electron experiences

a hyperfine interaction due to the surrounding nucleus.

3. The initial state of our model ρ_0 assigns a pure singlet state $|s\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$ to the

electrons, and a completely mixed state to the nucleus due to its interaction with the

neighboring soft matter environment, initial state $\rho(0) = \frac{I}{N} \otimes (|s\rangle\langle s|)$ where N is the dimension

of the Hilbert space of the nucleus (e.g., for one nucleus with spin 1/2, N=2, one nucleus

with spin 1, N=3, and so on).

4. Without loss of generality, the axial symmetry of the hyperfine tensor allows us to assume

that $\varphi = 0$ [50].

Our analyses are consistent with previous analyses of singlet yield for various

hyperfine interaction strengths for different spin quantum numbers, and reproduce the

sensitivity behaviour shown by Cai et al. [52] and Lee et al. [55] for cryptochrome. The

magnetic field effect was calculated as Φ_S, i.e., the fractional yield of the singlet reaction

product once all radical pairs have reacted. Φ_S lies in the range [0,1] and is related to the

triplet product yield by $\Phi_T = 1 - \Phi_S$. To quantify the effectiveness of a radical pair as a

magnetic compass, we define the anisotropy of the reaction yield, i.e. MS, as

$\Delta\Phi_S = \Phi_S^{max} - \Phi_S^{min}$ (the difference between the maximum and the minimum singlet yield as a

function of the inclination). S denotes the singlet yield and the max/min are with respect to

the magnetic field orientation, the difference between the maximum and minimum values of

Φ_S being calculated as a function of the direction of the magnetic field vector, following the

example of Lee et al. [55].

2.1. Isotope substitution

Substitution of one nucleus X, with spin quantum number I and factor g_n, by its isotope X'

with I' and g_n', changes the number of lines detected by Electron Spin Resonance

Spectroscopy from (2I+1) to (2I'+1) and the coupling constant from a_X to $a_{X'} = a_X \cdot g_n'/ g_n$

[56]. Thus, for example, replacing a proton $(X = {}^1H = H; I = 1/2; g_n = 5.5854)$ by a deuteron

$(X = {}^2H = D; I = 1; g_n' = 0.8574)$ increases the number of lines from 2*1/2 + 1 = 2 to 2*1 + 1

= 3 and decreases the coupling constant from a_H to $a_D = a_H$ (0.8574/5.5854) = 0.1535 a_H.

On the other hand, substituting a ^{14}N nucleus $({}^{14}N = N = H; I = 1; g_n = 0.4038)$ by its ^{15}N

isotope $({}^{15}N = N'; I = 1/2; g'_n = -0.5664)$ decreases the number of lines from 2*1 + 1 = 3 to

2*1/2 + 1 = 2 and converts the coupling constant a_N into $a_N' = a_N$ (- 0.5664)/0.4038 = -1.4027

a_N [56] (Figure 2). The properties of H and N and their isotopes are summarized in Table 1.

As previously stated, different isotopes of a given element can have different

magnetic effects on the dynamics of the system. Apart from the spin quantum number, the

strength of hyperfine interaction can be determined via the nuclear g-factor. By varying the

number of nuclear spins interacting with the RP, it is possible to investigate the effects of

different nuclear environments and thus draw some conclusions about the structure of cryptochrome, the biomolecule in which it is thought that the RP reaction takes place.

3. Results

3.1. Scenario 1. In this scenario, the only difference between the treatment of nuclei in the model lies in their spin quantum numbers, meaning that we investigate MS with respect to the hyperfine coupling of a single nucleus model for two different spin quantum numbers (I=1 and I=1/2). We also consider only the simplest type of anisotropy, with A = (0, 0, a). In this simplest case, we do not include noise in the model. The results of calculations of MS for different strengths of anisotropic hyperfine coupling are shown in Figure 3 and show a marked advantage for spin I=1/2 than I=1. These results are also useful in that they give us an idea of the best and worst hyperfine coupling strengths with respect to magnetic sensitivity, albeit in the limited case of simple anisotropy.

In Figure 4 we again investigate MS for the two different spin numbers across a range of hyperfine coupling strengths, but in this instance we have included noise in the model. As previously specified, our aim was not to investigate the effects of different noise models. Given the calculations by Cai et al. [52] it appears that a specific noise model, such as a perfectly correlated dephasing model, may increase the performance of a chemical compass. We were interested in the effects that this noise model would have on radical pair dynamics for different isotopes. With noise incorporated into the model according to Eq. (4) with $\Gamma = k - 3k$ for one nucleus with I=1/2, the maximum MS increases from 0.388 (corresponding to a hyperfine coupling strength of 17.4 µT) to 0.411 (corresponding to a hyperfine coupling strength of 20.9 µT). For one nucleus with I=1, the maximum MS increases from 0.259 (corresponding to a hyperfine coupling strength of 8.7 µT) to 0.274 (corresponding to a hyperfine coupling strength of 10.6 µT).

3.2. Scenario 2. For this second scenario we used the data provided by Jain et al. [43],

summarized in Table 2. These authors used three nuclei for each RP of cryptochrome (three

nuclei for FAD$^{\bullet-}$ and three nuclei for TrpH$^{\bullet+}$) instead of 11 nuclei for each RP as usually

considered when investigating hyperfine coupling. For our purposes we focus specifically

on the hydrogens H6 and H1 as well as the nitrogens N10 and N1 to compare results for

isotopic substitution, see Table 2 for details.

3.2.1. Scenario 2a. We investigated singlet yield with respect to the inclination (angle

describing the orientation of magnetic field to the basis of the hyperfine tensor) of different

N and H isotopes in the RP [FAD$^{\bullet-}$ - TrpH$^{\bullet+}$] of the cryptochrome molecule, and explored the

effect of isotope substitution. The results of these models are shown in Figure 5. It is clear

that isotopes with integer spin have consistently higher singlet yield than half integer, even

when taking into account their different coupling constants.

Similar conclusions are obtained when investigating a two-nuclei model instead of

single nuclear spin coupled to one of the electrons (Figure 6). In the two-nuclei model, there

are four different combinations of isotopes: ^{1}H^{14}N, ^{1}H^{15}N, ^{2}H^{14}N and ^{1}H^{15}N. The Hamiltonian

now gains an additional hyperfine term:

$$H = \gamma \mathbf{B}.(\hat{S}_1 + \hat{S}_2) + \hat{I}_1.\mathbf{A}_2.\hat{S}_1 + \hat{I}_2.\mathbf{A}_2.\hat{S}_1$$

Singlet yield is an interesting feature to consider with respect to RPs in the biological

context. In addition to magnetoreception, singlet yield has been used to investigate a

number of other biological functions which may depend on radical reactions. In reactions

involving reactive oxygen species, singlet yield might be used as an indication of oxidative

stress. The strong isotope dependence of singlet yield demonstrated in our results is

potentially interesting with respect to observations that isotopic changes in diet lead to

oxidative stress [9].

3.2.2. Scenario 2b. Here we investigated MS with respect to the strength of Earth's

magnetic field of different N and H isotopes in the RP [FAD$^{\bullet-}$ - TrpH$^{\bullet+}$] and explored the

effect of isotope substitution. Results are shown in Figure 7. Scenario 1 (see Figure 3)

showed a clear difference between how spin number changes MS, with $I=1/2$ conferring

greater MS than $I=1$. However, the effect of the spin number is also dependent on the

hyperfine coupling strength, with $I=1/2$ performing best at larger coupling constants than $I=1$.

In Scenario 2 it is more difficult to conclude that there is a favourable spin number with

respect to MS. This is likely due to the influence of the specific hyperfine coupling constants.

While the hyperfine coupling strength decreases with isotope substitution in hydrogen, it

increases with isotope substitution in nitrogen. If we consider only the magnetic field strength

relevant to migration, which is the geomagnetic field (25-65 µT), then for the case of both

hydrogens, $I=1$ gives greater magnetic sensitivity than $I=1/2$. The case is less clear for

nitrogen where both $I=1/2$ and $I=1$ show greater MS for specific magnetic field strengths

over the geomagnetic window (25-65 µT). This could be due to the fact that although the

coupling strengths for the nitrogen isotopes are further from the ideal values, the isotopic

substitution from $I=1$ to $I=1/2$ confers some advantage.

4. Discussion

Before any definite conclusions can be made, it should be acknowledged that the model we

have used is a toy model and is the simplest possible interaction of the RP. It serves,

however, to illustrate the hypothesis that isotopes may play a distinct and definite role in

macroscopic biological outcomes. This influence is exerted in various ways. Our theoretical

results show that the spin quantum number (I) of a nucleus in an RP strongly influences the

dependency of its MS on the strength of the hyperfine coupling. Considering a wide range

of hyperfine coupling values from 0 to 350 µT, following Lee et al. [55], the maximum MS is

considerably higher, and the asymptote is reached at larger hyperfine coupling values, when

I=1/2 (~20 µT) as compared to I=1 (~10 µT). These ideal hyperfine coupling strengths are

considerably smaller than those given for the specific nuclei of the RP mechanism, see

Table 2. Whether this distinction between I=1/2 and I=1 can be generalized to non-integer

and integer spin would depend on the inclusion of higher spin isotopes. RP performance

further improves when the noise model is perfectly correlated, suggesting that RPs in

cryptochrome, as well as other biological processes, might be optimized by the unavoidable

noise that is present in the environment, in accordance to Bandyopadhyay et al. [57].

Our results for the specific RP [FAD$^{\bullet-}$ - TrpH$^{\bullet+}$] of cryptochrome also show strong

differences in the dependency of the magnetic field effect (i.e., singlet yield) on inclination,

between isotopes with different spin numbers (^{1}H vs ^{2}H and ^{15}N vs ^{14}N). In both FAD$^{\bullet-}$ and

TrpH$^{\bullet+}$ radicals, the H and N isotope with I=1 (i.e., ^{2}H and ^{14}N) consistently experience higher

singlet yield along the whole inclination range than their counterparts with I=1/2.

The marked isotope-dependence of singlet yield might also offer a novel explanation

for isotope-induced oxidative stress, as reported in Zubarev [9]. The role that radical reaction

intermediates might play in the chemistry and biology of reactive oxygen species is a

growing field of research. Reactive oxygen species (ROS) include a wide variety of oxidant

molecules with different properties and biological functions that range from signalling to

oxidative damage [58]. In this context, the RP mechanism might give some insight.

Usselman et al. show how yields of ROS in live cells are changed by RP dynamics, in

particular coherent singlet-triplet mixing [59,60]. For an RP involving oxygen, the substitution

of isotopes, which we have here demonstrated to change singlet yield, could thus change

the balance of ROS, potentially causing oxidative stress. This is of direct relevance to the

migration evolution hypothesis posed here, because migration is a physiologically costly

strategy and, consequently, species of birds that conduct long-distance migration have

comparatively higher levels of antioxidant resources [61]. It is likely, then, that the possible

role of isotopes on the evolutionary origin of migration also leads to differences in oxidative

stress between migratory and non-migratory animals, something that should be explored in

the future. The role that isotopes play in RP-mediated changes to singlet yield have also

been investigated in a number of other biological contexts, including consciousness [62]. It

would thus seem important to determine how isotopes facilitate or inhibit the development

of biological functions that may confer some evolutionary advantage on an organism.

It is interesting to note, however, that the greater singlet yield of nuclei with I=1 does

not necessarily translate into greater magnetic sensitivity, as demonstrated by the results in

Figure 7, for the MS of the H and N isotopes investigated in $FAD^{\bullet-}$ and $TrpH^{\bullet+}$. We consider

in particular the range of external magnetic field values corresponding to the geomagnetic

field (25-65 µT). For both hydrogens, in contrast to the results given by the simple case

investigated in Scenario 1 (Figure 3), MS is improved by isotope substitution with I=1. This

is likely due to the hyperfine coupling environment. From the results in Figure 3 it appears

that nuclei with I=1 perform better at smaller coupling constants. Isotopic substitution in

hydrogen gives smaller coupling constants. However, further investigation would have to be

done before definite conclusions are drawn. The case for nitrogen and its isotopes is less

clear. Over the relevant geomagnetic window (25-65 µT), both I=1/2 and I=1 are more

favourable at certain values of the magnetic field. For the specific nitrogen N10, the isotopic

substitution with I=1/2 dramatically improves the MS at certain values of the field. While it is

difficult to draw any absolute conclusions, it is potentially interesting that for both hydrogen

and (at least partly) nitrogen, isotopic substitution acts to increase the MS. Although we have

investigated here only isotopes of H and N, differences in the potential to contribute to the

performance of cryptochrome as a magnetic compass are likely to exist also between

isotopes of the other constituent elements of the molecule (i.e., C and O), as well as any

element functioning in other biochemical processes (e.g., S and P). The fact that heavy

isotopes are less abundant in nature than their lighter counterparts (in the case of δ^2H and

δ^{15}N this being only 0.0115% and 0.364%, respectively, see Table 1) does not preclude the

possibility that differences in the isotopic composition of a biomolecule such as

cryptochrome between species of animals generate differences in the performance of the

involved biochemical process, despite the negligible effect of heavy isotopes on the speed

of chemical reactions that is predicted by conventional chemical kinetics. The Isotopic

Resonance Hypothesis, on the other hand, posits that optimal, 'resonance' abundances of

stable isotopes reduce the complexity of biochemical systems, affecting the kinetics of direct

and reversed reactions within the system and maximizing the efficiency of biochemical

reactions [13,14,63]. The Isotopic Resonance Hypothesis has received empirical support

from enzymatic reactions, also mediated by a quantum process such as quantum tunnelling

[64,65]. Thus, the enzymatic catalytic action of luciferase on its substrate luciferin with a

deuterium concentration in the local environment 2-4 times higher than the normal

concentration of 150 ppm but still considerably low (250-350 ppm) exhibits a significant

change that is not predicted by conventional chemical kinetics. This indicates that small

concentrations of heavy isotopes exert disproportionately strong effects on the kinetics of

enzymatic processes [66]. Similarly, animals with a cryptochrome composition enriched in

isotopes of, for example, hydrogen and nitrogen may have a superior magnetic sensitivity,

this not being hindered when such isotopes are heavy, such as deuterium (^{2}H).

Consequently, our results show that the isotopic composition of cryptochrome can

exert significant effects on the performance of RPs, hence influencing the likelihood of

developing an ability for magnetoreception and, thus, for migration. This means that the

evolution of migration may be favoured only under certain isotopic compositions of

cryptochrome. This has never been proposed before. But, indeed, non-adaptive origins of

adaptive traits are relevant sources of evolutionary innovations, particularly for metabolic

traits [67].

Physiology is considered to have a foundational role in migration, which has been

studied in detail (e.g., [68]). The evolutionary maintenance of migration is based on its

adaptive benefits, as it allows animals to get trophic resources that would otherwise be

inaccessible, and is also well understood. However, it is unknown whether differences in

migration among species are due to genetic or environmental effects [69,70]. Recently, a

study on a single migratory bird species (the peregrine falcon *Falco peregrinus*) has

revealed that divergence in the gene *ADCY8* explains differences in migration distance

between populations, but not why these birds always migrate in some extent [71]. As a

consequence, the ultimate factors that generally drive animal migration and that explain the

evolutionary appearance of this behaviour in some species and not in others remain

undeciphered [72]. Interspecific differences in the isotopic composition of cryptochrome may

therefore represent a general explanation for the origin of animal migration.

An effective test of our hypothesis will require empirical comparisons of cryptochrome

isotopologues between migratory and non-migratory species of birds or other animals.

Some already existing empirical results, however, may support our proposal. In fact, an

investigation on the isotopic composition of hair keratin in three species of coexisting bats

resulted in an isospace (δ^{13}C vs. δ^{15}N) of a long-distance migratory bat (*Pipistrellus nathusii*)

markedly distinct from the isospaces of non-migratory and mid-distance migratory bats

(*Pipistrellus pipistrellus* and *Nyctalus noctula*) [73]. Considering that the isotopic profile may

be consistent among all molecules of a species of animal (see Introduction), the latter results

may be in accordance with the hypothesis proposed here.

A causal association between animal isotopic profiles and their macroscopic

characteristics would represent a non-adaptive and non-organic source of evolutionary

innovations, and a connection between the mineral and the organic world, unveiling

attributes of organisms not carried by their genetic code. The transference of atomic

information across organisms would represent a DNA-independent inheritance that has

never been investigated, but future studies should investigate if this may in fact be the

evolutionary origin of animal migration and other biological processes.

Acknowledgements. We would like to thank Ilya Sinayskiy for his helpful discussions and comments during this work.

Funding. F.P. acknowledges support by the South African Research Chair Initiative of the Department of Science and Technology and the National Research Foundation.

Ethics. No ethical issues are associated with this study.

Data, accessibility. This article has no additional data.

Conflict of interest declaration. We declare we have no competing interests.

Authors' contributions. I.G. conceptualised the role of isotopes in organic evolution. A.H., B.A. and F.P. contributed to physics interpretation and model development. A.H. did the calculations and spin dynamics simulations. I.G., A.H., B.A. and F.P. wrote and reviewed the manuscript.

References

1. Simons JP. 2009 Good vibrations: probing biomolecular structure and interactions through spectroscopy in the gas phase. *Mol. Phys.* **107**, 2435-2458.

2. Yakushevich LV. 2004 *Nonlinear physics of DNA*. Weinheim, Germany: Wiley-VCH.

3. Lima-de-Faria A. 1997 The atomic basis of biological symmetry and periodicity. *BioSystems* **43**, 115-135.

4. Trixler F. 2013 Quantum tunnelling to the origin and evolution of life. *Curr. Org. Chem.* **17**, 1758-1770.

5. Galván I, Cerezo J, Jorge A, Wakamatsu K. 2018 Molecular vibration as a novel explanatory mechanism for the expression of animal colouration. *Integr. Biol.* **10**, 464-473.

6. Bigeleisen J. 1965 Chemistry of Isotopes: Isotope chemistry has opened new areas of chemical physics, geochemistry, and molecular biology. *Science* **147**, 463-471.

7. Epov VN, Malinovskiy D, Vanhaecke F, Bégué D, Donard OF. 2011 Modern mass
spectrometry for studying mass-independent fractionation of heavy stable isotopes in
environmental and biological sciences. *J. Anal. At. Spectrom.* **26**, 1142-1156.

8. Fry B. 2006 *Stable Isotope Ecology*. New York, NY: Springer Science+Business Media.

9. Zubarev RA. 2011 Role of stable isotopes in life—Testing isotopic resonance
hypothesis. *Genom. Proteom. Bioinform.* **9**, 15-20.

10. Caut S, Angulo E, Courchamp F. 2008 Caution on isotopic model use for analyses of
consumer diet. *Can. J. Zool.* **86**, 438-445.

11. Xie X, Zubarev RA. 2014 Effects of low-level deuterium enrichment on bacterial
growth. *PLoS ONE* **9**, e102071.

12. Basov A., Fedulova L., Baryshev M., Dzhimak S. 2019 Deuterium-depleted water
influence on the isotope ^{2}H/^{1}H regulation in body and individual adaptation. *Nutrients*
11, 1903.

13. Xie X, Zubarev RA. 2015 Isotopic resonance hypothesis: experimental verification by
Escherichia coli growth measurements. *Sci. Rep.* **5**, 9215.

14. Andriukonis E, Gorokhova E. 2017 Kinetic ^{15}N-isotope effects on algal growth. *Sci. Rep.*
7, 44181.

15. DeNiro MJ, Epstein S. 1978 Influence of diet on the distribution of carbon isotopes in
animals. *Geochim. Cosmochim. Acta* **42**, 495-506.

16. DeNiro MJ, Epstein S. 1981 Influence of diet on the distribution of nitrogen isotopes in
animals. *Geochim. Cosmochim. Acta* **45**, 341-351.

17. Olivar MP, Bode A, López-Pérez C, Hulley PA, Hernández-León S. 2019 Trophic
position of lanternfishes (Pisces: Myctophidae) of the tropical and equatorial Atlantic
estimated using stable isotopes. *ICES J. Mar. Sci.* **76**, 649-661.

18. Wißing C, Rougier H, Baumann C, Comeyne A, Crevecoeur I, Drucker DG et al. 2019.

Stable isotopes reveal patterns of diet and mobility in the last Neandertals and first

modern humans in Europe. *Sci. Rep.* **9**, 4433.

19. Manlick PJ, Petersen SM, Moriarty KM, Pauli JN. 2019 Stable isotopes reveal limited

Eltonian niche conservatism across carnivore populations. *Funct. Ecol.* **33**, 335-345.

20. Haak CR, Power M, Cowles GW, Danylchuk AJ. 2019 Hydrodynamic and isotopic niche

differentiation between juveniles of two sympatric cryptic bonefishes, *Albula vulpes* and

Albula goreensis. *Environ. Biol. Fishes* **102**, 129-145.

21. Moroz LL, Kocot KM, Citarella MR, Dosung S, Norekian TP, Povolotskaya IS et al. 2014.

The ctenophore genome and the evolutionary origins of neural systems. *Nature* **510**,

109-114.

22. Deng M, Zhang X, Luo J, Liu H, Wen W, Luo H et al. 2020 Metabolomics analysis

reveals differences in evolution between maize and rice. *Plant J.* **103**, 1710-1722.

23. Brookes JC 2017. Quantum effects in biology: golden rule in enzymes, olfaction,

photosynthesis and magnetodetection. *Proc. R. Soc. A* **473**, 20160822.

24. Marais A, Adams B, Ringsmuth AK, Ferretti M, Gruber JM, Hendrikx R et al. 2018. The

future of quantum biology. *J. R. Soc. Interface* **15**, 20180640.

25. Buchachenko AL. 2001 Magnetic isotope effect: Nuclear spin control of chemical

reactions. *J. Phys. Chem. A* **105**, 9995-10011.

26. Wiltschko W, Wiltschko R. 2005 Magnetic orientation and magnetoreception in birds

and other animals. *J. Comp. Physiol. A* **191**, 675-693.

27. Dreyer D, Frost B, Mouritsen H, Günther A, Green K, Whitehouse M et al. 2018. The

Earth's magnetic field and visual landmarks steer migratory flight behavior in the

nocturnal Australian Bogong moth. *Curr. Biol.* **28**, 2160-2166.

28. Ritz T, Thalau P, Phillips JB, Wiltschko R, Wiltschko W. 2004 Resonance effects

indicate a radical-pair mechanism for avian magnetic compass. *Nature* **429**, 177-180.

29. Günther A, Einwich A, Sjulstok E, Feederle R, Bolte P, Koch KW. et al. 2018 Double-cone localization and seasonal expression pattern suggest a role in magnetoreception for European robin cryptochrome 4. *Curr. Biol.* **28**, 211-223.

30. Xu J, Jarocha LE, Zollitsch T *et al.* 2021. Magnetic sensitivity of cryptochrome 4 from a migratory songbird. *Nature* **594,** 535–540.

31. Wiltschko R, Nießner C, Wiltschko W. 2021 The magnetic compass of birds: the role of cryptochrome. *Front. Physiol.* **12**, 667000.

32. Bazalova O, Kvicalova M, Valkova T, Slaby P, Bartos P, Netusil R et al. 2016. Cryptochrome 2 mediates directional magnetoreception in cockroaches. *Proc. Natl. Acad. Sci. USA* **113**, 1660-1665.

33. Foley LE, Gegear RJ, Reppert SM. 2011 Human cryptochrome exhibits light-dependent magnetosensitivity. *Nat. Commun.* **2**, 356.

34. Caspar KR, Moldenhauer K, Moritz RE, Němec P, Malkemper EP, Begall S. 2020 Eyes are essential for magnetoreception in a mammal. *J. R. Soc. Interface* **17**, 20200513.

35. Hogben HJ, Biskup T, Hore PJ. 2012 Entanglement and sources of magnetic anisotropy in radical-pair based avian magnetoreceptors. *Phys. Rev. Lett.* **109**, 220501.

36. Fay TP, Lindoy LP, Manolopoulos DE, Hore PJ. 2020 How quantum is radical pair magnetoreception? *Faraday Discuss.* **221**, 77-91.

37. Zhang Y, Berman GP, Kais S. 2015 The radical pair mechanism and the avian chemical compass: Quantum coherence and entanglement. *Int. J. Quantum Chem.* **115**, 1327-1341.

38. Hochstoeger T, Al Said T, Maestre D, Walter F, Vilceanu A, Pedron M. et al. 2020 The biophysical, molecular, and anatomical landscape of pigeon CRY4: A candidate light-based quantal magnetosensor. *Sci. Adv.* **6**, eabb9110.

39. Player TC, Hore PJ. 2019 Viability of superoxide-containing radical pairs as magnetoreceptors. *J. Chem. Phys.* **151**, 225101.

40. Tiersch M, Briegel HJ. 2012 Decoherence in the chemical compass: The role of
decoherence for avian magnetoreception. *Phil. Trans. R. Soc. A* **370**, 4517-4540.

41. Kattnig D. 2017 Radical-pair-based magnetoreception amplified by radical scavenging:
resilience to spin relaxation. *J. Phys. Chem. B* **121**, 10215-10227.

42. Hiscock HG, Worster S, Kattnig DR, Steers C, Jin Y, Manolopoulos DE, Mouritsen H,
Hore PJ. 2016 The quantum needle of the avian magnetic compass. *Proc. Natl. Acad.*
Sci. USA **113**, 4634-4639.

43. Jain R, Poonia VS, Saha K, Saha D, Ganguly S. 2021 The avian compass can be
sensitive even without sustained electron spin coherence. *Proc. R. Soc. A* **477**,
20200778.

44. Tiersch M, Steiner UE, Popescu S, Briegel HJ. 2012 Open quantum system approach
to the modeling of spin recombination reactions. *J. Phys. Chem. A* **116,** 4020-4028.

45. Ritz T, Wiltschko R, Hore PJ, Rodgers CT, Stapput K, Thalau P et al. 2009 Magnetic
compass of birds is based on a molecule with optimal directional sensitivity. *Biophys.*
J. **96**, 3451-3457.

46. Haberkorn R. 1976 Density matrix description of spin-selective radical pair reactions.
Mol. Phys. **32**, 1491–1493.

47. Jones JA, Hore PJ. 2010 Spin-selective reactions of radical pairs act as quantum
measurements. *Chem. Phys. Lett.* **488**, 90–93.

48. Kominis IK. 2009 Quantum Zeno effect explains magnetic-sensitive radical-ion-pair
reactions. *Phys. Rev. E* **80**, 056115.

49. Adams B, Sinayskiy I, Petruccione F. 2018 An open quantum system approach to the
radical pair mechanism. *Sci. Rep.* **8**, 15719.

50. Gauger EM, Rieper E, Morton JJL, Benjamin SC, Vedral V. 2011 Sustained quantum
coherence and entanglement in the avian compass. *Phys. Rev. Lett.* **106**, 040503.

51. Breuer HP, Petruccione F. 2002 *The theory of open quantum systems*. Oxford, UK: Oxford University Press.

52. Cai J, Caruso F, Plenio MB. 2012 Quantum limits for the magnetic sensitivity of a chemical compass. *Phys. Rev. A* **85**, 040304.

53. Johansson JR, Nation PD, Nori F. 2012 QuTiP: An open-source Python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **183**, 1760–1772.

54. Atkins PW, Friedman RS. 2010 *Molecular Quantum Mechanics. 5th Edition*. Oxford, UK: Oxford University Press.

55. Lee AA, Lau JCS, Hogben HJ, Biskup T, Kattnig DR, Hore PJ. 2014 Alternative radical pairs for cryptochrome-based magnetoreception. *J. R. Soc. Interface* **11**, 20131063.

56. Gerson F, Huber W. 2003 *Electron Spin Resonance Spectroscopy of Organic Radicals*. Weinheim, Germany: Wiley-VCH.

57. Bandyopadhyay JN, Paterek T, Kaszlikowski D. 2012 Quantum coherence and sensitivity of avian magnetoreception. *Phys. Rev. Lett.* **109**, 110502.

58. Sies H, Belousov VV, Chandel NS, Davies MJ, Jones DP, Mann GE et al. 2022 Defining roles of specific reactive oxygen species (ROS) in cell biology and physiology. *Nat. Rev. Mol. Cell Biol*. **23**, 499-515.

59. Usselman RJ, Hill I, Singel DJ, Martino CF. 2014 Spin biochemistry modulates reactive oxygen species (ROS) production by radio frequency magnetic fields. *PLoS ONE* **9**, e93065.

60. Usselman RJ, Chavarriaga C, Castello PR, Procopio M, Ritz T, Dratz EA et al. 2016 The quantum biology of reactive oxygen species partitioning impacts cellular bioenergetics. *Sci. Rep.* **6**, 38543.

61. Galván I, Erritzøe J, Karadaş F, Møller AP. 2012 High levels of liver antioxidants are associated with life-history strategies characteristic of slow growth and high survival rates in birds. *J. Comp. Physiol. B* **182**, 947-959.

62. Zadeh-Haghighi H, Simon C. 2022 Magnetic field effects in biology from the perspective of the radical pair mechanism. *J. R. Soc. Interface* **19**, 20220325.

63. Gorokhova E. 2017 Shifts in rotifer life history in response to stable isotope enrichment: testing theories of isotope effects on organismal growth. *R. Soc. Open Sci.* **4**, 160810.

64. Klinman JP, Kohen A. 2013 Hydrogen tunneling links protein dynamics to enzyme catalysis. *Ann. Rev. Biochem.* **82**, 471.

65. Pudney CR, Lane RS, Fielding AJ, Magennis SW, Hay S, Scrutton NS. 2013 Enzymatic single-molecule kinetic isotope effects. *J. Am. Chem. Soc.* **135**, 3855-3864.

66. Rodin S, Rebellato P, Lundin A, Zubarev RA. 2018 Isotopic resonance at 370 ppm deuterium negatively affects kinetics of luciferin oxidation by luciferase. *Sci. Rep.* **8**, 16249.

67. Barve A, Wagner A. 2013 A latent capacity for evolutionary innovation through exaptation in metabolic systems. *Nature* **500**, 203-206.

68. Lennox RJ, Chapman JM, Souliere CM, Tudorache C, Wikelski M, Metcalfe JD, Cooke SJ. 2016 Conservation physiology of animal migration. *Cons. Physiol.* **4**, cov072.

69. Cresswell KA, Satterthwaite WH, Sword GA. 2011 Understanding the evolution of migration through empirical examples. In: *Animal Migration: A Synthesis* (Eds. Milner-Gulland EJ, Fryxell JM, Sinclair ARE), pp. 7-16. Oxford, UK: Oxford University Press.

70. Dingle H. 2014 *Migration: the Biology of Life on the Move*. Oxford, UK: Oxford University Press.

71. Gu Z, Pan S, Lin Z, Hu L, Dai X, Chang J et al. 2021 Climate-driven flyway changes and memory-based long-distance migration. *Nature* **591**, 259-264.

72. Shaw AK, Couzin ID. 2013 Migration or residency? The evolution of movement behavior and information usage in seasonal environments. *Am. Nat.* **181**, 114-124.

73. Voigt CC, Lindecke O, Schönborn S, Kramer-Schadt S, Lehmann D. 2016 Habitat use of migratory bats killed during autumn at wind turbines. *Ecol. Appl.* **26**, 771-783.

Table 1. Nuclear isotope effect of nitrogen and hydrogen.

#	Name	Symbol	Z	N	A	Spin I parity	(μ in μ_N)	Nuclear g factor (g_n)	Natural abundance	Isotopic mass (u)
1	Hydrogen	H	1	0	1	1/2 +	+2.7928	+5.5857	99.9885 %	1.0078
2	Hydrogen	H (D)	1	1	2	1 +	0.8574	+0.8574	0.0115 %	2.0141
3	Nitrogen	N	7	7	14	1 +	0.4037	+0.4038	99.636 %	14.0030
4	Nitrogen	N	7	8	15	1/2 -	-0.2832	-0.5663	0.364 %	15.0001

μ in μ_N : Nuclear magnetic moment of spin I in h/2π.

Table 2. Hyperfine coupling tensors for the [FAD•− – TrpH+] radical system, taken from Hiscock et al. (2016). Following Jain et al. (2021), all hyperfine tensors are simultaneously diagonalized.

Nuclei	a_x (µT)	a_y (µT)	a_z (µT)
N10 in FAD•− radical (^{14}N)	- 24.1	- 14.4	604.6
N10 in FAD•− radical (^{15}N)	33.8	20.2	- 848.1
H6 in FAD•− radical (^{1}H)	- 530.4	- 433.6	- 197.6
H6 in FAD•− radical (^{2}H)	- 81.4	- 66.6	- 30.3
N1 in TrpH•+ radical (^{14}N)	- 63.6	- 53	1081.2
N1 in TrpH•+ radical (^{15}N)	89.2	74.3	- 1516.3
H1 in TrpH•+ radical (^{1}H)	1082.6	- 705.4	- 6.9
H1 in TrpH•+ radical (^{2}H)	166.2	- 108.3	- 1.06

Legends to figures:

Figure 1. The radical pair (RP) mechanism in cryptochrome may facilitate

magnetoreception in migratory animals. Within the protein cryptochome 4, a flavin

residue and a tryptophan form a dyad that, upon photoexcitation, gives rise to a flavin radical

($FH^{\bullet}$, or $FAD^{\bullet-}$) and a tryptophan radical ($W^{\bullet}$, or $TrpH^{\bullet+}$). Then the RP can either produce

photoproducts, or recombine to produce the original, zero-spin dyad molecule. The radical

pair is initially in singlet state, but hyperfine coupling leads to conversion into non-reactive

triplet states. The recombination reaction thus requires a triplet-singlet spin conversion,

which is a function of the magnetic field, the nuclear spin and its projection, the nuclear

magnetic moment, the hyperfine coupling constant, the frequency and amplitude of the

magnetic field, and the interaction between unpaired electrons [27]. As the nuclear spin

differs between isotopes, isotopes with specific I values experience specific hyperfine

coupling between the unpaired electron and the magnetic nucleus, potentially leading to fast

triplet-singlet conversion. This conversion is slower in isotopes with other I value. In the

flavin-tryptophan radical pair, the triplet-singlet conversion rate depends on the orientation

of the Earth's magnetic field, allowing orientation and migratory movements. Molecular and

Earth images taken from Dreyer et al. [27] (open access content). Swift image belongs to

Stuart Price (https://flic.kr/p/TLn66s) and is under CC BY-SA 2.0 license

(https://creativecommons.org/licenses/by-sa/2.0/).

Figure 2. Hyperfine patterns occurring during the replacement of a proton (H) by a

deuterium (D) and a ^{14}N nucleus (N) by its ^{15}N isotope. Following Gerson and Huber [56].

Figure 3. Modelling results for the magnetic sensitivity of cryptochrome ($\Delta\Phi_S$) for

different spin numbers along a range of anisotropic hyperfine coupling strengths (a).

a) Here, the only difference between the treatment of nuclei in the RP model lies in their spin

quantum numbers, meaning that we investigate MS with respect to the hyperfine coupling

of a single nucleus model for two different spin quantum numbers (I=1 and I=1/2). In this

simple case, we do not include noise in the model. The results demonstrate clear differences

between the two spin numbers, with I=1/2 giving the greatest magnetic sensitivity for the

greatest range of hyperfine coupling constants and the maximum sensitivity for also I=1/2 is

considerably greater than I=1. b) For hyperfine up to 1.5 mT, the sensitivity reaches a

plateau, which is equal to 0.16 for I=1 and 0.25 for I=1/2.

Figure 4. Modelling results for the magnetic sensitivity of cryptochrome ($\Delta\Phi_S$) for

different spin numbers along a range of anisotropic hyperfine coupling strength (a).

a) With noise incorporated into the model, it is shown that MS can be improved for certain

hyperfine coupling strengths. For a nucleus with I=1/2, the maximum MS increases from

0.388 (corresponding to a hyperfine coupling strength of 17.4 µT) to 0.411 (corresponding

to a hyperfine coupling strength of 20.9 µT). For a nucleus with I=1, the maximum MS

increases from 0.259 (corresponding to a hyperfine coupling strength of 8.7 µT) to 0.274

(corresponding to a hyperfine coupling strength of 10.6 µT). b) It is clear from the results

that the sensitivity of the avian compass relies on the recombination rate. A good sensitivity

requires a recombination rate that gives enough time for magnetic field effects to occur.

Here it is shown that sensitivity decreases when increasing the recombination rate, i.e.

shorter lifetime for the radical pair.

Figure 5. Isotope substitution effects on the magnetic effect (singlet yield, Φ_S) with

different inclinations of cryptochrome for a one-nucleus model. The results are

separately shown for the radicals of the RP system in cryptochrome (left panels: FAD$^{\bullet-}$, right

panels: TrpH$^{\bullet+}$) and for different hydrogen (a: H6, c: H1) and nitrogen (b: N10, d: N1) nuclei

and their isotopes. It is clear that isotopes with integer spin have consistently higher singlet

yield than half integer, even with different coupling constants. Singlet yield is an interesting

feature to consider with respect to RPs in the biological context. In addition to

magnetoreception, singlet yield has been used to investigate a number of other biological

functions which may depend on radical reactions [62]. In reactions involving reactive oxygen

species, singlet yield might be used as an indication of oxidative stress. The strong isotope

dependence of singlet yield demonstrated in our results is potentially interesting with respect

to observations made that isotopic changes in diet lead to oxidative stress [9].

Figure 6. Isotope substitution effects on the magnetic effect (singlet yield, Φ_S) with

different inclinations of cryptochrome for two-nuclei model. The results for the four

different combinations of hydrogen and nitrogen isotopes for the two-nuclei model are

shown. Results are relatively similar to those of the single-nucleus model (Figure 5), with a

minimum at about 90º singlet yield. Singlet yield is greatest for the case in which both nuclei

have integer spin, that is for N14 and deuterium.

Figure 7. Isotope substitution effects for the magnetic sensitivity of cryptochrome

($\Delta\Phi_S$) along a range of strength values of an external magnetic field (B). Examples of

different nuclei in the radicals of the RP system in cryptochrome are shown (from to down

panels: H6 in FAD$^{\bullet-}$, N1 in TrpH$^{\bullet+}$, H1 in TrpH$^{\bullet+}$, N10 in FAD$^{\bullet-}$). **Scenario 1** (see Figure 3)

showed a clear difference between how spin number changes MS, with $I=1/2$ conferring

greater MS than $I=1$. However, the effect of the spin number is also dependent on the

hyperfine coupling strength, with $I=1/2$ performing best at larger coupling constants than $I=1$.

In **Scenario 2** it is more difficult to conclude that there is a favourable spin number with

respect to MS. This is likely due to the influence of the specific hyperfine coupling constants.

While the hyperfine coupling strength decreases with isotope substitution in hydrogen, it

increases with isotope substitution in nitrogen. If we consider only the magnetic field strength

relevant to migration, which is the geomagnetic field (25-65 µT), then for the case of both

hydrogens, $I=1$ gives greater magnetic sensitivity than $I=1/2$. The case is less clear for

nitrogen where both $I=1/2$ and $I=1$ show greater MS for specific magnetic field strengths

over the geomagnetic window (25-65 µT). This could be due to the fact that although the

coupling strengths for the nitrogen isotopes are further from the ideal values, the isotopic

substitution from $I=1$ to $I=1/2$ confers some advantage.

Figure 1

singlet

P

triplet

θ

N

S

FH•

W•

FH•

W•

Figure 2

$H = {}^{1}H$ $|a_H|$

$D = {}^{2}H$ $|a_D|$

$N = {}^{14}N$ $|a_N|$

$N' = {}^{15}N$ $|a_{N'}|$

a)

One Nucleus model (Sensitivity vs HFC)

b)

One Nucleus model, I=1 vs I=1/2

Figure 4

Figure 6

Singlet Yield vs Theta (N10 + H6) in FAD^-