Dynamics of Pulsatile Activities of Arcuate Kisspeptin Neurons in Aging Female Mice

Teppei Goto[1,*] and Kazunari Miyamichi[1,*]

[1] Laboratory for Comparative Connectomics, RIKEN Center for Biosystems Dynamics Research, Kobe, Hyogo 650-0047, Japan.

Short title: Kisspeptin pulse in aging female mice
Keywords: Kisspeptin, Pulse, Fiber photometry, Aging, Menopause

Correspondence: Teppei Goto, E-mail: teppei.goto@riken.jp; Kazunari Miyamichi, E-mail: kazunari.miyamichi@riken.jp

Financial support: This study was supported by Grants-in-Aid to TG (Nos. 19K16270 and 21K15194) and KM (Nos. 20K20589 and 21H02587) from the Japan Society for the Promotion of Science (JSPS). TG is also supported by the RIKEN Special Postdoctoral Researchers Program.

Disclosures: The authors have nothing to disclose and no competing interests.

Categories: BIOLOGICAL SCIENCES, Neuroscience

Significance Statement
The transition to reproductive senescence significantly impacts the quality of life of women, but the underlying neural mechanisms remain poorly characterized. Goto and Miyamichi establish long-term (from the reproductive to acyclic phase) chronic imaging of the central pacemaker activities of reproductive functions by fiber photometry in female mice. In particular, they focus on kisspeptin neurons in the arcuate nucleus of the hypothalamus. Their data reveal that during the transition to reproductive senescence, the pulsatile activities of kisspeptin neurons show unexpected robustness in terms of frequency, but a tendency for the intensity to decline. Their findings exhibit the power of direct chronic visualization of hormonal regulators in the brain, which is generally applicable to facilitate studies of aging-associated malfunctions in neuroendocrine systems.

Abstract
Reproductive senescence is broadly observed across mammalian females, including humans, eventually leading to a loss of fertility. The pulsatile secretion of gonadotropin-releasing hormone (GnRH), which is essential for gonad function, is primarily controlled by kisspeptin neurons in the hypothalamic arcuate nucleus (ARC^{kiss}), the pulse generator of GnRH. The pulsatility of GnRH release, as assessed by the amount of circulating gonadotropin, is markedly reduced in aged animals, suggesting that the malfunctions of ARC^{kiss} may be responsible for reproductive aging and menopause-related disorders. However, the activity dynamics of ARC^{kiss} during the natural transition to reproductive senescence remain unclear. Herein, we introduce chronic *in vivo* Ca^{2+} imaging of ARC^{kiss} in female mice by fiber photometry to monitor the synchronous episodes of ARC^{kiss} (SEs^{kiss}), a known hallmark of GnRH pulse generator activity, from the fully reproductive to acyclic phase over 1 year. During the reproductive phase, we find that not only the frequency, but also the intensities and waveforms of individual SEs^{kiss}, vary depending on the stage of the estrus cycle. During the transition to reproductive senescence, the integrity of SEs^{kiss} patterns, including the frequency and waveforms, remains mostly unchanged, whereas the intensities tend to decline. These data illuminate the temporal dynamics of ARC^{kiss} activities in aging female mice. More generally, our findings demonstrate the utility of fiber-photometry-based chronic imaging of neuroendocrine regulators in the brain to characterize aging-associated malfunction.

Introduction
Reproductive aging occurs broadly across mammalian females, eventually leading to loss of fertility (Kermath and Gore, 2012). In humans, women undergo the perimenopausal transition to reproductive senescence, in which the menstrual cycles become irregular in length and timing before eventually ceasing at approximately 51 years of age (Gore, 2015). As this transition is associated with a risk of metabolic, physical, and mental disorders (Greendale et al., 1999), it is essential to understand the molecular and cellular mechanisms of reproductive aging for not only reproductive medicine, but also women's health in general.

Reproductive function in females is primarily controlled by coordinated interactions among the three layers of the hypothalamus–pituitary–ovary axis (Gore, 2015; Kermath and Gore, 2012). At the core, gonadotropin-releasing hormone (GnRH)-producing neurons in the anterior hypothalamus provide the driving force of the reproductive system. GnRH release occurs in a pulsatile manner at intervals of 30 minutes to several hours, depending on the species and reproductive (estrus) cycle (Herbison, 2018), and this pulsatility is essential for reproductive functions (Belchetz et al., 1978). GnRH released at the median eminence acts on the anterior pituitary gonadotrophs, which then release luteinizing hormone (LH) and follicle-stimulating hormone (FSH) into the peripheral circulation. Pulsatile (tonic) LH and FSH release facilitate follicular development and steroidogenesis in the ovary. The sex steroid hormones then exert

feedback control of the hypothalamic GnRH system and pituitary gonadotrophs. Although age-related changes can occur in each of these layers, classical evidence in rodent models suggests that modulation of GnRH neurons may be a major driving force of reproductive senescence. For example, pulsatile LH release as a proxy of GnRH activities decreases with age in rats (Scarbrough and Wise, 1990; Wise et al., 1988). As such, the preovulatory LH surge is delayed or attenuated with aging (Matt et al., 1998) with decreased c-Fos expression, a marker of neural activation, in the GnRH neurons (Rubin et al., 1994). Together with a classical transplantation study showing the presence of viable follicles in aged rodents (Krohn, 1962), these studies suggest that reproductive senescence is mediated by the neuroendocrine system independent of the depletion of the follicular reserve.

How is the homeostatic regulation of the GnRH system altered by aging? Of many candidates for GnRH regulation, kisspeptin, a neuropeptide crucial for puberty (d'Anglemont de Tassigny et al., 2007; Lapatto et al., 2007; Uenoyama et al., 2015), is thought to play a dominant role (Kermath and Gore, 2012). There are two populations of kisspeptin neurons in the rodent hypothalamus: one in the arcuate nucleus (ARC^{kiss}), which acts as the GnRH pulse generator (Clarkson et al., 2017; Nagae et al., 2021), and the other in the anteroventral periventricular nucleus ($AVPV^{kiss}$), which functions as the GnRH surge generator (Clarkson et al., 2008; Matsuda et al., 2019; Piet et al., 2018; Uenoyama et al., 2015). Previous studies have shown a decrease in kisspeptin signaling in both ARC^{kiss} (Kunimura et al., 2017) and $AVPV^{kiss}$ (Lederman et al., 2010; Neal-Perry et al., 2009) during aging in rats. Thus, malfunctions of the kisspeptin system may underpin the transition to reproductive senescence.

One critical step to characterizing the kisspeptin system for reproductive aging and menopause-related disorders (Padilla et al., 2018) is to visualize directly the dynamics of neural activities of kisspeptin neurons during aging. Until recently, activities of the GnRH pulse generator could only be indirectly assessed by repetitive measurement of circulating LH level (Steyn et al., 2013), which is not only labor-intensive, but also stressful to animals not fully suited for chronic analysis. Recent seminal studies have established cell-type-specific chronic Ca^{2+} imaging of ARC^{kiss} activities under a free-moving awake condition in mice by using fiber photometry (Clarkson et al., 2017; Han et al., 2019; McQuillan et al., 2019). ARC^{kiss} show remarkable pulsatile activities, termed synchronous episodes of elevated Ca^{2+} (hereafter referred to as SEs^{kiss} for simplicity). SEs^{kiss} are well correlated with pulsatile LH secretion (Clarkson et al., 2017), tightly regulated by gonadal negative feedback signals mediated by sex steroid hormones (Han et al., 2019), and undergo dynamic changes in frequency depending on the stage of the estrus cycle (McQuillan et al., 2019). However, the utility of this system is currently limited to only reproductive-phase or gonadectomized mice. It remains unknown how this system can facilitate the study of reproductive senescence by long-term imaging of SEs^{kiss} dynamics. This motivated us to apply fiber photometry to monitor SEs^{kiss} from the fully reproductive to acyclic phase over 1 year.

Results
Dynamics of SEs^{kiss} in the regular estrus cycle at the reproductive phase
Before characterizing SEs^{kiss} during aging, we revisited the detailed patterns and waveforms of SEs^{kiss} in the reproductive phase. We first established 7-day chronic imaging of ARC^{kiss} of cyclic *Kiss1-Cre* (Gottsch et al., 2011) female mice at age 4–6 months. An optical fiber was implanted above the ARC of *Kiss1-Cre* female mice that had been injected with an adeno-associated virus (AAV) driving Cre-dependent GCaMP6s (Chen et al., 2013) into the ARC. Their estrus cycle stages were determined by vaginal cytology (Byers et al., 2012; Caligioni, 2009), and Ca^{2+} imaging was performed twice per day for 6 hours each in the light and dark period with 6-hour inter-imaging intervals. Estrus cycle and Ca^{2+} imaging from ARC^{kiss} were monitored for 7 consecutive days. We detected sharp photometric peaks highly resembling the

previously reported SEs[kiss] (Clarkson et al., 2017) (Fig. 1A and Materials and Methods). The number of SEs[kiss] was dynamically altered in various stages of the estrus cycle and light and dark periods (Fig. 1B). To align data from multiple animals, we utilized the fact that SEs[kiss] are strongly suppressed by a high level of progesterone just before ovulation in the proestrus stage (McQuillan et al., 2019). Indeed, when the number of SEs[kiss] in the dark period was normalized to that of the preceding light period, proestrus just after the diestrus stage showed a stereotyped drop (<0.5) compared with other stages of the estrus cycle (Fig. 1C). Based on this profile, the estrus cycle was aligned starting from the proestrus stage. The number of SEs[kiss] during the estrus cycle was lowest at the dark period of the proestrus, and then gradually increased as the estrus cycle advanced, reaching the highest values at the dark period of the metestrus and diestrus (Fig. 1D). As expected from this frequency of SEs[kiss], the inter-peak intervals showed the opposite trend (Fig. S1A). We also found that the peak height of SEs[kiss] normalized to that in the light period of proestrus fluctuated during the regular estrus cycle, with the highest (approximately 1.5-fold) at the light period 2 days after proestrus (Fig. 1E). These data show that not only the frequency (McQuillan et al., 2019), but also the intensities of individual SEs[kiss], vary precisely depending on the stage of the estrus cycle. The higher peak during metestrus may be correlated with a higher amount of kisspeptin released to GnRH neurons to facilitate more LH/FSH secretion (Czieselsky et al., 2016).

We then quantified detailed waveforms of individual SEs[kiss] in each stage of the estrus cycle (Fig. 1F, Fig. S1B). Although the overall shape was similar, when the peak was aligned, the onset and offset in proestrus significantly advanced compared with the estrus/metestrus stage 2 days after proestrus (Fig. 1G). To confirm this trend further, we quantified left half-width at quarter maximum (LWQM) and right half-width at quarter maximum (RWQM). Whereas the full-width at quarter maximum (FWQM) was constant across all stages of the estrus cycle, LWQM in the proestrus stage was significantly longer than that in the other stages (Fig. 1H). RWQM in the proestrus stage tended to be shorter, while that in estrus/metestrus was significantly longer. Variations in waveforms correlated with the peak height: when the peak was higher, the upslope was steeper and the downslope was gentler. This difference likely reflects the activity dynamics of individual ARC[kiss] during each SE[kiss] (see Discussion).

Reduced estrus cyclicity with aging in *Kiss1-Cre* mice
Next, we characterized the progress of reproductive decline in our experimental animals (*Kiss1-Cre* females in C57BL/6 background). In rodents, the most revealing marker of the reproductive stage is the estrous cyclicity determined by a vaginal smear, which is an indicator of the ovarian and neuroendocrine environments (Byers et al., 2012; Caligioni, 2009). A longitudinal study of estrus cyclicity in aging rodents revealed age-related reduction of cycle frequency and changes in cytology (LeFevre and McClintock, 1988; Nelson et al., 1982). Middle-aged female rodents can be classified into three groups: regularly cycling, irregularly cycling, and acyclic categories. We first confirmed that the number of estrus cycles per 25 days of wild-type C57BL/6J females was gradually decreased in an age-dependent manner (Fig. 2A), generally consistent with a preceding study (Nelson et al., 1982). A similar trend was observed in *Kiss1-Cre* female mice that had been injected with an AAV driving Cre-dependent GCaMP6s into the ARC with an optic fiber implanted above it, indicating that the genetic manipulation and fiber photometry setting did not alter the reproductive decline process (Fig. 2B, $n = 12$). The average number of estrus cycles per 25 days substantially declined between age 12–13 months, supporting the notion that female mice begin to show irregular cyclicity at around 13 months of age. Based on these data, we set the threshold at three estrus cycles per 25 days and classified middle-aged female mice into one of three groups: regular cyclic (three or more), irregular cyclic (two or one), or acyclic (zero).

Dynamics of SEs^kiss during the transition to reproductive senescence

To characterize SEs^kiss during aging, we first chronically monitored the same female mice ($n =$ 2) from age 6 months (fully reproductive) to age 15–18 months (after reproductive senescence). To this end, we repeatedly conducted 7-day chronic imaging of ARC^kiss together with a vaginal smear-based analysis of the estrus cycle (Fig. 3). At a glance, the frequency of SEs^kiss was not drastically changed, whereas the intensity of SEs^kiss exhibited a decline across the transition to reproductive senescence (Fig. 3A and 3B). Quantitative analysis showed that the frequency of SEs^kiss as assessed by the average number per 6 hours was constant (Fig. 3C), with no difference in the cumulative probability of inter-peak intervals among data set from age 6 months (fully reproductive) to age 15–18 months (acyclic in aging) (Fig. 3D). Notably, the elevated frequency of SEs^kiss generally observed in ovariectomized females (Clarkson et al., 2017), which is thought to indicate a drastic decline of sex steroid hormone-mediated negative feedback (Padilla et al., 2018), was not found in the natural transition to reproductive senescence. We also found that the waveforms of individual SEs^kiss were grossly unchanged from the reproductive to acyclic phase, with only slight individual differences found in the aging phases (Fig. S2A and S2B). In contrast to the relatively stable frequency of SEs^kiss, quantitative analysis showed that the peak height as assessed by $\Delta F/F$ values was significantly decreased during the transition to the acyclic phase in these two females (Fig. 3E).

For further quantification, we collected more photometry data from 13–16-month-old *Kiss1-Cre* female mice and classified them into three groups: regular cyclic ($n = 4$), irregular cyclic ($n = 3$), and acyclic ($n = 3$) (Fig. 4A). Although a stereotyped drop in the ratio of SEs^kiss in the dark period normalized to those in the preceding light period was evident at the proestrus stage in young (reproductive) female mice (Fig. 1C), no such trend was observed in all three groups of aging mice (Fig. 4B), which suggests an alteration of negative feedback by sex steroid hormones. We also noticed a slight reduction in the frequency of SEs^kiss in the irregular compared with the regular cyclic group (Fig. 4C). In addition, in the irregular cyclic group, the number of SEs^kiss in the light period was significantly lower than that in the dark period. These subtle differences were not evident in the chronic analysis of $n = 2$ females (Fig. 3), probably because of high data variations. Consistent with the individual analysis (Fig. 3D), no difference in the cumulative probability of inter-peak intervals was found among the reproductive group (age 4–6 months) or three aging groups (Fig. 4D). Again, no increase in the frequency of SEs^kiss was observed in any of the three aging groups. Collectively, these data demonstrate that the frequency of SEs^kiss during the transition to reproductive senescence is overall stable.

We also analyzed whether the variability of the peak height would change in the aging female mice by assessing the coefficient of variance (Fig. S2C). We noticed slight individual differences in the peak height variance in the regular and irregular cyclic aging groups; however, no significant inter-group differences were found. Regarding the intensities of SEs^kiss, our additional analysis (Fig. 4E, and ID#5, 6 in Fig. 3E) showed that the peak height declined from the regular to the irregular cyclic (two of four individuals) and acyclic phases (all four individuals). Therefore, our data generally support the notion that in aged female mice, the intensity of SEs^kiss is reduced, whereas their frequency, to a lesser extent, is altered.

Discussion

Although reproductive aging significantly impacts the quality of life of women, its underlying neural mechanisms remain poorly characterized. We have established long-term chronic imaging of ARC^kiss by fiber photometry in female mice from the fully reproductive phase to the end of the transition to reproductive senescent over 1 year. Our data illuminate the temporal dynamics of SEs^kiss during the estrus cycle in the reproductive phase, and during the transition to reproductive senescence. Our strategy of direct chronic visualization of hormonal regulators in the brain is generally applicable to facilitate characterizations of aging-associated robustness

and/or malfunctions in each component of neuroendocrine systems. Here, we discuss specific insights and major limitations of the present to guide the design of future investigations.

We updated the frequency and waveforms of individual SEs^{kiss} in the young reproductive phase with a detailed analysis of the light/dark periods of all four stages of the estrus cycle (Fig. 1). First, we reproduced a previous report showing a stereotyped drop of SEs^{kiss} in response to a high level of progesterone just before ovulation in the proestrus stage (Fig. 1C) (McQuillan et al., 2019). Second, we found that the frequency of SEs^{kiss} varied gradually depending on the stage of the estrus cycle (Fig. 1D), in contrast to a bimodal frequency reported previously (McQuillan et al., 2019). Third, we noticed that the intensities of SEs^{kiss}, which have been thought to be constant across the estrus cycle (McQuillan et al., 2019), significantly varied during the estrus cycle (Fig. 1E); individual peak heights of SEs^{kiss} increased by almost 1.5-fold in the metestrus compared with the proestrus stage. This trend is consistent with the fact that the amplitude of the LH pulse is highest in the metestrus (Czieselsky et al., 2016). The enhanced activities of the GnRH pulse generator may be beneficial to provide LH/FSH more efficiently at the onset of follicular recruitment, which is sensitive to these hormones (McGee and Hsueh, 2000). What would be the underlying mechanism of these elevated activities? In one scenario, the number of ARC^{kiss} that contributes to SEs^{kiss} is constant, but individual ARC^{kiss} fire more. Alternatively, more ARC^{kiss} are dynamically recruited to the SEs^{kiss} in the metestrus. To distinguish between these two possibilities, future studies using new tools that allow imaging or recording of neural activities at single-cell resolution (Moore et al., 2022; Steinmetz et al., 2021) in gonad-intact cycling female mice are needed.

We also found that the waveforms of individual SEs^{kiss} varied depending on the stage of the estrus cycle. In the metestrus stage, when the peak was higher, the uphill was significantly steeper (Fig. 1F–1H). What would be the underlying mechanism? A recent Ca^{2+} imaging study of ARC^{kiss} at single-cell resolution in ovariectomized female mice revealed heterogeneity of ARC^{kiss} activations during each SE^{kiss}: some subsets of ARC^{kiss} act as "leaders" and others as "followers" (Moore et al., 2022). Given these temporal dynamics, in the metestrus stage, the number of "leaders" may be increased to achieve quicker activation of the entire pool. Alternatively, the "leaders" more efficiently activate "followers" in the metestrus stage so that the latency to the peak is short. Although the exact mechanism remains an open question, our data suggest that the molecular and/or cellular mechanisms by which synchronized activation of the ARC^{kiss} pool is achieved are dynamically modulated by the estrus stage, likely via feedback signals mediated by sex steroid hormones.

Our results also showed that the frequency of SEs^{kiss} during reproductive senescence was generally stable. Classical studies suggest that the reduced reproductive functions of aging rodents are primarily the result of the dysfunction of the neuroendocrine system, independent of follicular depletion (Kermath and Gore, 2012; Krohn, 1962). There is a reduction of intensities of pulsatile LH release (Scarbrough and Wise, 1990; Wise et al., 1988) and a reduction of kisspeptin expression in ARC^{kiss} (Kunimura et al., 2017) in aged rats. In addition, a marked increase in the frequency of SEs^{kiss} was observed in gonadectomized mice in the reproductive phase (Clarkson et al., 2017; Han et al., 2019) via a decline of sex steroid hormone-mediated feedback mechanisms. Based on these results, it was possible to assume an abnormal frequency of SEs^{kiss} in older females; however, our data did not support this prediction. How is the relatively unaltered frequency of SEs^{kiss} during the natural transition to reproductive senescence achieved? In one scenario, the levels of sex steroid hormones may be grossly intact, even in the acyclic phase (Mobbs et al., 1985; Nilsson et al., 2015). Alternatively, the reduction of sex steroid hormones in aging female mice (Gee et al., 1983; Nelson et al., 1981) may be counterbalanced by the attenuation of negative feedback mechanisms in ARC^{kiss}. The lack of the stereotyped drop of SEs^{kiss} during the proestrus stage in older females (Fig. 4B) generally supported a malfunction of negative feedback mechanisms by sex steroid hormones.

Whatever the scenario, the lack of increased SEs^{kiss} frequency in aged female mice might be adaptive in terms of thermoregulation because elevated SEs^{kiss} are thought to be causal to vasodilation (flushing) (Padilla et al., 2018; Rance et al., 2013), one of the most commonly observed symptoms of perimenopausal disorders in women (Casper et al., 1979). In this sense, our data generally support the limitations in the use of older female mice as a model of the human menopausal transition, because not only the ovarian environment, but also the adaptation of the hypothalamic regulators, may vary considerably between rodents and primates (Gore, 2015; Kermath and Gore, 2012).

Our data also suggest that the intensities of SEs^{kiss}, as assessed by the peak height, gradually declined from the regular to the acyclic phase (Fig. 3E and Fig. S2B). Although we do not exclude the possibility that the reduction of photometric peaks was observed owing to technical artifacts, such as degradation of GCaMP6s or a slight misaligning of the optic fiber during long-term chronic imaging, we noticed that a drop in the photometric signals occurred in association with the reproductive phase rather than the total duration of fiber photometry. For instance, the peak intensity of ID#5 was relatively stable over 10 months from the reproductive to the irregular phase, but quickly declined in the subsequent 2 months upon the transition to the acyclic phase (Fig. 3A, E). By contrast, ID#10 exhibited a quick drop of the peak intensity within just 3 months after the surgery upon entering into the acyclic phase (Fig. S2B). These observations support the interpretation that reduced photometric signals are associated with reduced neural activities of ARC^{kiss}—individual firing rates, number of active neurons, or both. This trend, together with reduced *Kiss1* mRNA expression (Kunimura et al., 2017), can explain reduced LH amounts in each pulse (Scarbrough and Wise, 1990; Wise et al., 1988), leading to inefficient follicular development and steroidogenesis in the ovary. The activity decline of individual ARC^{kiss} with the molecular underpinnings should be better characterized in future studies.

Materials and Methods
Animals
All animal experiments were approved by the Institutional Animal Care and Use Committee of the RIKEN Kobe Branch. C57BL/6J female mice were purchased from Japan SLC (Shizuoka, Japan), and *Kiss1*-Cre mice (Jax #017701) were purchased from the Jackson Laboratory. All animals were maintained at the animal facility of the RIKEN Center for Biosystems Dynamics Research (BDR) under an ambient temperature (18–23 °C) and a 12-h light/12-h dark cycle schedule. All mice had ad libitum access to a laboratory diet (MFG; Oriental Yeast, Shiga, Japan; 3.57 kcal/g) and water.

Estrus cycle monitoring by vaginal cytology
To prevent the induction of pseudopregnancy, 10-μl of tap water was gently pipetted once or twice into the vagina with minimum insertion. These vaginal smears were placed on slides and evaluated in the types of cells under the microscope described below. The stage of the estrous cycle was determined based on the presence or absence of leukocytes and cornified and nucleated epithelial cells in accordance with a previous study (Nelson et al., 1982). Mostly nucleated and some cornified epithelial cells are present in proestrus. Some leukocytes are present if the female is in early proestrus. As the stage where the estrus cycle progresses from proestrus to estrus, mostly cornified epithelial cells are present in a relatively larger amount of vaginal smear. These cornified epithelial cells with large amounts of leukocytes are observed in early metestrus. In late metestrus, there are some nucleated epithelial cells in large amounts of leukocytes. In diestrus, there is the lowest amount of vaginal smear that contains leukocytes dominantly and a few epithelial cells.

Stereotaxic injection

For injection of AAV and an optical fiber, *Kiss1*-Cre mice were anesthetized with an intraperitoneal injection of saline containing 65 mg/kg ketamine (Daiichi Sankyo) and 13 mg/kg xylazine (X1251; Sigma-Aldrich) and head-fixed to a stereotaxic apparatus (RWD; #68045). Cre-dependent AAV encoding GCaMP6s (AAV9 *CAG-FLEx-GCaMPs-WPRE-SV40*, 1.5×10^{13} gp/ml; Addgene, #100842-AAV9) was injected into the ARC using the following coordinates from the bregma: anteroposterior (AP) = –2.0 mm, mediolateral = –0.2 mm, and dorsoventral (DV) = –5.9 mm. A total of 200 nL of AAV was injected into the ARC at a speed of 50 nl/minute using a glass capillary regulated by a microsyringe pump (UMP3; World Precision Instruments). At 2–4 weeks after AAV injection, an optical fiber (numerical aperture = 0.50, core diameter = 400 μm; Kyocera) was placed over the ARC using the following coordinates from the bregma: AP = –1.8 mm, lateral = –0.2 mm, and DV = –5.7 mm defined on the brain atlas (Franklin and Paxinos, 2013). The injected fiber was fixed on the skull with dental cement (SUN MEDICAL, Shiga, Japan). Postsurgical mice were singly housed.

Fiber photometry recording

Fluorescence signals were acquired using a fiber photometry system based on a previously published design (Clarkson et al., 2017; Han et al., 2019; McQuillan et al., 2019). All optical components were purchased from Doric Lenses (Quebec, Canada). We performed chronic Ca^{2+} imaging by delivering excitation lights (470-nm modulated at 530.481 Hz) and collection of emitted fluorescence by using the integrated Fluorescence Mini Cube (iFMC4_IE(405)_E(460-490)_F(500-550)_S; Doric Lenses). Light collection, filtering, and demodulation were performed using Doric photometry setup and Doric Neuroscience Studio Software (Doric Lenses). The power output at the tip of the fiber was about 4–40 μW. The signals were initially acquired at 12 kHz and then decimated to 120 Hz for recording to disk.

Calcium signals were analyzed using a custom-made code in R (version 4.0.3, http://www.R-project.org/). Because the signal intensity of SEs^{kiss} varies widely owing to variations in surgery and individual variability, we first calculated the average ΔF/F height of the stereotyped SE^{kiss} peak of each mouse from several visually obvious peaks with the full width at half maximum threshold over 10 seconds (Clarkson et al., 2017; McQuillan et al., 2019). The ΔF/F was calculated by $(F_t – F_0) / F_0$, where F_t is the recorded signal at time = t and F_0 is the average of signals over the entire 6 hours of recording (Figs. 1A, 3A). Then, SEs^{kiss} were automatically detected by using the findpeaks function in R, with a peak height threshold of 40% of the estimated peak height. To analyze the reliable peak height and waveform further (Figs. 1E, 3E, 4E), we extracted the raw data from –240 to +120 seconds, when the peak was set to 0 seconds. We then recalculated the ΔF/F by $(F_t – F_0) / F_0$, where F_t is the recorded signal at time = t and F_0 is the average of signals from –240 to –120 seconds. The FWQM, LWQM, and RWQM were determined by the recalculated ΔF/F value (Fig. 1H). To show the averaged waveform (Figs. 1F, 1G; Figs. S1B, S2A), the recalculated ΔF/F data were downsized at 10 Hz.

Statistical analysis

The statistical details of each experiment, including the statistical tests used, the exact value of *n*, and what *n* represents, are described in each figure legend. Two-way analysis of variance (ANOVA) was applied to confirm the interaction effects in Fig. 4C. One-way repeated measures ANOVA followed by the Tukey–Kramer post hoc test were applied to determine significant differences in Figs. 1D, 1E, 1H, and Fig. S1A. One-way ANOVA followed by the Tukey–Kramer post hoc test was applied to determine significant differences in Figs. 2A, 3C, and 4C. The Kruskal–Wallis test followed by the Mann–Whitney *U* test was applied to determine significant differences in Figs. 3E, S2B, and S2C. The Mann–Whitney *U* test was applied to determine significant differences in Fig. 4E. The Kolmogorov–Smirnov test was

applied to determine significant differences in Figs. 3D and 4D. Two-tailed Student's t-test was applied in Fig. 1G. A paired t-test was applied in Fig. 4C. All statistical analyses were performed using R. Differences were considered to be significant when the P-values were < 0.05.

Data and code availability
- Data reported in this paper are available from the authors upon reasonable request.
- No original code was generated in the course of this study.
- Any additional information required to reanalyze the data reported in this work paper is available from the authors upon request.

Figures

Figure 1. Dynamics of SEs^{kiss} in the regular estrus cycle of the reproductive phase. (**A**) Representative 6-h photometry raw data of SEs^{kiss} in each stage of the estrus cycle. Red arrowheads indicate SEs^{kiss}. (**B**) Numbers of SEs^{kiss} (lower: line plots) along with the estrus cycle stages (upper: color bars) for 7 days (*n* = 4 animals). SEs^{kiss} were recorded for 6 h in both light and dark (LD) periods. (**C**) Ratio of SEs^{kiss} in each dark period normalized to those in the preceding light period. Dotted line = 0.5. (**D**) Numbers of SEs^{kiss} per 6 h in 5 days along with the estrus cycle anchoring at proestrus as Day 0. Different letters (a–c) in the upper part of the graph denote significant differences at *P* < 0.05 by one-way repeated measures ANOVA followed by the Tukey–Kramer post hoc test. (**E**) Intensities of SEs^{kiss} as measured by the ΔF/F values in each estrus stage normalized to that in the light period of the proestrus. **P* < 0.05 by one-way repeated measures ANOVA followed by the Tukey–Kramer post hoc test. (**F**)

Averaged traces of SE^{kiss} waveforms in the light period of each estrus stage. (**G**) Top, averaged traces magnifying the onset and offset of SEs^{kiss} in the proestrus and estrus/metestrus stages in the light period. Data are expressed as mean ± SEM. Bottom, *P*-value by *t*-test. Red dots represent *P*-values < 0.05. (**H**) Quantifications of detailed parameters of SE^{kiss} waveforms in each estrus stage: the normalized LWQM, FWQM, and RWQM. *P < 0.05 by one-way repeated measures ANOVA followed by the Tukey–Kramer post hoc test. P: proestrus, E: estrus, M: metestrus, D: diestrus.

Figure S1. Inter-pulse intervals and waveforms of SEs^kiss in each estrus stage, related to Fig. 1. (**A**) Inter-pulse intervals of SEs^kiss in each estrus stage. Different letters (a–c) in the upper part of the graph denote significant differences at $P < 0.05$ by one-way repeated measures ANOVA followed by the Tukey–Kramer post hoc test. (**B**) SE^kiss waveforms in the light and

dark periods of each estrus stage. Data are expressed as mean (black line) ± SEM (colored shadow).

Figure 2. Decrease in estrus cycle frequency during the transition to reproductive senescence. (**A**) Number of estrus cycles per 25 days at 3–4 ($n = 5$), 8–9 ($n = 8$), 11–12 ($n = 6$), and 14–15 ($n = 12$) months in old wild-type C57BL/6 mice. Different letters (a, b) in the upper part of the graph denote significant differences at $P < 0.05$ by one-way ANOVA followed by the Tukey–Kramer post hoc test. (**B**) Chronic monitoring of estrus cycle frequency from the individuals used for Ca^{2+} imaging of SEs^{kiss} (C57BL/6 *Kiss-Cre* mice with AAV-mediated GCaMP6s expression). Black: average number of estrus cycles, gray: individual data ($n = 12$ animals in total).

Figure 3. Chronic monitoring of SEs^kiss from the reproductive to acyclic phase. (**A**) Representative photometry raw data of SEs^kiss in the diestrus stage from the reproductive to acyclic phase (*n* = 2 animals). Red arrowheads indicate SEs^kiss. (**B**) Numbers of SEs^kiss in the estrus stages (color bars) for 7 days from the reproductive to acyclic phase. SEs^kiss were recorded for 6 h in both light and dark periods. (**C**) Number of SEs^kiss in individual estrus stages (shown in different colors) of individual female mice. ns: not significant by one-way ANOVA. (**D**) Cumulative probability of inter-pulse interval. ns: not significant by the Kolmogorov–Smirnov test. (**E**) Peak height as assessed by ΔF/F of SEs^kiss from the reproductive to acyclic phase. Different letters (a–d, w–z) in the lower part of the graph denote significant differences at $P < 0.05$ by the Kruskal–Wallis test followed by the Mann–Whitney *U* test. Re; reproductive, RC; regular cyclic, IrC; irregular cyclic, Ac; acyclic.

Figure 4. Characterization of SEs^kiss in aging female mice. (**A**) Representative numbers of SEs^kiss in the estrus stages (color bars) for 7 days from regular cyclic, irregular cyclic, and acyclic female mice. SEs^kiss were recorded for 6 h in both light and dark periods. (**B**) Ratio of SEs^kiss in the dark period normalized to those in the preceding light period of regular cyclic ($n = 4$), irregular cyclic ($n = 3$), and acyclic ($n = 3$) female mice, respectively. Dotted line = 0.5. (**C**) Average numbers of SEs^kiss in 6 h in the light (open boxes in the LD cycle) and dark (closed boxes in the LD cycle) periods of reproductive (Re), regular cyclic (RC), irregular cyclic (IrC), and acyclic (Ac) mice. Red horizontal lines indicate the mean. No interaction effect between estrus cyclicity and LD cycle was found by two-way ANOVA. *$P < 0.05$ by paired t-test. Of note, the Re group (age 4–6 months) represents a reanalysis of the 5 days of data reported in Fig. 1 (corresponding to one estrus cycle). (**D**) Cumulative probability of inter-peak intervals among the Re (black), RC (green), IrC (blue), and Ac (orange) groups. ns: not significant by the Kolmogorov–Smirnov test. (**E**) Intensities of SEs^kiss during reproductive aging as assessed by the fold change of ΔF/F values in the IrC (#ID8, 9) and Ac (ID#7, 10) phases normalized to those in the RC phase. *$P < 0.05$ by the Mann–Whitney U test. Ns: not significant.

Figure S2. Waveforms and intensities of SEs^{kiss}, related to Figs. 3 and 4. (**A**) Averaged traces of SE^{kiss} waveforms of the same female mice in the reproductive (Re), regular cyclic (RC), irregular cyclic (IrC), and acyclic (Ac) phases. (**B**) Quantifications of detailed parameters of SE^{kiss} waveforms in individual female mice: LWQM, FWQM, and RWQM as defined in Fig. 1. *$P < 0.05$ by the Kruskal–Wallis test followed by the Mann–Whitney U test. In ID#5, the shape of the waveform was changed in the IrC compared with the Re phase, and the shape reverted in the Ac phase. In ID#6, significant differences in the length of LWQM and RWQM were observed between the Re and Ac phases. The origin of these individual differences is unknown, but the bias of estrus cycles in aging female mice might underlay the variations. (**C**) Coefficient of variances of the intensities of SEs^{kiss} in a 6-h window from Re, RC, IrC, and Ac individual mice. *$P < 0.05$ by the Kruskal–Wallis test followed by the Mann–Whitney U test.

Acknowledgments

We wish to thank RIKEN BDR LARGE staff for animal care, and Dr. Uenoyama (Nagoya University), Dr. Murata (the University of Tokyo), and the members of the Miyamichi Laboratory for the critical reading of the manuscript. This study was supported by Grants-in-Aid to TG (Nos. 19K16270 and 21K15194) and KM (Nos. 20K20589 and 21H02587) from the Japan Society for the Promotion of Science (JSPS). TG is also supported by the RIKEN Special Postdoctoral Researchers Program.

References

Belchetz, P.E., Plant, T.M., Nakai, Y., Keogh, E.J., and Knobil, E. (1978). Hypophysial responses to continuous and intermittent delivery of hypopthalamic gonadotropin-releasing hormone. Science *202*, 631-633.

Byers, S.L., Wiles, M.V., Dunn, S.L., and Taft, R.A. (2012). Mouse estrous cycle identification tool and images. PLoS One *7*, e35538.

Caligioni, C.S. (2009). Assessing reproductive status/stages in mice. Curr Protoc Neurosci *Appendix 4*, Appendix 4I.

Casper, R.F., Yen, S.S., and Wilkes, M.M. (1979). Menopausal flushes: a neuroendocrine link with pulsatile luteninizing hormone secreation. Science *205*, 823-825.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V*., et al.* (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*, 295-300.

Clarkson, J., d'Anglemont de Tassigny, X., Moreno, A.S., Colledge, W.H., and Herbison, A.E. (2008). Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. J Neurosci *28*, 8691-8697.

Clarkson, J., Han, S.Y., Piet, R., McLennan, T., Kane, G.M., Ng, J., Porteous, R.W., Kim, J.S., Colledge, W.H., Iremonger, K.J*., et al.* (2017). Definition of the hypothalamic GnRH pulse generator in mice. Proc Natl Acad Sci U S A *114*, E10216-E10223.

Czieselsky, K., Prescott, M., Porteous, R., Campos, P., Clarkson, J., Steyn, F.J., Campbell, R.E., and Herbison, A.E. (2016). Pulse and Surge Profiles of Luteinizing Hormone Secretion in the Mouse. Endocrinology *157*, 4794-4802.

d'Anglemont de Tassigny, X., Fagg, L.A., Dixon, J.P., Day, K., Leitch, H.G., Hendrick, A.G., Zahn, D., Franceschini, I., Caraty, A., Carlton, M.B*., et al.* (2007). Hypogonadotropic hypogonadism in mice lacking a functional Kiss1 gene. Proc Natl Acad Sci U S A *104*, 10714-10719.

Franklin, K.B.J., and Paxinos, G. (2013). Paxinos and Franklin's The mouse brain in stereotaxic coordinates, Fourth edition. edn (Amsterdam: Academic Press, an imprint of Elsevier).

Gee, D.M., Flurkey, K., and Finch, C.E. (1983). Aging and the regulation of luteinizing hormone in C57BL/6J mice: impaired elevations after ovariectomy and spontaneous elevations at advanced ages. Biol Reprod *28*, 598-607.

Gore, A.C. (2015). Aging and Reproduction. Physiology of Reproduction *Chapter 37*, 1661-1693.

Gottsch, M.L., Popa, S.M., Lawhorn, J.K., Qiu, J., Tonsfeldt, K.J., Bosch, M.A., Kelly, M.J., Ronnekleiv, O.K., Sanz, E., McKnight, G.S*., et al.* (2011). Molecular properties of Kiss1 neurons in the arcuate nucleus of the mouse. Endocrinology *152*, 4298-4309.

Greendale, G.A., Lee, N.P., and Arriola, E.R. (1999). The menopause. Lancet *353*, 571-580.

Han, S.Y., Kane, G., Cheong, I., and Herbison, A.E. (2019). Characterization of GnRH Pulse Generator Activity in Male Mice Using GCaMP Fiber Photometry. Endocrinology *160*, 557-567.

Herbison, A.E. (2018). The Gonadotropin-Releasing Hormone Pulse Generator. Endocrinology *159*, 3723-3736.

Kermath, B.A., and Gore, A.C. (2012). Neuroendocrine control of the transition to reproductive senescence: lessons learned from the female rodent model. Neuroendocrinology *96*, 1-12.

Krohn, P.L. (1962). Review lectures on senescence. II. Heterochronic transplantation in the study of ageing. Proc R Soc Lond B Biol Sci *157*, 128-147.

Kunimura, Y., Iwata, K., Ishigami, A., and Ozawa, H. (2017). Age-related alterations in hypothalamic kisspeptin, neurokinin B, and dynorphin neurons and in pulsatile LH release in female and male rats. Neurobiol Aging *50*, 30-38.

Lapatto, R., Pallais, J.C., Zhang, D., Chan, Y.M., Mahan, A., Cerrato, F., Le, W.W., Hoffman, G.E., and Seminara, S.B. (2007). Kiss1-/- mice exhibit more variable hypogonadism than Gpr54-/- mice. Endocrinology *148*, 4927-4936.

Lederman, M.A., Lebesgue, D., Gonzalez, V.V., Shu, J., Merhi, Z.O., Etgen, A.M., and Neal-Perry, G. (2010). Age-related LH surge dysfunction correlates with reduced responsiveness of hypothalamic anteroventral periventricular nucleus kisspeptin neurons to estradiol positive feedback in middle-aged rats. Neuropharmacology *58*, 314-320.

LeFevre, J., and McClintock, M.K. (1988). Reproductive senescence in female rats: a longitudinal study of individual differences in estrous cycles and behavior. Biol Reprod *38*, 780-789.

Matsuda, F., Ohkura, S., Magata, F., Munetomo, A., Chen, J., Sato, M., Inoue, N., Uenoyama, Y., and Tsukamura, H. (2019). Role of kisspeptin neurons as a GnRH surge generator: Comparative aspects in rodents and non-rodent mammals. J Obstet Gynaecol Res *45*, 2318-2329.

Matt, D.W., Gilson, M.P., Sales, T.E., Krieg, R.J., Kerbeshian, M.C., Veldhuis, J.D., and Evans, W.S. (1998). Characterization of attenuated proestrous luteinizing hormone surges in middle-aged rats by deconvolution analysis. Biol Reprod *59*, 1477-1482.

McGee, E.A., and Hsueh, A.J. (2000). Initial and cyclic recruitment of ovarian follicles. Endocr Rev *21*, 200-214.

McQuillan, H.J., Han, S.Y., Cheong, I., and Herbison, A.E. (2019). GnRH Pulse Generator Activity Across the Estrous Cycle of Female Mice. Endocrinology *160*, 1480-1491.

Mobbs, C.V., Cheyney, D., Sinha, Y.N., and Finch, C.E. (1985). Age-correlated and ovary-dependent changes in relationships between plasma estradiol and luteinizing hormone, prolactin, and growth hormone in female C57BL/6J mice. Endocrinology *116*, 813-820.

Moore, A.M., Coolen, L.M., and Lehman, M.N. (2022). In vivo imaging of the GnRH pulse generator reveals a temporal order of neuronal activation and synchronization during each pulse. Proc Natl Acad Sci U S A *119*.

Nagae, M., Uenoyama, Y., Okamoto, S., Tsuchida, H., Ikegami, K., Goto, T., Majarune, S., Nakamura, S., Sanbo, M., Hirabayashi, M.*, et al.* (2021). Direct evidence that KNDy neurons maintain gonadotropin pulses and folliculogenesis as the GnRH pulse generator. Proc Natl Acad Sci U S A *118*.

Neal-Perry, G., Lebesgue, D., Lederman, M., Shu, J., Zeevalk, G.D., and Etgen, A.M. (2009). The excitatory peptide kisspeptin restores the luteinizing hormone surge and modulates amino acid neurotransmission in the medial preoptic area of middle-aged rats. Endocrinology *150*, 3699-3708.

Nelson, J.F., Felicio, L.S., Osterburg, H.H., and Finch, C.E. (1981). Altered profiles of estradiol and progesterone associated with prolonged estrous cycles and persistent vaginal cornification in aging C57BL/6J mice. Biol Reprod *24*, 784-794.

Nelson, J.F., Felicio, L.S., Randall, P.K., Sims, C., and Finch, C.E. (1982). A longitudinal study of estrous cyclicity in aging C57BL/6J mice: I. Cycle frequency, length and vaginal cytology. Biol Reprod *27*, 327-339.

Nilsson, M.E., Vandenput, L., Tivesten, A., Norlen, A.K., Lagerquist, M.K., Windahl, S.H., Borjesson, A.E., Farman, H.H., Poutanen, M., Benrick, A.*, et al.* (2015). Measurement of a

Comprehensive Sex Steroid Profile in Rodent Serum by High-Sensitive Gas Chromatography-Tandem Mass Spectrometry. Endocrinology *156*, 2492-2502.

Padilla, S.L., Johnson, C.W., Barker, F.D., Patterson, M.A., and Palmiter, R.D. (2018). A Neural Circuit Underlying the Generation of Hot Flushes. Cell Rep *24*, 271-277.

Piet, R., Kalil, B., McLennan, T., Porteous, R., Czieselsky, K., and Herbison, A.E. (2018). Dominant Neuropeptide Cotransmission in Kisspeptin-GABA Regulation of GnRH Neuron Firing Driving Ovulation. J Neurosci *38*, 6310-6322.

Rance, N.E., Dacks, P.A., Mittelman-Smith, M.A., Romanovsky, A.A., and Krajewski-Hall, S.J. (2013). Modulation of body temperature and LH secretion by hypothalamic KNDy (kisspeptin, neurokinin B and dynorphin) neurons: a novel hypothesis on the mechanism of hot flushes. Front Neuroendocrinol *34*, 211-227.

Rubin, B.S., Lee, C.E., and King, J.C. (1994). A reduced proportion of luteinizing hormone (LH)-releasing hormone neurons express Fos protein during the preovulatory or steroid-induced LH surge in middle-aged rats. Biol Reprod *51*, 1264-1272.

Scarbrough, K., and Wise, P.M. (1990). Age-related changes in pulsatile luteinizing hormone release precede the transition to estrous acyclicity and depend upon estrous cycle history. Endocrinology *126*, 884-890.

Steinmetz, N.A., Aydin, C., Lebedeva, A., Okun, M., Pachitariu, M., Bauza, M., Beau, M., Bhagat, J., Bohm, C., Broux, M.*, et al.* (2021). Neuropixels 2.0: A miniaturized high-density probe for stable, long-term brain recordings. Science *372*.

Steyn, F.J., Wan, Y., Clarkson, J., Veldhuis, J.D., Herbison, A.E., and Chen, C. (2013). Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice. Endocrinology *154*, 4939-4945.

Uenoyama, Y., Nakamura, S., Hayakawa, Y., Ikegami, K., Watanabe, Y., Deura, C., Minabe, S., Tomikawa, J., Goto, T., Ieda, N.*, et al.* (2015). Lack of pulse and surge modes and glutamatergic stimulation of luteinising hormone release in Kiss1 knockout rats. J Neuroendocrinol *27*, 187-197.

Wise, P.M., Dueker, E., and Wuttke, W. (1988). Age-related alterations in pulsatile luteinizing hormone release: effects of long-term ovariectomy, repeated pregnancies and naloxone. Biol Reprod *39*, 1060-1066.