Revealing the drivers of antibiotic resistance trends in *Streptococcus pneumoniae* amidst the 2020 COVID-19 pandemic: Insights from mathematical modeling

Aleksandra Kovacevic[1,2] David R M Smith[1,2,3,4], Eve Rahbé[1,2], Sophie Novelli[2], Paul Henriot[3,5], Emmanuelle Varon[6], Robert Cohen[7,8,9,10,11], Corinne Levy[7,8,10,11], Laura Temime[3,5], Lulla Opatowski[1,2]

1. Institut Pasteur, Université Paris Cité, Epidemiology and Modelling of Antibiotic Evasion (EMAE) unit, Paris, France
2. Université Paris-Saclay, Université de Versailles Saint-Quentin-en-Yvelines, Inserm U1018, CESP, Anti-infective evasion and pharmacoepidemiology team, Montigny-Le-Bretonneux, France
3. Modélisation, épidémiologie et surveillance des risques sanitaires (MESuRS), Conservatoire national des arts et métiers, Paris, France
4. Health Economics Research Centre, Nuffield Department of Health, University of Oxford, Oxford, United Kingdom
5. PACRI unit, Institut Pasteur, Conservatoire national des arts et métiers, Paris, France
6. Centre National de Référence des Pneumocoques, Centre Hospitalier Intercommunal de Créteil, Créteil, France
7. Institut Mondor de Recherche Biomédicale-Groupe de Recherche Clinique Groupe d'Etude des Maladies Infectieuses Néonatales et Infantiles (IMRB-GRC GEMINI), Université Paris Est, 94000 Créteil, France
8. Groupe de Pathologie Infectieuse Pédiatrique (GPIP), 06200 Nice, France
9. Unité Court Séjour, Petits Nourrissons, Service de Néonatologie, Centre Hospitalier Intercommunal de Créteil, France
10. Association Clinique et Thérapeutique Infantile du Val-de-Marne (ACTIV), 94000 Créteil, France
11. Association Française de Pédiatrie Ambulatoire (AFPA), 45000 Orléans, France

Keywords: *Streptococcus pneumoniae*; antibiotic resistance; AMR; invasive pneumococcal disease (IPD); COVID-19 pandemic; virus-bacteria interactions; pathogen interactions; pneumococcal carriage; co-infection model

Abstract

Non-pharmaceutical interventions implemented to block SARS-CoV-2 transmission in early 2020 led to global reductions in the incidence of invasive pneumococcal disease (IPD). By contrast, most European countries reported an increase in antibiotic resistance among invasive *Streptococcus pneumoniae* isolates from 2019 to 2020, while an increasing number of studies reported stable pneumococcal carriage prevalence over the same period. To disentangle the impacts of the COVID-19 pandemic on pneumococcal epidemiology in the community setting, we propose a mathematical model formalizing simultaneous transmission of SARS-CoV-2 and antibiotic-sensitive and -resistant strains of *S. pneumoniae*. To test hypotheses underlying these trends five mechanisms were built in into the model and examined: (1) a population-wide reduction of antibiotic prescriptions in the community, (2) lockdown effect on pneumococcal transmission, (3) a reduced risk of developing an IPD due to the absence of common respiratory viruses, (4) community azithromycin use in COVID-19 infected individuals, (5) and a longer carriage duration of antibiotic-resistant pneumococcal strains. Among 31 possible pandemic scenarios involving mechanisms individually or in combination, model simulations surprisingly identified only two scenarios that reproduced the reported trends in the general population. They included factors (1), (3), and (4). These scenarios replicated a nearly 50% reduction in annual IPD, and an increase in antibiotic resistance from 20% to 22%, all while maintaining a relatively stable pneumococcal carriage. Exploring further, higher SARS-CoV-2 R_0 values and synergistic within-host virus-bacteria interaction mechanisms could have additionally contributed to the observed antibiotic resistance increase. Our work demonstrates the utility of the mathematical modeling approach in unraveling the complex effects of the COVID-19 pandemic responses on AMR dynamics.

Introduction

In the early 2020, international responses to the coronavirus disease 2019 (COVID-19) pandemic led to unprecedented worldwide change in population mixing, healthcare-seeking behavior, and infection prevention and control practices. This modified the ecology and epidemiology of many infectious diseases at a global scale. Strong impacts of COVID-19 on infectious disease dynamics have been reported for common viral and bacterial respiratory infections, sexually transmitted pathogens like HIV, vector-borne diseases like dengue, and even non-communicable diseases (Braunstein et al., 2020; Brueggemann et al., 2021; Chen et al., 2022; Palmer et al., 2020). Antimicrobial resistance (AMR), however, remains one of the leading threats to global health. In 2019, estimates showed that AMR in clinically relevant bacteria was associated with 4.95 million deaths, of which 1.27 million were directly attributable to resistance (Murray et al., 2022). Impacts of the COVID-19 pandemic on AMR dynamics remain relatively poorly understood.

A joint report from the World Health Organization (WHO) and European Centre for Disease Prevention and Control (ECDC) has reported 2020 AMR trends across 29 European countries for eight antibiotic-resistant bacterial pathogens of concern, including *S. pneumoniae* (European Centre for Disease Prevention and Control and World Health Organization, 2022). While the situation varies widely across bacterial species, antimicrobial groups, and regions, most European countries, including France, documented an increase in pneumococcal resistance to both penicillin and macrolides between 2019 and 2020. The resistance rates rose from 12.2% in 2019 to 15.6% in 2020 for penicillin and from 14.5% in 2019 to 16.9% in 2020 for macrolides, as reported in the EU/EEA (European Centre for Disease Prevention and Control and World Health Organization, 2022). However, increased pneumococcal resistance was accompanied by a sharp worldwide decline in invasive pneumococcal disease (IPD) incidence (Brueggemann et al., 2021; Shaw et al., 2023).

Similar declines in bacterial disease during early waves of COVID-19 have been observed in the context of sentinel community-acquired infections in New Zealand (Duffy et al., 2021), IPDs in Taiwan (Chien et al., 2021) and Hong Kong (Teng et al., 2022), and lower respiratory tract infections in China (Chen et al., 2021). Yet, surprisingly, a growing number of studies have reported mostly stable pneumococcal carriage throughout the COVID-19 pandemic containment, including among infants in Belgium (Willen et al., 2022), children in Vietnam (Nation et al., 2023), Serbia (Petrović et al., 2022), France (Rybak et al., 2022), South Africa (Olwagen et al., 2023), and Israel (Dagan et al., 2023), adults in Connecticut (Wyllie et al., 2023), and households in Seattle (Bennett et al., 2023). In contrast, a study conducted in Denmark reported a decrease in pneumococcal carriage among older adults during the COVID-19 lockdown (Tinggaard et al., 2023).

Understanding the cause of these trends is not straightforward, as many responses to the COVID-19 pandemic, such as the implementation of non-pharmaceutical interventions (NPIs), changes in healthcare-seeking behavior, and alterations in antibiotic prescribing, occurred over the period (Knight et al., 2021). To gain a comprehensive understanding of the changes in AMR epidemiology during the COVID-19 pandemic, it is essential to simultaneously consider a range of scales and indicators. These include the rates of incidence of invasive bacterial diseases, the

138 proportion of antibiotic-resistant isolates among total invasive bacterial isolates, and the
139 prevalence of asymptomatic bacterial carriage in healthy individuals.
140
Several mechanisms may underlie the explanation of these contrasting observations. First,
NPIs implemented to block SARS-CoV-2 transmission, such as lockdowns and mask mandates,
may have led to reduced bacterial transmission. Containment measures also massively reduced
circulation of common respiratory viruses, which are known to be associated with invasive
bacterial disease (Domenech De Cellès et al., 2019; Smith and Opatowski, 2021). Second, the
lockdown was associated with reductions in primary care consultations (Homeniuk and Collins,
2021; Read et al., 2023; Zhang et al., 2021) leading to a global decrease of antibiotic prescriptions
(Högberg et al., 2021). In contrast, frequent antibiotic prescribing to COVID-19 outpatients may
have exacerbated AMR (Clancy et al., 2020; Knight et al., 2021). Differences in the duration of
pneumococcal carriage may have also played a role (Lehtinen et al., 2017). Finally, potential
within-host interactions between SARS-CoV-2 and *S. pneumoniae* could also have an impact on
infection risk (Amin-Chowdhury et al., 2021), although strong evidence for such interactions
remains limited (Wong et al., 2023).
154
Mathematical models incorporating the co-transmission of multiple pathogens within the
same host population provide a framework for investigating different hypotheses that underlie the
observed patterns in antibiotic resistance and incidence of IPD in *S. pneumoniae* and help to
enhance our understanding of the mechanisms involved. Co-circulation models have been used
previously to disentangle the public health consequences of interactions between pathogens such
as influenza and *S. pneumoniae* (Arduin et al., 2017; Domenech De Cellès et al., 2019; Shrestha
et al., 2013) and could similarly be used to understand impacts of the COVID-19 pandemic on
pathogens coinciding with SARS-CoV-2. However, in a systematic PubMed search conducted on
4 December 2023, we identified no epidemiological models describing the simultaneous
transmission of SARS-CoV-2 and antibiotic-resistant bacteria specific to the community setting
(Appendix 1).
166
Here, to disentangle how the COVID-19 pandemic has impacted the epidemiological
dynamics of antibiotic resistance in *S. pneumoniae*, we propose a mathematical model that
formalizes the transmission of SARS-CoV-2 and both antibiotic-sensitive and -resistant strains of
S. pneumoniae in the community setting, and which includes mechanistic impacts of COVID-19
burden on epidemiological parameters. Through simulation, we assess all possible combinations
of these mechanisms to evaluate their overall impact on IPD incidence, antibiotic resistance, and
the prevalence of pneumococcal carriage. Furthermore, we assess the changes in the incidence of
antibiotic-resistant IPD as we vary the basic reproduction number (R_0) of SARS-CoV-2 during the
first COVID-19 outbreak in Europe. We also consider assumed within-host pathogen interactions
between SARS-CoV-2 and *S. pneumoniae*.
177

Results

179
Antibiotic resistance trends and incidence of invasive pneumococcal disease in 2020
181 In routine surveillance data reported to the European Antimicrobial Resistance Surveillance
182 Network (EARS-Net), most European countries reported an increase in antibiotic resistance in *S.
183 pneumoniae* from 2019 to 2020, as indicated by increases in the proportion of invasive isolates

with phenotypic resistance to both penicillin and macrolides (Figure 1A). On the contrary, the total number of reported isolates in the EU/EEA decreased by 44.3% from 2019 to 2020 (European Centre for Disease Prevention and Control and World Health Organization, 2022) suggesting a decrease in incidence of invasive pneumococcal disease (Appendix 2 – Table 1).

Invasive pneumococcal isolate data for France provided by the French National Reference Center for Pneumococci (CNRP) revealed similar trends. In France, the total number of reported invasive pneumococcal isolates decreased by 45.1% from 2019 to 2020 (from 1119 to 614), while antibiotic resistance in *S. pneumoniae* isolates to penicillin and macrolides showed an increasing trend from 26.2% in 2019 to 35.5% in 2020 for penicillin, and from 20.9% in 2019 to 23.0% in 2020 for macrolides (Figure 1B). General decreasing trend in antibiotic resistance from 2017 to 2019 in *S. pneumoniae* was interrupted in 2020 (Figure 1 – figure supplement 1). These variations in antibiotic resistance manifested differently across age, with some age groups showing an increase in antibiotic resistance in 2020 compared to 2019, while others showed no significant change (Figure 1B).

Coinfection model of SARS-CoV-2 and *Streptococcus pneumoniae*
As mentioned above, several mechanisms may underlie the explanation of these contrasting observations (Figure 2A). COVID-19 NPIs may have led to reduced person-to-person bacterial transmission, potentially contributing to reduced rates of IPD incidence. These containment measures also massively reduced circulation of common respiratory viruses and the incidence of influenza-like-illnesses (ILIs). Respiratory viruses are known triggers and risk factors for developing an invasive bacterial disease from otherwise asymptomatic carriage; in that context, their reduction may have led to reduced infection risk (Domenech De Cellès et al., 2019; Smith and Opatowski, 2021). Due to reductions in primary care consultations in 2020, 26 European countries reported an estimated average decrease of 18.3% in overall antibiotic consumption, aligning with the global trend of reduced antibiotic prescriptions compared to 2019 (Högberg et al., 2021). On the other hand, frequent prescribing of azithromycin, a macrolide antibiotic initially hypothesized to be effective in COVID-19 treatment, has raised concerns for pandemic-associated antimicrobial overuse or misuse and may have exacerbated AMR during and following the first wave of the pandemic (Clancy et al., 2020; Knight et al., 2021; Kournoutou and Dinos, 2022; Langford et al., 2021; PRINCIPLE Trial Collaborative Group, 2021; Rusic et al., 2021). There are still uncertainties about pneumococcal ecology and the evolutionary processes that enable the robust coexistence of strains sensitive and resistant to antibiotics. The role of carriage duration, along with the impact of antibiotic consumption, is also not fully understood in this context. Longer carriage duration of antibiotic-resistant pneumococcal strains is a proposed explanation for this coexistence (Lehtinen et al., 2017). If so, antibiotic-resistant pneumococcal strains may have had an advantage during the lockdown period due to smaller clearance rates, ultimately leading to an increase in antibiotic resistance. Finally, among individuals with COVID-19, potential within-host interactions between SARS-CoV-2 and *S. pneumoniae* could also have had an impact on bacterial colonization and infection dynamics (Amin-Chowdhury et al., 2021).

To test mechanistic impacts of responses to the COVID-19 pandemic on pneumococcal epidemiology, we developed a compartmental, deterministic transmission model describing infection with SARS-CoV-2 being introduced on 1 Jan 2020 (Figure 2B) after colonization with *S. pneumoniae* reached an equilibrium in a large, well-mixed human population (Figure 2C). Two

lockdowns were implemented in the model in agreement with the two lockdowns implemented in
France in 2020. The model was parameterized to *S. pneumoniae* and five mechanisms were built
in into the model: (1) a population-wide reduction of antibiotic prescriptions in the community by
18% due to the reduced healthcare-seeking behavior, (2) lockdown reducing pneumococcal
transmission by 25%, (3) a reduced risk of developing an IPD from asymptomatic carriage due to
the absence of common respiratory viruses during the first lockdown (reduced by a factor IPD_{risk}
= 0.2), which continues after the first lockdown, albeit at a diminished level (IPD_{risk} = 0.4), (4)
community azithromycin use in 10% of COVID-19 infected individuals, (5) and a longer carriage
duration of antibiotic-resistant pneumococcal strains giving them a fitness advantage over
antibiotic-resistant strains (40 vs. 30 days).
240
Exploring the mechanisms and identifying the optimal scenario for explaining reported
trends
We conducted assessments on five distinct hypotheses, each characterized by a precise underlying
mechanism, and explored these hypotheses in combination within 31 pandemic scenarios, along
with two pre-pandemic (baseline) scenarios, which assume no SARS-CoV-2 circulation in the
population and allow for the same 30-day carriage duration (pre-pandemic 1) of both antibiotic-
sensitive and -resistant strains ($d_S = d_R$) or a longer, 40-day carriage duration (pre-pandemic 2)
of -resistant strains ($d_S > d_R$) (Table 1).
249
We assessed how different combinations of mechanisms may impact: (i) a change in the
annual IPD incidence as compared to the pre-pandemic (baseline) period, (ii) antibiotic resistance
rate in IPDs, defined as the annual number of antibiotic-resistant IPD cases over the total number
of IPD cases, and (iii) daily prevalences of antibiotic-resistant and total pneumococcal carriage in
a simulated population of 100,000 individuals (see Appendix 2 – Table 2 for parameter values).
To identify scenarios most compatible with the reported trends, results from model simulations
were compared to reported data trends from France in 2020 and more broadly to general EU/EEA
reported trends that followed similar patterns. Surprisingly only two scenarios were compatible
with reported trends. Scenarios S19 and S29 univocally reproduced increased antibiotic resistance
in the general population (AR%) accompanied by a reduction in the annual IPD incidence by
almost 50% (IPD inc.) with generally stable pneumococcal carriage prevalence in healthy
individuals during lockdown (Sp.). In contrast, model simulations revealed that a reduction in the
community antibiotic consumption alone (-18%) could not explain the reported trends and
generally led to a reduction of antibiotic resistance (Table 1, S1). Assuming a longer duration of
antibiotic-resistant pneumococcal carriage alone did not explain either the rise in antibiotic
resistance (Table 1, S5). Hypothesizing that lockdown reduced the transmission of pneumococcal
carriage (by 25%) in addition to a reduced community antibiotic prescribing did not seem probable
since, in simulations, this yielded a major reduction in pneumococcal carriage during containment
measures in all scenarios where this mechanism was implemented. On the other hand, considering
an indirect impact of lockdown on pneumococcal carriage where we implemented a reduction
factor for the risk of developing and IPD from otherwise asymptomatic carriage due to the absence
of viral respiratory infections during (*IPD_{risk}*=0.2) and after lockdown (*IPD_{risk}*=0.4) reproduced
the reported reduction in the annual IPD incidence while maintaining a stable prevalence of
pneumococcal carriage during lockdown (Table 1, S3). By itself however, this scenario did not
allow to observe an increase in antibiotic resistance.
275

When we combined reduced antibiotic prescribing and a reduced risk of developing an IPD
with community azithromycin use in a proportion of COVID-19 infected individuals, which
remains in the body for an additional 15.5 days after the last dose, in a single scenario, this scenario
satisfied the observed trends in AMR (Table 1, S19). Similar outcome was observed in scenario
S29 when adding a longer carriage duration of antibiotic resistant strains on top of this, however,
in the absence of community azithromycin use in COVID-19 infected (Table 1, S20) trends of
increasing antibiotic resistance cannot be reproduced. Therefore, our best model scenario for
describing the observed trends combined: (1) a reduction in the overall community antibiotic
consumption; (2) the assumption that lockdown effectively reduced SARS-CoV-2 transmission
including transmission of other respiratory viruses, but not pneumococcal carriage transmission,
indirectly reducing the risk of developing an IPD; (3) either identical or longer carriage durations
of antibiotic-resistant strains compared to antibiotic-sensitive strains, and (4) the community
azithromycin use in a proportion of COVID-19 infected individuals.

Effect of age

Next, we used the pandemic scenario S19 that best explains the reported trends to test the model
using different parameter combinations to mimic different subpopulations (children and the
elderly) considering that SARS-CoV-2 infection risk, pneumococcal disease risk, disease severity,
bacterial carriage prevalence, and antibiotic prescribing are all highly heterogeneous across age
groups. Using scenario S19, we initialized the model with lower and higher baseline carriage
prevalence (10%, 20%, and 30%) (Cohen et al., 2023; Rose et al., 2021; Rybak et al., 2022;
Tinggaard et al., 2023; Wang et al., 2017), we varied durations of pneumococcal carriage (20, 30,
and 45 days), pneumococcal invasion rate, and considered reductions of antibiotic consumption at
various levels (-13%, -18%, and -39%) consistent with the French data along with a range of
community azithromycin use in COVID-19 infected (0-20%). For a full list of parameters see
Appendix 2 – Table 2. Simulations showed that annual IPD incidence decreased between 43% and
51% compared to the pre-pandemic (baseline) scenario for children, the elderly, and the general
population (Figure 3, grey bars). Although the overall antibiotic prescribing in the community was
reduced (between 13% and 39%), antibiotic resistance is expected to increase (from 20.1% up to
23.6% in the elderly and from 32.8% up to 36.0% in children) compared to the pre-pandemic
period in all age groups and in all scenarios where azithromycin was used in COVID-19 infected
individuals (Figure 3, red bars). Daily prevalence of total pneumococcal carriage remained
relatively stable, exhibiting higher levels of decrease with increased azithromycin use, while the
prevalence of antibiotic-resistant pneumococcal carriage is expected to increase since clearance of
antibiotic-susceptible strains due to azithromycin use shifts the competitive balance in favor of the
existing resistant strains (Figure 3, third panel).

General trends produced in model simulations using scenario S19 remained unchanged
across different age groups. The extent of the impact depended on the combined magnitude of a
decrease in the general antibiotic use in the community and a degree of azithromycin use in
COVID-19 infected individuals belonging to a particular age group or a subpopulation. In the
elderly (≥65 years-old) and the general population, antibiotic resistance is expected to increase
due to azithromycin use in COVID-19 infected. Black arrows indicate model outcomes that
approximate the reported trends in antibiotic resistance in France for different age groups including
general population (Figure 3). Only in instances when there was no azithromycin use in COVID-
19 infected individuals, we observed a decrease in antibiotic resistance relative to the pre-

pandemic period (e.g., children <5 years-old). When combining the largest decrease in overall
antibiotic use with no or minimal azithromycin use in COVID-19 infected individuals, we expect
to see the largest decrease or no change in antibiotic resistance relative to the pre-pandemic period.
325
Effect of SARS-CoV-2 basic reproduction number (R$_0$) and within-host pathogen
interactions on AMR
Considering that model simulations reproduced an absolute increase in antibiotic resistance
comparable to that of 2% reported for macrolides in France but did not reproduce the reported
larger increase in penicillin resistance, which was more than a 9% rise (35.5% relative increase)
in France, we explored additional factors that may have amplified this increase. Using model
scenario S19, we show that an association between higher values of SARS-CoV-2 R$_0$ and a greater
percentage of COVID-19 infected individuals taking azithromycin leads to increased cumulative
incidence of antibiotic-resistant IPDs and elevated antibiotic resistance (Figure 4A). For example,
if pre-lockdown R$_0$ of SARS-CoV-2 was 3.8 instead of 3.2, model simulations predict an increase
of 3.5% (23.5%) in antibiotic resistance from the pre-pandemic levels instead of 2%. As the R$_0$
value increases, the impact of azithromycin use becomes more pronounced.
Assuming within-host interactions where SARS-CoV-2 infection favors progression from
pneumococcal colonization to disease ($\psi_c > 1$), we found that surges in COVID-19 cases
accompanied by increasing levels of azithromycin use lead to excess number of cases caused by
antibiotic-resistant strains. Indeed, a rate of disease progression increased by a factor $\psi_c = 40$ in
in scenario S19 with 10% of infected using azithromycin applied to the general population results
in approximately 0.75 additional cases of antibiotic-resistant disease per 100,000 inhabitants over
the course of one year compared to 0.06 additional cases if there are no within-host interactions
(Figure 4B). This represents 5% rise in resistance from the pre-pandemic levels (25% relative
increase).
347

348 # Discussion

349
We propose a novel co-circulation model describing the spread of SARS-CoV-2 and antibiotic-
resistant bacteria in a community setting to show how human behavioral responses to the COVID-
19 pandemic can differentially impact antibiotic resistance. Our model simulations assessed
different hypotheses proposed to explain the observed trends of antibiotic resistance, IPD
incidence, and pneumococcal carriage. We identified the most plausible mechanisms underlying
the observed patterns of resistance and disease incidence, showing how lockdowns indirectly
substantially reduce the incidence of IPD, while surges in COVID-19 cases accompanied by
antibiotic prescribing in COVID-19 infected individuals increase antibiotic resistance.
358
Many studies have reported trends on the incidence of community-acquired bacterial
infections since the onset of the pandemic (Brueggemann et al., 2021; Shaw et al., 2023). There
was a significant reduction in the risk of invasive disease caused by *S. pneumoniae* (risk ratio 0·47;
95% CI 0·40–0·55) (Shaw et al., 2023). Initially, this observation seemed to support the hypothesis
that NPIs implemented to control SARS-CoV-2 transmission may have simultaneously reduced
the incidence of bacterial infections by preventing bacterial transmission and acquisition
(Brueggemann et al., 2021; Kadambari et al., 2022). Indeed, the scenario of lockdown impact on
pneumococcal transmission reproduced such trends. However, incorporating a mechanism of
reduced risk for developing an IPD due to the absence of circulation of common respiratory viruses

led to similar estimates of the relative reduction in IPD incidence as reported in the EU/EEA for 2020 (Brueggemann et al., 2021; European Centre for Disease Prevention and Control and World Health Organization, 2022). This finding, coupled with the outcome of other studies that found a generally stable pneumococcal carriage prevalence in healthy individuals, both children and adults, during COVID-19 containment measures (Nation et al., 2023; Petrović et al., 2022; Rybak et al., 2022; Willen et al., 2022; Wyllie et al., 2023), supports the alternative hypothesis. This explanation accounts for the decreased incidence of IPD, rather than attributing it to reduced pneumococcal transmission, which resulted in a significant reduction in carriage according to the simulations (Smith and Opatowski, 2021). Furthermore, a study in Vietnam found that reductions in IPD associated with NPIs may be due to reductions in overall pneumococcal carriage density rather than carriage prevalence, driven by reductions in capsular pneumococcal carriage density frequently implicated in IPD (Nation et al., 2023). Considering that common respiratory viruses such as influenza increase pneumococcal carriage density, which contributes to transmission and disease, this hypothesis seems plausible (Alpkvist et al., 2015; Diavatopoulos et al., 2010; McCullers et al., 2010; Short et al., 2012; Wolter et al., 2014).

Globally, community antibiotic consumption dropped during the first year of the COVID-19 pandemic compared to the pre-pandemic period. Decreasing temporal trends were observed in England (Hussain et al., 2021), Canada (Mamun et al., 2021), the United States (Buehrle et al., 2021), China (Zhang et al., 2021), South Korea (Ryu et al., 2021), New Zealand (Duffy et al., 2021), and across European countries (Högberg et al., 2021). In France in particular, the number of antibiotic prescriptions decreased by 18.2% in the general population; however, this reduction ranged from 13% to 39% for the oldest and youngest age groups, respectively (Bara et al., 2022). These trends in antibiotic prescribing may largely be explained by reduced incidence of seasonal respiratory tract infections and reduced primary care consultations (Andrews et al., 2022; Homeniuk and Collins, 2021). On the other hand, the advent of telemedicine, pandemic-induced patient stress, and increased antibiotic demand may have partly offset prescription reductions due to decreased consultations and healthcare-seeking behavior (Hsu, 2020; Read et al., 2023). In a global analysis of antimicrobial sales, Khouja et al. found that antibiotic consumption initially increased by approximately 7% in March 2020, prior to subsequent declines through to August 2020 (Khouja et al., 2022). While overall antibiotic prescribing may have decreased, prescription of specific antibiotics has increased, particularly those associated with COVID-19 patient management. Across continents, a rise of 10% in monthly COVID-19 cases exhibited a correlative trend with elevated macrolide sales of 0.8%, 1.3%, and 1.5% in Europe, North America, and Africa, respectively (Nandi et al., 2023).

Community consumption of azithromycin, a macrolide, increased during the first year of the pandemic in multiple countries with significant variation across geographic locations and with greatest prescribing among older patients (Bara et al., 2022; Bednarčuk et al., 2023; Bogdanić et al., 2022; Crisafulli et al., 2022; Parveen et al., 2020; Weill et al., 2021). In an outpatient setting in southern Italy between February 2020 and January 2021, azithromycin represented 42.1% of all drug prescriptions to individuals diagnosed with COVID-19, while all other antibiotics combined represented just 20.9% (Crisafulli et al., 2022). A study in northwest London across two epidemic waves between January and August 2020 found that, among COVID-19 patients prescribed an antibiotic by a general practitioner during the study period, 31.5% received their prescription within 14 days of a positive SARS-CoV-2 test (Zhu et al., 2021). Two large USA-based studies

have also described early pandemic antibiotic prescribing among COVID-19 patients. From April 2020 to April 2021, approximately 30% of outpatient COVID-19–related visits among Medicare beneficiaries (≥65 years-old) have resulted in a filled antibiotic prescription, 50.7% of which were for azithromycin (Tsay et al., 2022). For 0-to-5 year-olds and 45-to-64 year-olds, 4% and 16% of outpatient COVID-19–related visits have resulted in a filled antibiotic prescription, respectively (Wittman et al., 2023). In the Alsace region in France, there was a clear peak azithromycin prescribing during the first wave of the COVID-19 (Danion et al., 2023). During the first lockdown in France, community azithromycin consumption increased by 25.9%, with the increase varying from 13.4% to 47.3% depending on the week (Weill et al., 2021), while the overall number of azithromycin prescriptions across France in 2020 increased by 10.1% relative to 2019 (Bara et al., 2022). Azithromycin treatment usually lasts 3-5 days depending on the disease, but the drug stays in the system for about 15.5 days after the last dose due to the long half-life of more than 60 hours (Foulds et al., 1990; Girard et al., 2005). On the other hand, penicillin has an elimination half-life of approximately 1.4 hours and leaves the body in 7.7 hours after the last dose. This suggests that if azithromycin consumption increased during the first year of the pandemic, antibiotic exposure time also increased as a result, although the overall number of antibiotic prescriptions decreased. Moreover, the use of azithromycin has been associated with selection of both macrolide and non-macrolide resistance (Doan et al., 2020). In a study investigating the direct effect of antibiotic exposure on resistance in the oral streptococcal flora of healthy volunteers, use of azithromycin (500 mg once daily for 3 days) significantly increased the proportion of macrolide-resistant streptococci in healthy individuals (Malhotra-Kumar et al., 2007). Resistance peaked at day four in the azithromycin group and this increase remained significantly higher in the azithromycin group than in the placebo group until day 180 (Malhotra-Kumar et al., 2007). A clinical trial of mass azithromycin distributions for treating trachoma in Ethiopia resulted in an increase in resistant *S. pneumoniae* isolates among children under the age of 10 (Keenan et al., 2018, 2015).

Our model simulations show that antibiotic resistance increases with surges in SARS-CoV-2 infections when there is a corresponding increase in azithromycin use, but that lockdowns can moderate this increasing trend by effectively limiting transmission of SARS-CoV-2 (Salje et al., 2020). Conversely, surges in azithromycin prescribing during SARS-CoV-2 outbreaks in the absence of effective measures to prevent transmission, as reported in certain regions and pandemic periods, may cause substantial increases in antibiotic resistance. Our model successfully captured the main trends of antibiotic resistance and IPD incidence observed in Europe in 2020 for *S. pneumoniae*. However, not all European countries reported an increase in antibiotic resistance. This inter-country heterogeneity may not be due only to heterogeneity of antibiotic use as shown in our model but may be attributed to other pandemic factors not directly implemented or assumed in the model scenario, such as different adherence to COVID-19 control measures across countries and different age groups, including impacts on disease surveillance and data reporting during the pandemic. Real-life scenarios are significantly more complicated and involve multiple alterations of many pandemic factors at different points in time and heterogeneity across populations (e.g., antibiotic prescribing increases in some demographic groups and decreases in others, multiple lockdowns, curfews, or telework).

In our model simulations, we used SARS-CoV-2 parameter value $R_0 = 3.2$ (Liu et al., 2020) in the absence of population immunity, best reflecting epidemiological dynamics from early in the pandemic. The most common estimates of SARS-CoV-2 R_0 in France and other European

countries ranged from $R_0 = 2$ to 4 (Flaxman et al., 2020; Liu et al., 2020). Modeling results suggest
that higher SARS-CoV-2 R_0 estimates combined with higher proportion of COVID-19 infected
individuals using azithromycin exacerbated impacts of COVID-19 on antibiotic resistance (Figure
4A). However, the overall impacts of COVID-19 on AMR are difficult to predict, likely vary over
the short, medium, and long term, and depend on the organism, setting, and subpopulation
considered.
466
SARS-CoV-2 bacterial coinfection has been reported relatively rarely over the course of
the pandemic, suggesting that most COVID-19 patients probably do not require antibiotic therapy
(Garcia-Vidal et al., 2021; Karami et al., 2021; Langford et al., 2020), although extensive
prophylactic antibiotic use may have limited observed co-infection incidence. The inflammatory
immune response resulting from COVID-19 likely predisposes patients to subsequent progression
to an invasive bacterial disease (IBD) to some extent (Sender et al., 2021), but antibiotic use may
also favor progression to IBD for patients colonized with drug-resistant strains (Baggs et al., 2018).
We do not explicitly model the dynamics of interaction since strong evidence for such interactions
remains limited (Wong et al., 2023). The results presented in Figure 4B suggest that such within-
host interactions could have important consequences for the resistant IPD incidence during
COVID-19 waves, especially in the elderly and high-risk groups. The model's structure allows for
easy integration of mechanistic interactions as more information becomes available on this
phenomenon.
480
Our study focused on the general community, but COVID-19 distinctly influenced AMR
in hospitals and long-term care facilities. Extensive antibiotic use in COVID-19 patients and
disruptions to antibiotic stewardship programs may have increased antibiotic-resistant carriage in
these settings. A meta-analysis conducted on studies published until June 2020 found that 68-81%
of hospitalized COVID-19 patients and 74-94% in intensive care received antibiotics (Monnet and
Harbarth, 2020). The disorganization in hospitals during the COVID-19 pandemic might have
reduced antibiotic resistance surveillance, allowing resistant organisms to spread. However, the
early implementation of antibiotic stewardship programs in March 2020, patient isolation, and
widespread use of personal protective equipment (PPE) have mitigated this increase to some
degree (Henig et al., 2021; Monnet and Harbarth, 2020; Seaton et al., 2020; Van Laethem et al.,
2021). Models analyzing these impacts in hospitals contribute to understanding COVID-19's
specific role in the antibiotic resistance burden in different settings (Smith et al., 2023).
493
A limitation of our model is the lack of age structure and contact patterns between age
groups, as SARS-CoV-2 infection risk, pneumococcal disease risk, disease severity, bacterial
carriage prevalence and antibiotic prescribing are all highly heterogeneous across age groups.
While this choice was made to keep the model as simple as possible, we tested the model using
different parameter combinations to mimic different subpopulations (children and ≥65 years-old).
This included varying durations of pneumococcal carriage, initializing the model with lower and
higher baseline carriage prevalence, considering reductions of general antibiotic consumption at
various levels, and varying a percentage of COVID-19 infected individuals using azithromycin.
Simulations of the different age groups individually interestingly reproduced realistic trends by
age.
504

In conclusion, we introduce the first epidemiological model outlining the impact of the
COVID-19 pandemic on the dynamics of AMR in the community. Our work demonstrates the
utility of mathematical modeling approach in unraveling the complex effects of the COVID-19
pandemic responses AMR dynamics. While our model was structured and parameterized based
upon *S. pneumoniae*, its adaptability allows for application to various bacteria and epidemiological
scenarios in the community (e.g., impacts of SARS-CoV-2-bacteria interactions in the context of
seasonal outbreaks of endemic pathogens). Future research would benefit from fitting the model
to real-world data for different bacterial species to enhance our understanding of AMR trends.
513
Methods

Streptococcus pneumoniae surveillance data

Antibiotic resistance trends reported in 2019 and 2020, provided by EARS-Net (European
Antimicrobial Resistance Surveillance Network) were acquired from a joint 2022 report on
antimicrobial resistance during 2020 by WHO and ECDC (European Centre for Disease
Prevention and Control and World Health Organization, 2022). The annual incidence of *S.*
pneumoniae invasive isolates for 2019 and 2020 was measured as the number of invasive isolates
from blood or cerebrospinal fluid. The proportion of resistant isolates represents the proportion of
isolates with phenotypic resistance to penicillin and macrolides using standardized bacterial
culture methods and EUCAST breakpoints. Out of 28 European countries that reported antibiotic
resistance data for *S. pneumoniae*, 24 countries had enough samples to establish 2019-2020
resistance trends for penicillin and macrolides. The resistance data for France, which were
subsequently analyzed, were provided by the CNRP (The French National Reference Center for
Pneumococci).

Model structure

We developed a pathogen co-circulation model (Appendix 2 – Figure 2) written using systems of
ordinary differential equations (ODEs) (Appendix 2; code available online at
https://github.com/alekskovacevic/antibiotic_resistance). The model simultaneously describes
potential infection with SARS-CoV-2 and colonization with antibiotic-sensitive and/or -resistant
strains of *S. pneumoniae* in a well-mixed community population. SARS-CoV-2 infection is
modeled by a Susceptible-Exposed-Infectious-Recovered (SEIR) process where individuals
become exposed to SARS-CoV-2 at rate β_C upon contact with other infected individuals. Infection
begins with a non-infectious exposed period lasting $1/\varepsilon$ days and is followed by an infectious
period lasting $1/\gamma^C$ days, eventually leading to recovery and immunization against future re-
infection. Waning immunity and competitive multi-strain SARS-CoV-2 dynamics are not
considered.

Individuals in S, E, I, and R compartments can be uncolonized with *S. pneumoniae* (U),
colonized with either a drug-sensitive (C^S) or a drug-resistant strain (C^R), or co-colonized with two
strains (C^{SS}, C^{RR}, C^{SR}). Colonization with each respective strain is acquired at rates β_S and $\beta_s f$
upon contact with other colonized individuals (Appendix 2 – Table 2). We assume a metabolic
cost of resistance, whereby the drug-resistant strain has a reduced intrinsic transmission rate
relative to the drug-sensitive strain due to reduced fitness, *f*. Bacterial carriage is cleared naturally

after an average duration of $\frac{1}{\gamma^S} = \frac{1}{\gamma^R} = \frac{1}{\gamma^{SR}} = \frac{1}{\gamma^{SS}} = \frac{1}{\gamma^{RR}}$ days, which we assume to be the same for

all types of carriers in our baseline scenario (in the scenarios assuming longer carriage duration of

antibiotic-resistant strains, $\frac{1}{\gamma^S} = \frac{1}{\gamma^{SS}}$ and $\frac{1}{\gamma^R} = \frac{1}{\gamma^{SR}} = \frac{1}{\gamma^{RR}}$). We further assume that some share of

the population is exposed to antibiotics at any given time, independent of bacterial carriage, with

individuals initiating antibiotic therapy at rate τ, which lasts for an average duration of $\frac{1}{r}$ days.

Another model assumption is that a proportion (p_{az}) of those infected with COVID-19 in the

community (between 0% and 20% of individuals in an I compartment) receive azithromycin

prescription from general practitioner reflecting azithromycin prescriptions in the early pandemic,

while the rest of the infectious individuals ($1 - p_{az}$) are exposed to the baseline antibiotic therapy.

We assume baseline treatment duration of seven days, on average, regardless of the antibiotic

prescribed and without any assumed persistence of the antibiotic in the system after the last dose

($\frac{1}{r}$). In case of antibiotic treatment with azithromycin for COVID-19 infected individuals we

assume the treatment lasts three days with antibiotics remaining in the system for additional 15.5

days after the last dose for a total of 18.5 days of antibiotic exposure where COVID-19 recovered

individuals (R_{az}) treated with azithromycin retain azithromycin in their system for an additional

11.5 days ($\frac{1}{r_{az}}$) after COVID-19 recovery. Individuals treated with antibiotics are unable to acquire

the sensitive strain. Antibiotics are assumed to clear colonization with sensitive strains at a rate ω

while having no direct impact on colonization with resistant strains. This bacterial colonization

process results in antibiotic selection for resistance via competition for limited hosts, facilitates

epidemiological coexistence between strains and is adapted from previous models of *S.*

pneumoniae (Colijn et al., 2010; Lipsitch et al., 2009; Mulberry et al., 2020). For a full list of

parameter values see Appendix 2 – Table 2.

Simulation in an early COVID-19 pandemic context

ODEs were integrated numerically using the R package deSolve to simulate and quantify

epidemiological dynamics (Soetaert et al., 2010). First, bacterial dynamics were simulated until

endemic equilibrium was achieved, under the assumption that *S. pneumoniae* was at endemic

equilibrium upon the emergence of COVID-19. Second, using equilibrium states as initial

conditions and re-initializing simulation time to *t*=0, a single SARS-CoV-2 infected individual

was introduced into the population and ODEs were again integrated numerically to *t*=365 days.

Parameter values used for simulation were taken from prior studies prioritizing French data and

are provided in Appendix 2 – Table 2.

These simulations were conducted in the context of an "early pandemic scenario"

coinciding with the implementation of population-wide NPIs to slow SARS-CoV-2 transmission.

This was conceived as the implementation of two 60-day lockdown periods starting on day 75 and

on day 305 in response to the simulated surge in COVID-19 cases. Lockdowns were assumed to

have three major potential impacts on population behavior and, in turn, the transmission dynamics

of SARS-CoV-2 and *S. pneumoniae*. These impacts were incorporated into simulations by

modifying epidemiological parameters in the model coincident with lockdowns. Three such

modifications were considered and switched on and off, considering all possible combinations.

First, lockdown led to reduced SARS-CoV-2 transmissibility by a factor θ_c. Second, lockdown led

to a population-wide change in antibiotic initiation rate by a factor *a* (representing modified

healthcare-seeking behavior leading to a reduction in the number of antibiotic prescriptions). Finally, lockdowns changed the pneumococcal disease risk by a factor IPD_{risk} (representing a reduced risk of developing an IPD due to the absence of other respiratory viruses).

Effect of SARS-CoV-2 basic reproduction number (R₀) on AMR

Impacts of SARS-CoV-2 on antibiotic-resistant IPD incidence may also depend on the characteristics of locally circulating SARS-CoV-2 R_0. To account for potential impacts of SARS-CoV-2 transmissibility and azithromycin use in the community, in simulations we varied (i) values of R_0 (basic reproduction number) for SARS-CoV-2 in France ($2 \leq R_0 \leq 4$) and (ii) the proportion of the COVID-19 infected individuals using azithromycin at simulation outset (from 0% to 20%).

Effect of within-host interactions on AMR

SARS-CoV-2 infection may impact progression from bacterial colonization to invasive bacterial disease at the within-host level. To incorporate this mechanism in our model, we included a within-host interaction term in scenario S19: the ecological interaction term (ψ_c) increases the rate of progression to invasive disease among colonized individuals who are also infected with SARS-CoV-2. The equations for calculating daily IPD incidence assuming within-host interactions due to SARS-CoV-2 co-infection with accompanying details can be found in Appendix 2.

Tables

Mechanisms:

1 Reduced community antibiotic prescribing.
2 Lockdown effect on reducing transmission of *S. pneumoniae*
3 Reduced risk of developing an IPD
4 Community azithromycin use in COVID-19 infected individuals
5 Longer carriage duration of antibiotic-resistant pneumococcal strains

Scenarios		Mechanisms					IPD inc.	AR (%)	Sp. (%)
		1	2	3	4	5			
Pre-pandemic 1 : ($d_S = d_R$)							10.8	20.0	NA
Pre-pandemic 2 : ($d_R > d_S$)						x	11.3	20.0	NA
Pandemic:	S1	x					10.9	19.2	+1.3
	S2		x				8.9	20.1	-36.1
	S3			x			5.9	20.0	0
	S4				x		9.9	23.7	-9.1
	S5					x	11.3	20.0	0
	S6	x	x				9.1	19.4	-35.2
	S7	x		x			6.0	19.4	+1.3
	S8	x			x		10.1	22.9	-8.0
	S9	x				x	11.5	19.3	+1.3
	S10		x	x			5.2	20.0	-36.1
	S11		x		x		8.9	20.1	-36.1
	S12		x			x	9.4	20.9	-34.3
	S13			x	x		5.6	22.5	-9.1
	S14			x		x	6.2	20.0	0
	S15				x	x	10.4	23.4	-9.1
	S16	x	x	x			5.3	19.6	-35.2
	S17	x	x		x		8.3	22.4	-41.3
	S18	x	x			x	9.6	20.3	-33.5
	S19	x		x	x		5.7	22.0	-8.0
	S20	x		x		x	6.3	19.5	+1.3
	S21	x			x	x	10.6	22.7	-7.9
	S22		x	x	x		5.0	22.0	-42.1
	S23		x	x		x	5.5	20.6	-34.3
	S24		x		x	x	8.7	23.9	-40.2
	S25			x	x	x	5.9	22.3	-9.1
	S26	x	x	x	x		5.0	21.6	-41.3
	S27	x	x	x		x	5.6	20.1	-33.5
	S28	x	x		x	x	8.8	23.2	-39.4
	S29	x		x	x	x	5.9	21.8	-7.9
	S30		x	x	x	x	5.2	22.5	-40.2
	S31	x	x	x	x	x	5.3	22.1	-39.4

REPORTED TRENDS:	IPD inc.	AR (%)	Sp. (%)
Pre-pandemic (FR, 2019)	10.5 [10.3-10.7]	26.2 (PENI) and 20.9 (ERY)	NA
Pandemic (FR, 2020)	5.8 [5.7-5.9]	35.5 (PENI) and 23.0 (ERY)	Stable
Pandemic (EU/EEA, 2020) General trends	*Decrease by 44.3% on avg.*	*Majority of EU countries report an increase*	*Generally stable*

Table 1. Five mechanisms implemented in 31 pandemic scenarios proposed to explain the
reported trends of IPD incidence, antibiotic resistance, and pneumococcal carriage in *S.*
pneumoniae. Scenarios explore all possible combinations of mechanisms proposed to test
hypotheses that can explain the reported trends of annual invasive pneumococcal disease incidence
(annual no. of cases per 100,000 inhabitants), antibiotic resistance (% of annual antibiotic-resistant
IPD cases among total IPD cases), and % change in the pneumococcal carriage prevalence at the
end of the first 60-day lockdown compared the prevalence before the lockdown. Model simulations
were initiated assuming the initial 20% antibiotic resistance. Two pre-pandemic scenarios assume
no SARS-CoV-2 circulation in the population and allow for the same 30-day carriage duration of
both antibiotic-sensitive and -resistant strains ($d_S = d_R$) or a longer, 40-day carriage duration of -
resistant strains ($d_R > d_S$). When implemented, these five mechanisms assume 18% reduction in
community antibiotic prescribing, a reduced risk of developing an IPD during (0.2) and after the
first lockdown (0.4), a 25% reduction in transmission of pneumococcal carriage during the first
lockdown, a 10% of azithromycin use among COVID-19 infected individuals, and a longer 40-
day carriage duration of -resistant strains. For a full list of parameters see Appendix 2 – Table 2.
Reported trends in European countries showed a decrease in annual IPD incidence by 44.3% on
average, an increase in antibiotic resistance, and generally stable asymptomatic pneumococcal
carriage in healthy individuals during the first lockdown period. Only scenarios S19 and S29 fulfill
all three reported trends during the COVID-19 pandemic in 2020 simultaneously while accounting
for the reported reduction in community antibiotic prescribing (d_S = carriage duration of antibiotic-
sensitive pneumococcal strains; d_R = carriage duration of antibiotic-resistant pneumococcal
strains; PENI = penicillin; ERY = erythromycin).

Figures

Figure 1. Antibiotic resistance trends in invasive *Streptococcus pneumoniae* isolates for the years 2019 and 2020. A. The proportion of invasive *S. pneumoniae* isolates resistant to penicillin and macrolides (azithromycin/ clarithromycin/ erythromycin) reported to EARS-Net (European Antimicrobial Resistance Surveillance Network) for 24 European countries. Error bars show 95% confidence intervals. **B.** The proportion of invasive *S. pneumoniae* isolates resistant to penicillin (MIC > 0.064 mg/L) and macrolides (erythromycin) according to age. Error bars show 95% confidence intervals. The total number of invasive pneumococcal isolates reported in France decreased by 45.1% from 2019 to 2020 (from 1119 to 614). Data are provided by the French National Reference Center for Pneumococci.

Figure 1 – figure supplement 1. Antibiotic resistance trends in invasive *Streptococcus pneumoniae* isolates in France, 2017-2020. The proportion of invasive *S. pneumoniae* isolates resistant to penicillin (A) and macrolides (B) according to age. Error bars show 95% confidence intervals. Across the period 2017-2020, a consistent decline in antibiotic resistance is observed for both penicillin and macrolides. Notably, this general trend experienced an anomaly in 2020, coinciding with the onset of the COVID-19 pandemic. Data are provided by the French National Reference Center for Pneumococci.

Figure 2. A modelling framework describing the transmission of SARS-CoV-2 and *Streptococcus pneumoniae* in the community setting, in the context of both general antibiotic prescribing and azithromycin prescribing for COVID-19 infected individuals. A. Non-pharmaceutical interventions (NPIs) implemented to control SARS-CoV-2 transmission (lockdown, face mask use, improved hygiene practices, travel restrictions, quarantine, telemedicine, and physical distancing) may also modify transmission of other pathogens, in addition to impacting antibiotic prescribing due to altered inter-individual contact and health-care seeking behavior. **B.** SEIR (Susceptible-Exposed-Infected-Recovered) model with antibiotic treatment compartments depicts interaction between SARS-CoV-2 infection and antibiotic prescribing, including both general community prescribing and azithromycin prescribing among individuals infected with SARS-CoV-2. **C.** Diagram depicting how pneumococcal colonization and the community antibiotic prescribing are affected by the COVID-19 pandemic impacts. Initiation of antibiotic treatment is assumed independent of bacterial carriage, reflecting widespread bystander selection for commensal bacteria like *S. pneumoniae*. For a complete modeling framework, see section S2 in Supporting Information.

Figure 3. Annual incidence of invasive pneumococcal disease (IPD), antibiotic resistance (AR%), and pneumococcal carriage prevalence for three different subpopulations. A. The elderly (≥65 years-old) **B.** general population (all ages), and **C.** children (< 5 years-old). Using pandemic scenario S19, which includes a combination of three different mechanisms: reduced community antibiotic prescribing, a reduced risk of developing an IPD, and community azithromycin use in COVID-19 infected individuals, we ran model simulations for three different subpopulations. For a full list of parameter values see Appendix 2 – Table 2. Annual IPD incidence (grey bars) decreased between 43% and 51% relative to the pre-pandemic (baseline) period with magnitude of a decrease depending on an age group and the level of azithromycin use in COVID-19 infected individuals. Antibiotic resistance (red bars) increased compared to the pre-pandemic (baseline) period in all age groups whenever azithromycin was used in COVID-19 infected. Black arrows indicate model outcomes that approximate the reported trends in antibiotic resistance in France for different age groups. Daily prevalence of total pneumococcal carriage remained relatively stable (solid-colored lines), exhibiting higher levels of decrease with increased azithromycin use. The prevalence of antibiotic-resistant pneumococcal carriage increased (dashed colored lines) over time in relation to SARS-CoV-2 outbreak (black dashed line) and higher azithromycin use. Highlighted time intervals (days 75-135 and 305-365) represent two lockdown periods.

Figure 4. The impact of varying SARS-CoV-2 R₀ and percentage of COVID-19 infected individuals taking azithromycin in scenario S19 on antibiotic resistance (%) and the annual incidence of antibiotic-resistant invasive pneumococcal disease (IPD). Hypothetical within-host interactions contribute to an excess incidence of antibiotic-resistant IPDs. (A) Cumulative incidence of antibiotic-resistant IPDs and antibiotic resistance increase with greater values of SARS-CoV-2 R₀ and higher percentage of the COVID-19 infected individuals taking azithromycin. The reproduction number for SARS-CoV-2 (R_0) in the community corresponds to the most common estimates of R_0 in France and other European countries ranging from $R_0 = 2$ to 4 (Allieta et al., 2022; D'Arienzo and Coniglio, 2020; Di Domenico et al., 2020; Flaxman et al., 2020; Liu et al., 2020; Roux et al., 2020; Salje et al., 2020). **(B) Annual excess in cumulative antibiotic-resistant IPD incidence in scenario S19 due to synergistic within-host ecological interactions compared to the same scenario with no within-host interactions and no azithromycin use (1.17 resistant IPD cases/100,000 inhabitants).** A rate of disease progression increased by a factor $\psi_c = 1$ (no within-host interaction) and $\psi_c = 40$ in scenario S19 applied to the general population assuming azithromycin use in 10% of the infected individuals resulted in approximately 0.06 and 0.75 additional cases of antibiotic-resistant disease per 100,000 inhabitants over the course of one year, respectively, compared to the scenario S19

814 assuming no within-host interaction and no azithromycin use (indicated by the black arrow). For
815 more details, see Appendix 2 - Figure 1.
816

References

Abdullahi O, Karani A, Tigoi CC, Mugo D, Kungu S, Wanjiru E, Jomo J, Musyimi R, Lipsitch
M, Scott JAG. 2012. Rates of acquisition and clearance of pneumococcal serotypes in the
nasopharynges of children in Kilifi District, Kenya. *J Infect Dis* **206**:1020–1029.
doi:10.1093/infdis/jis447
Allieta M, Allieta A, Rossi Sebastiano D. 2022. COVID-19 outbreak in Italy: estimation of
reproduction numbers over 2 months prior to phase 2. *J Public Health* **30**:2719–2727.
doi:10.1007/s10389-021-01567-1
Alpkvist H, Athlin S, Nauclér P, Herrmann B, Abdeldaim G, Slotved H-C, Hedlund J, Strålin K.
2015. Clinical and microbiological factors associated with high nasopharyngeal
pneumococcal density in patients with pneumococcal pneumonia. *PLOS ONE*
10:e0140112. doi:10.1371/journal.pone.0140112
Amin-Chowdhury Z, Aiano F, Mensah A, Sheppard CL, Litt D, Fry NK, Andrews N, Ramsay
ME, Ladhani SN. 2021. Impact of the coronavirus disease 2019 (COVID-19) pandemic
on invasive pneumococcal disease and risk of pneumococcal coinfection with severe
acute respiratory syndrome coronavirus 2 (SARS-CoV-2): prospective national cohort
study, England. *Clin Infect Dis* **72**:e65–e75. doi:10.1093/cid/ciaa1728
Andrews A, Bou-Antoun S, Guy R, Brown CS, Hopkins S, Gerver S. 2022. Respiratory
antibacterial prescribing in primary care and the COVID-19 pandemic in England, winter
season 2020–21. *J Antimicrob Chemother* **77**:799–802. doi:10.1093/jac/dkab443
Arduin H, Domenech De Cellès M, Guillemot D, Watier L, Opatowski L. 2017. An agent-based
model simulation of influenza interactions at the host level: insight into the influenza-
related burden of pneumococcal infections. *BMC Infect Dis* **17**:382. doi:10.1186/s12879-
017-2464-z
Baggs J, Jernigan JA, Halpin AL, Epstein L, Hatfield KM, McDonald LC. 2018. Risk of
subsequent sepsis within 90 days after a hospital stay by type of antibiotic exposure. *Clin*
Infect Dis **66**:1004–1012. doi:10.1093/cid/cix947
Bara W, Brun-Buisson C, Coignard B, Watier L. 2022. Outpatient antibiotic prescriptions in
France: patients and providers characteristics and impact of the COVID-19 pandemic.
Antibiotics **11**:643. doi:10.3390/antibiotics11050643
Bednarčuk N, Golić Jelić A, Stoisavljević Šatara S, Stojaković N, Marković Peković V,
Stojiljković MP, Popović N, Škrbić R. 2023. Antibiotic utilization during COVID-19: are
we over-prescribing? *Antibiotics* **12**:308. doi:10.3390/antibiotics12020308
Bennett JC, Emanuels A, Heimonen J, O'Hanlon J, Hughes JP, Han PD, Chow EJ, Ogokeh CE,
Rolfes MA, Lockwood CM, Pfau B, Uyeki TM, Shendure J, Hoag S, Fay K, Lee J,
Sibley TR, Rogers JH, Starita LM, Englund JA, Chu HY. 2023. *Streptococcus*
pneumoniae nasal carriage patterns with and without common respiratory virus detections
in households in Seattle, WA, USA before and during the COVID-19 pandemic. *Front*
Pediatr **11**.
Bhowmick S, Sokolov IM, Lentz HHK. 2023. Decoding the double trouble: a mathematical
modelling of co-infection dynamics of SARS-CoV-2 and influenza-like illness.
Biosystems **224**:104827. doi:10.1016/j.biosystems.2023.104827

Bogdanić N, Močibob L, Vidović T, Soldo A, Begovač J. 2022. Azithromycin consumption during the COVID-19 pandemic in Croatia, 2020. *PLoS ONE* **17**:e0263437. doi:10.1371/journal.pone.0263437

Braunstein SL, Lazar R, Wahnich A, Daskalakis DC, Blackstock OJ. 2020. COVID-19 infection among people with HIV in New York City: a population-level analysis of linked surveillance data. *Clin Infect Dis Off Publ Infect Dis Soc Am* ciaa1793. doi:10.1093/cid/ciaa1793

Brueggemann AB, Jansen Van Rensburg MJ, Shaw D, McCarthy ND, Jolley KA, Maiden MCJ, Van Der Linden MPG, Amin-Chowdhury Z, Bennett DE, Borrow R, Brandileone M-CC, Broughton K, Campbell R, Cao B, Casanova C, Choi EH, Chu YW, Clark SA, Claus H, Coelho J, Corcoran M, Cottrell S, Cunney RJ, Dalby T, Davies H, De Gouveia L, Deghmane A-E, Demczuk W, Desmet S, Drew RJ, Du Plessis M, Erlendsdottir H, Fry NK, Fuursted K, Gray SJ, Henriques-Normark B, Hale T, Hilty M, Hoffmann S, Humphreys H, Ip M, Jacobsson S, Johnston J, Kozakova J, Kristinsson KG, Krizova P, Kuch A, Ladhani SN, Lâm T-T, Lebedova V, Lindholm L, Litt DJ, Martin I, Martiny D, Mattheus W, McElligott M, Meehan M, Meiring S, Mölling P, Morfeldt E, Morgan J, Mulhall RM, Muñoz-Almagro C, Murdoch DR, Murphy J, Musilek M, Mzabi A, Perez-Argüello A, Perrin M, Perry M, Redin A, Roberts R, Roberts M, Rokney A, Ron M, Scott KJ, Sheppard CL, Siira L, Skoczyńska A, Sloan M, Slotved H-C, Smith AJ, Song JY, Taha M-K, Toropainen M, Tsang D, Vainio A, Van Sorge NM, Varon E, Vlach J, Vogel U, Vohrnova S, Von Gottberg A, Zanella RC, Zhou F. 2021. Changes in the incidence of invasive disease due to *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Neisseria meningitidis* during the COVID-19 pandemic in 26 countries and territories in the invasive respiratory infection surveillance initiative: a prospective analysis of surveillance data. *Lancet Digit Health* **3**:e360–e370. doi:10.1016/S2589-7500(21)00077-7

Buehrle DJ, Wagener MM, Nguyen MH, Clancy CJ. 2021. Trends in outpatient antibiotic prescriptions in the United States during the COVID-19 pandemic in 2020. *JAMA Netw Open* **4**:e2126114. doi:10.1001/jamanetworkopen.2021.26114

Cascante-Vega J, Galanti M, Schley K, Pei S, Shaman J. 2023. Inference of transmission dynamics and retrospective forecast of invasive meningococcal disease. *PLOS Comput Biol* **19**:e1011564. doi:10.1371/journal.pcbi.1011564

Chen C, Zhu P, Zhang Y, Liu B. 2021. Effect of the "Normalized epidemic prevention and control requirements" on hospital-acquired and community-acquired infections in China. *BMC Infect Dis* **21**:1178. doi:10.1186/s12879-021-06886-y

Chen Y, Li N, Lourenço J, Wang L, Cazelles B, Dong L, Li B, Liu Y, Jit M, Bosse NI, Abbott S, Velayudhan R, Wilder-Smith A, Tian H, Brady OJ, CMMID COVID-19 Working Group. 2022. Measuring the effects of COVID-19-related disruption on dengue transmission in southeast Asia and Latin America: a statistical modelling study. *Lancet Infect Dis* **22**:657–667. doi:10.1016/S1473-3099(22)00025-1

Chien Y-C, Lee Y-L, Liu P-Y, Lu M-C, Shao P-L, Lu P-L, Cheng S-H, Lin C-Y, Wu T-S, Yen M-Y, Wang L-S, Liu C-P, Lee W-S, Shi Z-Y, Chen Y-S, Wang F-D, Tseng S-H, Chen Yu-Hui, Sheng W-H, Lee C-M, Chen Yen-Hsu, Ko W-C, Hsueh P-R. 2021. National surveillance of antimicrobial susceptibilities to dalbavancin, telavancin, tedizolid, eravacycline, omadacycline and other comparator antibiotics and serotype distribution of invasive Streptococcus pneumoniae isolates in adults: results from the Surveillance of

Multicenter Antimicrobial Resistance in Taiwan (SMART) programme in 2017-2020. *J Glob Antimicrob Resist* **26**:308–316. doi:10.1016/j.jgar.2021.07.005

Clancy CJ, Buehrle DJ, Nguyen MH. 2020. PRO: The COVID-19 pandemic will result in increased antimicrobial resistance rates. *JAC-Antimicrob Resist* **2**:dlaa049. doi:10.1093/jacamr/dlaa049

Cohen R, Bidet P, Varon E, Béchet S, Cohen JF, Bonacorsi S, Levy C. 2023. Unprecedentedly high rates of Group A Streptococcus nasopharyngeal carriage in infants and toddlers in France, 2022–2023. *Infect Dis Now* **53**:104720. doi:10.1016/j.idnow.2023.104720

Colijn C, Cohen T, Fraser C, Hanage W, Goldstein E, Givon-Lavi N, Dagan R, Lipsitch M. 2010. What is the mechanism for persistent coexistence of drug-susceptible and drug-resistant strains of *Streptococcus pneumoniae*? *J R Soc Interface* **7**:905–919. doi:10.1098/rsif.2009.0400

Crisafulli S, Ientile V, L'Abbate L, Fontana A, Linguiti C, Manna S, Mercaldo M, Pagliaro C, Vezzaro M, Santacà K, Lora R, Moretti U, Reno C, Fantini MP, Corrao S, Barbato D, Tari M, Trifirò G, the ITA-COVID: COV-OUT Group. 2022. COVID-19 patient management in outpatient setting: a population-based study from Southern Italy. *J Clin Med* **11**:51. doi:10.3390/jcm11010051

Dagan R, Barkai G, Givon-Lavi N, Sharf AZ, Vardy D, Cohen T, Lipsitch M, Greenberg D. 2008. Seasonality of antibiotic-resistant *Streptococcus pneumoniae* that causes acute otitis media: a clue for an antibiotic-restriction policy? *J Infect Dis* **197**:1094–1102. doi:10.1086/528995

Dagan R, Beek BA van der, Ben-Shimol S, Greenberg D, Shemer-Avni Y, Weinberger DM, Danino D. 2023. The COVID-19 pandemic as an opportunity for unravelling the causative association between respiratory viruses and pneumococcus-associated disease in young children: a prospective study. *eBioMedicine* **90**. doi:10.1016/j.ebiom.2023.104493

Danion F, Margue M, Ruch Y, Séverac F, Hansmann Y. 2023. Seasonal variation in azithromycin prescription. *Lancet Infect Dis* **23**:277–278. doi:10.1016/S1473-3099(23)00009-9

Davies NG, Flasche S, Jit M, Atkins KE. 2019. Within-host dynamics shape antibiotic resistance in commensal bacteria. *Nat Ecol Evol* **3**:440–449. doi:10.1038/s41559-018-0786-x

D'Arienzo M, Coniglio A. 2020. Assessment of the SARS-CoV-2 basic reproduction number, R0, based on the early phase of COVID-19 outbreak in Italy. *Biosaf Health* **2**:57–59. doi:10.1016/j.bsheal.2020.03.004

Di Domenico L, Pullano G, Sabbatini CE, Boëlle P-Y, Colizza V. 2020. Impact of lockdown on COVID-19 epidemic in Île-de-France and possible exit strategies. *BMC Med* **18**:240. doi:10.1186/s12916-020-01698-4

Diavatopoulos DA, Short KR, Price JT, Wilksch JJ, Brown LE, Briles DE, Strugnell RA, Wijburg OL. 2010. Influenza A virus facilitates *Streptococcus pneumoniae* transmission and disease. *FASEB J* **24**:1789–1798. doi:10.1096/fj.09-146779

Doan T, Worden L, Hinterwirth A, Arzika AM, Maliki R, Abdou A, Zhong L, Chen C, Cook C, Lebas E, O'Brien KS, Oldenburg CE, Chow ED, Porco TC, Lipsitch M, Keenan JD, Lietman TM. 2020. Macrolide and nonmacrolide resistance with mass azithromycin distribution. *N Engl J Med* **383**:1941–1950. doi:10.1056/NEJMoa2002606

Domenech De Cellès M, Arduin H, Lévy-Bruhl D, Georges S, Souty C, Guillemot D, Watier L,
Opatowski L. 2019. Unraveling the seasonal epidemiology of pneumococcus. *Proc Natl*
Acad Sci **116**:1802–1807. doi:10.1073/pnas.1812388116
Duffy E, Thomas M, Hills T, Ritchie S. 2021. The impacts of New Zealand's COVID-19
epidemic response on community antibiotic use and hospitalisation for pneumonia,
peritonsillar abscess and rheumatic fever. *Lancet Reg Health West Pac* **12**:100162.
doi:10.1016/j.lanwpc.2021.100162
Ekdahl K, Ahlinder I, Hansson HB, Melander E, Mölstad S, Söderström M, Persson K. 1997.
Duration of nasopharyngeal carriage of penicillin-resistant *Streptococcus pneumoniae*:
experiences from the South Swedish pneumococcal intervention project. *Clin Infect Dis*
25:1113–1117. doi:10.1086/516103
Elias C, Sekri A, Leblanc P, Cucherat M, Vanhems P. 2021. The incubation period of COVID-
19: a meta-analysis. *Int J Infect Dis IJID Off Publ Int Soc Infect Dis* **104**:708–710.
doi:10.1016/j.ijid.2021.01.069
European Centre for Disease Prevention and Control, World Health Organization. 2022.
Antimicrobial resistance surveillance in Europe: 2022 : 2020 data. LU: Publications
Office.
Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, Whittaker C, Zhu H,
Berah T, Eaton JW, Monod M, Imperial College COVID-19 Response Team, Perez-
Guzman PN, Schmit N, Cilloni L, Ainslie KEC, Baguelin M, Boonyasiri A, Boyd O,
Cattarino L, Cooper LV, Cucunubá Z, Cuomo-Dannenburg G, Dighe A, Djaafara B,
Dorigatti I, Van Elsland SL, FitzJohn RG, Gaythorpe KAM, Geidelberg L, Grassly NC,
Green WD, Hallett T, Hamlet A, Hinsley W, Jeffrey B, Knock E, Laydon DJ, Nedjati-
Gilani G, Nouvellet P, Parag KV, Siveroni I, Thompson HA, Verity R, Volz E, Walters
CE, Wang H, Wang Y, Watson OJ, Winskill P, Xi X, Walker PGT, Ghani AC, Donnelly
CA, Riley S, Vollmer MAC, Ferguson NM, Okell LC, Bhatt S. 2020. Estimating the
effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* **584**:257–
261. doi:10.1038/s41586-020-2405-7
Foulds G, Shepard RM, Johnson RB. 1990. The pharmacokinetics of azithromycin in human
serum and tissues. *J Antimicrob Chemother* **25**:73–82. doi:10.1093/jac/25.suppl_A.73
Garcia-Vidal C, Sanjuan G, Moreno-García E, Puerta-Alcalde P, Garcia-Pouton N, Chumbita M,
Fernandez-Pittol M, Pitart C, Inciarte A, Bodro M, Morata L, Ambrosioni J, Grafia I,
Meira F, Macaya I, Cardozo C, Casals C, Tellez A, Castro P, Marco F, García F, Mensa
J, Martínez JA, Soriano A, Rico V, Hernández-Meneses M, Agüero D, Torres B,
González A, de la Mora L, Rojas J, Linares L, Fidalgo B, Rodriguez N, Nicolas D,
Albiach L, Muñoz J, Almuedo A, Camprubí D, Angeles Marcos M, Camprubí D,
Cilloniz C, Fernández S, Nicolas JM, Torres A. 2021. Incidence of co-infections and
superinfections in hospitalized patients with COVID-19: a retrospective cohort study.
Clin Microbiol Infect **27**:83–88. doi:10.1016/j.cmi.2020.07.041
Girard D, Finegan SM, Dunne MW, Lame ME. 2005. Enhanced efficacy of single-dose versus
multi-dose azithromycin regimens in preclinical infection models. *J Antimicrob*
Chemother **56**:365–371. doi:10.1093/jac/dki241
Grant J, Saux NL. 2021. Duration of antibiotic therapy for common infections. *J Assoc Med*
Microbiol Infect Dis Can **6**:181–197. doi:10.3138/jammi-2021-04-29

Henig O, Kehat O, Meijer SE, Chikly A, Weiss-Meilik A, Egoz E, Ben-Ami R, Paran Y. 2021.
Antibiotic use during the COVID-19 pandemic in a tertiary hospital with an ongoing
antibiotic stewardship program. *Antibiotics* **10**:1056. doi:10.3390/antibiotics10091056

Högberg LD, Vlahović-Palčevski V, Pereira C, Weist K, Monnet DL, ESAC-Net study group.
2021. Decrease in community antibiotic consumption during the COVID-19 pandemic,
EU/EEA, 2020. *Eurosurveillance* **26**. doi:10.2807/1560-7917.ES.2021.26.46.2101020

Homeniuk R, Collins C. 2021. How COVID-19 has affected general practice consultations and
income: general practitioner cross-sectional population survey evidence from Ireland.
BMJ Open **11**:e044685. doi:10.1136/bmjopen-2020-044685

Hsu J. 2020. How covid-19 is accelerating the threat of antimicrobial resistance. *BMJ*
369:m1983. doi:10.1136/bmj.m1983

Hussain AZ, Paudyal V, Hadi MA. 2021. Impact of the COVID-19 pandemic on the prescribing
patterns of first-line antibiotics in English primary care: a longitudinal analysis of
national prescribing dataset. *Antibiotics* **10**:591. doi:10.3390/antibiotics10050591

Kadambari S, Goldacre R, Morris E, Goldacre MJ, Pollard AJ. 2022. Indirect effects of the
covid-19 pandemic on childhood infection in England: population based observational
study. *BMJ* **376**:e067519. doi:10.1136/bmj-2021-067519

Jenness SM, Le Guillou A, Chandra C, Mann LM, Sanchez T, Westreich D, Marcus JL. 2021.
Projected HIV and bacterial sexually transmitted infection incidence following COVID-
19–related sexual distancing and clinical service interruption. *J Infect Dis* **223**:1019–
1028. doi:10.1093/infdis/jiab051

Karami Z, Knoop BT, Dofferhoff ASM, Blaauw MJT, Janssen NA, van Apeldoorn M,
Kerckhoffs APM, van de Maat JS, Hoogerwerf JJ, ten Oever J. 2021. Few bacterial co-
infections but frequent empiric antibiotic use in the early phase of hospitalized patients
with COVID-19: results from a multicentre retrospective cohort study in the Netherlands.
Infect Dis **53**:102–110. doi:10.1080/23744235.2020.1839672

Keenan JD, Bailey RL, West SK, Arzika AM, Hart J, Weaver J, Kalua K, Mrango Z, Ray KJ,
Cook C, Lebas E, O'Brien KS, Emerson PM, Porco TC, Leitman TM. 2018. Mass
azithromycin distribution for reducing childhood mortality in sub-Saharan Africa. *N Engl*
J Med **378**:1583–1592. doi:10.1056/NEJMoa1715474

Keenan JD, Klugman KP, McGee L, Vidal JE, Chochua S, Hawkins P, Cevallos V, Gebre T,
Tadesse Z, Emerson PM, Jorgensen JH, Gaynor BD, Lietman TM. 2015. Evidence for
clonal expansion after antibiotic selection pressure: pneumococcal multilocus sequence
types before and after mass azithromycin treatments. *J Infect Dis* **211**:988–994.
doi:10.1093/infdis/jiu552

Khouja T, Mitsantisuk K, Tadrous M, Suda KJ. 2022. Global consumption of antimicrobials:
impact of the WHO Global Action Plan on antimicrobial resistance and 2019 coronavirus
pandemic (COVID-19). *J Antimicrob Chemother* **77**:1491–1499.
doi:10.1093/jac/dkac028

Knight GM, Glover RE, McQuaid CF, Olaru ID, Gallandat K, Leclerc QJ, Fuller NM, Willcocks
SJ, Hasan R, Van Kleef E, Chandler CI. 2021. Antimicrobial resistance and COVID-19:
intersections and implications. *eLife* **10**:e64139. doi:10.7554/eLife.64139

Kournoutou GG, Dinos G. 2022. Azithromycin through the lens of the COVID-19 treatment.
Antibiotics **11**:1063. doi:10.3390/antibiotics11081063

Kuitunen I, Jääskeläinen J, Korppi M, Renko M. 2023. Antibiotic treatment duration for
community-acquired pneumonia in outpatient children in high-income countries—a

systematic review and meta-analysis. *Clin Infect Dis Off Publ Infect Dis Soc Am* **76**:e1123. doi:10.1093/cid/ciac374

Langford BJ, So M, Raybardhan S, Leung V, Soucy J-PR, Westwood D, Daneman N, MacFadden DR. 2021. Antibiotic prescribing in patients with COVID-19: rapid review and meta-analysis. *Clin Microbiol Infect* **27**:520–531. doi:10.1016/j.cmi.2020.12.018

Langford BJ, So M, Raybardhan S, Leung V, Westwood D, MacFadden DR, Soucy J-PR, Daneman N. 2020. Bacterial co-infection and secondary infection in patients with COVID-19: a living rapid review and meta-analysis. *Clin Microbiol Infect* **26**:1622–1629. doi:10.1016/j.cmi.2020.07.016

Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, Azman AS, Reich NG, Lessler J. 2020. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med* **172**:577–582. doi:10.7326/M20-0504

Lehtinen S, Blanquart F, Croucher NJ, Turner P, Lipsitch M, Fraser C. 2017. Evolution of antibiotic resistance is linked to any genetic mechanism affecting bacterial duration of carriage. *Proc Natl Acad Sci* **114**:1075–1080. doi:10.1073/pnas.1617849114

Lipsitch M, Colijn C, Cohen T, Hanage WP, Fraser C. 2009. No coexistence for free: neutral null models for multistrain pathogens. *Epidemics* **1**:2–13. doi:10.1016/j.epidem.2008.07.001

Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. 2020. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med* **27**:taaa021. doi:10.1093/jtm/taaa021

Malhotra-Kumar S, Lammens C, Coenen S, Van Herck K, Goossens H. 2007. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomised, double-blind, placebo-controlled study. *The Lancet* **369**:482–490. doi:10.1016/S0140-6736(07)60235-9

Mamun AA, Saatchi A, Xie M, Lishman H, Blondel-Hill E, Marra F, Patrick DM. 2021. Community antibiotic use at the population level during the SARS-CoV-2 pandemic in British Columbia, Canada. *Open Forum Infect Dis* **8**:ofab185. doi:10.1093/ofid/ofab185

McCullers JA, McAuley JL, Browall S, Iverson AR, Boyd KL, Henriques Normark B. 2010. Influenza enhances susceptibility to natural acquisition of and disease due to *Streptococcus pneumoniae* in ferrets. *J Infect Dis* **202**:1287–1295. doi:10.1086/656333

Melegaro A, Gay NJ, Medley GF. 2004. Estimating the transmission parameters of pneumococcal carriage in households. *Epidemiol Infect* **132**:433–441. doi:10.1017/S0950268804001980

Melnyk AH, Wong A, Kassen R. 2015. The fitness costs of antibiotic resistance mutations. *Evol Appl* **8**:273–283. doi:10.1111/eva.12196

Monnet DL, Harbarth S. 2020. Will coronavirus disease (COVID-19) have an impact on antimicrobial resistance? *Eurosurveillance* **25**. doi:10.2807/1560-7917.ES.2020.25.45.2001886

Mulberry N, Rutherford A, Colijn C. 2020. Systematic comparison of coexistence in models of drug-sensitive and drug-resistant pathogen strains. *Theor Popul Biol* **133**:150–158. doi:10.1016/j.tpb.2019.12.001

Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, Han C, Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G, Kashef Hamadani BH, Kumaran EAP, McManigal B, Achalapong S, Agarwal R, Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Babin F-

X, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Berkley JA, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalheiro C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Chiurchiù S, Chowdhury F, Clotaire Donatien R, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe S, Day NPJ, De Luca M, Dokova K, Dramowski A, Dunachie SJ, Duong Bich T, Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garcia C, Garrett D, Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Hsia Y, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Jenney AWJ, Khorana M, Khusuwan S, Kissoon N, Kobeissi E, Kostyanev T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Lim K, Limmathurotsakul D, Loftus MJ, Lunn M, Ma J, Manoharan A, Marks F, May J, Mayxay M, Mturi N, Munera-Huertas T, Musicha P, Musila LA, Mussi-Pinhata MM, Naidu RN, Nakamura T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Ochoa TJ, Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Ounchanum P, Pak GD, Paredes JL, Peleg AY, Perrone C, Phe T, Phommasone K, Plakkal N, Ponce-de-Leon A, Raad M, Ramdin T, Rattanavong S, Riddell A, Roberts T, Robotham JV, Roca A, Rosenthal VD, Rudd KE, Russell N, Sader HS, Saengchan W, Schnall J, Scott JAG, Seekaew S, Sharland M, Shivamallappa M, Sifuentes-Osornio J, Simpson AJ, Steenkeste N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Tigoi C, Turner C, Turner P, Van Doorn HR, Velaphi S, Vongpradith A, Vongsouvath M, Vu H, Walsh T, Walson JL, Waner S, Wangrangsimakul T, Wannapinij P, Wozniak T, Young Sharma TEMW, Yu KC, Zheng P, Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi M. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399**:629–655. doi:10.1016/S0140-6736(21)02724-0

Nandi A, Pecetta S, Bloom DE. 2023. Global antibiotic use during the COVID-19 pandemic: analysis of pharmaceutical sales data from 71 countries, 2020–2022. *eClinicalMedicine* **57**:101848. doi:10.1016/j.eclinm.2023.101848

Nation ML, Manna S, Tran HP, Nguyen CD, Vy LTT, Uyen DY, Phuong TL, Dai VTT, Ortika BD, Wee-Hee AC, Beissbarth J, Hinds J, Bright K, Smith-Vaughan H, Nguyen TV, Mulholland K, Temple B, Satzke C. 2023. Impact of COVID-19 nonpharmaceutical interventions on pneumococcal carriage prevalence and density in Vietnam. *Microbiol Spectr* **11**:e03615-22. doi:10.1128/spectrum.03615-22

Olesen SW, Lipsitch M, Grad YH. 2020. The role of "spillover" in antibiotic resistance. *Proc Natl Acad Sci* **117**:29063–29068. doi:10.1073/pnas.2013694117

Olwagen CP, Downs SL, Izu A, Tharasimbi L, Merwe LVD, Nunes MC, Madhi SA. 2023. Bacterial nasopharyngeal colonisation in children in South Africa before and during the COVID-19 pandemic: an observational study. *Lancet Microbe* **0**. doi:10.1016/S2666-5247(23)00260-4

Opatowski L, Varon E, Dupont C, Temime L, van der Werf S, Gutmann L, Boëlle P-Y, Watier L, Guillemot D. 2013. Assessing pneumococcal meningitis association with viral respiratory infections and antibiotics: insights from statistical and mathematical models. *Proc R Soc B Biol Sci* **280**:20130519. doi:10.1098/rspb.2013.0519

Palmer K, Monaco A, Kivipelto M, Onder G, Maggi S, Michel J-P, Prieto R, Sykara G, Donde S. 2020. The potential long-term impact of the COVID-19 outbreak on patients with non-communicable diseases in Europe: consequences for healthy ageing. *Aging Clin Exp Res* **32**:1189–1194. doi:10.1007/s40520-020-01601-4

Parveen M, Molla MMA, Yeasmin M, Nafisa T, Barna AA, Ghosh AK. 2020. Evidences on
irrational anti-microbial prescribing and consumption among COVID-19 positive patients
and possible mitigation strategies: a descriptive cross sectional study. *Bangladesh J Infect*
Dis S3–S7. doi:10.3329/bjid.v7i00.50155
Petrović V, Milosavljević B, Djilas M, Marković M, Vuković V, Andrijević I, Ristić M. 2022.
Pneumococcal nasopharyngeal carriage in children under 5 years of age at an outpatient
healthcare facility in Novi Sad, Serbia during the COVID-19 pandemic. *IJID Reg* **4**:88–
96. doi:10.1016/j.ijregi.2022.07.001
PRINCIPLE Trial Collaborative Group. 2021. Azithromycin for community treatment of
suspected COVID-19 in people at increased risk of an adverse clinical course in the UK
(PRINCIPLE): a randomised, controlled, open-label, adaptive platform trial. *Lancet Lond*
Engl **397**:1063–1074. doi:10.1016/S0140-6736(21)00461-X
Read B, McLeod M, Tonkin-Crine S, Ashiru-Oredope D, Quigley A, Brown CS, Lecky DM.
2023. Changes in public health-seeking behaviours for self-limiting respiratory tract
infections across England during the COVID-19 pandemic.
Rebelo JS, Domingues CPF, Dionisio F, Gomes MC, Botelho A, Nogueira T. 2021. COVID-19
lockdowns may reduce resistance genes diversity in the human microbiome and the need
for antibiotics. *Int J Mol Sci* **22**:6891. doi:10.3390/ijms22136891
Rhee C, Baker M, Vaidya V, Tucker R, Resnick A, Morris CA, Klompas M, CDC Prevention
epicenters program. 2020. Incidence of nosocomial COVID-19 in patients hospitalized at
a large US Academic Medical Center. *JAMA Netw Open* **3**:e2020498.
doi:10.1001/jamanetworkopen.2020.20498
Rose MA, Laurenz M, Sprenger R, Imöhl M, van der Linden M. 2021. Nasopharyngeal carriage
in children after the introduction of generalized infant pneumococcal conjugate vaccine
immunization in Germany. *Front Med* **8**.
Roux J, Massonnaud C, Crépey P. 2020. COVID-19: One-month impact of the French lockdown
on the epidemic burden (preprint). *Epidemiology*. doi:10.1101/2020.04.22.20075705
Rusic D, Vilovic M, Bukic J, Leskur D, Seselja Perisin A, Kumric M, Martinovic D, Petric A,
Modun D, Bozic J. 2021. Implications of COVID-19 pandemic on the emergence of
antimicrobial resistance: adjusting the response to future outbreaks. *Life* **11**:220.
doi:10.3390/life11030220
Rybak A, Levy C, Angoulvant F, Auvrignon A, Gembara P, Danis K, Vaux S, Levy-Bruhl D,
van der Werf S, Béchet S, Bonacorsi S, Assad Z, Lazzati A, Michel M, Kaguelidou F,
Faye A, Cohen R, Varon E, Ouldali N. 2022. Association of nonpharmaceutical
interventions during the COVID-19 pandemic with invasive pneumococcal disease,
pneumococcal carriage, and respiratory viral infections among children in France. *JAMA*
Netw Open **5**:e2218959. doi:10.1001/jamanetworkopen.2022.18959
Ryu S, Hwang Y, Ali ST, Kim D-S, Klein EY, Lau EHY, Cowling BJ. 2021. Decreased use of
broad-spectrum antibiotics during the coronavirus disease 2019 epidemic in South Korea.
J Infect Dis **224**:949–955. doi:10.1093/infdis/jiab208
Salje H, Tran Kiem C, Lefrancq N, Courtejoie N, Bosetti P, Paireau J, Andronico A, Hozé N,
Richet J, Dubost C-L, Le Strat Y, Lessler J, Levy-Bruhl D, Fontanet A, Opatowski L,
Boelle P-Y, Cauchemez S. 2020. Estimating the burden of SARS-CoV-2 in France.
Science **369**:208–211. doi:10.1126/science.abc3517
Seaton RA, Gibbons CL, Cooper L, Malcolm W, McKinney R, Dundas S, Griffith D, Jeffreys D,
Hamilton K, Choo-Kang B, Brittain S, Guthrie D, Sneddon J. 2020. Survey of antibiotic

and antifungal prescribing in patients with suspected and confirmed COVID-19 in

Scottish hospitals. *J Infect* **81**:952–960. doi:10.1016/j.jinf.2020.09.024

Sender V, Hentrich K, Henriques-Normark B. 2021. Virus-induced changes of the respiratory

tract environment promote secondary infections with *Streptococcus pneumoniae. Front*

Cell Infect Microbiol **11**.

Shaw D, Abad R, Amin-Chowdhury Z, Bautista A, Bennett D, Broughton K, Cao B, Casanova

C, Choi EH, Chu Y-W, Claus H, Coelho J, Corcoran M, Cottrell S, Cunney R, Cuypers

L, Dalby T, Davies H, Gouveia L de, Deghmane A-E, Demczuk W, Desmet S,

Domenech M, Drew R, Plessis M du, Duarte C, Erlendsdóttir H, Fry NK, Fuursted K,

Hale T, Henares D, Henriques-Normark B, Hilty M, Hoffmann S, Humphreys H, Ip M,

Jacobsson S, Johnson C, Johnston J, Jolley KA, Kawabata A, Kozakova J, Kristinsson

KG, Krizova P, Kuch A, Ladhani S, Lâm T-T, León ME, Lindholm L, Litt D, Maiden

MCJ, Martin I, Martiny D, Mattheus W, McCarthy ND, Meehan M, Meiring S, Mölling

P, Morfeldt E, Morgan J, Mulhall R, Muñoz-Almagro C, Murdoch D, Murphy J, Musilek

M, Mzabi A, Novakova L, Oftadeh S, Perez-Argüello A, Pérez-Vázquez M, Perrin M,

Perry M, Prevost B, Roberts M, Rokney A, Ron M, Sanabria OM, Scott KJ, Sheppard C,

Siira L, Sintchenko V, Skoczyńska A, Sloan M, Slotved H-C, Smith AJ, Steens A, Taha

M-K, Toropainen M, Tzanakaki G, Vainio A, Linden MPG van der, Sorge NM van,

Varon E, Vohrnova S, Gottberg A von, Yuste J, Zanella R, Zhou F, Brueggemann AB.

2023. Trends in invasive bacterial diseases during the first 2 years of the COVID-19

pandemic: analyses of prospective surveillance data from 30 countries and territories in

the IRIS Consortium. *Lancet Digit Health* **0**. doi:10.1016/S2589-7500(23)00108-5

Short KR, Reading PC, Wang N, Diavatopoulos DA, Wijburg OL. 2012. Increased

nasopharyngeal bacterial titers and local inflammation facilitate transmission of

Streptococcus pneumoniae. mBio **3**:e00255-12. doi:10.1128/mBio.00255-12

Shrestha S, Foxman B, Dawid S, Aiello AE, Davis BM, Berus J, Rohani P. 2013. Time and

dose-dependent risk of pneumococcal pneumonia following influenza: a model for

within-host interaction between influenza and *Streptococcus pneumoniae. J R Soc*

Interface **10**:20130233. doi:10.1098/rsif.2013.0233

Smith DRM, Opatowski L. 2021. COVID-19 containment measures and incidence of invasive

bacterial disease. *Lancet Digit Health* **3**:e331–e332. doi:10.1016/S2589-7500(21)00085-6

Smith DRM, Shirreff G, Temime L, Opatowski L. 2023. Collateral impacts of pandemic

COVID-19 drive the nosocomial spread of antibiotic resistance: a modelling study. *PLOS*

Med **20**:e1004240. doi:10.1371/journal.pmed.1004240

Soetaert K, Petzoldt T, Setzer RW. 2010. Solving differential equations in R: package deSolve. *J*

Stat Softw **33**:1–25. doi:10.18637/jss.v033.i09

Teng JLL, Fok KMN, Lin KPK, Chan E, Ma Y, Lau SKP, Woo PCY. 2022. Substantial decline

in invasive pneumococcal disease during coronavirus disease 2019 pandemic in Hong

Kong. *Clin Infect Dis Off Publ Infect Dis Soc Am* **74**:335–338. doi:10.1093/cid/ciab382

Tinggaard M, Slotved H-C, Petersen RF, Hovmand N, Benfield T. 2023. Decreased

pneumococcal carriage among older adults in Denmark during the COVID-19 lockdown.

Open Forum Infect Dis **10**:ofad365. doi:10.1093/ofid/ofad365

Tsay SV, Bartoces M, Gouin K, Kabbani S, Hicks LA. 2022. Antibiotic prescriptions associated

with COVID-19 outpatient visits among medicare beneficiaries, April 2020 to April

2021. *JAMA* **327**:2018. doi:10.1001/jama.2022.5471

Van Laethem J, Wuyts S, Van Laere S, Dirkx S, Seyler L, Mertens R, Ilsen B, Lacor P, Pierard D, Allard SD. 2021. Antibiotic prescriptions targeting bacterial respiratory infections in admitted patients with COVID-19: a prospective observational study. *Infect Dis Ther* **10**:2575–2591. doi:10.1007/s40121-021-00535-2

Wang L, Fu J, Liang Z, Chen J. 2017. Prevalence and serotype distribution of nasopharyngeal carriage of *Streptococcus pneumoniae* in China: a meta-analysis. *BMC Infect Dis* **17**:765. doi:10.1186/s12879-017-2816-8

Weill A, Drouin J, Desplas D, Cuenot F, Dray-Spira R, Zureik M. 2021. Usage des médicaments de ville en France durant l'épidémie de la Covid-19 – point de situation jusqu'au 25 avril 2021. Étude pharmaco-épidémiologique à partir des données de remboursement du SNDS. EPI-PHARE (Groupement d'intérêt scientifique ANSM-Cnam), 27 mai 2021.

Willen L, Ekinci E, Cuypers L, Theeten H, Desmet S. 2022. Infant pneumococcal carriage in Belgium not affected by COVID-19 containment measures. *Front Cell Infect Microbiol* **11**:825427. doi:10.3389/fcimb.2021.825427

Wittman SR, Martin JM, Mehrotra A, Ray KN. 2023. Antibiotic receipt during outpatient visits for COVID-19 in the US, From 2020 to 2022. *JAMA Health Forum* **4**:e225429. doi:10.1001/jamahealthforum.2022.5429

Wolter N, Tempia S, Cohen C, Madhi SA, Venter M, Moyes J, Walaza S, Malope-Kgokong B, Groome M, Du Plessis M, Magomani V, Pretorius M, Hellferscee O, Dawood H, Kahn K, Variava E, Klugman KP, Von Gottberg A. 2014. High nasopharyngeal pneumococcal density, increased by viral coinfection, is associated with invasive pneumococcal pneumonia. *J Infect Dis* **210**:1649–1657. doi:10.1093/infdis/jiu326

Wong A, Guevara LAB, Goult E, Briga M, Kramer SC, Kovacevic A, Opatowski L, Cellès MD de. 2023. The interactions of SARS-CoV-2 with cocirculating pathogens: epidemiological implications and current knowledge gaps. *PLOS Pathog* **19**:e1011167. doi:10.1371/journal.ppat.1011167

Wyllie AL, Mbodj S, Thammavongsa DA, Hislop MS, Yolda-Carr D, Waghela P, Nakahata M, Stahlfeld AE, Vega NJ, York A, Allicock OM, Wilkins G, Ouyang A, Siqueiros L, Strong Y, Anastasio K, Alexander-Parrish R, Arguedas A, Gessner BD, Weinberger DM. 2023. Persistence of pneumococcal carriage among older adults in the community despite COVID-19 mitigation measures. *Microbiol Spectr* **11**:e04879-22. doi:10.1128/spectrum.04879-22

Zhang T, Shen X, Liu R, Zhao L, Wang D, Lambert H, Cabral C. 2021. The impact of COVID-19 on primary health care and antibiotic prescribing in rural China: qualitative study. *BMC Health Serv Res* **21**:1048. doi:10.1186/s12913-021-07082-z

Zhou C, Jiang Y, Sun L, Li H, Liu X, Huang L. 2023. Secondary pulmonary infection and co-infection in elderly COVID-19 patients during the pandemics in a tertiary general hospital in Beijing, China. *Front Microbiol* **14**:1280026. doi:10.3389/fmicb.2023.1280026

Zhu N, Aylin P, Rawson T, Gilchrist M, Majeed A, Holmes A. 2021. Investigating the impact of COVID-19 on primary care antibiotic prescribing in North West London across two epidemic waves. *Clin Microbiol Infect* **27**:762–768. doi:10.1016/j.cmi.2021.02.007