***Ex vivo* midgut cultures of *Aedes aegypti* are efficiently infected by mosquito-borne**

alpha- and flaviviruses

Ana Lucia Rosales Rosas[a], Lanjiao Wang[a], Sara Goossens[a], Arno Cuvry[a], Li-Hsin Li[b],

Nanci Santos-Ferreira[a], Alina Soto[a], Kai Dallmeier[b], Joana Rocha-Pereira[a], Leen Delang[a]#

[a]KU Leuven Department of Microbiology, Immunology and Transplantation, Rega

Institute for Medical Research, Laboratory of Virology and Chemotherapy, Leuven,

Belgium.

[b]KU Leuven Department of Microbiology, Immunology and Transplantation, Rega

Institute for Medical Research, Molecular Vaccinology and Vaccine Discovery, Leuven,

Belgium.

Running Head: *Ex vivo* midgut cultures of *Aedes aegypti*

#Address correspondence to Prof. Leen Delang, leen.delang@kuleuven.be.

Abstract word count: 245

Text word count: 6586

Abstract (250 words)

Aedes aegypti mosquitoes can transmit several arboviruses, including chikungunya virus (CHIKV), dengue virus (DENV), and Zika virus (ZIKV). When blood-feeding on a virus-infected human, the mosquito ingests the virus into the midgut (stomach), where it replicates and must overcome the midgut barrier to disseminate to other organs and ultimately be transmitted via the saliva. Current tools to study mosquito-borne viruses (MBVs) include 2D-cell culture systems and *in vivo* mosquito infection models, which offer great advantages, yet have some limitations.

Here, we describe a long-term *ex vivo* culture of *Ae. aegypti* midguts. Cultured midguts were metabolically active for 7 days in a 96-well plate at 28°C and were permissive to ZIKV, DENV, Ross River virus (RRV) and CHIKV. *Ex vivo* midguts from *Culex pipiens* mosquitoes were found to be permissive to Usutu virus (USUV). Immunofluorescence staining confirmed viral protein synthesis in CHIKV-infected midguts of *Ae. aegypti*. Furthermore, fluorescence microscopy revealed replication and spread of a reporter DENV in specific regions of the midgut. In addition, two known antiviral molecules, β-D-N^4-hydroxycytidine (NHC) and 7-deaza-2'-*C*-methyladenosine (7DMA), were able to inhibit CHIKV and ZIKV replication, respectively, in the *ex vivo* model.

Together, our results show that *ex vivo* midguts can be efficiently infected with mosquito-borne alpha- and flaviviruses and employed to evaluate antiviral drugs. Furthermore, the setup can be extended to other mosquito species. *Ex vivo* midgut cultures could thus be a new model to study MBVs, offering the advantage of reduced biosafety measures compared to infecting living mosquitoes.

Importance (150 words)

Mosquito-borne viruses (MBVs) are a significant global health threat since they can cause

severe diseases in humans, such as hemorrhagic fever, encephalitis, and chronic arthritis.

MBVs rely on the mosquito vector to infect new hosts and perpetuate virus transmission.

No therapeutics are currently available. The study of arbovirus infection in the mosquito

vector can greatly contribute to elucidating strategies for controlling arbovirus

transmission. This work investigated the infection of midguts from *Aedes aegypti*

mosquitoes in an *ex vivo* platform. We found several MBVs capable of replicating in the

midgut tissue, including viruses of major health importance, such as dengue, chikungunya,

and Zika viruses. Additionally, antiviral compounds reduced arbovirus infection in the

cultured midgut tissue. Overall, the midgut model emerges as a useful tool for diverse

applications such as studying tissue-specific responses to virus infection and screening

potential anti-arboviral molecules.

1. Introduction

Arboviruses are a diverse group of viruses that rely on arthropod vectors such as

mosquitoes, ticks, and sandflies to infect susceptible vertebrate hosts and perpetuate

transmission. Three arboviral families/orders are of most clinical relevance: *Bunyavirales*,

Flaviviridae, and *Togaviridae*, as they encompass important arboviruses of global health

concern, such as dengue virus (DENV), chikungunya virus (CHIKV), yellow fever virus,

and Zika virus (ZIKV) (1). These four mosquito-borne viruses (MBVs) are mainly

transmitted by female *Aedes* (*Ae.*) mosquitoes (e.g., *Ae. aegypti* and *Ae. albopictus*) (2),

with *Ae. aegypti* mosquitoes considered highly competent vectors because of their

anthropophilic behavior (3).

When blood-feeding on a virus-infected human or animal, a female mosquito will ingest

the virus-containing blood in the midgut (i.e., mosquito stomach), where the virus must

replicate and overcome the midgut barrier to subsequently disseminate to mosquito

secondary organs. As the mosquito becomes systemically infected, resulting in a high viral

load, the virus will reach the salivary glands. The virus will be released in the saliva upon

the next bite, allowing virus transmission when the mosquito feeds on uninfected hosts (4,

5). The midgut is thus the initial site for infection and replication of any arbovirus in the

mosquito. Virus replication in and spread across midgut cells are a requisite for a

productive arbovirus infection in the mosquito (4).

In vitro cell cultures can be useful to study arbovirus infection and other relevant processes

in mosquitoes; however, they present several limitations. Most of the established mosquito-

derived cell lines are not derived from tissues relevant to specific stages of mosquito-virus

interactions (e.g., salivary glands or midgut), but were generated from larval or embryonic

tissue (e.g., *C6/36* [larvae] and *Aag-2* [embryos] cells) (6). Therefore, results obtained from

in vitro experiments are ought to be taken cautiously. Furthermore, cell lines are considered

homogenous cultures, both genetically and phenotypically, and single-cell type

populations might not well represent multicellular tissues such as a midgut (7, 8).

Moreover, some biological differences might arise among the same cell line coming from

different laboratories (7).

Another method commonly used to study mosquito-virus interactions is the mosquito

infection model. Through this approach, mosquito infection occurs artificially via a

bloodmeal (membrane filled with infectious blood), and mosquitoes are sacrificed to assess

viral infection and vector competence (9). Although living-infected mosquitoes provide

valuable transmission data, the value of these data is directly proportional to the labor-

intensive nature of the method. Transmission studies require skilled personnel for the

handling of infected mosquitoes and facilities with a high biosafety level that ensure proper

containment, and carry the risks of prick injury with virus-infected dissection tools or of

infected mosquitoes escaping. Additionally, considerable variability among mosquito

infection studies might exist as colonies bearing the same name can be highly genetically

divergent among diverse laboratories (10).

Ex vivo organ cultures have been arising as an advantageous tool in research and have also

been described for insects. For instance, *ex vivo* culture of insect organs has been

successfully and widely defined for ixodid ticks. The synganglion, midgut, and salivary

glands dissected from ticks were viable for 10 days and permissive to both Langat and

Powassan viruses. (8, 11). On the contrary, reports on *ex vivo* organ cultures from mosquito

tissues are scarce and mainly focused on germline and fat body tissues (12, 13). Despite

the essential implication of the mosquito midgut in arbovirus replication and dissemination,

no *ex vivo* mosquito midgut culture has been properly described. To date, only a short-term

ex vivo assay in mosquito midguts has been reported, in which the midguts were only

maintained for 35 hours post dissection in tissue culture plates (14).

Nevertheless, e*x vivo* cultures could offer a unique perspective to (i) study vector-virus

interactions in a relevant tissue, for example, the mosquito midgut, (ii) reveal tissue-

specific responses to virus infection, or (iii) study tissue-specific metabolic processes.

Furthermore, *ex vivo* mosquito midgut cultures could provide a convenient and relevant

platform to generate results that can be extrapolated to living mosquitoes. Additionally, *ex*

vivo midgut cultures provide a controlled environment in which many parameters can be

easily adjusted, and they constitute a safer option due to the low degree of containment

required while working with BSL2/3 arboviruses, compared to working with live mosquito

infection models.

On the lookout for novel tools that could be of use to study arbovirus infection within the

arthropod vector, we dissected *Ae. aegypti* midguts and established their viability for a 7-

day period. We also infected the midgut tissues with several mosquito-borne alpha- and

flaviviruses, confirming the production of infectious viral particles. Furthermore,

previously reported antiviral drugs were able to reduce arbovirus replication in the treated

midguts. We thus developed an *ex vivo* mosquito midgut model that constitutes a valuable

tool to study MBVs which complements the arsenal available to the mosquito arbovirus

research field.

2. Results

Viability of *Ae. aegypti* midguts cultured *ex vivo*

We first determined the viability of the dissected mosquito midguts in 96-well culture

plates. Since motility could be observed along the gut tissue following the dissection, a

bioassay was performed based on the number of contractions by the hindgut

(Supplementary material, Movie 1). The same area of the hindgut was analyzed for all

biological replicates (Fig. 1, b). Dissected midguts presented peristalsis in the hindgut for

up to 7 days post dissection (d.p.d.) (Fig. 1, d), indicating that the tissue was viable. At 10

d.p.d., the mean frequency in the *ex vivo* cultured midguts was significantly reduced

compared to day 0 measurements (day 0: 14.37 s vs day 10: 6.74 s) (Fig. 1, d), suggesting

that the tissue's viability had decreased.

We further assessed the metabolic activity of the *ex vivo* cultured midguts in a resazurin

salt-based assay. The dissected midguts remained metabolically active up until 7 d.p.d.

(Fig. 1, e), corresponding with the data obtained from gut peristalsis. Fluorescence intensity

in the tested midguts was still slightly higher than the "no organ" controls (NO, used to

measure background signal) and negative controls (NC, paraformaldehyde fixated

midguts) at 8, 9, and 10 d.p.d.. However, there was more variability among biological

replicates and therefore 7 d.p.d. was selected as the end time point for following

experiments.

Additionally, the effect of the neurotransmitter serotonin and the anticholinergic drug

atropine on peristalsis was assessed to rule out that this visual feature might be a reflex.

Midguts (day 0 post dissection) were exposed to either serotonin, atropine, or midgut

medium (mock-exposure), after which their gut peristalsis was recorded. The mean

frequency displayed by the mock-exposed midguts was 7.349 s (5 biological replicates,

median value: 7.48 s). Midguts exposed for 1 hour to serotonin displayed a slight increase

in gut motility, with a mean peristaltic period of 12.76 s (6 biological replicates, median

value: 10.61 s), whereas atropine-exposed midguts exhibit a somewhat reduced peristalsis,

with a mean peristaltic period of 6.72 s (6 biological replicates, median value: 7.58 s).

Despite the observed trends, there were no statistically significant differences among the

conditions evaluated (Supplementary Fig. S1).

Infection of dissected *Ae. aegypti* midguts with mosquito-borne alpha- and

flaviviruses

Next, *ex vivo* virus infection was performed in the cultured midguts to test their capacity

to support the replication of several arboviruses. Replication kinetics were evaluated for

Ross River virus (RRV) and CHIKV as representative members of the alphavirus genus of

the *Togaviridae* family. Both viruses were able to efficiently replicate in the *ex vivo*

cultured midguts with an increase in viral RNA levels between 1 and 3 days post infection

(d.p.i.), compared to viral RNA levels at 2 hours p.i. (h.p.i.) (Fig. 2, a). Infectious virus

titers of RRV and CHIKV increased accordingly in the midgut tissue (Fig 2, b), while no

infectious virus was observed yet at 2 h.p.i. (data not shown). The peak of viral RNA and

viral titer levels was detected at 3 d.p.i. for both alphaviruses.

To evaluate whether *Aedes*-borne flaviviruses could replicate in the *ex vivo* midguts, the

replication kinetics of DENV-2 and ZIKV were studied. As both viruses needed a longer

period to replicate in the midgut tissue (data not shown), viral RNA and infectious virus

levels were measured starting at 5 d.p.i.. Viral titers were increased at 5 d.p.i. and further

peaked at 7 d.p.i., with DENV-2 reaching RNA levels up to 2.5×10^8 genome

copies/midgut and a viral titer of 21 PFU/midgut. On the contrary, ZIKV replication and

infection presented more variability among individual midguts. The maximum amount of

ZIKV RNA and infectious virus quantified at 7 d.p.i. were 2.1×10^6 genome copies/midgut

and 4.2×10^2 PFU/midgut, respectively (Fig 3).

Usutu virus (USUV) can modestly replicate in *ex vivo* cultured midguts from *Culex*

***pipiens* mosquitoes**

As a proof-of-concept, the *ex vivo* midgut model was applied to another mosquito species:

Culex (Cx.) pipiens. The midguts of *Cx. pipiens* thrived in culture and showed peristaltic

movements during their incubation, as *Ae. aegypti* midguts did. Therefore, we evaluated

the susceptibility of these *Cx.* midguts to USUV, as *Cx. pipiens* mosquitoes are considered

a competent vector for this virus (15). USUV infection was assessed in the midguts at 7

d.p.i. by qRT-PCR and plaque assay. Only 50% (3 out of 6) of the midguts became infected

with USUV, reaching up to 1.8×10^6 genome copies/midgut. However, only 2 out of these

3 USUV-infected midguts contained infectious virus, amounting to 6.4×10^2 PFU/mL. Of

note, a considerable amount of viral RNA was detected in samples corresponding to 2 h.p.i.

and in 2 out of the 3 negative controls included in the assay, yet no infectious virus was

detected in these samples (Fig. 4, a, b).

CHIKV viral protein synthesis in infected *ex vivo* midguts

Previous infection experiments indicated the susceptibility of the *ex vivo* midguts to

infection with several arboviruses. To further corroborate these results, the CHIKV E2

glycoprotein was visualized in CHIKV-infected midguts at day 3 p.i. by immunostaining.

Specific staining for this viral protein could be observed in the infected guts, while no

signal was observed in the negative controls (fixated guts that followed the same infection

protocol; Fig. 5, a). At day 3 p.i., E2 protein synthesis was mainly detected and spread

along the posterior midgut region (Fig. 5, b, c), with some infection foci located in the

hindgut region (Fig. 5, c). The E2 signal was also present in the tracheal tubes that remained

attached to the midgut after dissection (Fig. 5, b, arrow heads).

Infection of *ex vivo* cultured midguts with an mCherry-expressing DENV-2

To follow the progression of virus infection in the midguts by imaging, cultured mosquito

midguts were infected with DENV-2 expressing the red fluorescent protein mCherry

(DV2/mCherry,(16)) and imaged using the FLoid™ Cell Imaging Station (Life

Technologies) at selected time points p.i.. The mCherry signal was used as a proxy for

infection. DENV-2 mCherry infection was observed initially at day 3 p.i. as single or few

foci in the posterior midgut region (Supplementary Fig. S2, a). Over time, these focal

infection points increased in number and spread to neighboring areas, primarily along the

posterior region of the midgut and the hindgut (Supplementary Fig. S2, b-e). No mCherry

expression was observed in the negative control or mock-infected midguts at day 7 p.i.

(Fig. 6, a). Following 7 days of infection, the mCherry signal had spread mostly along the

posterior midgut region, forming infection foci comprising multiple cells (Fig. 6, b, c).

Moreover, some infected cells were found in the tracheal tubes that remained in the midgut

tissue after dissection (Supplementary Fig. S2, h).

Arbovirus replication is reduced upon treatment with antiviral drugs in midgut

cultures

We next assessed whether the *ex vivo* mosquito midguts could be used to evaluate the

antiviral activity of inhibitors against arboviruses. To this end, the antiviral activity of β-

D-N^4-hydroxycytidine, also known as EIDD-1931 or NHC, was tested against CHIKV at a

concentration of 50 µM. At day 3 p.i., no difference in the CHIKV RNA levels was

observed between the untreated and NHC-treated groups (Fig. 7, a). In contrast, NHC

significantly reduced the infectious virus levels by 1 log (mean values, control: 1.0×10^4;

50 µM NHC: 1.1×10^3 PFU/midgut) (Fig. 7, b).

As the *ex vivo* cultured midguts also supported flavivirus infection, the viral polymerase

inhibitor 7-Deaza-2'-*C*-Methyladenosine (7-DMA) was tested against ZIKV. Viral RNA

levels were reduced for all treated midguts (25, 100, and 200 µM of 7-DMA) compared to

the untreated group at day 7.p.i.. However, only the groups treated with 7-DMA at 100 and

200 µM showed a significant difference compared to the untreated group, with 1.5 log and

2.4 log reductions (virus control: 2.6×10^6; 100 µM: 7.0×10^4; 200 µM: 9.5×10^3 mean

genome copies/midgut), respectively (Fig. 7, c).

3. Discussion

The study of arbovirus infection in the mosquito vector relies mainly on a combination of *in vitro* and *in vivo* approaches, which has yielded great progress in the knowledge regarding arbovirus biology, virus-vector interactions, and vector competence. While convenient and handy, *in vitro* cell culture systems have the shortcoming of not being as biologically relevant as mosquito infection models, aside from the limited selection of mosquito cell lines available. On the other hand, working with living infected mosquitoes grants useful information when studying virus-vector interactions or vector competence for a specific virus, but such *in vivo* models are not always accessible and require the implementation of cumbersome safety measures ensuring adequate containment while working with BSL2/3 pathogens (17). *Ex vivo* organ culture methods have not yet been described in-depth for the study of MBVs. Hence, in this study, we have established an *ex vivo* mosquito midgut model using dissected midguts from *Ae. aegypti* mosquitoes. When cultured *ex vivo,* the midguts displayed peristaltic contractions, mainly observed as waves along the hindgut, which is a frequent occurrence for this type of tissue (18). Here, we used this visual feature as a proxy for viability, paired with a custom script for gut motility in zebrafish (19) to measure the peristaltic period (time between contractions) in the *ex vivo* cultured guts, which remained stable for 7 days p.d.. This observation was further corroborated when measuring the metabolic activity of the dissected midguts over time in culture with PrestoBlue™ (PB). This resazurin-based reagent has been successfully used to assess cell viability and cytotoxicity in both two-dimensional cell monolayers and 3D cellular interfaces (including organ explants) (20–23). Here, we found that only after 7

hours of incubation of the *ex vivo* midguts (using multiple midgut organs per well) in

presence of the PB reagent yielded a colorimetric change.

The *ex vivo* cultured mosquito midguts supported infection with four arboviruses, despite

the unusual route of infection (through incubation with virus inoculum) that differs from

what occurs naturally in mosquitoes (infection via blood feeding). Both viral RNA and

infectious virus particles were detected in the mosquito midgut organs following a

progressive increase over time, which also confirmed the viability of the midguts. These

replication kinetics results indicated the preservation of midgut tissue in an *ex vivo* culture

set-up and, more specifically, the presence of midgut cells that could constitute the target

for arbovirus infection.

With *Ae. aegypti* being the main vector for both DENV-2 and ZIKV, it was however

unexpected that infectious virus titers for both flaviviruses detected in the *ex vivo* midguts

did not reach higher levels than what was inoculated (1×10^4 PFU/mL), in contrast to viral

RNA levels. Kinetics of DENV replication in mosquitoes have reported a steady increase

in infectious virus until 8 d.p.i., after which it normally declined without affecting viral

RNA levels (24, 25). Such results could explain the discrepancy seen between flavivirus

RNA and infectious levels. More importantly, it has been described that within 2 hours of

infection with DENV or ZIKV (both *in vivo* and *ex vivo*), there is a rapid induction of

apoptosis in the *Ae. aegypti* midgut epithelium in an attempt of the host to control flavivirus

infection (26). Nonetheless, this process might cause tissue damage, which correlates with

the high midgut cell turnover rate in mosquitoes during a bloodmeal digestion (27). As

such, we cannot disregard that the *ex vivo* midguts in our experiments undergo the same

process during DENV or ZIKV infection, consequently damaging the tissue and thus

limiting flaviviral replication. This hypothesis may also apply to *ex vivo* midguts of *Cx.*

pipiens mosquitoes, as a comparable replication pattern can be observed when assessing

USUV (flavivirus) infection at day 7 p.i.. Further addition of an apoptosis inhibitor to the

virus inoculum used to infect the *ex vivo* midguts might elucidate whether rapid induction

of apoptosis in the midgut epithelium is indeed a limiting factor for flavivirus replication

in this setup.

Active virus replication was confirmed for CHIKV through an immunofluorescence assay.

Envelope (E2) protein synthesis was localized in the posterior midgut region, where it

would also occur when the mosquito ingests a CHIKV-infectious bloodmeal (28),

regardless of the infection method employed with the *ex vivo* cultured midguts. Moreover,

CHIKV infection of tracheal tubes in the tissue could be observed, consistent with *in vivo*

reports (28). Similar results were obtained when analyzing the progression of

DENV2/mCherry infection in the *ex vivo* midguts. Previously, DENV-2 infection in *Ae.*

aegypti (Chetumal strain) midguts was reported to start with individual infected epithelial

cells detected as early as 2 d.p.i., slowly progressing to infection foci consisting of multiple

cells until the whole midgut organ was infected at 7-10 d.p.i. (24). In agreement with this,

a less strong but akin infection pattern was observed in DENV2/mCherry-infected midguts,

with few foci of several infected cells by 7 d.p.i.. Together, these data show that arbovirus

infection occurred at considerable levels in *ex vivo* cultured midguts and therefore it could

be used to study other aspects of arbovirus infection and facilitate the collection of

preliminary data before experimenting with mosquitoes *in vivo*.

The use of antiviral compounds to inhibit virus infection in the mosquito vector is an

innovative concept that might reduce or block arbovirus transmission from mosquitoes to

humans. This idea involves antiviral molecules being ingested by adult mosquitoes when

they take a bloodmeal on a mammalian host undergoing antiviral treatment. As the

mosquito midgut is the entry point and key replication site for MBVs, studying the effect

of antiviral compounds in the midgut tissue would be of great interest. For this purpose,

the antiviral activity of two inhibitors was assessed in the *ex vivo* mosquito midguts. NHC

is a nucleoside analog that has been characterized as a potent antiviral drug against

alphaviruses *in vitro*, including CHIKV and Venezuelan equine encephalitis virus (VEEV)

(29, 30). Various assays point to the compound acting as a pyrimidine analog that may

target the viral polymerase domain of nsP4, provoking chain-termination (31). In addition,

NHC also induces a high level of mutations in virus-specific RNAs, resulting in lethal

mutagenesis (32). Consistent with these findings, CHIKV RNA levels were not reduced in

NHC-treated midguts, but there was a marked decrease in virus infectivity. The viral

polymerase inhibitor 7DMA has shown potent anti-ZIKV activity *in vitro* and delayed

disease progression in mice (33). In the *ex vivo* mosquito midguts, 7DMA significantly

reduced ZIKV RNA loads. Although the RNA levels presented some variability among the

ZIKV-infected midguts, the inhibitory effect of 7DMA in the *ex vivo* mosquito midguts

was significant and followed a dose-response relationship. These results indicate that *ex*

vivo midgut cultures could be used to rationally select potential arbovirus-blocking

molecules to be tested in living mosquitoes at a later stage.

In summary, we have established a long-term *ex vivo* mosquito midgut culture. To support

this model, we have (a) assessed the viability of the *ex vivo* cultured midguts, (b)

determined the replication kinetics of two alphaviruses (RRV and CHIKV) and two

flaviviruses (DENV-2 and ZIKV), (c) detected viral protein synthesis and followed live

arbovirus infection in infected midguts, (d) demonstrated that the *ex vivo* protocol can be

translated to other mosquito genera, and, lastly, (e) evaluated the antiviral activity of two

inhibitors against CHIKV and ZIKV. Altogether, we have provided a reference for the use

of *ex vivo* mosquito midguts as a tool to study arbovirus infection and related processes,

while offering groundwork for developing other *ex vivo* mosquito organ cultures that can

potentially provide insightful data, like salivary glands.

4. Materials and Methods

4.1. Cells

Mammalian cells

African green monkey kidney cells (Vero cells, ATCC CCL-81) and Vero E6 cells (ATCC

CRL-1586) were cultured in minimum essential medium (MEM 1X) enriched with 10%

Fetal Bovine Serum (FBS), 1% sodium bicarbonate, 1% L-glutamine and 1% non-essential

amino acids (NEAA). Baby hamster kidney cells (BHK, ATCC CCL-10) were maintained

in Dulbecco's Modified Eagle's Medium (DMEM) containing 10% FBS, 1% sodium

bicarbonate and 1% L-glutamine. Mammalian cell cultures were incubated at 37°C, with

5% CO_2.

Mosquito cells

Ae. albopictus larval cells (C6/36, obtained from ATCC, CRL-1660) were maintained in

Leibovitz's L-15 medium containing 10% FBS, 1% Penicillin-Streptomycin (PenStrep),

1% NEAA, and 1% HEPES buffer. Mosquito-derived cell lines were incubated at 28°C,

without CO_2.

For cell culture assays containing virus or virus-infected material, the concentration of FBS

in the medium was reduced to 2%, for both mammalian and mosquito cells. All cell culture

media and supplements were obtained from Gibco^{TM}, ThermoFisher Scientific (Aalst,

Belgium).

4.2. Viruses

Flaviviruses

Dengue virus serotype 2 (DENV-2/TH/1974, isolated in 1974 from human serum collected

in Bangkok, Thailand, GenBank MK268692.1) was kindly provided by Prof. A. Failloux

(Institut Pasteur, Paris, France) (34); ZIKV (SL1602, Suriname strain, GenBank

KY348640.1) was acquired via the EVAg consortium (https://www.european-virus-

archive.com). The infectious clone DENV-2 pDVWS601 used for the construction of the

DENV-2 reporter virus expressing the red fluorescent protein mCherry (DV2/mCherry,

New Guinea C strain, NGC, GenBank AF038403.1) was kindly provided by Prof. Andrew

Davidson (University of Bristol, Bristol, UK) (16).

Alphaviruses

Ross River virus (RRV) was received from the National Collection of Pathogenic Viruses

(UK; catalog number 0005281v); and CHIKV (Indian Ocean strain 899, GenBank

FJ959103.1) was generously provided by Prof. Drosten (University of Bonn, Bonn,

Germany) (35).

Virus stocks were prepared by passaging the isolates on Vero (for CHIKV and USUV) or

C6/36 cells (for ZIKV, DENV-2, and RRV). Viral titers of the stocks were determined via

plaque assay or end-point titration on Vero or BHK cells.

4.3. Compounds

364 7-Deaza-2'-*C*-Methyladenosine (7-DMA) was purchased from Carbosynth (Berkshire,

365 UK) and dissolved in DMSO. β-D-N^4-hydroxycytidine (EIDD-1931 or NHC) was

purchased from MedChemExpress (Monmouth Junction, NJ, USA) and dissolved in DMSO.

4.4. *Ae. aegypti* rearing

Ae. aegypti Paea (Papeete, Tahiti, collected in 1994) were obtained via the Infavec2 consortium. For each rearing, eggs were hatched in dechlorinated tap water. After hatching, groups of ±400 larvae were transferred into trays containing 3 L of dechlorinated tap water and fed every day with a yeast tablet (Gayelord Hauser, Saint-Genis-Laval, France) until the pupae stage. Pupae were placed in small plastic containers inside cardboard cups for their emergence. Adult mosquitoes were supplied with cotton balls soaked in a 10% sucrose solution supplemented with 100 U/mL and 100 µg/mL of PenStrep. Cardboard cups containing adults were maintained at 28 ± 1 °C with a light/dark cycle of 16/8 h and 80% relative humidity.

4.5. *Cx. pipiens* rearing

Cx. pipiens biotype *pipiens* were kindly provided by Prof. Sander Koenraadt (Wageningen University & Research, Wageningen, Netherlands) (15). For each rearing, eggs rafts were hatched in trays containing 2 L of Milli-Q water (Synergy® UV, Merck, Germany). Larvae were fed continuously until the pupae stage with TetraMin® baby fish food (Tetra, Spectrum Brands, Germany). Pupae were collected as described in 4.4. Cardboard cups containing adults were maintained at 25 ± 1 °C with a light/dark cycle of 16/8 h and 70% relative humidity.

4.6. Mosquito midgut dissection and *ex vivo* culture

Unfed, antibiotic-treated female mosquitoes (3 – 7 days old) were cold-anaesthetized and surface sterilized by soaking in 70% ethanol for 20 s followed by soaking in Dulbecco's

phosphate-buffered saline (PBS) for 20 s. Next, mosquitoes were dissected in PBS on a

petri dish using the stereomicroscope (VisiScope®, VWR). In brief, the midgut was

exposed by carefully pulling the second to last segment of the mosquito abdomen. The gut

of the mosquito was excised in its entirety (foregut, anterior and posterior midgut, and

hindgut) to keep the tissue of interest (midgut) from degradation, and tracheal tubes were

removed as much as possible without damaging the tissue. Following dissection, mosquito

guts were washed twice in midgut medium, consisting of: Leibovitz's L-15 medium

supplemented with 2% FBS, 100 U/mL of PenStrep, 50 µg/mL of kanamycin, and 0.25

µg/mL of amphotericin B (Sigma Aldrich, USA), and finally placed in a 96-well tissue

culture plate with a clear bottom (PerkinElmer®, USA) containing midgut medium. One

mosquito gut was placed in each testing well. Plates were maintained at 28°C, without CO_2.

Of note, handling and optimization of the *ex vivo* mosquito midguts culture are extensively

described in Supplementary Appendix 1.

4.7. Videography of gut peristalsis and analysis

Ex vivo cultured mosquito midguts were recorded with the Leica DMi8 microscope to

quantify the peristalsis observed *ex vivo*. A drop of carboxymethyl cellulose (CMC) 0.8%

diluted in Leibovitz's L-15 medium was deposited on a microscope glass slide and one

mosquito midgut was soaked in the drop. Forceps were used to gently arrange the hindgut

in a position suitable for analysis (horizontally positioned on the slide). Each hindgut was

recorded for a total of 1 minute 7 s (1 frame/0.3 s) and further discarded. This procedure

was repeated at several time points during incubation starting from day 0 (right after

dissection) to assess the contractibility, and hence viability, of the *ex vivo* cultured midguts

over time.

For the analysis of the gut peristalsis, frames generated for each midgut were processed in

IgorPro (Wavemetrics, USA) using a custom script described for the measurement of

zebrafish gut motility (19). This script quantifies the changes in pixel intensity in a

designated area of analysis as it sequentially goes through all frames generated during the

video recording. As output, the software indicates individual peristaltic periodicity

(seconds in between contractions) per point of evaluation and overall averages of peristaltic

periodicity for each midgut analyzed. These outputs were annotated manually to check for

artifacts that could be generated by debris or tracheal tubes.

Serotonin hydrochloride (Sigma Aldrich, USA) and atropine sulfate salt monohydrate

(Sigma Aldrich, USA) were used to test the responsiveness of the mosquito midgut tissue.

Midguts were incubated with either serotonin (20 µM), atropine (20 µM), or midgut

medium for 1 hour, after which they were video recorded for further analysis.

4.8. PrestoBlue™ (PB) assay

The metabolic activity of the dissected midguts was further assessed using a resazurin salt-

based cell viability reagent PrestoBlue™ (Invitrogen, USA). For these experiments, three

dissected midguts were placed per well in a 96-well tissue culture plate. Each trio of

midguts was considered a biological replicate. Using more than one midgut per well was

necessary to ensure both a robust readout and a relatively rapid change of color in the

testing medium. Midguts fixated in 4% paraformaldehyde (PFA, Sigma Aldrich, USA)

were included in the assay as negative controls (NC). No organ (NO) controls were

composed of wells where one tip of the forceps was dipped in during dissection.

The live midguts and the corresponding controls were tested with PB every day for a period

of 10 days. In brief, the medium was carefully removed from the wells and replaced with

100 µL of a 1:10 dilution of PB prepared in midgut medium (without phenol red). Next,

the midguts and their controls were incubated at 28°C without CO_2 for 7 hours. Once the

incubation time was completed, the PB-containing medium was transferred to a 96-well

tissue culture plate and fluorescence intensity was measured at wavelengths 560 nm

excitation and 590 nm emission with a Spark® Multimode Microplate Reader (Tecan

Trading AG, Switzerland). New midgut medium was added to wells containing midguts

and they were returned to the incubator (28°C, no CO_2) until the next time point

assessment. Two independent assays were carried out with at least 3 biological replicates

per time point.

4.9. Infection and antiviral assays in *ex vivo* mosquito midguts

Mosquito guts were dissected as described in 4.6 and placed into a 96-well tissue culture

plate filled with midgut medium. After removing the medium, a total of 1×10^4 PFU/mL

of the virus was added to each well. Of note, inocula of 1×10^5 PFU/mL and 1×10^6

PFU/mL were used for DENV-2 and USUV, respectively. Negative controls consisted of

dissected guts fixated in 4% PFA for 30 minutes. Midguts were incubated for two hours at

28°C, without CO_2. The virus inoculum was carefully removed, and midguts were washed

two times with midgut medium. Fresh medium was finally added to all wells for incubation

at 28°C, without CO_2.

To test the activity of antiviral drugs in the *ex vivo* mosquito midguts, compound dilutions

were prepared in midgut medium and added to the wells containing midguts to be treated,

after which they were infected with $1 \times 10^{4-5}$ PFU/mL of virus. Following two hours of

incubation at 28°C, without CO_2, the midguts were washed twice before adding midgut

medium alone (for virus control midguts) or containing compound (for treated midguts).

Midguts were returned to the incubator at 28°C, without CO_2.

Incubation time for alphavirus-infected midguts was 3 days, while flavivirus-infected

midguts were maintained for 7 days. Midguts were collected at several time points after

infection for further analysis.

4.10. Determination of viral RNA levels and detection of infectious virus

replication

Collected midguts were homogenized individually in 250 µL of PBS using bead disruption

(2.8 mm beads, Precellys). The midgut homogenate was filtered using 0.8 µm MINI

column filters (Sartorius, Germany) to remove debris, bacteria, and fungi. The filtered

homogenate was used for further viral RNA isolation and qRT-PCR to determine viral

RNA levels, and plaque assay to assess infectious virus particles.

Viral RNA isolation was performed with the NucleoSpin RNA Virus kit (Macherey-Nagel,

Germany) following the manufacturer's protocol. The sequences of primers and probes

used for each virus are compiled in Table 1. One-Step, quantitative RT-PCR was performed

for CHIKV and ZIKV in a total volume of 25 µL, consisting of 13.94 µL of RNase free

water (Promega, USA), 6.25 µL of master mix (Eurogentec, Belgium), 0.375 µL of each

forward and reverse primer (to a final concentration for each primer: 150 nM [CHIKV and

USUV]; 900 nM [ZIKV]), 1 µL of probe (to a final concentration of 400 nM [CHIKV and

USUV]; 200 nM [ZIKV]), 0.0625 µL of reverse transcriptase (Eurogentec, Belgium), and

3 µL of RNA sample. For RRV, the reaction mixture was prepared in a total volume of 20

µL, containing 5.2 µL of RNase free water, 10 µL of SYBR Green master mix (BioRad,

USA), 1 µL of each forward and reverse primer (to a final concentration of 125 nM for

each primer), 0.3 µL of reverse transcriptase (BioRad, USA), and 4 µL of RNA sample.

For DENV-2, the reaction mixture was prepared to a final volume of 20 µL, consisting of

3 µL of RNase free water, 10 µL of SYBR Green master mix (BioRad, USA), 0.3 µL of

each forward and reverse primer (to a final concentration of 900 nM for each primer), 0.4

µL of reverse transcriptase (BioRad, USA) and 6 µL of RNA sample.

The qRT-PCR assays were performed using the QuantStudio^{TM} 5 Real-Time PCR System

(ThermoFisher Scientific, USA) with the following cycling conditions for CHIKV, ZIKV,

and USUV: 30 min at 48°C, 10 min at 95°C, followed by 40 cycles of 15 s at 95°C, and 1

min at 60°C (55 °C for USUV). Cycling conditions for RRV were as follows: 10 min at

50°C, 3 min at 95°C, followed by 40 cycles of 15 s at 95°C, and 30 s at 60°C. For DENV-

2, the following cycling conditions were used: 30 min at 48°C, 3 min at 95°C, followed by

40 cycles of 15 s at 95°C, and 1 min at 62°C.

For absolute quantification, standard curves were generated each run using 10-fold serial

dilutions of: cDNA of CHIKV-nsP1, viral RNA isolated from RRV and DENV-2 Bangkok

virus stocks or synthesized gBlocks^{TM} gene fragments (Integrated DNA Technologies,

USA) for ZIKV and USUV.

Plaque assays were performed to quantify infectious virus particles from infected midgut

samples. Mammalian cells (Vero cells for RRV and CHIKV; BHK cells for ZIKV, DENV-

2, and USUV) were seeded at a density of 2.5×10^5 cells per well in 24-well tissue culture

plates. One-day confluent monolayers of cells were inoculated with serial 10-fold dilutions

of midgut samples, dilutions were prepared in 2% assay medium, and the inoculum was

allowed to infect the cells for 2 hours at 37°C, with 5% CO_2. A negative control consisting

of 2% assay medium used for preparing the dilutions was included in all plaque assays.

The inoculum was removed and replaced with CMC 0.8% overlay diluted in RPMI

medium (supplemented with 1% HEPES, 1% sodium bicarbonate, 1% L-glutamine, 1%

PenStrep and 2% FBS). Infected cells were incubated for 3 (*Alphaviruses*) or 7

(*Flaviviruses*) days. After incubation, 1 mL of 4% PFA was added to each well on top of

the overlay medium and allowed for fixation for 1 hour, after which the overlay mixture

was discarded, and the wells were carefully washed with water. The culture plates were

allowed to air dry, and then the wells were stained with 1% crystal violet solution (Sigma

Aldrich, USA). Plaques were counted and titer of each sample was calculated.

4.11.   Immunofluorescence staining of CHIKV viral proteins and imaging

Mosquito guts were dissected and infected with CHIKV as described in 4.6 and 4.9,

respectively. At day 3 p.i., midguts were fixated with 4% PFA for one hour at room

temperature. Fixated midguts were transferred to a μ-slide 18-well (Ibidi GmbH, Germany)

plate and rinsed with PBS five times, for 5 minutes each time. Midguts were then saturated

with PBS-T (0.1% Triton X-100 [Sigma Aldrich, USA] and 1% bovine serum albumin

[BSA, Sigma Aldrich, USA] in PBS 1X) for 2 hours at room temperature. Primary antibody

anti-E2 protein [Chk265] (Absolute Antibody, Wilton, UK) was diluted in PBS-T (1:500)

and added to the fixed midguts for overnight incubation at 4°C. After incubation, midguts

were rinsed five times with PBS-T, for 5 to 10 minutes each rinse, followed by incubation

with Alexa Fluor 594 donkey anti-rabbit secondary antibody (ThermoFisher Scientific,

USA, A-21207, diluted in PBS-T, 1:500) for 2 hours at room temperature. Midguts were

rinsed five times with PBS-T, for 5 to 10 minutes each rinse. Phalloidin (diluted in 1%

BSA in PBS 1X solution, final concentration 100 nm) incubation followed for 10 minutes,

after which DAPI (diluted in PBS 1X, final concentration 100 nm) was added on top of the

phalloidin solution and allowed to incubate for 10 minutes. Finally, the phalloidin/DAPI

solution was removed, and midguts were rinsed with PBS. Midgut samples were mounted

in microscope slides using Glycergel® mounting medium (Agilent, USA) and allowed to

dry before imaging with a Leica DMi8 microscope (Leica Microsystems, Germany) and

the Andor Dragonfly Confocal Spinning Disk microscope (Oxford Instruments, UK). All

incubation steps were carried out on rotation, at 300 rpm.

4.12.   **Infection of *ex vivo* cultured midguts with DENV-2 expressing**

mCherry and imaging

Mosquito guts were dissected and infected with a DENV-2 expressing mCherry (16) as

described in 4.6 and 4.9, respectively, using an inoculum of 1×10^5 PFU/mL. Negative

controls and mock-infected midguts were included. Mock-infected midguts were incubated

in midgut medium instead of virus dilution. Midguts were monitored every day during the

incubation period and checked under the FLoid™ Cell Imaging Station (Life

Technologies) for any signal that indicated mCherry expression. At day 7 p.i., the mosquito

midguts were fixated with 4% PFA for one hour at room temperature. After fixation,

midguts followed the protocol described in 4.11, skipping the addition of any primary and

secondary antibodies.

5. Acknowledgements

Special thanks to Prof. Johan Neyts (University of Leuven, Belgium) for allowing us to use

his laboratory space, equipment, and compounds for our experimental work. We are

grateful to Prof. Julia Dallman and Prof. Mason Klein (University of Miami, USA) for

kindly providing the custom IgorPro script used for gut motility analysis. Many thanks to

Prof. Sarah Merkling and Elodie Couderc (Institut Pasteur, Paris, France) for their helpful

advice regarding the immunostaining of mosquito midguts. We would also like to thank the European virus archive, EVAg, supported by the European Union's Horizon 2020 research and innovation program under grant agreement No. 871029, for supplying the ZIKV SL1602, Suriname strain. Thanks to Prof. Anna Bella Failloux (Institut Pasteur, Paris, France) for providing the DENV-2/ Bangkok strain. We thank Prof. Christian Drosten (University of Bonn, Germany) for providing the CHIKV 899 strain. We also thank Prof. Andrew Bristol (University of Bristol, UK) for providing the infectious clone DENV-2 pDVWS601, which served to construct the reporter DV2/mCherry used in this study. Many thanks to Prof. Pedro Elias Marques (University of Leuven, Belgium) for granting us the use of the Andor Dragonfly Confocal Spinning Disk microscope. Our sincere thanks to Prof. Sander Koenraadt (Wageningen University & Research, Netherlands) for kindly sharing *Cx. pipiens* mosquito egg rafts. Finally, we thank the Infravec2 Consortium for providing the *Ae. aegypti* mosquito eggs. This project was funded by KU Leuven (C22/18/007 and starting grant STG/19/008).

Supplementary material

Appendix I. Handling and optimization of *ex vivo* mosquito midguts.

Figure S1. *Ex vivo* cultured mosquito guts were not significantly affected by external incubation with serotonin or atropine.

Figure S2. Replication of DENV-2 expressing mCherry in the *ex vivo* cultured midguts.

Figure S3. ZIKV RNA loads in the mosquito midguts when cultured in medium or when using a foam substrate for support.

Figure S4. CHIKV replication kinetics in mosquito midguts cultured in medium and in carboxymethyl cellulose (CMC).

Movie 1. Representative mosquito hindgut videography used for the gut motility

analysis. Peristalsis is observed as waves along the hindgut region.

6. References

1.   Becker N, Petrić D, Zgomba M, Boase C, Madon M, Dahl C, Kaiser A. 2010.

Mosquitoes and Their Control, 2nd ed. Springer-Verlag Berlin Heidelberg.

2.   Patterson J, Sammon M, Garg M. 2016. Dengue, zika and chikungunya: Emerging

arboviruses in the new world. Western Journal of Emergency Medicine.

eScholarship https://doi.org/10.5811/westjem.2016.9.30904.

3.   Harrington LC, Fleisher A, Ruiz-Moreno D, Vermeylen F, Wa C v., Poulson RL,

Edman JD, Clark JM, Jones JW, Kitthawee S, Scott TW. 2014. Heterogeneous

Feeding Patterns of the Dengue Vector, Aedes aegypti, on Individual Human Hosts

in Rural Thailand. PLoS Negl Trop Dis 8:e3048.

4.   Franz A, Kantor A, Passarelli A, Clem R. 2015. Tissue Barriers to Arbovirus

Infection in Mosquitoes. Viruses 7:3741–3767.

5.   Vogels CB, Göertz GP, Pijlman GP, Koenraadt CJ. 2017. Vector competence of

European mosquitoes for West Nile virus. Emerg Microbes Infect 6:1–13.

6.   Walker T, Jeffries CL, Mansfield KL, Johnson N. 2014. Mosquito cell lines:

History, isolation, availability and application to assess the threat of arboviral

transmission in the United Kingdom. Parasit Vectors. BioMed Central Ltd.

https://doi.org/10.1186/1756-3305-7-382.

7.   Pena LJ, Guarines KM, Duarte Silva AJ, Sales Leal LR, Félix DM, Silva A, de

Oliveira SA, Junqueira Ayres CF, Silva JúNior A, de Freitas AC. 2018. In vitro and

in vivo models for studying Zika virus biology. Journal of General Virology

99:1529–1550.

8.   Grabowski JM, Tsetsarkin KA, Long D, Scott DP, Rosenke R, Schwan TG, Mlera

L, Offerdahl DK, Pletnev AG, Bloom ME. 2017. Flavivirus infection of ixodes

scapularis (Black-legged tick) Ex vivo organotypic cultures and applications for

disease control. mBio 8:1255–1272.

9.   Delang L, Yen P-S, Vallet T, Vazeille M, Vignuzzi M, Failloux A-B. 2018.
602   Differential Transmission of Antiviral Drug-Resistant Chikungunya Viruses by
603   Aedes Mosquitoes . mSphere 3:2020.

604   10.   Gloria-Soria A, Soghigian J, Kellner D, Powell JR. 2019. Genetic diversity of
605   laboratory strains and implications for research: The case of Aedes aegypti. PLoS
606   Negl Trop Dis 13.

607   11.   Grabowski JM, Offerdahl DK, Bloom ME. 2018. The Use of Ex Vivo Organ
608   Cultures in Tick-Borne Virus Research. ACS Infect Dis. American Chemical
609   Society https://doi.org/10.1021/acsinfecdis.7b00274.

610   12.   Hughes GL, Pike AD, Xue P, Rasgon JL. 2012. Invasion of Wolbachia into
611   Anopheles and Other Insect Germlines in an Ex vivo Organ Culture System. PLoS
612   One 7:e36277.

613   13.   Chung HN, Rodriguez SD, Carpenter VK, Vulcan J, Bailey CD, Nageswara-Rao M,
614   Li Y, Attardo GM, Hansen IA. 2017. Fat body organ culture system in aedes aegypti,
615   A vector of zika virus. Journal of Visualized Experiments 2017:55508.

616   14.   Eggleston H, Adelman ZN. 2020. Transcriptomic analyses of Aedes aegypti
617   cultured cells and ex vivo midguts in response to an excess or deficiency of heme: a
618   quest for transcriptionally-regulated heme transporters. BMC Genomics 21.

619   15.   Fros JJ, Miesen P, Vogels CB, Gaibani P, Sambri V, Martina BE, Koenraadt CJ, van
620   Rij RP, Vlak JM, Takken W, Pijlman GP. 2015. Comparative Usutu and West Nile
621   virus transmission potential by local Culex pipiens mosquitoes in north-western
622   Europe. One Health 1:31.

623   16.   Li LH, Kaptein SJF, Schmid MA, Zmurko J, Leyssen P, Neyts J, Dallmeier K. 2020.
624   A dengue type 2 reporter virus assay amenable to high-throughput screening.
625   Antiviral Res 183:104929.

626   17.   Pondeville E, Veronesi E, Kohl A. 2021. Guidelines for the design and operation of
627   containment level 2 and 3 insectaries.

628   18.   Gubler DJ. 1968. A method for the in vitro cultivation of ovarian and midgut cells
629   from the adult mosquito. Am J Epidemiol 87:502–508.

630   19.   James DM, Kozol RA, Kajiwara Y, Wahl AL, Storrs EC, Buxbaum JD, Klein M,
631   Moshiree B, Dallman JE. 2019. Intestinal dysmotility in a zebrafish (Danio rerio)
632   shank3a;shank3b mutant model of autism. Mol Autism 10:1–15.

633   20.   Luzak B, Siarkiewicz P, Boncler M. 2022. An evaluation of a new high-sensitivity
634   PrestoBlue assay for measuring cell viability and drug cytotoxicity using EA.hy926
635   endothelial cells. Toxicology in Vitro 83:105407.

636   21.   Xu M, McCanna DJ, Sivak JG. 2015. Use of the viability reagent PrestoBlue in
637   comparison with alamarBlue and MTT to assess the viability of human corneal
638   epithelial cells. J Pharmacol Toxicol Methods 71:1–7.

639   22.   Rettinger CL, Wang HC. 2018. Quantitative Assessment of Retina Explant Viability
640   in a Porcine Ex Vivo Neuroretina Model. https://home.liebertpub.com/jop 34:521–
641   530.

642   23.   Terrasso AP, Silva AC, Filipe A, Pedroso P, Ferreira AL, Alves PM, Brito C. 2017.
643   Human neuron-astrocyte 3D co-culture-based assay for evaluation of
644   neuroprotective compounds. J Pharmacol Toxicol Methods 83:72–79.

645   24.   Salazar MI, Richardson JH, Sánchez-Vargas I, Olson KE, Beaty BJ. 2007. Dengue
646   virus type 2: Replication and tropisms in orally infected Aedes aegypti mosquitoes.
647   BMC Microbiol 7:1–13.

648   25.   Choy MM, Sessions OM, Gubler DJ, Ooi EE. 2015. Production of Infectious
649   Dengue Virus in Aedes aegypti Is Dependent on the Ubiquitin Proteasome Pathway.
650   PLoS Negl Trop Dis 9.

651   26.   Ayers JB, Coatsworth HG, Kang S, Dinglasan RR, Zhou L. 2021. Clustered rapid
652   induction of apoptosis limits ZIKV and DENV-2 proliferation in the midguts of
653   Aedes aegypti. Communications Biology 2021 4:1 4:1–9.

654   27.   Okuda K, de Almeida F, Mortara RA, Krieger H, Marinotti O, Tania Bijovsky A.
655   2007. Cell death and regeneration in the midgut of the mosquito, Culex
656   quinquefasciatus. J Insect Physiol 53:1307–1315.

657   28.   Dong S, Kantor AM, Lin J, Passarelli AL, Clem RJ, Franz AWE. 2016. Infection
658   pattern and transmission potential of chikungunya virus in two New World
659   laboratory-adapted Aedes aegypti strains. Scientific Reports 2016 6:1 6:1–13.

29. Ehteshami M, Tao S, Zandi K, Hsiao HM, Jiang Y, Hammond E, Amblard F, Russell
OO, Merits A, Schinazi RF. 2017. Characterization of β-D-N4-hydroxycytidine as
a novel inhibitor of Chikungunya virus. Antimicrob Agents Chemother 61.

30. Urakova N, Kuznetsova V, Crossman DK, Sokratian A, Guthrie DB, Kolykhalov
AA, Lockwood MA, Natchus MG, Crowley MR, Painter GR, Frolova EI, Frolov I.
2017. D-N 4-Hydroxycytidine Is a Potent Anti-alphavirus Compound That Induces
a High Level of Mutations in the Viral Genome https://doi.org/10.1128/JVI.

31. Battisti V, Urban E, Langer T. 2021. Antivirals against the Chikungunya Virus.
Viruses 2021, Vol 13, Page 1307 13:1307.

32. Hucke FIL, Bugert JJ. 2020. Current and Promising Antivirals Against
Chikungunya Virus. Front Public Health 8:916.

33. Zmurko J, Marques RE, Schols D, Verbeken E, Kaptein SJF, Neyts J. 2016. The
Viral Polymerase Inhibitor 7-Deaza-2'-C-Methyladenosine Is a Potent Inhibitor of
In Vitro Zika Virus Replication and Delays Disease Progression in a Robust Mouse
Infection Model. PLoS Negl Trop Dis 10.

34. Vazeille-Falcoz M, Mousson L, Rodhain F, Chungue E, Failloux AB. 1999.
Variation in oral susceptibility to dengue type 2 virus of populations of Aedes
aegypti from the islands of Tahiti and Moorea, French Polynesia. American Journal
of Tropical Medicine and Hygiene 60:292–299.

35. Panning M, Grywna K, van Esbroeck M, Emmerich P, Drosten C. 2008.
Chikungunya fever in travelers returning to Europe from the Indian Ocean region,
2006. Emerg Infect Dis 14:416–422.

36. Dash PK, Agarwal A, Sharma S, Saha A, Joshi G, Gopalan N, Sukumaran D, Parida
MM. 2016. Development of a SYBR green I-based quantitative RT-PCR for Ross
River virus: Application in vector competence studies and antiviral drug evaluation.
J Virol Methods 234:107–114.

37. Delang L, Guerrero NS, Tas A, Quérat G, Pastorino B, Froeyen M, Dallmeier K,
Jochmans D, Herdewijn P, Bello F, Snijder EJ, de Lamballerie X, Martina B, Neyts
J, van Hemert MJ, Leyssen P. 2014. Mutations in the chikungunya virus non-
structural proteins cause resistance to favipiravir (T-705), a broad-spectrum
antiviral. Journal of Antimicrobial Chemotherapy 69:2770–2784.

38.   Jacobs S, Delang L, Verbeken E, Neyts J, Kaptein SJF. 2019. A Viral Polymerase

Inhibitor Reduces Zika Virus Replication in the Reproductive Organs of Male Mice.

Int J Mol Sci 20.

39.   Cavrini F, Pepa ME della, Gaibani P, Pierro AM, Rossini G, Landini MP, Sambri

V. 2011. A rapid and specific real-time RT-PCR assay to identify Usutu virus in

human plasma, serum, and cerebrospinal fluid. Journal of Clinical Virology 50:221–

223.

Figure 1. *Ex vivo* **cultured mosquito guts were viable for 7 days post dissection, based on gut peristalsis and a resazurin-based assay. a**, Representative image of a mosquito gut placed in a carboxymethyl cellulose (CMC) drop for videography. **b**, Close-up of the hindgut which comprises the area of analysis (AOA) manually set and outlined by the red lines. Cyan boxes denoted the regions to be analyzed along the AOA. The number and size of the regions (cyan boxes) were adjusted to be 25 boxes on each side of the hindgut. **c**, Representative kymograph showing the output for one mosquito gut analysis. Each line in the kymograph corresponded to an individual region, which was further investigated in a plot. In the figure, the red highlighted line in the kymograph correlates to the presented plot displaying a representative analysis readout for this region. **d**, Contractions observed per minute in function of time post dissection in days. Each point represents one individual mosquito midgut. The black line represents the median value. The graph shows data from three independent experiments. Statistical significance was assessed with a Kruskal-Wallis test. Significantly different values are indicated by asterisks: *, $P < 0.05$. **e**, Viability readouts exhibit metabolic activity for a period of 10 days for tested midguts, no organ and negative controls. Assays were performed with 2 or 3 biological replicates, each replicate composed of 3 midguts per well, along with its corresponding controls. Error bars represent the standard errors of the means per time point. The graph shows data from three independent assays.

Figure 2. *Ex vivo* **cultured midguts supported Ross River (RRV) and chikungunya virus (CHIKV) infection. a**, Viral RNA levels in the mosquito midguts were quantified at 2 hours (day 0), 1, 2 and 3 days p.i. by means of qRT-PCR. **b**, Infectious virus loads in the mosquito midguts were quantified by means of plaque assay. Each dot represents an individual midgut organ. The black line represents the median value. Data correspond to at least two independent replication kinetics assays. LOD: Limit of detection of the corresponding assay.

Figure 3. *Ex vivo* **cultured midguts supported dengue virus serotype 2 (DENV-2) and Zika virus (ZIKV) infection. a**, **c**, Viral RNA levels in the mosquito midguts were quantified at 2 hours, 5 and 7 days p.i. by means of qRT-PCR. **b, d**, Infectious virus loads in the mosquito midguts were quantified by means of plaque assay. Each symbol represents an individual midgut organ. The black line represents the median value. Data correspond to two independent replication kinetics assays. LOD: Limit of detection of the corresponding assay.

Figure 4. *Ex vivo* **cultured midguts were partially permissive to Usutu virus (USUV) infection. a**, Viral RNA levels in the mosquito midguts were quantified at 2 hours and 5 days p.i. by means of qRT-PCR. **b**, Infectious virus loads in the mosquito midguts were quantified by means of plaque assay. Each symbol represents an individual midgut organ. The black line represents the median value. LOD: Limit of detection.

Figure 5. **Detection of CHIKV protein synthesis in infected** *ex vivo* **midgut cultures. a**, Magnification, 25X. Overlay of blue, green, and magenta filter imaging are shown for two infected midgut organs and one negative control midgut. Imaging of each individual channel as follows: in blue, DAPI-stained cell nuclei; in green, actin filaments stained with phalloidin; and in magenta, E2 viral protein. No E2 expression was detected in the negative control midguts. The scale bar in panels, represented by the white line, corresponds to 400 µM. pmg: posterior midgut. hg: hindgut. **b**, Close-up panel showing infection foci in the posterior midgut region corresponding to the yellow squares indicated in **a** for "Midgut 1". Arrow heads indicate E2 expression localized in tracheal tubes of the midgut. **c**, Close-up panel showing infection foci in the posterior midgut and hindgut region corresponding to the yellow squares indicated in **a** for "Midgut 2".

Figure 6. **Replication of DENV-2 expressing mCherry in the *ex vivo* cultured midguts. a**, Magnification, 25X. Confocal microscopy displays the DENV-2 infection in the *ex vivo* midguts at day 7 p.i., as seen by the mCherry (red) signal. Overlay of blue, green, and red filter imaging are shown for two midgut organs and one negative control midgut. Imaging of each individual channel as follows: in blue, DAPI-stained cell nuclei; in green, actin filaments stained with phalloidin; and in red, mCherry signal. No mCherry expression was detected in fixated or mock-infected midguts. The scale bar in panels, represented by the white line, corresponds to 400 µM. pmg: posterior midgut. hg: hindgut. **b**, Close-up panel showing several infection foci in the posterior midgut region corresponding to the yellow squares indicated in **a** for "Midgut 1". **c**, Close-up panel showing infection foci in the posterior midgut and hindgut region corresponding to the yellow squares indicated in **a** for "Midgut 2".

Figure 7. **Antiviral activity of NHC and 7DMA in the *ex vivo* cultured mosquito midguts. a**, CHIKV RNA loads in the mosquito midguts were quantified at day 3 p.i. by qRT-PCR. **b**, CHIKV infectious virus loads were quantified at day 3 p.i. by plaque assay. Statistical significance was assessed with a Mann-Whitney test. Significantly different values are indicated by asterisks: *, $P<0.05$. **c**, ZIKV RNA levels in the mosquito midguts were quantified at day 7 p.i. by qRT-PCR. Statistical significance was assessed with a Kruskal-Wallis test. Significantly different values are indicated by asterisks: **, $P<0.05$. ns: not significant. Each dot represents an individual midgut. The black line represents the mean value. LOD: Limit of detection of the corresponding assay.

715 **Table 1. Sequences of primers and probes for qRT-PCR used in this study**

Virus [target protein]	Forward primer (5' → 3')	Reverse primer (5' → 3')	Probe (5' → 3')	Reference
Ross River virus [E2]	TACAAGCACGACCCATTGCCG	GATAGTCCTGCCGCCTGCTGT	N/A	(36)
Chikungunya virus [nsP1]	CCGACTCAACCATCCTGGAT	GGCAGACGCAGTGGTACTTCCT	FAM-TCCGACATCATCCTCCTTGCTGGC-TAMRA	(37)
Zika virus [E]	CCGCTGCCCAACACAAG	CCACTAACGTTCTTTTGCAGACAT	FAM-AGCCTACCT-ZEN-TGACAAGCAATCAGACACTCAA-IABkFQ	(38)
Dengue virus serotype 2 [NS3]	CAGATCGGAGCTGGAGTTTAC	TTTTGACGTCCGCCCATGAA	N/A	*In-house*
Usutu virus [NS5]	AAAAATGTACGCGGATGACACA	TTTGGCCTCGTTGTCAAGATC	FAM-TGGGACACCCGGATACCAGAG-TAMRA	(39)