Long-lasting impact of chito-oligosaccharide application on strigolactone biosynthesis and fungal accommodation promotes arbuscular mycorrhiza in *Medicago truncatula*

Veronica Volpe[1], Matteo Chialva[1], Teresa Mazzarella[1], Andrea Crosino[1], Lorenzo Costamagna[1], Wouter Kohlen[2], Andrea Genre[1*]

[1]Department of Life Science and Systems Biology, University of Turin, Viale Mattioli 25, 10125, Torino, Italy. [2]Laboratory of Molecular Biology, Wageningen University & Research, Wageningen 6708, PB, The Netherlands.

*Andrea Genre
Telephone/Email: +390116705083; andrea.genre@unito.it

This PDF file includes:
      Summary
      Main text: 5974 words
      Introduction: 468
      Materials and Methods: 2053
      Results: 2589
      Discussion: 864
      References
      Figures: 6 (1,2,3,4,6 in colour)
      Tables: 1

Summary

• The establishment of arbuscular mycorrhiza (AM) between plants and Glomeromycotina fungi is preceded by the exchange of chemical signals: fungal released Myc-factors, including chitoligosaccharides (CO) and lipo-chitooligosaccharides (LCO), activate plant symbiotic responses, while root exuded strigolactones stimulate hyphal branching and boost CO release.

• In this frame, the reinforcement of fungal signaling through CO application was shown to promote AM development in *Medicago truncatula*, but the cellular and molecular bases of this effect remained unclear.

• Here we demonstrate that CO treatment impacted the transcriptome of both mycorrhizal and non-mycorrhizal *M. truncatula* roots over several weeks, providing a novel insight into the mechanistic bases of the CO-dependent promotion of AM colonization.

• CO treatment caused the long-lasting regulation of strigolactone biosynthesis and fungal accommodation related genes. This was further validated by mass spectrometry analyses showing an increase in root didehydro-orobanchol content, and live cell imaging demonstrating the promotion of accommodation responses to AM fungi in root epidermal cells. Lastly, an advanced down-regulation of AM symbiosis marker genes was observed at our latest time point in CO-treated plants, in line with an increased number of senescent arbuscules.

• Overall, CO treatment triggered molecular, metabolic and cellular responses underpinning a protracted acceleration of AM development.

Keywords: Arbuscular mycorrhiza, Chito-oligosaccharides, *Medicago truncatula*, Strigolactones, Symbiosis

Introduction

Mineral nutrition of most plants is supported by the mutualistic root symbiosis with Glomeromycotina, an ancient group of soil fungi (Spatafora et al., 2016) that grant their host plants preferential access to soil inorganic nutrients, in change for plant-photosynthesized sugars and lipids (Smith and Read, 2008; Wewer et al., 2014; Keymer et al., 2017).

Plant-fungus recognition is essential for AM establishment and is based on an exchange of chemical signals (Bonfante and Requena, 2011; Zipfel and Oldroyd, 2017). On the one hand, root-exuded strigolactones (SL), a class of terpenoid lactones, signal host proximity (Akiyama et al., 2005) activating hyphal metabolism and branching, and eventually promoting physical encounter with the root surface (Besserer et al., 2006; Waters et al., 2017). On the other hand, AM fungi release diffusible molecules (Myc-factors) that activate a symbiotic signal transduction pathway (Zipfel and Oldroyd, 2017; Choi et al., 2018). Downstream responses include local and systemic changes in gene expression and metabolism, overall preparing the host plant to symbiosis establishment (MacLean et al., 2017; Choi et al., 2018; Pimprikar and Gutjarh, 2018).

Myc-factors include two classes of molecules: lipo-chito-oligosaccharides, or LCO (Maillet et al., 2011), structurally similar to rhizobial Nod-factors and composed of a short chitin chain with a few lateral substitutions; and chito-oligosaccharides, or CO (Genre et al., 2013), where only the chitin backbone is present. While LCO bioactivity has mostly been studied in legumes (Maillet et al, 2011; Sun et al., 2015; Bonhomme et al., 2021), CO are active in all tested AM host plants (Nasir et al., 2021) and their release is boosted upon strigolactone perception (Genre et al., 2013). Furthermore, CO can easily be produced through chitin hydrolysis (Crosino et al., 2021) making them particularly interesting for large scale agricultural applications (Volpe et al., 2020). In this context, we recently demonstrated that plant treatment with exogenous CO prior to fungal inoculation promotes AM colonization with a marked increase in arbuscule development, biomass accumulation and total photosynthetic surface, compared to untreated mycorrhizal plants (Volpe et al., 2020).

In order to understand the molecular bases of this CO-dependent promotion of AM development, we used RNA-seq to investigate genome-wide changes in root gene expression after CO treatment over four weeks, and further validated our results with functional insights. CO treatment changed the expression pattern of the whole strigolactone biosynthetic pathway and increased the didehydro-orobanchol content in root tissue. Moreover, an early, CO-dependent stimulation of intracellular accommodation processes consistently emerged from RNA-seq data, targeted gene expression analyses and live imaging of epidermal cell reorganization. Lastly, the down-regulation of AM symbiosis marker genes was consistent with the increased number of senescent arbuscules at the end of our experimental time frame. In conclusion, our results revealed that CO treatment

85 impacted on molecular, cellular and metabolic mechanisms that converged towards a global
86 acceleration in AM development.
87
88
89 **Materials and Methods**
90
Plant and fungal materials
The model legume *Medicago truncatula* cv 'Jemalong' (line A17) was used for this study. Seeds
were collected from the pods, oxygenated for 2 days at room temperature and scarified on
sandpaper in order to break the seed coat. Seeds were then sterilized using 5% (v/v) sodium
hypochlorite in sterile water and rinsed in sterile distilled water, each step for 5 min keeping the
falcon tubes under constant stirring. To break dormancy and allow rapid and synchronized
germination, seeds were placed on 0.6% Plant-Agar (Duchefa, Haarlem, The Netherlands) plates,
maintained at 4°C in the dark for two days and then moved at 23°C until germination.
A commercial inoculum of the AM fungus *Funneliformis mosseae* (strain BEG 12) from MycAgroLab
(www.mycagrolab.com; France) was used for all experiments.
101
102 **Chitin oligomers**
103 A mixture of short-chain chito-oligosaccharides (Volpe et al. 2020) obtained from crustacean food
104 manufacturing waste (Zhengzhou Sigma Chemical Co., Ltd. Zhengzhou, Henan, China) was used
105 for all plant treatments as a 1 g/L solution in sterile distilled water supplemented with Tween 20
106 (0.005%) as a surfactant. Sterile water with Tween 20 (0.005%) was used in control treatments.
107
Pot culture
Young *M. truncatula* seedlings were transferred into 9 cm x 9 cm x 12 cm pots filled with sterile
coarse sand (0,4-0,8 mm; Valle Po, Revello, CN, Italy), supplemented or not with 22.5 ml of fungal
inoculum. A plastic bag was placed on the pots during the first week, to protect the young plants
from desiccation. Plants were grown in phytochambers under controlled conditions (23°C day/21°C
night temperature, 16/8 h light/dark photoperiod) and fertilized once per week using a modified
Long Ashton nutrient solution (Hewitt, 1966) with a low phosphate content (3.2 µM). During the rest
of the week they were watered when necessary with tap water.
Four experimental conditions were set up: control (CTR) plants, lacking both CO treatment and AM
inoculation; CO-treated control (CTR+CO); mycorrhizal (MYC), where AM inoculum was added in
the absence of CO treatments; and CO-treated mycorrhizal (MYC+CO), where plants were both
inoculated and exposed to the CO solution. CO treatment and fungal inoculum were applied as
previously described in Volpe et al. (2020). In short, 5 ml of CO solution were sprayed over the pot
substrate surface 4 and 2 days before inoculation with *Funnelliformis mosseae.*

Plants were harvested at 10, 14, 21, 28 days post-inoculation (dpi) and carefully washed under tap water to remove sand. AM colonization was assessed based on the presence of root-bound extraradical mycelium prior to rapid sampling, dip-freezing in liquid nitrogen and storage at -80°C until RNA extraction. We used 5 plants for each experimental condition at each time point, except at 10 dpi, when the limited extension of root systems required the sampling of a larger number of plants (12). We collected at least three biological replicates (plants) for each condition, selecting the best developed plants, with no visible sign of pathogenesis or stress and an extensive extraradical fungal development.

RNA isolation and sequencing

Total RNA was extracted from all samples using the RNeasy™ Plant Mini kit (Qiagen, Hilden, Germany). Samples were mechanically homogenized using a TissueLyser system for 2 minutes at 18 Hz. An aliquot of each sample was mixed with RLT buffer (Qiagen), and further treated following the manufacturer's protocol. RNA quantity and quality were spectrophotometrically checked using a NanoDrop ND-1000 instrument. RNA samples were further quantified and tested for integrity by capillary electrophoresis using Agilent 2100 Bioanalyzer instrument with the Agilent RNA 6000 Nano Kit following manufacturer's instructions. As requested by the sequencing company (IGATech, Udine, Italy) samples dedicated to RNA-seq were not treated with DNase.

RNA-seq analysis on root RNA samples was performed at IGA Technology services. TruSeq stranded mRNA kit (Illumina, San Diego, CA) was used for library preparation and sequencing performed using Illumina NextSeq 500 platform (Illumina, San Diego, CA) in single-end mode at 75 bp read-length and with a sequencing depth ranging from 20 to 30 M of reads per sample (Table S1).

Bioinformatics

Adapter sequences were masked with Cutadapt v1.11 (Martin, 2011) from raw reads using the following parameters: --anywhere (on both adapter sequences) --overlap 5 -times 2 –minimum-length 35 -mask-adapter. Raw reads were then trimmed removing lower quality bases and adapters using ERNE software (Del Fabbro et al., 2013). The expression level of each gene in each library was calculated by mapping filtered reads on Mt4.1 reference genome (Young et al., 2011; Tang, et al., 2014) using STAR splice-aware aligner (Dobin et al., 2013) with default parameters. Reads overlapping with annotated exons were counted using HTSeq (Anders et al., 2015) and differential expression analysis performed for each comparison (CTR+CO *vs* CTR, MYC vs CTR, MYC+CO *vs* CTR and MYC+CO *vs* MYC) at each time points (10, 14, 21, 28 dpi) using DESeq2 R package (Love et al., 2014) at a False discovery rate (FDR or adjusted p-value) (Benjamini et al., 1995) threshold of 0.05 (SI_DATA1-4).

Differentially expressed genes were plotted into KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway maps using the 'pathview' R package (Luo et al., 2013) and the *M. truncatula*

annotation available at the KEGG website. Metabolic pathways visualization was achieved using MapMan software (Usadel et al., 2005). MapMan functional annotation for *M. truncatula* (Table S2) was obtained from the proteome using Mercator4 annotation tool (Schwacke et al., 2019).

Gene-ontology (GO) enrichment analyses were carried out on each of the previously generated contrasts using GOseq R package (Young et al., 2010) at a FDR cutoff of 0.1. The $-\log_{10}$ of the corrected p-value was plotted.

Variance Partitioning Analysis (VPA) was performed using the 'varpart' function in R package 'vegan' (Oksanen et al., 2019) using gene counts normalized with DESeq2 through the Variance Stabilizing Transformation (vst). Principal component analysis (PCA) was calculated with the 'prcomp' function in 'base' R package, using same normalization used for VPA. Statistical and graphical elaborations were performed in R programming environment (R Core Team 2020) using 'ggplot2' package (Wickham et al., 2022).

Gene expression analysis by real-time qPCR

To remove any trace of genomic DNA before cDNA synthesis, RNA samples were treated with TURBO™ DNase (QIAGEN, Hilden, Germany) according to the manufacturer's instructions, followed by second NanoDrop quantification. The RNA samples were routinely checked for DNA contamination by PCR analysis, using primers *MtTEF-F*: 5'-AAGCTAGGAGGTATTGAAAG- 3' and *MtTEF-R*: 5'- ACTGTGCAGTAGTACTTGGTG-3' for *MtTEF* (Elongation Factor, NCBI Ref. Seq.: XM_013595882).

For single-strand cDNA synthesis, approximately 700 ng of total RNA was denatured at 65°C for 5 min and then reverse-transcribed using the Super-Script II kit (Invitrogen, Carlsbad, CA, USA) at 25°C for 10 min, 42°C for 50 min and 70°C for 15 min. The final volume was 20 μl volume and contained 10 μM of random primers, 0.5 mM deoxynucloside triphosphates, 4 μl 5x buffer, 2 μl 0.1M dithiothreitol (DTT) and 1 μl Superscript II enzyme Control PCR was set with TEF primers to test the presence of cDNA.

RNA-seq data was validated by quantitative real-time PCR (qRT-PCR) analysis of the expression profile of 10 differentially-expressed genes (DEGs) across all treatments and time points (Fig. S1). Specific primers (listed in Table S3) were designed using PerlPrimer software (http://perlprimer.sourceforge.net) on *M. truncatula* CDS sequences from NCBI (http://www.ncbi.nlm.nih.gov/) and *M. truncatula* genome database (http://www.medicagogenome.org). All primers were first tested by PCR on genomic *M. truncatula* and *F. mosseae* DNA, to confirm their plant-specificity.

qRT-PCR was also used to investigate the regulation of marker genes for AM fungal accommodation and early signaling, using the primers listed in Table S4.

qRT-PCR analyses were done according to Rotor-Gene SYBR Green PCR Kit instructions (QIAGEN, Hilden, Germany) and were run in a final volume of 15 μl for each tube containing 7.5 μl of Rotor-Gene SYBR Green PCR Master Mix, 5.5 μl of 3 μM specific primers mix, and 10 ng of

cDNA. A Rotor Gene machine (Qiagen) was used with the following program: 3 min pre-incubation
at 95°C, followed by 40 cycles of 15s at 95°C, and 30s at 60°C. Each amplification was followed
by melting curve analysis (60–94°C) with a heating rate of 0.5°C every 15s. All reactions were
performed with two technical replicates and only Ct values with a standard deviation that did not
exceed 0.5 were considered. Relative RNA levels were calibrated using the elongation factor (TEF)
mRNA as endogenous reference and normalized to the control line. Results were validated
statistically using the unpaired Student's t-test to compare two means: differences were considered
significant at $P<0.05$.
206

207 Exudates collection and germination test with parasitic plants

CTR and CTR+CO plants were twice respectively treated with water or CO solution, and then grown
for 21 days. 21 day-old plants were carefully sampled and washed free of sand. They were put
inside a tube (2 plants for tube) with 40 mL of LA solution covering the whole root system and
closed with Parafilm; the tubes were placed into a closed Magenta box and the root apparatus was
covered with a black shield. Plants (4 biological replicates for each treatment) were placed in a
growth chamber for 36 h for exudate collection. After 36 hours, the root exudate was recovered
and stored at -20°C for the next step. Whole plant fresh weight, as well as shoot and root biomass
separately, were measured for each biological replicate, before storage at -20°C.
216

217 Extraction and analysis of strigolactones from root apparatus and exudates

CTR and CTR+CO plants were treated twice with water or CO solution respectively, and then grown
in quartz sand, as described above. After 21 days, plants were carefully sampled and washed free
of sand. Whole plant, shoot and root biomass were measured for each biological replicate before
storage at -20°C.
For SL analysis, each root was ground to a fine powder in a mortar using liquid nitrogen and kept
at -80°C before further use. 500 mg of each ground sample was transferred to 2 mL eppendorf
tubes and SL extracted with 2 mL of ethyl acetate with 10^{-8} M GR24 as internal standard (end
concentration 10^{-7} M). Tubes were vortexed, sonicated for 20 sec (sonication bath, Branson
Ultrasonic) and centrifuged for 10 min at 10,000 rpm at RT. Subsequently, the organic phases were
transferred to 4 mL glass vials and the solvent was evaporated in a speed vacuum system
(SPD121P, ThermoSavant, Hastings, UK). Root exudates were purified and concentrated as
previously described for tomato (Kohlen et al., 2012); SL from root tissues and exudates were
quantified according to Liu et al. (2011).
231

232 Local CO treatment of root organ cultures

In order to monitor the local occurrence of early gene regulation in response to CO perception, 1
g/L, 1 mg/L or 1 µg/L CO solutions were applied to *Agrobacterium rhizogenes*-transformed root
organ cultures (ROCs) expressing the membrane-bound fluorescent protein GFP::LjLTI6b (Suzaki

et al., 2012), which was used as a visual reporter of cell viability. ROC lines were propagated on M
medium (Bècard & Fortin, 1988) at 25°C in the dark, in vertically oriented petri dishes to favor the
regular fishbone-shaped root system (Chabaud et al., 2002) and grown for 3 weeks. For local CO
treatment, small discs of filter paper (Ø 4mm) soaked in 20 µl of CO solution were applied on
several lateral roots, 10-15 mm from the root tip, in agreement with Chabaud et al. (2011). Six hours
later, 1 cm-long root segments underlying each disc were excised, immediately frozen and stored
at -80°C until RNA extraction.
To further investigate whether CO treatment altered cell responses to fungal contact, we modified
the targeted AM inoculation technique used by Genre et al. (2005; 2008) for visualizing the pre-
penetration apparatus (PPA). In short, *Agrobacterium rhizogenes*-transformed root organ cultures
(ROCs) expressing the endoplasmic reticulum-targeted GFP-HDEL (Haseloff et al., 1997) were
grown on M medium (Boisson-Dernier et al., 2001) in vertically-oriented Petri dishes and inoculated
with pre-germinated spores of *Gigaspora margarita*. 1 ml of filtered CO solution (1mg/L) or sterilized
water (as control) was then added over the root culture before covering it with a gas-permeable
plastic film (bioFOLIE 25; Sartorius, Goettingen, Germany). Fungus-root interaction was monitored
daily using a stereomicroscope and contact sites were imaged at 7, 10 and 14 dpi using a Leica
TCS SP2 confocal microscope fitted with a long distance 40X water-immersion objective (HCX Apo
0.80). GFP fluorescence was excited with the argon laser band at 488 nm and recorded with an
emission window set at 500-525 nm.
255

Morphological and functional analysis of AM fungus
Mycorrhizal plants, treated or not with CO, were sampled at 28 dpi and were used to determine the
fresh weight of the entire plants and of the root apparatus. Biomass data were compared and the
results were validated statistically using the unpaired student's t-test used to compare two means:
differences were considered significant at $P<0,05$. For microscope analysis, root portion with fungal
external hyphae were selected and treated for 1 h in phosphate (Pi) buffer, pH 7, containing 3%
(w/v) paraformaldehyde. After washing in Pi buffer, roots were embedded in agarose (5%). Agarose
blocks were cut into 100 µm-vibratome slices, which were put on a slide. Slices were treated for 5
min in 0.5% commercial bleach, diluted in Pi buffer, washed again, and then incubated for overnight
with wheat germ agglutinin-fluorescein isothiocyanate (WGA-FITC) (Sigma-Aldrich, Milan, Italy), at
a final concentration of 10 µg/mL, to detect the chitin of fungal cell walls. For observation and
image acquisition we used a Leica TCS SP2 confocal microscope (Leica Microsystems GmbH,

Wetzlar, Germany) equipped with a 40X water immersion objective and we set up the excitation at
488 nm and recording the FITC fluorescence at 500-550 nm.
270
271
272
273 **Results**
274
CO treatment impacted on the root transcriptome for several weeks
In a first global analysis of the RNA-seq dataset, principal component analysis (PCA; Fig. 1A) and
variance partitioning analysis (VPA; Fig. 1B) revealed a high level of variation in the transcriptome
between conditions and time points. As shown in Figure 1A, CTR, CTR+CO, MYC and MYC+CO
transcriptomes clustered in different areas of the PCA plot, indicating that both AM inoculation and
CO treatment had a strong influence on the root gene expression profiles. At 10 dpi, CTR+CO,
MYC and MYC+CO samples were clearly separated from CTR along the PC1 axis (explaining
33.38% of variance), highlighting the distance in the transcriptional profile of CTR samples
compared to the remaining treatments. A major distinction along the PC1 axis became evident
since 14 dpi between inoculated (MYC and MYC+CO) and non-inoculated samples (CTR and
CTR+CO), as expected in relation with AM establishment. A specific effect of the CO treatment
was evident at 10 dpi (with a clear separation between CTR and CTR+CO, and a milder but evident
separation between MYC and MYC+CO), and partially rebounded at 28 dpi between MYC and
MYC+CO samples.
VPA analysis (Fig. 1B) highlighted time as the most influential variable in our time course study, in
line with the expected major changes in gene expression throughout AM development (Handa et
al., 2015). VPA was subsequently used to address individual time points: this correlated CO
treatment to the highest amount of variance in gene regulation at 10 dpi, an early time point in root
colonization in our experimental conditions, while AM inoculation became the most influential
variable in the later time points. Also in this case the CO effect extended across all four time points,
with a partial reinforcement at 28 dpi.
In conclusion, both PCA and VPA indicated an impact of CO treatment across our whole time
course analysis, with the strongest effect at the earliest time point in both inoculated and non-
inoculated plants.
The number and proportion of differentially expressed genes (DEGs) among the four conditions at
each of the four time points is summarized in Figure 1C. The MYC *vs* CTR comparison highlighted
a progressive increase in gene regulation, in line with the ongoing colonization of the root apparatus
by the AM fungus. In the CTR+CO *vs* CTR comparison, the CO effect was strongest at 10 dpi (with
2906 DEGs), intermediate at 14 and 21 dpi, and minimal at 28 dpi (with only 7 DEGs). This pattern
can be explained with the progressive degradation/elution of CO from the pot over the weeks that
follow the initial treatments (4 and 2 days before inoculation). The strongest differences in gene

regulation were observed when comparing MYC+CO *vs* CTR samples (which represent the most dissimilar conditions in our experiment). Here, the strongest impact of the CO treatment was recorded at 10 dpi (5526 DEGs), with a progressive decrease thereafter. This is suggestive of a synergistic effect of exogenous CO and fungal-secreted Myc-factors during early root colonization, and is in line with the progressive decrease of CO effect over time (as highlighted in CTR+CO *vs* CTR). Lastly, the MYC+CO *vs* MYC comparison aimed at dissecting the effect of exogenous CO on the AM interaction. Also in this case, a major impact was evident at 10 dpi (3709 DEGs), in line with the reinforcement of fungal signaling by the applied CO. Nevertheless, differential gene regulation peaked again at 28 days (1468 DEGs) with a large majority of down-regulated sequences, whereas intermediate time points displayed a very similar gene regulation scenario in the two conditions.

In summary, differential gene expression analysis convincingly supported the evidence emerging from multivariate analysis, indicating an impact of CO on plant gene regulation throughout our experimental time frame, with a major effect during early root colonization.

Functional enrichment analysis suggests that CO treatment mimics fungal presence

To gain insights into the molecular processes impacted by CO treatment, we performed a Gene Ontology (GO) enrichment analysis of DEGs (SI_DATA5-8; http://geneontology.org/docs/ontology-documentation/). We will focus here on the two most strongly regulated time points: 10 and 28 dpi, and the most over-represented GO terms for each comparison, as presented in Figure S2-S5. Interestingly, in the MYC *vs* CTR comparison (Fig. S2) *Defense responses* were over-represented both at 10 and 28 dpi, in agreement with the well documented regulation of defense-related genes during AM interaction (Salzer et al., 2000; Jung et al., 2012; Cameron et al., 2013). *Protein serine/threonine kinase activity* emerged at 10 and 28 dpi and *DNA-binding transcription factor activity* at 28 dpi, reflecting the extensive impact of symbiosis on regulatory pathways (Liu et al., 2003; Küster et al., 2004; Sanchez et al., 2004; Hohnjec et al., 2005; Hogekamp and Küster, 2013). The whole ubiquitination system (*ubiquitin ligase complex*, *protein ubiquitination*, *ubiquitin protein transferase activity*) was regulated at 28 dpi, possibly indicating the onset of senescence-related mechanisms. Lastly, *fatty acid biosynthesis* appeared at 28 dpi, likely related to both the extensive synthesis of perifungal membranes and the intense production of lipids feeding the fungus (Wewer et al., 2014; Luginbuehl et al., 2017; MacLean et al. 2017).

Since no enriched GO category emerged at 28 dpi for the CTR+CO *vs* CTR comparison (Fig. S3), we only comment here on the 10 dpi time point. In this case, a remarkable analogy emerged with the same time point of the MYC *vs* CTR comparison (Fig. S6): in fact, the global pattern of gene regulation was very similar for *kinase activity*, *signal transduction*, *defense response* and *acyl*

transferase activity. This supports the role of CO as mimics of fungal presence and elicitors of
symbiotic signaling and gene regulation in the host root.
In the MYC+CO *vs* CTR comparison at 10 dpi (Fig. S4), a massive activation was observed in
protein biosynthesis (*translation, ribosome, ribosome constituents*), cytoskeleton-associated
processes (*microtubule associated complex, microtubule-based movement and microtubule motor*
activity) and chromatin reorganization (*nucleosome, nucleosome assembly*). This is suggestive of
intense cell reorganization and is supported by our knowledge of the fungal accommodation
process that takes place inside each colonized cell (Gutjahr and Parniske, 2013; Carotenuto et al.,
2019). At 28 dpi (Fig. S4), GO terms related to transport (*transporter activity*, *transport*, *heme*
binding, *sulfate transport*, *transmembrane transport*, *secondary active sulfate transmembrane*
transport activity, *membrane*) were highly represented, likely related to the extensive AM
colonization promoted by CO treatment (Benedito et al., 2010; Gaude et al., 2012; Handa et al.,
2015).
Finally, the MYC+CO *vs* MYC comparison (Fig. S5) suggested that exogenous CO enhanced *cell*
cycle regulation, *chromatin rearrangement* and *cytoskeleton-associated processes* at 10 dpi,
hinting at an intensification of the fungal accommodation responses in CO-treated plants. By
contrast, a major impact on regulatory and proteolytic processes appeared at 28 dpi in CO-treated
plants, suggesting the appearance of senescence-related processes.
In conclusion, our analysis of GO enrichment indicated a global consistency between the regulatory
responses induced by AM fungi and CO treatment, with a reinforcement and an advancement of
symbiotic processes in plants that were exposed to both stimuli.
362
CO treatment stimulated strigolactone signaling and fungal accommodation
Among the various gene pathways related to AM establishment that were highlighted by GO
enrichment analysis (Table 1), we chose to validate the observed regulation of signaling- and cell
reorganization-related genes with functional analyses.
Firstly, we focused on the regulation of Strigolactone (SL) signalling. SL are carotenoid-derived
plant hormones (Gomez-Roldan et al., 2008; Umehara et al., 2008) that also function as
extraradical signals to activate AM fungi (Waters et al., 2017). SL biosynthesis (*CCD8-1* and
CCD8-2) and transport (*PDR1a*) marker genes were down-regulated at 28 dpi in MYC compared
to CTR plants (Fig. 2A), which is in line with the inhibition of rhizospheric SL signaling upon
maximum root colonization (Lopez-Raez et al., 2011). Remarkably, this down-regulation of SL
biosynthesis and transport occurred as early as 10-14 dpi in CO-treated plants (MYC+CO *vs* CTR).
A rather different scenario was recorded in CO-treated plants that were not inoculated with AMF
(CTR+CO *vs* CTR, SI_DATA1). Under these conditions, SL biosynthesis and transport was found
to be up-regulated between 14 and 21 dpi. Such an up-regulation of SL biosynthesis suggests a
CO-dependent boosting of fungus-directed signaling in the absence of an effective interaction with
the symbiont.

To correlate the observed regulation of SL related gene expression to SL metabolomics, we
analyzed SL content in root tissues and exudates from 21-day-old CTR and CTR+CO plants. In our
samples only one SL, didehydro-orobanchol (DDO), was above the detection level. This SL has
been identified in roots of several AM host plants (Lopez-Raez et al., 2008; Yoneyama et al., 2008;
Kohlen et al., 2012), and characterized as the major strigolactone in *M. truncatula*, with a role in
stimulating hyphal branching in the AM fungus *Gigaspora margarita* (Liu et al., 2011, Tokunaga et
al., 2015). Interestingly, CTR+CO roots contained a significantly higher level of DDO compared to
CTR roots and a comparable trend (albeit statistically not significant) was also observed in root
exudates (Fig. 2B).
Secondly, MYC+CO *vs* CTR and MYC+CO *vs* MYC comparisons (Table 1; SI_DATA4, 5)
highlighted the up-regulation of a consistent group of functional categories related to cell
remodelling, such as *Cytoskeleton Organization*, and *Cell Wall Organization*. This is in line with the
activation of intracellular fungal accommodation by the host tissues (Luginbuehl and Oldroyd, 2017;
Pimprikar and Gutjarh, 2018), requiring the reorganization of cytoskeletal elements (Genre et al.,
2005), the onset of massive exocytic processes (Genre et al., 2012), cell wall material deposition
and remodelling (Balestrini and Bonfante, 2014) to generate the symbiotic interface (Balestrini et
al., 2005). Furthermore, a significant up-regulation was also observed in *Cell Cycle Organization*
for numerous cell cycle regulators (*CYCA*, *CYCB*, *alpha-AURORA kinase activator*) and cell plate-
associated proteins (*KNOLLE*), in strong agreement with the described activation of cell division
(Russo et al., 2018) and endoreduplication (Carotenuto et al., 2019) as part of the fungal
accommodation response.
In the light of this data, and because fungal accommodation responses such as prepenetration
apparatus (PPA) development are known to occur in the range of a few hours (Genre et al., 2005),
we decided to further investigate earlier responses to CO, by applying local, 6 hour long treatments
(Fig. 3) and analyzing the regulation of *MtKNOLLE* (Richter, et al., 2014; Russo et al., 2019) and
two additional cell cycle-related markers known to be expressed in early AM development, *MtAPC2*,
the *alpha subunit of the Adaptor Protein complex2* (Van Damme et al., 2011; Russo et al., 2018)
and *MtCYCL3*, a cyclin-like F-box protein (Russo et al, 2019). Two additional early symbiotic
markers were also analyzed: *MtPUB1* (*Plant U-box protein1*), an E3 ubiquitin ligase (Vernié et al.,
2016) and *MtCBF3*, a CAAT box-binding transcription factor (Hogekamp et al., 2011).
As shown in Figure 3, each gene was upregulated upon either 1 g/L, 1 mg/L or 1 µg/L CO treatment,
with 1 mg/L CO solution resulting to significantly up-regulate *MtPUB1*, *MtCBF3*, *MtVapyrin*,
MtKNOLLE and *MtAPC2*, whereas *MtCYCL3* was only overexpressed in response to 1g/L
treatment.
In conclusion, our targeted investigation of gene expression confirmed the CO-dependent
activation of several early AM markers and revealed that a 6h CO application is sufficient to activate
a set of genes related to prepenetration responses. Together, this gene expression pattern
corroborated our RNA-seq data and was fully compatible with the observed acceleration of AM

development in CO-treated plants, providing molecular evidence in favor of a CO-dependent
advance of pre-penetration responses in root cells.

CO treatment stimulated pre-penetration responses upon AM inoculation
We then decided to investigate the effect of CO treatment on prepenetration responses in the
presence of an AM fungal inoculum. To this aim, we grew *M. truncatula* root organ cultures (ROCs)
expressing GFP-HDEL in the presence of 1 mg/L CO solution (the most active concentration in our
gene expression analyses) or sterile water as a control. GFP-HDEL is a fluorescent marker for the
endoplasmic reticulum (ER) that has previously been used to track cytoplasmic aggregations and
PPA development during AM colonization (Genre et al., 2005; 2008; 2012). The asynchronous
progression of hyphal development and root colonization limits our ability to fully control the timing
of fungus-plant contact even using the so-called targeted inoculation method that we here adapted
(Chabaud et al., 2002). For this reason, observations were done at 7, 10 or 14 dpi with pre-
germinated *G. margarita* spores, a time frame that corresponds to hyphopodium formation and
initial epidermis colonization. As expected, broad ER aggregations were observed at all time points,
extending between the hyphopodium contact site and the epidermal cell nucleus, as previously
described during PPA formation (Genre et al., 2005). Such aggregations were observed both in the
presence or absence of the CO treatment. Nevertheless, such PPA-related ER aggregations
appeared to be more frequent in CO-treated roots, as confirmed by quantitative analysis (Fig. 4).
Indeed, the percent of cells showing ER aggregations in hyphopodium-contacted areas was
significantly higher in treated than untreated roots both at 7 and 10 dpi, while the two data were
comparable at 14 dpi. The major impact of CO treatment on earlier time points can be explained
with the progressive decrease in residual CO concentration and/or the inhibition of new
hyphopodium formation as arbuscules start developing in the inner root tissues. In this scenario,
the observed significant stimulation of prepenetration responses in early time points provides a
convincing functional validation of our gene expression data and the first cellular basis for the
observed CO-dependent promotion of AM colonization (Volpe et al., 2020).

Long term effects of the CO treatment
The 28 dpi time point was associated with a significant increase in plant development for MYC+CO
compared to MYC plants (Fig. S7). According to our previous studies (Volpe et al., 2020), this is
also the time of maximum AM development in our pot cultured *M. truncatula*. Nevertheless, when
comparing MYC+CO to MYC root transcriptome, we observed a surprising decline in the
expression of several AM markers, including transcription factors (*MtMYB* and *MtRAM2*) and
arbuscule-specific phosphate (*MtPT4*) and ammonium (*MtAMT1*) transporters (Fig. 5A;
SI_DATA4). We suspected that this could indicate the inception of symbiosis senescence, possibly
related to space limitations to root development in our pot cultures: in fact, the root apparatus of

MYC+CO plants had extended to the whole substrate volume at 28 dpi, which was not the case for
MYC plants.
We therefore used confocal microscopy to image and compare arbuscule morphology in MYC and
MYC+CO root samples at 28 dpi. Following image acquisition, individual arbuscules were classified
into four developmental stages based on their morphological features to generate a quantitative
comparison (Fig. 5B): Stage I (developing arbuscules with a limited number of large branches);
Stage II (intermediate maturity, with several fine branching occupying part of the host cell lumen);
Stage III (mature arbuscules, with fine branches in most of the host cell volume); Stage IV
(senescent arbuscules displaying clusters of collapsed branches). As shown in the histogram of
Figure 5B, the arbuscule population in MYC+CO plants had a maximum frequency in Stage III,
whereas the highest frequency was observed in Stage II for MYC plants. Furthermore, the direct
comparison of each stage of arbuscule development revealed that Stage I arbuscules were only
detected in MYC plants, where Stage II arbuscules were also significantly more abundant. By
contrast, a significant increase in Stage III and Stage IV arbuscules was recorded in MYC+CO
plants. We interpret this developmental shift as the consequence of the CO-induced advancement
in fungal accommodation responses - as indicated by our molecular analyses and further
demonstrated in epidermal cells by live cell imaging - culminating in a significant advance in
arbuscule development and senescence in CO-treated plants.
472
473
474 **Discussion**
475
The main aim of this investigation was to test the hypothesis that CO stimulate symbiotic responses
in the host root throughout AM development. In fact, we recently demonstrated that early CO
treatment promotes root infection by AM fungi over 28-48 days (Volpe et al., 2020), but a major
gap was left in our understanding of the impact of CO treatment on root transcriptome, metabolism
and cellular responses throughout AM colonization.
Indeed, several studies have investigated early plant responses to exogenous treatment with AM
fungal raw exudates or purified LCO and CO, revealing the rapid (< 1h) activation of plant symbiotic
signaling processes (Maillet et al., 2011; Genre et al., 2013) and short-term (1-48 hours)
transcriptional reprogramming (Czaja et al., 2012; Giovannetti et al., 2015; Feng et al., 2019). All
of these investigations were anyway analyzing plant responses occurring in the absence of any
fungal inoculation.
By analyzing the root transcriptome alongside symbiosis development in a period of over four
weeks following initial CO treatment, the present study breaks through this limitation. The
consolidation of our gene expression analysis with functional insights generated a first, consistent
outlook on CO-dependent plant responses. At any rate, our large transcriptomic dataset, combining

four different conditions and four time points, remains available (link in Data Availability Statement)
to drive further investigations.
493
Exogenous CO stimulate plant symbiotic responses throughout AM development
Our present results demonstrate that, besides the well-characterized activation of early symbiotic
signaling and gene regulation, CO also influenced key symbiotic features of the host tissues from
as soon as 6 hours after treatment to 28 dpi. Altogether, CO impacted on root SL metabolism,
transport and composition; repressed defense responses (pathogenesis-related genes and actors
of effector-triggered immunity) and secondary metabolism (isoflavonoid and terpenoid synthesis),
and increased protein biosynthesis (lipid and mineral nutrient transport), globally reinforcing the
host response to the AM fungus and prompting a symbiosis-oriented metabolic and physiological
context. Furthermore, CO also promoted cell-cycle reactivation and fungus accommodation-related
reorganization of the host cell cytoplasm, and caused an advance in the root colonization process
that extended to 28 dpi.
In short, our experimental results indicate a general and long-lasting stimulation of symbiotic
responses in both inoculated and uninoculated CO-treated plants, demonstrating that CO have all
the predicted features of Myc-factors, intended as fungal-secreted molecules that prepare the host
plant to symbiosis establishment (Gutjahr & Parniske 2013) and actively promote it.
509
The strange case of an unattainable love
Root exposure to fungal signaling molecules in the absence of the fungus (our CTR+CO condition)
represents a markedly artificial situation. In nature, as AM fungal hyphae approach a host root,
Myc-factor perception takes place alongside (or shortly before) additional, largely uncharacterized
signals (Bonfante and Requena, 2011), including the physical contact between hyphae and root
epidermal cells, and the subsequent development of intraradical colonization (Choi et al., 2018).
The progression of AM development has an impact on root responses, by transiently triggering
defense reactions (Garcia-Garrido and Ocampo, 2002; Jung et al., 2012; Cameron et al., 2013),
progressively reducing SL synthesis and secretion (Lopez-Raez et al., 2011; this paper) and overall
outlining the context of a successful symbiotic interaction. By contrast, our CTR+CO plants lack
this second set of fungal stimuli, which are crucial modulators of the host response. When
interpreting the gene regulation pattern in these root samples, the presence of a certain degree of
overreaction should therefore be taken into account. With this caveat, the CTR+CO condition
provided the experimental tool to dissect otherwise unseen CO effects. A remarkable example of
this artificial amplification of plant responses is the boosting of SL signaling, which extended for a
few weeks after CO treatment in CTR+CO plants. We interpret this as a climax in plant presymbiotic
signaling, in a condition where physical plant-fungus interaction is not going to occur. While the
biological significance of this observation can be limited, with reference to a natural context, it

convincingly indicates that CO effects extend over several weeks and impact precisely on SL
content, a root feature that is directly linked to AM symbiosis establishment.
530
Closing the loop
Plants exude diverse mixtures of SL, depending on plant species, growth conditions and
developmental stages (Wang and Bouwmeester, 2018). With reference to AM signaling, DDO has
been characterized as the most active molecule in *M. truncatula* (Lopez-Raez et al., 2008;
Yoneyama et al., 2008; Kohlen et al., 2012). In this frame, our novel observation of SL metabolism
promotion upon CO treatment is further reinforced by the specific accumulation of DDO, suggesting
a redirection of SL metabolism towards AM fungus-directed signals in CO-treated roots.
Furthermore, previous investigations have recorded an increase of CO release by SL-treated AM
fungi (Genre et al., 2013). By demonstrating that exogenous CO treatment promotes AM-specific
SL production, our present data close the loop, suggesting the existence of a positive feedback
mechanism that reinforces reciprocal plant-fungus signaling in support of symbiosis establishment.
To summarize our conclusions, we propose a model - schematized in Figure 6 - where the
exogenous application of CO does not only boost signal exchange in this symbiotic signaling loop
by promoting SL production, but also stimulates intracellular accommodation responses in the host
root, facilitating fungal colonization and eventually accelerating symbiosis development.
546
547

548 **Acknowledgments**
549 We are grateful to David Barker and Paola Bonfante for constructive discussion and their critical
550 revision of the manuscript draft. This research was funded by Fondazione Cassa di Risparmio di
551 Cuneo (Bando Ricerca Scientifica 2015 – Project AMforQuality 2015/17271). We are grateful to
552 Erik Limpens for hosting LC in his lab in Wageningen for an Erasmus+ traineeship.

553
554

555 **Author Contributions**
556 VV, TM performed experiments and data analysis. MC performed data analysis and contributed
557 figure editing and manuscript writing. AC, LC and WK performed experiments. VV, AG conceived
558 experiments and wrote the manuscript.

559
560
561 **Data Availability**
562 Raw RNA-seq data were deposited in the Sequence Read Archive (SRA, NCBI) under accession
563 PRJNA813377.
564
565

References

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**:166-9.

Akiyama K, Matsuzaki K, Hayashi H. 2005. Plant sesquiterpenes induce hyphal branching in
arbuscular mycorrhizal fungi. *Nature* **43**: 824–827.

Balestrini R, Cosgrove DJ, Bonfante P. 2005. Differential location of α-expansin proteins during
the accommodation of root cells to an arbuscular mycorrhizal fungus. *Planta* **220**: 889–899.

Balestrini R, Bonfante P. 2014. Cell wall remodeling in mycorrhizal symbiosis: a way towards
biotrophism. *Frontiers in Plant Science* **5**: 237.

Bécard G, Fortin JA. 1988. Early events of vesicular-arbuscular mycorrhiza formation on Ri T-
DNA transformed roots. *New Phytologist* **108**: 211-218.

Benedito VA, Li H, Dai X, Wandrey M, He J, Kaundal R, Torres-Jerez I, Gomez SK, Harrison
MJ, Tang Y et al. 2010. Genomic inventory and transcriptional analysis of *Medicago truncatula*
transporters. *Plant Physiology* **152**:1716-1730.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *J R Statist Soc B.* **57**:289–300.

Besserer A, Puech-Pagès V, Kiefer P, Gomez-Roldan V, Jauneau A, Roy S, Portais JC, Roux
C, Bécard G, Séjalon-Delmas N. 2006. Strigolactones stimulate arbuscular mycorrhizal fungi by
activating mitochondria. *PLoS Biology* **4**:e226.

Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker DG. 2001.
Agrobacterium rhizogenes-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing
and endomycorrhizal symbiotic associations. *Molecular Plant Microbe Interactions* 14:695-700.

Bonfante P, Requena N. 2011. Dating in the dark: how roots respond to fungal signals to establish
arbuscular mycorrhizal symbiosis. *Current Opinion in Plant Biology* **14**: 451-457.

Bonhomme M, Bensmihen S, André O, Amblard E, Garcia M, Maillet F, Puech-Pagès V,
Gough C, Fort S, Cottaz S et al. 2021. Distinct genetic basis for root responses to lipo-

chitooligosaccharide signal molecules from different microbial origins. *Journal of Experimental*

Botany. **72**: 3821-3834.

Cameron DD, Neal AL, van Wees SC, Ton J. 2013. Mycorrhiza-induced resistance: more than

the sum of its parts. *Trends in Plant Science* **18**:539–545.

Carotenuto G, Volpe V, Russo G, Politi M, Sciascia I, de Almeida-Engler J, Genre A. 2019.

Local endoreduplication as a feature of intracellular fungal accommodation in arbuscular

mycorrhizas. *New Phytologist* **223**: 430-446.

Chabaud M, Venard C, Defaux-Petras A, Bécard G, Barker DG. 2002. Targeted inoculation of

Medicago truncatula in vitro root cultures reveals *MtENOD11* expression during early stages of

infection by arbuscular mycorrhizal fungi. *New Phytologist* **156**: 265-273.

Chabaud M, Genre A, Sieberer BJ, Faccio A, Fournier J, Novero M, Barker DG, Bonfante P.

2011. Arbuscular mycorrhizal hyphopodia and germinated spore exudates trigger Ca2+ spiking in

the legume and nonlegume root epidermis. *New Phytologist* **189**: 347-355.

Choi J, Summers W, Paszkowski U. 2018. Mechanisms underlying establishment of arbuscular

mycorrhizal symbioses. *Annual Review of Phytopathology* **56**: 135–160.

Crosino A, Moscato E, Blangetti M, Carotenuto G, Spina F, Bordignon S, Puech-Pagès V,

Anfossi L, Volpe V, Prandi C et al. 2021. Extraction of short chain chitooligosaccharides from

fungal biomass and their use as promoters of arbuscular mycorrhizal symbiosis. *Scientific Reports*

11: 3798.

Czaja LF, Hogekamp C, Lamm P, Maillet F, Martinez EA, Samain E, Dénarié J, Küster H,

Hohnjec N. 2012. Transcriptional responses toward diffusible signals from symbiotic microbes

reveal *MtNFP-* and *MtDMI3*-dependent reprogramming of host gene expression by arbuscular

mycorrhizal fungal lipochitooligosaccharides. *Plant Physiology* **159**:1671-85.

Del Fabbro C, Scalabrin S, Morgante M, Giorgi FM. 2013. An extensive evaluation of read

trimming effects on Illumina NGS data analysis. *PLoS One* **8**: e85024.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15-21.

Feng F, Sun J, Radhakrishnan GV, Lee T, Bozsóki Z, Fort S, Gavrin A, Gysel K, Thygesen
MB, Andersen KR et al. 2019. A combination of chitooligosaccharide and lipochitooligosaccharide
recognition promotes arbuscular mycorrhizal associations in *Medicago truncatula*. *Nature*
Communications **10**: 5047.

García- Garrido JM, Ocampo JA. 2002. Regulation of the plant defence response in arbuscular
mycorrhizal symbiosis. *Journal of Experimental Botany* **53**: 1377-1386.

Gaude N, Bortfeld S, Duensing N, Lohse M, Krajinski F. 2012. Arbuscule-containing and non-
colonized cortical cells of mycorrhizal roots undergo a massive and specific reprogramming during
arbuscular mycorrhizal development. *Plant Journal* **69**: 510–528.

Genre A, Chabaud M, Timmers T, Bonfante P, Barker DG. 2005. Arbuscular mycorrhizal fungi
elicit a novel intracellular apparatus in *Medicago truncatula* root epidermal cells before infection.
Plant Cell **17**: 3489–3499.

Genre A, Chabaud M, Faccio A, Barker DG, Bonfante P. 2008. Prepenetration apparatus
assembly precedes and predicts the colonization patterns of arbuscular mycorrhizal fungi within
the root cortex of both *Medicago truncatula* and *Daucus carota*. *Plant Cell* **20**:1407–1420.

Genre A, Chabaud M, Balzergue C, Puech-Pagès V, Novero M, Rey T, Fournier J, Rochange
S, Bécard G, Bonfante P et al. 2013. Short-chain chitin oligomers from arbuscular mycorrhizal
fungi trigger nuclear Ca2+spiking in *Medicago truncatula* roots and their production is enhanced by
strigolactone. *New Phytologist* **19**:190-202.

Genre A, Ivanov S, Fendrych M, Faccio A, Zársky V, Bisseling T, Bonfante P. 2012. Multiple
exocytotic markers accumulate at the sites of perifungal membrane biogenesis in arbuscular
mycorrhizas. *Plant and Cell Physiology* **53**: 244–255.

Giovannetti M, Mari A, Novero M, Bonfante P. 2015. Early *Lotus japonicus* root transcriptomic
responses to symbiotic and pathogenic fungal exudates. *Frontiers in Plant Science* **6**: 480.

Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pagès V, Dun EA, Pillot JP, Letisse F,
Matusova R, Danoun S, Portais JC et al. 2008. Strigolactone inhibition of shoot branching. *Nature*
455: 189-194.

Gutjahr C, Parniske M. 2013. Cell and developmental biology of arbuscular mycorrhiza symbiosis.
Annual Review of Cellular and Developmental Biology **29**: 593-617.

Handa Y, Nishide H, Takeda N, Suzuki Y, Kawaguchi M, Saito K. 2015. RNA-seq Transcriptional Profiling of an Arbuscular Mycorrhiza Provides Insights into Regulated and Coordinated Gene Expression in *Lotus japonicus* and *Rhizophagus irregularis*, *Plant and Cell Physiology* **56**: 1490-1511.

Haseloff J, Siemering KR, Prasher DC, Hodge S. 1997. Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic Arabidopsis plants brightly. *Proceedings of the National Academy of Science USA* **94**: 2122–2127.

Sand and Water Culture Methods Used in the Study of Plant Nutrition. Technical Communication No. 22. Commonwealth Bureau, London

Hewitt EJ. 1966. Sand and Water Culture Methods Used in the Study of Plant Nutrition. *Experimental Agriculture* **3**: 104-104.

Hogekamp C, Arndt D, Pereira PA, Becker JD, Hohnjec N, Küster H. 2011. Laser microdissection unravels cell-type-specific transcription in arbuscular mycorrhizal roots, including CAAT-box transcription factor gene expression correlating with fungal contact and spread. *Plant Physiology* **157**: 2023-2043.

Hogekamp C, Küster H. 2013. A roadmap of cell-type specific gene expression during sequential stages of the arbuscular mycorrhiza symbiosis. *BMC genomics* **14**: 306.

Hohnjec N, Vieweg MF, Pühler A, Becker A, Küster H. 2005. Overlaps in the transcriptional profiles of *Medicago truncatula* roots inoculated with two different *Glomus* fungi provide insights into the genetic program activated during arbuscular mycorrhiza. *Plant Physiology* **137**:1283–1301.

Keymer A, Pimprikar P, Wewer V, Huber C, Brands M, Bucerius SL, Delaux PM, Klingl V, Röpenack-Lahaye EV, Wang TL et al. 2017. Lipid transfer from plants to arbuscular mycorrhiza fungi. *Elife* **6**: e29107.

Kohlen W, Charnikhova T, Lammers M, Pollina T, Tóth P, Haider I, Pozo MJ, de Maagd RA, Ruyter-Spira C, Bouwmeester HJ et al. 2012. The tomato CAROTENOID CLEAVAGE DIOXYGENASE8 (SlCCD8) regulates rhizosphere signaling, plant architecture and affects reproductive development through strigolactone biosynthesis. *New Phytologist* **196**: 535-547.

715 **Küster H, Hohnjec N, Krajinski F, El Yahyaoui F, Manthey K, Gouzy J,Dondrup M, Meyer F,**
716 **Kalinowski J, Brechenmacher L, et al. 2004.** Construction and validation of cDNA-based Mt6k-
717 RIT macro- and microarrays to explore root endosymbioses in the model legume *Medicago*
718 *truncatula. Journal of Biotechnology* **108**: 95–113.
719
720 **Jung SC, Martinez-Medina A, Lopez-Raez JA, Pozo MJ. 2012.** Mycorrhiza-induced resistance
721 and priming of plant defenses. *Journal of Chemical Ecology* **38:** 651–664.
722
723 **Liu J, Blaylock LA, Endre G, Cho J, Town CD, VandenBosch KA, Harrison MJ. 2003.**
724 Transcript profiling coupled with spatial expression analyses reveals genes involved in distinct
725 developmental stages of an arbuscular mycorrhizal symbiosis. *Plant Cell* **15**: 2106–2123
726
727 **Liu W, Kohlen W, Lillo A, Op den Camp R, Ivanov S, Hartog M, Limpens E, Jamil M,**
728 **Smaczniak C, Kaufmann K. 2011.** Strigolactone biosynthesis in *Medicago truncatula* and rice
729 requires the symbiotic GRAS-type transcription factors NSP1 and NSP2. *Plant Cell* **23**: 3853-3865.
730
731 **López-Ráez JA, Charnikhova T, Fernández I, Bouwmeester H, Pozo MJ. 2011.** Arbuscular
732 mycorrhizal symbiosis decreases strigolactone production in tomato. *Journal of Plant Physiology*
733 **168**: 294-297.
734
735 **López-Ráez JA, Charnikhova T, Mulder P, Kohlen W, Bino R, Levin I, Bouwmeester H. 2008.**
736 Susceptibility of the tomato mutant high pigment-2dg (hp-2dg) to *Orobanche spp.* infection. *Journal*
737 *of Agricultural and Food Chemistry* **56**: 6326–6332.
738
739 **Love MI, Huber W, Anders S. 2014.** Moderated estimation of fold change and dispersion for RNA-
740 seq data with DESeq2. *Genome Biology* **15**: 550.
741
742 **Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A, Oldroyd**
743 **GED, Eastmond PJ. 2017.** Fatty acids in arbuscular mycorrhizal fungi are synthesized by the host
744 plant. *Science* **356**: 1175-1178.
745
746 **Luginbuehl LH, Oldroyd G. 2017.** Understanding the Arbuscule at the Heart of Endomycorrhizal
747 Symbioses in Plants. *Current Biology* **27**: R952–R963.
748
749 **Luo W, Brouwer C. 2013.** Pathview: an R/Bioconductor package for pathway-based data
750 integration and visualization. *Bioinformatics* **29**: 1830-1831.
751

752 **MacLean AM, Bravo A, Harrison MJ. 2017.** Plant Signaling and Metabolic Pathways Enabling
753 Arbuscular Mycorrhizal Symbiosis. *Plant Cell* **29**: 2319-2335.
754
755 **Maillet F, Poinsot V, André O, Puech-Pagès V, Haouy A, Gueunier M, Cromer L, Giraudet D,**
756 **Formey D, Niebel A et al. 2011.** Fungal lipochitooligosaccharide symbiotic signals in arbuscular
757 mycorrhiza. *Nature* **469**: 58-63.
758
759 **Martin M. 2011.** Cutadapt removes adapter sequences from high-throughput sequencing reads.
760 *EMBnet Journal* **17**: 10-12.
761
762 **Müller LM, Flokova K, Schnabel E, Sun X, Fei Z, Frugoli J, Bouwmeester HJ, Harrison MJ.**
763 **2019.** A CLE-SUNN module regulates strigolactone content and fungal colonization in arbuscular
764 mycorrhiza. *Nature Plants* **5:** 933-939.
765
766 **Nasir F, Bahadur A, Lin X, Gao Y, Tian C. 2021.** Novel insights into host receptors and receptor-
767 mediated signaling that regulate arbuscular mycorrhizal symbiosis. *Journal of Experimental Botany*
768 **72**:1546-1557.
769
770 **Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara**
771 **RB, Simpson GL, Solymos P et al. 2019.** Vegan: community ecology pakage. R package version
772 2.5–5. https://CRAN.R-project.org/package=vegan
773
774 **Pimprikar P, Gutjahr C. 2018.** Transcriptional Regulation of Arbuscular Mycorrhiza Development.
775 *Plant and Cell Physiology* **59:** 673–690.
776
777 **Richter S, Kientz M, Brumm S, Nielsen ME, Park M, Gavidia R, Krause C, Voss U, Beckmann**
778 **H, Mayer U et al . 2014.** Delivery of endocytosed proteins to the cell-division plane requires change
779 of pathway from recycling to secretion. *eLife* **3**: e02131.
780
781 **Russo G, Carotenuto G, Fiorilli V, Volpe V, Faccio A, Bonfante P, Chabaud M, Chiapello M,**
782 **Van Damme D, Genre A. 2019.** TPLATE Recruitment Reveals Endocytic Dynamics at Sites of
783 Symbiotic Interface Assembly in Arbuscular Mycorrhizal Interactions. *Frontiers in Plant Science* **10:**
784 1628.
785
786 **Russo G, Carotenuto G, Fiorilli V, Volpe V, Chiapello M, Van Damme D, Genre A. 2018.** Ectopic
787 activation of cortical cell division during the accommodation of arbuscular mycorrhizal fungi. *New*
788 *Phytologist* **221**: 1036-1048.
789

790 **Salzer P, Bonanomi A, Beyer K, Voˈgeli-Lange R, Aeschbacher RA, Lange JAW, Kim D, Cook**
791 **DR, Boller T. 2000.** Differential expression of eight chitinase genes in *Medicago truncatula* roots
792 during mycorrhiza formation, nodulation, and pathogen infection. *Molecular Plant Microbe*
793 *Interactions* **13**: 763–777.
794
795 **Sanchez L, Weidmann S, Brechenmacher L, Batoux M, van Tuinen D, Lemanceau P,**
796 **Gianinazzi S, Gianinazzi-Pearson V. 2004.** Common gene expression in *Medicago truncatula*
797 roots in response to *Pseudomonas fluorescens* colonization, mycorrhiza development and
798 nodulation. *New Phytologist* **161**: 855–863.
799
800 **Schwacke R, Ponce-Soto GY, Krause K, Bolger AM, Arsova B, Hallab A, Gruden K, Stitt M,**
801 **Bolger ME, Usadel B. 2019.** MapMan4: A Refined Protein Classification and Annotation
802 Framework Applicable to Multi-Omics Data Analysis. *Molecular Plant* **12**: 879-892.
803
804 **Smith SE, Read DJ. 2008.** Mycorrhizal symbiosis. Academic Press, Inc., San Diego, CA
805
806 **Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N,**
807 **Grigoriev I, Gryganskyi A et al. 2016.** A phylum-level phylogenetic classification of zygomycete
808 fungi based on genome-scale data. *Mycologia* **108**: 1028-1046.
809
810 **Sun J, Miller JB, Granqvist E, Wiley-Kalil A, Gobbato E, Maillet F, Cottaz S, Samain E,**
811 **Venkateshwaran M, Fort S et al. 2015.** Activation of Symbiosis Signaling by Arbuscular
812 Mycorrhizal Fungi in Legumes and Rice. *Plant Cell* **27**: 823-838.
813
814 **Suzaki T, Yano K, Ito M, Umehara Y, Suganuma N, Kawaguchi M. 2012.** Positive and negative
815 regulation of cortical cell division during root nodule development in *Lotus japonicus* is
816 accompanied by auxin response. *Development* **139**: 3997-4006.
817
818 **Tang H, Krishnakumar V, Bidwell S, Rosen B, Chan A, Zhou S, Gentzbittel L, Childs KL,**
819 **Yandell M, Gundlach H et al. 2014.** An improved genome release (version Mt4.0) for the model
820 legume *Medicago truncatula*. *BMC Genomics* **15**: 312
821
822 **Tokunaga T, Hayashi H, Akiyama K. 2015.** Medicaol, a strigolactone identified as a putative
823 didehydro-orobanchol isomer, from *Medicago truncatula*. *Phytochemistry* **111**: 91-97.
824
825 **Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya**
826 **Y, Shirasu K, Yoneyama K et al. 2008.** Inhibition of shoot branching by new terpenoid plant
827 hormones. *Nature* **455**: 195-200.

Usadel B, Nagel A, Thimm O, Redestig H, Blaesing OE, Palacios-Rojas N, Selbig J, Hannemann J, Piques MC, Steinhauser D et al. 2005. Extension of the visualization tool MapMan to allow statistical analysis of arrays, display of corresponding genes, and comparison with known responses. *Plant Physiology* **138**: 1195-204.

Van Damme D, Gadeyne A, Vanstraelen M, Inze D, Van Montagu MC, De Jaeger G, Russinova E, Geelen D. 2011. Adaptin-like protein TPLATE and clathrin recruitment during plant somatic cytokinesis occurs via two distinct pathways. *Proceedings of the National Academy of Sciences USA* **108**: 615–620.

Vernié T, Camut S, Camps C, Rembliere C, de Carvalho-Niebel F, Mbengue M, Timmers T, Gasciolli V, Thompson R, le Signor C et al. 2016. PUB1 Interacts with the Receptor Kinase DMI2 and Negatively Regulates Rhizobial and Arbuscular Mycorrhizal Symbioses through Its Ubiquitination Activity in *Medicago truncatula*. *Plant Physiology* **170**: 2312–2324.

Volpe V, Carotenuto G, Berzero C, Cagnina L, Puech-Pagès V, Genre A. 2020. Short chain chito-oligosaccharides promote arbuscular mycorrhizal colonization in *Medicago truncatula*. *Carbohydrate Polymers* **229**: 115505.

Wang Y, Bouwmeester HJ. 2018. Structural diversity in the strigolactones. *Journal of Experimental Botany* **69**: 2219-2230.

Waters MT, Gutjahr C, Bennett T, Nelson DC. 2017. Strigolactone Signaling and Evolution. *Annual Review of Plant Biology* **68**: 291-322.

Wewer V, Brands M, Dörmann P. 2014. Fatty acid synthesis and lipid metabolism in the obligate biotrophic fungus *Rhizophagus irregularis* during mycorrhization of *Lotus japonicus*. *Plant Journal* **79**: 398-412.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. 2nd ed. Springer International Publishing; Cham, Switzerland.

Yoneyama K, Xie X, Sekimoto H, Takeuchi Y, Ogasawara S, Akiyama K, Hayashi H, Yoneyama K. 2008. Strigolactones, host recognition signals for root parasitic plants and arbuscular mycorrhizal fungi, from Fabaceae plants. *New Phytologist* **179**: 484-494.

865 **Young MD, Wakefield MJ, Smyth GK, Oshlack A. 2010.** Gene ontology analysis for RNA-seq:
866 accounting for selection bias. *Genome Biology* **11**: R14.
867
868 **Young ND, Debellé F, Oldroyd GE, Geurts R, Cannon SB, Udvardi MK, Benedito VA, Mayer**
869 **KF, Gouzy J, Schoof H et al. 2011.** The Medicago genome provides insight into the evolution of
870 rhizobial symbioses. *Nature* **480**: 520-524
871
872 **Zipfel C, Oldroyd GE. 2017.** Plant signaling in symbiosis and immunity. *Nature* **543**: 328-336.
873
874

Figures and Tables

877
878
Fig. 1. Global overview of transcriptomic data. A) Principal component analysis (PCA) plots
highlighted a clear separation of the expression profiles between AM inoculated (dark and light
orange) and non-inoculated (dark and light blue) samples, as well as between treated (dark orange
e dark blue) and non-treated (light orange and light blue) ones. Three biological replicates
(individual dots) are plotted per condition and time point. B) Variance Partitioning Analysis (VPA)
considered three factors: AM fungus presence (inoculation), CO treatment and time (10, 14, 21, 28
dpi). The strongest influence on gene regulation resulted to depend on time, while CO treatment
had a major effect at the earliest time point (10 dpi). C) Differentially expressed genes (DEGs) in
the four comparisons and at different time points. Number and expression trend of DEGs. Up-
regulation is shown in red, down-regulation in blue (FDR <0.05).

889
890

Fig. 2. CO effect on SL-related gene expression and SL accumulation. A) Heatmap plot of strigolactone-related genes in each comparison and time points. A progressive decrease in strigolactone biosynthesis (blue) is in line with the progress of AM development in MYC *vs* CTR. This down-regulation was earlier and more extensive in MYC+CO *vs* CTR comparison, in line with the acceleration of colonization observed in the presence of CO treatment. Remarkably, a strong up-regulation (red) was evident in both strigolactone biosynthesis and transport until 21 dpi In the CTR+CO *vs* CO comparison. Genes are clustered within each category according to Spearman's distance (Müller et al., 2019). B) Didehydro-orobanchol content in root tissues and exudates of CTR and CTR+CO plants sampled at 21 dpi. A significant increase in DDO root content correlated with CO treatment. A similar trend was observed in the exudates, even if the difference between CTR and CTR+CO plants was statistically not significant. Mean values ± SDs of four biological replications for each treatment are shown. Each biological replication was composed by two plants. Asterisks indicate statistically significant differences (Student's t-test, *P < 0.05, **P < 0.01, and ***P < 0.001).

909
910
911
Fig. 3. Local regulation of early AM marker genes. Local CO treatment stimulated concentration-
independent expression of several genes involved in fungal accommodation. The highest number
of significantly upregulated genes was recorded upon 1mg/L CO application. Grey bars = control,
water treated roots; black bars = CO treated roots. Mean values ± SDs of five biological replicates
for each treatment are shown. Asterisks indicate statistically significant differences (Student's t-
test, *P < 0.05, **P < 0.01, and ***P < 0.001).

925
Fig. 4. PPA formation in epidermal cells of *M. truncatula*. Root organ cultures expressing GFP-
HDEL in the endoplasmic reticulum (ER) were treated with 1mg/L CO or water (H_2O) as control at
the time of inoculation with pre-germinated *G. margarita* spores. Epidermal cells in the vicinity of
fungal hyphopodia were then imaged at 7, 10, 14 dpi and categorized based on the absence or
presence of ER aggregations (A). A statistically significant increase in the percentage of epidermal
cells with prominent ER aggregation was recorded in CO-treated roots at 7 and 10 dpi. At least 100
independent infection events were visualized for each treatment. Asterisks indicate statistically
significant differences (Student's t-test, $*P < 0.05$, $**P < 0.01$, and $***P < 0.001$).
934
935

A)

Gene ID	Log2 FoldChange at 28 dpi		Annotation
Medtr1g028600	8,49	6,87	*MtPT4*
Medtr7g068600	8,16	6,90	*MtMYB*
Medtr1g036410	7,44	6,06	*MtAMT1*
Medtr1g040500	7,22	6,47	*MtRAM2*
Medtr4g093845	6,46	5,16	*MtABC*
Medtr5g011320	7,60	6,26	*MtSubtilisin*
Medtr5g076900	8,11	6,46	*MtGST*
Medtr5g031090	8,09	6,29	*MtLectin*
Medtr8g006790	7,48	6,36	*MtHA1*
	MYC vs CTR	**MYC+CO vs CTR**	

B) Arbuscule morphology at 28 dpi

Fig. 5. Quantitative analyses of mycorrhizal phenotypes between MYC and MYC+CO plants.
A) Expression of AM-induced genes at 28 dpi was declined in the comparison MYC+COvsCTR respect to MYCvsCTR, suggesting an arbuscule senescence. B) Arbuscules were imaged at 28 dpi in MYC and MYC+CO plants and classified based on their developmental stage: developing arbuscules with a trunk and very few branches (stage I), intermediate arbuscules with thick branches (stage II), fully mature arbuscules with abundant fine branches (stage III) and senescent arbuscules with clusters of collapsed branches (stage IV). A significantly higher presence of stage II arbuscules was recorded in MYC compared to MYC+CO plants. By contrast, MYC+CO plants had significantly more Stage III and Stage IV arbuscules. This indicates that the advance in AM development caused by CO treatment extends to four weeks post inoculation. Data are based on the measurements of 30 infection units for each of four biological replicates. Mean values ± SDs of four biological replications for each treatment are shown. Asterisks indicate statistically significant differences (Student's t-test, *P < 0.05, **P < 0.01, and ***P < 0.001).

952
953
954

955
956
Fig. 6. Schematic representation of exogenous CO effects on AM symbiosis development.
The perception of root-released SL by AM fungi is known to induce hyphal branching and boost
CO release. In turn, CO perception activates root symbiotic responses leading to intracellular fungal
accommodation and symbiosis establishment. Our current results (red arrows) indicate that
exogenous CO application further stimulates SL production in the host root, suggesting the
presence of a positive feedback mechanism that we propose here as a presymbiotic signaling loop.
As a consequence of this reinforcement of presymbiotic signaling, we also observed a stimulation
of intracellular fungal accommodation, supporting the observed advance in root colonization and
symbiosis development.
966

967 **Table 1.** Main AM symbiosis-related pathways impacted at 10 dpi by CO treatment (n.d.= not
968 detected).

Gene ID	MYC vs CTR	MYC+CO vs CTR	CTR+CO vs CTR	MYC+CO vs MYC	Annotation
	Log2 fold-change value				*Strigolactone signaling*
Medtr3g045440	n.d.	n.d	1.21	n.d	sigma factor sigb regulation rsbq-like protein DLK2
Medtr3g110195	n.d.	0.35	1.00	n.d	retinal pigment epithelial membrane protein CCD8b
Medtr3g104560	n.d.	0.47	1.13	n.d	cytochrome P450 family protein MAX1a
Medtr7g045370	n.d.	n.d.	0.66	n.d	carotenoid cleavage dioxygenase CCD7
Medtr1g471050	n.d.	0.75	0.63	n.d	beta-carotene isomerase D27
Medtr3g109610	n.d.	n.d	0.94	n.d	carotenoid cleavage dioxygenase CCD8-1
Medtr3g107870	n.d.	n.d	1.41	n.d	drug resistance transporter-like ABC domain protein PDR1a
Medtr7g063800	n.d.	n.d	1.52	n.d	carotenoid cleavage dioxygenase CCD8-2
					External stimuli response
Medtr5g076900	n.d.	5.82	n.d.	n.d.	glutathione S-transferase amino-terminal domain protein
Medtr5g035640	n.d.	4.73	n.d.	n.d.	polygalacturonase 1 beta-like protein 3
Medtr7g068600	n.d.	3.91	n.d.	n.d.	myb-like transcription factor family protein MYB1
Medtr1g040500	1.04	3.73	n.d.	n.d.	glycerol-3-phosphate acyltransferase RAM2
Medtr1g109110	0.82	3.02	n.d.	n.d.	palmitoyl-acyl carrier thioesterase FatM
Medtr6g011490	n.d.	2.56	n.d.	n.d.	AP2 domain transcription factor RAM1
Medtr4g104020	0.77	2.26	n.d.	n.d.	GRAS family transcription factor RAD1
Medtr5g030910	1.13	1.19	n.d.	-0.94	lipid transporter STR
					Cytoskeleton organization
Medtr6g082470	n.d.	0.93	n.d.	1.20	ATP-binding microtubule motor family protein
Medtr1g082920	n.d.	0.79	n.d.	1.20	kinesin motor domain protein
Medtr5g094310	n.d.	0.79	n.d.	1.15	kinesin motor domain protein
Medtr7g091290	n.d.	0.71	n.d.	1.07	kinesin motor catalytic domain protein
Medtr1g075680	n.d.	0.74	n.d.	1.06	kinesin motor domain protein
Medtr6g053680	n.d.	0.75	n.d.	1.03	ATP-binding microtubule motor family protein
Medtr7g061020	n.d.	0.80	n.d.	0.96	ATP-binding microtubule motor family protein
Medtr2g010760	n.d.	0.73	n.d.	0.96	kinesin motor catalytic domain protein
Medtr4g092140	n.d.	0.68	n.d.	0.95	kinesin motor catalytic domain protein
Medtr1g017540	n.d.	0.62	n.d.	0.93	kinesin motor catalytic domain protein
Medtr5g031470	n.d.	0.68	n.d.	0.93	kinesin motor domain Di-glucose-binding protein
Medtr3g107650	0.74	1.65	n.d.	-0.91	kinesin-associated protein
					Cell cycle organization
Medtr3g102530	-0.62	0.59	n.d.	1.21	carboxy-terminal domain cyclin, Cyclin A
Medtr7g113510	n.d.	0.94	n.d.	1.19	cyclin family protein, putative; mitosis division plane proteinTAN1
Medtr3g110405	n.d.	0.88	n.d.	1.16	Serine/Threonine-kinase aurora-like protein, homolog to AtAUR1
Medtr5g088980	n.d.	0.80	n.d.	1.15	carboxy-terminal domain cyclin, Cyclin B
Medtr3g088415	n.d.	0.85	n.d.	1.11	carboxy-terminal domain cyclin, Cyclin A
Medtr5g012010	n.d.	0.73	n.d.	1.10	syntaxin of plants 122 protein, KNOLLE protein
Medtr8g074000	n.d.	0.84	n.d.	1.06	carboxy-terminal domain cyclin, Cyclin B
Medtr7g089080	n.d.	0.71	n.d.	1.05	Cyclin B
Medtr5g023790	n.d.	0.73	n.d.	1.02	carboxy-terminal domain cyclin, Cyclin B
Medtr5g011390	n.d.	0.71	n.d.	0.86	cell cycle regulated microtubule-associated protein, AURORA
					Cell wall organization
Medtr8g009560	n.d.	1.73	1.45	n.d.	fatty acid hydroxylase superfamily protein
Medtr7g062250	n.d.	1.52	n.d.	1.15	laccase/diphenol oxidase family protein
Medtr5g075320	n.d.	1.30	n.d.	0.93	expansin A10
Medtr2g097030	n.d.	1.20	0.88	0.68	expansin A10
Medtr4g053380	n.d.	1.12	n.d.	1.41	fasciclin-like arabinogalactan protein
Medtr4g081950	n.d.	1.04	n.d.	n.d.	expansin-A1-like protein
Medtr5g079950	n.d.	1.03	0.90	0.62	expansin A10
Medtr2g086070	n.d.	-1.00	-0.72	n.d.	gland-specific fatty acyl-CoA reductase
Medtr4g019225	n.d.	-1.07	n.d.	n.d.	laccase/diphenol oxidase family protein
Medtr3g098980	n.d.	-1.10	-0.70	-0.83	omega-hydroxypalmitate O-feruloyl transferase
Medtr6g086365	-1.16	n.d.	-2.20	n.d.	arabinogalactan protein
Medtr6g086390	n.d.	n.d.	-1.59	n.d.	arabinogalactan peptide-like protein
Medtr4g059720	n.d.	n.d.	n.d.	0.83	fasciclin-like arabinogalactan protein
Medtr5g081810	n.d.	n.d.	n.d.	1.18	laccase/diphenol oxidase family protein

969
970