1   Aidan Foo[1], Louise Cerdeira[2], Grant L. Hughes[1], Eva Heinz[3*]
2
3   **Author affiliations**
4
5   [1]Departments of Vector Biology and Tropical Disease Biology, Centre for Neglected Tropical
6   Disease, Liverpool School of Tropical Medicine, Liverpool, UK
7
8   [2]Department of Vector Biology, Liverpool School of Tropical Medicine, Liverpool, UK
9
10 [3]Departments of Vector Biology and Clinical Sciences, Liverpool School of Tropical Medicine,
11 Liverpool, UK
12
13 *Corresponding author: eva.heinz@lstmed.ac.uk
14

15 **Title**

16 Recovery of Metagenomic Data from the *Aedes aegypti* Microbiome using a Reproducible
17 Snakemake Pipeline: MINUUR
18

Abstract

Background

Ongoing research of the mosquito microbiome aims to uncover novel strategies to reduce pathogen transmission. Sequencing costs, especially for metagenomics, are however still significant. A resource that is increasingly used to gain insights into host-associated microbiomes is the large amount of publicly available genomic data based on whole organisms like mosquitoes, which includes sequencing reads of the host-associated microbes and provides the opportunity to gain additional value of these initially host-focused sequencing projects.

Methods

To analyse non-host reads from existing genomic data, we developed a snakemake workflow called MINUUR (**M**icrobial **IN**sights **U**sing **U**nmapped **R**eads). Within MINUUR, reads derived from the host-associated microbiome were extracted and characterised using taxonomic classifications and metagenome assembly followed by binning and quality assessment. We applied this pipeline to five publicly available *Aedes aegypti* genomic datasets, consisting of 62 samples with a broad range of sequencing depths.

Results

We demonstrate that MINUUR recovers previously identified phyla and genera and is able to extract bacterial metagenome assembled genomes (MAGs) associated to the microbiome. Of these MAGS, 42 are high-quality representatives with >90% completeness and <5% contamination. These MAGs improve the genomic representation of the mosquito microbiome and can be used to facilitate genomic investigation of key genes of interest. Furthermore, we show that samples with a high number of KRAKEN2 assigned reads produce more MAGs.

Conclusions

Our metagenomics workflow, MINUUR, was applied to a range of *Aedes aegypti* genomic samples to characterise microbiome-associated reads. We confirm the presence of key mosquito-associated symbionts that have previously been identified in other studies and recovered high-

quality bacterial MAGs. In addition, MINUUR and its associated documentation are freely

available on GitHub (https://github.com/aidanfoo96/MINUUR) and provide researchers with a

convenient workflow to investigate microbiome data included in the sequencing data for any

applicable host genome of interest.

Keywords

Mosquito, *Aedes aegypti,* Microbiome, Metagenomics, Snakemake, MAGs, Read Classification.

Introduction

Aedes aegypti is an important vector of human pathogens including dengue virus (DENV), yellow fever virus, chikungunya virus and Zika virus. DENV cases alone are estimated to cause 10,000 deaths and 100 million infections per year, contributing to significant burden of human morbidity and mortality worldwide (1). Interest in the mosquito microbiome has emerged due to evidence of its influence in vectorial capacity (2,3), offering potential for novel approaches to reduce pathogen transmission from mosquitoes to vertebrate hosts (2,4–7).

Mosquito microbiomes are highly variable, dependent on multiple deterministic processes such as the environment (8–12), season (13), host factors (14,15), microbial interactions (16–18) and mosquito-microbe interactions (19–23). These findings from mosquito microbiome studies are largely driven by amplicon based 16S rRNA sequencing approaches (18,24,25). Metagenomic approaches for mosquito microbiome characterisation are limited in number (26,27), but would facilitate our understanding by providing the genomic context of symbionts (28). An attractive resource for gaining additional insights into microbiomes is to make use of the microbiome reads derived from whole genome sequencing (WGS), especially in cases where preparation of the host for sequencing included its associated microbiome. Studies in *Drosophila*, bumble bees, moths and nematodes have shown existing WGS data is a rich source to characterize associated symbionts (29–34). Whilst some studies include specific enrichment of non-host with bait sequences targeting a specific taxon of interest (29,32); it is also possible to assess microbes present in the sequencing experiment without prior enrichment, taking into account that in the latter case the microbiome recovered is likely a biased representation and lack of presence does not prove absence.

Whole genome shotgun sequencing is commonly used to study mosquito genomics at the individual (35,36) and population (37,38) level; meaning non-mosquito sequence data (we refer to these as non-reference reads, since they do not map to the reference genome of interest) are

a source to identify mosquito microbiome members using metagenomics. Genomic surveillance

programs such as the *Anopheles gambiae* 1000 Genomes Project contain a large number of

genomic samples with each release (39) and, at time of writing, currently 100,514 *Aedes aegypti*

whole-genome sequencing runs are deposited on the European Nucleotide Archive. As such,

there is great potential to leverage existing mosquito WGS data to explore members of their

microbiomes from non-reference reads.

To make use of this resource and streamline future large-scale analysis of mosquito

metagenomes, we developed a Snakemake pipeline to provide **M**icrobial **IN**sight **U**sing

Unmapped **R**eads from WGS data. MINUUR uses short read, whole genome sequencing data as

input and performs an analysis of non-reference reads derived from a host sequencing

experiment. We used MINUUR to study five published *Aedes aegypti* sequencing projects (in total

124 FASTQ files) currently available on the European Nucleotide Archive. We gained insight of

associated microbes based on taxonomic read classifications and recover high-quality

metagenome assembled genomes (MAGs) of mosquito-associated bacteria. To assess the

suitability of samples for MAG recovery, we also investigated patterns between KRAKEN2

assigned reads and the number of MAGs within samples. We show samples with taxa assigned

high numbers of KRAKEN2 classified reads produce relatively more high and medium quality

MAGs. The pipeline is open source and available on GitHub

(https://github.com/aidanfoo96/MINUUR) with an accompanying JupyterBooks page

(https://aidanfoo96.github.io/MINUUR/).

Materials and Methods

Specifications

To undertake this analysis, we developed a metagenomics workflow called MINUUR (**M**icrobial

INsights Using Unmapped Reads), using the workflow manager Snakemake (40) (Figure 1). This

pipeline was developed to facilitate the following analysis and future studies aiming to

characterise non-reference reads in mosquitoes or other organisms. A breakdown of the pipeline

that produced each result of this study is provided in the following section, with details of how

each step was configured in the final section.

Database Setup

MINUUR requires several databases to perform the analysis. This includes a high quality bowtie2-

indexed reference genome (41) to separate host and non-host reads, and a KRAKEN2 (42),

BRACKEN (43) and MetaPhlAn3 (44) database for taxonomic read classifications. All databases

are available in their respective GitHub repositories. The databases used in this study are the

default MetaPhlAn3 marker gene database, KRAKEN2 and BRAKEN indexes from the Ben

Langmead repository located here: https://benlangmead.github.io/aws-indexes/k2. For our study,

we downloaded and compiled these default databases on the 08/09/2022.

Data Preparation

MINUUR accepts either BAM or paired FASTQ inputs. For FASTQ inputs, MINUUR performs

quality control (QC) using FASTQC (v0.11.9) (45), providing a QC report per sample. MINUUR

does not use the FASTQC report in subsequent steps, but only as a quality assurance metric for

the user and to estimate if read trimming is required. Read trimming can be performed within

MINUUR using Cutadapt (v1.5) (46) with user defined parameters for minimum read length, base

quality and adapter content (default: minimum base length = 50, average base quality = 30). To

separate host and non-host, reads are aligned to a user defined indexed reference genome (the

relevant host genome) using Bowtie2 (v2.4.4) (41). Alignment sensitivity and type can be adjusted

within the pipeline at the user's discretion. A high quality, chromosome level assembled, reference

genome is recommended if available. In situations where this is not possible, users should be

aware that read alignment will likely result in mismatches between the reference and target

sequence and produce alignments with poor coverage (47). As a result, non-reference reads used

in subsequent steps are likely to contain a substantial number of host data. In this instance, we

included a feature within MINUUR to extract KRAKEN2 assigned reads pertaining to known

microbes and potentially improve metagenome assemblies. Non-reference reads within the

coordinate sorted binary alignment (BAM) file are extracted using samtools using the command

"samtools -view -f 4" (v.1.14) (48) and converted to FASTQ format using bedtools (v2.3.0) (49)

("–bamToFastq"). As users might already have their data in BAM format mapped against the host

reference (e.g. in large-scale sequencing projects like the Ag1000G), we also included the option

of a BAM input. Here, any non-reference reads within the BAM file will be extracted, converted to

FASTQ and used in downstream steps.

For this study, we used five published genome datasets of *Aedes aegypti* publicly available on

the European Nucleotide Archive (35,37,50–53). We selected the data sets to cover a range of

sequencing depths, DNA extraction method and sequencing platform (Figure 2C). All raw FASTQ

files of published sequencing data were retrieved from the European Nucleotide Archive (ENA)

under the project accession numbers PRJEB33044 (37)**,** PRJNA255893 (50), PRJNA385349

(51), PRJNA718905 (35), PRJNA776956 (52) and PRJNA992905 (53).

Read Classification

MINUUR uses two read classification approaches to infer taxonomy. KRAKEN2 (v2.1.2) (42),

which uses a k-mer based approach to map read fragments of k-length against a taxonomic

genome library of k-mer sequences, and MetaPhlAn3 (v3.0.13) (44) to align reads against a

library of marker genes using Bowtie2 (41). MINUUR also provides the option to use KRAKEN2

classified reads, parsed from KrakenTools (v1.2), to select a specific set of reads (for example

bacterial) for metagenome assembly. To estimate the relative taxonomic abundance from

KRAKEN2 classifications, MINUUR will parse KRAKEN2 read classifications to BRACKEN

(v2.6.2) (43) which uses a Bayesian probability approach to redistribute reads assigned at higher

taxonomic levels to lower (species) taxonomic levels.

MINUUR outputs classified and unclassified reads to paired FASTQ files and generates

BRACKEN estimated taxonomic abundance profiles for further analysis. An additional feature we

included within MINUUR is the option to extract a specific taxon or group of taxa from KRAKEN2,

using the program KrakenTools (42). This option is useful in situations where a specific group of

taxa are of interest or to exclude groups of taxa such as viral or archaeal reads. Alternatively, if

alignment to a reference is poor, this option can be used to remove host reads that did not map

to its reference.

Metagenome Assembly, Binning and Quality Assurance

MINUUR uses the non-reference reads to perform *de novo* metagenome assemblies (the same

reads used for KRAKEN2 and MetaPhlAn3 taxonomic classifications). Reads are parsed to

MEGAHIT (v1.2.9) (54), a rapid and memory efficient metagenome assembler, for *de novo*

metagenome assembly. Assembled contigs are quality checked using QUAST (v5.0.2) (55) to

assess contig N50 and L50 scores. The resultant contigs, which are fasta files with sequences

pertaining to genomic regions of a microbe, need to be placed within defined taxonomic groups -

referred to as a bin. For this, contigs are indexed using the Burrows Wheeler Aligner (BWA)

(v0.7.17) (56), and the original non-reference reads are aligned to the indexed contigs using "–

bwa-mem". The subsequent coordinate sorted BAM file is parsed to the

"jgi_summarize_bam_contig_depth" script from MetaBAT2 (v2.12.1) (57) to produce a depth file

of contig coverage. The depth file and assembled contigs are input to the metagenome binner

MetaBAT2 (v2.12.1) (57), to group contigs within defined genomic bins. Each bin is a predicted

metagenome assembled genome (MAG). CheckM (v1.1.3) (58) is then used for quality assurance

of each bin by identifying single copy core genes. Specifically, bin contamination is assessed by

looking for one single copy core gene within each bin, and completeness by calculating a required

set of single copy core genes.

Pipeline Configuration

All sample names were listed in the samples.tsv file of the configuration directory and paths to

the FASTQ directories given in the samples_table.tsv file. To implement the pipeline, the

configuration file was set to the following parameters: FASTQ = True, QC = True,

CutadaptParams = "–minimum-length 50 -q 30", RemoveHostFromFastqGz = True,

AlignmentSensitivity = "–sensitive-local", ProcessBam = True, From-Fastq = True,

KrakenClassification = True, ConfidenceScore = 0, KrakenSummaries = True,

GenusReadThreshold = 1000, SpeciesReadThreshold = 30000, MetagenomeAssm = True,

MetagenomeBinning = True, MinimumContigLength = 1500, CheckmBinQA = True. All databases

were installed from their respective repositories on 08/09/22. The pipeline was run on an Ubuntu

Linux system with 660gb of available memory and 128 CPUs. For our analysis, with the above

settings and 10 cores available, runtime was two weeks; the maximum Resident Set Size (RSS)

of an individual sample during this run was 9771 RSS (occurring during metagenome assembly);

and total storage used (including temporary files) was 5.18Tb (terabytes) across all samples used

in this study.

Taxonomic Classification of MAGs with GTDB-Tk

Separate from MINUUR, all bins produced from MetaBAT2 were taxonomically classified with

GTDB-Tk (59) (v2.1.1) using "–classify-wf" against the Genome Taxonomy Database (GTDB)

(release 07-R207 08/04/22, downloaded on 08/09/22). GTDB-Tk assigns genes to MAGs using

Prodigal (v2.6.3) (60) and ranks the taxonomic domain of each MAG using a database of 120

bacteria and 122 archaea marker genes (61) using HMMER3 (62). With this information, MAGs

are then placed into domain specific reference trees with pplacer (v1.1) (63). Taxonomic

classifications with GTDB-Tk are based on placement within the GTDB reference tree, relative

evolutionary divergence, and average nucleotide identity (ANI) scores with its closest reference

genome. The relative evolutionary divergence score is used to refine ambiguous taxonomic rank

assignments and ANI scores are used to define species classifications (59).

Results

Extraction of Non-Reference Reads Post-Alignment to *Aedes aegypti*

In total, we retrieved 62 samples (124 FASTQ files) across six sequencing experiments (Figure

2A) and parsed them through MINUUR. After alignment to the *Aedes aegypti* reference genome

(AaegL5.3) (64) with Bowtie2, the proportion of mapped and non-reference reads were calculated.

Of our initial 62 samples, on average, 176,310,722 reads aligned to the AaegL5.3 reference

genome, ranging between 20,655,064 and 543,378,958 reads (Figure 2B). To estimate the

number of reads associated to the microbiome among the non-reference reads, the overall

number of KRAKEN2 classifications were counted. The average proportion of KRAKEN2

classified reads from all non-reference reads was 17.9%. The number and proportion of

KRAKEN2 classifications per sample is given in Supplementary Table 1 within the extended data.

Taxonomic Classifications Produced from Non-Reference Reads of *Aedes aegypti*

The KRAKEN2 classifications of non-reference reads were counted from each sequencing

project. Four out of the six sequencing projects showed a much lower number of classifications

compared to sequencing projects PRJNA776596 and PRJEB33044 (Figure 3A). PRJEB33044

gave the highest average proportion of taxonomic classification (81.3%), while PRJNA255893

was the lowest (0.893%).

Multiple phyla were identified across sequencing projects. *Bacteroidetes, Proteobacteria,*

Firmicutes and Actinobacteria were the most common phylum present across all samples, with

varied relative abundance between projects (Figure 3C). For example, *Bacteroidetes* and

Proteobacteria were dominant in PRJEB33044 and PRJNA882905, *Firmicutes* and

Proteobacteria in PRJNA255893, and *Bacteroidetes, Actinobacteria* and *Proteobacteria* in

PRJNA385349 and PRJNA718905. At the genera level, there are several dominant members

across samples, including *Wolbachia, Pseudomonas, Serratia* and *Elizabethkingia.* Generally,

however, there is considerable variation at the genus level within and between sequencing

experiments (Figure 3C). We summarised all KRAKEN2 classifications across 62 samples. The

most common genera, identified in >50% of all samples, were *Pseudomonas, Elizabethkingia,*

Clostridium, Bacillus and *Chryseobacterium.* Genera identified in 25-50% of samples were

Acinetobacter, *Enterobacter, Serratia* and *Delftia* (Figure 3B). All KRAKEN2 taxonomic

classifications are given in Supplementary Table 2.

Metagenome Assembly and Binning

We used MINUUR to further parse non-reference reads through assembly, binning and quality

assurance steps, with the aim to recover MAGs associated to the *Aedes aegypti* microbiome.

Assembly was conducted with MEGAHIT and binning with MetaBat2. We used CheckM to assess

MAG completeness and contamination through the presence and copy number of single copy

core genes, and recovered 105 MAGs (Figure 4A). Using the standards of MAG quality set by the

genome standards consortium (GSC) (65), 42 MAGs met the criteria of high quality with

completeness >90% and contamination <5%; and 20 MAGs classify as medium quality with

completeness >50% and contamination <5%. Overall, 62 high and medium quality MAGs were

recovered. The remaining MAGs were low-quality (<50% complete and <10% contamination) or

contaminated >10%, and therefore excluded from MAG classification. Of the high and medium-

quality MAGs, the mean N50 (the minimum contig length of an assembled contig that covers 50%

of the genome) was 177.2 kilobase pairs (KBP), ranging between 4.72KBP and 471.6KBP (Figure

4B). The average genome size was 3.36 megabase pairs (MBP), ranging between 1.07MBP and

6.19MBP (Figure 4C), while low-quality MAGs showed a wider range of genome sizes between

0.24MBP and 106MBP (Supplementary Figure 1).

Relationship Between KRAKEN2 Taxonomic Classifications and MAG Recovery

The suitability of a sample for MAG recovery would be of interest to estimate in advance, and we

investigated if there was a correlation between the proportion of KRAKEN2 classifications from

non-reference reads and MAG recovery. In one project, PRJNA255893, we recovered no high-

quality MAGs, (Figure 5A) and the reads used for assembly contained no taxa exceeding 100,000

KRAKEN2 assigned reads (Figure 5B). In contrast, PRJNA33044, PRJNA776596 and

PRJNA882905 allowed retrieval of more high and medium quality MAGs, and within these

projects, multiple samples contained >100,000 reads assigned to a taxon (Figure 5A, B). As such,

MAG recovery is likely linked to a sufficient number of classified reads to a taxon (Figure 5A,

Figure 5B), rather than overall number of KRAKEN2 assigned reads within a sample

(Supplementary Figure 2). In accordance with this, the total number of classified taxa totaling or

greater than 100,000 KRAKEN2 assigned reads for PRJEB33044 is 33 taxa (summed across all

samples) and 22 high and medium quality MAGs were recovered from this sample. Similarly,

PRJNA776956 shows 19 taxa with associated reads totaling or greater than 100,000 KRAKEN2

assigned reads, resulting in 16 high and medium-quality MAGs. Furthermore, taxonomic

classifications with a high number of KRAKEN2 assigned reads (>100,000) are akin to the

taxonomic classifications of medium and high-quality MAGs (Supplementary Figure 3). There are,

however, exceptions to these patterns; PRJNA385349 contained five taxa with over 100,000

KRAKEN2 assigned reads, yet no high-quality MAGs could be recovered from this project. As

such, applicable for future application of this approach, the number of KRAKEN2 classifications

assigned to a specific taxon is one factor that can help estimate high-quality MAG recovery.

Taxonomic Classification of MAGs with GTDB-Tk

Following MAG recovery using MINUUR, we used the taxonomic classifier GTBD-Tk to classify

high and medium quality MAGs against the Genome Taxonomy Database (GTDB). We compared

the genome size of each MAG to its closest reference genome in the GTDB (Figure 6A, Figure

6B). Of the high-quality MAGs, these were larger than their reference genome by mean = 183kb,

whereas medium quality MAGs deviated from their reference genomes by 1768kb (Figure 6A).

Congruent with pairwise size differences between MAG and reference genome, we found the

overall distribution of high-quality MAGs compared to their reference genome size to be similar,

but significantly different between medium-quality MAGs (Figure 6B). Of these, 48 MAGs were

classified to the species level with a mean FastANI score of 98.4%, ranging between 95.6% and

100%. No MAGs were identified <95% ANI to a known species, indicating no undescribed *Aedes*

aegypti associated bacterial species were present within these MAGs (Figure 6C).

Discussion

Metagenomic datasets of mosquito microbiomes are so far limited in number (66). In this study,

we developed a metagenomics like workflow called MINUUR to facilitate recovery and use of

host-associated bacterial sequences using non-reference reads from existing host WGS projects.

We demonstrate that MINUUR can be used to recover genus level taxonomic classifications and

draft high and medium quality MAGs from host WGS projects. The recovery of metagenomic

information, such as MAGs, are applied in large scale metagenomic studies from chickens (67),

humans (68) to cows (69–71), with these studies yielding between 400 to 92,000 MAGs per study.

We apply a similar approach with non-reference *Aedes aegypti* sequencing reads across a range

of different studies and can demonstrate that using MINUUR expands the genomic representation

of known mosquito-associated bacterial symbionts. Overall, these provide a valuable resource for

researchers in the field and can be used in further work such as facilitating biosynthetic gene

cluster discovery (71) or to identify genetic targets in symbiont pathogen blocking approaches (2).

The data retrieved in this study agrees with published insights; the phylum level classifications

are consistent with findings from other mosquito microbiome studies (8,9,11,18,72,73), showing

that *Proteobacteria, Bacteroidetes* and *Firmicutes* are dominant phyla of the mosquito

microbiome; and our taxonomic classifications highlight the inherent variability of the *Aedes*

aegypti microbiome (18,24,74). These findings give us confidence that taxonomic classifications

with KRAKEN2, within the pipeline, can accurately predict the presence of microbes associated

to the *Aedes aegypti* microbiome from non-reference sequences. At the genus level, we also find

consistent observations of taxa previously identified in other studies (16,19,23,75,76). The

KRAKEN2 classifications show within the two most common phyla, *Proteobacteria* and

Bacteroidetes; *Elizabethkingia, Pseudomonas* and *Serratia* are the most common. All three of

these symbionts are documented to play key roles in either blood digestion (19,77), iron-

acquisition (75) and microbial interactions (16). Notably *Elizabethkingia* has previously been

implicated in responses to iron fluxes in *Anopheles gambiae* (75), and blood meals in *Aedes*

albopictus (25) and *Aedes aegypti* (24). Similarly, *Pseudomonas* has shown to interact with

Elizabethkingia, triggering the expression of *hemS* to break down heme into biliverdin catabolites

(17). It is encouraging to note the presence of these two bacteria within our taxonomic

classifications. We recovered high and medium quality MAGs associated to these taxa (*Serratia,*

Elizabethkingia and *Pseudomonas*), which should allow further interrogation of genes associated

to these biological processes. To note, the presence of *Wolbachia* in the projects we have

analysed is expected since these mosquitoes were transinfected with high titers of this bacterium,

further validating the pipeline results (37).

Whilst the nature of the samples can act as confounder given differences in sample handling

preparation, we note the relative abundance of the key phyla, and constituent genera, are varied

across these projects. Putative biological causes of this variation are likely multifactorial,

supported by studies showing environment (9,15), host genetics (23) and competitive

mechanisms amongst bacteria (16) to be influential for bacterial colonization in the mosquito.

Conclusion

In summary, we present a reproducible workflow to analyse host-associated microbial sequence

data derived from host WGS experiments, leveraging a vast resource of data for additional

insights. Our work focuses on the mosquito microbiome, where future considerations and

prospects were recently established by the Mosquito Microbiome Consortium (66). A key point

highlighted in this statement was the need for (meta)genomics approaches with solid

reproducibility for data analysis within the field. Our pipeline provides a workflow to assess non-

host reads from existing mosquito genome sequence data and increases our knowledge of

mosquito-associated bacterial genomes. This approach and accompanying workflow will facilitate

more analyses of existing WGS data within *Aedes aegypti* and other organisms of interest for the

scientific community.

Data and Software Availability

All original sequencing projects can be accessed under the following accession numbers:

PRJEB33044 (https://www.ebi.ac.uk/ena/browser/view/PRJEB33044), PRJNA255893

(https://www.ebi.ac.uk/ena/browser/view/PRJNA255893), PRJNA385349

(https://www.ebi.ac.uk/ena/browser/view/PRJNA385349), PRJNA718905

(https://www.ebi.ac.uk/ena/browser/view/PRJNA718905), PRJNA776956

(https://www.ebi.ac.uk/ena/browser/view/PRJNA776956), PRJNA882905

(https://www.ebi.ac.uk/ena/browser/view/PRJNA882905). The source code for our workflow,

MINUUR, is available here: https://github.com/aidanfoo96/MINUUR with an accompanying

jupyter books page to run the analysis available here: https://aidanfoo96.github.io/MINUUR/.

Extended Data

Supplementary Figures and Tables are available in the FigShare repository under the title

"Recovery of Metagenomic Data from the *Aedes aegypti* Microbiome Using a Reproducible

Snakemake Pipeline" and can be accessed using the following URL:

https://figshare.com/projects/Recovery_of_Metagenomic_Data_from_the_Aedes_aegypti_Micro

[biome_using_a_Reproducible_Snakemake_Pipeline/158210](https://figshare.com/projects/Recovery_of_Metagenomic_Data_from_the_Aedes_aegypti_Microbiome_using_a_Reproducible_Snakemake_Pipeline/158210). This repository contains the

following data:

Supplementary Table 1: KRAKEN2 Summary Report

Supplementary Table 2: KRAKEN2 Per Sample Genera Classifications

Supplementary Figure 1: Genome Size Comparison Between High, Medium and Low-Quality

MAGs

Supplementary Figure 2: Correlation Between Classified Read Number and Number of MAGs

Supplementary Figure 3: GTDB Classifications vs KRAKEN2 Assigned Taxa > 100,000 Reads

Author Contributions

AF, EH, GLH conceived the project. AF and EH designed the methodology. AF performed the

analyses and wrote the pipeline. LC provided technical expertise. EH and GLH provided oversight

throughout the project. AF and EH drafted the manuscript, and all authors contributed to and

approved the final version.

Grant Information

AF was supported by the DTP scholarship (Medical Research Council MR/N013514/1). GLH was

supported by the BBSRC (BB/T001240/1, BB/V011278/1, and BB/W018446/1), the UKRI (20197

and 85336), the EPSRC (V043811/1), a Royal Society Wolfson Fellowship (RSWF\R1\180013),

the NIHR (NIHR2000907) and the Bill and Melinda Gates Foundation (INV-048598). EH

acknowledges funding from Wellcome (217303/Z/19/Z) and the BBSRC (BB/V011278/1).

References

1. Messina JP, Brady OJ, Pigott DM, Brownstein JS, Hoen AG, Hay SI. A global
compendium of human dengue virus occurrence. Sci Data. 2014 Dec;1(1):140004.

2. Cansado-Utrilla C, Zhao SY, McCall PJ, Coon KL, Hughes GL. The microbiome and
mosquito vectorial capacity: rich potential for discovery and translation. Microbiome.
2021 Dec;9(1):111.

3. Utarini A, Indriani C, Ahmad RA, Tantowijoyo W, Arguni E, Ansari MR, et al. Efficacy
of *Wolbachia*-Infected Mosquito Deployments for the Control of Dengue. N Engl J
Med. 2021 Jun 10;384(23):2177–86.

4. Scolari F, Casiraghi M, Bonizzoni M. *Aedes* spp. and Their Microbiota: A Review.
Front Microbiol. 2019 Sep 4;10:2036.

5. Steven B, Hyde J, LaReau JC, Brackney DE. The Axenic and Gnotobiotic Mosquito:
Emerging Models for Microbiome Host Interactions. Front Microbiol. 2021 Jul
12;12:714222.

6. Bongio NJ, Lampe DJ. Inhibition of *Plasmodium berghei* Development in Mosquitoes
by Effector Proteins Secreted from *Asaia* sp. Bacteria Using a Novel Native
Secretion Signal. PLOS ONE. 2015 Dec 4;10(12):e0143541.

7. Romoli O, Gendrin M. The tripartite interactions between the mosquito, its
microbiota and *Plasmodium*. Parasit Vectors. 2018 Dec;11(1):200.

8. Dada N, Jumas-Bilak E, Manguin S, Seidu R, Stenström TA, Overgaard HJ.
Comparative assessment of the bacterial communities associated with *Aedes*
aegypti larvae and water from domestic water storage containers. Parasit Vectors.
2014;7(1):391.

9. David MR, Santos LMB dos, Vicente ACP, Maciel-de-Freitas R. Effects of
environment, dietary regime and ageing on the dengue vector microbiota: evidence
of a core microbiota throughout *Aedes aegypti* lifespan. Mem Inst Oswaldo Cruz.
2016 Aug 25;111(9):577–87.

10. Saab SA, Dohna H zu, Nilsson LKJ, Onorati P, Nakhleh J, Terenius O, et al. The
environment and species affect gut bacteria composition in laboratory co-cultured
Anopheles gambiae and *Aedes albopictus* mosquitoes. Sci Rep. 2020
Dec;10(1):3352.

11. Onyango GM, Bialosuknia MS, Payne FA, Mathias N, Ciota TA, Kramer DL.
Increase in temperature enriches heat tolerant taxa in *Aedes aegypti* midguts. Sci
Rep. 2020 Dec;10(1):19135.

12. Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J. Dynamic Gut Microbiome across
Life History of the Malaria Mosquito *Anopheles gambiae* in Kenya. PLoS ONE. 2011
Sep 21;6(9):e24767.

13. Sandeu MM, Maffo CGT, Dada N, Njiokou F, Hughes GL, Wondji CS. Seasonal
variation of microbiota composition in *Anopheles gambiae* and *Anopheles coluzzii* in
two different eco-geographical localities in Cameroon. Med Vet Entomol. 2022
Sep;36(3):269–82.

14. Kakani P, Gupta L, Kumar S. Heme-Peroxidase 2, a Peroxinectin-Like Gene,
Regulates Bacterial Homeostasis in *Anopheles stephensi* Midgut. Front Physiol.
2020 Sep 8;11:572340.

15. Minard G, Tran FH, Tran Van V, Fournier C, Potier P, Roiz D, et al. Shared larval
rearing environment, sex, female size and genetic diversity shape *Ae. albopictus*
bacterial microbiota. PLOS ONE. 2018 Apr 11;13(4):e0194521.

16. Kozlova EV, Hegde S, Roundy CM, Golovko G, Saldaña MA, Hart CE, et al.
Microbial interactions in the mosquito gut determine *Serratia* colonization and blood-
feeding propensity. ISME J. 2021 Jan;15(1):93–108.

17. Ganley JG, D'Ambrosio HK, Shieh M, Derbyshire ER. Coculturing of Mosquito-
Microbiome Bacteria Promotes Heme Degradation in *Elizabethkingia anophelis*.
ChemBioChem. 2020 May 4;21(9):1279–84.

18. Hegde S, Khanipov K, Albayrak L, Golovko G, Pimenova M, Saldaña MA, et al.
Microbiome Interaction Networks and Community Structure From Laboratory-
Reared and Field-Collected *Aedes aegypti*, *Aedes albopictus*, and *Culex*
quinquefasciatus Mosquito Vectors. Front Microbiol. 2018 Sep 10;9:2160.

19. Heu K, Romoli O, Schönbeck JC, Ajenoe R, Epelboin Y, Kircher V, et al. The Effect
of Secondary Metabolites Produced by *Serratia marcescens* on *Aedes aegypti* and
Its Microbiota. Front Microbiol. 2021 Jul 7;12:645701.

20. Mitri C, Bischoff E, Belda Cuesta E, Volant S, Ghozlane A, Eiglmeier K, et al.
Leucine-Rich Immune Factor APL1 Is Associated With Specific Modulation of
Enteric Microbiome Taxa in the Asian Malaria Mosquito *Anopheles stephensi*. Front
Microbiol. 2020 Feb 26;11:306.

21. Vogel KJ, Valzania L, Coon KL, Brown MR, Strand MR. Transcriptome Sequencing
Reveals Large-Scale Changes in Axenic *Aedes aegypti* Larvae. PLoS Negl Trop
Dis. 2017 Jan 6;11(1):e0005273.

22. Short SM, Mongodin EF, MacLeod HJ, Talyuli OAC, Dimopoulos G. Amino acid
metabolic signaling influences *Aedes aegypti* midgut microbiome variability. PLoS
Negl Trop Dis. 2017 Jul 28;11(7):e0005677.

23. Stathopoulos S, Neafsey DE, Lawniczak MKN, Muskavitch MAT, Christophides GK.
Genetic Dissection of *Anopheles gambiae* Gut Epithelial Responses to *Serratia*
marcescens. PLoS Pathog. 2014 Mar 6;10(3):e1003897.

24. Muturi EJ, Dunlap C, Ramirez JL, Rooney AP, Kim CH. Host blood meal source has
a strong impact on gut microbiota of *Aedes aegypti*. FEMS Microbiol Ecol. 2019 Jan
1;95(1).

25. Chen S, Zhang D, Augustinos A, Doudoumis V, Bel Mokhtar N, Maiga H, et al.
Multiple Factors Determine the Structure of Bacterial Communities Associated With
Aedes albopictus Under Artificial Rearing Conditions. Front Microbiol. 2020 Apr
15;11:605.

26. Zhao T, Li B qi, Gao H ting, Xing D, Li M jin, Dang Y qi, et al. Metagenome
Sequencing Reveals the Microbiome of *Aedes albopictus* and Its Possible
Relationship With Dengue Virus Susceptibility. Front Microbiol. 2022 May
11;13:891151.

27. Wang Y ting, Shen R xin, Xing D, Zhao C pei, Gao H ting, Wu J hong, et al.
Metagenome Sequencing Reveals the Midgut Microbiota Makeup of *Culex pipiens*
quinquefasciatus and Its Possible Relationship With Insecticide Resistance. Front
Microbiol. 2021 Feb 25;12:625539.

28. Pérez-Cobas AE, Gomez-Valero L, Buchrieser C. Metagenomic approaches in
microbial ecology: an update on whole-genome and marker gene sequencing
analyses. Microb Genomics. 2020 Aug 1;6(8).

29. Hooper R, Brealey JC, Valk T, Alberdi A, Durban JW, Fearnbach H, et al. Host-
derived population genomics data provides insights into bacterial and diatom
composition of the killer whale skin. Mol Ecol. 2019 Jan;28(2):484–502.

30. Ghanavi HR, Twort VG, Duplouy A. Exploring bycatch diversity of organisms in
whole genome sequencing of *Erebidae* moths (*Lepidoptera*). Sci Rep. 2021
Dec;11(1):24499.

31. LaBonte NR, Jacobs J, Ebrahimi A, Lawson S, Woeste K. Data mining for discovery
of endophytic and epiphytic fungal diversity in short-read genomic data from
deciduous trees. Fungal Ecol. 2018 Oct;35:1–9.

32. Salzberg SL, Hotopp JCD, Delcher AL, Pop M, Smith DR, Eisen MB, et al.
Serendipitous discovery of *Wolbachia* genomes in multiple *Drosophila* species.
Genome Biol. 2005;8.

33. Martinson VG, Magoc T, Koch H, Salzberg SL, Moran NA. Genomic Features of a
Bumble Bee Symbiont Reflect Its Host Environment. Appl Environ Microbiol. 2014
Jul;80(13):3793–803.

34. Fierst JL, Murdock DA, Thanthiriwatte C, Willis JH, Phillips PC. Metagenome-
Assembled Draft Genome Sequence of a Novel Microbial *Stenotrophomonas*
maltophilia Strain Isolated from *Caenorhabditis remanei* Tissue. Genome Announc.
2017 Feb 16;5(7).

35. Chen C, Compton A, Nikolouli K, Wang A, Aryan A, Sharma A, et al. Marker-
assisted mapping enables effective forward genetic analysis in the arboviral vector
Aedes aegypti, a species with vast recombination deserts. Genetics; 2022 Nov
1;222(3).

36. Crava C, Varghese FS, Pischedda E, Halbach R, Palatini U, Marconcini M, et al.
Immunity to infections in arboviral vectors by integrated viral sequences: an
evolutionary perspective. http://biorxiv.org/lookup/doi/10.1101/2020.04.02.022509

37. Ford SA, Allen SL, Ohm JR, Sigle LT, Sebastian A, Albert I, et al. Selection on
Aedes aegypti alters *Wolbachia*-mediated dengue virus blocking and fitness. Nat
Microbiol. 2019 Nov;4(11):1832–9.

38. Faucon F, Dusfour I, Gaude T, Navratil V, Boyer F, Chandre F, et al. Identifying
genomic changes associated with insecticide resistance in the dengue mosquito
Aedes aegypti by deep targeted sequencing. Genome Res. 2015 Sep;25(9):1347–
59.

39. The Anopheles gambiae 1000 Genomes Consortium. Genome variation and
population structure among 1142 mosquitoes of the African malaria vector species
Anopheles gambiae and *Anopheles coluzzii*. Genome Res. 2020 Oct;30(10):1533–
46.

40. Köster M. Sustainable data analysis with Snakemake. F1000Research. 2022 Apr
21;10(33).

41. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012 Apr;9(4):357–9.

42. Lu J, Salzberg SL. Ultrafast and accurate 16S rRNA microbial community analysis
using Kraken 2. Microbiome. 2020 Dec;8(1):124.

43. Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species
abundance in metagenomics data. PeerJ Comput Sci. 2017 Jan 2;3:e104.

44. Truong D. T, Franzosa A. E, Tickle T.L, Scholz M, Weingart G, et al. MetaPhlAn2 for enhanced metagenomic taxonomic profiling. Nat Methods. 2015 Oct;12(10):902-3.

45. Andrew, S A S. FASTQC: A Quality Control Tool for High Throughput Sequence Data. www.bioinformatics.babraham.ac.uk/projects/fastqc/

46. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet. 2011 May; 17(1):10-12.

47. Valiente-Mullor C, Beamud B, Ansari I, Francés-Cuesta C, García-González N, Mejía L, et al. One is not enough: On the effects of reference genome for the mapping and subsequent analyses of short-reads. PLOS Comput Biol. 2021 Jan 27;17(1):e1008678.

48. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078–9.

49. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. 2010 Mar 15;26(6):841–2.

50. Hall AB, Basu S, Jiang X, Qi Y, Timoshevskiy VA, Biedler JK, et al. A male-determining factor in the mosquito *Aedes aegypti*. Science. 2015 Jun 12;348(6240):1268-1270.

51. Lee Y, Schmidt H, Collier TC, Conner WR, Hanemaaijer MJ, Slatkin M, et al. Genome-wide divergence among invasive populations of *Aedes aegypti* in California. BMC Genomics. 2019 Dec;20(1):204.

52. Lau MJ, Schmidt TL, Yang Q, Chung J, Sankey L, Ross PA, et al. Genetic stability of *Aedes aegypti* populations following invasion by wMel *Wolbachia*. BMC Genomics. 2021 Dec 14;22(1):894.

53. Rose NH, Dabo S, da Veiga Leal S, Sylla M, Diagne CT, Faye O, et al. Enhanced mosquito vectorial capacity underlies the Cape Verde Zika epidemic. PLOS Biol. 2022 Oct 26;20(10):e3001864.

54. Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 2015 May 15;31(10):1674–6.

55. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

56. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009 Jul 15;25(14):1754–60.

57. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome
assemblies. PeerJ. 2019 Jul 26;7:e7359.

58. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells,
and metagenomes. Genome Res. 2015 Jul;25(7):1043–55.

59. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Bioinformatics. 2019 Nov
15;btz848.

60. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics. 2010 Dec;11(1):119.

61. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, et al.
A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. Nat Biotechnol. 2018 Nov;36(10):996–1004.

62. Finn RD, Clements J, Eddy SR. HMMER web server: interactive sequence similarity
searching. Nucleic Acids Res. 2011 Jul 1;39:W29–37.

63. Matsen FA, Kodner RB, Armbrust EV. pplacer: linear time maximum-likelihood and
Bayesian phylogenetic placement of sequences onto a fixed reference tree. BMC
Bioinformatics. 2010 Dec;11(1):538.

64. Matthews BJ. Improved reference genome of *Aedes aegypti* informs arbovirus
vector control. 2018 Nov;563(7732):501-507.

65. The Genome Standards Consortium, Bowers RM, Kyrpides NC, Stepanauskas R,
Harmon-Smith M, Doud D, et al. Minimum information about a single amplified
genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and
archaea. Nat Biotechnol. 2017 Aug;35(8):725–31.

66. Dada N, Jupatanakul N, Minard G, Short SM, Akorli J, Villegas LM. Considerations
for mosquito microbiome research from the Mosquito Microbiome Consortium.
Microbiome. 2021 Feb 1;9(1):36.

67. Glendinning L, Stewart RD, Pallen MJ, Watson KA, Watson M. Assembly of
hundreds of novel bacterial genomes from the chicken caecum. Genome Biol. 2020
Dec;21(1):34.

68. Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, et al. A new
genomic blueprint of the human gut microbiota. Nature. 2019 Apr
25;568(7753):499–504.

69. Wilkinson T, Korir D, Ogugo M, Stewart RD, Watson M, Paxton E, et al. 1200 high-
quality metagenome-assembled genomes from the rumen of African cattle and their
relevance in the context of sub-optimal feeding. Genome Biol. 2020 Dec;21(1):229.

70. Watson M. New insights from 33,813 publicly available metagenome-assembled-
genomes (MAGs) assembled from the rumen microbiome. Microbiology; 2021 Apr.
http://biorxiv.org/lookup/doi/10.1101/2021.04.02.438222

71. Stewart RD, Auffret MD, Warr A, Wiser AH, Press MO, Langford KW, et al.
Assembly of 913 microbial genomes from metagenomic sequencing of the cow
rumen. Nat Commun. 2018 Dec;9(1):870.

72. Mancini MV, Damiani C, Accoti A, Tallarita M, Nunzi E, Cappelli A, et al. Estimating
bacteria diversity in different organs of nine species of mosquito by next generation
sequencing. BMC Microbiol. 2018 Dec;18(1):126.

73. Coon KL, Hegde S, Hughes GL. Inter-species microbiota transplantation
recapitulates microbial acquisition and persistence in mosquitoes. Microbiology.
2022 Apr 11;10(1):58.

74. Muturi EJ, Njoroge TM, Dunlap C, Cáceres CE. Blood meal source and mixed
blood-feeding influence gut bacterial community composition in *Aedes aegypti*.
Parasit Vectors. 2021 Dec;14(1):83.

75. Chen S, Johnson BK, Yu T, Nelson BN, Walker ED. *Elizabethkingia anophelis*:
Physiologic and Transcriptomic Responses to Iron Stress. Front Microbiol. 2020
May 7;11:804.

76. Onyango MG, Lange R, Bialosuknia S, Payne A, Mathias N, Kuo L, et al. Zika virus
and temperature modulate *Elizabethkingia anophelis* in *Aedes albopictus*. Parasit
Vectors. 2021 Dec;14(1):573.

77. Gaio A de O, Gusmão DS, Santos AV, Berbert-Molina MA, Pimenta PF, Lemos FJ.
Contribution of midgut bacteria to blood digestion and egg production in *Aedes*
aegypti. Parasit Vectors. 2011 Dec;4(1):105.

78. Dahl EM, Neer E, Bowie KR, Leung ET, Karstens L. microshades: An R Package for
Improving Color Accessibility and Organization of Microbiome Data. 2022;11(11).

Figures

623
Figure 1: Study Workflow. Workflow describing the main steps of characterising non-reference reads from *Aedes aegypti* whole genome sequencing data. 62 samples (124 FASTQ files) were input to our metagenomics workflow MINUUR (**M**icrobial **IN**sight **U**sing **U**nmapped **R**eads). Reads were mapped to the *Aedes aegypti* reference genome (AaegL5.3) using Bowtie2 (v2.4.4), and unmapped (=non-reference) reads extracted, parsed for taxonomic classification with KRAKEN2

(v2.1.2) and metagenome assembly with MEGAHIT (v1.2.9). The resulting contigs are indexed
with burrows wheeler aligner (BWA) (v0.7.17) and used to produce a depth file generated using
the jgi_summarize_bam_contig_depth script from MetaBAT2 (v2.12.1). Bins are produced using
the assembled contigs and depth file with MetaBAT2. The resulting metagenome assembled
genomes (MAGs) are quality checked using CheckM (v1.1.3). High and medium quality MAGs,
based on definitions set by the genome standards consortium (65), are taxonomically classified
using GTDB-Tk (v1.5.0). Blue boxes denote the steps within MINUUR, red boxes denote steps
outside of MINUUR. Green starts denote key outputs of the analyses.
637

Study Accession	TaxID	Mosquito Species	Sequencing Platform	Isolation Source	Study Date	Location	DNA Extraction Method	Number of Samples
PRJEB33044	7159	aedes_aegypti	Hiseq_3000	lab	6/14/19	Pennsylvania	TRIzol_Reagent	10
PRJNA255893	7159	aedes_aegypti	HiSeq 2000	lab	28/06/16	Virginia	QiaGEN_blood_tissue_kit	4
PRJNA385349	7159	aedes_aegypti	HiSeq_4000	field	7/1/17	California	MagMAX_protocol	28
PRJNA718905	7159	aedes_aegypti	HiSeq X Ten	lab	4/13/21	Austria	QIAGEN_GenomicTip_DNA	6
PRJNA718905	7159	aedes_aegypti	Illumina NovaSeq 6000	lab	4/13/21	Austria	QIAGEN_GenomicTip_DNA	4
PRJNA776956	7159	aedes_aegypti	Hiseq_2500	field	11/6/21	Australia	QiaGEN_blood_tissue_kit	6
PRJNA882905	7159	aedes_aegypti	Novaseq_6000	lab	10/7/22	Cape_Verde	QIAGEN_Dneasy	15

Figure 2: Sequencing Project Selection and Alignment. **A.** Bar chart depicting the total
number of whole genome sequencing projects referred to as "*Aedes aegypti*" on the European
Nucleotide Archive. Colour bars show the target sequencing samples we used in our study, colour
coded in the legend. **B.** Facetted bar charts showing the total number of reads (right, green),
aligned reads (left, blue) and non-reference reads (middle, dark blue) of our selected samples.
Mapped and non-reference reads result from the alignment to the *AaegL5.3* reference genome
using Bowtie2 (v2.4.4) (41) within our metagenomics workflow MINUUR. **C.** Table showing (left
to right) the original sequencing project, taxonomic id, mosquito species, sequencing platform,
isolation source, study date, location of the sequencing project, DNA extraction method and
number of samples.

Figure 3: KRAKEN2 (v1.2) Genera Classifications of Non-Reference Reads After Alignment to the *AaegL5.3* Reference Genome. **A.** Box plot depicting the number of KRAKEN2 (42) classifications from non-reference reads across 6 publicly available *Aedes aegypti* sequencing projects. Each dot represents one sample within the sequencing project. **B.** Bar chart showing the proportion of genera summed across all samples. **C.** Facetted heatmaps showing phylum and genus level KRAKEN2 assignments of taxa identified from non-reference reads. Colour of each phylum is denoted in the right-hand legend panel, with the most abundant genera depicted by a colour gradient corresponding to their respective phylum. The graph was generated using the microshades R package (78).

Figure 4: Recovered *Aedes aegypti* Associated Bacterial Metagenome Assembled Genomes (MAGs). A. MAGs assembled from non-reference reads using MEGAHIT (v1.2.9) and binned with MetaBAT2 (v2.12.1). Colours denote MAG genome standards consortium (GSC) high, medium and low MAG quality; green = high-quality MAGs (>90% completeness, <5% contamination), grey = medium-quality MAGs (>50% completeness, <5% contamination) and orange = low-quality MAGs. **B.** Bar graph depicting N50 score of medium and high-quality MAGs. Green = high-quality MAG and grey = medium-quality MAGs. **C.** Bar graph depicting genome size in base pairs of medium and high-quality MAGs. Green = high-quality MAG, grey = medium-quality MAG.

Figure 5: Relationship Between KRAKEN2 classifications and MAG Recovery. **A.** Bar chart depicting the recovery of high, medium and low-quality MAGs from *Aedes aegypti* non-reference reads across each sequencing project. Y-axis = number of recovered MAGs, X-axis = MAG quality ranked using CheckM completeness and contamination. **B.** KRAKEN2 classifications depicted through increased stringency of a KRAKEN2 classification threshold. Each line represents a sample within the sequencing project, Y-axis shows the number of distinct KRAKEN2 classified genera and X-axis represents the number of KRAKEN2 classifications. Each graph represents the filtering of samples towards taxa with a high number of assigned reads.

Figure 6: *Aedes aegypti* Associated Bacterial MAG Classifications with GTDB-Tk (v2.1.1). **A.**
Genome size (base pairs) of *Aedes aegypti* associated MAGs (orange), split between high-quality
(CheckM completeness > 90%, contamination <5%) and medium-quality (CheckM completeness
>50%, contamination <5%) rankings, compared to their closest GTDB-Tk reference genome
(blue). Each bar denotes the origin sample and the bin number (sample.bin). **B.** Violin plot
showing the distribution of *Aedes aegypti* MAG genome size (orange) compared to their GTDB-
Tk classified reference genome. Each point denotes a MAG and graphs are facetted between
high and medium quality MAGs (see above). **C.** Taxonomic classifications from the Genome
Taxonomy Database (GTDB) of MAGs obtained from non-reference *Aedes aegypti* reads. X-axis
= FastANI (Average Nucleotide Identity) (%) score to the closest related reference genome in the
GTDB, Y-axis = GTDB-Tk species classification.