Multivalent BARD1-nucleosome interactions facilitate H2A ubiquitylation by BRCA1/BARD1

Samuel R. Witus[1], Lisa M. Tuttle[1], Alex Zelter[1], Klaiten E. Kermoade[1], Damien B. Wilburn[2,3], Weixing Zhao[4], Trisha N. Davis[1], Peter S. Brzovic[1], Rachel E. Klevit[1,*]

1. Department of Biochemistry, University of Washington, Seattle, WA.
2. Department of Genome Sciences, University of Washington, Seattle, WA.
3. Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH.
4. University of Texas Health Science Center at San Antonio, Department of Biochemistry and Structural Biology. San Antonio, TX.
* Corresponding author email: klevit@uw.edu

Abstract

BRCA1/BARD1 is a tumor suppressor E3 ubiquitin (Ub) ligase complex with roles in DNA damage repair and in transcriptional regulation. The N-terminal RING domains of BRCA1/BARD1 interact with nucleosomes to facilitate mono-ubiquitylation of distinct lysine residues on the extreme C-terminal tail of histone H2A. These enzymatic domains constitute only a small fraction of the total sequence of the heterodimer, raising the possibility of functional interactions with chromatin that are mediated through other parts of the protein. Possible interaction sites include the BARD1 C-terminal domains that bind to nucleosomes containing the DNA damage signal H2A K15-Ub and H4 K20me0, as well as portions of the expansive intrinsically disordered regions found in both subunits that have been reported to bind to DNA with unknown functional significance. Using full-length and truncated forms of BRCA1/BARD1 and a collection of designer chromatin substrates, we reveal novel interactions that support robust H2A ubiquitylation activity mediated through a high-affinity, intrinsically disordered DNA-binding region of BARD1. These interactions can be specifically inhibited by the presence of specialized DNA structures that represent intermediates during DNA damage and repair processes. We also identify distinct complexes between BRCA1/BARD1 and chromatin that depend on the presence of H2A K15-Ub, including a complex that spans adjacent nucleosome units. Together, our findings reveal an extensive network of multivalent BARD1-nucleosome interactions that serve as a platform for BRCA1/BARD1-associated functions on chromatin.

Introduction

Mutations in *BRCA1* and *BARD1* increase lifetime risk of developing breast and ovarian cancer. BRCA1 and BARD1 form a large, obligate heterodimeric complex (BRCA1/BARD1) that has distinct roles in DNA double-stranded break (DSB) repair by homologous recombination (HR), transcriptional regulation, and several other nuclear processes (Densham & Morris, 2019; Mullan et al., 2006; Tarsounas & Sung, 2020). These functions are mediated, at least in part, through a direct association with chromatin. Upon DNA damage, BRCA1/BARD1 is recruited to damaged chromatin where it segregates with DNA repair factors (Kolas Nadine K. et al., 2007; Mattiroli et al., 2012; Scully et al., 1997; Sobhian Bijan et al., 2007). Cancer-predisposing mutations that disrupt BRCA1/BARD1 functions on chromatin lead to DNA-damage hypersensitivity and genomic instability(Alenezi et al., 2020; Gudmundsdottir & Ashworth, 2006; Krais & Johnson, 2020). BRCA1/BARD1 also has distinct roles in both stimulation and repression of transcription through as yet incompletely defined mechanisms (Mullan et al., 2006). A full understanding of how BRCA1/BARD1 interfaces with chromatin, where it exerts its major biological functions, is therefore critical.

Nucleosomes, the organizing unit of chromatin, directly interact with BRCA1/BARD1 through multiple direct binding interfaces (Witus et al., 2022). The N-terminal RING domains of BRCA1 and BARD1 form a heterodimer that constitute a RING-type E3 ubiquitin (Ub) ligase and provide the only known enzymatic activity of BRCA1/BARD1 (Brzovic et al., 2001; Lorick et al., 1999; Witus, Stewart, et al., 2021). The RING heterodimer binds to the histone surface of one pseudo-symmetrical 'face' of a nucleosome to facilitate site-specific transfer of mono-Ub to a cluster of lysine residues on the extreme C-terminal tail of canonical histone H2A variants (K125, K127, and K129; collectively referred to henceforth as K127) (Hu et al., 2021; Kalb et al., 2014; Witus, Burrell, et al., 2021). H2A lysine specificity is largely determined by the RING domain of BARD1. This domain binds to a unique histone surface compared to structurally similar Ub ligase

complexes that target other lysine residues on H2A (e.g., K119 by RING1B/BMI1) (McGinty et

al., 2014). The H2A ubiquitylation ("H2A-Ub") activity of BRCA1/BARD1 is thought to contribute

to its function in DNA damage repair via HR by promoting long-range resection of broken DNA

ends (Densham et al., 2016; Uckelmann et al., 2018). BRCA1/BARD1-dependent H2A-Ub

activity is also implicated in the transcriptional repression of certain estrogen-metabolizing

cytochrome P450 genes and the constitutive repression of α-satellite DNA regions (Stewart et

al., 2018; Q. Zhu et al., 2011, 2018).

The BARD1 C-terminal domains (CTDs; **Figure 1A**) also bind to nucleosomes to facilitate DNA

DSB repair via HR (Becker et al., 2021; Dai et al., 2021; Hu et al., 2021; Krais et al., 2021;

Nakamura et al., 2019). The BARD1 CTDs specifically recognize nucleosomes that contain H2A

K15-Ub (the product of the Ub ligase RNF168) and unmethylated H4 K20 (H4K20me0).

Together, these signals serve as a binding platform for the recognition of damaged chromatin in

S/G2 phases when a sister chromatid is available to template high-fidelity DNA repair via HR.

The BARD1 CTDs and N-terminal BRCA1/BARD1 RING domains bind to a fully overlapping

histone surface, precluding their simultaneous binding to one nucleosome face. Despite this,

mono-nucleosomes with ubiquitin pre-installed at H2A K15 are better substrates for

BRCA1/BARD1-dependent ubiquitylation of H2A K127 *in vitro* (Hu et al., 2021). This

observation implies that both domains may bind to the same nucleosome unit, with each

occupying one face. However, a cellular mechanism for histone modification crosstalk between

H2A K15-Ub marks deposited by RNF168 and H2A K127-Ub marks deposited by

BRCA1/BARD1 has yet to be established. Additionally, regions throughout the expansive

intrinsically disordered segments of both BRCA1 and BARD1 are reported to bind to DNA, but

the functional significance of such interactions in chromatin binding and H2A-Ub activity remains

to be determined (Mark et al., 2005; Masuda et al., 2016; Simons et al., 2006; Zhao et al.,

2017).

Although a minimal RING/RING heterodimer is sufficient to direct the transfer of mono-Ub to

H2A K127 in nucleosomes, full-length BRCA1/BARD1 heterodimers form stronger complexes

with unmodified nucleosome substrates and exhibit increased H2A-Ub activity (Hu et al., 2021;

Witus, Burrell, et al., 2021). This suggests that additional interactions between full-length

BRCA1/BARD1 and nucleosomes can occur in the absence of the DNA-damage specific H2A

K15-Ub histone marks. Because BRCA1/BARD1 performs diverse biological functions in the

nucleus including but not limited to DNA repair via HR, transcriptional regulation, stalled

replication fork protection, centrosome regulation, R-loop resolution, and cell-cycle regulation, it

is likely that the complex can be recruited to chromatin via signals not limited to H2A K15-Ub. In

pursuit of a wholistic understanding of BRCA1/BARD1 recruitment to chromatin, we sought to

characterize interactions with nucleosomes that lead to enhanced chromatin binding and H2A-

Ub activity in both the absence and presence of H2A K15-Ub marks.

Here, we identify an intrinsically disordered region of BARD1 adjacent to its RING domain that

strongly binds to both nucleosomal and extra-nucleosomal DNA, dramatically enhancing the

affinity of the complex and its H2A-Ub activity. The interactions can be modulated by specialized

DNA structures that are derived from the DNA damage repair process and can compete directly

with BARD1 binding to nucleosomal DNA. We incorporate our findings into a molecular

mechanism of recognition. Additionally, we provide evidence for multiple, distinct higher-order

chromatin complexes that contain H2A K15-Ub nucleosomes. These include the presence of

both a 'wrapped' complex with the BRCA1/BARD1 RING domains and BARD1 CTDs bound to

opposite sides of one nucleosome unit, as well a complex where these domains bind to

adjacent nucleosome units. Throughout, we evaluate the contribution of BARD1 DNA binding to

nucleosome affinity and H2A-Ub activity. Together, our findings reveal an extensive network of

multivalent BARD1-nucleosome interactions that serve as a platform for BRCA1/BARD1-

associated functions on chromatin.

Results

Identification of a BARD1 region that supports increased H2A ubiquitylation activity

BRCA1/BARD1 is a large, obligate heterodimeric complex with distinct functional domains and

long intrinsically disordered regions (IDRs**; Figure 1A)**. A minimal complex composed of the two

N-terminal RINGs of BRCA1 (1-110) and BARD1 (26-140; BRCA1^{RING}/BARD1^{RING}) is sufficient

to direct site-specific mono-ubiquitylation of nucleosomal histone H2A at K127 (Kalb et al.,

2014). We and others previously reported that full-length BRCA1/BARD1 (~300 kDa) exhibits

faster H2A-Ub kinetics and stronger binding to mono-nucleosomes than the minimal

RING/RING complex (~25 kDa; **Figure 1—figure supplement 1A-C**) (Hu et al., 2021; Witus,

Burrell, et al., 2021). To uncover regions of BRCA1 or BARD1 responsible for increased H2A-

Ub activity and binding affinity, we generated a series of heterodimers with truncations in one or

the other subunit **(Figure 1—figure supplement 2 and Extended Data Table I)**.

To measure contributions from the BRCA1 subunit, heterodimers containing full-length BARD1

and three different BRCA1 constructs were compared: full-length BRCA1 (BRCA1^{FL}), a clinically

relevant allele missing ~1100 residues in the expansive IDR region (BRCA1$^{\Delta 11q}$), and the

minimal BRCA1 RING fragment **(**BRCA1^{RING}**)**. All three heterodimers that contained full-length

BARD1 exhibited faster H2A-Ub kinetics than the minimal BRCA1^{RING}/BARD1^{RING} complex

(Figure 1—figure supplement 1C). Complexes containing truncated BRCA1 (BRCA1^{RING} and

BRCA1$^{\Delta 11q}$) exhibited faster H2A-Ub kinetics than the full-length heterodimer. In contrast,

BARD1^{RING}-heterodimers containing BRCA1^{FL} or BRCA1$^{\Delta 11q}$ exhibited diminished H2A-Ub

activity compared to the full-length complex **(Figure 1—figure supplement 1D)**. Together these

observations suggest that there may be inhibitory elements or alternative interactions involving

regions of BRCA1 in the full-length complex. Overall, the data indicate that BARD1 is

predominantly responsible for the enhanced H2A-Ub activity that is observed in the full-length

BRCA1/BARD1 complex.

To identify regions of BARD1 that contribute to increased H2A-Ub activity, the H2A-Ub kinetics

and nucleosome binding affinity of BRCA1^{RING}-heterodimers containing BARD1 truncations

were compared. Notably, BRCA1^{RING}/BARD1^{FL} heterodimers bind nucleosomes with ~1000-fold

higher affinity than the minimal BRCA1^{RING}/BARD1^{RING} complex and had markedly higher H2A-

Ub activity **(Figure 1B, C)**. Removal of the BARD1 Ank-BRCT C-terminal domains (BARD1$^{\Delta CTD}$)

or truncation back to residue 216 in the putative intrinsically disordered region (BARD1^{216}) had

minimal impact on H2A-Ub activity and nucleosome binding **(Figure 1B, C)**. However,

truncation to BARD1 residue 193 (BARD1^{193}) caused a substantial decrease in H2A-Ub activity

and a ~10-fold reduction in binding affinity, implying a role for residues between 194 and 216.

Consistent with this, deletion of the first half of the BARD1 IDR (BARD1$^{\Delta 140-270}$) greatly

diminished H2A-Ub activity **(Figure 1—figure supplement 1E)**. A more targeted deletion

(BARD1$^{\Delta 194-216}$) only modestly decreased H2A-Ub activity, suggesting the presence of additional

or compensatory nucleosome interaction sites in BARD1. Although deletion of residues 140-270

decreased nucleosome binding affinity by ~3-fold relative to full-length BARD1, the affinity is still

considerably stronger than the minimal RING/RING heterodimer or BARD1^{193} truncations

(Figure 1—figure supplement 1F). To assess intrinsic nucleosome-binding function of the

BARD1 IDR, constructs containing only either the N-terminal half (residues 124-270) or the C-

terminal half (residues 269-424) were analyzed. Each IDR-only construct exhibited nucleosome

binding by EMSA **(Figure 1—figure supplement 1G).** The apparent dispensability of BARD1

IDR residues past 216 in H2A-Ub activity may indicate a requirement for specific types of

interactions that our binding assay is not sensitive to **(Figure 1B)**. In total, a region within the

IDR of BARD1 (BARD1 194-216) was identified that is important for productive nucleosome

binding that supports increased H2A-Ub activity. Importantly, these data also demonstrate that

the BARD1 CTDs are dispensable for mono-nucleosome binding and for H2A-Ub activity on

nucleosomes that lack H2A K15-Ub marks.

Additional BARD1-nucleosome interactions are mediated through DNA binding

Having identified non-RING regions of BARD1 that contribute to increased H2A-Ub activity and

nucleosome binding, we sought to characterize the molecular interactions involved. BARD1

residues 141-216 are in a large internal segment predicted to be intrinsically disordered (124-

424) (McGuffin et al., 2000). To our knowledge, whether this region of BARD1 is fully disordered

has not been explicitly tested. Nuclear magnetic resonance (NMR) $^{1}H^{15}N$-HSQC spectra of ^{15}N-

labelled BARD1 fragments containing residues 124-270, 269-424, and 124-424 all exhibited

narrow ^{1}H chemical shift dispersion that is consistent with an IDR **(Figure 2—figure**

supplement 1A, B). Furthermore, the spectra corresponding to the two halves (124-270 and

269-424) overlay well with the spectrum of the full region, indicating that the two halves behave

independently of one another **(Figure 2—figure supplement 1B)**.

The BARD1 region that enhances nucleosome binding and H2A-Ub activity identified above is

contained within the first half of the IDR, spanning residues 124-270. This region has been

reported to bind to double-stranded DNA and specialized DNA structures containing base-pair

mismatch (discussed below) (Zhao et al., 2017). The sequence of BARD1 124-270 is largely

conserved across orthologs and its amino acid content is strongly basic **(Figure 2—figure**

supplement 1C). We hypothesized that observed contributions to H2A-Ub and nucleosome

binding activity from the disordered region outside the RING domain (BARD1 residues 141-216)

could be mediated through interactions with nucleosomal DNA. Several nucleosome-modifying

enzymes such as the methyltransferases DOT1L and COMPASS contain DNA-binding regions

adjacent to their catalytic domains that are required for full activity (Hsu et al., 2019; McGinty &

Tan, 2016, 2021; Worden et al., 2019, 2020). Indeed, we observed strong binding of BARD1

124-270 to nucleosomes and to free 147-bp '601' dsDNA by EMSA **(Figure 2—figure**

supplement 2A).

For residue-level information, we performed NMR binding experiments with ^{15}N-labelled BARD1

124-270 and either a 36-bp dsDNA fragment or a 147-bp nucleosome **(Figure 2A and Figure**

2—figure supplement 2B, respectively). Between these two titrants, only a full titration of the

dsDNA fragment was possible due to its smaller molecular weight than a nucleosome and

complex solubility considerations. The resulting spectral series identified a subset of resonances

that shift continuously as a function of DNA concentration **(Figure 2A)**. A similar set of

resonances were perturbed (broadened in this case) upon addition of 147-bp nucleosome,

indicating that BARD1 employs similar residues to recognize nucleosomes and free dsDNA

(Figure 2—figure supplement 2B). To overcome spectral overlap in the spectrum of BARD1

124-270, a smaller construct encompassing the putative DNA binding region identified in H2A-

Ub activity and nucleosome binding assays (BARD1 141-216, where residue 141 is the first to

extend beyond BARD1^{RING}) was assigned. Assignments from the smaller construct could be

transferred to the longer segment, enabling binding analysis by NMR in both contexts **(Figure**

2A, B and Figure 2—figure supplement 2B, C). Secondary structure propensity (SSP)

predicted from the resonance assignments are consistent with the region being primarily

disordered, with helical propensity predicted for residues 191-201 and a weaker prediction of

extended structure towards the beginning of the construct **(Figure 2—figure supplement 3A).**

The largest backbone amide chemical shift perturbations (CSPs) due to dsDNA addition are

clustered in BARD1 residues 194-216, with smaller perturbations observed for residues 146-161

(Figure 2B). As well, ^{15}N dynamics measurements reveal that residues 146-161 undergo

changes in dynamics in the presence of dsDNA **(Figure 2—figure supplement 3B)**. Predicted

secondary structure from chemical shifts of BARD1 in the presence of 1 molar equivalent of

dsDNA indicate reduced helical propensity in residues 191-199 upon DNA binding **(Figure 2—**

figure supplement 3A). Importantly, residues between 162-191 of BARD1 were relatively

unaffected by addition of dsDNA, consistent with the existence of specific DNA-binding regions.

Due to solubility and concentration limitations, we were unable to perform similar analysis on

nucleosome-bound samples. Nevertheless, the similar ^{1}H^{15}N-HSQC spectral perturbations

observed in the presence of low molar equivalents of nucleosomes or dsDNA suggest similar

binding modes (**Figure 2—figure supplement 2B)**. Thus, the data identify two regions of

BARD1 (146-161 and 194-216) in the IDR adjacent to the enzymatic RING domains that bind

DNA when presented as free dsDNA or in the context of a nucleosome. The identified regions

are consistent with the H2A-Ub activity and nucleosome binding data presented above.

Having established that isolated BARD1 IDR fragments bind to free DNA and nucleosomes

similarly by NMR, we sought to observe the interactions within the context of an active

enzymatic complex. As such a complex would prove too large for NMR analysis, we turned to

targeted crosslinking of BRCA1/BARD1 to nucleosomal DNA. A photoactivatable crosslinker p-

Benzoyl-L-phenylalanine (Bpa) was incorporated using amber-codon suppression in *E. coli* at

individual locations throughout the BARD1 IDR in the context of RING-containing constructs

(L120Bpa, W146Bpa, Y180Bpa, A195Bpa and W218Bpa**; Figure 2—figure supplement 4A)**.

Due to the need to co-express a Bpa-specific aminoacyl tRNA transferase, we designed a

genetic fusion in which BRCA1 residues 1-104 are fused to BARD1 residues 26-221 through a

12-residue GlySer-linker (BRCA1-f-BARD1^{221}). Wild-type BRCA1-f-BARD1^{221} and Bpa-

incorporated mutants retained similar H2A-Ub activity to each other and to unfused

heterodimers **(Figure 2—figure supplement 4B)**. A nucleosome substrate with extra-

nucleosomal linker DNA that carried a fluorescent probe was used for crosslinking experiments

(NCP[185]; **Figure 2—figure supplement 4C)**. SDS-PAGE analysis of crosslinking reaction

products revealed shifted bands consistent with a covalent DNA-protein adduct for all Bpa

variants **(Figure 2C)**. Quantitation revealed enhanced crosslinks for the Bpa variants at BARD1

W146 and A195 positions, consistent with the NMR dsDNA binding data presented above

(Figure 2C). Weaker crosslinks from the Y180 and W218 positions were comparable to

crosslinks observed from L120 that is located close to the RING domains and not expected to

contribute to productive DNA interactions. A similar crosslinking pattern was observed in

reactions containing free DNA **(Figure 2—figure supplement 4D)**. Together, the data support a

direct interaction between the BARD1 IDR and nucleosomal DNA in the context of an

enzymatically active complex.

Contribution of DNA-binding to enhanced H2A-Ub activity

Data presented thus far establish that the DNA-binding region of the BARD1 IDR is necessary

for high H2A-Ub activity, but is it sufficient? To address this question, we asked whether the

DNA-binding properties of full-length BARD1 can support H2A-Ub activity in the presence of

highly disruptive BARD1 RING histone-interface mutations **(**BARD1 P89A/W91A; **Figure 3A)**

(Hu et al., 2021; Witus, Burrell, et al., 2021). Although high-affinity nucleosome binding and

auto-ubiquitylation activity were retained, heterodimers containing full-length BARD1

(P89A/W91A) do not possess H2A-Ub activity **(Figure 3B, Figure 3—figure supplement 1A,**

B). They also do not transfer Ub to other (off-target) histone sites **(Figure 3—figure**

supplement 1C). Thus, the RING-histone interface is absolutely required for H2A-Ub activity,

while the DNA-binding properties of the BARD1 IDR serve to promote the formation of a high-

affinity complex that leads to enhanced H2A-Ub activity.

The disordered nature of the BARD1 DNA-binding region(s) raises the question of whether its

sequence identity is critical to support increased H2A-Ub activity. In other words, can any DNA-

binding region placed adjacent to the RING domain serve the same purpose? We generated a

chimeric BARD1 construct in which a previously identified DNA-binding region from the BRCA1

IDR (498-663) was appended onto the BARD1 RING (Mark et al., 2005). A heterodimer

composed of the BRCA1 RING and the BARD1 chimera supported increased H2A-Ub activity

as compared to the minimal RING/RING complex, but not to the levels of the native BARD1

sequence **(Figure 3C)**. The difference in activity is unlikely to be due solely to a difference in

DNA-binding affinity of the native BARD1 region versus the installed BRCA1-derived region, as

the measured affinities for nucleosomes are quite similar **(Figure 3—figure supplement 1D)**.

The small differences notwithstanding, the findings indicate that a major contributor to increased

H2A-Ub activity is the presence of a disordered, high-affinity DNA-binding region near the

enzymatic RING domains.

Specialized DNA structures bind to the BARD1 IDR and inhibit H2A-Ub activity

The BARD1 DNA-binding region enhances BRCA1/BARD1-mediated H2A-Ub activity by

interacting with nucleosomal DNA and increasing the affinity of the complex. It was previously

reported that this region of BARD1 preferentially binds to partly unwound DNA structures

(bubble-DNA) and homologous recombination (HR) intermediates (D-loop DNA) over double- or

single-stranded DNA (Zhao et al., 2017). The interactions with specialized DNA structures help

to promote HR through assembly of a functional complex consisting of BARD1, the

recombinase RAD51, and DNA. We hypothesized that the interaction between the BARD1

DNA-binding region and bubble- or D-loop DNA may alter its ability to interact with nucleosomal

DNA and consequently affect nucleosome binding and/or H2A-Ub activity. The molecular

features that BARD1 recognizes in bubble- and D-loop- DNA substrates are likely similar based

on the near-identical affinities of the BARD1-DNA complexes (Zhao et al., 2017), so we chose

to use the simpler bubble-DNA substrate in our assays.

Equal concentrations of DNA fragments containing various patterns of base-pair mismatch were

added to nucleosome ubiquitylation reactions carried out by full-length BRCA1/BARD1 **(Figure**

4A, B and Figure 4—figure supplement 1A, B). Notably, among the DNA species tested, only

bubble-DNA substantially inhibited H2A-Ub activity. We hypothesized that the observed

inhibition is mediated through competing interactions between bubble-DNA and nucleosomal

DNA for binding to BARD1 141-216. Consistent with this notion, the minimal RING heterodimer

exhibits no sensitivity of its H2A-Ub activity to bubble-DNA **(Figure 4—figure supplement 1C,**

D). Further support for a competitive binding mechanism is provided from UV-induced Bpa

crosslinking reactions that contained Bpa variants of BRCA1^{RING}-f-BARD1^{221}, nucleosomes, and

increasing concentrations of either bubble-DNA or dsDNA **(Figure 4C and Figure 4—figure**

supplement 2). Both types of DNA structure inhibited crosslink formation in a concentration-

dependent manner, but the bubble-DNA was substantially more effective.

To characterize the interactions between BARD1 141-216 and bubble-DNA, NMR spectra of

BARD1 IDR-DNA complexes bound to bubble-DNA and dsDNA fragments of similar sequence

and size were compared. CSPs and changes in dynamics within BARD1 194-216, the region we

previously showed to be the major site of dsDNA and nucleosomal DNA binding, were similar

between the bubble-DNA and dsDNA complexes **(Figure 4D and Figure 4—figure**

supplement 3A-C). In addition, specific effects were produced by bubble-DNA: backbone

amide NMR resonances of BARD1 residues 143-157 exhibited severe broadening, with many

signals disappearing, indicative of large changes in chemical environment and/or dynamics. A

notable exception to the pattern of signals broadening beyond detection in this region is a large

CSP for the sidechain resonance of W146 that was observed only in the presence of bubble-

DNA **(Figure 4—figure supplement 3A, B)**. Further support for bubble-DNA-specific effects

comes from $^1H^{13}C$-HSQC spectra of BARD1 141-216. In the region of $^1H^{13}C$-HSQC spectra that

contains methyl group resonances, signals from residues within BARD1 143-157 exhibit large

CSPs that are not observed in the spectrum of dsDNA-bound BARD1 **(Figure 4E and Figure**

4—figure supplement 3D). The data suggest that bubble-DNA binds to BARD1 194-216

similarly to dsDNA, but also strongly and specifically engages an additional DNA-binding site

near the beginning of the BARD1 IDR, namely residues 143-157. Consistent with this finding,

concomitant deletion of basic residues in both IDR regions (BARD1$^{\Delta 150-155/\Delta 194-216}$) was required

to alleviate H2A-Ub activity inhibition by bubble-DNA **(Figure 4F and Figure 4—figure**

supplement 1E). Based on these findings, we conclude that BARD1 specifically recognizes

bubble-DNA via BARD1 residues 143-157 and that interactions with bubble-DNA can compete

with BARD1 IDR binding to nucleosomal DNA. This may lead to the observed repression of

BRCA1/BARD1 H2A-Ub activity in the presence of bubble-DNA competitor fragments.

Extra-nucleosomal linker DNA enhances H2A ubiquitylation

The extended domain topology of both BRCA1 and BARD1 raises the possibility of additional

interactions in higher-order chromatin substrates that are not present in minimal mono-

nucleosome substrates. For example, chromatin factors such as PRC2, LSD1, and VRK1

require linker DNA or adjacent nucleosome units for optimal binding and enzymatic activity

(Budziszewski et al., 2022; Kim et al., 2015; Poepsel et al., 2018; Wang et al., 2017). To test for

additional chromatin substrate requirements for BRCA1/BARD1-dependent H2A ubiquitylation,

nucleosome substrates with extra-nucleosomal linker DNA (NCP^{185}) and tri-nucleosomes (tri-

NCP) were used as substrates in H2A-Ub assays **(Figure 5A and Figure 5—figure**

supplement 1A). In reactions performed using full-length BRCA1/BARD1, faster H2A-Ub

kinetics were observed for the NCP^{185} substrate with linker DNA compared to the minimal

NCP^{147} substrate **(Figure 5B)**. Notably, no additional increase in H2A-Ub activity was observed

for a tri-NCP substrate. The data are consistent with two conclusions: 1) the presence of linker

DNA enhances H2A-Ub activity and 2) functional interactions are limited to one mono-

nucleosome unit, at least in the absence of histone PTMs.

To confirm that the enhanced activity observed for the NCP[185] and tri-NCP substrates described

above stem from the BARD1 IDR DNA-binding region, the H2A-Ub activity of the

BRCA1[RING]/BARD1[216] truncation was tested **(Figure 5C)**. This species exhibited considerably

faster H2A-Ub kinetics on either a NCP[185] or a tri-NCP substrate compared to a minimal NCP[147]

substrate, again consistent with a dependence on the presence of linker DNA. To ask whether

the activity difference between the nucleosomes with and without linker DNA was due to

differences in binding affinity, we measured nucleosome binding affinity at two different ionic

strengths using the NCP[147] and NCP[185] substrates **(**100 mM vs. 150 mM NaCl; **Figure 5—figure**

supplement 1B). Although nucleosome complexes exhibited considerably weaker binding (~5-

fold reduction) at higher buffer salt concentration, there was only a modest increase in binding

affinity with the NCP[185] substrate at both ionic strengths. This behavior contrasts with other

chromatin factors that exhibit affinity and activity differences on the order of magnitudes in the

presence or absence of linker DNA and may reflect the different types of interactions (i.e.,

ordered vs disordered DNA-binding regions) (Kim et al., 2015; Wang et al., 2017, p. 2).

Together, our results indicate that the BARD1 DNA-binding region association with either

histone-associated nucleosomal DNA or extra-nucleosomal linker DNA leads to similar apparent

binding affinity. However, interactions with linker DNA support an additional enhancement of

H2A-Ub activity.

H2A K15-Ub-mediated enhancement of BRCA1/BARD1 H2A-Ub activity

In addition to the RING domains and the BARD1 DNA-binding region, the ordered C-terminal

domains (CTDs) of BARD1 bind to nucleosomes that contain both H2A K15-Ub and H4K20me0

(Dai et al., 2021; Hu et al., 2021). The BARD1 CTDs-nucleosome interactions serve to recruit

BRCA1/BARD1 to damaged chromatin to facilitate DNA DSB repair via HR (Becker et al., 2021;

Krais et al., 2021; Nakamura et al., 2019). The BARD1 CTDs use a histone-binding surface that

overlaps with the RING domain site, precluding binding of both to the same nucleosome face.

However, it was previously shown that nucleosomes containing H2A K15-Ub and H4K20me0

form a stronger complex with BRCA1^{RING}/BARD1^{FL} and are better substrates for ubiquitylation of

H2A K127, thus implying that the RING domains and CTDs can simultaneously bind to opposite

faces of a nucleosome (Hu et al., 2021). We sought to further characterize the complexes using

our suite of BRCA1/BARD1 truncation constructs and chromatin substrates.

To ascertain the effects of pre-existing mono-Ub at H2A positions known to be ubiquitylated by

different ligases, homogenously mono-ubiquitylated nucleosomes containing ubiquitin

preinstalled at H2A positions K15, K119, or K127 were generated using a non-hydrolyzable

dichloroacetone crosslinker **(Figure 6—figure supplement 1A, B)** (Morgan et al., 2019). In

agreement with previous observations, ubiquitin preinstalled at H2A K15 uniquely increased the

H2A-Ub activity of full-length BRCA1/BARD1 **(Figure 6—figure supplement 1C, D)**. Ubiquitin

preinstalled at either H2A C-terminal position (K119 or K127) decreased BRCA1/BARD1-

dependent H2A-Ub activity, hinting at E3 exclusivity between BRCA1/BARD1 and RING1/PCGF

complexes that modify the H2A C-terminal tail. As expected, the BARD1 CTDs were required

for enhancement of BRCA1/BARD1-dependent H2A-Ub activity using H2A K15-Ub nucleosome

substrates **(Figure 6A)**. Importantly, this contrasts with our findings for unmodified nucleosome

substrates where the BARD1 CTDs are not required for enhanced H2A-Ub activity.

Are the DNA-binding properties of BARD1 important in the context of H2A K15-Ub nucleosome

substrates? Ubiquitylation reactions were carried out on H2A K15-Ub nucleosomes with a

selection of BARD1 truncation species. The internal BARD1 IDR deletion construct

(BRCA1^{RING}/BARD1$^{\Delta140-270}$) exhibited decreased H2A-Ub kinetics compared with the

BRCA1^{RING}/BARD1^{FL} complex **(Figure 6A)**. Deletion of BARD1 140-270 still supported greater

activity than removal of the BARD1 CTDs (BRCA1^{RING}/BARD1$^{\Delta CTD}$). This indicates that BARD1-

DNA interactions are secondary to BARD1 CTDs-nucleosome binding on this substrate or can

be otherwise compensated by additional DNA-binding regions of BARD1. Additionally, the

histone-binding BARD1 RING mutant BRCA1^{RING}/BARD1$^{FL-P89A/W91A}$ does not support

BRCA1/BARD1-dependent H2A-Ub activity on H2A K15-Ub nucleosomes, supporting a model

where the BRCA1/BARD1 RINGs and CTDs bind to either face of a nucleosome. These

findings support a 'wrapped' complex model in which three points of contact between BARD1

and H2A K15-Ub nucleosomes (RING-histone, IDR-DNA, and CTDs-histone/Ub) are required

for full BRCA1/BARD1-dependent H2A-Ub activity **(Figure 7E)**.

To obtain information regarding complexes formed on unmodified mono-nucleosomes versus

H2A K15-Ub-modified nucleosomes, chemical crosslinking experiments were performed using

the amine-reactive crosslinkers DSS and BS3 **(Figure 6—figure supplement 2)**. To focus on

the interactions involving the BARD1 subunit, BRCA1^{RING}/BARD1^{FL} was used; products formed

in the crosslinking reactions were analyzed by mass spectrometry. Numerous intermolecular

crosslinks (judged by unique crosslinks and peptide spectral mapping counts) were detected

between the BARD1 CTDs and histones from nucleosomes containing H2A K15-Ub, while such

crosslinks were sparse in complexes composed of unmodified nucleosomes **(Figure 6B and**

Figure 6—figure supplement 2C). The crosslinks are consistent with high-resolution structures

of the BARD1 CTDs bound to nucleosomes containing H2A K15-Ub and the lack of these in the

unmodified nucleosome reaction is consistent with the interaction being dependent on the

presence of Ub at H2A K15 **(Figure 6—figure supplement 2D)**. There was also increased

crosslinking between BARD1 RING and IDR domains to histones in samples containing H2A

K15-Ub nucleosomes **(Figure 6B, gray crosslinks and Figure 6—figure supplement 2C, E)**,

consistent with the reported higher binding affinity of that complex. Thus, binding of the BARD1

CTDs in response to the presence of H2A K15-Ub facilitates BARD1 RING and IDR interactions

that lead to the observed increase in H2A K127-Ub activity.

The BARD1 N-terminal RING domain and the CTD are separated by ~300 intrinsically

disordered residues, allowing for as much as 1000 Å of separation between the two ordered

domains. It is therefore possible that in addition to a 'wrapped' complex with H2A K15-Ub

nucleosomes, the BARD1 CTDs may anchor the heterodimeric complex by binding to one

nucleosome unit, while the RING domains are recruited to a nearby nucleosome unit on

chromatin to facilitate H2A-Ub activity. To test this possibility explicitly, asymmetric di-

nucleosome substrates (di-NCPs) where one nucleosome unit contained H2A K15-Ub (H2A^{silent})

and an adjacent nucleosome unit contained unmodified H2A with a fluorophore conjugated near

its N-terminus to facilitate specific detection in a nucleosome ubiquitylation assay were

assembled **(**H2A^{observe}; **Figure 6C and Figure 6—figure supplement 1E-G)**. Using

BRCA1^{RING}/BARD1^{FL} heterodimers, we observed that an asymmetric di-NCP substrate

containing H2A K15-Ub as H2A^{silent} uniquely facilitates enhanced H2A-Ub modification of an

adjacent nucleosome unit containing unmodified H2A **(**H2A^{observe}; **Figure 6D and Figure 6—**

figure supplement 1H). Removal of the BARD1 CTDs led to a large decrease in H2A-Ub

activity that was comparable to that of a di-NCP substrate where both nucleosome units contain

unmodified H2A. Consistent with observations made using H2A K15-Ub mono-nucleosomes,

BRCA1^{RING}/BARD1$^{FL-P89A/W91A}$ did not support H2A-Ub activity using asymmetric di-NCPs

(Figure 6E). Additionally, deletion of BARD1 DNA-binding residues (Δ140-270) substantially

decreased H2A-Ub activity. The results support the formation of a higher-order chromatin

complex where BARD1 spans adjacent nucleosome units. The complex is mediated through

binding of the BRCA1/BARD1 RING domains and BARD1 CTDs to neighboring nucleosome

units and of the BARD1 IDR to intervening linker or nucleosomal DNA **(Figure 7F)**.

<u>Discussion</u>

Here, we describe distinct complexes between BRCA1/BARD1 and chromatin substrates that

depend on the status of histone PTMs on nucleosomes (e.g., unmodified or H2A K15-Ub)

(Figure 7). Importantly, although the BARD1 CTDs are critical for enhanced interactions with

nucleosomes that contain H2A K15-Ub, they are dispensable for high-affinity interactions with

unmodified nucleosomes **(Figure 7C, D)**. Instead, additional interactions that enhance H2A-Ub

activity are mediated through a short and highly basic intrinsically disordered region of BARD1

(primarily residues 194-216) adjacent to the enzymatic RING domains that engages

nucleosomal and extra-nucleosomal linker DNA. These strong, electrostatic-based interactions

increase the binding affinity of the complex from the micromolar to the nanomolar binding

regime that is likely critical to support the physiological functions of BRCA1/BARD1 on

chromatin. Our findings indicate that DNA binding plays an auxiliary role in the ability of

BRCA1/BARD1 to ubiquitylate H2A by boosting the affinity of the Ub ligase/nucleosome

complex. Supporting this, a BARD1 RING-domain mutant (BARD1 P89A/W91A) that disrupts

histone binding is unable to support H2A-Ub activity with any chromatin substrate tested despite

retaining strong nucleosome binding affinity and intrinsic Ub ligase activity. This indicates that

high-affinity binding to nucleosomal DNA by the BARD1 IDR is not sufficient for H2A-Ub activity,

while the positioning of the RING domains on the histone surface is absolutely required.

Therefore, BARD1 P89A/W91A may constitute a separation-of-function mutant for cellular

studies to investigate the specific roles of BRCA1/BARD1-dependent H2A-Ub.

BRCA1/BARD1 functions in many distinct chromatin-associated complexes and also binds to

nucleic acids (Mark et al., 2005; Savage & Harkin, 2015; Zhao et al., 2017). Highlighting the

importance of other BRCA1/BARD1 binding factors, we identified specific DNA structures

(bubble-DNA) that inhibit nucleosome binding and H2A-Ub activity by binding to the BARD1 IDR

and competing with nucleosome binding. The recognition mechanism is likely similar for D-loop

DNA structures that are intermediates in HR (Zhao et al., 2017), and possibly R-loop structures

that have similar chemical properties (Santos-Pereira & Aguilera, 2015). It is possible that

binding of the BARD1 IDR region to either nucleosomal DNA or specialized nucleic acid

structures serves to control its spatial and temporal involvement in HR and other processes

where such specialized DNA structures may be present (e.g., stalled replication fork protection,

R-loop resolution) (Hatchi et al., 2015; Schlacher et al., 2012). Additionally, BRCA1/BARD1 is

present in higher-order protein assemblies that are known to bind to chromatin via distinct

nucleosome interfaces that may be formed by other proteins in the complex (Belotserkovskaya

et al., 2020, p. 2; Bochar et al., 2000; Huen et al., 2010; Savage & Harkin, 2015). Understanding

the combined binding contributions of BRCA1/BARD1 and other proteins in such complexes to

nucleosome association and H2A-Ub activity is an important future goal.

BRCA1/BARD1 nucleosome binding and H2A-Ub activity is also responsive to pre-existing

histone PTMs, as in the case of H2A K15-Ub. In addition to a wrapped complex where the

BARD1 CTDs and BRCA1/BARD1 RINGs occupy opposite nucleosome faces, we provide

evidence for a complex where the BARD1 CTDs and the enzymatic RING domains occupy

adjacent nucleosome units **(Figure 7E, F)**. The BARD1 IDR region enables extensive

separation between the enzymatic RING domains and BARD1 CTDs while forming high-affinity

interactions with nucleosomal DNA. This may allow the RING domains to reach nucleosome

units considerably further away than an adjacent unit while anchored to chromatin via the

BARD1 CTDs. This ability could lead to BRCA1/BARD1-dependent H2A-Ub of multiple

nucleosomes in the vicinity of a H2A K15-Ub anchor nucleosome. Additionally, the ability to

reach across nucleosome units in chromatin provides a platform to scaffold unique higher-order

complexes across long molecular distances. As H2A K15-Ub is a highly specific mark of DNA

damage, it is unlikely to serve as a universal recruitment signal for other BRCA1/BARD1-

dependent processes such as transcriptional regulation. An important unanswered question is

how other histone PTMs influence the chromatin binding and H2A-Ub activity of

BRCA1/BARD1. Previously, we observed that methylation at H3 K79, a mark associated with

actively transcribed chromatin, inhibits BRCA1/BARD1-mediated H2A-Ub activity (Witus,

Burrell, et al., 2021). This may be attributable to the location of H3 K79 near the BARD1-histone

interface. A comprehensive understanding of histone PTMs that influence BRCA1/BARD1

nucleosome binding, higher-order chromatin complex formation, and H2A-Ub activity will

provide valuable insight into its functions and regulation in different biological processes.

In sum, our results establish an extensive multivalent network of interactions that facilitate

BRCA1/BARD1 chromatin binding and H2A-Ub activity. These interactions are likely critical for

many functions of BRCA1/BARD1 in the nucleus that are not limited to its H2A-Ub activity and

may be disrupted by mutations that cause cancer.

Ideas and Speculation:

Because BRCA1/BARD1 serves as a large protein interaction hub that forms many distinct

complexes, it is possible that the accessibility of DNA-binding regions in BARD1 (and perhaps in

BRCA1) is determined by the landscape of binding partners with which it is associated. On this

point, we observed that full-length BRCA1/BARD1 has lower H2A-Ub activity than complexes

containing truncated BRCA1 (Δ11q or RING). This is despite the presence of additional DNA-

binding regions within the BRCA1 IDR that can support increased H2A-Ub activity. A possible

explanation is a competition among the various DNA-binding regions from BRCA1 and BARD1

for binding to nucleosomes, with binding of the BRCA1 region resulting in lower H2A-Ub activity.

Alternatively, possible inhibitory interactions between BRCA1 and BARD1 subunits might be

alleviated upon BRCA1 truncation. Occupancy of different regions of BRCA1 and BARD1 by

unique binding partners may occlude certain nucleosome-binding regions, resulting in inhibition

or enhancement of chromatin binding and H2A-Ub activity. Further investigation into chromatin

complexes containing full-length BRCA1/BARD1 and the many protein assemblies and binding

partners that are required to execute its cellular functions is warranted.

<u>Acknowledgements</u>

We thank K. Luger (University of Colorado Boulder) for sharing plasmids for the 147-bp and

NLE-trimer '601' DNA, S. Tan (Penn State University) for sharing the plasmid for the 185-bp

'601' DNA, and G. Debelouchina (UC San Diego) for the MMTV DNA plasmid. We thank D.

Veelser for providing access to equipment, and M. Morgan for insights into dichloroacetone

crosslinking of H2A-Ub. S.R.W., L.M.T., K.EK., P.S.B., and R.E.K. were supported by the NIH

(R01CA260834). R.E.K. is the Edmond H. Fischer/Washington Research Foundation Endowed

Chair in Biochemistry. This work and A.Z. was supported in part by the University of

Washington's Proteomics Resource (UWPR95794). A.Z. and T.N.D. were also supported by the

NIH (P41GM103533). W.Z. was supported by a Young Investigator Award from Max and Minnie

Tomerlin Voelcker Fund, the Cancer Prevention and Research Institute of Texas (RP210102),

and the NIH (R01GM141091). D.B.W. was supported by the NIH (R00-HD090201).

<u>Author Contributions</u>

540   S.R.W., W.Z., P.S.B., and R.E.K. conceived of the study. S.R.W., L.M.T., and K.E.K. performed

541   experiments and analyzed data. D.B.W. analyzed data. A.Z. performed chemical crosslinking

542   and mass spectrometry and associated data processing under the supervision of T.N.D. S.R.W.

543   wrote the initial draft of the manuscript and generated all figures. The final manuscript was

544   written by S.R.W., P.S.B., and R.E.K., with input from all co-authors.

Competing Interests Statement

The authors of this article declare no competing interests.

References

Alenezi, W. M., Fierheller, C. T., Recio, N., & Tonin, P. N. (2020). Literature Review of BARD1 as a Cancer Predisposing Gene with a Focus on Breast and Ovarian Cancers. *Genes*, *11*(8). https://doi.org/10.3390/genes11080856

Becker, J. R., Clifford, G., Bonnet, C., Groth, A., Wilson, M. D., & Chapman, J. R. (2021). BARD1 reads H2A lysine 15 ubiquitination to direct homologous recombination. *Nature*, *596*(7872), 433–437. https://doi.org/10.1038/s41586-021-03776-w

Belotserkovskaya, R., Raga Gil, E., Lawrence, N., Butler, R., Clifford, G., Wilson, M. D., & Jackson, S. P. (2020). PALB2 chromatin recruitment restores homologous recombination in BRCA1-deficient cells depleted of 53BP1. *Nature Communications*, *11*(1), 819. https://doi.org/10.1038/s41467-020-14563-y

Bochar, D. A., Wang, L., Beniya, H., Kinev, A., Xue, Y., Lane, W. S., Wang, W., Kashanchi, F., & Shiekhattar, R. (2000). BRCA1 Is Associated with a Human SWI/SNF-Related Complex: Linking Chromatin Remodeling to Breast Cancer. *Cell*, *102*(2), 257–265. https://doi.org/10.1016/S0092-8674(00)00030-1

Brzovic, P. S., Keeffe, J. R., Nishikawa, H., Miyamoto, K., Fox, D., 3rd, Fukuda, M., Ohta, T., & Klevit, R. (2003). Binding and recognition in the assembly of an active BRCA1/BARD1 ubiquitin-ligase complex. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(10), 5646–5651. PubMed. https://doi.org/10.1073/pnas.0836054100

Brzovic, P. S., Rajagopal, P., Hoyt, D. W., King, M. C., & Klevit, R. E. (2001). Structure of a BRCA1-BARD1 heterodimeric RING-RING complex. *Nature Structural Biology*, *8*(10), 833–837. PubMed. https://doi.org/10.1038/nsb1001-833

Budziszewski, G. R., Zhao, Y., Spangler, C. J., Kedziora, K. M., Williams, M. R., Azzam, D. N., Skrajna, A., Koyama, Y., Cesmat, A. P., Simmons, H. C., Arteaga, E. C., Strauss, J. D., Kireev, D., & McGinty, R. K. (2022). Multivalent DNA and nucleosome acidic patch interactions specify VRK1 mitotic localization and activity. *Nucleic Acids Research*, *50*(8), 4355–4371. https://doi.org/10.1093/nar/gkac198

Chambers, M. C., Maclean, B., Burke, R., Amodei, D., Ruderman, D. L., Neumann, S., Gatto, L., Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen, B., Baker, T. A., Brusniak, M.-Y., Paulse, C., Creasy, D., … Mallick, P. (2012). A cross-platform toolkit for mass spectrometry and proteomics. *Nature Biotechnology*, *30*(10), 918–920. https://doi.org/10.1038/nbt.2377

Chin Jason W., Martin Andrew B., King David S., Wang Lei, & Schultz Peter G. (2002). Addition of a photocrosslinking amino acid to the genetic code of Escherichia coli. *Proceedings of the National Academy of Sciences*, *99*(17), 11020–11024. https://doi.org/10.1073/pnas.172226299

Christensen, D. E., Brzovic, P. S., & Klevit, R. E. (2007). E2-BRCA1 RING interactions dictate synthesis of mono- or specific polyubiquitin chain linkages. *Nature Structural & Molecular Biology*, *14*(10), 941–948. https://doi.org/10.1038/nsmb1295

Dai, L., Dai, Y., Han, J., Huang, Y., Wang, L., Huang, J., & Zhou, Z. (2021). Structural insight into BRCA1-BARD1 complex recruitment to damaged chromatin. *Molecular Cell*, *81*(13), 2765-2777.e6. https://doi.org/10.1016/j.molcel.2021.05.010

Dao, H. T., Dul, B. E., Dann, G. P., Liszczak, G. P., & Muir, T. W. (2020). A basic motif anchoring ISWI to nucleosome acidic patch regulates nucleosome spacing. *Nature Chemical Biology*, *16*(2), 134–142. https://doi.org/10.1038/s41589-019-0413-4

Densham, R. M., Garvin, A. J., Stone, H. R., Strachan, J., Baldock, R. A., Daza-Martin, M., Fletcher, A., Blair-Reid, S., Beesley, J., Johal, B., Pearl, L. H., Neely, R., Keep, N. H., Watts, F. Z., & Morris, J. R. (2016). Human BRCA1-BARD1 ubiquitin ligase activity counteracts chromatin barriers to DNA resection. *Nature Structural & Molecular Biology*, *23*(7), 647–655. PubMed. https://doi.org/10.1038/nsmb.3236

Densham, R. M., & Morris, J. R. (2019). Moving Mountains—The BRCA1 Promotion of DNA Resection. *Frontiers in Molecular Biosciences*, *6*, 79. https://doi.org/10.3389/fmolb.2019.00079

Dyer, P. N., Edayathumangalam, R. S., White, C. L., Bao, Y., Chakravarthy, S., Muthurajan, U. M., & Luger, K. (2003). Reconstitution of Nucleosome Core Particles from Recombinant Histones and DNA. In *Methods in Enzymology* (Vol. 375, pp. 23–44). Academic Press. https://doi.org/10.1016/S0076-6879(03)75002-2

Eng, J. K., Jahan, T. A., & Hoopmann, M. R. (2013). Comet: An open-source MS/MS sequence database search tool. *PROTEOMICS*, *13*(1), 22–24. https://doi.org/10.1002/pmic.201200439

Gudmundsdottir, K., & Ashworth, A. (2006). The roles of BRCA1 and BRCA2 and associated proteins in the maintenance of genomic stability. *Oncogene*, *25*(43), 5864–5874. https://doi.org/10.1038/sj.onc.1209874

Hatchi, E., Skourti-Stathaki, K., Ventz, S., Pinello, L., Yen, A., Kamieniarz-Gdula, K., Dimitrov, S., Pathania, S., McKinney, K. M., Eaton, M. L., Kellis, M., Hill, S. J., Parmigiani, G., Proudfoot, N. J., & Livingston, D. M. (2015). BRCA1 recruitment to transcriptional pause sites is required for R-loop-driven DNA damage repair. *Molecular Cell*, *57*(4), 636–647. PubMed. https://doi.org/10.1016/j.molcel.2015.01.011

Hoopmann, M. R., Zelter, A., Johnson, R. S., Riffle, M., MacCoss, M. J., Davis, T. N., & Moritz, R. L. (2015). Kojak: Efficient Analysis of Chemically Cross-Linked Protein Complexes. *Journal of Proteome Research*, *14*(5), 2190–2198. https://doi.org/10.1021/pr501321h

Hsu, P. L., Shi, H., Leonen, C., Kang, J., Chatterjee, C., & Zheng, N. (2019). Structural Basis of H2B Ubiquitination-Dependent H3K4 Methylation by COMPASS. *Molecular Cell*, *76*(5), 712-723.e4. https://doi.org/10.1016/j.molcel.2019.10.013

Hu, Q., Botuyan, M. V., Zhao, D., Cui, G., Mer, E., & Mer, G. (2021). Mechanisms of BRCA1– BARD1 nucleosome recognition and ubiquitylation. *Nature*, *596*(7872), 438–443. https://doi.org/10.1038/s41586-021-03716-8

Huen, M. S. Y., Sy, S. M. H., & Chen, J. (2010). BRCA1 and its toolbox for the maintenance of genome integrity. *Nature Reviews Molecular Cell Biology*, *11*(2), 138–148. https://doi.org/10.1038/nrm2831

Kalb, R., Mallery, D. L., Larkin, C., Huang, J. T. J., & Hiom, K. (2014). BRCA1 is a histone-H2A-specific ubiquitin ligase. *Cell Reports*, *8*(4), 999–1005. PubMed. https://doi.org/10.1016/j.celrep.2014.07.025

Käll, L., Canterbury, J. D., Weston, J., Noble, W. S., & MacCoss, M. J. (2007). Semi-supervised
learning for peptide identification from shotgun proteomics datasets. *Nature Methods*,
4(11), 923–925. https://doi.org/10.1038/nmeth1113

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Molecular Biology and Evolution*, *30*(4),
772–780. https://doi.org/10.1093/molbev/mst010

Kim, S.-A., Chatterjee, N., Jennings, M. J., Bartholomew, B., & Tan, S. (2015).
Extranucleosomal DNA enhances the activity of the LSD1/CoREST histone demethylase
complex. *Nucleic Acids Research*, *43*(10), 4868–4880.
https://doi.org/10.1093/nar/gkv388

Kjaergaard, M., & Poulsen, F. M. (2011). Sequence correction of random coil chemical shifts:
Correlation between neighbor correction factors and changes in the Ramachandran
distribution. *Journal of Biomolecular NMR*, *50*(2), 157–165.
https://doi.org/10.1007/s10858-011-9508-2

Kolas Nadine K., Chapman J. Ross, Nakada Shinichiro, Ylanko Jarkko, Chahwan Richard,
Sweeney Frédéric D., Panier Stephanie, Mendez Megan, Wildenhain Jan, Thomson
Timothy M., Pelletier Laurence, Jackson Stephen P., & Durocher Daniel. (2007).
Orchestration of the DNA-Damage Response by the RNF8 Ubiquitin Ligase. *Science*,
318(5856), 1637–1640. https://doi.org/10.1126/science.1150034

Krais, J. J., & Johnson, N. (2020). BRCA1 Mutations in Cancer: Coordinating Deficiencies in
Homologous Recombination with Tumorigenesis. *Cancer Research*, *80*(21), 4601–4609.
https://doi.org/10.1158/0008-5472.CAN-20-1830

Krais, J. J., Wang, Y., Patel, P., Basu, J., Bernhardy, A. J., & Johnson, N. (2021). RNF168-
mediated localization of BARD1 recruits the BRCA1-PALB2 complex to DNA damage.
Nature Communications, *12*(1), 5016. https://doi.org/10.1038/s41467-021-25346-4

Lazar, G. A., Desjarlais, J. R., & Handel, T. M. (1997). De novo design of the hydrophobic core
of ubiquitin. *Protein Science : A Publication of the Protein Society*, *6*(6), 1167–1178.
PubMed. https://doi.org/10.1002/pro.5560060605

Lee, Y.-T., Gibbons, G., Lee, S. Y., Nikolovska-Coleska, Z., & Dou, Y. (2015). One-pot refolding
of core histones from bacterial inclusion bodies allows rapid reconstitution of histone
octamer. *Protein Expression and Purification*, *110*, 89–94. PubMed.
https://doi.org/10.1016/j.pep.2015.02.007

Lorick, K. L., Jensen, J. P., Fang, S., Ong, A. M., Hatakeyama, S., & Weissman, A. M. (1999).
RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination.
Proceedings of the National Academy of Sciences, *96*(20), 11364.
https://doi.org/10.1073/pnas.96.20.11364

Lowary, P. T., & Widom, J. (1998). New DNA sequence rules for high affinity binding to histone
octamer and sequence-directed nucleosome positioning11Edited by T. Richmond.
Journal of Molecular Biology, *276*(1), 19–42. https://doi.org/10.1006/jmbi.1997.1494

Luger, K., Rechsteiner, T. J., & Richmond, T. J. (1999). Preparation of nucleosome core particle
from recombinant histones. *Methods in Enzymology*, *304*, 3–19.
https://doi.org/10.1016/s0076-6879(99)04003-3

Mark, W.-Y., Liao, J. C. C., Lu, Y., Ayed, A., Laister, R., Szymczyna, B., Chakrabartty, A., &
Arrowsmith, C. H. (2005). Characterization of segments from the central region of
BRCA1: An intrinsically disordered scaffold for multiple protein-protein and protein-DNA

interactions? *Journal of Molecular Biology*, *345*(2), 275–287. PubMed.
https://doi.org/10.1016/j.jmb.2004.10.045

Masuda, T., Xu, X., Dimitriadis, E. K., Lahusen, T., & Deng, C.-X. (2016). "DNA Binding Region"
of BRCA1 Affects Genetic Stability through modulating the Intra-S-Phase Checkpoint.
International Journal of Biological Sciences, *12*(2), 133–143. PubMed.
https://doi.org/10.7150/ijbs.14242

Mattiroli, F., Vissers, J. H. A., van Dijk, W. J., Ikpa, P., Citterio, E., Vermeulen, W., Marteijn, J.
A., & Sixma, T. K. (2012). RNF168 ubiquitinates K13-15 on H2A/H2AX to drive DNA
damage signaling. *Cell*, *150*(6), 1182–1195. https://doi.org/10.1016/j.cell.2012.08.005

McGinty, R. K., Henrici, R. C., & Tan, S. (2014). Crystal structure of the PRC1 ubiquitylation
module bound to the nucleosome. *Nature*, *514*(7524), 591–596. PubMed.
https://doi.org/10.1038/nature13890

McGinty, R. K., & Tan, S. (2016). Recognition of the nucleosome by chromatin factors and
enzymes. *Current Opinion in Structural Biology*, *37*, 54–61.
https://doi.org/10.1016/j.sbi.2015.11.014

McGinty, R. K., & Tan, S. (2021). Principles of nucleosome recognition by chromatin factors and
enzymes. *Catalysis and Regulation ● Protein Nucleic Acid Interactions*, *71*, 16–26.
https://doi.org/10.1016/j.sbi.2021.05.006

McGuffin, L. J., Bryson, K., & Jones, D. T. (2000). The PSIPRED protein structure prediction
server. *Bioinformatics*, *16*(4), 404–405. https://doi.org/10.1093/bioinformatics/16.4.404

Morgan, M., Jbara, M., Brik, A., & Wolberger, C. (2019). Chapter One—Semisynthesis of
ubiquitinated histone H2B with a native or nonhydrolyzable linkage. In M. Hochstrasser
(Ed.), *Methods in Enzymology* (Vol. 618, pp. 1–27). Academic Press.
https://doi.org/10.1016/bs.mie.2019.01.003

Morgan Michael T., Haj-Yahya Mahmood, Ringel Alison E., Bandi Prasanthi, Brik Ashraf, &
Wolberger Cynthia. (2016). Structural basis for histone H2B deubiquitination by the
SAGA DUB module. *Science*, *351*(6274), 725–728.
https://doi.org/10.1126/science.aac5681

Mullan, P. B., Quinn, J. E., & Harkin, D. P. (2006). The role of BRCA1 in transcriptional
regulation and cell cycle control. *Oncogene*, *25*(43), 5854–5863.
https://doi.org/10.1038/sj.onc.1209872

Nakamura, K., Saredi, G., Becker, J. R., Foster, B. M., Nguyen, N. V., Beyer, T. E., Cesa, L. C.,
Faull, P. A., Lukauskas, S., Frimurer, T., Chapman, J. R., Bartke, T., & Groth, A. (2019).
H4K20me0 recognition by BRCA1–BARD1 directs homologous recombination to sister
chromatids. *Nature Cell Biology*, *21*(3), 311–318. https://doi.org/10.1038/s41556-019-
0282-9

Poepsel, S., Kasinath, V., & Nogales, E. (2018). Cryo-EM structures of PRC2 simultaneously
engaged with two functionally distinct nucleosomes. *Nature Structural & Molecular
Biology*, *25*(2), 154–162. https://doi.org/10.1038/s41594-018-0023-y

Riffle, M., Jaschob, D., Zelter, A., & Davis, T. N. (2016). ProXL (Protein Cross-Linking
Database): A Platform for Analysis, Visualization, and Sharing of Protein Cross-Linking
Mass Spectrometry Data. *Journal of Proteome Research*, *15*(8), 2863–2870.
https://doi.org/10.1021/acs.jproteome.6b00274

Riffle, M., Jaschob, D., Zelter, A., & Davis, T. N. (2019). Proxl (Protein Cross-Linking Database): A Public Server, QC Tools, and Other Major Updates. *Journal of Proteome Research*, *18*(2), 759–764. https://doi.org/10.1021/acs.jproteome.8b00726

Santos-Pereira, J. M., & Aguilera, A. (2015). R loops: New modulators of genome dynamics and function. *Nature Reviews Genetics*, *16*(10), 583–597. https://doi.org/10.1038/nrg3961

Savage, K. I., & Harkin, D. P. (2015). BRCA1, a "complex" protein involved in the maintenance of genomic stability. *The FEBS Journal*, *282*(4), 630–646. https://doi.org/10.1111/febs.13150

Schlacher, K., Wu, H., & Jasin, M. (2012). A Distinct Replication Fork Protection Pathway Connects Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. *Cancer Cell*, *22*(1), 106–116. https://doi.org/10.1016/j.ccr.2012.05.015

Scully, R., Chen, J., Ochs, R. L., Keegan, K., Hoekstra, M., Feunteun, J., & Livingston, D. M. (1997). Dynamic Changes of BRCA1 Subnuclear Location and Phosphorylation State Are Initiated by DNA Damage. *Cell*, *90*(3), 425–435. https://doi.org/10.1016/S0092-8674(00)80503-6

Simons, A. M., Horwitz, A. A., Starita, L. M., Griffin, K., Williams, R. S., Glover, J. N. M., & Parvin, J. D. (2006). BRCA1 DNA-Binding Activity Is Stimulated by BARD1. *Cancer Research*, *66*(4), 2012–2018. https://doi.org/10.1158/0008-5472.CAN-05-3296

Sobhian Bijan, Shao Genze, Lilli Dana R., Culhane Aedín C., Moreau Lisa A., Xia Bing, Livingston David M., & Greenberg Roger A. (2007). RAP80 Targets BRCA1 to Specific Ubiquitin Structures at DNA Damage Sites. *Science*, *316*(5828), 1198–1202. https://doi.org/10.1126/science.1139516

Stewart, M. D., Zelin, E., Dhall, A., Walsh, T., Upadhyay, E., Corn, J. E., Chatterjee, C., King, M.-C., & Klevit, R. E. (2018). BARD1 is necessary for ubiquitylation of nucleosomal histone H2A and for transcriptional regulation of estrogen metabolism genes. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(6), 1316–1321. PubMed. https://doi.org/10.1073/pnas.1715467115

Tan, W., Murphy, V. J., Charron, A., van Twest, S., Sharp, M., Constantinou, A., Parker, M. W., Crismani, W., Bythell-Douglas, R., & Deans, A. J. (2020). Preparation and purification of mono-ubiquitinated proteins using Avi-tagged ubiquitin. *PLOS ONE*, *15*(2), e0229000. https://doi.org/10.1371/journal.pone.0229000

Tarsounas, M., & Sung, P. (2020). The antitumorigenic roles of BRCA1–BARD1 in DNA repair and replication. *Nature Reviews Molecular Cell Biology*, *21*(5), 284–299. https://doi.org/10.1038/s41580-020-0218-z

Uckelmann, M., Densham, R. M., Baas, R., Winterwerp, H. H. K., Fish, A., Sixma, T. K., & Morris, J. R. (2018). USP48 restrains resection by site-specific cleavage of the BRCA1 ubiquitin mark from H2A. *Nature Communications*, *9*(1), 229. https://doi.org/10.1038/s41467-017-02653-3

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, *17*(3), 261–272. https://doi.org/10.1038/s41592-019-0686-2

Vizcaíno, J. A., Côté, R., Reisinger, F., M. Foster, J., Mueller, M., Rameseder, J., Hermjakob, H., & Martens, L. (2009). A guide to the Proteomics Identifications Database proteomics data repository. *PROTEOMICS*, *9*(18), 4276–4283. https://doi.org/10.1002/pmic.200900402

Wang, X., Paucek, R. D., Gooding, A. R., Brown, Z. Z., Ge, E. J., Muir, T. W., & Cech, T. R. (2017). Molecular analysis of PRC2 recruitment to DNA in chromatin and its inhibition by RNA. *Nature Structural & Molecular Biology*, *24*(12), 1028–1038. https://doi.org/10.1038/nsmb.3487

Witus, S. R., Burrell, A. L., Farrell, D. P., Kang, J., Wang, M., Hansen, J. M., Pravat, A., Tuttle, L. M., Stewart, M. D., Brzovic, P. S., Chatterjee, C., Zhao, W., DiMaio, F., Kollman, J. M., & Klevit, R. E. (2021). BRCA1/BARD1 site-specific ubiquitylation of nucleosomal H2A is directed by BARD1. *Nature Structural & Molecular Biology*, *28*(3), 268–277. https://doi.org/10.1038/s41594-020-00556-4

Witus, S. R., Stewart, M. D., & Klevit, R. E. (2021). The BRCA1/BARD1 ubiquitin ligase and its substrates. *Biochemical Journal*, 3467–3483. https://doi.org/10.1042/BCJ20200864

Witus, S. R., Zhao, W., Brzovic, P. S., & Klevit, R. E. (2022). BRCA1/BARD1 is a nucleosome reader and writer. *Trends in Biochemical Sciences*. https://doi.org/10.1016/j.tibs.2022.03.001

Worden, E. J., Hoffmann, N. A., Hicks, C. W., & Wolberger, C. (2019). Mechanism of Cross-talk between H2B Ubiquitination and H3 Methylation by Dot1L. *Cell*, *176*(6), 1490-1501.e12. PubMed. https://doi.org/10.1016/j.cell.2019.02.002

Worden, E. J., Zhang, X., & Wolberger, C. (2020). Structural basis for COMPASS recognition of an H2B-ubiquitinated nucleosome. *ELife*, *9*, e53199. https://doi.org/10.7554/eLife.53199

Zelter, A., Bonomi, M., Kim, J. ook, Umbreit, N. T., Hoopmann, M. R., Johnson, R., Riffle, M., Jaschob, D., MacCoss, M. J., Moritz, R. L., & Davis, T. N. (2015). The molecular architecture of the Dam1 kinetochore complex is defined by cross-linking based structural modelling. *Nature Communications*, *6*(1), 8673. https://doi.org/10.1038/ncomms9673

Zhao, W., Steinfeld, J. B., Liang, F., Chen, X., Maranon, D. G., Jian Ma, C., Kwon, Y., Rao, T., Wang, W., Sheng, C., Song, X., Deng, Y., Jimenez-Sainz, J., Lu, L., Jensen, R. B., Xiong, Y., Kupfer, G. M., Wiese, C., Greene, E. C., & Sung, P. (2017). BRCA1-BARD1 promotes RAD51-mediated homologous DNA pairing. *Nature*, *550*(7676), 360–365. PubMed. https://doi.org/10.1038/nature24060

Zhu, G., Xia, Y., Nicholson, L. K., & Sze, K. H. (2000). Protein Dynamics Measurements by TROSY-Based NMR Experiments. *Journal of Magnetic Resonance*, *143*(2), 423–426. https://doi.org/10.1006/jmre.2000.2022

Zhu, Q., Hoong, N., Aslanian, A., Hara, T., Benner, C., Heinz, S., Miga, K. H., Ke, E., Verma, S., Soroczynski, J., Yates, J. R., Hunter, T., & Verma, I. M. (2018). Heterochromatin-Encoded Satellite RNAs Induce Breast Cancer. *Molecular Cell*, *70*(5), 842-853.e7. https://doi.org/10.1016/j.molcel.2018.04.023

Zhu, Q., Pao, G. M., Huynh, A. M., Suh, H., Tonnu, N., Nederlof, P. M., Gage, F. H., & Verma, I. M. (2011). BRCA1 tumour suppression occurs via heterochromatin-mediated silencing. *Nature*, *477*(7363), 179–184. https://doi.org/10.1038/nature10371

 Figures

811
Figure 1. Contributions of BARD1 regions to nucleosome binding and H2A-Ub activity.
(A) Domain organization of BRCA1 and BARD1. The domain names are indicated above the
cartoons (RING, really interesting new gene; IDR, intrinsically disordered region; CC, coiled-coil;
BRCT, BRCA1 C-terminal; Ank, Ankyrin repeat domain; CTD, C-terminal domain). Binding sites
of interacting partners previously described and relevant to this work are indicated below the
domain illustrations. The heterodimeric interface is denoted by an asterisk. **(B)** Quantification of
time-course nucleosome H2A-Ub assays using the indicated BRCA1/BARD1 truncations. Data
show the mean; error bars are ± 1-s.d. of n=3 independent experiments. **(C)** Comparison of
nucleosome binding affinity from fluorescence-quenching based measurements using the
indicated BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of n=4
independent experiments. The reported difference in affinities to BRCA1^{RING}/BARD1^{RING} is likely
underestimated, as the minimal RING/RING heterodimer binding data was collected at a lower
ionic strength than the rest of the constructs to obtain well-behaved binding data (50 mM vs.
100 mM NaCl).

826
Figure 1—figure supplement 1. Characterization of BRCA1/BARD1 constructs. (A)
Representative time-course nucleosome ubiquitylation assays monitoring H2A-Ub efficiency of
different BRCA1/BARD1 constructs quantified in Panel E and Figure 1B. These data are
representative of the basic assay set-up for other time-course H2A-Ub assays in this study. **(B)**
Electrophoretic mobility shift assay (EMSA) comparing the nucleosome binding of full-length
BRCA1/BARD1 (top) to the minimal RING heterodimer (bottom). Data shown are representative
of n=2 independent experiments. **(C-E)** Quantification of H2A-Ub assays using the indicated
BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of n=3 independent
experiments. **(F)** Comparison of nucleosome binding affinity from fluorescence-quenching
based measurements using the indicated BRCA1/BARD1 constructs. Data show the mean;
error bars are ± 1-s.d. of n=4 independent experiments. **(G)** EMSA monitoring nucleosome
binding by the indicated BARD1 IDR fragments. Data shown are representative of n=2
independent experiments.
840
841
842

1. BRCA1^{FL}/BARD1^{FL}
2. BRCA1$^{\Delta 11q}$/BARD1^{FL}
3. BRCA1^{RING}/BARD1^{FL}
4. BRCA1^{RING}/BARD1$^{\Delta CTDs}$
5. BRCA1^{FL}/BARD1^{RING}
6. BRCA1$^{\Delta 11q}$/BARD1^{RING}
7. BRCA1^{RING}/BARD1$^{FL\text{-}P89A/W91A}$
8. BRCA1^{RING}/BARD1$^{\Delta 194\text{-}216}$
9. BRCA1^{RING}/BARD1$^{\Delta 150\text{-}155/\Delta 194\text{-}216}$
10. BRCA1^{RING}/BARD1$^{\Delta 140\text{-}270}$
11. BRCA1^{RING}/BARD1^{216}
12. BRCA1^{RING}/BARD1^{193}
13. BRCA1^{RING}/BARD1^{RING}

843
844 **Figure 1—figure supplement 2. Coomassie-stained SDS-PAGE gels of BRCA1/BARD1**
845 **constructs used in this study.**

Figure 2. The BARD1 IDR binds to nucleosomes via DNA. (A) $^1H^{15}N$-HSQC NMR titration spectra of ^{15}N-BARD1 124-270 with a 36-bp dsDNA fragment. Darker shades of blue correspond to increasing amounts of DNA added. A subset of highly affected signals is labelled with their residue identities, with arrows showing their trajectories over the course of the titration. **(B)** $^1H^{15}N$-HSQC NMR amide chemical shift perturbations (CSPs) observed to ^{15}N-BARD1 141-216 signals when bound to a 36-bp dsDNA fragment (1:1 molar equivalent complex). **(C)** Representative SDS-PAGE gel monitoring in-gel fluorescence (labelled DNA) of UV-induced Bpa crosslinking of BRCA1-f-BARD1[221] to nucleosomes (left). Quantification of Bpa crosslinking experiments with nucleosomes (right). The intensity of each crosslinked band was normalized to the intensity of the L120Bpa crosslinked band for each replicate experiment. Graph bars show the mean; error bars are ± 1-s.d. and the open circles are the values of individual replicates for of n=3 independent experiments.

892
893
Figure 2—figure supplement 1. Characterization of the BARD1 intrinsically disordered
region. (A) Coomassie-stained SDS-PAGE gel of BARD1 fragments used in NMR experiments.
(B) $^1H^{15}N$-HSQC NMR spectra of the indicated ^{15}N-labelled BARD1 fragments. **(C)** Multiple-
sequence alignment of BARD1 124-270 with a representative subset of homologs. Residue
identities that are conserved to the human sequence are indicated by the presence of a period
in place of a residue letter. Basic residues (Lys, Arg, His) are color coded in blue, and basic
positions that are fully conserved across all species are highlighted with a gray background.

909
910
Figure 2—figure supplement 2. NMR analysis of BARD1 IDR-DNA and IDR-nucleosome
complexes. (A) EMSA comparing DNA and nucleosome binding by BARD1 124-270. Data
shown are representative of n=2 independent experiments. **(B)** ^{1}H^{15}N-HSQC NMR spectra of
^{15}N-BARD1 124-270, titrating a 36-bp dsDNA fragment (left), and a single titration point with
nucleosomes (right). The red arrows point to signals that are similarly affected between the
dsDNA- and nucleosome-bound samples. **(C)** ^{1}H^{15}N-HSQC NMR spectra of ^{15}N-BARD1 141-
216 with a 36-mer dsDNA fragment. Data from panel C are also shown in Figure 4—figure
supplement 3A.

928
Figure 2—figure supplement 3. Changes in dynamics and secondary structure of the
BARD1 IDR bound to DNA. (A) Changes in secondary structural propensity (SSP) of [15]N-
BARD1 141-216 when bound to a 36-bp dsDNA fragment. The black bars for apo BARD1 are
behind the gray bars where DNA was added. **(B)** T1/T2 dynamics measurements for amide
signals of [15]N-BARD1 141-216 in apo form and bound to dsDNA (1:1 molar equivalent complex).
Data from panel B are also shown in Figure 4—figure supplement 3C.

954
955
Figure 2—figure supplement 4. UV-induced Bpa crosslinking of BRCA1/BARD1 to
nucleosomal DNA. (A) Coomassie-stained SDS-PAGE gel of unfused BRCA1[RING]/BARD1[216]
heterodimers and BRCA1-f-BARD1[221] constructs with and without Bpa incorporation. **(B)** H2A-
Ub activity of unfused heterodimers compared to wild-type and Bpa-incorporated BRCA1-f-
BARD1[221]. Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments. **(C)**
Native gel of fluorescently labelled DNA and nucleosomes used in Bpa crosslinking
experiments. **(D)** Representative data of in-gel fluorescence signal from of a UV-induced
crosslinking reaction to free 185-bp '601' DNA (DNA[185], top) and quantification of crosslinking
experiments (bottom). The intensity of each crosslinked band was normalized to the L120Bpa
crosslinked band intensity for each replicate experiment. Data bars show the mean; error bars
are $\pm$ 1-s.d. and the open circles are the values of individual replicates for of n=3 independent
experiments.

975
976
Figure 3. Contribution of BARD1 DNA-binding and RING-histone binding to H2A-Ub
activity. (A) The BARD1 RING-histone interface showing the locations of BARD1 P89 and W91
relative to the histone surface (PDB: 7JZV). **(B, C)** Quantification of time-course H2A-Ub assays
using the indicated BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of
n=3 independent experiments. Two single time-points were determined to be outliers in panel C
and excluded from analysis. Values for these excluded points are included as source data.

1014
1015
Figure 3—figure supplement 1. Nucleosome binding and Ub ligase activity of BARD1
RING mutants and DNA-binding chimeric constructs. (A) Comparison of nucleosome
binding affinity using the indicated BRCA1/BARD1 constructs. Data show the mean; error bars
are ± 1-s.d. of n=3 independent experiments. **(B)** Auto-ubiquitylation assay using
BRCA1^{RING}/BARD1^{FL} (wild-type or P89A/W91A). The assay monitors depletion of Ub and
appearance of high molecular weight auto-Ub products in the presence of BRCA1/BARD1. Data
shown is representative of n=2 independent experiments. **(C)** Nucleosome ubiquitylation assay
using the indicated BRCA1/BARD1 constructs and blotting for VSVG-tagged H2A (top) and H3
(middle and bottom). The asterisk denotes a proteolysis product of H3 that appears over the
time-course. **(D)** Comparison of nucleosome binding affinity using the indicated BRCA1/BARD1
constructs. Data show the mean; error bars are ± 1-s.d. of n=2 independent experiments.

1042
1043
Figure 4. Inhibition of nucleosome binding and H2A-Ub by specialized DNA structures.
(A) Design of 36-mer DNA competitor fragments used in H2A-Ub activity, Bpa crosslinking, and
NMR assays. Red indicates non-base-paired regions. The number of unpaired bases or base-
pairs is: ss – 36; ds – 0; bubble – 8; frayed – 8; ssOverhang – 24. **(B)** Single time-point H2A-Ub
inhibition assays using BRCA1[FL]/BARD1[FL] (100 nM) and the indicated competitor DNA (2.5 µM).
Data show the mean; error bars are $\pm$ 1-s.d.; and open circles are individual data points of n=4
independent experiments. One outlier point for "frayed-DNA" was excluded from the analysis.
(C) Inhibition of UV-induced Bpa crosslinking between the indicated BRCA1-f-BARD1[221] Bpa-
incorporated constructs and NCP[185] substrates in the presence of increasing amounts of dsDNA
or bubble-DNA competitor. Data show the mean; error bars are $\pm$ 1-s.d. of n=3 (W146Bpa) or
n=4 (A195Bpa) independent experiments. **(D)** [1]H[15]N-HSQC NMR CSPs observed to [15]N-BARD1
141-216 signals when bound to a 36-mer dsDNA (blue, front) or bubble-DNA (gray, back)
fragment (1:1 molar equivalent complex). Signals for residues 143-157 are broadened beyond
detection in the bubble-DNA-bound spectrum, and gray bars are equal to the CSP value for the
W146Nε CSP that was observed (see Extended Data Figure 3c, d for spectra). **(E)** Selected
region of [13]C-HSQC spectra of [13]C-BARD1 141-216 in 1:1 molar equivalent complex with

dsDNA fragment or bubble-DNA fragment. Signal trajectories in bound spectra are indicated by
arrows. **(F)** Single time-point H2A-Ub inhibition assays using heterodimers containing
BRCA1^{RING}/BARD1^{FL} (left, 12 min endpoint) or the indicated BARD1 internal deletion mutant
(middle and right, 20 min endpoint) and increasing amounts of dsDNA or bubble-DNA
competitor. The same amount of E3 (50 nM) was used for each BRCA1/BARD1 truncation.
Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments. One time-point
from Panels B and C each were determined to be outliers and excluded from analysis. Values
for these points are included as source data.

1081
1082
Figure 4—figure supplement 1. Assembly of competitor DNA fragments and inhibition of
H2A-Ub activity. (A) DNA-stained native-gel of annealed DNA competitor fragments. Amounts
loaded were normalized to show similar band-intensity for each fragment. **(B)** Representative
gel data of H2A-Ub inhibition assay using full-length BRCA1/BARD1 (100 nM) and the indicated
DNA competitor fragment (2.5 mM) **(C)** Representative gel data of single time-point (20 min)
H2A-Ub inhibition assays using BRCA1^{RING}/BARD1^{216} (top, 50 nM E3) and
BRCA1^{RING}/BARD1^{140} (bottom, 750 nM E3) and indicated amounts of dsDNA and bubble-DNA
competitors. **(D)** Quantification of H2A-Ub DNA-competitor inhibition assays from panel b using
the indicated BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of n=3
independent experiments. **(E)** Representative gel data of single time-point H2A-Ub inhibition
assays using BRCA1^{RING}/BARD1^{FL} (top, 50 nM E3, 12 min endpoint), BRCA1^{RING}/BARD1$^{\Delta194-216}$
(middle, 50 nM E3, 20 min endpoint), and BRCA1^{RING}/BARD1$^{\Delta150-155/\Delta194-216}$ (bottom, 50 nM E3,
20 min endpoint) and indicated amounts of dsDNA and bubble-DNA competitors.

1096
1097
1098

1099
1100
1101 **Figure 4—figure supplement 2. Inhibition of UV-induced Bpa crosslinking of**
1102 **BRCA1/BARD1 to nucleosomal DNA.** Representative data of in-gel fluorescence signal from a
1103 UV-induced Bpa crosslinking reaction between W146Bpa (top) and A195Bpa (bottom) BRCA1-
1104 f-BARD1[221] to 185-bp '601' nucleosomes (NCP[185]) in the presence of dsDNA and bubble-DNA
1105 competitors. Quantified data from n=3 replicate experiments are shown in Figure 4C.

1127

Figure 4—figure supplement 3. Comparison of dsDNA and bubble-DNA binding to the BARD1 IDR by NMR. (A) $^1H^{15}N$-HSQC NMR spectra of ^{15}N-BARD1 141-216 with a 36-mer dsDNA fragment or **(B)** a 36-mer bubble-DNA fragment with 8-bp of mismatch. Both complexes were assembled at 1:1 molar equivalent. The W146Nε signals are inlaid, and the red arrows point to a signal that is present exclusively in the DNA-bound spectra. **(C)** T1/T2 dynamics measurements for amide signals of ^{15}N-BARD1 141-216 in apo form and bound to dsDNA or bubble-DNA (1:1 molar equivalent complex). **(D)** $^1H^{13}C$-HSQC NMR spectra of ^{13}C-BARD1 141-216 with a 36-mer dsDNA fragment (blue), or a 36-mer bubble-DNA fragment with 8-bp of mismatch (gray). Both complexes were assembled at 1:1 molar equivalent. Data from panel A are also shown in Figure 2—figure supplement 2C. A subset of data from panel C (apo and +dsDNA) are also shown in Figure 2—figure supplement 3B.

1179
1180
Figure 5. An unmodified mono-nucleosome with linker DNA is an optimal substrate for
BRCA1/BARD1. (A) Schematic of chromatin substrates used for H2A-Ub activity and
nucleosome binding assays in panels b and c. The length of linker-DNA is indicated next to the
NCP[185] and tri-NCP. **(B, C)** Quantification of time-course H2A-Ub assays using the indicated
BRCA1/BARD1 constructs and chromatin substrates. Data show the mean; error bars are $\pm$ 1-
s.d. of n=3 independent experiments.

1187
1188
1189

1190
Figure 5—figure supplement 1. Analysis of chromatin substrates (A) DNA-stained (left)
and fluorescence (right) native gel of chromatin substrates used for H2A-Ub activity assays in
Figure 5b and c. EcoRI digestion of tri-NCPs is used to confirm octamer binding saturation. **(B)**
Comparison of nucleosome binding affinity between BRCA1[RING]/BARD1[216] and NCP[147] and
NCP[185] substrates at 100 mM (top) and 150 mM (bottom) buffer-salt concentration (NaCl). The
solid vertical lines denote the position of K_d values for the indicated buffer-salt concentration,
and the dotted vertical lines denote the position of K_d values from the other buffer-salt
concentration. Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments
using both NCP[147] and NCP[185], with an additional replicate (total of n=4) for NCP[147].

1206
1207
Figure 6. Enhancement of H2A-Ub activity using chromatin substrates with pre-installed
H2A K15-Ub. (A) Quantification of time-course H2A-Ub assays using homogenously modified
H2A K15-Ub mono-nucleosome substrates and indicated BRCA1/BARD1 constructs. **(B)**
Intermolecular crosslinks observed by chemical crosslinking and MS analysis between BARD1
and histones using wild-type (left) and H2A K15-Ub (right) 147-bp '601' nucleosomes and
BRCA1[RING]/BARD1[FL] heterodimers. Crosslinks to histones emanating from the CTDs of BARD1
are shown in blue, and the RING-IDR region in gray. A lysine-depleted region of the BARD1
IDR is labelled and indicated by a black bar. **(C)** Design of asymmetric di-NCP substrates. **(D,**
E) Quantification of time-course H2A-Ub assays using the indicated combinations of di-NCP
substrate and BRCA1/BARD1 heterodimer. For panels A, D, and E, data show the mean; error
bars are ± 1-s.d. of n=3 independent experiments. One time-point from Panel A was determined
to be an outlier and excluded from analysis. The excluded value is reported as source data.

1220
1221
Figure 6—figure supplement 1. Design, assembly, and quality control of H2A K15-Ub
chromatin substrates. (A) Coomassie-stained SDS-PAGE gel of wild-type histones and
dichloroacetone linked H2A-Ub/H2B dimers. **(B)** DNA-stained native gel of mono-nucleosomes
with and without H2A-Ub incorporation at the indicated positions. **(C)** Example data of H2A-Ub
activity assay using full-length BRCA1/BARD1 and preinstalled H2A-Ub NCP substrates.
Western blots were probed for H2A. **(D)** Quantification of time-course H2A-Ub assays using
unmodified or pre-installed H2A-Ub chromatin substrates and BRCA1^{FL}/BARD1^{FL}. **(E)**
Schematic of asymmetric di-NCP assembly strategy. **(F)** DNA-stained native gel of di-NCP
ligation reaction. **(G)** DNA-stained (left) and fluorescence (right) native gels of di-NCP
substrates after purification by MgCl$_2$ precipitation. **(H)** Example gel data of H2A-Ub activity
assay using di-NCP substrates and indicated BRCA1/BARD1 constructs.

1233
1234

1235
1236
Figure 6—figure supplement 2. Analysis of BRCA1/BARD1 chromatin complexes by
chemical crosslinking and mass-spectrometry. (A) Coomassie-stained SDS-PAGE gel of
chemical crosslinking reactions used in mass spectrometry analysis **(B)** Crosslinks observed
from the RING domain of BRCA1 to ordered regions histones mapped on a high-resolution
structure of the complex (PDB: 7JZV). Cα-Cα distances are indicated in the figure. **(C)**
Quantitative comparison of crosslinking from reactions with unmodified and H2A K15-Ub NCPs.
Different subsets of crosslinks were quantified for the number of unique observed crosslinks and
total peptide spectral mapping counts for all crosslinks within a given subset. **(D)** Crosslinks
observed from the CTDs of BARD1 to ordered regions histones in the H2A K15-Ub NCP sample
mapped on a high-resolution structure (PDB: 7E8I). Cα-Cα distances are indicated. The P117
Cα serves as a proxy for the true crosslinking sites of K118 and K119 that are not modelled,
denoted by an asterisk. **(E)** Intermolecular crosslinks observed by chemical crosslinking and MS

1249   analysis between BARD1 and histones using wild-type (left) and H2A K15-Ub (right, separated
1250   by dashed line) 147-bp '601' nucleosomes and BRCA1^{RING}/BARD1^{FL} heterodimers. Crosslinks
1251   to histones emanating from the BARD1 RING-IDR region are highlighted in blue. A lysine-
1252   depleted region of the BARD1 IDR is labelled and indicated by a black bar.

Figure 7. Models of BRCA1/BARD1/chromatin complexes supported by this study. (A, B)
Model of the BRCA1/BARD1 minimal RING heterodimer binding to an unmodified mono-
nucleosome or higher-order chromatin substrate. **(C, D)** Model of BRCA1/BARD1 (BARD1 full-
length or RING-IDR) binding with an unmodified mono-nucleosome or higher-order chromatin
substrate. **(E, F)** Model of BRCA1/BARD1[FL] binding to ubiquitylated chromatin substrates in a
'wrapped complex (panel E) or in a complex that spans nucleosome units (panel F). The relative
H2A-Ub activity and affinity of the complexes is indicated below each set of panels.

1300 **Tables**

1301

1302 **Extended Data Table I. Details about BRCA1/BARD1 constructs used in this study.**

1303

Construct	BRCA1 bounds*	BRCA1 tags	BARD1 bounds*	BARD1 tags	Expression system
BRCA1^{FL}/ BARD1^{FL}	1-1863	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1$^{\Delta 11q}$/ BARD1^{FL}	1-264, 1366-1863	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1^{FL}	1-112	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta CTDs}$	1-112	N-FLAG	1-424	N-TwinStrepII	*Sf9*
BRCA1^{FL}/ BARD1^{RING}	1-1863	N-FLAG	1-140	N-TwinStrepII	*Sf9*
BRCA1$^{\Delta 11q}$/ BARD1^{RING}	1-264, 1366-1863	N-FLAG	1-140	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 194-216}$	1-112	N-FLAG	1-193, 217-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 150-155/\Delta 194-216}$	1-112	N-FLAG	1-149, 156-193, 217-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 140-270}$	1-112	N-FLAG	1-139, 271-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1^{216}	1-110	N-6xHis	26-216	none	*E. coli*
BRCA1^{RING}/ BARD1^{193}	1-110	N-6xHis	26-193	none	*E. coli*
BRCA1^{RING}/ BARD1^{RING}	1-110	N-6xHis	26-140	none	*E. coli*

1304
1305 * BRCA1 UniProt sequence ID: P38398-1; BARD1 UniProt sequence ID: Q99728-1

<u>Methods</u>

Protein, nucleosome, and experimental reagent production

Ubiquitylation machinery: Human E1 (UBA1), E2 (UBE2D3), and ubiquitin (wild-type and Q2C

mutant) were expressed and purified as previously described(Brzovic et al., 2003; Christensen

et al., 2007; Lazar et al., 1997). For a full list of BRCA1/BARD1 truncation constructs, their

residue bounds, purification/epitope tags, and expression system, see Extended Data Table I.

1320

BRCA1/BARD1 from E. coli: For purification of truncated BRCA1/BARD1 heterodimers from *E.*

coli expression (BRCA1^{RING}/BARD1^{RING}, BRCA1^{RING}/BARD1^{193}, and BRCA1^{RING}/BARD1^{216}),

plasmids containing 6xHis-tagged BRCA1 (pCOT7n) and untagged BARD1 (pET28n) were co-

transformed into *E. coli* BL21 (DE3) cells, grown in LB media at 37 °C to OD_{600nm} of 0.6-0.8,

supplemented with 100 μM $ZnCl_2$ and induced with 0.2 mM IPTG for ~16 hours at 16 °C. The

cell pellets were resuspended in 30 mL of Ni^{2+} start buffer (25 mM Tris-HCl pH 7.5, 500 mM

NaCl, 10 mM imidazole) supplemented with DNase, RNase, EDTA-free protease inhibitors

(Roche) and 1 mM PMSF, lysed by French press, centrifuged, and the supernatant was applied

to a 5 mL HisTrap FF crude column (Cytiva). After extensive washing with Ni^{2+} start buffer

containing 10 mM imidazole and 30 mM imidazole, protein was eluted using the same buffer

supplemented with 500 mM imidazole. The BRCA1^{RING}/BARD1^{RING} and BRCA1^{RING}/BARD1^{193}

constructs were concentrated and further purified using a 120 mL Superdex 75 column on an

AKTA FPLC system in SEC buffer (25 mM HEPES-NaOH pH 7.5, 150 mM NaCl, 1 mM DTT;

GE Healthcare). For the BRCA1^{RING}/BARD1^{216}, HisTrap elutions were dialyzed into ion-

exchange buffer (25 mM Tris-HCl pH 7.5, 250 mM NaCl, 1 mM DTT) and applied to a 5 mL

HiTrap SP HP column (Cytiva) and eluted over a 12 column-volume (CV) gradient (0.25-1 M

NaCl). Peak fractions were concentrated, aliquoted, and flash frozen.

1338

For Bpa crosslinking experiments, 6xHis-BRCA1$^{1-104}$-(GS)$_6$-BARD1$^{26-221}$ was sub-cloned using

Gibson assembly into a pET28n vector. Constructs for Bpa incorporation were generated via

site-directed mutagenesis to install amber stop codons in place of BARD1 L120, W146, Y180,

A195, and W218. Bpa-mutant Pet28n-6xHis-BRCA1$^{1-104}$-(GS)$_6$-BARD1$^{26-221}$ plasmids were co-

transformed with a pEVOL-pBpF plasmid encoding an Bpa aminoacyl tRNA synthetase (gift

from P. Schultz, Addgene #31190) in *E. coli* BL21 (DE3) cells(Chin Jason W. et al., 2002). At

OD$_{600nm}$=0.4-0.5, cells were shifted from 37 °C to 16 °C and supplemented with 1 mM Bpa

(dissolved in 1M NaOH; Bachem) and 100 µM ZnCl$_2$. After 30 minutes, protein expression was

induced by addition of 0.02% L-arabinose and 0.2 mM IPTG for ~16 hours at 16 °C. Proteins

were purified as described for the BRCA1^{RING}/BARD1^{216} construct. Truncations that did not

incorporate Bpa were separated via the SP column.

1350

BRCA1/BARD1 from insect cells: For insect cell expression, truncation and deletion constructs

were derived from full-length BRCA1 (pFastBac-FLAG-BRCA1$^{1-1863}$; gift from W. Zhao, UT

Health Sciences Center at San Antonio) and insect cell codon-optimized full-length BARD1

(pFastBac-Twin-StrepTagII-BARD1$^{1-777}$; gift from A. Deans, Addgene plasmid #137166)(Tan et

al., 2020) using Gibson assembly cloning (NEB). Baculovirus was generated using the Bac-to-

Bac system in suspension culture *Sf9* cells according to the manufacturers protocols

(Invitrogen). For general *Sf9* growth and virus amplification, SF900-II media (ThermoFisher)

was supplemented with 5% FBS (HyClone) and 1x antibiotic-antimycotic (Gibco). For protein

expression, 15 mL of BRCA1- and 15 mL of BARD1-containing P3 baculoviruses were added to

650 mL of *Sf9* cells at ~1.5x10^6 cells/mL in SF900-II media supplemented with 1% FBS and 1x

antibiotic-antimycotic. Protein expression was carried out for 48-72 hours at 27 °C, shaking at

110 RPM. Following expression, cells were spun down at 150 x g for 10 minutes. The pellet was

resuspended in 30 mL of Strep-start buffer (50 mM Tris-HCl pH 8, 10% glycerol, 150 mM NaCl,

1 mM DTT) supplemented with EDTA-free protease inhibitors and 1 mM PMSF and flash-

frozen. Following freeze-thawing in a 37 °C water bath, additional EDTA-free protease inhibitors

and PMSF were added in addition to 2 µL of benzonase (Sigma Aldrich). All subsequent steps

were performed on ice or in a cold room over a short timeframe to prevent proteolysis (<3 hours

total). Lysis was performed by sonication using a Branson Sonifier 250 (VWR Scientific) by

applying 2 rounds of 15 pulses at 70% duty on power level 8, ensuring that the cell mixture

stayed cold. PMSF was added throughout the lysis protocol. The lysed cells were centrifuged at

41,000 x g for 15 minutes, filtered through a 0.45 mM low protein-binding PVDF filter (Millipore)

and applied to a 1 mL StrepTrap HP column pre-equilibrated in Strep-start buffer (Cytiva). The

column was washed with 8 mL Strep-start buffer supplemented with protease inhibitors, 8 mL

wash buffer (25 mM Tris-HCl pH 7.5, 300 mM NaCl, 0.01% Igepal-CA630, 1 mM DTT, 5 mM

MgCl2, 2 mM ATP, 10% glycerol, EDTA-free protease inhibitors), and an additional 5 mL of

Strep-start buffer. The column was eluted using 6 mL of Strep-start buffer supplemented with 3

mM d-Desthiobiotin (Sigma). The elution was diluted with 20 mL of ion-exchange buffer (25 mM

HEPES-NaOH pH 7.5, 10% glycerol, 1 mM DTT) and applied to a 1 mL SP HP column pre-

equilibrated in ion-exchange buffer containing 75 mM NaCl. The SP column was eluted over 12-

CV (75-500 mM NaCl). Peak fractions were combined and concentrated using a 30K MWCO

concentrator (Amicon), and flash-frozen in small aliquots. Concentrations were estimated by

measuring absorbance at 280 nM using molar extinction coefficients obtained from ProtParam

(Expasy). Concentration estimates were verified by SDS-PAGE and Coomassie staining, and

slight adjustments in concentration were applied to best equalize the amounts of E3 in activity

and binding assays.

1386

BARD1 intrinsically disordered region: BARD1 124-270, 141-216, 269-424, and 124-424 were

sub-cloned from pFastBac-6xHis-BARD1[1-777] (gift from W. Zhao, UTHCSCA) into a modified

pET28 vector containing an N-terminal 6xHis-SUMO tag using Gibson assembly cloning.

Plasmids were transformed into BL21 (DE3) *E. coli*, grown at 37 °C in M9 media supplemented

with appropriate combination of isotopes (^{15}N-NH$_4$Cl, ^{13}C-glucose; Cambridge Isotopes) to an

OD$_{600nm}$ of 0.6-0.8, and induced with 0.2 mM IPTG for ~16 hours at 16 °C. HisTrap Ni^{2+}

purification was performed as described above for BRCA1/BARD1 constructs expressed in *E.*

coli. Elutions were dialyzed into ion exchange buffer (25 mM Tris pH 8.0, 200 mM NaCl, 0.5 mM

EDTA, 1 mM DTT) in the presence of GST-SENP1 (produced in-house) to cleave the SUMO

tag, leaving a non-native serine residue preceding the N-terminus except in the case of 269-

424. The cleaved product was applied at a HiTrap Q column preceding a HiTrap SP column in-

tandem to capture the non-specifically bound DNA (Cytiva). After application, the Q column was

removed, and the SP column was eluted using a 12-CV gradient (0.2-1M NaCl). The peak

fractions were combined, concentrated, and further purified using a Superdex 75 column

equilibrated in NMR buffer (25 mM MOPS-NaOH pH 7.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM

TCEP (tris(2- carboxyethyl)phosphine)).

1403

Histone purification: Wild-type and mutant human histones were transformed in *E. coli* BL21

(DE3) *pLysS* cells, grown at 37 °C to an OD$_{600nm}$ of 0.5-0.6, and induced with 0.5 mM IPTG for 2

hours (H4) or 3 hours (H2A-2A, H2B-1K, H3.2) at 37 °C. Histone octamers used for

fluorescence labeling were purified using the one-pot refolding protocol as described previously

using 6xHis-TEV-H2A(Lee et al., 2015; Witus, Burrell, et al., 2021). For nucleosome

ubiquitylation assays, a glycine within the TEV cleavage scar was mutated to a cysteine for

fluorophore conjugation (H2A G-1C) via site-directed mutagenesis. For fluorescence based

binding assays, a cysteine was introduced at H2B D51 via site-directed mutagenesis.

1412

For reconstituting chemically modified H2A-Ub nucleosomes, histones were purified individually

essentially as described(Luger et al., 1999). Briefly, inclusion bodies from 6-8L of histone

expression in LB media were lysed in buffer T (50 mM Tris-HCl pH 7.4, 1 mM EDTA, 100 mM

NaCl), washed by sonication twice in buffer TW (50 mM Tris-HCl pH 7.4, 1 mM EDTA, 100 mM

NaCl, 1% v/v Triton-X100), and then twice again in buffer T. Inclusion bodies were softened by

stirring with a small volume of DMSO (2-3 mL), and extracted with 60 mL of extraction buffer (25

mM Tris-HCl pH 7.5, 7 M guanidinium-HCl, 0.5 mM EDTA, 10 mM DTT). After centrifugation,

the supernatant was dialyzed overnight into ion-exchange buffer (25 mM Tris-HCl pH 7.5, 7 M

deionized urea, 100 mM NaCl, 0.5 mM EDTA, 10 mM DTT), and applied to 2x5 mL HiTrap Q

columns preceding 2x5 mL HiTrap SP columns in-tandem to capture the non-specifically bound

DNA (Cytiva). After sample application, the Q columns were removed, and the SP columns

were eluted using a 12-CV salt gradient (0.1-1M NaCl for H2A and H2B, 0.2-1M NaCl for H3

and H4). The peak fractions were assessed by Coomassie stained SDS-PAGE gel and

Abs_{260}/Abs_{280} value, and the purest fractions were extensively dialyzed into water, aliquoted,

flash frozen, and lyophilized. Lyophilized histones were reconstituted into H2A/H2B dimers,

H3/H4 tetramers, and octamers as previously described(Dyer et al., 2003), and purified via SEC

using Superdex 75 (H2A/H2B) or Superdex 200 (H3/H4 and octamers) columns in SEC buffer

(25 mM Tris-HCl pH 7.5, 2M NaCl, 1 mM DTT).

1431

Fluorophore conjugation to histones and ubiquitin: For fluorophore conjugation, Alexa Fluor 680

C_2 maleimide (ThermoFisher), IRDye 680LT maleimide (Li-Cor), and Oregon green 488

maleimide (AAT Bioquest) were reconstituted in DMSO to 10 mM. Fluorophore maleimides

were mixed with octamers containing single cysteine mutants H2A G-1C for H2A-Ub activity

assays (with Alexa Fluor 680 C_2 maleimide or IRDye 680LT maleimide) or H2B D51C for

binding assays (with Oregon green 488 maleimide). Conjugation reactions were performed

using 20 µM octamer and 100 µM fluorophore overnight at 4 °C in conjugation buffer (25 mM

Tris-HCl pH 7.5, 2M NaCl, 1 mM EDTA, 0.5 mM TCEP), and stopped by addition of 10 mM DTT

and flash frozen. Excess fluorophore was removed after nucleosome reconstitution by SEC

using a Superdex 200 increase 10/300 GL column or extensive buffer exchange in a 30K

MWCO concentrator (Amicon). Typical labelling efficiency was ~70%. For labelling of Ub Q2C,

100 µM Alexa Fluor 680 C_2 maleimide was mixed with ~1 mM of Ub Q2C in buffer (25 mM Tris-

HCl pH 7.5, 1 mM EDTA, 0.5 mM TCEP) at 4 °C overnight. The reaction was quenched with 10

mM DTT, and buffer exchanged into reducing buffer (25 mM Tris-HCl pH 7.5, 1 mM EDTA, 10

mM DTT) in a concentrator to remove unconjugated dye, and stored at -80 °C. The fluorophore

labelling efficiency of Ub was calculated to be ~5%.

1448

Ubiquitin G76C and dichloroacetone crosslinking of H2A-Ub: H2A K15C, K119C, and K127C

were generated by site-directed mutagenesis and purified as described above for individual

histones. His-TEV-ubiquitin G76C was a gift from C. Wolberger (Addgene plasmid #75299) and

purified as previously described(Morgan et al., 2019; Morgan Michael T. et al., 2016).

Dichloroacetone crosslinking of Ub G76C to H2A single cysteine mutants was performed as

previously described in detail (Morgan et al., 2019), with the exception that the final product

containing Ub, Ub-Ub dimers, and H2A-Ub was not purified by RP-HPLC to remove

unconjugated Ub. Instead, it was lyophilized, and mixed with a slight excess of H2B in unfolding

buffer (25 mM Tris-HCl pH 7.5, 7 M guanidinium-HCl, 0.5 mM EDTA, 1 mM DTT), refolded into

H2A/H2B dimers in refolding buffer (25 mM Tris-HCl pH 7.5, 2M NaCl, 1 mM DTT), and purified

by SEC using a 24 mL Superdex 75 column (GE Healthcare) to separate the H2A-Ub from Ub

species.

1461

'601' and MMTV DNA: For large scale purification of 147-bp '601' (gift from K. Luger, CU

Boulder), 185-bp '601' (gift from S. Tan, Penn State Univ.), MMTV (gift from G. Debelouchina,

(UC San Diego), and NLE-trimer '601' (for tri-NCP; gift from K. Luger) DNA, repeat plasmids

from 8-12L of DH5a cells in LB media were subjected to alkaline lysis, phenol-chloroform

extraction, PEG precipitation, EcoRV digestion (NEB), and ion exchange of the excised

fragment by HiTrap DEAE (147-bp '601', 185-bp '601', and MMTV; Cytiva) or mono-Q (NLE-

trimer '601'; GE Healthcare) according to established protocols(Dyer et al., 2003).

1469

For UV-induced Bpa crosslinking and EMSA experiments, fluorophore labelled '601' DNA was

generated using large-scale PCR with Phusion polymerase (produced in-house) from a pGEM-

3z/601 plasmid containing one copy of '601' DNA (gift from J. Widom, Addgene plasmid

#26656)(Lowary & Widom, 1998) with 5'-IRdye700 labelled forward primers (IDT). Following

PCR, the reactions were purified using a 1 mL mono-Q column equilibrated in ion-exchange

buffer (25 mM Tris-HCl pH 8.0, 200 mM NaCl, 1 mM EDTA) over a 10-CV gradient (0.2-1M

NaCl). Peak fractions containing purified 147- and 185-bp DNA fragments were dialyzed into

storage buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA), concentrated, and stored at -20 °C. The

primer sequences were as follows: 147-fwd: ctggagaatcccggtgccgagg; 147-rev:

acaggatgtatatatctgacacg; 185-fwd: atccctatacgcggccgccctggagaatcccggtgccgagg

; 185-rev: atcgctgttcaatacatgcacaggatgtatatatctgacacg.

1481

For generation of DNA fragments with DraIII sticky ends for dinucleosome assembly, large scale

PCR was performed as described for the fluorophore-labelled DNA fragments above. Primer

sequences containing DraIII cut-sites were based off those used by Poepsel *et al*. to make

dinucleosomes with 35-bp of linker DNA(Poepsel et al., 2018). Fragments used to reconstitute

nucleosomes containing fluorophore-labelled H2A nucleosomes (H2A[observe]) were generated

using the "147-fwd" primer (sequence above) and the di-NCP reverse primer sequence

(taggtatcgtatCACGGGGTGagatcgctacaggatgtatatatctgacacg). The fragment for the H2A[silent]

nucleosomes were generated using the "147-rev" and the di-NCP-forward primer sequence

(ctgacttattgaCACCCCGTGatgctcgatactgtcatactggagaatcccggtgccgag). The DraIII sites in the

primers are underlined and uppercase. Following PCR purification by 1 mL mono-Q, DNA

fragments were dialyzed into TE/0.1 buffer (10 mM Tris-HCl, 0.1 mM EDTA) and digested with

DraIII-HF in 1x CutSmart buffer for ~20 hr at 37 °C (NEB). Fragments with DraIII sticky ends

were re-purified by 1 mL mono-Q ion-exchange, dialyzed into storage buffer (10 mM Tris-HCl, 1

mM EDTA), concentrated, and stored at -20 °C.

1496

Nucleosome reconstitution: mono-nucleosomes (NCPs) were reconstituted by the standard salt-

dialysis method by mixing histone octamers (or H3/H4 tetramers and H2A/H2B dimers) with

'601' DNA in high salt buffer (25 mM Tris-HCl pH 7.5, 2M NaCl, 0.1 mM EDTA, 1 mM DTT) in a

slide-a-lyzer mini dialysis unit (0.1 mL, ThermoFisher)(Dyer et al., 2003). Low-salt buffer (25 mM

Tris-HCl pH 7.5, 0.1 mM EDTA, 1 mM DTT) was pumped in over ~24 hours in a cold room with

stirring. A final dialysis was performed into NCP storage buffer (25 mM HEPES-NaOH, 10 mM

NaCl, 0.1 mM EDTA, 1 mM DTT). Fluorophore-labelled nucleosomes were further purified by

SEC using a 24 mL Superdex 200 increase 10/300 column equilibrated in NCP storage buffer or

by extensive buffer exchange in a concentrator to remove excess fluorophore. For tri-NCPs, salt

dialysis was performed as described for mono-NCPs using ~1.4x octamer to '601' sites in the

presence of 0.2 equivalents of MMTV competitor DNA. Following dialysis into TEK10 (10 mM

Tris-HCl pH 7.5, 0.1 mM EDTA, 10 mM KCl), trimers were precipitated by adding 4 mM $MgCl_2$,

incubating on ice for 10 minutes and pelleted by centrifugation at 17,000 x g at 4 °C. The

supernatant was discarded, and the pellet was gently resuspended in TEK10, and dialyzed

overnight into NCP storage buffer. Tri-NCP assembly was verified by EcoRI digestion of sites

located between nucleosome units.

1513

For di-NCP assembly, mono-NCPs with DraIII sticky ends were assembled as described above.

The strategy to make asymmetric di-NCPs was based on previously described methods(Dao et

al., 2020; Poepsel et al., 2018). 100 nM of each NCP was mixed in T4 ligase buffer (50 mM

Tris-HCl pH 7.5, 10 mM MgCl$_2$, 1 mM ATP, 10 mM DTT) on ice. T4 ligase (NEB) was added to a

final concentration of 20 U/μL, the reaction was incubated for 30 minutes at 16 °C, and dialyzed

into TEK10 for 4 hours at 4 °C. Following dialysis, MgCl$_2$ was added to a final concentration of

18 mM, incubated at room temperature for 15 minutes, and pelleted at 17,000 x g at 4 °C. The

pellet containing di-NCPs was resuspended in 50 μL of TEK10 and dialyzed overnight into NCP

storage buffer. All chromatin substrates were analyzed for quality on 5% polyacrylamide 0.5x

TBE gels monitoring fluorescence and/or DNA-staining, ensuring minimal free DNA or

contaminating species. Chromatin substrates were stored on ice for no longer than one month.

1525

Design and assembly of DNA competitor fragments: single-stranded, double-stranded, bubble,

frayed, and single-stranded overhang DNA fragments were assembled by annealing ssDNA

oligos (IDT) in IDT duplex buffer (30 mM HEPES-NaOH pH 7.5, 100 mM potassium acetate).

Oligo mixtures were heated to 95 °C and slowly cooled to 25 °C stepwise over 1 hour in a

thermocycler. For NMR experiments, dsDNA and bubble-DNA were buffer exchanged into NMR

buffer in a spin concentrator (Amicon). The sequences of oligos used are as follows. For

ssDNA, dsDNA, bubbleDNA, and ssOverhang a common "bottom" oligo was used

(ggtacacaattgcgctggtaccccaggcgtcgtagg). The oligos annealed to the "bottom" oligo were as

follows: (dsDNA-top: cctacgacgcctggggtaccagcgcaattgtgtacc; bubbleDNA-top:

cctacgacgcctggctctttcccgcaattgtgtacc; ssOverhang-top: ccgcaattgtgtac). For the frayed DNA,

the annealed oligos were as follows: (frayedDNA-top: ggtacacaattgcgccaggcgtcgtaggctggtacc;

frayedDNA-bottom: ctctttcccctacgacgcctggcgcaattgtgtacc).

1538

Nucleosome ubiquitylation assays

Time-course H2A-Ub assays: The general setup for time-course H2A-Ub assays was a reaction

mixture containing 0.2 μM E1 (UBA1), 1 μM E2 (UBE2D3), 30 μM ubiquitin, 0.5 μM NCP

substrate, and 12.5-100 nM E3 (depending on the assay), assembled in 30 µL of reaction buffer

(25 mM HEPES-NaOH pH 7.5, 150 mM NaCl). For assays containing di-NCPs, 10 nM of

fluorophore-labelled di-NCP substrate was mixed with 0.5 µM unlabeled NCP[147] substrate in the

same reaction tube to facilitate quantifiable reaction kinetics. Critically, the salt concentration of

each reaction within an assay was rigorously controlled, as H2A-Ub kinetics are extremely

sensitive to differences in buffer ionic strength. 5 µL of each reaction was mixed 1:1 with 2x

SDS-PAGE load dye as a zero time-point. The mixtures were brought to 37 °C in a heat block,

and 1 µL of 100 mM $MgCl_2$/ATP was added (4 mM final concentration) to initiate the reaction,

and the indicated time-points were obtained by diluting 5 µL the reaction 1:1 with 2x SDS-PAGE

load dye. For experiments with tri-NCPs, the concentration of nucleosome units in the reaction

was normalized to NCP[147] and NCP[185] (0.5 µM NCP unit or 0.17 µM tri-NCP), and reactions

were initiated with 2 mM ATP and 1 mM $MgCl_2$.

1554

Gel samples were run on 15% SDS-PAGE gels and visualized by in-gel fluorescence monitoring

a fluorophore conjugated to the N-terminus of H2A. Alternatively, some assays were visualized

by Western blot for H2A (EMD Millipore, 07-146) or a VSVG epitope tag on the N-terminal tag of

H2A (Sigma, V4888) and a fluorescently labelled secondary antibody (Cell Signaling, 5151S).

Gels and blots were imaged on a Li-Cor Odyssey imager (Li-Cor). Band quantification was

performed using Image Studio software (Li-Cor) by drawing a box encompassing the unmodified

H2A band to measure its intensity, performing background subtraction, and normalizing to the

zero time-point of a given reaction.

1563

Inhibition of H2A-Ub by competitor DNA: A reaction mixture containing 0.2 µM E1 (UBA1), 3 µM

E2 (UBE2D3), 30 µM ubiquitin, 0.5 µM NCP substrate, and 50 nM E3 (or 750 nM for

BRCA1[RING]/BARD1[RING]) was assembled in 9 µL of reaction buffer (25 mM HEPES-NaOH pH

7.5, 100 mM NaCl) in PCR strips. To that, 1 µL of DNA competitor at 10x concentration was

added. Reactions were brought to 37 °C in a thermocycler, started by addition of 5 µL of 10 mM

ATP/MgCl$_2$ in reaction buffer, and allowed to proceed for 12-20 minutes depending on the E3.

Reactions were stopped by addition of 10 µL of 4x fluorescence sample load dye (Li-Cor),

resolved on a 15% SDS-PAGE gel, and visualized using a Li-Cor Odyssey imager.

Quantification was performed as described above, monitoring the intensity of unmodified H2A,

and normalizing each reaction to a -ATP reaction.

1574

Auto-ubiquitylation assay: Reactions were assembled with 0.2 µM E1 (UBA1), 1 µM E2

(UBE2D3), 10 µM ubiquitin (Q2C-Alexa Fluor 680), and 0.2 µM of the indicated E3 in buffer (25

mM HEPES-NaOH pH 7.5, 50 mM NaCl), and were initiated by addition of 5 mM ATP/MgCl$_2$.

After 60 minutes at 37 °C, reactions were quenched by mixing 1:1 with 4x fluorescence sample

load dye (Li-Cor), resolved on a 4-20% SDS-PAGE gel (BioRad), and visualized using a Li-Cor

Odyssey imager.

1581

Nucleosome binding assays

Electrophoretic mobility shift assay: For EMSA experiments, 5 µL of 10 nM of substrate (NCPs

with either H2A G-1C labelled with Alexa Fluor 680 or DNA/NCPs labelled with IRDye700) were

mixed with 5 µL of indicated RING/RING heterodimer or BARD1 IDR constructs (2x

concentrated) in EMSA buffer (15 mM Tris-HCl pH 7.5, 90 mM NaCl, 1 mM DTT, 0.1 mg/mL

BSA, 0.05% triton X-100) in PCR strips. Binding was allowed to proceed for 10 minutes at room

temperature, followed by 5 minutes on ice. 5 µL of 25% sucrose was added to each tube, and

15 µL of sample was loaded onto 5% polyacrylamide native gels (0.5x TBE) in pre-chilled 0.5x

TBE buffer. Gels were run at 4 °C for 90-120 minutes at 110V and visualized using a Li-Cor

Odessey scanner (Li-Cor).

1592

Fluorescence-quenching binding experiments: Assay set up was performed essentially as

previously described(Hu et al., 2021). 10 µL of 10 nM NCPs with Oregon green 488 conjugated

to H2B D51C in low-salt fluorescence buffer (25 mM Tris-HCl pH 7.5, 10 mM NaCl, 0.01%

CHAPS, 0.01% NP40, 0.1 mg/ml BSA, 1 mM DTT) were mixed with the 10 µL of the indicated

BRCA1/BARD1 construct at 2x concentration in high-salt fluorescence buffer containing either

90 mM, 190 mM, or 290 mM NaCl, depending on the assay, in a 384-well low volume black

round bottom polystyrene non-binding surface microplate (Corning). Plates were briefly mixed,

incubated at 22 °C for 10-15 minutes, and scanned using a BioTek Synergy Neo2 plate reader

equipped with a fluorescence filter (excitation: 485/20 nm, emission: 528/20 nm). The gain was

set to 100 for all experiments. Two to three technical replicates for each independent replicate

experiment (fresh NCP and BRCA1/BARD1 dilutions) were performed, and the readings from

two scans were averaged. BRCA1/BARD1/nucleosome binding interactions were quantitatively

analyzed by fluorescence quenching assuming the following systems of equations:

$$\theta = \frac{K_D + [NCP] + [BCBD] - \sqrt{(K_D + [NCP] + [BCBD])^2 - 4 \times [NCP] \times [BCBD]}}{2 \times [NCP]}$$

$$F = [NCP] \times F_{NCP} \times (1 - q_{BCBD} \times \theta) + N(0, error)$$

where θ is the fraction of nucleosome bound by BRCA1/BARD1 (BCBD), F_{NCP} is the

fluorescence per nucleosome, q_{BCBD} is the fraction of fluorescence quenching per

BRCA1/BARD1 binding, and fluorescence measurements are gaussian distributed with mean 0.

Maximum likelihood parameter estimates were determined using custom Python scripts with

optimization by a differential evolution algorithm implemented in Scipy(Virtanen et al., 2020). K_d

and q_{BCBD} were separately estimated for each BRCA1/BARD1 construct, and NCP

concentration was treated as a random effect per individual dilutions from higher concentration

stock solutions, with the total number of model parameters being equal to number of NCP

dilutions + (2 x number of BCBD constructs) + 2. The 95% confidence limits were estimated by

computing the profile likelihood with an applied Bonferroni correction.

1618

UV-induced Bpa crosslinking

NCPs and DNA for UV-induced crosslinking assays were reconstituted using a 185-bp '601'

sequence labeled with an IRdye700 fluorophore (described above). 50 nM NCPs or DNA were

incubated with 500 nM of the BRCA1$^{1-104}$-(GS)$_6$-BARD1$^{26-221}$ fusion (wild-type or with different

Bpa-incorporated sites L120Bpa, W146Bpa, Y180Bpa, A195Bpa, W218Bpa) for 15 minutes at

4°C in 30 uL UV Crosslinking buffer (20 mM HEPES-NaOH pH 7.5, 100 mM NaCl) in a 96-well

clear round bottom polystyrene non-treated microplate (Corning). Following incubation, 5 µL

was added to 5 µL 2X SDS-PAGE gel-loading dye for the -UV sample. Samples were then

subjected to UV irradiation (365 nm) using a Blak-Ray B-100AP/R High Intensity UV Lamp

(UVP) from 20 cm for 30 minutes at 4°C. Following irradiation, 5 µL was added to 5 µL of 2x

SDS-PAGE gel-loading dye. Samples were heated at 60°C for 5 minutes and separated on 10%

SDS-PAGE gels. Gels were visualized using a Li-Cor Odyssey scanner (Li-Cor). Band

quantification was performed using Image Studio software (Li-Cor) by drawing a box measuring

the total band intensity for each crosslinked product, subtracting the background signal, and

normalizing to the crosslinked band of the L120Bpa fusion mutant. For the Bpa UV crosslinking

DNA competition assays, 50 nM NCPs were incubated with 500 nM BRCA1-f-BARD1^{221}

(W146Bpa or A195Bpa) with varying amounts of competitor 36-mer dsDNA or bubble-DNA (0,

0.3125, 0.625, 1.25, 2.5, 5, or 10 µM) for 15 minutes at 4°C, and assays were performed as

described above.

1638

Chemical Crosslinking and Mass Spectrometry Analysis

Sample preparation: Reactions were 90 µL of protein/nucleosome mix containing 2 mM

BRCA1^{RING}/BARD1^{FL} and 2 mM of unmodified H2A or H2A K15-Ub nucleosomes wrapped with

147-bp '601' DNA in crosslinking bufffer (25 mM HEPES-NaOH pH 7.5, 150 mM NaCl, 1%

glycerol, 1 mM DTT) plus 3.06 µL of 14.5 mM DSS or BS3 (final concentration 0.5 mM) (DSS:

disuccinimidyl suberate, BS3: bis(sulfosuccinimidyl)suberate, ThermoFisher). Crosslinking was

carried out for 10, 30 or 45 mins at room temperature before removing 30 µL and quenching by

addition of 3 µL of 1 M ammonium bicarbonate at room temperature for 30 minutes. Four

separate reaction conditions were performed: (1) WT-NCP DSS; (2) WT-NCP BS3; (3) K15Ub-

NCP DSS; (4) K15Ub-NCP BS3. After quenching, 9 µL of each sample at each timepoint was

analyzed by SDS-PAGE **(see Extended Data Fig 10a)** and the remainder was stored at -80°C

until processing for mass spectrometry analysis.

1651

Reactions were prepared for MS analysis by bringing them up to 0.1% PPS silent surfactant

(Expedion Inc.), 5 mM TCEP. Samples were reduced for 60 mins at 60°C in an Eppendorf

Thermomixer with shaking (1200 rpm). Alkylation was performed at room temperature in the

dark for 30 mins with 6 mM iodoacetamide. Excess iodoacetamide was quenched by addition of

5 mM DTT and samples were then digested by trypsin digestion (Sequencing Grade Modified

Trypsin, Promega Corp) at 37°C for 6 hours in a Thermomixer with shaking (1000 rpm) at a

substrate to enzyme ratio of 15:1 prior to acidification with 250 mM HCl (final concentration).

Acidified samples were incubated for 1 hour at room temperature, spun at max speed in a

microfuge and supernatants containing peptides were transferred to autosampler vials and

stored at -80°C until analysis.

1662

Chromatography: Mass spectrometry and data analysis was based on previously described

methods(Zelter et al., 2015). For each injection, 3 µl of protein digests were loaded by

autosampler onto a 150-µm Kasil fritted trap packed with 2 cm of Reprosil-Pur C18-AQ (3-µm

bead diameter, Dr. Maisch) at a flow rate of 2.5 µl/min. After desalting with 8 µl of 0.1% formic

acid plus 2% acetonitrile, the trap was brought online with a Self-Packed PicoFrit Column (New

Objective part number PF360-75-10-N-5, 75 µm i.d.) packed with 30 cm of Reprosil-Pur

C18-AQ (3-µm bead diameter, Dr. Maisch) mounted to a heated nanospray ionization source

(CorSolutions LLC) set at 50°C and placed in line with a Thermo Scientific EASY-nLC 1200

UPLC pump plus autosampler.

1672

1673 Peptides were eluted from the column at 0.25 µL/min using an acetonitrile gradient consisting of

1674 the following steps: (1) 0-10 mins; 6-10% B; (2) 10-90 mins; 10-32% B; (3) 100-130 mins; 32-

1675 75% B; (4) 130-135 mins; 75% B; (5) 135-136 mins; 75-100% B; (6) 136-151 mins; 100% B,

1676 followed by re-equilibration to 0% buffer B prior to the subsequent injection. Buffer A was: 0.1%

1677 formic acid in water and buffer B was 0.1% formic acid 80% acetonitrile.

1678

Data acquisition: A Thermo Fisher Scientific Exploris 480 was used to perform mass

spectrometry in positive ion mode with the following settings. Data dependent acquisition (DDA)

mode was used with a maximum of 20 tandem MS (MS/MS) spectra acquired per MS spectrum

(scan range of *m/z* 400–1,600). The resolution for MS and MS/MS was 60,000 at *m/z* 200. The

normalized automatic gain control targets for both MS and MS/MS were set to 100%, and the

maximum injection times were 50 and 100 ms for MS and MS/MS scans, respectively. MS/MS

spectra were acquired using an isolation width of 2 *m/z* and a normalized collision energy of 27.

MS/MS acquisitions were prevented for +1, +2, ≥+6 or undefined precursor charge states.

Dynamic exclusion was set for 10 seconds. MS and MS/MS spectra were collected in centroid

mode.

1689

MS data processing – identification of crosslinked peptides: Raw mass spectra were converted

into mzML using ProteoWizard's msconvert(Chambers et al., 2012). All proteins in the sample

were identified using Comet(Eng et al., 2013) searching against a database including the entire

S. frugiperda, *E. coli* and *H. sapiens* proteomes (Uniprot 7108, 83333 and 9606, respectively)

plus common contaminants (https://www.thegpm.org/crap/) along with the sequences of all

heterologously expressed proteins (FLAG-BRCA1, P38398; TwinStrepII-BARD1, Q99728; H2A-

2A, Q99728; H2B-1K, O60814 ; H3.2 (C110A), Q71Dl3; H4, P62805; Ub G76C, sequence

derived from P0CG48). Smaller search databases were made for subsequent XL searching

consisting only of proteins identified in initial comet searches by at least 3 peptides with a

Percolator assigned q-value of ≤ 0.01. Decoy databases consisted of the corresponding set of

reversed protein sequences. Cross-linked peptides were identified within these proteins by

Kojak version 2.0.0-alpha8 available at (http://www.kojak-ms.org/)(Hoopmann et al., 2015). A

statistically meaningful q-value was assigned to each peptide spectrum match (PSM) through

analysis of the target and decoy PSM distributions using Percolator version 2.08(Käll et al.,

2007). All data reported in this paper were filtered to show hits to the target proteins that had a

Percolator assigned peptide level q value ≤ 0.05.

1706

NMR experiments and resonance assignments.

All spectra were collected at 25° C in NMR buffer (25 mM MOPS-NaOH, 100 mM NaCl, 1 mM

EDTA, and 0.5 mM TCEP, pH 7.0) with 7% D_2O. NMR HSQC and titration data were completed

on a Bruker 800 MHz Avance III spectrometer with cyroprobe. ($^1H,^{15}N$)-HSQC spectra were

collected on the following ^{15}N-BARD1 IDR constructs: BARD1 Ser-124-424, Ser-124-270, 269-

424, and Ser-141-216.

1713

($^1H,^{15}N$)- and ($^1H,^{13}C$)-HSQC titration experiments were completed for BARD1 IDR + 36mer-

dsDNA or 36mer-bubDNA by adding unlabeled DNA to 150 µM [$^{13}C,^{15}N$]-BARD1 124-270 or

150 µM [$^{13}C,^{15}N$]-BARD1 Ser-141-216, maintaining a constant concentration of the labeled

species. Only apo and 1:1 spectra for the [$^{13}C,^{15}N$] BARD1 Ser-141-216 construct were

obtained since intermediate titration points resulted in loss of sample. Additionally, titration

spectra with nucleosome were collected for 150 µM [$^{13}C,^{15}N$]-BARD1 124-270 + 0.05x NCP[147].

A ($^1H,^{15}N$)-HSQC was collected on 150 µM ^{15}N-BARD1 269-424 + 150 µM 28mer-dsDNA.

1721

Dynamics experiments were collected on a Bruker 500 MHz Avance III spectrometer with a

room temperature probe. T_1 and T_2 ^{15}N-Trosy HSQC experiments for 150 µM [^{13}C,^{15}N]-BARD1

Ser-141-216, 150 µM [^{13}C,^{15}N]-BARD1 Ser-141-216 + 150 µM 36mer-dsDNA, and 150 µM

[^{13}C,^{15}N]-BARD1 Ser-141-216 + 150 µM 36mer-bubble-DNA were collected in an interleaved

manner with 8 points each(G. Zhu et al., 2000). T_1 delays were 10, 40, 80, 120, 160, 320, 640,

and 1000 ms; T_2 cpmg loop delays were 8.48, 16.96, 25.44, 25.44, 42.40, 50.88, 67.84, and

84.80 ms. T_1 and T_2 for each residue were fitted to a single exponential with errors reflecting the

quality in the fit. Residues corresponding to peaks with overlapping intensities were excluded

from the analysis.

1731

Backbone chemical shift assignments were determined from standard backbone triple-

resonance experiments (HNCA, HNCOCA, HNCOCACB, HNCACB, HNCO, and HNCACO) and

an HNHA experiment using a Bruker 800 MHz Avance III spectrometer with cryoprobe. The 3D

data were collected for 500 µM [^{13}C,^{15}N]-BARD1 Ser-124-270 and 500 µM [^{13}C,^{15}N]-BARD1

Ser-141-216. Assignments from the shorter construct were transferred as appropriate to the

longer BARD1 construct. HNCACB and HNCOCACB spectra were collected on the 1:1 samples

of 150 µM [^{13}C,^{15}N]-BARD1 Ser-141-216 with either 36mer dsDNA or 36mer bubble-DNA.

1739

The NH chemical shift perturbation was calculated according to $\Delta\delta_{NH}$ (ppm) = sqrt [$\Delta\delta_H^2$ +

$(\Delta\delta_N/5)^2$]. Component ^{1}H and ^{15}N chemical shift differences are (free – bound) in ppm. The

CaCb CSP was calculated according to $\Delta\delta_{CaCb}$ (ppm) = 0.25*[(Cα-Cβ)$_{free}$-(Cα-Cβ)$_{bound}$].

Secondary structure propensity (SSP) was determined from $\Delta\delta$(Cα-Cβ) = (Cα-Cβ)$_{measured}$ - (Cα-

Cβ)$_{random\ coil}$, where the random coil shifts were generated from BARD1 IDR sequence using a

webserver (https://spin.niddk.nih.gov/bax/nmrserver/Poulsen_rc_CS/)(Kjaergaard & Poulsen,

2011).

1747

Multiple Sequence Alignment

Mammalian BARD1 ortholog sequences were downloaded from Ensembl (v106), globally

aligned using MAFFT(Katoh & Standley, 2013), and trimmed to BARD1 residues 124-270.

Materials, Data, and Code availability Statement

Source data for H2A-Ub and binding assays as well as uncropped gels and blots are included

as supplemental files. For crosslinking mass-spectrometry analysis, the complete unfiltered list

of all PSMs and their Percolator assigned q values are available on the ProXL web

application(Riffle et al., 2016, 2019) https://proxl.yeastrc.org/proxl/p/bard1nucleosome along

with the raw MS spectra and search parameters used. In addition, complete search algorithm

configuration files, fasta search databases, raw search output and raw MS data files were

deposited to the ProteomeXchange Consortium via the PRIDE(Vizcaíno et al., 2009) partner

repository with the dataset identifier PXD035345. NMR chemical shift assignments for BARD1

Ser-141-216 and associated DNA-bound assignments have been deposited to the Biological

Magnetic Resonance Bank with dataset identifiers 51523 (apo), 51524 (with dsDNA), 51525

(with bubble-DNA). Python scripts used to analyze and plot fluorescence-based binding data

are included with this manuscript. Plasmid reagents generated in this study can be obtained

upon request from the corresponding author Dr. Rachel Klevit.