BRCA1/BARD1 intrinsically disordered regions facilitate chromatin recruitment and ubiquitylation.

Samuel R. Witus[1], Lisa M. Tuttle[1,#], Wenjing Li[4,#], Alex Zelter[1], Meiling Wang[4], Klaiten E. Kermoade[1], Damien B. Wilburn[2,3], Trisha N. Davis[1], Peter S. Brzovic[1,*], Weixing Zhao[4,*], Rachel E. Klevit[1,*]

1. Department of Biochemistry, University of Washington, Seattle, WA.
2. Department of Genome Sciences, University of Washington, Seattle, WA.
3. Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH.
4. University of Texas Health Science Center at San Antonio, Department of Biochemistry and Structural Biology, San Antonio, TX.
* Corresponding author email: brzovic@uw.edu, zhaow2@uthscsa.edu, klevit@uw.edu
# Equal contributions

Abstract

BRCA1/BARD1 is a tumor suppressor E3 ubiquitin (Ub) ligase with roles in DNA damage repair and in transcriptional regulation. BRCA1/BARD1 RING domains interact with nucleosomes to facilitate mono-ubiquitylation of distinct residues on the C-terminal tail of histone H2A. These enzymatic domains constitute a small fraction of the heterodimer, raising the possibility of functional chromatin interactions involving other regions such as the BARD1 C-terminal domains that bind nucleosomes containing the DNA damage signal H2A K15-Ub and H4 K20me0, or portions of the expansive intrinsically disordered regions found in both subunits. Herein, we reveal novel interactions that support robust H2A ubiquitylation activity mediated through a high-affinity, intrinsically disordered DNA-binding region of BARD1. These interactions support BRCA1/BARD1 recruitment to chromatin and sites of DNA damage in cells and contribute to their survival. We also reveal distinct BRCA1/BARD1 complexes that depend on the presence of H2A K15-Ub, including a complex where a single BARD1 subunit spans adjacent nucleosome units. Our findings identify an extensive network of multivalent BARD1-nucleosome interactions that serve as a platform for BRCA1/BARD1-associated functions on chromatin.

Introduction

Mutations in *BRCA1* and *BARD1* increase lifetime risk of breast and ovarian cancer. BRCA1 and BARD1 form a large, obligate heterodimeric complex (BRCA1/BARD1) that has distinct roles in DNA double-stranded break (DSB) repair by homologous recombination (HR), transcriptional regulation, and several other nuclear processes (Densham & Morris, 2019; Mullan et al., 2006; Tarsounas & Sung, 2020). These functions are mediated, at least in part, through a direct association with chromatin. Upon DNA damage, BRCA1/BARD1 is recruited to damaged chromatin where it segregates with DNA repair factors (Kolas Nadine K. et al., 2007; Mattiroli et al., 2012; Scully et al., 1997; Sobhian Bijan et al., 2007). Cancer-predisposing mutations that disrupt BRCA1/BARD1 functions on chromatin lead to DNA-damage hypersensitivity and genomic instability (Alenezi et al., 2020; Gudmundsdottir & Ashworth, 2006; Krais & Johnson, 2020). BRCA1/BARD1 also has distinct roles in both stimulation and repression of transcription through as yet incompletely defined mechanisms (Mullan et al., 2006). Full knowledge of how BRCA1/BARD1 interfaces with chromatin, and where it exerts its major biological functions, is critical for understanding the etiology of BRCA1/BARD1-mutant associated breast and ovarian cancer.

Nucleosomes, the organizing unit of chromatin, directly interact with BRCA1/BARD1 through multiple direct binding interfaces (Witus et al., 2022). The N-terminal RING domains of BRCA1 and BARD1 form a heterodimer that constitute a RING-type E3 ubiquitin (Ub) ligase and provide the only known enzymatic activity of BRCA1/BARD1 (Brzovic et al., 2001; Lorick et al., 1999; Witus, Stewart, et al., 2021). The RING heterodimer binds to the histone surface of one pseudo-symmetrical 'face' of a nucleosome to facilitate site-specific transfer of mono-Ub to a cluster of lysine residues on the extreme C-terminal tail of canonical histone H2A variants (K125, K127, and K129; collectively referred to henceforth as K127) (Hu et al., 2021; Kalb et al., 2014; Witus, Burrell, et al., 2021). H2A lysine specificity is largely determined by the RING domain of BARD1.

This domain binds to a unique histone surface compared to structurally similar Ub ligase

complexes that target other H2A lysine residues (e.g., K119 by RING1B/BMI1) (McGinty et al.,

2014). The H2A ubiquitylation ("H2A-Ub") activity of BRCA1/BARD1 is thought to promote long-

range resection of broken DNA ends during DSB repair via HR (Densham et al., 2016;

Uckelmann et al., 2018). BRCA1/BARD1-dependent H2A-Ub activity is also implicated in the

transcriptional repression of certain estrogen-metabolizing cytochrome P450 genes and the

constitutive repression of α-satellite DNA regions (Stewart et al., 2018; Thapa et al., 2022; Q.

Zhu et al., 2011, 2018).

The BARD1 C-terminal domains (CTDs; **Figure 1A**) also bind to nucleosomes to facilitate DNA

DSB repair (Becker et al., 2021; Dai et al., 2021; Hu et al., 2021; Krais et al., 2021; Nakamura et

al., 2019; Sherker et al., 2021). The BARD1 CTDs specifically recognize nucleosomes that

contain H2A K15-Ub (the product of the Ub ligase RNF168) and unmethylated H4 K20

(H4K20me0). Together, these signals serve as a binding platform for the recognition of

damaged chromatin in S/G2 phases when a newly replicated sister chromatid is available to

template high-fidelity DNA repair via HR. The BARD1 CTDs and N-terminal BRCA1/BARD1

RING domains bind to histone surfaces that overlap, precluding their simultaneous binding to

one nucleosome face. Despite this, mono-nucleosomes with ubiquitin pre-installed at both

copies of H2A K15 residues are better substrates for BRCA1/BARD1-dependent ubiquitylation

of H2A K127 *in vitro* (Hu et al., 2021). This observation implies that both domains may bind to

the same nucleosome unit, with each occupying one face. Additionally, regions throughout the

expansive intrinsically disordered segments of both BRCA1 and BARD1 are reported to bind to

DNA, but the functional significance of such interactions in chromatin binding and H2A-Ub

activity remains to be determined (Mark et al., 2005; Masuda et al., 2016; Simons et al., 2006;

Zhao et al., 2017).

Although a minimal RING/RING heterodimer is sufficient to direct the transfer of mono-Ub to

H2A K127 in nucleosomes, full-length BRCA1/BARD1 heterodimers form stronger complexes

with unmodified nucleosome substrates and exhibit increased H2A-Ub activity (Hu et al., 2021;

Witus, Burrell, et al., 2021). This suggests that additional interactions between full-length

BRCA1/BARD1 and nucleosomes can occur in the absence of the DNA-damage specific H2A

K15-Ub histone marks. Because BRCA1/BARD1 performs diverse biological functions in the

nucleus including, but not limited to, DNA repair via HR, transcriptional regulation, stalled

replication fork protection, centrosome regulation, R-loop resolution, and cell-cycle regulation, it

is likely that the complex can be recruited to chromatin via signals not limited to H2A K15-Ub. In

pursuit of a wholistic understanding of BRCA1/BARD1 recruitment to chromatin, we sought to

characterize interactions with nucleosomes that lead to enhanced chromatin binding and H2A-

Ub activity in both the absence and presence of H2A K15-Ub marks.

Here, we identify an intrinsically disordered region of BARD1 adjacent to its RING domain that

binds strongly to both nucleosomal and extra-nucleosomal DNA, dramatically enhancing the

affinity of the complex and its H2A-Ub activity. The interactions can be modulated by specialized

DNA structures that are derived from the DNA damage repair process and can compete directly

with BARD1 binding to nucleosomal DNA. We incorporate our findings into a molecular

mechanism of recognition and establish a role for BARD1-DNA interactions in chromatin

recruitment and DNA damage repair in cells. Additionally, we provide evidence for multiple,

distinct higher-order chromatin complexes that contain H2A K15-Ub nucleosomes. These

include both a 'wrapped' complex with the BRCA1/BARD1 RING domains and BARD1 CTDs

bound to opposite sides of one nucleosome unit, as well as an extended complex where the

domains bind to adjacent nucleosome units. Throughout, we evaluate the contribution of

BARD1 DNA binding to nucleosome affinity and H2A-Ub activity. Our findings reveal a network

of multivalent BARD1-nucleosome interactions that serve as a platform for BRCA1/BARD1-

associated functions on chromatin and hint at novel modes of DNA recognition by intrinsically

disordered regions of proteins.

Results

BARD1 IDR supports increased H2A ubiquitylation activity

The BRCA1/BARD1 heterodimer contains both structured domains and long intrinsically

disordered regions (IDRs**; Figure 1A)**. A minimal complex composed of the N-terminal RINGs

of BRCA1 (1-112) and BARD1 (26-140; BRCA1^{RING}/BARD1^{RING}) is sufficient to direct site-

specific mono-ubiquitylation of nucleosomal histone H2A at K127 (Kalb et al., 2014). We and

others have observed that full-length BRCA1/BARD1 (~300 kDa) exhibits faster H2A-Ub

kinetics and stronger binding to unmodified mono-nucleosomes than the minimal RING/RING

complex (~25 kDa; **Supplemental Figure 1A-D**) (Hu et al., 2021; Witus, Burrell, et al., 2021).

To identify regions responsible for increased H2A-Ub activity and binding affinity, we generated

heterodimers with truncations in either BRCA1 or BARD1 **(Supplemental Figure 2 and**

Extended Data Table I).

Heterodimers containing full-length BARD1 and three BRCA1 constructs were compared to

survey BRCA1 regions that may enhance H2A-Ub activity: full-length BRCA1 (BRCA1^{FL}), a

clinically relevant allele missing ~1100 residues in the expansive IDR (BRCA1$^{\Delta 11q}$), and the

minimal BRCA1 RING fragment **(**BRCA1^{RING}**)**. Notably, in the context of BARD1^{FL}, the RING

domain of BRCA1 is sufficient to generate fast H2A-Ub kinetics **(Supplemental Figure 1C)**.

The presence of additional regions of BRCA1 do not further enhance activity (as in

BRCA1$^{\Delta 11q}$/BARD1^{FL}) or appear to inhibit activity (e.g., BRCA1^{FL}/BARD1^{FL}). These results

indicate that the BARD1 subunit is largely responsible for enhanced H2A-Ub activity. In the

context of heterodimers that contain only the RING of BARD1, modest activity enhancements

are afforded by inclusion of additional BRCA1 regions, suggesting that, in the absence of FL-

BARD1, regions of BRCA1 may provide some redundancy in function **(Supplemental Figure**

1D).

To identify regions of BARD1 that contribute to increased H2A-Ub activity, the H2A-Ub kinetics

and nucleosome binding affinity of BRCA1^{RING}-heterodimers containing BARD1 truncations

were compared. Notably, BRCA1^{RING}/BARD1^{FL} heterodimers bind nucleosomes with ~100-fold

higher affinity than minimal BRCA1^{RING}/BARD1^{RING} and have markedly higher H2A-Ub activity

(Figure 1B, C). Removal of the BARD1 C-terminal domains (BARD1$^{\Delta CTD}$) or truncation back to

residue 216 in the putative intrinsically disordered region (BARD1^{216}) had minimal impact on

H2A-Ub activity and nucleosome binding **(Figure 1B, C)**. However, truncation to residue 193

(BARD1^{193}) caused a substantial decrease in H2A-Ub activity and a ~10-fold reduction in

binding affinity, suggesting a nucleosome-binding interface for residues between 194 and 216.

Consistent with this, deletion of the first half of the BARD1 IDR (BARD1$^{\Delta 140-270}$) greatly

diminished H2A-Ub activity **(Supplemental Figure 1E)**. Although deletion of residues 140-270

decreased nucleosome binding affinity by ~3-fold relative to full-length BARD1, the affinity is still

considerably stronger than the minimal RING/RING heterodimer or BARD1^{193} truncations

(Supplemental Figure 1F). A more targeted deletion (BARD1$^{\Delta 194-216}$) only modestly decreased

H2A-Ub activity, suggesting the presence of additional or compensatory nucleosome interaction

sites in BARD1 *in vitro*. Intrinsic nucleosome-binding properties of BARD1 IDRs were analyzed

using constructs containing only the N-terminal half (residues 124-270) or only the C-terminal

half (residues 269-424). Each IDR-only construct exhibited nucleosome binding by EMSA, with

modest differences in affinity **(Supplemental Figure 1G).** In summary, only the RING domain of

BRCA1 is required for high H2A-Ub activity in vitro while a small region within the BARD1 IDR

(BARD1 141-216, referred to in the discussion as IDR[Prox]) that engages in nucleosome binding

is required in addition to the BARD1 RING to support increased H2A-Ub activity.

Characterization of the BARD1 IDR reveals that the additional BARD1-nucleosome

interactions are mediated through DNA binding

BARD1 residues 141-216 identified as contributing to increased H2A-Ub activity and

nucleosome binding are in the large central segment predicted to be intrinsically disordered

(124-424) (McGuffin et al., 2000). Nuclear magnetic resonance (NMR) ^{1}H^{15}N-HSQC spectra of

^{15}N-BARD1 fragments containing residues 124-270, 269-424, and 124-424 each exhibited

narrow ^{1}H chemical shift dispersion consistent with intrinsic disorder **(Supplemental Figure 3A,**

B). Furthermore, individual spectra of BARD1 IDRs 124-270 and 269-424 overlay well with the

spectrum of the full region (124-424), indicating that the two regions behave independently of

one another **(Supplemental Figure 3B)**.

Despite being intrinsically disordered, the BARD1 124-270 sequence is largely conserved

across orthologs **(Supplemental Figure 3C)**. The region was previously reported to bind to

double-stranded DNA and specialized DNA structures containing base-pair mismatch

(discussed below) (Zhao et al., 2017). We therefore hypothesized that the contributions of the

BARD1 IDR (residues 141-216) to increased binding and activity are mediated through

interactions with nucleosomal DNA. Indeed, BARD1 124-270 binds strongly to nucleosomes

and to free 147-bp '601' dsDNA by EMSA **(Supplemental Figure 4A)**.

For residue-level information, we performed NMR binding experiments with ^{15}N-labelled BARD1

124-270 and either a 36-bp dsDNA fragment or mono-nucleosomes wrapped with 147-bp

dsDNA **(Figure 2A and Supplemental Figure 4B)**. A full titration was possible only with the

dsDNA fragment due to its lower molecular weight and higher solubility. The spectral series

displayed a subset of resonances that shift continuously as a function of DNA concentration

(Figure 2A). A similar set of resonances was perturbed (broadened in this case) upon addition

of 147-bp nucleosome, indicating that similar BARD1 residues engage nucleosomes and free

dsDNA **(Supplemental Figure 4B)**. To overcome spectral overlap in the spectrum of BARD1

124-270, the spectrum of a smaller construct containing the putative DNA binding region

(BARD1 141-216, where residue 141 is the first to extend beyond BARD1^{RING}) was assigned

and assignments were transferred to the longer segment, enabling analysis in both contexts

(Figure 2A, B and Supplemental Figure 4B, C). Secondary structure propensity (SSP)

predicted from the resonance assignments are consistent with the region being primarily

disordered, with some helical propensity predicted for residues 191-201 and a weaker prediction

of extended structure towards the beginning of the construct **(Supplemental Figure 5A).**

BARD1 residues 194-216 exhibit the largest backbone amide chemical shift perturbations

(CSPs) due to dsDNA addition, with smaller perturbations observed for residues 146-161

(Figure 2B). Residues between 162-191 were relatively unaffected by dsDNA, consistent with

the existence of distinct DNA-binding regions within the IDR. Secondary structure predicted

from chemical shifts in the presence of dsDNA (1 molar equivalent) indicate reduced helical

propensity in residues 191-199 upon DNA binding **(Supplemental Figure 5A)**. As well, ^{15}N

dynamics measurements reveal that residues 146-161 undergo changes in dynamics in the

presence of dsDNA **(Supplemental Figure 5B)**. Although low solubility limited similar analysis

on nucleosome-bound samples, the similar spectral perturbations observed in the presence of

low molar equivalents of nucleosomes or dsDNA suggest similar protein-DNA binding modes

(Supplemental Figure 4B). Thus, the data reveal two IDR regions adjacent to the RING

domain (146-161 and 194-216) that bind DNA when presented as free dsDNA or in the context

of a nucleosome. These regions are consistent with the H2A-Ub activity and nucleosome

binding results presented above.

Having established that isolated BARD1 IDR fragments bind to nucleosomes, we sought to

observe the interactions within the context of an active enzymatic complex. As such a complex

is too large for NMR analysis, we employed targeted crosslinking of BRCA1/BARD1 to

nucleosomal DNA. Photoactivatable crosslinker p-Benzoyl-L-phenylalanine (Bpa) was

incorporated at individual positions in the BARD1 IDR in the context of RING-containing

constructs using amber-codon suppression in *E. coli* (L120Bpa, W146Bpa, Y180Bpa, A195Bpa

and W218Bpa**; Supplemental Figure 6A)**. Due to the need to co-express a Bpa-specific

aminoacyl tRNA transferase, we designed a genetic fusion in which BRCA1 residues 1-104 are

fused to BARD1 residues 26-221 through a 12-residue GlySer-linker (BRCA1-f-BARD1[221]).

Wild-type BRCA1-f-BARD1[221] and Bpa-incorporated mutants retained similar H2A-Ub activity to

each other and to unfused heterodimers **(Supplemental Figure 6B)**. Nucleosomes with extra-

nucleosomal linker DNA that carried a fluorescent probe enabled visualization of covalent

BARD1-DNA crosslinked products via SDS-PAGE **(**NCP[185]; **Supplemental Figure 6C)**. Such

bands were detected for every Bpa variant, with enhanced crosslinking by variants with Bpa at

BARD1 W146 and A195 positions, consistent with the NMR dsDNA binding data presented

above **(Figure 2C)**. Crosslink bands were weaker from Y180 and W218 positions and

comparable to crosslinks from L120 which is located close to the RING domains and not

expected to contribute to productive DNA interactions. A similar pattern was observed in

crosslinking reactions containing free DNA **(Supplemental Figure 6D)**. As crosslinks can only

form from Bpa to sites that are within ~3Å, the data support a direct interaction between the

BARD1 IDR and nucleosomal DNA in the context of an enzymatically active complex.

Contribution of DNA binding to enhanced H2A-Ub activity

Results presented thus far establish that the DNA-binding region of the BARD1 IDR is

necessary for high H2A-Ub activity, but is it sufficient? To address this question, we asked

whether the DNA-binding properties of full-length BARD1 can support H2A-Ub activity in the

presence of BARD1 RING mutations that disrupt the histone-binding interface **(**BARD1

P89A/W91A; **Figure 3A)** (Hu et al., 2021; Witus, Burrell, et al., 2021). Heterodimers containing

BRCA1[RING] and full-length BARD1 (P89A/W91A) retain high-affinity nucleosome binding and

auto-ubiquitylation activity but do not transfer Ub to H2A or any other histone sites **(Figure 3B,**

Supplemental Figure 7A-C). Thus, the BARD1 RING-histone interface is absolutely required

for H2A-Ub activity while the BARD1 IDR-DNA interactions promote formation of a high-affinity

complex that supports enhanced H2A-Ub activity.

The BARD1 DNA-binding region(s) are disordered. To test whether any DNA-binding region

placed adjacent to the RING domain can serve the same purpose, we created a chimeric

BARD1 construct in which a previously identified DNA-binding region from the BRCA1 IDR

(498-663) was appended onto the BARD1 RING (Mark et al., 2005). A heterodimer composed

of BRCA1[RING] and the BARD1 chimera displayed increased H2A-Ub activity compared to the

minimal RING/RING complex, but not to the level of the native BARD1 sequence **(Figure 3C)**.

The nucleosome-binding affinities for these species are quite similar, suggesting that the

modest activity difference between the native BARD1 region and the installed BRCA1-derived

region arises from additional features **(Supplemental Figure 7D)**. The findings indicate that a

major contributor to increased H2A-Ub activity is the presence of a disordered, high-affinity

DNA-binding region near the enzymatic RING domains, but indicate that the BARD1 IDR is

especially effective.

Specialized DNA structures bind to the BARD1 IDR and inhibit H2A-Ub activity

Based on a previous report that the BARD1 IDR preferentially binds to partly unwound DNA

structures (bubble-DNA) and homologous recombination (HR) intermediates (D-loop DNA) over

double- or single-stranded DNA (Zhao et al., 2017), we hypothesized that interaction of the

265 BARD1 DNA-binding region with bubble- or D-loop DNA might affect its ability to interact with

266 nucleosomal DNA. Bubble- and D-loop- DNA substrates are likely similar based on the near-

267 identical affinities of the BARD1-DNA complexes (Zhao et al., 2017), so we used the simpler

268 bubble-DNA substrate in our assays. In nucleosome ubiquitylation reactions carried out with full-

269 length BRCA1/BARD1 in the presence of equal concentrations of DNA fragments containing

270 various patterns of base-pair mismatch, only bubble-DNA substantially inhibited H2A-Ub

271 activity. **(Figure 4A, B and Supplemental Figure 8A, B)**. The minimal RING heterodimer

272 activity was unaffected by bubble-DNA, consistent with the inhibition being due to competing

273 interactions of bubble- and nucleosomal DNA for binding to BARD1 141-216, **(Supplemental**

274 **Figure 8C, D)**. BARD1-nucleosome DNA crosslink formation was inhibited by either bubble-

275 DNA or dsDNA inhibited in a concentration-dependent manner, with bubble-DNA being

276 substantially more effective **(Figure 4C and Supplemental Figure 9).**

278 We compared the interactions of BARD1 141-216 with bubble- and dsDNA by NMR using DNA

279 species of similar sequence and size. CSPs and changes in dynamics were similar within

280 BARD1 194-216, the region identified as the major site for dsDNA and nucleosomal DNA

281 binding **(Figure 4D and Supplemental Figure 10A-C)**. Much stronger effects were produced

282 by bubble-DNA on backbone resonances of residues 143-157 with severe peak

283 broadening/disappearance, indicative of large changes in chemical environment and/or

284 dynamics. Only in the presence of bubble-DNA was a large CSP observed for the sidechain

285 resonance of W146 **(Supplemental Figure 10A, B).** Additional effects on sidechains were

286 observed in ^{1}H^{13}C-HSQC spectra of BARD1 141-216, where methyl group peaks from BARD1

287 residues 143-157 exhibit large CSPs that are not observed in the presence of dsDNA **(Figure**

288 **4E and Supplemental Figure 10D)**. Together, the data show that bubble DNA engages

289 strongly and specifically at a site near the beginning of the BARD1 IDR (residues 143-157) and

290 that both forms of DNA engage BARD1 194-216. Both IDR regions appear to be functionally

important, as only concomitant deletion of basic residues in both IDR regions (BARD1$^{\Delta150\text{-}155/\Delta194\text{-}}$

^{216}) alleviated H2A-Ub activity inhibition by bubble-DNA **(Figure 4F and Supplemental Figure**

8E).

DNA-binding IDR facilitates chromatin recruitment and DNA damage repair in cells

To test the importance of BARD1 IDR DNA binding in a cellular context, we established

doxycycline-inducible HeLa cell lines that deplete endogenous BARD1 via doxycycline-induced

expression of shRNA against BARD1 (HeLa-shBARD1) and stably re-express shBARD1-

resistant HA-tagged BARD1 (wild-type or Δ194-216; **Supplemental Figure 11A**). Endogenous

BARD1 is largely depleted (>90%) in these cells and both wild-type BARD1 and BARD1$^{\Delta194\text{-}216}$

localize to the nucleus and are expressed at similar levels **(Supplemental Figure 11A, B)**. In

cellular fractionation assays, less BARD1$^{\Delta194\text{-}216}$ is associated with chromatin than wild-type

BARD1 under both basal and olaparib-induced DNA-damage conditions **(Figure 5A)**.

Significantly reduced HA-BARD1$^{\Delta194\text{-}216}$ foci formation was observed in both untreated and

olaparib-induced DNA-damage conditions, indicating a defect in BRCA1/BARD1 recruitment to

chromatin and DNA damage sites **(Figure 5B and Supplemental Figure 11C)**. To assess the

ability of the BARD1 IDR mutant to facilitate DNA break repair, we performed clonogenic

survival assays upon treatment with DNA-damaging agents olaparib and cisplatin **(Figure 5C**

and Supplemental Figure 11D). Survival of BARD1$^{\Delta194\text{-}216}$ cells was significantly reduced

compared to wild-type controls under both drug treatments – a hallmark of defective HR **(Figure**

5C). Altogether, the results demonstrate that the short intrinsically disordered stretch of BARD1

with DNA-binding ability contributes to chromatin association in cells under both basal and

exogenous DNA-damage conditions and contributes to DNA break repair required for cell

survival.

Extra-nucleosomal linker DNA enhances H2A ubiquitylation

Chromatin is composed of adjacent nucleosome units separated by linker DNA. The extended

domain topology of both BRCA1 and BARD1 with their long IDRs raises the possibility of

interactions in higher-order chromatin substrates that are not present in minimal mono-

nucleosome substrates. Nucleosome substrates with extra-nucleosomal linker DNA (NCP[185])

and tri-nucleosomes (tri-NCP) were used as substrates in H2A-Ub assays to test for additional

chromatin involvement **(Figure 6A and Supplemental Figure 12A)**. In reactions using full-

length BRCA1/BARD1, faster H2A-Ub kinetics were observed for NCP[185] with linker DNA

compared to the minimal NCP[147] substrate **(Figure 6B)**. Notably, no additional increase in H2A-

Ub activity was observed for a tri-NCP substrate despite containing a longer linker DNA and

multiple histone binding interfaces. The data are consistent with two conclusions: 1) the

presence of linker DNA enhances BRCA1/BARD1 H2A-Ub activity and 2) functional interactions

are limited to one mono-nucleosome unit, at least in the absence of histone PTMs.

To explicitly test the BARD1 IDR DNA-binding region as the source of the enhanced activity

observed for NCP[185] and tri-NCP substrates, the H2A-Ub activity of the BRCA1[RING]/BARD1[216]

truncation was tested **(Figure 6C)**. This species exhibited considerably faster H2A-Ub kinetics

with either NCP[185] or tri-NCP compared to a minimal NCP[147] substrate, confirming that the

dependence on linker DNA arises from the BARD1 IDR. The presence of linker DNA provides

less than a two-fold increase in binding affinity for BRCA1[RING]/BARD1[216] to NCPs under salt

conditions used in H2A ubiquitylation assays **(Supplemental Figure 12B)**. Thus, in the absence

of linker DNA, BARD1 IDR can interact nearly as well with nucleosomal DNA, but the interaction

with linker DNA supports enhanced activity.

Effects of H2A-Ub marks on BRCA1/BARD1 H2A-Ub activity

Effects of mono-Ub at H2A positions known to be ubiquitylated by different E3 ligases were

assessed using homogenously mono-ubiquitylated nucleosomes containing ubiquitin

preinstalled at H2A positions K15, K119, or K127 via a non-hydrolyzable dichloroacetone

crosslinker **(Supplemental Figure 13A, B)** (Morgan et al., 2019). Ubiquitin at either H2A C-

terminal position (K119 or K127) decreased BRCA1/BARD1-dependent H2A-Ub activity, hinting

at exclusivity between BRCA1/BARD1 and RING1/PCGF E3 complexes that both modify the

H2A C-terminal tail. Consistent with previous observations, Ub preinstalled at H2A K15

increased the H2A-Ub activity of full-length BRCA1/BARD1, and this enhancement requires

BARD1 CTDs **(Figure 6D** and **Supplemental Figure 13C, D)**. This contrasts with unmodified

nucleosome substrates where BARD1 CTDs are not required for enhanced H2A-Ub activity (Hu

et al., 2021).

H2A K15-Ub-mediated enhancement of BRCA1/BARD1 H2A-Ub activity

Binding of BARD1 CTDs to nucleosomes that contain both H2A K15-Ub and H4K20me0 (Dai et

al., 2021; Hu et al., 2021) recruits BRCA1/BARD1 to damaged chromatin to facilitate DNA DSB

repair via HR (Becker et al., 2021; Krais et al., 2021; Nakamura et al., 2019). The BARD1 CTDs

use a binding surface that overlaps with the RING domain site, precluding binding of both

domains to the same nucleosome face. However, the observation that nucleosomes containing

H2A K15-Ub and H4K20me0 are better substrates in the context of BARD1^{FL} implies that RING

domains and CTDs can bind simultaneously to opposite faces of a nucleosome (Hu et al.,

2021). We used BARD1 truncation constructs to determine what features of BARD1 contribute

to enhanced activity on H2A K15-Ub nucleosome substrates. As expected, deletion of the CTDs

or histone-binding RING mutations of BARD1 do not support activity **(Figure 6D).** Deletion of

the BARD1 DNA-binding IDR (BRCA1^{RING}/BARD1$^{\Delta140-270}$) led to decreased H2A-Ub kinetics

compared with the BRCA1^{RING}/BARD1^{FL} complex **(Figure 6D)**. Thus, even with the additional

nucleosome-binding contribution provided by the CTDs on H2A K15-Ub nucleosomes, the DNA-

binding IDR still plays a role in enhancing ubiquitylation kinetics. Altogether, our findings support

a model in which three points of contact between BARD1 and H2A K15-Ub nucleosomes

(RING-histone, IDR-DNA, and CTDs-histone/Ub) are required for full BRCA1/BARD1-

dependent H2A-Ub activity **(Figure 7E)**.

Chemical crosslinking experiments using amine-reactive crosslinkers DSS and BS3 and

analyzed by mass spectrometry were carried out to compare complexes formed on unmodified

versus H2A K15-Ub-modified nucleosomes **(Supplemental Figure 14)**. To focus on the BARD1

subunit, BRCA1^{RING}/BARD1^{FL} was used. There were clear differences in cross-linked products

involving BARD1 CTDs, with numerous intermolecular crosslinks (judged by unique crosslinks

and peptide spectral mapping counts) to nucleosomes containing H2A K15-Ub but only sparse

crosslinks from reactions that included unmodified nucleosomes **(Figure 6E and Supplemental**

Figure 14C). The crosslinks detected are consistent with high-resolution structures of BARD1

CTDs bound to nucleosomes containing H2A K15-Ub and the lack of these in the unmodified

nucleosome reaction is consistent with the interaction being critically dependent on the

presence of Ub at H2A K15 **(Supplemental Figure 14D)**. There was also increased

crosslinking between BARD1 RING and IDR domains to histones in samples containing H2A

K15-Ub nucleosomes **(Figure 6E, gray crosslinks and Supplemental Figure 14C, E)**,

consistent with the reported higher binding affinity of that complex. Thus, binding of the BARD1

CTDs in response to the presence of H2A K15-Ub facilitates BARD1 RING and IDR interactions

that lead to the observed increase in H2A K127-Ub activity.

The BARD1 RING domain and CTDs are separated by ~300 intrinsically disordered residues,

allowing for as much as 1000 Å of separation between the two structured domains. It is

therefore possible that the CTDs bind to one nucleosome unit, while the RING domains are

recruited to a nearby nucleosome unit on chromatin to facilitate H2A-Ub activity. To test this

possibility explicitly, asymmetric di-nucleosome substrates (di-NCPs) where one nucleosome

unit contained H2A K15-Ub (H2A[silent]) and an adjacent nucleosome unit contained unmodified

H2A with a fluorophore conjugated near its N-terminus to facilitate specific detection in a

nucleosome ubiquitylation assay were assembled **(**H2A[observe]; **Figure 6F and Supplemental**

Figure 13E-G). The asymmetric di-NCP substrate containing H2A K15-Ub as H2A[silent] was

rapidly ubiquitylated by BRCA1[RING]/BARD1[FL] and this level of activity requires the BARD1 CTDs

(H2A[observe]; **Figure 6G and Supplemental Figure 13H)**. Unmodified di-NCPs were ubiquitylated

with similar kinetics regardless of the presence or absence of the CTDs. The results confirm that

BARD1 CTDs bound to an H2A K15-Ub-containing nucleosome enable RING binding to a

neighboring nucleosome and that this configuration leads to high ubiquitylation activity.

Furthermore, deletion of the BARD1 DNA-binding IDR yields a substantial loss of H2A-Ub

activity, indicating that all three binding functionalities are required for full activity. Altogether, the

results support formation of a higher-order chromatin complex where BARD1 spans adjacent

nucleosome units in which at least one contains H2A K15-Ub. The complex is mediated through

binding of the BRCA1/BARD1 RING domains and BARD1 CTDs to neighboring nucleosome

units and of the BARD1 IDR to intervening linker DNA **(Figure 7F)**.

Discussion

Structural and biochemical studies have revealed that the two RING domains of heterodimeric

BRCA1/BARD1 bind to the histone face of a nucleosome and that this interaction is essential for

the ability to modify histone H2A at its C-terminal tail (Hu et al., 2021; Witus, Burrell, et al.,

2021). Other studies revealed that BARD1 CTD also binds to the histone face, but this

interaction requires Ub-modified H2A, a modification that occurs in response to DNA damage

(Becker et al., 2021; Hu et al., 2021; Krais et al., 2021). Herein we have identified an additional

region of BARD1 that engages with chromatin, namely the BARD1 IDR proximal to the RING

(IDR[Prox]). In concert with the RING domains, BARD1 IDR[Prox] is sufficient to support highly

enhanced H2A-Ub activity in vitro. BARD1 IDR^{Prox} increases the binding affinity of the

BRCA1/BARD1 RINGs from a micromolar to nanomolar binding regime, likely critical in cells

where BRCA1/BARD1 is in low abundance. Indeed, our cellular data reveal an important role for

this region in chromatin recruitment under both basal and DNA-damage conditions and in DNA

damage repair. Remarkably, this critical function is carried out by a short stretch of intrinsically

disordered BARD1 residues that are highly conserved.

We propose distinct complexes of BRCA1/BARD1 and chromatin substrates that depend on the

status of nucleosomal histone PTMs (e.g., unmodified or H2A K15-Ub) **(Figure 7)**. Importantly,

while BARD1 CTDs are critical for interactions with nucleosomes that contain H2A K15-Ub, they

are dispensable for high-affinity interactions with unmodified nucleosomes **(Figure 7C, D)**.

Additional interactions that enhance H2A-Ub activity are mediated through BARD1 IDR^{Prox}

which engages nucleosomal and extra-nucleosomal linker DNA. Our findings indicate that DNA

binding plays an auxiliary role in the ability of BRCA1/BARD1 to ubiquitylate H2A by boosting

the affinity of the Ub ligase/nucleosome complex. Supporting this, a BARD1 RING-domain

mutant (BARD1 P89A/W91A) that disrupts histone binding is unable to support H2A-Ub activity

with any chromatin substrate tested despite retaining strong nucleosome binding affinity and

intrinsic Ub ligase activity. This indicates that high-affinity binding to nucleosomal DNA by

BARD1 IDR^{Prox} is not sufficient for H2A-Ub activity, while the positioning of the RING domains

on the histone surface to orient the RING-E2~Ub complex for site-specific ubiquitin transfer to

histone H2A is absolutely required.

BARD1 IDR^{Prox} contains two discrete regions with DNA-binding properties, separated by ~30

residues. Each DNA-binding region has a high Lys/Arg content while the intervening region

contains more negatively charged residues (**Supplemental Figure 3C**). The more C-terminal

residues of BARD1 IDR^{Prox} (190-216) appear to be the dominant site for dsDNA engagement

while the portion of BARD1 IDR^{Prox} (145-165) directly adjacent to the RING domain also

engages dsDNA but shows a pronounced preference for specialized DNA structures such as

bubble-DNA that have features found in intermediates in HR (Zhao et al., 2017) and possibly in

R-loop structures formed during transcription (Santos-Pereira & Aguilera, 2015). The two

regions display different secondary structure propensities, and we note a conserved

hydrophobic trio (145-MWF) within the RING-adjacent site that is highly affected by bubble-

DNA. Structural information regarding IDR-DNA interactions is sparse, but our findings hint at

distinct modes of binding by IDRs rather than merely electrostatic, non-specific associations. To

that point, BARD1 IDR^{Prox} appears to possess small regions with specialized binding modes that

are linked to BRCA1/BARD1 function.

Preferential binding to bubble-DNA serves to inhibit H2A-Ub activity in our assays where the

specialized DNA species was present in *trans*. This suggests the intriguing possibility that

proximity or high local concentrations of such species might modulate the modification of H2A

by BRCA1/BARD1 in cells. Furthermore, binding of BARD1 IDR^{Prox} to nucleosomal DNA or to

specialized nucleic acid structures could serve to control its spatial and temporal involvement in

HR and other processes where such specialized DNA structures may be present (e.g., stalled

replication fork protection, R-loop resolution) (Hatchi et al., 2015; Schlacher et al., 2012).

Additionally, BRCA1/BARD1 is present in higher-order protein assemblies that are known to

bind to chromatin via distinct nucleosome interfaces that may be formed by other proteins in the

complex (Belotserkovskaya et al., 2020; Bochar et al., 2000; Huen et al., 2010; Savage &

Harkin, 2015). Understanding the combined binding contributions of BRCA1/BARD1 and other

proteins in such complexes to nucleosome association and H2A-Ub activity is an important

future goal.

Data presented here provide direct experimental evidence in support of an extended

BRCA1/BARD1/chromatin complex that spans nucleosomal units. Such a complex is enabled

by the presence of H2A K15-Ub, an early mark of DNA damage, which engages the BARD1

CTDs while the enzymatic RING domains occupy an adjacent/nearby nucleosome unit **(Figure**

7E, F). The 300-residue BARD1 IDR allows extensive separation between the enzymatic RING

domains and BARD1 CTDs while itself engaging in high-affinity interactions with

nucleosomal/linker DNA. Such a configuration may allow the RING domains to reach

nucleosome units even further away than an adjacent unit while anchored to chromatin via the

BARD1 CTDs. This situation could lead to BRCA1/BARD1-dependent H2A-Ub of multiple

nucleosomes in the vicinity of a H2A K15-Ub anchor nucleosome. Additionally, the ability to

reach across nucleosome units in chromatin provides a platform to scaffold higher-order

complexes across long molecular distances.

As H2A K15-Ub is specifically a mark of DNA damage, this model will not be in play for other

BRCA1/BARD1-dependent processes on chromatin such as its transcriptional repression and

activation activities. Results presented here show that the DNA-binding BARD1 IDR^{Prox} is

required for enhanced H2A-Ub activity on unmodified nucleosomes. An important question

going forward is what other histone PTMs influence the chromatin binding and H2A-Ub activity

of BRCA1/BARD1. We previously reported that methylation at H3 K79, a mark associated with

actively transcribed chromatin, inhibits BRCA1/BARD1-mediated H2A-Ub activity (Witus,

Burrell, et al., 2021). This may be attributable to the location of H3 K79 near the BARD1-histone

interface. A comprehensive understanding of histone PTMs that influence BRCA1/BARD1

nucleosome binding, higher-order chromatin complex formation, and H2A-Ub activity will

provide valuable insight into its functions and regulation in different biological processes.

Mostly missing from our discussion is the huge BRCA1 subunit. Its only established direct point-

of-contact to chromatin is via its N-terminal RING domain, although at least one section of its

enormous IDR has DNA binding activity in vitro (Mark et al., 2005). BRCA1/BARD1 serves as a

large protein interaction hub that forms numerous distinct complexes, many of which involve

BRCA1 as the binding partner. In our in vitro investigation, the entirety of BRCA1 apart from its

enzymatic RING domain, was dispensable for H2A-Ub activity. That said, we note that full-

length BRCA1/BARD1 displayed lower H2A-Ub activity than complexes containing truncated

BRCA1 (Δ11q or RING) and offer two hypotheses for the observation. First, BRCA1^{FL} but

neither truncation contains a DNA-binding region that supports lower H2A-Ub activity than

BARD1 IDR^{Prox}. The presence of strong DNA-binding regions in both BRCA1 and BARD1 in the

full-length complex could set up a competition between these regions for binding to nucleosomal

DNA, leading to lower apparent H2A-Ub activity. Alternatively, or in addition, possible inhibitory

interactions between BRCA1 and BARD1 subunits might have been alleviated upon BRCA1

truncation. Although such effects could be artifactual, we suggest that the status of the BRCA1

subunit in terms of its binding partners could provide an additional layer of modulation of

BRCA1/BARD1 H2A-Ub activity. Occupancy of regions of BRCA1 and BARD1 by unique

binding partners may occlude certain nucleosome-binding regions, resulting in inhibition or

enhancement of chromatin binding and H2A-Ub activity. Further investigation into chromatin

complexes containing full-length BRCA1/BARD1 and the many protein assemblies and binding

partners that are required to execute its cellular functions is warranted.

In conclusion, our results establish an extensive multivalent network of interactions that facilitate

BRCA1/BARD1 chromatin association and H2A-Ub activity. These interactions are likely critical

for many functions of BRCA1/BARD1 in the nucleus that are not limited to its H2A-Ub activity

and may be disrupted by mutations that cause cancer.

Acknowledgements

We thank K. Luger (University of Colorado Boulder) for sharing plasmids for the 147-bp and

NLE-trimer '601' DNA, S. Tan (Penn State University) for sharing the plasmid for the 185-bp

'601' DNA, and G. Debelouchina (UC San Diego) for the MMTV DNA plasmid. We thank D.

Veelser for providing access to equipment, and M. Morgan for insights into dichloroacetone

crosslinking of H2A-Ub. S.R.W., L.M.T., K.E.K., P.S.B., and R.E.K. were supported by the NIH

(R01CA260834)**.** R.E.K. is the Edmond H. Fischer/Washington Research Foundation Endowed

Chair in Biochemistry**.** This work and A.Z. was supported in part by the University of

Washington's Proteomics Resource (UWPR95794). A.Z. and T.N.D. were also supported by the

NIH (P41GM103533). W.Z. was supported by a Young Investigator Award from Max and Minnie

Tomerlin Voelcker Fund, the Cancer Prevention and Research Institute of Texas (RP210102),

and the NIH (R01GM141091). D.B.W. was supported by the NIH (R00-HD090201).

Author Contributions

S.R.W., W.Z., P.S.B., and R.E.K. conceived of the study. S.R.W., W.L., L.M.T., M.W., W.Z., and

K.E.K. performed experiments and analyzed data. D.B.W. analyzed data. A.Z. performed

chemical crosslinking and mass spectrometry and associated data processing under the

supervision of T.N.D. The manuscript was written by S.R.W., P.S.B., and R.E.K., with input from

all co-authors.

Competing Interests Statement

The authors of this article declare no competing interests.

References

Alenezi, W. M., Fierheller, C. T., Recio, N., & Tonin, P. N. (2020). Literature Review of BARD1 as a
Cancer Predisposing Gene with a Focus on Breast and Ovarian Cancers. *Genes*, *11*(8).
https://doi.org/10.3390/genes11080856

Becker, J. R., Clifford, G., Bonnet, C., Groth, A., Wilson, M. D., & Chapman, J. R. (2021). BARD1 reads
H2A lysine 15 ubiquitination to direct homologous recombination. *Nature*, *596*(7872), 433–437.
https://doi.org/10.1038/s41586-021-03776-w

Belotserkovskaya, R., Raga Gil, E., Lawrence, N., Butler, R., Clifford, G., Wilson, M. D., & Jackson, S.
P. (2020). PALB2 chromatin recruitment restores homologous recombination in BRCA1-
deficient cells depleted of 53BP1. *Nature Communications*, *11*(1), 819.
https://doi.org/10.1038/s41467-020-14563-y

Bochar, D. A., Wang, L., Beniya, H., Kinev, A., Xue, Y., Lane, W. S., Wang, W., Kashanchi, F., &
Shiekhattar, R. (2000). BRCA1 Is Associated with a Human SWI/SNF-Related Complex:
Linking Chromatin Remodeling to Breast Cancer. *Cell*, *102*(2), 257–265.
https://doi.org/10.1016/S0092-8674(00)00030-1

Brzovic, P. S., Keeffe, J. R., Nishikawa, H., Miyamoto, K., Fox, D., 3rd, Fukuda, M., Ohta, T., & Klevit,
R. (2003). Binding and recognition in the assembly of an active BRCA1/BARD1 ubiquitin-ligase
complex. *Proceedings of the National Academy of Sciences of the United States of America*,
100(10), 5646–5651. PubMed. https://doi.org/10.1073/pnas.0836054100

Brzovic, P. S., Rajagopal, P., Hoyt, D. W., King, M. C., & Klevit, R. E. (2001). Structure of a BRCA1-
BARD1 heterodimeric RING-RING complex. *Nature Structural Biology*, *8*(10), 833–837.
PubMed. https://doi.org/10.1038/nsb1001-833

Chambers, M. C., Maclean, B., Burke, R., Amodei, D., Ruderman, D. L., Neumann, S., Gatto, L., Fischer,
B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen, B., Baker, T.
A., Brusniak, M.-Y., Paulse, C., Creasy, D., … Mallick, P. (2012). A cross-platform toolkit for
mass spectrometry and proteomics. *Nature Biotechnology*, *30*(10), 918–920.
https://doi.org/10.1038/nbt.2377

Chin Jason W., Martin Andrew B., King David S., Wang Lei, & Schultz Peter G. (2002). Addition of a
photocrosslinking amino acid to the genetic code of Escherichia coli. *Proceedings of the National
Academy of Sciences*, *99*(17), 11020–11024. https://doi.org/10.1073/pnas.172226299

Christensen, D. E., Brzovic, P. S., & Klevit, R. E. (2007). E2-BRCA1 RING interactions dictate synthesis
of mono- or specific polyubiquitin chain linkages. *Nature Structural & Molecular Biology*,
14(10), 941–948. https://doi.org/10.1038/nsmb1295

Dai, L., Dai, Y., Han, J., Huang, Y., Wang, L., Huang, J., & Zhou, Z. (2021). Structural insight into
BRCA1-BARD1 complex recruitment to damaged chromatin. *Molecular Cell*, *81*(13), 2765-
2777.e6. https://doi.org/10.1016/j.molcel.2021.05.010

Dao, H. T., Dul, B. E., Dann, G. P., Liszczak, G. P., & Muir, T. W. (2020). A basic motif anchoring ISWI
to nucleosome acidic patch regulates nucleosome spacing. *Nature Chemical Biology*, *16*(2), 134–
142. https://doi.org/10.1038/s41589-019-0413-4

Densham, R. M., Garvin, A. J., Stone, H. R., Strachan, J., Baldock, R. A., Daza-Martin, M., Fletcher, A.,
Blair-Reid, S., Beesley, J., Johal, B., Pearl, L. H., Neely, R., Keep, N. H., Watts, F. Z., & Morris,
J. R. (2016). Human BRCA1-BARD1 ubiquitin ligase activity counteracts chromatin barriers to
DNA resection. *Nature Structural & Molecular Biology*, *23*(7), 647–655. PubMed.
https://doi.org/10.1038/nsmb.3236

Densham, R. M., & Morris, J. R. (2019). Moving Mountains—The BRCA1 Promotion of DNA
589   Resection. *Frontiers in Molecular Biosciences*, *6*, 79. https://doi.org/10.3389/fmolb.2019.00079

590   Dyer, P. N., Edayathumangalam, R. S., White, C. L., Bao, Y., Chakravarthy, S., Muthurajan, U. M., &
591   Luger, K. (2003). Reconstitution of Nucleosome Core Particles from Recombinant Histones and
592   DNA. In *Methods in Enzymology* (Vol. 375, pp. 23–44). Academic Press.
593   https://doi.org/10.1016/S0076-6879(03)75002-2

594   Eng, J. K., Jahan, T. A., & Hoopmann, M. R. (2013). Comet: An open-source MS/MS sequence database
595   search tool. *PROTEOMICS*, *13*(1), 22–24. https://doi.org/10.1002/pmic.201200439

596   Gudmundsdottir, K., & Ashworth, A. (2006). The roles of BRCA1 and BRCA2 and associated proteins in
597   the maintenance of genomic stability. *Oncogene*, *25*(43), 5864–5874.
598   https://doi.org/10.1038/sj.onc.1209874

599   Hatchi, E., Skourti-Stathaki, K., Ventz, S., Pinello, L., Yen, A., Kamieniarz-Gdula, K., Dimitrov, S.,
600   Pathania, S., McKinney, K. M., Eaton, M. L., Kellis, M., Hill, S. J., Parmigiani, G., Proudfoot, N.
601   J., & Livingston, D. M. (2015). BRCA1 recruitment to transcriptional pause sites is required for
602   R-loop-driven DNA damage repair. *Molecular Cell*, *57*(4), 636–647. PubMed.
603   https://doi.org/10.1016/j.molcel.2015.01.011

604   Hoopmann, M. R., Zelter, A., Johnson, R. S., Riffle, M., MacCoss, M. J., Davis, T. N., & Moritz, R. L.
605   (2015). Kojak: Efficient Analysis of Chemically Cross-Linked Protein Complexes. *Journal of
606   Proteome Research*, *14*(5), 2190–2198. https://doi.org/10.1021/pr501321h

607   Hu, Q., Botuyan, M. V., Zhao, D., Cui, G., Mer, E., & Mer, G. (2021). Mechanisms of BRCA1–BARD1
608   nucleosome recognition and ubiquitylation. *Nature*, *596*(7872), 438–443.
609   https://doi.org/10.1038/s41586-021-03716-8

610   Huen, M. S. Y., Sy, S. M. H., & Chen, J. (2010). BRCA1 and its toolbox for the maintenance of genome
611   integrity. *Nature Reviews Molecular Cell Biology*, *11*(2), 138–148.
612   https://doi.org/10.1038/nrm2831

613   Kalb, R., Mallery, D. L., Larkin, C., Huang, J. T. J., & Hiom, K. (2014). BRCA1 is a histone-H2A-
614   specific ubiquitin ligase. *Cell Reports*, *8*(4), 999–1005. PubMed.
615   https://doi.org/10.1016/j.celrep.2014.07.025

616   Käll, L., Canterbury, J. D., Weston, J., Noble, W. S., & MacCoss, M. J. (2007). Semi-supervised learning
617   for peptide identification from shotgun proteomics datasets. *Nature Methods*, *4*(11), 923–925.
618   https://doi.org/10.1038/nmeth1113

619   Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:
620   Improvements in Performance and Usability. *Molecular Biology and Evolution*, *30*(4), 772–780.
621   https://doi.org/10.1093/molbev/mst010

622   Kjaergaard, M., & Poulsen, F. M. (2011). Sequence correction of random coil chemical shifts: Correlation
623   between neighbor correction factors and changes in the Ramachandran distribution. *Journal of
624   Biomolecular NMR*, *50*(2), 157–165. https://doi.org/10.1007/s10858-011-9508-2

625   Kolas Nadine K., Chapman J. Ross, Nakada Shinichiro, Ylanko Jarkko, Chahwan Richard, Sweeney
626   Frédéric D., Panier Stephanie, Mendez Megan, Wildenhain Jan, Thomson Timothy M., Pelletier

627   Laurence, Jackson Stephen P., & Durocher Daniel. (2007). Orchestration of the DNA-Damage
628   Response by the RNF8 Ubiquitin Ligase. *Science*, *318*(5856), 1637–1640.
629   https://doi.org/10.1126/science.1150034

630   Krais, J. J., & Johnson, N. (2020). BRCA1 Mutations in Cancer: Coordinating Deficiencies in
631   Homologous Recombination with Tumorigenesis. *Cancer Research*, *80*(21), 4601–4609.
632   https://doi.org/10.1158/0008-5472.CAN-20-1830

633   Krais, J. J., Wang, Y., Patel, P., Basu, J., Bernhardy, A. J., & Johnson, N. (2021). RNF168-mediated
634   localization of BARD1 recruits the BRCA1-PALB2 complex to DNA damage. *Nature*
635   *Communications*, *12*(1), 5016. https://doi.org/10.1038/s41467-021-25346-4

636   Lazar, G. A., Desjarlais, J. R., & Handel, T. M. (1997). De novo design of the hydrophobic core of
637   ubiquitin. *Protein Science : A Publication of the Protein Society*, *6*(6), 1167–1178. PubMed.
638   https://doi.org/10.1002/pro.5560060605

639   Lee, Y.-T., Gibbons, G., Lee, S. Y., Nikolovska-Coleska, Z., & Dou, Y. (2015). One-pot refolding of core
640   histones from bacterial inclusion bodies allows rapid reconstitution of histone octamer. *Protein*
641   *Expression and Purification*, *110*, 89–94. PubMed. https://doi.org/10.1016/j.pep.2015.02.007

642   Lorick, K. L., Jensen, J. P., Fang, S., Ong, A. M., Hatakeyama, S., & Weissman, A. M. (1999). RING
643   fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proceedings of the*
644   *National Academy of Sciences*, *96*(20), 11364. https://doi.org/10.1073/pnas.96.20.11364

645   Lowary, P. T., & Widom, J. (1998). New DNA sequence rules for high affinity binding to histone
646   octamer and sequence-directed nucleosome positioning11Edited by T. Richmond. *Journal of*
647   *Molecular Biology*, *276*(1), 19–42. https://doi.org/10.1006/jmbi.1997.1494

648   Luger, K., Rechsteiner, T. J., & Richmond, T. J. (1999). Preparation of nucleosome core particle from
649   recombinant histones. *Methods in Enzymology*, *304*, 3–19. https://doi.org/10.1016/s0076-
650   6879(99)04003-3

651   Mark, W.-Y., Liao, J. C. C., Lu, Y., Ayed, A., Laister, R., Szymczyna, B., Chakrabartty, A., &
652   Arrowsmith, C. H. (2005). Characterization of segments from the central region of BRCA1: An
653   intrinsically disordered scaffold for multiple protein-protein and protein-DNA interactions?
654   *Journal of Molecular Biology*, *345*(2), 275–287. PubMed.
655   https://doi.org/10.1016/j.jmb.2004.10.045

656   Masuda, T., Xu, X., Dimitriadis, E. K., Lahusen, T., & Deng, C.-X. (2016). "DNA Binding Region" of
657   BRCA1 Affects Genetic Stability through modulating the Intra-S-Phase Checkpoint.
658   *International Journal of Biological Sciences*, *12*(2), 133–143. PubMed.
659   https://doi.org/10.7150/ijbs.14242

660   Mattiroli, F., Vissers, J. H. A., van Dijk, W. J., Ikpa, P., Citterio, E., Vermeulen, W., Marteijn, J. A., &
661   Sixma, T. K. (2012). RNF168 ubiquitinates K13-15 on H2A/H2AX to drive DNA damage
662   signaling. *Cell*, *150*(6), 1182–1195. https://doi.org/10.1016/j.cell.2012.08.005

663   McGinty, R. K., Henrici, R. C., & Tan, S. (2014). Crystal structure of the PRC1 ubiquitylation module
664   bound to the nucleosome. *Nature*, *514*(7524), 591–596. PubMed.
665   https://doi.org/10.1038/nature13890

McGuffin, L. J., Bryson, K., & Jones, D. T. (2000). The PSIPRED protein structure prediction server.
Bioinformatics, *16*(4), 404–405. https://doi.org/10.1093/bioinformatics/16.4.404

Morgan, M., Jbara, M., Brik, A., & Wolberger, C. (2019). Chapter One—Semisynthesis of ubiquitinated
histone H2B with a native or nonhydrolyzable linkage. In M. Hochstrasser (Ed.), *Methods in*
Enzymology (Vol. 618, pp. 1–27). Academic Press. https://doi.org/10.1016/bs.mie.2019.01.003

Morgan Michael T., Haj-Yahya Mahmood, Ringel Alison E., Bandi Prasanthi, Brik Ashraf, & Wolberger
Cynthia. (2016). Structural basis for histone H2B deubiquitination by the SAGA DUB module.
Science, *351*(6274), 725–728. https://doi.org/10.1126/science.aac5681

Mullan, P. B., Quinn, J. E., & Harkin, D. P. (2006). The role of BRCA1 in transcriptional regulation and
cell cycle control. *Oncogene*, *25*(43), 5854–5863. https://doi.org/10.1038/sj.onc.1209872

Nakamura, K., Saredi, G., Becker, J. R., Foster, B. M., Nguyen, N. V., Beyer, T. E., Cesa, L. C., Faull, P.
A., Lukauskas, S., Frimurer, T., Chapman, J. R., Bartke, T., & Groth, A. (2019). H4K20me0
recognition by BRCA1–BARD1 directs homologous recombination to sister chromatids. *Nature*
Cell Biology, *21*(3), 311–318. https://doi.org/10.1038/s41556-019-0282-9

Poepsel, S., Kasinath, V., & Nogales, E. (2018). Cryo-EM structures of PRC2 simultaneously engaged
with two functionally distinct nucleosomes. *Nature Structural & Molecular Biology*, *25*(2), 154–
162. https://doi.org/10.1038/s41594-018-0023-y

Riffle, M., Jaschob, D., Zelter, A., & Davis, T. N. (2016). ProXL (Protein Cross-Linking Database): A
Platform for Analysis, Visualization, and Sharing of Protein Cross-Linking Mass Spectrometry
Data. *Journal of Proteome Research*, *15*(8), 2863–2870.
https://doi.org/10.1021/acs.jproteome.6b00274

Riffle, M., Jaschob, D., Zelter, A., & Davis, T. N. (2019). Proxl (Protein Cross-Linking Database): A
Public Server, QC Tools, and Other Major Updates. *Journal of Proteome Research*, *18*(2), 759–
764. https://doi.org/10.1021/acs.jproteome.8b00726

Santos-Pereira, J. M., & Aguilera, A. (2015). R loops: New modulators of genome dynamics and
function. *Nature Reviews Genetics*, *16*(10), 583–597. https://doi.org/10.1038/nrg3961

Savage, K. I., & Harkin, D. P. (2015). BRCA1, a "complex" protein involved in the maintenance of
genomic stability. *The FEBS Journal*, *282*(4), 630–646. https://doi.org/10.1111/febs.13150

Schlacher, K., Wu, H., & Jasin, M. (2012). A Distinct Replication Fork Protection Pathway Connects
Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. *Cancer Cell*, *22*(1), 106–116.
https://doi.org/10.1016/j.ccr.2012.05.015

Scully, R., Chen, J., Ochs, R. L., Keegan, K., Hoekstra, M., Feunteun, J., & Livingston, D. M. (1997).
Dynamic Changes of BRCA1 Subnuclear Location and Phosphorylation State Are Initiated by
DNA Damage. *Cell*, *90*(3), 425–435. https://doi.org/10.1016/S0092-8674(00)80503-6

Sherker, A., Chaudhary, N., Adam, S., Heijink, A. M., Noordermeer, S. M., Fradet-Turcotte, A., &
Durocher, D. (2021). Two redundant ubiquitin-dependent pathways of BRCA1 localization to
DNA damage sites. *EMBO Reports*, *n/a*(n/a), e53679. https://doi.org/10.15252/embr.202153679

Simons, A. M., Horwitz, A. A., Starita, L. M., Griffin, K., Williams, R. S., Glover, J. N. M., & Parvin, J.
D. (2006). BRCA1 DNA-Binding Activity Is Stimulated by BARD1. *Cancer Research*, *66*(4),
2012–2018. https://doi.org/10.1158/0008-5472.CAN-05-3296

Sobhian Bijan, Shao Genze, Lilli Dana R., Culhane Aedín C., Moreau Lisa A., Xia Bing, Livingston
David M., & Greenberg Roger A. (2007). RAP80 Targets BRCA1 to Specific Ubiquitin
Structures at DNA Damage Sites. *Science*, *316*(5828), 1198–1202.
https://doi.org/10.1126/science.1139516

Stewart, M. D., Zelin, E., Dhall, A., Walsh, T., Upadhyay, E., Corn, J. E., Chatterjee, C., King, M.-C., &
Klevit, R. E. (2018). BARD1 is necessary for ubiquitylation of nucleosomal histone H2A and for
transcriptional regulation of estrogen metabolism genes. *Proceedings of the National Academy of*
Sciences of the United States of America, *115*(6), 1316–1321. PubMed.
https://doi.org/10.1073/pnas.1715467115

Tan, W., Murphy, V. J., Charron, A., van Twest, S., Sharp, M., Constantinou, A., Parker, M. W.,
Crismani, W., Bythell-Douglas, R., & Deans, A. J. (2020). Preparation and purification of mono-
ubiquitinated proteins using Avi-tagged ubiquitin. *PLOS ONE*, *15*(2), e0229000.
https://doi.org/10.1371/journal.pone.0229000

Tarsounas, M., & Sung, P. (2020). The antitumorigenic roles of BRCA1–BARD1 in DNA repair and
replication. *Nature Reviews Molecular Cell Biology*, *21*(5), 284–299.
https://doi.org/10.1038/s41580-020-0218-z

Thapa, I., Vahrenkamp, R., Witus, S. R., Lightle, C., Falkenberg, O., Sellin Jeffries, M. K., Klevit, R. E.,
& Stewart, M. D. (2022). Conservation of transcriptional regulation by BRCA1 and BARD1 in
Caenorhabditis elegans. *Nucleic Acids Research*, gkac877. https://doi.org/10.1093/nar/gkac877

Uckelmann, M., Densham, R. M., Baas, R., Winterwerp, H. H. K., Fish, A., Sixma, T. K., & Morris, J. R.
(2018). USP48 restrains resection by site-specific cleavage of the BRCA1 ubiquitin mark from
H2A. *Nature Communications*, *9*(1), 229. https://doi.org/10.1038/s41467-017-02653-3

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E.,
Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J.,
Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … SciPy 1.0 Contributors. (2020).
SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, *17*(3),
261–272. https://doi.org/10.1038/s41592-019-0686-2

Vizcaíno, J. A., Côté, R., Reisinger, F., M. Foster, J., Mueller, M., Rameseder, J., Hermjakob, H., &
Martens, L. (2009). A guide to the Proteomics Identifications Database proteomics data
repository. *PROTEOMICS*, *9*(18), 4276–4283. https://doi.org/10.1002/pmic.200900402

Witus, S. R., Burrell, A. L., Farrell, D. P., Kang, J., Wang, M., Hansen, J. M., Pravat, A., Tuttle, L. M.,
Stewart, M. D., Brzovic, P. S., Chatterjee, C., Zhao, W., DiMaio, F., Kollman, J. M., & Klevit, R.
E. (2021). BRCA1/BARD1 site-specific ubiquitylation of nucleosomal H2A is directed by
BARD1. *Nature Structural & Molecular Biology*, *28*(3), 268–277.
https://doi.org/10.1038/s41594-020-00556-4

Witus, S. R., Stewart, M. D., & Klevit, R. E. (2021). The BRCA1/BARD1 ubiquitin ligase and its
substrates. *Biochemical Journal*, 3467–3483. https://doi.org/10.1042/BCJ20200864

Witus, S. R., Zhao, W., Brzovic, P. S., & Klevit, R. E. (2022). BRCA1/BARD1 is a nucleosome reader
and writer. *Trends in Biochemical Sciences*. https://doi.org/10.1016/j.tibs.2022.03.001

Zelter, A., Bonomi, M., Kim, J. ook, Umbreit, N. T., Hoopmann, M. R., Johnson, R., Riffle, M., Jaschob,
D., MacCoss, M. J., Moritz, R. L., & Davis, T. N. (2015). The molecular architecture of the
Dam1 kinetochore complex is defined by cross-linking based structural modelling. *Nature*
Communications, *6*(1), 8673. https://doi.org/10.1038/ncomms9673

Zhao, W., Steinfeld, J. B., Liang, F., Chen, X., Maranon, D. G., Jian Ma, C., Kwon, Y., Rao, T., Wang,
W., Sheng, C., Song, X., Deng, Y., Jimenez-Sainz, J., Lu, L., Jensen, R. B., Xiong, Y., Kupfer, G.
M., Wiese, C., Greene, E. C., & Sung, P. (2017). BRCA1-BARD1 promotes RAD51-mediated
homologous DNA pairing. *Nature*, *550*(7676), 360–365. PubMed.
https://doi.org/10.1038/nature24060

Zhu, G., Xia, Y., Nicholson, L. K., & Sze, K. H. (2000). Protein Dynamics Measurements by TROSY-
Based NMR Experiments. *Journal of Magnetic Resonance*, *143*(2), 423–426.
https://doi.org/10.1006/jmre.2000.2022

Zhu, Q., Hoong, N., Aslanian, A., Hara, T., Benner, C., Heinz, S., Miga, K. H., Ke, E., Verma, S.,
Soroczynski, J., Yates, J. R., Hunter, T., & Verma, I. M. (2018). Heterochromatin-Encoded
Satellite RNAs Induce Breast Cancer. *Molecular Cell*, *70*(5), 842-853.e7.
https://doi.org/10.1016/j.molcel.2018.04.023

Zhu, Q., Pao, G. M., Huynh, A. M., Suh, H., Tonnu, N., Nederlof, P. M., Gage, F. H., & Verma, I. M.
(2011). BRCA1 tumour suppression occurs via heterochromatin-mediated silencing. *Nature*,
477(7363), 179–184. https://doi.org/10.1038/nature10371

Main Figures

Figure 1. Contributions of BARD1 regions to nucleosome binding and H2A-Ub activity.

A. Domain organization of BRCA1 and BARD1. The domain names are indicated above the
cartoons (RING, really interesting new gene; IDR, intrinsically disordered region; CC, coiled-coil;
BRCT, BRCA1 C-terminal; Ank, Ankyrin repeat domain; CTD, C-terminal domain). Constructs
used to generate data in panels b and c are shown, and residue bounds are reported in
Extended Data Figure I.

B. Quantification of time-course nucleosome H2A-Ub assays using the indicated
BRCA1/BARD1 truncations. Data points represent mean values and error bars are ± 1-s.d. of
n=3 independent experiments.

C. Comparison of nucleosome binding affinity from fluorescence-quenching based
measurements using the indicated BRCA1/BARD1 constructs. Data show mean values and
error bars are ± 1-s.d. of n=4 independent experiments. The reported difference in affinities to
BRCA1^{RING}/BARD1^{RING} is likely underestimated, as the minimal RING/RING heterodimer binding
data was collected at a lower ionic strength than the rest of the constructs to obtain well-
behaved binding data (50 mM vs. 100 mM NaCl).

Figure 2. The BARD1 IDR binds to nucleosomes via DNA.

A. $^1H^{15}N$-HSQC NMR titration spectra of ^{15}N-BARD1 124-270 with a 36-bp dsDNA fragment. Darker shades of blue correspond to increasing amounts of DNA added. A subset of highly affected signals is labelled with their residue identities, with arrows showing their trajectories over the course of the titration.

B. $^1H^{15}N$-HSQC NMR amide chemical shift perturbations (CSPs) observed to ^{15}N-BARD1 141-216 signals when bound to a 36-bp dsDNA fragment (1:1 molar equivalent complex).

C. Representative SDS-PAGE gel monitoring in-gel fluorescence (labelled DNA) of UV-induced Bpa crosslinking of BRCA1-f-BARD1[221] to nucleosomes (left). Quantification of Bpa crosslinking experiments with nucleosomes (right). The intensity of each crosslinked band was normalized to the intensity of the L120Bpa crosslinked band for each replicate experiment. Graph bars show the mean; error bars are $\pm$ 1-s.d. and the open circles are the values of individual replicates for of n=3 independent experiments.

828
829 **Figure 3. Contribution of BARD1 DNA-binding and RING-histone binding to H2A-Ub**
830 **activity.**
831
832 **A.** The BARD1 RING-histone interface showing the locations of BARD1 P89 and W91 relative
833 to the histone surface (PDB: 7JZV).
834
835 **B, C.** Quantification of time-course H2A-Ub assays using the indicated BRCA1/BARD1
836 constructs. Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments. Two
837 single time-points were determined to be outliers in panel C and excluded from analysis. Values
838 for these excluded points are included as source data.

Figure 4. Inhibition of nucleosome binding and H2A-Ub by specialized DNA structures.

A. Design of 36-mer DNA competitor fragments used in H2A-Ub activity, Bpa crosslinking, and NMR assays. Red indicates non-base-paired regions. The number of unpaired bases or base-pairs is: ss – 36; ds – 0; bubble – 8; frayed – 8; ssOverhang – 24.

B. Single time-point H2A-Ub inhibition assays using BRCA1^{FL}/BARD1^{FL} (100 nM) and the indicated competitor DNA (2.5 μM). Data show the mean; error bars are ± 1-s.d.; and open circles are individual data points of n=4 independent experiments. One outlier point for "frayed-DNA" was excluded from the analysis.

C. Inhibition of UV-induced Bpa crosslinking between the indicated BRCA1-f-BARD1^{221} Bpa-incorporated constructs and NCP^{185} substrates in the presence of increasing amounts of dsDNA or bubble-DNA competitor. Data show the mean; error bars are ± 1-s.d. of n=3 (W146Bpa) or n=4 (A195Bpa) independent experiments.

856   **D.** ^{1}H^{15}N-HSQC NMR CSPs observed to ^{15}N-BARD1 141-216 signals when bound to a 36-mer
857   dsDNA (blue, front) or bubble-DNA (gray, back) fragment (1:1 molar equivalent complex).
858   Signals for residues 143-157 are broadened beyond detection in the bubble-DNA-bound
859   spectrum, and gray bars are equal to the CSP value for the W146Ne CSP that was observed
860   (see Supplemental Figure 10A, B for spectra).
861
862   **E.** Selected region of ^{13}C-HSQC spectra of ^{13}C-BARD1 141-216 in 1:1 molar equivalent complex
863   with dsDNA fragment or bubble-DNA fragment. Signal trajectories in bound spectra are
864   indicated by arrows.
865
866   **F.** Single time-point H2A-Ub inhibition assays using heterodimers containing
867   BRCA1^{RING}/BARD1^{FL} (left, 12 min endpoint) or the indicated BARD1 internal deletion mutant
868   (middle and right, 20 min endpoint) and increasing amounts of dsDNA or bubble-DNA
869   competitor. The same amount of E3 (50 nM) was used for each BRCA1/BARD1 truncation.
870   Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments.
871
872   One time-point from Panels B and C each were determined to be outliers and excluded from
873   analysis. Values for these points are included as source data.
874

875
876
Figure 5. Cellular role for BARD1 DNA-binding in chromatin recruitment and DNA
damage repair.

A. Western blot analysis to detect HA-BARD1 in nuclei isolated from HeLa-shBARD1 cells
stably expressing wild type or Δ194-216 mutant of HA-BARD1, where endogenous BARD1 was
depleted by doxycycline-induced shBARD1 expression. The nuclei were salt-fractionated into
NS100, NS300, NS420, and NP420 to assess the amount of HA-BARD1 associated with
chromatin.

B. Quantification of cells with >5 HA-BARD1 foci (left panel) and number of HA-BARD1 foci per
nucleus (right panel) with and without Olaparib treatment. *p<0.05, **p<0.01, ***p<0.001, by
Student's t-test.

C. Clonogenic survival of HeLa-shBARD1 cells stably expressing wild type or Δ194-216 mutant
of HA-BARD1 upon their treatment with indicated amount of olaparib and cisplatin. Error bars
represent SEM (n=3).
890
891

892
893
894 **Figure 6. H2A-Ub activity using chromatin substrates with linker DNA and pre-installed**
895 **H2A K15-Ub.**
896
897 **A.** Schematic of chromatin substrates used for H2A-Ub activity and nucleosome binding assays
898 in panels b and c. The length of linker-DNA is indicated next to the NCP[185] and tri-NCP.
899

B, C. Quantification of time-course H2A-Ub assays using the indicated BRCA1/BARD1 constructs and chromatin substrates.

D. Quantification of time-course H2A-Ub assays using homogenously modified H2A K15-Ub mono-nucleosome substrates and indicated BRCA1/BARD1 constructs.

E. Intermolecular crosslinks observed by chemical crosslinking and MS analysis between BARD1 and histones using wild-type (left) and H2A K15-Ub (right) 147-bp '601' nucleosomes and BRCA1^{RING}/BARD1^{FL} heterodimers. Crosslinks to histones emanating from the CTDs of BARD1 are shown in blue, and the RING-IDR region in gray. A lysine-depleted region of the BARD1 IDR is labelled and indicated by a black bar.

F. Design of asymmetric di-NCP substrates.

G, H. Quantification of time-course H2A-Ub assays using the indicated combinations of di-NCP substrate and BRCA1/BARD1 heterodimer. For panels A, D, and E, data show the mean; error bars are ± 1-s.d. of n=3 independent experiments. One time-point from Panel A was determined to be an outlier and excluded from analysis. The excluded value is reported as source data.

925
926
927
928 **Figure 7. Models of BRCA1/BARD1/chromatin complexes supported by this study.**
929
930 **A, B.** Model of the BRCA1/BARD1 minimal RING heterodimer binding to an unmodified mono-
931 nucleosome or higher-order chromatin substrate.
932
933 **C, D.** Model of BRCA1/BARD1 (BARD1 full-length or RING-IDR) binding with an unmodified
934 mono-nucleosome or higher-order chromatin substrate.
935
936 **E, F.** Model of BRCA1/BARD1^{FL} binding to ubiquitylated chromatin substrates in a 'wrapped'
937 complex (panel E) or in a complex that spans nucleosome units (panel F). The relative H2A-Ub
938 activity and affinity of the complexes is indicated below each set of panels.
939

Supplemental Figures

Supplemental Figure 1. Characterization of BRCA1/BARD1 constructs.

A. Representative time-course nucleosome ubiquitylation assays monitoring H2A-Ub efficiency of different BRCA1/BARD1 constructs quantified in Panel E and Figure 1B. These data are representative of the basic assay set-up for other time-course H2A-Ub assays in this study.

B. Electrophoretic mobility shift assay (EMSA) comparing the nucleosome binding of full-length BRCA1/BARD1 (top) to the minimal RING heterodimer (bottom). Data shown are representative of n=2 independent experiments.

C-E. Quantification of H2A-Ub assays using the indicated BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of n=3 independent experiments.

960 **F.** Comparison of nucleosome binding affinity from fluorescence-quenching based
961 measurements using the indicated BRCA1/BARD1 constructs. Data show the mean; error bars
962 are ± 1-s.d. of n=4 independent experiments.
963
964 **G.** EMSA monitoring nucleosome binding by the indicated BARD1 IDR fragments. Data shown
965 are representative of n=2 independent experiments.
966

1. BRCA1^{FL}/BARD1^{FL}
2. BRCA1$^{\Delta 11q}$/BARD1^{FL}
3. BRCA1^{RING}/BARD1^{FL}
4. BRCA1^{RING}/BARD1$^{\Delta CTDs}$
5. BRCA1^{FL}/BARD1^{RING}
6. BRCA1$^{\Delta 11q}$/BARD1^{RING}
7. BRCA1^{RING}/BARD1$^{FL-P89A/W91A}$
8. BRCA1^{RING}/BARD1$^{\Delta 194-216}$
9. BRCA1^{RING}/BARD1$^{\Delta 150-155/ \Delta 194-216}$
10. BRCA1^{RING}/BARD1$^{\Delta 140-270}$
11. BRCA1^{RING}/BARD1^{216}
12. BRCA1^{RING}/BARD1^{193}
13. BRCA1^{RING}/BARD1^{RING}

968
969 **Supplemental Figure 2. Coomassie-stained SDS-PAGE gels of BRCA1/BARD1 constructs**
970 **used in this study.**

971
972
973 **Supplemental Figure 3. Characterization of the BARD1 intrinsically disordered region.**
974
975 **A.** Coomassie-stained SDS-PAGE gel of BARD1 fragments used in NMR experiments.
976
977 **B.** ^{1}H^{15}N-HSQC NMR spectra of the indicated ^{15}N-labelled BARD1 fragments.
978
979 **C.** Multiple-sequence alignment of BARD1 124-270 with a representative subset of homologs.
980 Residue identities that are conserved to the human sequence are indicated by the presence of a
981 period in place of a residue letter. Basic residues (Lys, Arg, His) are color coded in blue, and
982 basic positions that are fully conserved across all species are highlighted with a gray
983 background.
984

985
986
987 **Supplemental Figure 4. NMR analysis of BARD1 IDR-DNA and IDR-nucleosome**
988 **complexes.**
989
990 **A.** EMSA comparing DNA and nucleosome binding by BARD1 124-270. Data shown are
991 representative of n=2 independent experiments.
992
993 **B.** ^{1}H^{15}N-HSQC NMR spectra of ^{15}N-BARD1 124-270, titrating a 36-bp dsDNA fragment (left),
994 and a single titration point with nucleosomes (right). The red arrows point to signals that are
995 similarly affected between the dsDNA- and nucleosome-bound samples.
996
997 **C.** ^{1}H^{15}N-HSQC NMR spectra of ^{15}N-BARD1 141-216 with a 36-mer dsDNA fragment. Data from
998 panel C are also shown in Supplemental Figure 10A.
999

1000
1001 **Supplemental Figure 5. Changes in dynamics and secondary structure of the BARD1 IDR**
1002 **bound to DNA.**
1003
1004 **A.** Changes in secondary structural propensity (SSP) of ^{15}N-BARD1 141-216 when bound to a
1005 36-bp dsDNA fragment. The black bars for apo BARD1 are behind the gray bars where DNA
1006 was added.
1007
1008 **B.** T1/T2 dynamics measurements for amide signals of ^{15}N-BARD1 141-216 in apo form and
1009 bound to dsDNA (1:1 molar equivalent complex). Data from panel B are also shown in
1010 Supplemental Figure 10C.
1011

1012
1013 **Supplemental Figure 6. UV-induced Bpa crosslinking of BRCA1/BARD1 to nucleosomal**
1014 **DNA.**
1015
1016 **A.** Coomassie-stained SDS-PAGE gel of unfused BRCA1^{RING}/BARD1^{216} heterodimers and
1017 BRCA1-f-BARD1^{221} constructs with and without Bpa incorporation.
1018
1019 **B.** H2A-Ub activity of unfused heterodimers compared to wild-type and Bpa-incorporated
1020 BRCA1-f-BARD1^{221}. Data show the mean; error bars are ± 1-s.d. of n=3 independent
1021 experiments.
1022
1023 **C.** Native gel of fluorescently labelled DNA and nucleosomes used in Bpa crosslinking
1024 experiments.
1025
1026 **D.** Representative data of in-gel fluorescence signal from of a UV-induced crosslinking reaction
1027 to free 185-bp '601' DNA (DNA^{185}, top) and quantification of crosslinking experiments (bottom).
1028 The intensity.y of each crosslinked band was normalized to the L120Bpa crosslinked band
1029 intensity for each replicate experiment. Data bars show the mean; error bars are ± 1-s.d. and
1030 the open circles are the values of individual replicates for of n=3 independent experiments.
1031
1032

1033
1034 **Supplemental Figure 7. Nucleosome binding and Ub ligase activity of BARD1 RING**
1035 **mutants and DNA-binding chimeric constructs.**
1036
1037 **A.** Comparison of nucleosome binding affinity using the indicated BRCA1/BARD1 constructs.
1038 Data show the mean; error bars are $\pm$ 1-s.d. of n=3 independent experiments.
1039
1040 **B.** Auto-ubiquitylation assay using BRCA1^{RING}/BARD1^{FL} (wild-type or P89A/W91A). The assay
1041 monitors depletion of Ub and appearance of high molecular weight auto-Ub products in the
1042 presence of BRCA1/BARD1. Data shown is representative of n=2 independent experiments.
1043
1044 **C.** Nucleosome ubiquitylation assay using the indicated BRCA1/BARD1 constructs and blotting
1045 for VSVG-tagged H2A (top) and H3 (middle and bottom). The asterisk denotes a proteolysis
1046 product of H3 that appears over the time-course.
1047
1048 **D.** Comparison of nucleosome binding affinity using the indicated BRCA1/BARD1 constructs.
1049 Data show the mean; error bars are $\pm$ 1-s.d. of n=2 independent experiments.
1050

1051
1052
1053 **Supplemental Figure 8. Assembly of competitor DNA fragments and inhibition of H2A-Ub**
1054 **activity.**
1055
1056 **A.** DNA-stained native-gel of annealed DNA competitor fragments. Amounts loaded were
1057 normalized to show similar band-intensity for each fragment.
1058
1059 **B.** Representative gel data of H2A-Ub inhibition assay using full-length BRCA1/BARD1 (100
1060 nM) and the indicated DNA competitor fragment (2.5 mM)
1061
1062 **C.** Representative gel data of single time-point (20 min) H2A-Ub inhibition assays using
1063 BRCA1[RING]/BARD1[216] (top, 50 nM E3) and BRCA1[RING]/BARD1[140] (bottom, 750 nM E3) and
1064 indicated amounts of dsDNA and bubble-DNA competitors.
1065
1066 **D.** Quantification of H2A-Ub DNA-competitor inhibition assays from panel b using the indicated
1067 BRCA1/BARD1 constructs. Data show the mean; error bars are ± 1-s.d. of n=3 independent
1068 experiments.

1069
1070 **E.** Representative gel data of single time-point H2A-Ub inhibition assays using
1071 BRCA1^{RING}/BARD1^{FL} (top, 50 nM E3, 12 min endpoint), BRCA1^{RING}/BARD1$^{\Delta194\text{-}216}$ (middle, 50
1072 nM E3, 20 min endpoint), and BRCA1^{RING}/BARD1$^{\Delta150\text{-}155/\Delta194\text{-}216}$ (bottom, 50 nM E3, 20 min
1073 endpoint) and indicated amounts of dsDNA and bubble-DNA competitors.
1074
1075

1076
1077
1078 **Supplemental Figure 9. Inhibition of UV-induced Bpa crosslinking of BRCA1/BARD1 to**
1079 **nucleosomal DNA.**
1080
1081 Representative data of in-gel fluorescence signal from a UV-induced Bpa crosslinking reaction
1082 between W146Bpa (top) and A195Bpa (bottom) BRCA1-f-BARD1^{221} to 185-bp '601'
1083 nucleosomes (NCP^{185}) in the presence of dsDNA and bubble-DNA competitors. Quantified data
1084 from n=3 replicate experiments are shown in Figure 4C.

1085

Supplemental Figure 10. Comparison of dsDNA and bubble-DNA binding to the BARD1 IDR by NMR.

A, B. ^{1}H^{15}N-HSQC NMR spectra of ^{15}N-BARD1 141-216 with a 36-mer dsDNA fragment (A) or a 36-mer bubble-DNA fragment with 8-bp of mismatch (B). Both complexes were assembled at 1:1 molar equivalent. The W146Nε signals are inlaid, and the red arrows point to a signal that is present exclusively in the DNA-bound spectra. Data from panel A are also shown in Supplemental Figure 4C.

C. T1/T2 dynamics measurements for amide signals of ^{15}N-BARD1 141-216 in apo form and bound to dsDNA or bubble-DNA (1:1 molar equivalent complex). A subset of data from panel C (apo and +dsDNA) are also shown in Supplemental Figure 5B.

D. ^{1}H^{13}C-HSQC NMR spectra of ^{13}C-BARD1 141-216 with a 36-mer dsDNA fragment (blue), or a 36-mer bubble-DNA fragment with 8-bp of mismatch (gray). Both complexes were assembled at 1:1 molar equivalent.

Supplemental Figure 11. Cellular studies of BARD1 DNA-binding mutant.

A. Western blot analysis to detect BRCA1, BARD1 and HA-BARD1 of the whole-cell lysate from HeLa-shBARD1 cells stably expressing wild type or Δ194-216 mutant of HA-BARD1, where endogenous BARD1 was depleted by doxycycline induced shBARD1 expression. The tubulin blot was included as a loading control.

B. Western blot analysis to detect BRCA1 and HA-BARD1 in the whole-cell lysate (WCL),
cytoplasmic fraction (Cyto) and nuclear extraction (NC) from HeLa-shBARD1 cells stably
expressing wild type or Δ194-216 mutant of HA-BARD1, where endogenous BARD1 was
depleted by doxycycline treatment. The further fractionations of the nuclei isolated are shown in
Figure 5A. Tubulin and Lamin B1 were included as loading controls and as indication of the
Cyto and NC fractions, respectively.
1117
1118 **C.** Representative micrographs of HA-BARD1 foci (red) in the nucleus of HeLa-shBARD1 cells
1119 stably expressing wild type or Δ194-216 mutant of HA-BARD1 with or without the treatment of
1120 10 mM olaparib 24h (left panels). Blue: DAPI. Cells with >5 HA-BARD1 foci and number of HA-
1121 BARD1 foci per nuclear were quantified (reported in Figure 5B).
1122
1123 **D.** Representative images for cell survival data quantified in Figure 5C.
1124
1125

1126
1127 **Supplemental Figure 12. Analysis of chromatin substrates.**
1128
1129 **A.** DNA-stained (left) and fluorescence (right) native gel of chromatin substrates used for H2A-
1130 Ub activity assays in Figure 5b and c. EcoRI digestion of tri-NCPs is used to confirm octamer
1131 binding saturation.
1132
B. Comparison of nucleosome binding affinity between BRCA1^{RING}/BARD1^{216} and NCP^{147} and
NCP^{185} substrates at 100 mM (top) and 150 mM (bottom) buffer-salt concentration (NaCl). The
solid vertical lines denote the position of K_d values for the indicated buffer-salt concentration,
and the dotted vertical lines denote the position of K_d values from the other buffer-salt
concentration. Data show the mean; error bars are ± 1-s.d. of n=3 independent experiments
using both NCP^{147} and NCP^{185}, with an additional replicate (total of n=4) for NCP^{147}.

1139
1140
1141 **Supplemental Figure 13. Design, assembly, and quality control of H2A K15-Ub chromatin**
1142 **substrates.**
1143
1144 **A.** Coomassie-stained SDS-PAGE gel of wild-type histones and dichloroacetone linked H2A-
1145 Ub/H2B dimers.
1146
1147 **B.** DNA-stained native gel of mono-nucleosomes with and without H2A-Ub incorporation at the
1148 indicated positions.
1149
1150 **C.** Example data of H2A-Ub activity assay using full-length BRCA1/BARD1 and preinstalled
1151 H2A-Ub NCP substrates. Western blots were probed for H2A.
1152

1153   **D.** Quantification of time-course H2A-Ub assays using unmodified or pre-installed H2A-Ub
1154   chromatin substrates and BRCA1[FL]/BARD1[FL].
1155
1156   **E.** Schematic of asymmetric di-NCP assembly strategy.
1157
1158   **F.** DNA-stained native gel of di-NCP ligation reaction.
1159
1160   **G.** DNA-stained (left) and fluorescence (right) native gels of di-NCP substrates after purification
1161   by $MgCl_2$ precipitation.
1162
1163   **H.** Example gel data of H2A-Ub activity assay using di-NCP substrates and indicated
1164   BRCA1/BARD1 constructs.
1165
1166

1167
1168 **Supplemental Figure 14. Analysis of BRCA1/BARD1 chromatin complexes by chemical**
1169 **crosslinking and mass-spectrometry.**
1170
1171 **A.** Coomassie-stained SDS-PAGE gel of chemical crosslinking reactions used in mass
1172 spectrometry analysis
1173
1174 **B.** Crosslinks observed from the RING domain of BRCA1 to ordered regions histones mapped
1175 on a high-resolution structure of the complex (PDB: 7JZV). Cα-Cα distances are indicated in the
1176 figure.
1177
1178 **C.** Quantitative comparison of crosslinking from reactions with unmodified and H2A K15-Ub
1179 NCPs. Different subsets of crosslinks were quantified for the number of unique observed
1180 crosslinks and total peptide spectral mapping counts for all crosslinks within a given subset.

1181
1182 **D.** Crosslinks observed from the CTDs of BARD1 to ordered regions histones in the H2A K15-
1183 Ub NCP sample mapped on a high-resolution structure (PDB: 7E8I). $C\alpha$-$C\alpha$ distances are
1184 indicated. The P117 $C\alpha$ serves as a proxy for the true crosslinking sites of K118 and K119 that
1185 are not modelled, denoted by an asterisk.
1186
1187 **E.** Intermolecular crosslinks observed by chemical crosslinking and MS analysis between
1188 BARD1 and histones using wild-type (left) and H2A K15-Ub (right, separated by dashed line)
1189 147-bp '601' nucleosomes and BRCA1[RING]/BARD1[FL] heterodimers. Crosslinks to histones
1190 emanating from the BARD1 RING-IDR region are highlighted in blue. A lysine-depleted region
1191 of the BARD1 IDR is labelled and indicated by a black bar.
1192
1193

1194

1195

1229 **Tables**

1230

1231 **Extended Data Table I. Details about BRCA1/BARD1 constructs used in this study.**

1232

Construct	BRCA1 bounds*	BRCA1 tags	BARD1 bounds*	BARD1 tags	Expression system
BRCA1^{FL}/ BARD1^{FL}	1-1863	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1$^{\Delta 11q}$/ BARD1^{FL}	1-264, 1366-1863	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1^{FL}	1-112	N-FLAG	1-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta CTDs}$	1-112	N-FLAG	1-424	N-TwinStrepII	*Sf9*
BRCA1^{FL}/ BARD1^{RING}	1-1863	N-FLAG	1-140	N-TwinStrepII	*Sf9*
BRCA1$^{\Delta 11q}$/ BARD1^{RING}	1-264, 1366-1863	N-FLAG	1-140	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 194-216}$	1-112	N-FLAG	1-193, 217-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 150-155/\Delta 194-216}$	1-112	N-FLAG	1-149, 156-193, 217-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1$^{\Delta 140-270}$	1-112	N-FLAG	1-139, 271-777	N-TwinStrepII	*Sf9*
BRCA1^{RING}/ BARD1^{216}	1-110	N-6xHis	26-216	none	*E. coli*
BRCA1^{RING}/ BARD1^{193}	1-110	N-6xHis	26-193	none	*E. coli*
BRCA1^{RING}/ BARD1^{RING}	1-110	N-6xHis	26-140	none	*E. coli*

1233

1234 * BRCA1 UniProt sequence ID: P38398-1; BARD1 UniProt sequence ID: Q99728-1

1235

1236

1237

1238

Methods

Protein, nucleosome, and experimental reagent production

Ubiquitylation machinery: Human E1 (UBA1), E2 (UBE2D3), and ubiquitin (wild-type and Q2C

mutant) were expressed and purified as previously described(Brzovic et al., 2003; Christensen

et al., 2007; Lazar et al., 1997). For a full list of BRCA1/BARD1 truncation constructs, their

residue bounds, purification/epitope tags, and expression system, see Extended Data Table I.

1245

BRCA1/BARD1 from E. coli: For purification of truncated BRCA1/BARD1 heterodimers from *E.*

coli expression (BRCA1^{RING}/BARD1^{RING}, BRCA1^{RING}/BARD1^{193}, and BRCA1^{RING}/BARD1^{216}),

plasmids containing 6xHis-tagged BRCA1 (pCOT7n) and untagged BARD1 (pET28n) were co-

transformed into *E. coli* BL21 (DE3) cells, grown in LB media at 37 °C to OD$_{600nm}$ of 0.6-0.8,

supplemented with 100 µM ZnCl$_2$ and induced with 0.2 mM IPTG for ~16 hours at 16 °C. The

cell pellets were resuspended in 30 mL of Ni^{2+} start buffer (25 mM Tris-HCl pH 7.5, 500 mM

NaCl, 10 mM imidazole) supplemented with DNase, RNase, EDTA-free protease inhibitors

(Roche) and 1 mM PMSF, lysed by French press, centrifuged, and the supernatant was applied

to a 5 mL HisTrap FF crude column (Cytiva). After extensive washing with Ni^{2+} start buffer

containing 10 mM imidazole and 30 mM imidazole, protein was eluted using the same buffer

supplemented with 500 mM imidazole. The BRCA1^{RING}/BARD1^{RING} and BRCA1^{RING}/BARD1^{193}

constructs were concentrated and further purified using a 120 mL Superdex 75 column on an

AKTA FPLC system in SEC buffer (25 mM HEPES-NaOH pH 7.5, 150 mM NaCl, 1 mM DTT;

GE Healthcare). For the BRCA1^{RING}/BARD1^{216}, HisTrap elutions were dialyzed into ion-

exchange buffer (25 mM Tris-HCl pH 7.5, 250 mM NaCl, 1 mM DTT) and applied to a 5 mL

HiTrap SP HP column (Cytiva) and eluted over a 12 column-volume (CV) gradient (0.25-1 M

NaCl). Peak fractions were concentrated, aliquoted, and flash frozen.

1263

For Bpa crosslinking experiments, 6xHis-BRCA1$^{1\text{-}104}$-(GS)$_6$-BARD1$^{26\text{-}221}$ was sub-cloned using

Gibson assembly into a pET28n vector. Constructs for Bpa incorporation were generated via

site-directed mutagenesis to install amber stop codons in place of BARD1 L120, W146, Y180,

A195, and W218. Bpa-mutant Pet28n-6xHis-BRCA1$^{1\text{-}104}$-(GS)$_6$-BARD1$^{26\text{-}221}$ plasmids were co-

transformed with a pEVOL-pBpF plasmid encoding an Bpa aminoacyl tRNA synthetase (gift

from P. Schultz, Addgene #31190) in *E. coli* BL21 (DE3) cells(Chin Jason W. et al., 2002). At

OD$_{600nm}$=0.4-0.5, cells were shifted from 37 °C to 16 °C and supplemented with 1 mM Bpa

(dissolved in 1M NaOH; Bachem) and 100 μM ZnCl$_2$. After 30 minutes, protein expression was

induced by addition of 0.02% L-arabinose and 0.2 mM IPTG for ~16 hours at 16 °C. Proteins

were purified as described for the BRCA1^{RING}/BARD1^{216} construct. Truncations that did not

incorporate Bpa were separated via the SP column.

1275

BRCA1/BARD1 from insect cells: For insect cell expression, truncation and deletion constructs

were derived from full-length BRCA1 (pFastBac-FLAG-BRCA1$^{1\text{-}1863}$; gift from W. Zhao, UT

Health Sciences Center at San Antonio) and insect cell codon-optimized full-length BARD1

(pFastBac-Twin-StrepTagII-BARD1$^{1\text{-}777}$; gift from A. Deans, Addgene plasmid #137166)(Tan et

al., 2020) using Gibson assembly cloning (NEB). Baculovirus was generated using the Bac-to-

Bac system in suspension culture *Sf9* cells according to the manufacturers protocols

(Invitrogen). For general *Sf9* growth and virus amplification, SF900-II media (ThermoFisher)

was supplemented with 5% FBS (HyClone) and 1x antibiotic-antimycotic (Gibco). For protein

expression, 15 mL of BRCA1- and 15 mL of BARD1-containing P3 baculoviruses were added to

650 mL of *Sf9* cells at ~1.5x10^6 cells/mL in SF900-II media supplemented with 1% FBS and 1x

antibiotic-antimycotic. Protein expression was carried out for 48-72 hours at 27 °C, shaking at

110 RPM. Following expression, cells were spun down at 150 x g for 10 minutes. The pellet was

resuspended in 30 mL of Strep-start buffer (50 mM Tris-HCl pH 8, 10% glycerol, 150 mM NaCl,

1 mM DTT) supplemented with EDTA-free protease inhibitors and 1 mM PMSF and flash-

frozen. Following freeze-thawing in a 37 °C water bath, additional EDTA-free protease inhibitors

and PMSF were added in addition to 2 μL of benzonase (Sigma Aldrich). All subsequent steps

were performed on ice or in a cold room over a short timeframe to prevent proteolysis (<3 hours

total). Lysis was performed by sonication using a Branson Sonifier 250 (VWR Scientific) by

applying 2 rounds of 15 pulses at 70% duty on power level 8, ensuring that the cell mixture

stayed cold. PMSF was added throughout the lysis protocol. The lysed cells were centrifuged at

41,000 x g for 15 minutes, filtered through a 0.45 mM low protein-binding PVDF filter (Millipore)

and applied to a 1 mL StrepTrap HP column pre-equilibrated in Strep-start buffer (Cytiva). The

column was washed with 8 mL Strep-start buffer supplemented with protease inhibitors, 8 mL

wash buffer (25 mM Tris-HCl pH 7.5, 300 mM NaCl, 0.01% Igepal-CA630, 1 mM DTT, 5 mM

MgCl2, 2 mM ATP, 10% glycerol, EDTA-free protease inhibitors), and an additional 5 mL of

Strep-start buffer. The column was eluted using 6 mL of Strep-start buffer supplemented with 3

mM d-Desthiobiotin (Sigma). The elution was diluted with 20 mL of ion-exchange buffer (25 mM

HEPES-NaOH pH 7.5, 10% glycerol, 1 mM DTT) and applied to a 1 mL SP HP column pre-

equilibrated in ion-exchange buffer containing 75 mM NaCl. The SP column was eluted over 12-

CV (75-500 mM NaCl). Peak fractions were combined and concentrated using a 30K MWCO

concentrator (Amicon), and flash-frozen in small aliquots. Concentrations were estimated by

measuring absorbance at 280 nM using molar extinction coefficients obtained from ProtParam

(Expasy). Concentration estimates were verified by SDS-PAGE and Coomassie staining, and

slight adjustments in concentration were applied to best equalize the amounts of E3 in activity

and binding assays.

1311

BARD1 intrinsically disordered region: BARD1 124-270, 141-216, 269-424, and 124-424 were

sub-cloned from pFastBac-6xHis-BARD1[1-777] (gift from W. Zhao, UTHCSCA) into a modified

pET28 vector containing an N-terminal 6xHis-SUMO tag using Gibson assembly cloning.

Plasmids were transformed into BL21 (DE3) *E. coli*, grown at 37 °C in M9 media supplemented

with appropriate combination of isotopes (^{15}N-NH$_4$Cl, ^{13}C-glucose; Cambridge Isotopes) to an

OD$_{600nm}$ of 0.6-0.8, and induced with 0.2 mM IPTG for ~16 hours at 16 °C. HisTrap Ni^{2+}

purification was performed as described above for BRCA1/BARD1 constructs expressed in *E.*

coli. Elutions were dialyzed into ion exchange buffer (25 mM Tris pH 8.0, 200 mM NaCl, 0.5 mM

EDTA, 1 mM DTT) in the presence of GST-SENP1 (produced in-house) to cleave the SUMO

tag, leaving a non-native serine residue preceding the N-terminus except in the case of 269-

424. The cleaved product was applied at a HiTrap Q column preceding a HiTrap SP column in-

tandem to capture the non-specifically bound DNA (Cytiva). After application, the Q column was

removed, and the SP column was eluted using a 12-CV gradient (0.2-1M NaCl). The peak

fractions were combined, concentrated, and further purified using a Superdex 75 column

equilibrated in NMR buffer (25 mM MOPS-NaOH pH 7.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM

TCEP (tris(2- carboxyethyl)phosphine)).

1328

Histone purification: Wild-type and mutant human histones were transformed in *E. coli* BL21

(DE3) *pLysS* cells, grown at 37 °C to an OD$_{600nm}$ of 0.5-0.6, and induced with 0.5 mM IPTG for 2

hours (H4) or 3 hours (H2A-2A, H2B-1K, H3.2) at 37 °C. Histone octamers used for

fluorescence labeling were purified using the one-pot refolding protocol as described previously

using 6xHis-TEV-H2A(Lee et al., 2015; Witus, Burrell, et al., 2021). For nucleosome

ubiquitylation assays, a glycine within the TEV cleavage scar was mutated to a cysteine for

fluorophore conjugation (H2A G-1C) via site-directed mutagenesis. For fluorescence based

binding assays, a cysteine was introduced at H2B D51 via site-directed mutagenesis.

1337

For reconstituting chemically modified H2A-Ub nucleosomes, histones were purified individually

essentially as described(Luger et al., 1999). Briefly, inclusion bodies from 6-8L of histone

expression in LB media were lysed in buffer T (50 mM Tris-HCl pH 7.4, 1 mM EDTA, 100 mM

NaCl), washed by sonication twice in buffer TW (50 mM Tris-HCl pH 7.4, 1 mM EDTA, 100 mM

NaCl, 1% v/v Triton-X100), and then twice again in buffer T. Inclusion bodies were softened by

stirring with a small volume of DMSO (2-3 mL), and extracted with 60 mL of extraction buffer (25

mM Tris-HCl pH 7.5, 7 M guanidinium-HCl, 0.5 mM EDTA, 10 mM DTT). After centrifugation,

the supernatant was dialyzed overnight into ion-exchange buffer (25 mM Tris-HCl pH 7.5, 7 M

deionized urea, 100 mM NaCl, 0.5 mM EDTA, 10 mM DTT), and applied to 2x5 mL HiTrap Q

columns preceding 2x5 mL HiTrap SP columns in-tandem to capture the non-specifically bound

DNA (Cytiva). After sample application, the Q columns were removed, and the SP columns

were eluted using a 12-CV salt gradient (0.1-1M NaCl for H2A and H2B, 0.2-1M NaCl for H3

and H4). The peak fractions were assessed by Coomassie stained SDS-PAGE gel and

Abs_{260}/Abs_{280} value, and the purest fractions were extensively dialyzed into water, aliquoted,

flash frozen, and lyophilized. Lyophilized histones were reconstituted into H2A/H2B dimers,

H3/H4 tetramers, and octamers as previously described(Dyer et al., 2003), and purified via SEC

using Superdex 75 (H2A/H2B) or Superdex 200 (H3/H4 and octamers) columns in SEC buffer

(25 mM Tris-HCl pH 7.5, 2M NaCl, 1 mM DTT).

1356

Fluorophore conjugation to histones and ubiquitin: For fluorophore conjugation, Alexa Fluor 680

C_2 maleimide (ThermoFisher), IRDye 680LT maleimide (Li-Cor), and Oregon green 488

maleimide (AAT Bioquest) were reconstituted in DMSO to 10 mM. Fluorophore maleimides

were mixed with octamers containing single cysteine mutants H2A G-1C for H2A-Ub activity

assays (with Alexa Fluor 680 C_2 maleimide or IRDye 680LT maleimide) or H2B D51C for

binding assays (with Oregon green 488 maleimide). Conjugation reactions were performed

using 20 μM octamer and 100 μM fluorophore overnight at 4 °C in conjugation buffer (25 mM

Tris-HCl pH 7.5, 2M NaCl, 1 mM EDTA, 0.5 mM TCEP), and stopped by addition of 10 mM DTT

and flash frozen. Excess fluorophore was removed after nucleosome reconstitution by SEC

using a Superdex 200 increase 10/300 GL column or extensive buffer exchange in a 30K

MWCO concentrator (Amicon). Typical labelling efficiency was ~70%. For labelling of Ub Q2C,

100 μM Alexa Fluor 680 C_2 maleimide was mixed with ~1 mM of Ub Q2C in buffer (25 mM Tris-

HCl pH 7.5, 1 mM EDTA, 0.5 mM TCEP) at 4 °C overnight. The reaction was quenched with 10

mM DTT, and buffer exchanged into reducing buffer (25 mM Tris-HCl pH 7.5, 1 mM EDTA, 10

mM DTT) in a concentrator to remove unconjugated dye, and stored at -80 °C. The fluorophore

labelling efficiency of Ub was calculated to be ~5%.

1373

Ubiquitin G76C and dichloroacetone crosslinking of H2A-Ub: H2A K15C, K119C, and K127C

were generated by site-directed mutagenesis and purified as described above for individual

histones. His-TEV-ubiquitin G76C was a gift from C. Wolberger (Addgene plasmid #75299) and

purified as previously described(Morgan et al., 2019; Morgan Michael T. et al., 2016).

Dichloroacetone crosslinking of Ub G76C to H2A single cysteine mutants was performed as

previously described in detail (Morgan et al., 2019), with the exception that the final product

containing Ub, Ub-Ub dimers, and H2A-Ub was not purified by RP-HPLC to remove

unconjugated Ub. Instead, it was lyophilized, and mixed with a slight excess of H2B in unfolding

buffer (25 mM Tris-HCl pH 7.5, 7 M guanidinium-HCl, 0.5 mM EDTA, 1 mM DTT), refolded into

H2A/H2B dimers in refolding buffer (25 mM Tris-HCl pH 7.5, 2M NaCl, 1 mM DTT), and purified

by SEC using a 24 mL Superdex 75 column (GE Healthcare) to separate the H2A-Ub from Ub

species.

1386

'601' and MMTV DNA: For large scale purification of 147-bp '601' (gift from K. Luger, CU

Boulder), 185-bp '601' (gift from S. Tan, Penn State Univ.), MMTV (gift from G. Debelouchina,

(UC San Diego), and NLE-trimer '601' (for tri-NCP; gift from K. Luger) DNA, repeat plasmids

from 8-12L of DH5a cells in LB media were subjected to alkaline lysis, phenol-chloroform

extraction, PEG precipitation, EcoRV digestion (NEB), and ion exchange of the excised

fragment by HiTrap DEAE (147-bp '601', 185-bp '601', and MMTV; Cytiva) or mono-Q (NLE-

trimer '601'; GE Healthcare) according to established protocols(Dyer et al., 2003).

1394

For UV-induced Bpa crosslinking and EMSA experiments, fluorophore labelled '601' DNA was

generated using large-scale PCR with Phusion polymerase (produced in-house) from a pGEM-

3z/601 plasmid containing one copy of '601' DNA (gift from J. Widom, Addgene plasmid

#26656)(Lowary & Widom, 1998) with 5'-IRdye700 labelled forward primers (IDT). Following

PCR, the reactions were purified using a 1 mL mono-Q column equilibrated in ion-exchange

buffer (25 mM Tris-HCl pH 8.0, 200 mM NaCl, 1 mM EDTA) over a 10-CV gradient (0.2-1M

NaCl). Peak fractions containing purified 147- and 185-bp DNA fragments were dialyzed into

storage buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA), concentrated, and stored at -20 °C. The

primer sequences were as follows: 147-fwd: ctggagaatcccggtgccgagg; 147-rev:

acaggatgtatatatctgacacg; 185-fwd: atccctatacgcggccgccctggagaatcccggtgccgagg

; 185-rev: atcgctgttcaatacatgcacaggatgtatatatctgacacg.

1406

For generation of DNA fragments with DraIII sticky ends for dinucleosome assembly, large scale

PCR was performed as described for the fluorophore-labelled DNA fragments above. Primer

sequences containing DraIII cut-sites were based off those used by Poepsel *et al*. to make

dinucleosomes with 35-bp of linker DNA(Poepsel et al., 2018). Fragments used to reconstitute

nucleosomes containing fluorophore-labelled H2A nucleosomes (H2A[observe]) were generated

using the "147-fwd" primer (sequence above) and the di-NCP reverse primer sequence

(taggtatcgtat<u>CACGGGGTG</u>agatcgctacaggatgtatatatctgacacg). The fragment for the H2A[silent]

nucleosomes were generated using the "147-rev" and the di-NCP-forward primer sequence

(ctgacttattga<u>CACCCCGTG</u>atgctcgatactgtcatactggagaatcccggtgccgag). The DraIII sites in the

primers are underlined and uppercase. Following PCR purification by 1 mL mono-Q, DNA

fragments were dialyzed into TE/0.1 buffer (10 mM Tris-HCl, 0.1 mM EDTA) and digested with

DraIII-HF in 1x CutSmart buffer for ~20 hr at 37 °C (NEB). Fragments with DraIII sticky ends

were re-purified by 1 mL mono-Q ion-exchange, dialyzed into storage buffer (10 mM Tris-HCl, 1

mM EDTA), concentrated, and stored at -20 °C.

1421

Nucleosome reconstitution: mono-nucleosomes (NCPs) were reconstituted by the standard salt-

dialysis method by mixing histone octamers (or H3/H4 tetramers and H2A/H2B dimers) with

'601' DNA in high salt buffer (25 mM Tris-HCl pH 7.5, 2M NaCl, 0.1 mM EDTA, 1 mM DTT) in a

slide-a-lyzer mini dialysis unit (0.1 mL, ThermoFisher)(Dyer et al., 2003). Low-salt buffer (25 mM

Tris-HCl pH 7.5, 0.1 mM EDTA, 1 mM DTT) was pumped in over ~24 hours in a cold room with

stirring. A final dialysis was performed into NCP storage buffer (25 mM HEPES-NaOH, 10 mM

NaCl, 0.1 mM EDTA, 1 mM DTT). Fluorophore-labelled nucleosomes were further purified by

SEC using a 24 mL Superdex 200 increase 10/300 column equilibrated in NCP storage buffer or

by extensive buffer exchange in a concentrator to remove excess fluorophore. For tri-NCPs, salt

dialysis was performed as described for mono-NCPs using ~1.4x octamer to '601' sites in the

presence of 0.2 equivalents of MMTV competitor DNA. Following dialysis into TEK10 (10 mM

Tris-HCl pH 7.5, 0.1 mM EDTA, 10 mM KCl), trimers were precipitated by adding 4 mM $MgCl_2$,

incubating on ice for 10 minutes and pelleted by centrifugation at 17,000 x g at 4 °C. The

supernatant was discarded, and the pellet was gently resuspended in TEK10, and dialyzed

overnight into NCP storage buffer. Tri-NCP assembly was verified by EcoRI digestion of sites

located between nucleosome units.

1438

For di-NCP assembly, mono-NCPs with DraIII sticky ends were assembled as described above.

The strategy to make asymmetric di-NCPs was based on previously described methods(Dao et

al., 2020; Poepsel et al., 2018). 100 nM of each NCP was mixed in T4 ligase buffer (50 mM

Tris-HCl pH 7.5, 10 mM MgCl$_2$, 1 mM ATP, 10 mM DTT) on ice. T4 ligase (NEB) was added to a

final concentration of 20 U/µL, the reaction was incubated for 30 minutes at 16 °C, and dialyzed

into TEK10 for 4 hours at 4 °C. Following dialysis, MgCl$_2$ was added to a final concentration of

18 mM, incubated at room temperature for 15 minutes, and pelleted at 17,000 x g at 4 °C. The

pellet containing di-NCPs was resuspended in 50 µL of TEK10 and dialyzed overnight into NCP

storage buffer. All chromatin substrates were analyzed for quality on 5% polyacrylamide 0.5x

TBE gels monitoring fluorescence and/or DNA-staining, ensuring minimal free DNA or

contaminating species. Chromatin substrates were stored on ice for no longer than one month.

1450

Design and assembly of DNA competitor fragments: single-stranded, double-stranded, bubble,

frayed, and single-stranded overhang DNA fragments were assembled by annealing ssDNA

oligos (IDT) in IDT duplex buffer (30 mM HEPES-NaOH pH 7.5, 100 mM potassium acetate).

Oligo mixtures were heated to 95 °C and slowly cooled to 25 °C stepwise over 1 hour in a

thermocycler. For NMR experiments, dsDNA and bubble-DNA were buffer exchanged into NMR

buffer in a spin concentrator (Amicon). The sequences of oligos used are as follows. For

ssDNA, dsDNA, bubbleDNA, and ssOverhang a common "bottom" oligo was used

(ggtacacaattgcgctggtaccccaggcgtcgtagg). The oligos annealed to the "bottom" oligo were as

follows: (dsDNA-top: cctacgacgcctggggtaccagcgcaattgtgtacc; bubbleDNA-top:

cctacgacgcctggctctttcccgcaattgtgtacc; ssOverhang-top: ccgcaattgtgtac). For the frayed DNA,

the annealed oligos were as follows: (frayedDNA-top: ggtacacaattgcgccaggcgtcgtaggctggtacc;

frayedDNA-bottom: ctctttcccctacgacgcctggcgcaattgtgtacc).

1463

Nucleosome ubiquitylation assays

Time-course H2A-Ub assays: The general setup for time-course H2A-Ub assays was a reaction

mixture containing 0.2 µM E1 (UBA1), 1 µM E2 (UBE2D3), 30 µM ubiquitin, 0.5 µM NCP

substrate, and 12.5-100 nM E3 (depending on the assay), assembled in 30 µL of reaction buffer

(25 mM HEPES-NaOH pH 7.5, 150 mM NaCl). For assays containing di-NCPs, 10 nM of

fluorophore-labelled di-NCP substrate was mixed with 0.5 µM unlabeled NCP[147] substrate in the

same reaction tube to facilitate quantifiable reaction kinetics. Critically, the salt concentration of

each reaction within an assay was rigorously controlled, as H2A-Ub kinetics are extremely

sensitive to differences in buffer ionic strength. 5 µL of each reaction was mixed 1:1 with 2x

SDS-PAGE load dye as a zero time-point. The mixtures were brought to 37 °C in a heat block,

and 1 µL of 100 mM MgCl$_2$/ATP was added (4 mM final concentration) to initiate the reaction,

and the indicated time-points were obtained by diluting 5 µL the reaction 1:1 with 2x SDS-PAGE

load dye. For experiments with tri-NCPs, the concentration of nucleosome units in the reaction

was normalized to NCP[147] and NCP[185] (0.5 µM NCP unit or 0.17 µM tri-NCP), and reactions

were initiated with 2 mM ATP and 1 mM MgCl$_2$.

1479

Gel samples were run on 15% SDS-PAGE gels and visualized by in-gel fluorescence monitoring

a fluorophore conjugated to the N-terminus of H2A. Alternatively, some assays were visualized

by Western blot for H2A (EMD Millipore, 07-146) or a VSVG epitope tag on the N-terminal tag of

H2A (Sigma, V4888) and a fluorescently labelled secondary antibody (Cell Signaling, 5151S).

Gels and blots were imaged on a Li-Cor Odyssey imager (Li-Cor). Band quantification was

performed using Image Studio software (Li-Cor) by drawing a box encompassing the unmodified

H2A band to measure its intensity, performing background subtraction, and normalizing to the

zero time-point of a given reaction.

1488

Inhibition of H2A-Ub by competitor DNA: A reaction mixture containing 0.2 µM E1 (UBA1), 3 µM

E2 (UBE2D3), 30 µM ubiquitin, 0.5 µM NCP substrate, and 50 nM E3 (or 750 nM for

BRCA1[RING]/BARD1[RING]) was assembled in 9 µL of reaction buffer (25 mM HEPES-NaOH pH

7.5, 100 mM NaCl) in PCR strips. To that, 1 µL of DNA competitor at 10x concentration was

added. Reactions were brought to 37 °C in a thermocycler, started by addition of 5 µL of 10 mM

ATP/MgCl$_2$ in reaction buffer, and allowed to proceed for 12-20 minutes depending on the E3.

Reactions were stopped by addition of 10 µL of 4x fluorescence sample load dye (Li-Cor),

resolved on a 15% SDS-PAGE gel, and visualized using a Li-Cor Odyssey imager.

Quantification was performed as described above, monitoring the intensity of unmodified H2A,

and normalizing each reaction to a -ATP reaction.

1499

Auto-ubiquitylation assay: Reactions were assembled with 0.2 µM E1 (UBA1), 1 µM E2

(UBE2D3), 10 µM ubiquitin (Q2C-Alexa Fluor 680), and 0.2 µM of the indicated E3 in buffer (25

mM HEPES-NaOH pH 7.5, 50 mM NaCl), and were initiated by addition of 5 mM ATP/MgCl$_2$.

After 60 minutes at 37 °C, reactions were quenched by mixing 1:1 with 4x fluorescence sample

load dye (Li-Cor), resolved on a 4-20% SDS-PAGE gel (BioRad), and visualized using a Li-Cor

Odyssey imager.

1506

Nucleosome binding assays

Electrophoretic mobility shift assay: For EMSA experiments, 5 µL of 10 nM of substrate (NCPs

with either H2A G-1C labelled with Alexa Fluor 680 or DNA/NCPs labelled with IRDye700) were

mixed with 5 µL of indicated RING/RING heterodimer or BARD1 IDR constructs (2x

concentrated) in EMSA buffer (15 mM Tris-HCl pH 7.5, 90 mM NaCl, 1 mM DTT, 0.1 mg/mL

BSA, 0.05% triton X-100) in PCR strips. Binding was allowed to proceed for 10 minutes at room

temperature, followed by 5 minutes on ice. 5 µL of 25% sucrose was added to each tube, and

15 µL of sample was loaded onto 5% polyacrylamide native gels (0.5x TBE) in pre-chilled 0.5x

TBE buffer. Gels were run at 4 °C for 90-120 minutes at 110V and visualized using a Li-Cor

Odessey scanner (Li-Cor).

1517

Fluorescence-quenching binding experiments: Assay set up was performed essentially as

previously described (Hu et al., 2021). 10 μL of 10 nM NCPs with Oregon green 488 conjugated

to H2B D51C in low-salt fluorescence buffer (25 mM Tris-HCl pH 7.5, 10 mM NaCl, 0.01%

CHAPS, 0.01% NP40, 0.1 mg/ml BSA, 1 mM DTT) were mixed with the 10 μL of the indicated

BRCA1/BARD1 construct at 2x concentration in high-salt fluorescence buffer containing either

90 mM, 190 mM, or 290 mM NaCl, depending on the assay, in a 384-well low volume black

round bottom polystyrene non-binding surface microplate (Corning). Plates were briefly mixed,

incubated at 22 °C for 10-15 minutes, and scanned using a BioTek Synergy Neo2 plate reader

equipped with a fluorescence filter (excitation: 485/20 nm, emission: 528/20 nm). The gain was

set to 100 for all experiments. Two to three technical replicates for each independent replicate

experiment (fresh NCP and BRCA1/BARD1 dilutions) were performed, and the readings from

two scans were averaged. BRCA1/BARD1/nucleosome binding interactions were quantitatively

analyzed by fluorescence quenching assuming the following systems of equations:

1531
$$\theta = \frac{K_D + [NCP] + [BCBD] - \sqrt{(K_D + [NCP] + [BCBD])^2 - 4 \times [NCP] \times [BCBD]}}{2 \times [NCP]}$$

1532
$$F = [NCP] \times F_{NCP} \times (1 - q_{BCBD} \times \theta) + N(0, error)$$

where θ is the fraction of nucleosome bound by BRCA1/BARD1 (BCBD), F_{NCP} is the

fluorescence per nucleosome, q_{BCBD} is the fraction of fluorescence quenching per

BRCA1/BARD1 binding, and fluorescence measurements are gaussian distributed with mean 0.

Maximum likelihood parameter estimates were determined using custom Python scripts with

optimization by a differential evolution algorithm implemented in Scipy (Virtanen et al., 2020). K_d

and q_{BCBD} were separately estimated for each BRCA1/BARD1 construct, and NCP

concentration was treated as a random effect per individual dilutions from higher concentration

stock solutions, with the total number of model parameters being equal to number of NCP

dilutions + (2 x number of BCBD constructs) + 2. The 95% confidence limits were estimated by

computing the profile likelihood with an applied Bonferroni correction.

1543

UV-induced Bpa crosslinking

NCPs and DNA for UV-induced crosslinking assays were reconstituted using a 185-bp '601'

sequence labeled with an IRdye700 fluorophore (described above). 50 nM NCPs or DNA were

incubated with 500 nM of the BRCA1$^{1-104}$-(GS)$_6$-BARD1$^{26-221}$ fusion (wild-type or with different

Bpa-incorporated sites L120Bpa, W146Bpa, Y180Bpa, A195Bpa, W218Bpa) for 15 minutes at

4°C in 30 uL UV Crosslinking buffer (20 mM HEPES-NaOH pH 7.5, 100 mM NaCl) in a 96-well

clear round bottom polystyrene non-treated microplate (Corning). Following incubation, 5 μL

was added to 5 μL 2X SDS-PAGE gel-loading dye for the -UV sample. Samples were then

subjected to UV irradiation (365 nm) using a Blak-Ray B-100AP/R High Intensity UV Lamp

(UVP) from 20 cm for 30 minutes at 4°C. Following irradiation, 5 μL was added to 5 μL of 2x

SDS-PAGE gel-loading dye. Samples were heated at 60°C for 5 minutes and separated on 10%

SDS-PAGE gels. Gels were visualized using a Li-Cor Odyssey scanner (Li-Cor). Band

quantification was performed using Image Studio software (Li-Cor) by drawing a box measuring

the total band intensity for each crosslinked product, subtracting the background signal, and

normalizing to the crosslinked band of the L120Bpa fusion mutant. For the Bpa UV crosslinking

DNA competition assays, 50 nM NCPs were incubated with 500 nM BRCA1-f-BARD1^{221}

(W146Bpa or A195Bpa) with varying amounts of competitor 36-mer dsDNA or bubble-DNA (0,

0.3125, 0.625, 1.25, 2.5, 5, or 10 μM) for 15 minutes at 4°C, and assays were performed as

described above.

1563

Cellular Studies

Mammalian cell culture and transfection: HeLa cells were grown in Dulbecco's modified Eagles

medium (DMEM) supplemented with 10% fetal bovine serum (Sigma), 100 μg/ml streptomycin,

and 100 U/ml penicillin (Sigma). The cells were tested for mycoplasma contamination by

Bionique testing labs (http://www.bionique.com/). To create a FLP-in version of HeLa, we stably

integrated a flippase recognition target (FRT) sequence into the cells by using the pFRT/lacZeo

plasmid (Thermo Fisher Scientific). We tested Zeocin-resistant clones that had a single

integration site detected by Southern blot for high-activity integration sites by using the

mammalian b-galactosidase activity assay (Gal-Screen, Thermo Fisher Scientific). Clonal

expansion of the selected colony established the HeLa-FRT cell line. To generate stable HeLa-

FRT shBARD1 cells, Dharmacon™ TRIPZ™ lentiviral shRNAs against BARD1 (RHS4696) were

transfected with their respective plasmids and individual clones were selected with 2 µg/ml

puromycin. To generate HeLa-FRT shBARD1 cell lines expressing HA-BARD1 WT or mutant

(Δ194-216), cells were transfected with their respective plasmids and individual clones were

selected with 200 µg/ml hygromycin. BARD1 was cloned into pcDNA5/FRT-HA vector.

QuikChange site-directed mutagenesis was used to construct the mutant variant (Δ194-216) of

BARD1.

1581

Immunoblot analysis: HeLa-shBARD1 cells stably expressing wild type or Δ194-216 mutant of

HA-BARD1 were pretreated with doxycycline(1µg/ml) for three days to knockdown the

endogenous BARD1, and then incubated with or without Olaparib (10 µM) for another 24 h.

Cells were trypsinized and pellets were washed by PBS. Protein was extracted using NETN

buffer (20 mM Tris-HCl pH 8, 420 mM NaCl, 1 mM EDTA, 0.5% IGEPAL, 1 mM DTT, and

Roche Protease Inhibitor Cocktail) and were resolved in 8% Bis-Tris SDS-PAGE gels, and blots

(20-50 µg of total protein) were probed with the following antibodies: HA (3724S, Cell Signaling;

1:1500), BRCA1 (SC6954, Santa Cruz; 1:500), BARD1 (ab50984, Abcam; 1:1000), Tubulin

(2128S, Cell Signaling; 1:2000), Lamin B1 (SC374015, Santa Cruz, 1:500), according to the

instructions provided by the manufacturers. The blots were incubated with HRP-conjugated

secondary antibodies (Pierce 31450 for rabbit anti-mouse IgG-HRP; Sigma A6154 for goat anti-

rabbit IgG-HRP) before being visualization of protein signals using the ECL max kit (Bio-rad).

1594

Preparation of cytoplasmic and nuclear extracts: The REAP method for the preparation of

cytoplasmic and nuclear extracts was followed. Briefly, HeLa cells from 10-cm dishes were

washed with ice-cold phosphate buffer saline (PBS) pH 7.4, collected by centrifugation,

resuspended in 900 μL of ice-cold PBS with 0.1% NP40 and protease inhibitors, and triturated 5

times using a p1000 micropipette. The lysed cell suspension was centrifuged for supernatant

(this is the cytoplasmic fraction) and the pelleted nuclei was washed once with PBS with 0.1%

NP40 and lysed by NETN buffer with protease inhibitors and different NaCl concentrations (100,

300 and 420 mM) to yield the nuclear extract fractions. Each soluble fraction was labeled as

NS100, NS300 and NS420, while the final pellets were resuspended with NETN420 buffer and

sonicated for 10 seconds for NP420 fraction. The cytoplasmic and nuclear fractions, 20 μg

each, were analyzed by immunoblotting for their content of BRCA1, BARD1, Tubulin and Lamin

B1.

Immunofluorescence microscopy and image analysis: Cells were pretreated with doxycycline (1

μg/ml) for three days to knockdown the endogenous BARD1. Cells were trypsinized and 4×10^{4}

cells seeded in the 4-chambers slide with or without Olaparib (10 μM) for another 24 h. Cells

were pre-extracted on ice for 10 min with cold cytoskeleton buffer (10 mM PIPES pH 6.8, 100

mM NaCl, 300 mM sucrose, 3 mM MgCl2, 1 mM EGTA, 0.5% Triton X-100) followed by 10 min

with cytoskeleton stripping buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 1%

Tween 20 (v/v), 0.5% sodium deoxycholate). Pre-extracted cells were then fixed at room

temperature for 10 min with 4% paraformaldehyde, washed with PBS, and treated for 10 min

with 1% Triton X-100 in PBS. Primary anti-HA antibody (3724S, Cell Signaling; 1:500) was

incubated overnight at 4°C in 5% goat serum in PBS. Alexa Fluor-568 conjugated secondary

antibody (Jackson Immunno Research Labs) was incubated for 1 h at room temperature, and

1619   DAPI were incubated for another 15 mins in room temperature. Then slides mounted using

1620   antifade mounting media (9071, Cell signaling). A random selection of 10-11 areas were imaged

1621   using an Evos M5000 Microscope. HA-BARD1 foci>5 and HA-BARD1 foci per nuclear were

1622   quantified by the Celleste software and Prism GraphPad software was used for the final figure

1623   and significance analysis.

1624

Clonogenic survival assay: HeLa cell lines stably expressing HA-BARD1 wild type or mutant

were pretreated with doxycycline for 3 days. Then, 400 cells/well were seeded into 12-well

plates, treated with indicated amount of Olaparib (Selleckchem) or Cisplatin (Selleckchem) in

regular growth medium for 11-12 days. Cells were fixed with methanol and stained with 0.5%

crystal violet in methanol before colonies were counted. Clonogenic survival was determined for

a given concentration of cells that were plated by dividing the number of colonies on each

treated plate by the number of colonies on the untreated plate, taking the plating efficiency of

untreated cells into account.

1633

Chemical Crosslinking and Mass Spectrometry Analysis

Sample preparation: Reactions were 90 µL of protein/nucleosome mix containing 2 mM

BRCA1[RING]/BARD1[FL] and 2 mM of unmodified H2A or H2A K15-Ub nucleosomes wrapped with

147-bp '601' DNA in crosslinking bufffer (25 mM HEPES-NaOH pH 7.5, 150 mM NaCl, 1%

glycerol, 1 mM DTT) plus 3.06 µL of 14.5 mM DSS or BS3 (final concentration 0.5 mM) (DSS:

disuccinimidyl suberate, BS3: bis(sulfosuccinimidyl)suberate, ThermoFisher). Crosslinking was

carried out for 10, 30 or 45 mins at room temperature before removing 30 µL and quenching by

addition of 3 µL of 1 M ammonium bicarbonate at room temperature for 30 minutes. Four

separate reaction conditions were performed: (1) WT-NCP DSS; (2) WT-NCP BS3; (3) K15Ub-

NCP DSS; (4) K15Ub-NCP BS3. After quenching, 9 µL of each sample at each timepoint was

analyzed by SDS-PAGE **(see Extended Data Fig 10a)** and the remainder was stored at -80°C

until processing for mass spectrometry analysis.

1646

Reactions were prepared for MS analysis by bringing them up to 0.1% PPS silent surfactant

(Expedion Inc.), 5 mM TCEP. Samples were reduced for 60 mins at 60°C in an Eppendorf

Thermomixer with shaking (1200 rpm). Alkylation was performed at room temperature in the

dark for 30 mins with 6 mM iodoacetamide. Excess iodoacetamide was quenched by addition of

5 mM DTT and samples were then digested by trypsin digestion (Sequencing Grade Modified

Trypsin, Promega Corp) at 37°C for 6 hours in a Thermomixer with shaking (1000 rpm) at a

substrate to enzyme ratio of 15:1 prior to acidification with 250 mM HCl (final concentration).

Acidified samples were incubated for 1 hour at room temperature, spun at max speed in a

microfuge and supernatants containing peptides were transferred to autosampler vials and

stored at -80°C until analysis.

1657

Chromatography: Mass spectrometry and data analysis was based on previously described

methods(Zelter et al., 2015). For each injection, 3 µl of protein digests were loaded by

autosampler onto a 150-µm Kasil fritted trap packed with 2 cm of Reprosil-Pur C18-AQ (3-µm

bead diameter, Dr. Maisch) at a flow rate of 2.5 µl/min. After desalting with 8 µl of 0.1% formic

acid plus 2% acetonitrile, the trap was brought online with a Self-Packed PicoFrit Column (New

Objective part number PF360-75-10-N-5, 75 µm i.d.) packed with 30 cm of Reprosil-Pur

C18-AQ (3-µm bead diameter, Dr. Maisch) mounted to a heated nanospray ionization source

(CorSolutions LLC) set at 50°C and placed in line with a Thermo Scientific EASY-nLC 1200

UPLC pump plus autosampler.

1667

1668   Peptides were eluted from the column at 0.25 µL/min using an acetonitrile gradient consisting of

1669   the following steps: (1) 0-10 mins; 6-10% B; (2) 10-90 mins; 10-32% B; (3) 100-130 mins; 32-

1670 75% B; (4) 130-135 mins; 75% B; (5) 135-136 mins; 75-100% B; (6) 136-151 mins; 100% B,

1671 followed by re-equilibration to 0% buffer B prior to the subsequent injection. Buffer A was: 0.1%

1672 formic acid in water and buffer B was 0.1% formic acid 80% acetonitrile.

1673

Data acquisition: A Thermo Fisher Scientific Exploris 480 was used to perform mass

spectrometry in positive ion mode with the following settings. Data dependent acquisition (DDA)

mode was used with a maximum of 20 tandem MS (MS/MS) spectra acquired per MS spectrum

(scan range of *m/z* 400–1,600). The resolution for MS and MS/MS was 60,000 at *m/z* 200. The

normalized automatic gain control targets for both MS and MS/MS were set to 100%, and the

maximum injection times were 50 and 100 ms for MS and MS/MS scans, respectively. MS/MS

spectra were acquired using an isolation width of 2 *m/z* and a normalized collision energy of 27.

MS/MS acquisitions were prevented for +1, +2, ≥+6 or undefined precursor charge states.

Dynamic exclusion was set for 10 seconds. MS and MS/MS spectra were collected in centroid

mode.

1684

MS data processing – identification of crosslinked peptides: Raw mass spectra were converted

into mzML using ProteoWizard's msconvert(Chambers et al., 2012). All proteins in the sample

were identified using Comet(Eng et al., 2013) searching against a database including the entire

S. frugiperda, *E. coli* and *H. sapiens* proteomes (Uniprot 7108, 83333 and 9606, respectively)

plus common contaminants (https://www.thegpm.org/crap/) along with the sequences of all

heterologously expressed proteins (FLAG-BRCA1, P38398; TwinStrepII-BARD1, Q99728; H2A-

2A, Q99728; H2B-1K, O60814 ; H3.2 (C110A), Q71DI3; H4, P62805; Ub G76C, sequence

derived from P0CG48). Smaller search databases were made for subsequent XL searching

consisting only of proteins identified in initial comet searches by at least 3 peptides with a

Percolator assigned q-value of ≤ 0.01. Decoy databases consisted of the corresponding set of

reversed protein sequences. Cross-linked peptides were identified within these proteins by

Kojak version 2.0.0-alpha8 available at (http://www.kojak-ms.org/)(Hoopmann et al., 2015). A

statistically meaningful q-value was assigned to each peptide spectrum match (PSM) through

analysis of the target and decoy PSM distributions using Percolator version 2.08(Käll et al.,

2007). All data reported in this paper were filtered to show hits to the target proteins that had a

Percolator assigned peptide level q value ≤ 0.05.

1701

NMR experiments and resonance assignments.

All spectra were collected at 25° C in NMR buffer (25 mM MOPS-NaOH, 100 mM NaCl, 1 mM

EDTA, and 0.5 mM TCEP, pH 7.0) with 7% D_2O. NMR HSQC and titration data were completed

on a Bruker 800 MHz Avance III spectrometer with cyroprobe. (^{1}H,^{15}N)-HSQC spectra were

collected on the following ^{15}N-BARD1 IDR constructs: BARD1 Ser-124-424, Ser-124-270, 269-

424, and Ser-141-216.

1708

(^{1}H,^{15}N)- and (^{1}H,^{13}C)-HSQC titration experiments were completed for BARD1 IDR + 36mer-

dsDNA or 36mer-bubDNA by adding unlabeled DNA to 150 µM [^{13}C,^{15}N]-BARD1 124-270 or

150 µM [^{13}C,^{15}N]-BARD1 Ser-141-216, maintaining a constant concentration of the labeled

species. Only apo and 1:1 spectra for the [^{13}C,^{15}N] BARD1 Ser-141-216 construct were

obtained since intermediate titration points resulted in loss of sample. Additionally, titration

spectra with nucleosome were collected for 150 µM [^{13}C,^{15}N]-BARD1 124-270 + 0.05x NCP[147].

A (^{1}H,^{15}N)-HSQC was collected on 150 µM ^{15}N-BARD1 269-424 + 150 µM 28mer-dsDNA.

1716

Dynamics experiments were collected on a Bruker 500 MHz Avance III spectrometer with a

room temperature probe. T_1 and T_2 ^{15}N-Trosy HSQC experiments for 150 µM [^{13}C,^{15}N]-BARD1

Ser-141-216, 150 µM [^{13}C,^{15}N]-BARD1 Ser-141-216 + 150 µM 36mer-dsDNA, and 150 µM

[^{13}C,^{15}N]-BARD1 Ser-141-216 + 150 µM 36mer-bubble-DNA were collected in an interleaved

manner with 8 points each(G. Zhu et al., 2000). T_1 delays were 10, 40, 80, 120, 160, 320, 640,

and 1000 ms; T_2 cpmg loop delays were 8.48, 16.96, 25.44, 25.44, 42.40, 50.88, 67.84, and

84.80 ms. T_1 and T_2 for each residue were fitted to a single exponential with errors reflecting the

quality in the fit. Residues corresponding to peaks with overlapping intensities were excluded

from the analysis.

1726

Backbone chemical shift assignments were determined from standard backbone triple-

resonance experiments (HNCA, HNCOCA, HNCOCACB, HNCACB, HNCO, and HNCACO) and

an HNHA experiment using a Bruker 800 MHz Avance III spectrometer with cryoprobe. The 3D

data were collected for 500 µM [^{13}C,^{15}N]-BARD1 Ser-124-270 and 500 µM [^{13}C,^{15}N]-BARD1

Ser-141-216. Assignments from the shorter construct were transferred as appropriate to the

longer BARD1 construct. HNCACB and HNCOCACB spectra were collected on the 1:1 samples

of 150 µM [^{13}C,^{15}N]-BARD1 Ser-141-216 with either 36mer dsDNA or 36mer bubble-DNA.

1734

The NH chemical shift perturbation was calculated according to $\Delta\delta_{NH}$ (ppm) = sqrt [$\Delta\delta_H^2$ +

$(\Delta\delta_N/5)^2$]. Component ^{1}H and ^{15}N chemical shift differences are (free – bound) in ppm. The

CaCb CSP was calculated according to $\Delta\delta_{CaCb}$ (ppm) = 0.25*[(Cα-Cβ)$_{free}$-(Cα-Cβ)$_{bound}$].

Secondary structure propensity (SSP) was determined from $\Delta\delta$(Cα-Cβ) = (Cα-Cβ)$_{measured}$ - (Cα-

Cβ)$_{random coil}$, where the random coil shifts were generated from BARD1 IDR sequence using a

webserver (https://spin.niddk.nih.gov/bax/nmrserver/Poulsen_rc_CS/)(Kjaergaard & Poulsen,

2011).

1742

1743   **Multiple Sequence Alignment**

1744   Mammalian BARD1 ortholog sequences were downloaded from Ensembl (v106), globally

1745   aligned using MAFFT(Katoh & Standley, 2013), and trimmed to BARD1 residues 124-270.

1746

Materials, Data, and Code availability Statement

Source data for H2A-Ub and binding assays as well as uncropped gels and blots and original

microscopy images are included as supplemental files. For crosslinking mass-spectrometry

analysis, the complete unfiltered list of all PSMs and their Percolator assigned q values are

available on the ProXL web application(Riffle et al., 2016, 2019)

https://proxl.yeastrc.org/proxl/p/bard1nucleosome along with the raw MS spectra and search

parameters used. In addition, complete search algorithm configuration files, fasta search

databases, raw search output and raw MS data files were deposited to the ProteomeXchange

Consortium via the PRIDE(Vizcaíno et al., 2009) partner repository with the dataset identifier

PXD035345. NMR chemical shift assignments for BARD1 Ser-141-216 and associated DNA-

bound assignments have been deposited to the Biological Magnetic Resonance Bank with

dataset identifiers 51523 (apo), 51524 (with dsDNA), 51525 (with bubble-DNA). Python scripts

used to analyze and plot fluorescence-based binding data are included with this manuscript.

Plasmid reagents generated in this study can be obtained upon request from the corresponding

author Dr. Rachel Klevit.