1 **Inhibition of PI3K p110δ activity reduces IgE production in IL-4 and anti-**

2 **CD40 stimulated human tonsil B cell cultures**

4 Anna Cutrina-Pons[1*], Aloka De Sa[1*], David J Fear[2], and Faruk Ramadani[1]

Author Affiliations: [1] Randall Centre of Cell & Molecular Biophysics, and [2]

Division of Asthma, Allergy and Lung Biology, King's College London, United

Kingdom. *The first 2 authors contributed equally to this article, and both

should be considered first author.

Correspondence: Faruk Ramadani, Randall Centre of Cell and Molecular

Biophysics, King's College London, London, SE1 1UL, UK; e-mail:

faruk.ramadani@kcl.ac.uk or Phone: 44 (0)20 7848 6442; Fax: +44 (0)20 7848

6435.

This study was supported by Asthma UK career development award (AUK-

CDA-2019-412) to F.R.

Abstract

Background: Phosphoinositide 3-kinase (PI3K) p110δ signalling negatively regulates the production of mouse IgE. However, there are disparities between the mouse and human IgE biology, and the role of PI3K p110δ in the production of human IgE is yet to be determined.

Objective: To investigate the effect of PI3K p110δ inhibition in the production of human IgE.

Methods: Human tonsil B cells were stimulated with IL-4 and anti-CD40 antibody to induce class switching to IgE and IgG1 in the presence or absence of IC87114, a small molecule inhibitor of PI3K p110δ. Using FACS, RT-PCR and ELISA we examined the effect of PI3K p110δ inhibition on IgE production and also determined the mechanisms involved.

Results: Unlike in mice, we observed that PI3K p110δ inhibition significantly reduces the number of IgE+ switched cells and the amounts of secreted IgE in IL4 and anti-CD40 cultures. However, the number of IgG1+ cells and secreted IgG1 were largely unaffected by PI3K p110δ inhibition. The expression levels of AID, ε and γ1 germinal transcripts or other factors involved in the regulation of CSR to IgE and IgG1 were also unaffected by IC87114. However, we found that IC87114 significantly decreases the proliferation of tonsil B cells stimulated with IL-4 and anti-CD40, specifically reducing the frequency of cells that had undergone 4 divisions or more.

Conclusion: Our data identifies proliferation as the key mechanism through which PI3K p110δ regulates CSR of human B cells to IgE and highlights PI3K p110δ as a pharmacological target for the treatment of allergic disease.

Keywords

IgE, allergy, class-switching, human B cells, Phosphoinositide 3-kinases

p110δ, signal transduction

Abbreviations

AID   Activation-induced cytidine deaminase

CFSE   Carboxyfluorescein diacetate succinimidyl ester

CSR   Class switch recombination

FACS   Fluorescence activated cell sorting

GLT   Germline transcription

NFκB   Nuclear factor-kappa B

PI3K   Phosphoinositide 3-kinase

PTEN.   Phosphatase and tensin homolog

STAT6   Signal transducer and activator of transcription-6

Introduction

Immunoglobulin (IgE) antibodies play a fundamental role in the pathogenies of allergic disease. They bind to the high affinity IgE receptor (FcεRI) on mast cells and basophils, which, when cross-linked by multivalent allergens, trigger cell activation and the release of various mediators of immediate hypersensitivity and allergic inflammation [1,2]. Since its discovery, over a half a century ago, we have made significant progress not only in understanding the immunological mechanisms associated with type 1 hypersensitivity reactions[1,2] but also in our understanding of the mechanisms involved in the regulation of IgE production[3–9].

The commitment of a B cell to undergo CSR to IgE is dependent on signals downstream of the tumour necrosis factor receptor superfamily member CD40, after its interaction with the CD40 ligand (CD40L) on the surface of T cells, and IL4/L13 binding to their receptors on B cells[2,10]. These signals activate the nuclear factor-κ B (NFκB) and signal transducer and activator of transcription 6 (STAT6) transcription factors that act synergistically to regulate CSR to IgE[10–12]. Both NFkB and STAT6 bind to the Iε promoter and initiate the synthesis of ε germline transcripts (εGLT) and make ε switch region available for genetic recombination. Furthermore, NFκB and STAT6 can also activate the expression of the enzyme activation-induced cytidine deaminase (AID)[13], an important regulator of CSR and SMH, which produces DNA breaks in the switch regions that precede the Cε exon[10].

Several molecular mechanisms have evolved to ensure rigorous regulation of IgE production[10,14]. In mice, they also include signalling by PI3K p110δ[15–17], a

member of the class IA PI3K catalytic isoforms, which is activated downstream

of various receptors on B cells[18]. Inhibition of the PI3K p110δ signalling in

mouse B cells, both *in vitro* and *in vivo,* increases IgE production[15–17]. This

effect of PI3K p110δ inhibition was found to be associated with increased εGLT

and AID expression levels, suggesting that under normal conditions PI3K

p110δ signalling supresses IgE production by repressing εGLT and AID

expression. This was surprising since signalling by PI3K p110δ is also required

for the production of type 2 cytokines such as IL4 and IL13[19,20]. It was later

suggested that PI3K p110δ signalling limits IgE production in a B cell-intrinsic

manner, a mechanism that involves the activation of Bcl6[16], a known negative

regulator of CSR to IgE[21].

However, the B cell regulatory systems of IgE production in humans differ from

those in mice[7], and the role of PI3K p110δ signalling in the production of human

IgE is yet to be determined. Here, we isolated human tonsil B cells and

stimulated CSR to IgE with IL4 and anti-CD40. To investigate the involvement

of PI3K p110δ signalling in human IgE production, the B cell cultures were

treated with IC87114, a small molecular inhibitor of p110δ also used in mouse

B cell cultures[15,17]. The data show that IC87114 reduces the IgE production in

tonsil B cell cultures stimulated with IL-4 and anti-CD40 antibody. Interestingly,

this effect was shown to be specific for IgE, with the production of IgG1 being

largely unaffected. This result is in striking contrast with the effect of PI3K p110δ

inhibition in mouse B cells[19,20].

Materials and methods

Ethics

Following full informed written consent, tonsils were obtained from patients undergoing routine tonsillectomies. The study was conducted at and in accordance with the recommendations of King's College London and Guy's and St Thomas's NHS Fundation Trust and the protocol was approved by the London Bridge Research Ethics Committee (REC number 08/H0804/94).

Cell culture

Tonsils were obtained from patients undergoing routine tonsillectomies. The protocol was approved by the London Bridge Research Ethics Committee (REC number 08/H0804/94), and the study was conducted at and in accordance with the recommendations of Guy's and St Thomas's NHS Fundation Trust and King's College London. The untouched resting B cells were isolated from the human tonsils by depletion of other cells using MojoSort Human B Cell (CD43-) Isolation Kit (BioLegend), whereas the naïve B cells were isolated using the EasySep Human Naïve B cell isolation kit (Stem Cell Technologies). Purified B cells were then cultured at 0.5×10^6 cells/mL in RPMI 1640 (Lonza) containing penicillin (100 IU/ml), streptomycin (100μg/ml) and glutamine (2mM, Invitrogen) and 10% Foetal Calf Serum (Hyclone; Perbio Biosciences). To induce CSR to IgE, cultures were stimulated with IL-4 (200 IU/ml; R&D Europe Systems Ltd) and anti-CD40 antibody (0.5 µg/ml; G28.5; American Type Culture Collection). To investigate the role of PI3K p110δ signalling in human IgE production, cell cultures were also treated with different concentration of IC87114 (LKT

Laboratories Inc), a small molecule inhibitor of PI3K p110δ signalling used in

the mouse studies. Considering that IC87114 has an IC_{50} of 0.5µM, it was used

at concentrations below and above this IC_{50}; 0.125µM, 0.25µM, 0.5µM, 1µM,

and 2µM. The cells were then incubated for up to 10 days at 37°C with 5%

CO_2.

Flow cytometry

To determine the effect of the PI3K p110δ inhibition on STAT6 and NFκB

phosphorylation levels, cells were harvested 24 hours after culture with IL-4

and anti-CD40 and fixed with 2% paraformaldehyde (PFA), washed with PBS

and permeabilised with 1 ml of ice cold methanol for 20 mins on ice. The

permeabilised cells were then intracellularly stained for 1 hour at room

temperature (RT) with anti-STAT6^{Tyr641} Alexa Fluor 647 (A15137E;

Biolegend) and anti-NFκB p65^{Ser536} Alexa Fluor 488 (93H1; Cell Signalling

Technology). To determine the effect of the PI3K p110δ inhibition on CSR to

IgE, cells were harvested on day 10 of culture and stained with a live/dead

fixable stain dye (Life Technologies Ltd) followed by fixation with 2%

paraformaldehyde and permeabilization with PBS containing 0.5% Triton X-

100, and 0.5% saponin. The cells were then stained with anti-human IgE FITC

(IgE21; eBioscience) and anti-IgG1-PE human (IS11-3B2.2.3; Miltenyi Biotec)

for 30 minutes at room temperature in the dark. Following the staining cells

were washed and resuspended in FACS buffer and acquired with BD FACS

Canto.

Proliferation

Following the CFSE labelling, performed using CellTrace™ CFSE Cell proliferation kit (C34554, Molecular Probes™ Invitrogen), the cells were cultured as above, with IL-4 and anti-CD40 antibody, in the presence or absence of 2µM IC87114. After 10 days of culture, cells were harvested and acquired using FACSCanto. The data was analysed using FlowJo and the proliferative capacity of cultured tonsil B cells was determined based on the dilution of the CFSE labelling.

ELISA

Maxisorp plates (Nunc) were coated with polyclonal mouse anti-human IgE (Dako Agilent Technologies) or mouse anti-human IgG1 (BD Pharmingen) in pH 9.8 carbonate buffer overnight at 4°C. Unbound sites were blocked with 2.5% BSA (Sigma-Aldrich) for 2 hours at RT and then washed 4 times with PBS+0.05% Tween-20. Supernatants from day 10 cultures were then added at appropriate dilutions and plates incubated overnight at 4°C. Human serum IgE (NIBSC) or IgG1 (Sigma-Aldrich) were used to construct standard curves. Binding was detected by polyclonal goat anti-human IgE-HRP (Sigma-Aldrich) or mouse anti-human IgG HRP (BD Pharmingen) in 1% PBS/BSA for 2 h at 37°C. The color reaction was developed using the 1- StepTM Ultra TMB-ELISA substrate solution (Thermo Scientific) and plates were read at 450nm using the Labtech LT 4500 microplate reader. IgE and IgG1 concentrations were then calculated from the standard curve using Prism 9.2 software (GraphPad).

qRT-PCR

Total RNA was isolated from cells using the Qiagen RNeasy Plus Mini Kit

(Qiagen) and the RNA concentrations were measured using the NanoDrop

2000 (Thermo Scientific). The isolated RNA was then reverse transcribed into

cDNA using SuperScript III First-Strand Synthesis SuperMix for qRT-PCR

(Invitrogen). To evaluate the expression levels of certain genes we used

TaqMan MGB gene expression assays and TaqMan Universal PCR Master Mix

on Viia7 RT-PCR machine (Applied Biosystems). Gene expression was

standardised to an endogenous reference gene 18s rRNA (Hs99999901_s1,

Applied Biosystems). AID (Hs00757808_m1) and BCL6 (Hs00153368_m1)

gene specific qPCR assays (Applied Biosystems) were used with Taqman

MGB chemistry. εGLT and γ1GLT were analysed using previously published

Taqman primers and probes [22]. The sequences of forward and reverse primers

for εGLT (fw: 5′-CTGTCCAGGAACCCGACAGA-3′; rev: 5′-TGCA

GCAGCGGGTCAAG-3′) and γ1GLT (fw: 5′-CCAGGGCAGGGTCAGCA-3′;

rev: 5′-GGTGCTCTTGGA GGAGGGT-3′) (Sigma-Aldrich) and their

corresponding FAM-labelled MGB probes (εGLT: 5′-AGGCACCAAATG-3′) and

γ1GLT (γ1GLT: 5 ′-CTCAGCCAGGACCAAG-3′) (Applied Biosystems) were

designed in house. All gene specific assays were multiplexed with the 18s

endogenous control assay and run-in triplicate. SDS software was used to

quantify the target cDNA relative to 18s according to the $2^{(-\Delta ct)}$ method.

STATISTICAL ANALYSIS

Statistical analysis was performed using the One-Way ANOVA, with Tuky's

correction, or unless otherwise stated. A *p* value of < 0.05 was considered

significant (**p* < 0.05,***p* < 0.01,****p* < 0.001).

RESULTS

IC87114 inhibits CSR of human tonsil B cells to IgE *in vitro*

Studies in mice have shown that CSR to IgE can be enhanced both *in vivo* and

in vitro when the activity of PI3K p110δ is inhibited[15–17]. This effect of p110δ

inhibition is not exclusive to CSR to IgE, with CSR to IgG1 also being enhanced

in mouse B cells[15,17,23]. To investigate whether CSR to IgE and IgG1 in human

B cells is also affected by the inhibition of the p110δ signaling, as in the mouse

system, the isolated untouched tonsil (CD43⁻) B cells were cultured with IL-4

and anti-CD40 and different concentrations of the IC87114. After 10 days of

culture, we observed an IC87114 dose dependent reduction in the frequency

of IgE⁺ cells (Figure 1A, B). While the number of IgE⁺ cells in cultures treated

with 0.125µM and 0.25µM of IC87114 were not significantly different to the IL-

4 and anti-CD40 cultures only, cultures treated with 0.5µM, 1µM and 2µM of

IC87114 had a significant reduction in the number of IgE⁺ cells (Figure 1A, B).

When quantifying the amounts of secreted IgE antibody in culture supernatants

we observed that the effects of IC87114 on secreted IgE were much more

drastic (Figure 1D). Indeed, cultures treated with 1µM and 2µM of IC87114 had

almost 80% of their secreted IgE reduced compared to the IL-4 and anti-CD40

cultures only. Furthermore, we can also observe a significant reduction of the

secreted IgE in cultures treated with 0.25µM of IC87114. Surprisingly, IC87114

had a negligible effect on the number of IgG1[+] cells, and we could only observe

a significant IgG1[+] cell reduction in cultures treated with 2µM of IC87114

(Figure 1A, C). Similar effects of IC87114 on CSR were also observed in naïve

B cell cultures stimulated with IL-4 and anti-CD40 (Figure 1F, G). These

observations are in striking contrast to those reported by the mouse studies and

suggests that PI3K p110δ regulation of CSR to IgE and IgG1 differs between

the mouse and the human system.

PI3K p110δ inhibition does not affect the class switch signals

downstream of the IL-4R and CD40

To identify the mechanism by which PI3K p110δ regulates CSR to IgE and IgG1

in our cultured tonsil B cells, we first examined the expression levels of

phosphorylated and thus activated, STAT6 and NF-κB. As it can be seen in

Figure 2A, 24 hours post culture with IL-4 and anti-CD40, the phosphorylation

levels of STAT6 and NF-κB were unaffected by the inhibition of p110δ PI3K with

2µM IC87114. However, previous mouse studies have reported that IC87114

regulates CSR to IgE and IgG1 by upregulating the expression of AID, εGLT

and γ1GLT. Therefore, to confirm that the class switch signals downstream of

the IL-4R and CD40 in human B cells are unaffected by the p110δ PI3K

inhibition we also evaluated the expression levels of AID, εGLT and γ1GLT on

day 5 of the cell culture. The RT-PCR data showed that the treatment of the

cell culture with 2µM IC87114 has no significant effect in the expression levels

of AID, εGLT and γ1GLT (Figure 2B). This is consistent with the unchanged

STAT6 and NFκB phosphorylation levels and suggests that unlike in the mouse

system, the effect of PI3K p110δ inhibition on CSR of human B cells to IgE and

IgG1 occurs independently of class switch signals downstream of the IL-4R and

CD40.

PI3K p110δ inhibition does not affect the expression levels of BCL6 in

human B cells.

Previously, it was reported that PI3K p110δ regulates CSR of mouse B cells

to IgE in a B cell intrinsic manner by modulating BCL6 expression[16]. BCL6 is

a transcription factor that is important for GC reaction but also is known to

compete with STAT6 at the εGLT binding site leading to the repression of its

transcription and that of CSR to IgE[10,21]. Therefore, in mice the reduction of

the BCL6 by the inhibition of p110δ can lead to enhanced CSR to IgE[16].

To test if this is also true in the human B cells, we evaluated the expression

levels of BCL6 by RT-PCR. As can be seen from Figure 3, we did not observe

any changes in the BCL6 expression levels between cell cultures that were

untreated and those treated with 2µM IC87114.

Overall, this data, and that from above, suggests that there are fundamental

differences in the mechanisms through which PI3K p110δ signalling regulates

CSR to IgE in human and mouse B cells.

286 **Inhibition of the PI3K p110δ activity reduces the proliferative capacity of**

287 **human B cells stimulated to undergo CSR to IgE.**

Cell division plays an important role in CSR[24,25]. For CSR to IgE to happen B

cells are required to divide a minimum of 5 times, whereas CSR to IgG1

requires only 2 cell divisions[24,25]. Mouse studies investigating the role of p110δ

PI3K in IgE production did not report any significant effect of the p110δ PI3K

inhibition on the IL-4 and anti-CD40 induced B cell proliferation[15,17,23]. To

determine if the p110δ PI3K inhibition reduces the proliferative capacity of the

cultured tonsil B cells, thus, reducing the CSR of human B cells to IgE, we first

labelled the isolated tonsil B cells with CFSE and analyzed cell division after 10

days of culture. The representative FACS histograms and cumulative data

(Figure 4A, B) show that the number of proliferating B cells in cell cultures

treated with 2µM IC87114 was reduced by half. Considering that the minimum

number (5) of cell divisions required for CSR to IgE to occur, we also

investigated the number of cell divisions by measuring the halving of the CFSE

intensity. We found that the number of B cells undergoing 4 cell divisions, or

more is significantly reduced when cell cultures were treated with 2µM IC87114

(Figure 4C, D). In contrast, the number of B cells undergoing 2 and 3 divisions,

which are decisive for CSR to IgG1 (22) was slightly, but not significantly,

reduced.

The data suggests that the modulation of the B cell proliferation is the main

mechanism through which PI3K p110δ signalling regulates CSR to IgE in

human B cells stimulated with IL-4 and anti-CD40.

Discussion

Mast cells lacking the p110δ activity have a considerable reduction in IgE-FcεRI

mediated degranulation and release of proinflammatory mediators[26,27].

Inhibition of the p110δ signalling in mouse models of asthma is also associated

with reduced Th2 cytokines as well as reduced lung cellular influx, including

tissue esonophilia, airway mucus production and airway hyper-

responsiveness[28–30]. These studies confirm PI3K p110δ as a potential target for

the treatment of allergic disease. Surprisingly, however, other mouse studies

have reported that PI3K p110δ signaling negatively regulates CSR to IgE, thus

its inhibition enhances IgE production[15–17]. In this study, we investigated the

effect of PI3K p110δ inhibition in cultures of human tonsil B cells stimulated to

undergo CSR to IgE. We find that the inhibition of the PI3K p110δ signaling in

these human B cell cultures reduces IgE production. This contradiction

between the human and mouse responses to PI3K p110δ inhibition highlights

the differences in their B cell regulatory systems of IgE production.

PI3K p110δ is activated downstream of various receptors on B cells and plays

critical roles in the development and function of B cells[18,31–33]. Several studies

have also implicated PI3K signalling in the regulation of CSR[15,17,23,34]. The first

line of evidence came from mouse B cells lacking PTEN, a phosphatase that

negatively regulates PI3K activity[34]. These PTEN-deficient mouse B cells,

which have enhanced PI3K signalling, fail to undergo CSR both *in vivo* and *in*

vitro. IC87114 reverses the effect of PTEN-deficiency on CSR, suggesting that

CSR is predominantly regulated by the PI3K p110δ isoform. Furthermore,

inhibition of PI3K p110δ was able to also enhance CSR in wild type mouse B

cells, suggesting that under normal circumstances PI3K p110δ negatively

regulates CSR. In the case of mouse IgE and IgG1, PI3K p110δ negatively

regulates their IL4 and anti-CD40 induced CSR *in vitro* by downregulating the

levels of AID as well as that of ϵGLT and γ1GLT, respectively[15,17]. Interestingly,

our data from the human B cell cultures were inconsistent with these

observations made in the mouse systems. IC87114 treatment leads to a

significant reduction in the number of IgE[+] cells in the IL-4 and anti-CD40

stimulated tonsil B cell cultures. The effect of IC87114 on IgG1[+] cells was not

as marked, and a significant reduction was observed only with the highest

concentration of 2μM IC87114. It has also been reported that upon

immunisation with T dependent antigens, both total and antigen specific IgE

titters are dramatically reduced in mice treated with IC87114 and those with B

cell deficient PI3K p110δ (CD19[Cre]p110δ[flox/flox])[15,35]. In contrast, the titers of

IgG1 and IgM are unaffected, suggesting that the PI3K p110δ selectively

regulates IgE production *in vivo.* Further investigations indicated that PI3K

p110δ regulates CSR to IgE in a B cell-intrinsic manner by modulating the

expression levels of BCL6 [16]. However, we did not observe an effect of IC87114

on the class switch signals downstream of IL4R and CD40 and the levels of

BCL6 expression were unchanged, suggesting that the mechanisms through

which PI3K signalling regulates CSR to IgE differs between the human and the

mouse systems.

PI3K p110δ signalling plays a key role in B cell proliferation as it regulates the

expression levels of different components of the cell cycle machinery[26,32,33]. Our

findings show that regulation of the B cell proliferative capacity is the key

mechanism through which PI3K p110δ signalling regulates CSR to IgE in our

IL4 and anti-CD40 stimulated B cell cultures. Indeed, IC87114 significantly

reduces the frequency of cells that have undergone 4 or more cell divisions.

This contradicts the previous observations made in mouse, which showed that

the proliferation of mouse B cells, stimulated with IL-4 and anti-CD40, is

unaffected by the PI3K p110δ inhibition[3,15,17,23]. Furthermore, the effect of the

IC87114 on the number of B cells undergoing 4 or more cell divisions may

explain why we see a more pronounced effect on CSR to IgE, which requires a

minimum of 5 divisions, compared to CSR to IgG1, which only requires two cell

divisions.

An important feature of both mouse and human IgE+ B cells is their tendency

to differentiate into the PC lineage[6,8,9,36]. Recent studies in mice revealed that

this process is promoted by the mIgE signalling independently of antigen and

involves the PI3K signalling pathway[4,5,37]. The drastic effects of IC87114 on the

amounts of secreted IgE suggests that in addition to inhibiting the CSR to

human IgE, the loss of PI3K p110δ signalling might also inhibit the generation

of IgE producing PCs. Our future studies will aim to elucidate the role of PI3K

p110δ signalling in the differentiation of human IgE+ B cells.

Overall, our data have demonstrated that the regulation of IgE production by

PI3K p110δ differs between the mouse and the human systems. In human B

cell cultures, the PI3K p110δ regulates IgE production by modulating the

proliferative capacity of the IL-4 and anti-CD40 stimulated B cells. In addition,

these data highlights the therapeutic potential of small molecule inhibitors of

PI3K p110δ in the treatment of allergic disease.

References

1. Gould HJ, Sutton BJ. IgE in allergy and asthma today. Vol. 8, Nature Reviews Immunology. 2008. p. 205–17.

2. Dullaers M, de Bruyne R, Ramadani F, Gould HJ, Gevaert P, Lambrecht BN. The who, where, and when of IgE in allergic airway disease. Vol. 129, Journal of Allergy and Clinical Immunology. Mosby Inc.; 2012. p. 635–45.

3. Newman R, Tolar P. Chronic calcium signaling in IgE+ B cells limits plasma cell differentiation and survival. Immunity. 2021;54:2756-2771.e10.

4. Yang Z, Robinson MJ, Chen X, Smith GA, Taunton J, Liu W, et al. Regulation of B cell fate by chronic activity of the IgE B cell receptor. Elife. 2016;5.

5. Haniuda K, Fukao S, Kodama T, Hasegawa H, Kitamura D. Autonomous membrane IgE signaling prevents IgE-memory formation. Nature Immunology. 2016;17:1109–17.

6. He JS, Meyer-Hermann M, Xiangying D, Zuan LY, Jones LA, Ramakrishna L, et al. The distinctive germinal center phase of IgE+ B lymphocytes limits their contribution to the classical memory response. Journal of Experimental Medicine. 2013;210:2755–71.

7. Gould HJ, Ramadani F. IgE responses in mouse and man and the persistence of IgE memory. Vol. 36, Trends in Immunology. Elsevier Ltd; 2015. p. 40–8.

8. Ramadani F, Bowen H, Upton N, Hobson PS, Chan YC, Chen JB, et al. Ontogeny of human IgE-expressing B cells and plasma cells. Allergy [Internet]. 2017 [cited 2022 Apr 26];72:66–76. Available from: https://pubmed.ncbi.nlm.nih.gov/27061189/

9. Yang Z, Sullivan BM, Allen CDC. Fluorescent In Vivo Detection Reveals that IgE+ B Cells Are Restrained by an Intrinsic Cell Fate Predisposition. Immunity. 2012;36:857–72.

414   10.   Geha RS, Jabara HH, Brodeur SR. The regulation of immunoglobulin E
415   class-switch recombination. Vol. 3, Nature Reviews Immunology. 2003.
416   p. 721–32.

417   11.   Delphin S, Stavnezer J. Characterization of an Interleukln 4 (IL-4)
418   Responsive Region in the Immunoglobulln Heavy Chain Germ!ine E
419   Promoter: Regulation by NF-IL-4, a C/EBP Family Member and NF-
420   KB/p50 [Internet]. Available from: http://rupress.org/jem/article-
421   pdf/181/1/181/1106000/181.pdf

422   12.   Takeda H, Tanaka T, Shi W, Matsumoto M, Minami M, Kashiwamura SI,
423   et al. Essential role of Stat6 in IL-4 signalling. Nature 1996 380:6575
424   [Internet]. 1996 [cited 2022 Apr 26];380:627–30. Available from:
425   https://www.nature.com/articles/380627a0

426   13.   Dedeoglu F, Horwitz B, Chaudhuri J, Alt FW, Geha RS. Induction of
427   activation-induced cytidine deaminase gene expression by IL-4 and
428   CD40 ligation is dependent on STAT6 and NFκB. International
429   Immunology. 2004;16:395–404.

430   14.   Hellman L. Regulation of IgE homeostasis, and the identification of
431   potential targets for therapeutic intervention. Biomedicine and
432   Pharmacotherapy. 2007;61:34–49.

433   15.   Zhang T ting, Okkenhaug K, Nashed BF, Puri KD, Knight ZA, Shokat KM,
434   et al. Genetic or pharmaceutical blockade of p110δ phosphoinositide 3-
435   kinase enhances IgE production. Journal of Allergy and Clinical
436   Immunology. 2008;122.

437   16.   Zhang T ting, Makondo KJ, Marshall AJ. p110δ Phosphoinositide 3-
438   Kinase Represses IgE Switch by Potentiating BCL6 Expression. The
439   Journal of Immunology. 2012;188:3700–8.

440   17.   Doi T, Obayashi K, Kadowaki T, Fujii H, Koyasu S. PI3K is a negative
441   regulator of IgE production. International Immunology. 2008;20:499–508.

442   18.   Okkenhaug K, Vanhaesebroeck B. PI3K in lymphocyte development,
443   differentiation and activation. Vol. 3, Nature Reviews Immunology.
444   European Association for Cardio-Thoracic Surgery; 2003. p. 317–30.

445   19.   Soond DR, Bjørgo E, Moltu K, Dale VQ, Patton DT, Torgersen KM, et al.
446   PI3K p110delta regulates T-cell cytokine production during primary and
447   secondary immune responses in mice and humans. Blood [Internet].

2010 [cited 2022 Apr 26];115:2203–13. Available from:
https://pubmed.ncbi.nlm.nih.gov/20081091/

20. Nashed BF, Zhang T, Al-Alwan M, Srinivasan G, Halayko AJ, Okkenhaug
K, et al. Role of the phosphoinositide 3-kinase p110delta in generation of
type 2 cytokine responses and allergic airway inflammation. Eur J
Immunol [Internet]. 2007 [cited 2022 Apr 26];37:416–24. Available from:
https://pubmed.ncbi.nlm.nih.gov/17236236/

21. Harris MB, Chang CC, Berton MT, Danial NN, Zhang J, Kuehner D, et al.
Transcriptional Repression of Stat6-Dependent Interleukin-4-Induced
Genes by BCL-6: Specific Regulation of Iε Transcription and
Immunoglobulin E Switching. Vol. 19, MOLECULAR AND CELLULAR
BIOLOGY. 1999.

22. Ramadani F, Upton N, Hobson P, Chan YC, Mzinza D, Bowen H, et al.
Intrinsic properties of germinal center-derived B cells promote their
enhanced class switching to IgE. Allergy: European Journal of Allergy
and Clinical Immunology. 2015;70:1269–77.

23. Chiu H, Mallya S, Nguyen P, Mai A, Jackson L v., Winkler DG, et al. The
Selective phosphoinoside-3-kinase p110δ inhibitor IPI-3063 potently
suppresses b cell survival, proliferation, and differentiation. Frontiers in
Immunology. 2017;8.

24. McCall MN, Hodgkin PD. Switch recombination and germ-line
transcription are division-regulated events in B lymphocytes. Biochimica
et Biophysica Acta (BBA) - Gene Structure and Expression.
1999;1447:43–50.

25. Hasbold J, Lyons AB, Kehry MR, Hodgkin PD. Cell division number
regulates IgG1 and IgE switching of B cells following stimulation by CD40
ligand and IL-4. European Journal of Immunology. 1998;28:1040–51.

26. Ali K, Bilancio A, Thomas M, Pearce W, Gilfillan AM, Tkaczyk C, et al.
Essential role for the p110delta phosphoinositide 3-kinase in the allergic
response. Nature [Internet]. 2004 [cited 2022 Apr 26];431:1007–11.
Available from: https://pubmed.ncbi.nlm.nih.gov/15496927/

27. Ali K, Camps M, Pearce WP, Ji H, Rückle T, Kuehn N, et al. Isoform-
specific functions of phosphoinositide 3-kinases: p110 delta but not p110
gamma promotes optimal allergic responses in vivo. J Immunol [Internet].

2008   [cited   2022   Apr   26];180:2538–44.   Available   from:
https://pubmed.ncbi.nlm.nih.gov/18250464/

28.   Kim JS, Jeong JS, Kwon SH, Kim SR, Lee YC. Roles of PI3K pan-
inhibitors and PI3K-δ inhibitors in allergic lung inflammation: a systematic
review and meta-analysis. Scientific Reports. 2020;10.

29.   Park SJ, Min KH, Lee YC. Phosphoinositide 3-kinase δ inhibitor as a
novel therapeutic agent in asthma. Vol. 13, Respirology. 2008. p. 764–
71.

30.   Lee KS, Lee HK, Hayflick JS, Lee YC, Puri KD. Inhibition of
phosphoinositide 3-kinase δ attenuates allergic airway inflammation and
hyperresponsiveness in murine asthma model. The FASEB Journal.
2006;20:455–65.

31.   Turner M, Hodson DJ. The role of PI3K signalling in the b cell response
to antigen. In: Advances in Experimental Medicine and Biology. 2009. p.
43–53.

32.   Clayton E, Bardi G, Bell SE, Chantry D, Peter Downes C, Gray A, et al.
A crucial role for the p110δ subunit of phosphatidylinositol 3-kinase in B
cell development and activation. Journal of Experimental Medicine.
2002;196:753–63.

33.   Okkenhaug K, Bilancio A, Farjot G, Priddle H, Sancho S, Peskett E, et al.
Impaired B and T cell antigen receptor signaling in p110delta PI 3-kinase
mutant mice. Science [Internet]. 2002 [cited 2022 Apr 26];297:1031–4.
Available from: https://pubmed.ncbi.nlm.nih.gov/12130661/

34.   Omori SA, Cato MH, Anzelon-Mills A, Puri KD, Shapiro-Shelef M, Calame
K, et al. Regulation of Class-Switch Recombination and Plasma Cell
Differentiation by Phosphatidylinositol 3-Kinase Signaling. Immunity.
2006;25:545–57.

35.   Rolf J, Bell SE, Kovesdi D, Janas ML, Soond DR, Webb LMC, et al.
Phosphoinositide 3-Kinase Activity in T Cells Regulates the Magnitude of
the Germinal Center Reaction. The Journal of Immunology [Internet].
2010   [cited   2022   Apr   26];185:4042–52.   Available   from:
https://www.jimmunol.org/content/185/7/4042

36.   Ramadani F, Bowen H, Gould HJ, Fear DJ. Transcriptional analysis of
515   the human IgE-expressing plasma cell differentiation pathway. Frontiers
516   in Immunology. 2019;10:402.

517   37.   Newman R, Tolar P. Chronic calcium signaling in IgE+ B cells limits
518   plasma cell differentiation and survival. Immunity. 2021;54:2756-
519   2771.e10.

 Figures

 Figure 1. PI3K p110δ inhibition reduces class switching of human B cells to

 IgE.

 (A) IL4 and anti-CD40 stimulated cells were harvested from the day 10 cultures

 containing different concentrations of IC87114 and intracellularly stained for IgE

 and IgG1. The percentages of total IgE$^+$ cells (**B**) and IgG1$^+$ cells (**C**) relative to

 the IL-4 and anti-CD40 only cultures. Secreted IgE (**D**) and IgG1(**E**) analyzed

 by ELISA. Data represent the mean +/- standard error of mean (SEM) and

 show the amounts of secreted IgE and IgG1 [ng/mL] made relative to the IL-4

 and anti-CD40 only cultures. The percentages of total IgE$^+$ cells (**F**) and IgG1$^+$

 cells (**G**) from naïve B cell cultures made relative to the IL-4 and anti-CD40 only

 cultures.

 Figure 2. IC87114 has no effect on the class switch signalling downstream of

 IL-4R and CD40.

 (**A**) After 24h of IL-4 and anti-CD40 stimulation, B cells were fixed,

 permeabilized and subsequently stained with anti-STAT6 phospho (Tyr641)

 and anti-NFκB p65 (Ser536). The data show the fold change in median

 fluorescence intensity (MFI) relative to IL-4 and anti-CD40 only cultures. **(B)**

 The expressions of AID, εGLT and γ1GLT were quantified with qRT-PCR from

 RNA isolated from day 5 of the tonsillar B cell culture. Gene expression was

 normalised with 18s rRNA.

Figure 3. PI3K p110δ inhibition does not affect the expression levels of BCL6

in IL-4 and anti-CD40 stimulated human B cells.

The RNA was isolated from cell pellets collected on day 5 of the B cell culture

and used to generate cDNA, which was used to quantify the expression levels

of BCL6 by RT-PCR. Values on the graphs are expressed in relation to

endogenous control, 18s x 10,000. Data represent means and SEM from 7

independent experiments. NOTE: I used RNA from 3 other experiments by Dr.

Ramadani that used the same culture conditions.

Figure 4. IC87114 inhibits the proliferative capacity of the IL-4 and anti-CD40

stimulated B cells.

Isolated tonsil B cells were stained with CFSE and cultured for 10 days with IL-

4 and anti-CD40 alone or with 2μM IC87114. **(A)** The FACS histograms show

the percentage of proliferating B cells after 10 days of culture, whereas **(B)**

shows the fold change in proliferation relative to the IL-4 and anti-CD40 only

culture. The statistical comparisons were performed using the paired t-test. ****

$P < 0.0001$ (n= 5). (**C**) The FACS plots show the number of divisions and the

percentage of cells at each division that the IL-4 and anti-CD40 stimulated B

cells, with or without IC87114, were undergoing. Based on the CFSE dilution,

six numbers of cell divisions (D1-D6+) were identified and gated. (D) Graph

shows the frequency of B cells at divisions 1 to 6+ (D1 to D6+). The data

represent the mean and SEM

3

BCL6

□ IL4 + anti-CD40

■ IL4 + anti-CD40
+ 2μM IC87114

4A

IL4 + anti-CD40

Proliferating
44.8

IL4 + anti-CD40
+ 2µM IC87114

Proliferating
21.2

Counts

CSFE

4B

Cell proliferation

% fold change relative
to IL4 + aantiCD40

****

□ IL4 + anti-CD40
■ IL4 + anti-CD40
+ 2µM IC87114

4C

IL4 + anti-CD40

D6+	D5	D4	D3	D2	D1
10.4	7.77	4.97	4.64	4.1	3.5

IL4 + anti-CD40
+ 2µM IC87114

D6+	D5	D4	D3	D2	D1
1.89	3.88	3.5	3.52	4.1	4.2

CSFE

4D

Effect of IC87114 on cell division

% fold change relative
to IL4 + anti-CD40

*
****
***
**
*
**

IL4 +
anti-CD40

IL4+ anti-CD40
+2µm IC87114

■ D1
■ D2
■ D3
■ D4
■ D5
■ D6+