Deciphering the Inter-domain Decoupling in the Gram-negative Bacterial Membrane Insertase

Adithya Polasa and Mahmoud Moradi*

Department of Chemistry and Biochemistry, University of Arkansas, Fayetteville, AR 72701

E-mail: moradi@uark.edu

Abstract

YidC is a protein found in membranes that plays an important role in the process of inserting newly generated proteins into lipid membranes. The SecYEG-dependent complex is responsible for inserting proteins into the lipid bilayer, and this process is facilitated by YidC. In addition, YidC acts as a chaperone during the folding processes of proteins. Multiple investigations have conclusively shown that the gram-positive bacterium YidC has SecY-independent insertion mechanisms. Through the use of microsecond level all-atom molecular dynamics simulations, we have carried out the first in-depth investigation of the YidC protein originating from gram-negative bacteria. This research sheds light on the significance of several structural areas related to YidC at an atomic level by utilizing equilibrium molecular dynamics (MD) simulations. In this research, multiple models of YidC inside the lipid bilayer were constructed in order to achieve a deeper understanding of the critical role of the C2 loop and the extra periplasmic domain present in gram-negative YidC. According to the results of our research, the C2 loop is responsible for the overall stabilization of the protein, most notably in the transmembrane region, and it also has an allosteric influence on the periplasmic domain. We have found critical interactions that contribute to the stability of the protein as well as its functional aspect. Finally, our study provides a hypothetical SecY-independent insertion mechanism for gram-negative bacterial YidC.

Keywords: YidC, Molecular Dynamics Simulations, Membrane Insertion, SecY-independent pathway, Insertase

Introduction

Membrane proteins participate in fundamentally crucial biological processes such as signaling, transcriptional regulation, ion and macro-molecule transport, proteolysis, motility, metabolism, energy creation, and energy transfer. Specialized cellular machinery enables proper membrane protein folding and insertion into the lipid bilayer. A group of membrane proteins may be inserted through the Sec apparatus in bacteria. YidC is one of the proteins working with Sec in order to introduce client proteins into the membrane[1–7].

YidC is a member of the Oxa/Alb3/YidC family of insertases found in mitochondria, chloroplasts, and bacteria[8–14]. YidC catalyzes the transmembrane insertion of freshly produced membrane proteins in the absence of an energy supply domain, such as an ATPase[15]. It also plays a vital role in insertion in the insertion and positioning of membrane proteins in bacteria[16–18]. Insertase proteins, such as YidC, have been exhaustively investigated to determine their importance for the insertion of proteins into membranes[19]. They are versatile proteins that, in conjunction with the SecYEG pathway[1–7], act to insert peptides into the membrane through the Signal Recognition Particle (SRP) mechanism. YidC may fold and insert polypeptides independently of the SecY pathway[2,20–27]. YidC is a crucial protein for the insertion of small phage coat proteins like Pf3 coat and M13 in a SecY-independent pathway[20,28–33].

A few experimental studies have explored the role of YidC in various microbial organisms. The genome of most gram-positive microscopic organisms encode two YidC proteins: YidC1 and YidC2[34,35]. Although YidC typically exists as a dimer or tetramer[36] under physiological conditions, it is discovered that YidC can also exist as a monomer in lipid bilayers[15,37]. The gram-negative YidC protein possesses an additional transmembrane (TM) segment at the N-terminus and a large periplasmic domain (PD)[38,39]. Although the PD region and the additional N-terminus TM segment are not required for YidC activity, PD region inteacts with SecY machinery and helps creating stable complex.[38]. The area with the C-terminal

five TM segments are vital for the membrane insertase activity of gram-negative YidC[40]. In both gram-negative and gram-positive strains of bacteria, the protein is firmly anchored in the lipid bilayer by interfacial aromatic residues, a cytoplasmic salt-bridge group, and a periplasmic helix enhanced with aromatic residues. The highly conserved arginine residue (R366) in the hydrophilic groove was found in the same locations as in gram-positive (R72), implying that the arginine residue is as important for gram-negative function as it is for gram-positive function[38]. A group of aromatic residues around R72/R366 may bind with incoming peptide during insertion into the lipid bilayer[19,34,35,40–43]. The C–terminus of monomeric YidC cooperates with the ribosomes, and the short interhelical loops come into contact with the ribosomal proteins[44].

Fig. 1. The cartoon representation of YidC (PDB:6AL2) (A) includes periplasmic domain (PD), transmembrane (TM) domain, C1, and C2 loops on the cytoplasmic side. (B) The cartoon representation of YidC's individual TM helices TM1 (blue), TM2 (Red), TM3 (purple), TM4 (orange), and TM5 (black).

Both in gram-negative and gram-positive bacteria, YidC is believed to promote membrane insertion simply by binding nascent chains and promoting their insertion into the lipid bilayer using cytoplasmic loop interactions, hydrophobic force, and interactions in the groove[15,19,45,46]. The hydrophilic groove inside the membrane core of YidC increases the rate of accepting the hydrophilic moieties of a substrate into the membrane[47,48]. During the process of its independent insertion mechanism, YidC also goes through a number of confor-

mational changes, including widening of the TM region and hydration and dehydration of the hydrophilic groove. Additionally, a broad range of interactions with the incoming protein are engaged at each step of the insertion process, for example the salt bridge interaction with R72 aromatic residue[19].

In the past, gram-positive bacterial YidC insertase has been widely researched[15,19,45,46]. The significance of the extra PD region in gram-negative YidC is not fully understood, yet. We don't know if the PD area of YidC affects protein stability and function, nor do we know if the cytoplasmic loops serve the same purpose in gram-positive and gram-negative bacterial YidC. Here, gram-negative YidC's structure was investigated using microsecond level molecular dynamics simulations. We examined the regional and global conformational changes in YidC brought on by the loss of the periplasmic domain and the cytoplasmic loop.

Methods

From the Protein Data Bank, the crystal structure of the gram-negative bacterial YidC (PDB:6al2[49]) was downloaded. The CHARMM36m[50] force field[51], together with the NAMD 2.14[52] software package were used for all molecular dynamics simulations. The protein was solvated in TIP3P[53] water. Using the membrane builder on CHARMM-GUI[54], YidC was introduced into the lipid bilayer, solvated, and ionized. In these MD investigations, YidC was placed in a lipid bilayer of 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoethanolamine (POPC) lipids. A 90 Å × 90 Å membrane layer surface was constructed along the XY plane. The protein-lipid assembly was solvated in TIP3[53] water with 18 Å thick layers of water on top and bottom. To neutralize the system, 0.15 M of Na^+ and Cl^- ions were added to the solution with a slight modification in the number of ions to neutralize the system. There were about ≈143000 atoms in the final solvated system. Utilizing the conjugate gradient technique [55], each system was energy minimized before the equilibrium simulation. The systems were then gradually relaxed using constrained MD simulations in accordance with the standard CHARMM-GUI[54] procedure. In the NPT ensemble at 310 K, 1 μs of equilibrium MD simulations were performed under periodic boundary conditions for each system. In the simulations, a Langevin integrator with a damping coefficient of γ =0.5 ps^{-1} and 1 atm pressure was maintained using the Nose-Hoover Langevin piston method[56,57 18,58–67].

We are interested in the significance of the cytoplasmic C2 loop and extracellular periplasmic domain (PD). It has been shown in the past that the C2 loop is crucial to YidC's overall conformation and function in gram-positive bacteria[19,68]. We are interested in learning more about the functions of the PD (Fig. 1A), which is absent in gram-positive bacterial YidC and present in gram-negative bacterial YidC. Accordingly, we created four systems: YidC with PD and C2 loop (YidC), YidC without C2 loop (YidC ΔC2), YidC without PD region (YidC ΔPD), and YidC without PD region and C2 loop (YidC ΔPD ΔC2).

The preliminary simulations were run on TACC Stampede supercomputer. Subsequently,

the production run for each model was increased to 400 ns with a timestep of 2.5 fs on Anton2[58]. Every 240 picoseconds, conformations were gathered on Anton2. Initial processing of the Anton2 simulation trajectories was carried out on Kollman[58]. Later, the simulations were extended to 1 microsecond on TACC Stampede and one additional set of 1 μs simulation was performed for all systems.

All trajectories were visualized and examined using the VMD software[69]. A VMD plugin was used to analyze salt bridge interactions by measuring the distance between the oxygen atoms of acidic residues and the nitrogen atoms of basic residues with a cut-off distance of 4 Å. The interhelical angles were determined as the angle between the third main axes of the respective helices[19,60,67,70,71]. The selection of the TM helices and other sub-domains are as indicated: TM1 (355-388); TM2 (423-442); TM3 (466-479); TM4 (497-508); TM5 (511-528); C1 area (380-420); C2 loop (480-492); and PD (49-326) (Fig. 1). To analyze the water inside the groove region, we counted the number of water molecules within 5 Å of R366. Principle component analysis (PCA) was performed for each trajectory using PRODY[19,60,72], taking only protein C_α atoms into account.

More quantitative data on the coordinated movements of the C_α atoms was obtained using Dynamic Network Analysis (DNA) of the associated motions of the protein[58,73]. The correlation coefficient for the motion of each C_α atom with respect to the other C_α atoms was determined using MD-TASK[74], a software package of MD analysis tools. For each of the TM regions in all the simulated trajectories, a correlation matrix M was created.

To quantify the differences in correlation between a system and some reference, a difference matrix Δ was calculated,

$$\Delta = |M_i - M_{Ref.}| \tag{1}$$

where M_i is the correlation matrix of interest, and M_{Ref} is the correlation matrix of a reference conformation. In this work, the difference between a TM region in complete YidC

structure conformation and other YidC structures was our point of interest. For this reason, the TM region in the wild type YidC simulations were compared with TM region in the other YidC simulation system described above.

Results and Discussion

The Overall Protein Conformation is Stabilized by the Presence of the C2 loop and the PD Domain.

Fig. 2. Analysis of YidC's structural stability in presence and absence of PD and C2 loop. (A–D) The root mean square deviation of the YidC in different systems. Based on RMSD data, we have observed that YidC fluctuates more in the system with the C2 loop removed compare to systems with C2 loop. The simulations for each system were run twice, and the dashed lines in the graphs reflect the second run of those simulations.

The root mean square deviation (RMSD) of the protein was first calculated in order to assess the stability during equilibrium simulations. The RMSD of the YidC TM region was computed, and the results are shown as a function of the simulation run-time (Fig. 2).

According to the C_α RMSD of the four different systems, the presence of the PD and C2 loop helps to stabilize the protein when it is in its native state (Fig. 2A). The system without the C2 loop is, beyond a shadow of a doubt, less stable than the system with the loop (Fig. 2B & D). The combined effect of removing the PD and the C2 loop has a higher impact on RMSD (Fig. 2D), which suggests that the PD could be responsible for preserving the stability of the protein. However, the effect of just removing the C2 loop (Fig. 2B) on protein RMSD is slightly higher than the native system (Fig. S1), but not as high as the system without PD and the C2 loop (Fig. 2D). We believe in between the C2 loop and the PD region, the C2 loop is more important for the stability of the protein compared to the PD region, as you can see in system YidC ΔPD (Fig. 2C) the RMSD of the TM region is very similar to the wild type YidC structure (Fig. 2A). Therefore, we have arrived at the conclusion that the impact of eliminating the C2 loop on gram-negative bacterial YidC is far more substantial than the effect of removing just PD regions (Fig. S1).

Fig. 3. Projections of the principal components (PC's) 1 and 2. (A–H) PCA findings of PC1 vs PC2 for YidC systems from set 1 simulations are shown in the top row, while the results of set 2 simulations are displayed in the bottom row. The gradation of colors in the image denotes a timeline, with lighter shades reflecting earlier points in the simulation and darker hues denoting later points. Only the PCA analysis of the TM area that is present in all systems is shown here for the sake of consistency. For easier comparison, the green line in the plot is to show the difference in projections of PC's.

The principal component analysis (PCA) was used in order to determine the most impor-

tant distinctions between the systems could be determined. The projections onto principle components (PCs) 1 and 2 allowed for a straightforward differentiation between the YidC wild type and the other systems (Fig. 3). In this particular analysis, only the YidC C_α atoms that are located in the TM area are taken into account. PC1 contributed 27.5% of the total variance, while PC2 contributed 19.1% of the total variance. It was anticipated that the structural analysis of YidC ΔC2 and YidC ΔC2 ΔPD models would contradict that of YidC and YidC ΔPD in PC1 and PC2 (Fig. 3). This contradiction is rational, considering the large conformational discrepancies that were found earlier in the RMSD analysis (Fig. S1). In general, the most important thing that came out of the principal components analysis was the realization that the behavior of the YidC ΔC2 ΔPD (Fig. 3 D & H) protein system was quite different from the other systems (Fig. 3). The results of this study lend credence to the theory that the C2 loop play a significant part in the conformational dynamics of YidC.

Fig. 4. Inter-helical angles between transmembrane helices of YidC. (A–D) The inter-helical angle between the transmembrane helix 1 and helix 3 of the protein. (E–H) The overall inter-helical angle between helix 2 region and helix 4 of the protein.

Previous studies have revealed that the YidC transmembrane (TM) region is crucial for membrane protein insertion mechanism[19,75,76]. In order to examine the impact of deleting the PD and C2 loop on the TM helices, the helical angle between each pair of TMs was

measured in this work (Fig. 4). When compared to the wild type, we found that the local shape of the TM helices was altered in all of the other systems. Similar to the analysis above, the local conformation was more affected by the removal of both the PD and the C2 loop (Fig. 4D & H). However, when just the PD is removed, we do see some changes in the helical angle (Fig. 4C & G), despite the fact that the impact is not as large as what is seen in systems that do not have a C2 loop (Fig. 4B, F, D, & H). This demonstrates the critical role that the C2 loop plays in maintaining the structural stability of transmembrane region of YidC. We do observe a similar trend in other transmembrane helices combinations (Fig. S2).

Fig. 5. The DNA analysis revealed differences in correlation between the YidC set 1 control system and YidC ΔC2 ΔPD system examined. The theoretical maximum for correlation difference is 2, but the observed maximum was less than 1. (A–B) Differences in correlation are shown as a red gradient, with darker red indicating a larger difference.

Furthermore, we used dynamic network analysis (DNA), which finds the linear connection between various residue pairs, in order to conduct a comprehensive study on the allosteric interactions of the C2 loop and the PD region with the various protein domains. The correlation coefficient of each residue pair is shown in Figure 5. This coefficient was calculated from the trajectory with the C2 loop and PD region, then subtracted from the same quantity that was calculated from the trajectory without the loop and PD region systems, and the

result was reported as its absolute value. The amount that is presented for each pair of residues measures the size of the difference in the correlation behavior of the two residues that is brought on by the presence of the C2 loop and the PD domain in the TM region. The presence or absence of the C2 loop has been demonstrated to create significant variations in the correlations between YidC's distinct domains. The inter-domain correlations, notably between the TM/C1 loop region, vary greatly between the YidC and other systems (Fig. S3). This difference in the cross-correlation is more pronounced, especially between TM1 and TM4 (Fig. 5). Based on this, we believe that the C2 loop does play a crucial role in the YidC conformational dynamics and that the conformational dynamics of the TM region are affected in its absence. The DNA findings (Fig. S3) are consistent with the early evidence for global and regional structural alterations. Overall, the results show that the C2 loop affects the behavior of the functionally essential areas of gram-negative YidC.

C2 loop and the Periplasmic Domain Allosterically Influence YidC's Other Functionally Important Regions

Fig. 6. Analyses of the water inside the YidC groove. (A–D) The number of water molecules that are located within 5 Å of the R366 residue inside the hydrophilic groove (HG) region of YidC in each and every system. (E) The graphical illustration of the residue R366, which may be found in the central part of the hydrophilic groove (HG) of YidC

YidC's U-shaped hydrophilic groove, which is exposed on the cytoplasmic side of the membrane bilayer, is essential for the insertion process[47,48]. The membrane proteins enter

the YidC groove through the cytoplasmic side of the membrane bilayer during the insertion process. The YidC groove area is filled with water to provide a smooth sliding motion for the entering protein. As a membrane protein advances through the insertion processes, the groove water molecules are expelled from the TM groove. These two variables produce a change in the region's hydrophobicity, making it more vulnerable to membrane insertion[19,41,77].

To analyze the water content of the groove inside the TM region, the amount of water molecules within the groove region of the YidC protein was determined and plotted against simulation time. The lack of the C2 loop had a significant impact on the amount of water that was present inside the groove area (Fig. 6B & D). Because hydrophilic contacts initially serve to keep membrane proteins in place inside the groove whilst the insertion process is being carried out[19]. This provides us with strong evidence that the C2 loop not only contributes to the conformational dynamics of the protein but also plays a significant role in the function of the protein. The water quantity is significantly reduced in system YidC ΔC2 ΔPD loop compared to system YidC. Our findings lead us to infer that the removal of just PD does have a marginal impact on the conformational dynamics of YidC. However, when this modification is coupled with the removal of the C2 loop, the effect is significantly amplified, which could ultimately effect the insertion process.

We also found an intradomain hydrogen bond in the TM region between Y516 and G429 that is only stable in the wild type system compared to other systems (Fig. 7). Especially in the YidC ΔC2 ΔPD system, this bond is completely broken. We think that this hydrogen bond is unstable in the system YidC ΔC2 ΔPD because fluctuation of the TM region caused due to absence of C2 loop (Fig 7B & D). These results clearly show that there is a link between the functionally important C2 loop and the PD region on the TM side of YidC. We also think this hydrogen bond (Fig 7) could be important to the insertion process. As the incoming membrane protein moves along the groove and toward the periplasmic side, it breaks this link (Fig 7), which causes widening of the TM region and leads the water to

Fig. 7. Interaction analysis of a hydrogen bond between Y516 and G429 (YidC), which is situated inside the groove area. (A–D) Analysis of the distance between the hydrogen bonds formed by Y516 and G429. (E) The graphical representation of the residues that participate in the interaction involving hydrogen bonds

leave the hydrophilic groove, resulting in a hydrophobic shift and increasing the likelihood of membrane insertion. To this point, we have shown that the C2 loop and PD directly alter the structural behavior of YidC, while the PD region also slightly affects the conformational dynamics of the protein. However, the absence of both the PD and the C2 loop has an allosteric influence on the behavior of the YidC conformational dynamics, although the effect of the C2 loop absence is substantially larger than the PD absence. This leads us to the conclusion that the C2 loop is much more important to the structure and function of YidC than the PD region, which is in support of previous research[38,39]. On the other hand, the removal of just PD did affect the conformational dynamics of the TM region. To determine what caused this effect, we analyzed interactions between the PD and the TM region that play a significant role in the stability of the protein.

Inter-domain Amino Acids Interactions Play a Key Role in the Stabilization of the Transmembrane Domain

Previous experimental findings lead researchers to the hypothesis that the interactions between the PD region of YidC and the TM region of YidC are crucial for maintaining the

Table 1: Occupancy (%) of Inter-domain H-Bonds between PD and TM region

System	Y259-M441 (%)		A214-Y437 (%)		T327-E312 (%)	
	set1	set2	set1	set2	set1	set2
YidC	40	40	66	42	81	43
YidC ΔC2	0	0	0	0	0	0
YidC ΔPD	NA	NA	NA	NA	NA	NA
YidC ΔC2 ΔPD	NA	NA	NA	NA	NA	NA

protein's stable state in the membrane[38]. Through this research, we were able to identify critical hydrogen bonds and salt-bridle interactions that take place between the PD region and the TM region. It is interesting to note that the hydrogen bonds that had been formed between the PD region and the TM region were only observed in wild type YidC system and entirely disrupted in the YidC ΔC2 system. However, there is no direct interaction between the PD area and the C2 loop region which does have an effect on stability of TM region. We also identified two salt bridge interaction between PD and TM which are contributing to stability of TM region. The salt bridge between D315 and K345 is stably formed in both systems of YidC even without C2 loop this salt bridge is unaffected (Fig. 8A & B). However the salt bridge between K232 and D329 (Fig. 9) is disrupted in YidC ΔC2 (Fig. 9B).

We believe that the removal of the C2 loop has caused an effect on the hydrogen bond (Fig. 7) in the TM core region, causing instability of the TM region and ultimately causing disruption of hydrogen bonds and salt bridge interactions between the PD and the TM region. Based on the analysis presented above, we postulate that the the hydrogen bond (Fig. 7) in the protein's groove region is essential for preserving the structural stability of the protein. The C2 loop is crucial for this hydrogen bond to remain stable in the structure. Although the YidC PD region does have some influence on the protein's structure, its contribution to the protein's overall function is noticeably less substantial than that of the C2 loop. Even without the PD region, it is still possible to have a normal Sec-independent insertase mechanism; however, the absence of the C2 loop may have a detrimental influence on the protein function.

Fig. 8. Salt–bridge interaction between D315 and K345 (YidC), which is located between the PD and TM regions. Graphical illustration of the important salt-bridge interactions that take place between the PD region and the TM area (right). (A–B) Distance analysis between D315–K345 salt-bridge and we also reported the occupancy of salt bridge interaction.

Proposed Independent Insertion Mechanism of Gram-negative Bacterial YidC

According to the findings and earlier hypotheses, gram-negative bacterial YidC must likewise go through significant conformational changes during the SecY-independent insertion procedure, much like gram-positive bacterial YidC. During the SecY-independent insertion process, the entering membrane protein would first make contact with the cytoplasmic loops and then squeak into the hydrophilic groove, where it would join forces with R366 to create a salt bridge. Incoming protein must be moved into YidC's hydrophilic groove by the YidC loops on the cytoplasmic side of the bilayer. The salt bridge between the incoming protein

Fig. 9. Salt–bridge formed between D329 and K232 (YidC), which is located between the PD and TM regions. The cartoon representation of the salt-bridge interactions that take place between the PD region and the TM area (right). (A–B) Distance analysis between D329–K232 salt-bridge and we also reported the occupancy of salt-bridge interaction.

and R366 of YidC also contributes to the passage of the protein towards the periplasmic side, stabilizing its position within the groove. As the incoming protein moves through the groove and approaches the periplasmic side, it breaks the hydrogen bond between Y516 and G429, causing widening of the TM region that results in a hydrophobic shift through dehydration of the groove. The protein will finally make contact with lipid tails. After that, with the aid of the proton motive force and the membrane's hydrophobic interaction, the protein moves in the direction of the membrane. Then, via the groove, the protein enters the membrane.

Conclusions

We have come to the conclusion that the C2 cytoplasmic loop of YidC could be an important part of the protein. This could be achieved, for example, by the C2 cytoplasmic loop of YidC helping to stabilize the protein structure via its indirect effects on interactions in the transmembrane core region, as well as its indirect effects on other periplasmic domains, in particular on interactions between the PD and the TM region. It also indicates that the existence of the C2 loop has an influence on the functional features of YidC, such as the hydration of the groove. In order to further understand the C2 loop's role in the sec-independent insertion process of small single-spanning membrane proteins like the pf3 coat protein, whose interactions with the cytoplasmic region are thought to be essential, more study is required. In the context of molecular dynamics simulations, the results of our research show that C2 loops may be required for the structural stability and SecY-independent function of the gram-negative YidC membrane protein dynamics.

Acknowledgement

This research is supported by the National Science Foundation grant CHE 1945465, the National Institutes of Health grant R15GM139140, and the Arkansas Biosciences Institute. This work used the Extreme Science and Engineering Discovery Environment (allocation MCB150129), which is supported by National Science Foundation grant number ACI-1548562. Anton 2 computer time was provided by the Pittsburgh Supercomputing Center (PSC) through Grant R01GM116961 from the National Institutes of Health. The Anton 2 machine at PSC was generously made available by D.E. Shaw Research.

Supporting Information Available

Movies S1–S2 and Figures S1-S3 in Supporting Information provide additional analysis based on our MD as discussed in the manuscript.

References

(1) Nagamori, S.; Smirnova, I. N.; Kaback, H. R. Role of YidC in folding of polytopic membrane proteins. *Journal of Cell Biology* **2004**, *165*, 53–62.

(2) Lewis, N. E.; Brady, L. J. Breaking the bacterial protein targeting and translocation model: Oral organisms as a case in point. *Molecular Oral Microbiology* **2015**, *30*, 186–197.

(3) Serdiuk, T.; Balasubramaniam, D.; Sugihara, J.; Mari, S. A.; Kaback, H. R.; Müller, D. J. YidC assists the stepwise and stochastic folding of membrane proteins. *Nature Chemical Biology* **2016**, *12*, 911–917.

(4) Kol, S.; Turrell, B. R.; De Keyzer, J.; Van Der Laan, M.; Nouwen, N.; Driessen, A. J. YidC-mediated membrane insertion of assembly mutants of subunit c of the F1F0 ATPase. *Journal of Biological Chemistry* **2006**, *281*, 29762–29768.

(5) Van der Laan, M.; Urbanus, M. L.; Ten Hagen-Jongman, C. M.; Nouwen, N.; Oudega, B.; Harms, N.; Driessen, A. J.; Luirink, J. A conserved function of YidC in the biogenesis of respiratory chain complexes. *Proceedings of the National Academy of Sciences of the United States of America* **2003**, *100*, 5801–5806.

(6) Yi, L.; Dalbey, R. E. Oxa1/Alb3/YidC system for insertion of membrane proteins in mitochondria, chloroplasts and bacteria. *Molecular Membrane Biology* **2005**, *22*, 101–111.

(7) Van Bloois, E.; Jan Haan, G.; De Gier, J. W.; Oudega, B.; Luirink, J. F1F0 ATP synthase subunit c is targeted by the SRP to YidC in the E. coli inner membrane. *FEBS Letters* **2004**, *576*, 97–100.

(8) Dalbey, R. E.; Kuhn, A. Membrane Insertases Are Present in All Three Domains of Life. *Structure* **2015**, *23*, 1559–1560.

(9) McDowell, M. A.; Heimes, M.; Sinning, I. Structural and molecular mechanisms for membrane protein biogenesis by the Oxa1 superfamily. *Nature Structural & Molecular Biology* **2021**, *28*, 234–239.

(10) Jiang, F.; Chen, M.; Yi, L.; De Gier, J. W.; Kuhn, A.; Dalbey, R. E. Defining the regions of Escherichia coli YidC that contribute to activity. *Journal of Biological Chemistry* **2003**, *278*, 48965–48972.

(11) Nass, K. J.; Ilie, I. M.; Saller, M. J.; Driessen, A. J. M.; Caflisch, A.; Kammerer, R. A.; Li, X. The role of the N-terminal amphipathic helix in bacterial YidC: Insights from functional studies, the crystal structure and molecular dynamics simulations. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2022**, *1864*, 183825.

(12) Güngör, B.; Flohr, T.; Garg, S. G.; Herrmann, J. M. The ER membrane complex (EMC) can functionally replace the Oxa1 insertase in mitochondria. *PLOS Biology* **2022**, *20*, e3001380.

(13) Rapoport, T. A. Protein translocation across the eukaryotic endoplasmic reticulum and bacterial plasma membranes. *Nature* **2007**, *450*, 663–669.

(14) Krogh, A.; Larsson, B.; Von Heijne, G.; Sonnhammer, E. L. Predicting transmembrane protein topology with a hidden Markov model: Application to complete genomes. *Journal of Molecular Biology* **2001**, *305*, 567–580.

(15) Dalbey, R. E.; Kuhn, A.; Zhu, L.; Kiefer, D. The membrane insertase YidC. *Biochimica et Biophysica Acta - Molecular Cell Research* **2014**, *1843*, 1489–1496.

(16) Borowska, M. T.; Dominik, P. K.; Anghel, S. A.; Kossiakoff, A. A.; Keenan, R. J. A YidC-like Protein in the Archaeal Plasma Membrane. *Structure* **2015**, *23*, 1715–1724.

(17) Kuhn, A.; Kiefer, D. Membrane protein insertase YidC in bacteria and archaea. *Molecular Microbiology* **2017**, *103*, 590–594.

(18) Chen, H.; Ogden, D.; Pant, S.; Cai, W.; Tajkhorshid, E.; Moradi, M.; Roux, B.; Chipot, C. A Companion Guide to the String Method with Swarms of Trajectories: Characterization, Performance, and Pitfalls. *Journal of Chemical Theory and Computation* **2022**, *18*, 1406–1422.

(19) Polasa, A.; Hettige, J.; Immadisetty, K.; Moradi, M. An Investigation of the YidC-Mediated Membrane Insertion of Pf3 Coat Protein Using Molecular Dynamics Simulations. *bioRxiv* **2022**,

(20) Dalbey, R. E.; Chen, M. Sec-translocase mediated membrane protein biogenesis. *Biochimica et Biophysica Acta - Molecular Cell Research* **2004**, *1694*, 37–53.

(21) Spann, D.; Pross, E.; Chen, Y.; Dalbey, R. E.; Kuhn, A. Each protomer of a dimeric YidC functions as a single membrane insertase. *Scientific Reports* **2018**, *8*, 589.

(22) Scotti, P. A.; Urbanus, M. L.; Brunner, J.; De Gier, J. W. L.; Von Heijne, G.; Van Der Does, C.; Driessen, A. J.; Oudega, B.; Luirink, J. YidC, the Escherichia coli homologue of mitochondrial Oxa1p, is a component of the Sec translocase. *EMBO Journal* **2000**, *19*, 542–549.

(23) Facey, S. J.; Kuhn, A. Membrane integration of E. coli model membrane proteins. *Biochimica et Biophysica Acta - Molecular Cell Research* **2004**, *1694*, 55–66.

(24) Samuelson, J. C.; Chen, M.; Jiang, F.; Möller, I.; Wiedmann, M.; Kuhn, A.; Phillips, G. J.; Dalbey, R. E. YidC mediates membrane protein insertion in bacteria. *Nature* **2000**, *406*, 637–641.

(25) Kiefer, D.; Kuhn, A. YidC-mediated membrane insertion. *FEMS Microbiology Letters* **2018**, *365*.

(26) Laskowski, P. R.; Pluhackova, K.; Haase, M.; Lang, B. M.; Nagler, G.; Kuhn, A.; Müller, D. J. Monitoring the binding and insertion of a single transmembrane protein by an insertase. *Nature Communications* **2021**, *12*, 7082.

(27) Lewis, A. J. O.; Hegde, R. S. A unified evolutionary origin for the ubiquitous protein transporters SecY and YidC. *BMC Biology* **2021**, *19*, 266.

(28) Endo, Y.; Shimizu, Y.; Nishikawa, H.; Sawasato, K.; Nishiyama, K.-i. Interplay between MPIase, YidC, and PMF during Sec-independent insertion of membrane proteins. *Life Science Alliance* **2022**, *5*.

(29) Xin, Y.; Zhao, Y.; Zheng, J.; Zhou, H.; Zhang, X. C.; Tian, C.; Huang, Y. Structure of YidC from Thermotoga maritima and its implications for YidC-mediated membrane protein insertion. *FASEB Journal* **2018**, *32*, 2411–2421.

(30) Yuan, J.; Phillips, G. J.; Dalbey, R. E. Isolation of cold-sensitive yidC mutants provides insights into the substrate profile of the YidC insertase and the importance of transmembrane 3 in YidC function. *Journal of Bacteriology* **2007**, *189*, 8961–8972.

(31) Klenner, C.; Kuhn, A. Dynamic disulfide scanning of the membrane-inserting Pf3 coat protein reveals multiple YidC substrate contacts. *Journal of Biological Chemistry* **2012**, *287*, 3769–3776.

(32) Kol, S.; Nouwen, N.; Driessen, A. J. Mechanisms of YidC-mediated insertion and as-

sembly of multimeric membrane protein complexes. *Journal of Biological Chemistry* **2008**, *283*, 31269–31273.

(33) Ernst, S.; Schönbauer, A. K.; Bär, G.; Börsch, M.; Kuhn, A. YidC-driven membrane insertion of single fluorescent Pf3 coat proteins. *Journal of Molecular Biology* **2011**, *412*, 165–175.

(34) Funes, S.; Kauff, F.; Van Der Sluis, E. O.; Ott, M.; Herrmann, J. M. Evolution of YidC/Oxa1/Alb3 insertases: Three independent gene duplications followed by functional specialization in bacteria, mitochondria and chloroplasts. *Biological Chemistry* **2011**, *392*, 13–19.

(35) Funes, S.; Hasona, A.; Bauerschmitt, H.; Grubbauer, C.; Kauff, F.; Collins, R.; Crowley, P. J.; Palmer, S. R.; Brady, L. J.; Herrmann, J. M. Independent gene duplications of the YidC/Oxa/Alb3 family enabled a specialized cotranslational function. *Proceedings of the National Academy of Sciences of the United States of America* **2009**, *106*, 6656–6661.

(36) Kohler, R.; Boehringer, D.; Greber, B.; Bingel-Erlenmeyer, R.; Collinson, I.; Schaffitzel, C.; Ban, N. YidC and Oxa1 Form Dimeric Insertion Pores on the Translating Ribosome. *Molecular Cell* **2009**, *34*, 344–353.

(37) He, H.; Kuhn, A.; Dalbey, R. E. Tracking the Stepwise Movement of a Membrane-inserting Protein In Vivo. *Journal of Molecular Biology* **2020**, *432*, 484–496.

(38) Kumazaki, K.; Kishimoto, T.; Furukawa, A.; Mori, H.; Tanaka, Y.; Dohmae, N.; Ishitani, R.; Tsukazaki, T.; Nureki, O. Crystal structure of Escherichia coli YidC, a membrane protein chaperone and insertase. *Scientific Reports* **2014**, *4*, 7299.

(39) Kumazaki, K.; Chiba, S.; Takemoto, M.; Furukawa, A.; Nishiyama, K. I.; Sugano, Y.; Mori, T.; Dohmae, N.; Hirata, K.; Nakada-Nakura, Y. et al. Structural basis of Sec-independent membrane protein insertion by YidC. *Nature* **2014**, *509*, 516–519.

(40) Chen, Y.; Soman, R.; Shanmugam, S. K.; Kuhn, A.; Dalbey, R. E. The role of the strictly conserved positively charged residue differs among the gram-positive, gram-negative, and chloroplast YidC homologs. *Journal of Biological Chemistry* **2014**, *289*, 35656–35667.

(41) Chen, Y.; Capponi, S.; Zhu, L.; Gellenbeck, P.; Freites, J. A.; White, S. H.; Dalbey, R. E. YidC Insertase of Escherichia coli: Water Accessibility and Membrane Shaping. *Structure* **2017**, *25*, 1403–1414.e3.

(42) Schulze, R. J.; Komar, J.; Botte, M.; Allen, W. J.; Whitehouse, S.; Gold, V. A. M.; a Nijeholt, J. A. L.; Huard, K.; Berger, I.; Schaffitzel, C. et al. Membrane protein insertion and proton-motive-force-dependent secretion through the bacterial holo-translocon SecYEG-SecDF-YajC-YidC. *Proceedings of the National Academy of Sciences* **2014**, *111*, 4844–4849.

(43) Shimokawa-Chiba, N.; Kumazaki, K.; Tsukazaki, T.; Nureki, O.; Ito, K.; Chiba, S. Hydrophilic microenvironment required for the channel-independent insertase function of YidC protein. *Proceedings of the National Academy of Sciences of the United States of America* **2015**, *112*, 5063–5068.

(44) Kedrov, A.; Wickles, S.; Crevenna, A. H.; van der Sluis, E. O.; Buschauer, R.; Berninghausen, O.; Lamb, D. C.; Beckmann, R. Structural Dynamics of the YidC:Ribosome Complex during Membrane Protein Biogenesis. *Cell Reports* **2016**, *17*, 2943–2954.

(45) Kiefer, D.; Kuhn, A. B. T. I. R. o. C. YidC as an Essential and Multifunctional Component in Membrane Protein Assembly. *Academic Press* **2007**, *259*, 113–138.

(46) Gallusser, A.; Kuhn, A. Initial steps in protein membrane insertion. Bacteriophage M13 procoat protein binds to the membrane surface by electrostatic interaction. *The EMBO journal* **1990**, *9*, 2723–2729.

(47) Wickles, S.; Singharoy, A.; Andreani, J.; Seemayer, S.; Bischoff, L.; Berninghausen, O.; Soeding, J.; Schulten, K.; van der Sluis, E. O.; Beckmann, R. A structural model of the active ribosome-bound membrane protein insertase YidC. *eLife* **2014**, *3*, 1–17.

(48) Ito, K.; Shimokawa-Chiba, N.; Chiba, S. Sec translocon has an insertase-like function in addition to polypeptide conduction through the channel. *F1000Research* **2019**, *8*, F1000 Faculty Rev–2126.

(49) Tanaka, Y.; Izumioka, A.; Abdul Hamid, A.; Fujii, A.; Haruyama, T.; Furukawa, A.; Tsukazaki, T. 2.8-Å crystal structure of Escherichia coli YidC revealing all core regions, including flexible C2 loop. *Biochemical and Biophysical Research Communications* **2018**, *505*, 141–145.

(50) Huang, J.; Rauscher, S.; Nawrocki, G.; Ran, T.; Feig, M.; de Groot, B. L.; Grubmüller, H.; MacKerell Jr, A. D. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nature methods* **2017**, *14*, 71–73.

(51) Klauda, J. B.; Venable, R. M.; Freites, J. A.; O'Connor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell, A. D.; Pastor, R. W. Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *Journal of Physical Chemistry B* **2010**, *114*, 7830–7843.

(52) Phillips, J. C.; Braun, R.; Wang, W.; Gumbart, J.; Tajkhorshid, E.; Villa, E.; Chipot, C.; Skeel, R. D.; Kalé, L.; Schulten, K. Scalable molecular dynamics with NAMD. *Journal of Computational Chemistry* **2005**, *26*, 1781–1802.

(53) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of simple potential functions for simulating liquid water. *The Journal of Chemical Physics* **1983**, *79*, 926–935.

(54) Jo, S.; Kim, T.; Im, W. Automated builder and database of protein/membrane complexes for molecular dynamics simulations. *PLoS ONE* **2007**, *2*.

(55) Reid, J. K. In *Large Sparse Sets of Linear Equations*; Reid, J. K., Ed.; Academic Press: London, 1971; pp 231–254.

(56) Martyna, G. J.; Tobias, D. J.; Klein, M. L. Constant pressure molecular dynamics algorithms. *J. Chem. Phys.* **1994**, *101*, 4177–4189.

(57) Feller, S. E.; Zhang, Y.; Pastor, R. W.; Brooks, B. R. Constant pressure molecular dynamics simulation: The Langevin piston method. *J. Chem. Phys.* **1995**, *103*, 4613–4621.

(58) Govind Kumar, V.; Ogden, D. S.; Isu, U. H.; Polasa, A.; Losey, J.; Moradi, M. Prefusion Spike Protein Conformational Changes Are Slower in SARS-CoV-2 than in SARS-CoV-1. *Journal of Biological Chemistry* **2022**, *298*.

(59) Immadisetty, K.; Polasa, A.; Shelton, R.; Moradi, M. Elucidating the Molecular Basis of pH Activation of an Engineered Mechanosensitive Channel. *bioRxiv* **2021**,

(60) Polasa, A.; Mosleh, I.; Losey, J.; Abbaspourrad, A.; Beitle, R.; Moradi, M. Developing a Rational Approach to Designing Recombinant Proteins for Peptide-Directed Nanoparticle Synthesis. *Nanoscale Adv.* **2022**, *4*, 3161–3171.

(61) Moradi, M.; Enkavi, G.; Tajkhorshid, E. Atomic-level characterization of transport cycle thermodynamics in the glycerol-3-phosphate:phosphate transporter. *Nat. Commun.* **2015**, *6*, 8393.

(62) Moradi, M.; Sagui, C.; Roland, C. Calculating relative transition rates with driven nonequilibrium simulations. *Chem. Phys. Lett.* **2011**, *518*, 109–113.

(63) Moradi, M.; Sagui, C.; Roland, C. Invstigating rare events with nonequilibrium work measurements: I. nonequilibrium transition path probabilities. *J. Chem. Phys.* **2014**, *140*, 034114.

(64) Moradi, M.; Sagui, C.; Roland, C. Invstigating rare events with nonequilibrium work measurements: II. transition and reaction rates. *J. Chem. Phys.* **2014**, *140*, 034115.

(65) Moradi, M.; Tajkhorshid, E. Mechanistic Picture for Conformational Transition of a Membrane Transporter at Atomic Resolution. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 18916–18921.

(66) Ogden, D.; Moradi, M. *Structure and Function of Membrane Proteins*; Springer US: New York, NY, 2021; pp 289–309.

(67) Immadisetty, K.; Polasa, A.; Shelton, R.; Moradi, M. Elucidating the Molecular Basis of Spontaneous Activation in an Engineered Mechanosensitive Channel. *Computational and Structural Biotechnology Journal* **2022**,

(68) Harkey, T.; Govind Kumar, V.; Hettige, J.; Tabari, S. H.; Immadisetty, K.; Moradi, M. The Role of a Crystallographically Unresolved Cytoplasmic Loop in Stabilizing the Bacterial Membrane Insertase YidC2. *Scientific Reports* **2019**, *9*, 14451.

(69) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual molecular dynamics. *Journal of Molecular Graphics* **1996**, *14*, 33–38.

(70) Immadisetty, K.; Hettige, J.; Moradi, M. What Can and Cannot Be Learned from Molecular Dynamics Simulations of Bacterial Proton-Coupled Oligopeptide Transporter GkPOT? *J. Phys. Chem. B* **2017**, *121*, 3644–3656.

(71) Immadisetty, K.; Hettige, J.; Moradi, M. Lipid-Dependent Alternating Access Mechanism of a Bacterial Multidrug ABC Exporter. *ACS Central Science* **2019**, *5*, 43–56.

(72) Bakan, A.; Meireles, L. M.; Bahar, I. ProDy: Protein dynamics inferred from theory and experiments. *Bioinformatics* **2011**, *27*, 1575–1577.

(73) Sethi, A.; Eargle, J.; Black, A. A.; Luthey-Schulten, Z. Dynamical networks in tRNA:protein complexes. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 6620–6625.

(74) Brown, D. K.; Penkler, D. L.; Amamuddy, O. S.; Ross, C.; Atilgan, A. R.; Atilgan, C.; Bishop, Ö. T. MD-TASK: a software suite for analyzing molecular dynamics trajectories. *Bioinformatics* **2017**, *33*, 2768 – 2771.

(75) Chen, M.; Samuelson, J. C.; Jiang, F.; Muller, M.; Kuhn, A.; Dalbey, R. E. Direct interaction of YidC with the Sec-independent Pf3 coat protein during its membrane protein insertion. *Journal of Biological Chemistry* **2002**, *277*, 7670–7675.

(76) Yu, Z.; Koningstein, G.; Pop, A.; Luirink, J. The conserved third transmembrane segment of YidC contacts nascent Escherichia coli inner membrane proteins. *Journal of Biological Chemistry* **2008**, *283*, 34635–34642.

(77) Shimokawa-Chiba, N.; Kumazaki, K.; Tsukazaki, T.; Nureki, O.; Ito, K.; Chiba, S. Hydrophilic microenvironment required for the channel-independent insertase function of YidC protein. *Proceedings of the National Academy of Sciences of the United States of America* **2015**, *112*, 5063–5068.