**Sex differences in inter-individual gene expression variability across human tissues**

**Authors:**

Samuel Khodursky[1], Caroline S. Jiang[2], Eric B. Zheng[1], Roger Vaughan[2], Daniel R. Schrider[3], Li Zhao[1,*]

**Affiliations:**

[1]Laboratory of Evolutionary Genetics and Genomics, The Rockefeller University, New York, NY 10065,

USA

[2]Department of Biostatistics, The Rockefeller University, New York, NY 10065, USA

[3]Department of Genetics, University of North Carolina, Chapel Hill, NC 27514, USA

Correspondence to: lzhao@rockefeller.edu

**Abstract**

Understanding phenotypic sex differences has long been a goal of biology from both a medical and

evolutionary perspective. Although much attention has been paid to mean differences in phenotype

between the sexes, little is known about sex differences in phenotypic variability. To gain insight into sex

differences in inter-individual variability at the molecular level, we analyzed RNA-seq data from 43

tissues from the Genotype-Tissue Expression project (GTEx). Within each tissue, we identified genes that

show sex differences in gene expression variability. We found that these sex-differentially variable (SDV)

genes are associated with various important biological functions, including sex hormone response,

immune response, and other signaling pathways. By analyzing single-cell RNA sequencing data collected

from breast epithelial cells, we found that genes with sex differences in gene expression variability in

breast tissue tend to be expressed in a cell-type-specific manner. We looked for an association between

SDV expression and Graves' disease, a well-known heavily female-biased disease, and found a

significant enrichment of Graves' associated genes among genes with higher variability in females in

thyroid tissue. This suggests a possible role for SDV expression in the context of sex-biased disease. We

then examined the evolutionary constraints acting on genes with sex differences in variability and found

that they exhibit evidence of increased selective constraint. Through analysis of sex-biased eQTL data, we

found evidence that SDV expression may have a genetic basis. Finally, we propose a simple evolutionary

model for the emergence of sex-differentially variable expression from sex-specific constraints.

**Introduction**

Identifying, characterizing, and understanding phenotypic differences between the sexes has long been a

major goal of biology. A very wide variety of traits in humans show significant sex differences including

morphological, behavioral, metabolic, immune, and disease traits (Chella Krishnan et al., 2018; Choleris

et al., 2018; Klein and Flanagan, 2016; Ober et al., 2008). These trait differences are likely in part due to

sex differences in gene expression and regulation (Naqvi et al., 2019; Parsch and Ellegren, 2013;

Williams and Carroll, 2009). As a phenotype, gene expression is both relatively straightforward to

measure and highly quantitative. As a result, many studies have examined differences in mean gene

expression levels between the sexes in many species (Catalán et al., 2012; Gershoni and Pietrokovski,

2017; Khodursky et al., 2020; Lopes-Ramos et al., 2020; Naqvi et al., 2019; Oliva et al., 2020; Parsch and

Ellegren, 2013). Sex-biased expression (SBE) is thought to arise from sex-specific genetic architectures

which are ultimately established by sex-determination systems initiated during early development

(Williams and Carroll, 2009). In addition to underlying sexually dimorphic phenotypes, genes with SBE

have been found to possess many interesting evolutionary and genomic properties (Parsch and Ellegren

2013). For example, SBE has been associated with increased rates of evolution at both the coding and

expression levels in *Drosophila* and other animals (Khodursky et al., 2020; Meisel et al., 2012; Parsch

and Ellegren, 2013; Ranz et al., 2003). In humans, genes with SBE have been associated with reduced

selective constraint (Gershoni and Pietrokovski, 2017; Naqvi et al., 2019). Additionally, genes with SBE

have been shown to be distributed across the genome in a non-random fashion with respect to the X

chromosome – thereby linking the evolution of sex-biased genes to the unique evolutionary forces

experienced by the X chromosome (Khodursky et al., 2020; Oliva et al., 2020; Parisi et al., 2003).

Relative to sex differences in phenotypic means, sex differences in phenotypic variabilities have

been less thoroughly studied. In the biomedical sciences there has been a long-standing assumption that

females must generally be more phenotypically variable due to the menstrual or estrous cycles; however,

there has been little empirical evidence to support this assumption (Beery and Zucker, 2011). In fact, a

meta-analysis of microarray studies in mice and humans revealed that males may have slightly higher

expression variability (Itoh and Arnold, 2015) and a meta-analysis of various traits in mice indicated that

sex differences in phenotypic variability are trait dependent (Zajitschek et al., 2020). Despite growing

interest in understanding sex differences in phenotypic variability, the biological and evolutionary

properties of genes with sex differences in gene expression variability have not, to our knowledge, been

explored in humans. Here, we define genes with sex-differentially variable (SDV) expression to be genes

which show significant differences in inter-individual gene expression variability between the sexes.

From a medical standpoint, genes with SDV expression may underlie diseases for which there are sex

differences in prevalence or outcome. For example, one could imagine a disease partially caused by the

aberrant expression of a certain gene (akin to the liability threshold model; (Falconer, 1967)). If one sex

has more variable expression of the gene in a normal healthy population, this may help explain a higher

prevalence of that disease in that sex. Additionally, since more variable expression has been associated

with reduced effectiveness of drug targets it is possible that SDV expression may account for sex

differences in drug effectiveness, which are known to exist (Simonovsky et al., 2019; Soldin and

Mattison, 2009).

Understanding the evolutionary properties of these genes may give us novel insight into the

evolution of sex differences. For example, genes with sex differences in expression variability may evolve

under sex-specific constraints (e.g., purifying or stabilizing selection) and be of particular functional

importance for a given sex. Here we examine the biological properties and evolutionary constraints

associated with SDV expression across 43 human tissues.

**Results**

**Identification and characterization of genes with sex differences in gene expression variability**

We obtained RNA-seq count data for 43 tissues from the GTEx project (version 8). Since we sought to identify genes with sex differences in gene expression variability, we first compared several statistical methods using power simulations. Specifically, we compared three tests: the Fligner-Killeen test, Levene's test, and a test based on negative binomial generalized additive models for location, scale and shape (GAMLSS) ( Rigby and Stasinopoulos, 2005). Since RNA-seq count data is known to follow an approximately negative binomial distribution (Robinson and Smyth, 2007; Robinson et al., 2010), we performed power simulations by drawing random samples from negative binomial distributions with identical means but different variances by varying the overdispersion term (see methods). At each parameter combination, 1000 pairs of samples were drawn, and tests were performed at an alpha of 0.05. We found that GAMLSS had the highest power across a range of parameters, and this was especially true when testing for differences in variance between samples of unequal size (**Supplementary Figure 1A-D**). Both GAMLSS and Levene's test managed to maintain an adequate false positive rate (FPR) across a range of parameters. Beyond increased power, GAMLSS offers other advantages over the other two tests: GAMLSS negative binomial models allow for the independent parametrization of both the mean and the non-Poisson variation (overdispersion) terms as a function of variables and covariates of interest, including library size (de Jong et al., 2019; Rigby and Stasinopoulos, 2005).

When considering the inter-sample variation of gene expression, it is natural to consider the coefficient of variation: the standard deviation divided by the mean. This provides a mean-normalized statistic for variability. Notably, it is thought that the Poisson component of the coefficient of variation in RNA-seq data is due to technical variation while the overdispersion component of the coefficient of variation contains the true biological variation (McCarthy et al., 2012). The overdispersion parameter fit by GAMLSS can therefore be considered to be the square of the biological coefficient of variation (de Jong et al., 2019; McCarthy et al., 2012). Observed overdispersion values and effect sizes in GTEx data were similar to the simulated values, as determined by models fit using GAMLSS (**Supplementary Figure 1E**).

To determine the effect of sex on gene expression variation we fit three negative binomial

generalized additive models using the R library 'gamlss' for each expressed gene in each tissue (Rigby

and Stasinopoulos, 2005). To reduce the risk of confounding due to unequal sampling across various

populations, we limited our analysis to European-American individuals. Significance of the effect of sex

on overdispersion and mean expression level was determined through a pair of likelihood-ratio tests (see

methods). All models included covariates for age, RNA integrity number (RIN), ischemic time, and the

first five genotype principal components (to control for population structure) for both mean and

overdispersion terms. We initially considered genes to be sex-differentially variable (SDV) if the effect of

sex on overdispersion was significant at FDR<0.05. Since we observed that outliers dramatically

influenced estimates of variation in some genes, as a secondary test we performed randomization tests by

randomly permuting the sex labels on genes with initially significant coefficients for the effect of sex on

overdispersion and refitting the GAMLSS models (see methods) (**Figure 1A**). Coefficients for the effect

of sex on overdispersion were saved after each permutation. We used 1000 rounds of permutation to

generate empirical null distributions for the effect of sex on overdispersion. Initially significant genes

with FDR-adjusted empirical p-values greater than 0.05 in the permutation test were no longer considered

to be SDV (see methods). Results from all tested genes are included in **Supplementary Table 1**.

Across 43 tissues, we identified 3895 unique genes with SDV expression in at least one tissue out

of a total 22054 tested genes (**Figure 1A, B and Supplementary Table 1**). We were able to identify

SDV genes in 42 out of the 43 tested tissues (**Figure 1B**). Examples of two genes with SDV expression in

the pituitary gland and breast tissue are shown in **Figure 1C**. A total of 2792 unique genes showed higher

variability in males in at least one tissue, while 1382 genes showed higher variability in females in at least

one tissue. This imbalance between the sexes was largely driven by breast tissue. Surprisingly, 279 genes

had higher variability in males in one tissue and higher variability in females in at least one other tissue.

The vast majority (3347) of SDV genes only showed SDV expression in one tissue while 548 genes

showed SDV expression in multiple tissues.

Breast tissue had the largest number of SDV genes with 2049 genes showing significantly higher

variation in males and 125 genes showing significantly higher variation in females (**Figure 1B).**  Apart

from a few tissues**,** SDV genes were not consistently expressed at higher or lower levels than non-SDV

genes across tissues (in terms of both CPM and TPM), suggesting that our discovery of SDV genes was

not significantly biased towards genes of high or low expression (**Supplementary Figure 2**).

Given that differences in variation are generally difficult to detect between groups, we performed

a replication analysis using data from breast tissue to help validate our statistical approach. First, we

randomly split individuals into discovery and replication sets – 70% of males and females were placed in

the discovery set while the rest were placed in the replication set. We identified genes with sex

differences in overdispersion in the discovery set using the combined likelihood ratio test and permutation

approach described above (both at FDR<0.05). These genes were then tested in the replication set: genes

with p-values for the overdispersion term less than 0.05 in both the likelihood ratio test and permutation

test were considered 'replicated' if the sex with higher variability in the discovery set was also more

variable in the replication set. We observed a replication rate of 21.9% (**Supplementary Figure 3**). This

replication rate is limited by reduced power due to the relatively small size of the replication set. To

determine the significance of this replication rate we generated an empirical null distribution under the

null hypothesis that our statistical approach was simply randomly sampling genes. Therefore, we

randomly sampled genes from the discovery set and calculated the replication rate for each sample. The

median replication rate under the null hypothesis was found to be 8.4% and out of 10,000 random

samples none had a replication rate higher than 11.3% (**Supplementary Figure 3**). This null replication

rate should not be considered as a type 1 error rate. Since we are randomly sampling genes from the set of

all expressed genes, we are also capturing some true SDV genes in each sample. One would expect the

replication rate under random sampling to be higher than 5% even if the FDR is properly controlled at

5%.

Since our GAMLSS framework allowed us to simultaneously test for sex differences in both

mean and overdispersion we examined the relationship between SDV genes and genes with sex-biased

expression (SBE). In several tissues a majority of SDV genes also displayed sex-biased expression

(**Supplementary Figure 4**). Although a minority of SDV genes in breast tissue, 466 genes with

significant sex differences in both mean and overdispersion showed discordance, with higher

overdispersion but lower mean expression in one sex versus the other (**Supplementary Figure 4**).

Specifically, 387 of those genes simultaneously had significantly higher overdispersion in males and

significantly higher mean expression in females.

Given that genes with sex-biased expression had been shown to be overrepresented on the X

chromosome in a variety of tissues and species (Khodursky et al., 2020; Oliva et al., 2020; Parisi et al.,

2003), we wondered if SDV genes were similarly enriched on the X chromosome. However, we found no

consistent enrichment of SDV genes on the X chromosome; the enrichment patterns were highly tissue-

dependent (**Supplementary Figure 5**).

To identify biological processes associated with SDV gene expression, we performed an

overrepresentation analysis using gene sets obtained from the hallmark gene set collection of MSigDB

(Liberzon et al., 2015; Subramanian et al., 2005). The hallmark gene set collection consists of non-

redundant and refined gene sets associated with various biological processes and states (Liberzon et al.,

2015). The hallmark gene set collection is ideal for our analysis since we identified few SDV genes in

most tissues – a larger, more redundant, and less refined collection would introduce more noise while

increasing the multiple testing burden for our overrepresentation analysis. We first performed an

enrichment analysis at the tissue level and aggregated results across tissues using an empirical approach

based on Fisher's combined probability method (**Figure 2A, B, C**) (see methods). Following the meta-

analysis, we found that genes with higher variation in males were overrepresented in sets related to

estrogen and androgen response, immune response, and cancer-associated pathways (**Figure 2A**). Genes

with increased variation in females were similarly overrepresented in sets related to immune response,

estrogen response, and cancer-associated pathways (**Figure 2A)**. Notably, there was overlap in multiple

sets between the two sexes even though the lists of SDV genes did not show much overlap. At the single-

tissue level, genes with higher variation in males in breast tissue were overrepresented in sets related to

estrogen response and immune response (**Figure 2B**). This suggests that genes of particular biological

importance to one sex (genes related to estrogen response in female breast tissue) may show higher

variation in the opposite sex. To identify transcription factors whose targets are overrepresented in the set

of SDV genes we downloaded the GTRD dataset from MSigDB (Liberzon et al., 2015; Subramanian et

al., 2005). The GTRD collection contains sets of transcription factors and their predicted binding targets

(within the promoter region) (Kolmykov et al., 2021). We first performed an enrichment analysis at the

individual tissue level, followed by a meta-analysis across tissues using the same modified Fisher's

combined probability approach mentioned above (see methods). Following the meta-analysis, we found

that genes with higher variability in females were significantly enriched for targets of MAML1 and GLI1

(**Figure 2C**).  Genes with higher variability in males were significantly enriched for targets of KMT2D.

GLI1 is a transcription factor that is part of the Hedgehog signaling pathway, which is involved in

differentiation during development. However, in fully differentiated tissues GLI1 can act as an oncogene

has been implicated in a wide variety of cancers, including glioblastoma, breast cancer, and colon cancer

(Avery et al., 2021; Clement et al., 2007; Sun et al., 2014). MAML1 is a transcription factor involved in

Notch signaling and is a transcriptional co-activator of Notch receptors (Wu et al., 2000). Additionally,

MAML1 has been shown to interact cooperatively with GLI1 to regulate hedgehog signaling (Quaranta et

al., 2017). KMT2D is a broadly expressed histone methyltransferase that is believed to function as a

tumor suppressor and is frequently mutated in a wide variety of cancers (Rao and Dou, 2015).

**Single-cell RNA-seq data from breast epithelial cells shows that genes with SDV expression tend to**
**be expressed in a cell type-specific manner**

To get further biological insight into genes with SDV expression in breast tissue, which had the most

SDV genes identified out of any tissue by far, we analyzed a previously published single cell RNA-seq

dataset (Nguyen et al., 2018). With count data from individual 4 (female), we applied a standard single

cell RNA-seq pipeline using the 'R' library 'Seurat' (Hao et al., 2020)(see methods). Our analysis

recovered the three major cell types found in the breast epithelium: type 1 and type 2 luminal cells along

with basal cells (Nguyen et al., 2018)(**Supplementary Figure 6**). To determine the cell type specificity of

each gene we combined raw count data from each cell to generate a pseudo-bulk count matrix for each

cell type and calculated CPM values for each gene. We then applied a standard measure of tissue

specificity, $\tau$, to the log-transformed CPM values across the three cell types (see methods).

We found that genes exhibiting SDV expression were expressed in a significantly more cell-type-

specific manner than non-SDV genes (p=1.9e-41, Mann-Whitney U test)(**Figure 2D**).   If there exist

significant sex differences in inter-individual cell-type composition variation, such cell-type-specific

expression could lead to sex differences in expression variability in bulk tissue samples. Overall, these

results suggest that SDV expression may be partly due to sex differences in inter-individual cell-type

composition variation.

**SDV genes in the thyroid with higher variability in females are associated with Graves' disease**

Given the preponderance of sex-biased diseases (Mauvais-Jarvis et al., 2020), we wondered if genes with

SDV expression could be associated with a classically sex-biased disease. Hence, we decided to examine

Graves' disease: a common autoimmune disease of the thyroid which results in hyperthyroidism (McIver

and Morris, 1998). The incidence of Graves' disease is almost ten-fold higher in females than in males

(McIver and Morris, 1998). We obtained GWAS summary data from the NHGRI-EBI GWAS catalog

compiled from 5 studies on the genetics of Graves' disease in various populations (Buniello et al., 2019;

Chu et al., 2011; Cooper et al., 2012; Nakabayashi et al., 2011; Sakaue et al., 2021; Zhao et al., 2013).  Of

the 36 genes implicated in Graves' disease which were expressed in thyroid tissue, 6 displayed SDV

expression with higher variation in females. This corresponds to a significant 9-fold enrichment of

Graves' associated genes among SDV genes with higher variability in females (p=3.5e-5, hypergeometric

test) (Figure 2E). Although many of the genes implicated in Graves' disease are expressed in immune

cells, a characteristic of Graves' disease is the infiltration of immune cells into the thyroid (Dayan et al.,

1991; McIver and Morris, 1998; Totterman et al., 1979). Overall, this analysis suggests a role for SDV

expression in sex-biased diseases.

**Genes with sex-differentially variable expression show evidence of increased selective constraint**

**acting on gene expression and sequence**

Given that genes with sex-biased expression (SBE) have been shown to be associated with

relaxed selective constraint at both the sequence and expression levels, we asked whether SDV genes

would similarly be associated with reduced constraint (Gershoni and Pietrokovski, 2017; Naqvi et al.,

2019). Stabilizing selection acting on gene expression is thought to reduce *cis*-regulatory variation and

reduce the minor allele frequency (MAF) of *cis*-regulatory variants (Glassberg et al., 2019; Josephs et al.,

2015). Therefore, if SDV expression is associated with reduced selective constraint, one would expect an

enrichment of *cis*-regulatory variation in SDV genes. To determine whether SDV genes were enriched or

depleted for *cis*-regulatory variation, we obtained expression quantitative trait loci (eQTL) summary data

for each tissue from GTEx (v8). For each tissue with at least 20 autosomal SDV genes, we tested for

enrichment of genes with significant eQTLs (eGenes, q-value<0.1) within that tissue's set of SDV genes.

Only genes that were tested in both studies in a given tissue were considered. Contrary to expectation, we

observed a depletion of eGenes within the set of SDV genes in most tissues, and this depletion was

significant in three tissues: sun-exposed skin, breast tissue, and pituitary glands (**Figure 3A**). We also

tested whether the median fold enrichment of eGenes across the 20 tested tissues (median fold enrichment

= 0.93) was lower than expected by chance. To this end, we randomly permuted the SDV labels in each

tissue (genes were either SDV or not SDV in each tissue) and calculated a median fold enrichment across

all tissues. In total 10,000 permutations were performed and for each round of permutation, the median

fold enrichment of eGenes in the set of SDV genes was recorded. We found that the observed median fold

enrichment of eGenes among SDV genes was significantly lower than expected by chance suggesting that

expression levels of SDV genes are evolving under elevated constraint (p=1.3e-03, permutation test)

(**Figure 3B).**

We examined if the observed depletion of *cis*-regulatory variation in SDV genes in breast tissue

could be explained by their elevated cell-type specificity. For example, one could imagine that most

expression variation (across bulk samples) in cell-type specific genes is driven by sample-to-sample

differences in cell-type composition and not by *cis*-regulatory genetic factors. Therefore, one might

expect to observe a depletion of eQTLs in cell-type specific genes. Upon examination, we found no

significant relationship between cell-type specificity and the presence or absence of a significant eQTL

(p=0.99, Mann-Whitney *U* test)(**Supplementary Figure 7**).

Although our analysis of eQTL data provided some evidence that expression levels of SDV genes

are evolving under increased constraint, a depletion of *cis*-regulatory variation might also be expected if

the gene expression variation observed in SDV genes was due to environmental factors and/or *trans*-

regulatory genetic variation to a larger extent than in non-SDV genes. We therefore sought to quantify

selective constraint using a complementary—and more direct—approach. To this end, we used 20-species

phastCons scores to look for further evidence that the *cis*-regulatory regions of SDV genes were

evolutionary highly constrained (Siepel et al., 2005).  In particular, we examined the sequence

conservation in the proximal promoter regions of genes –100 bp upstream from the transcription start

sites (TSSs). The phastCons score of a nucleotide can be interpreted as the probability that the nucleotide

resides in a conserved element (Siepel et al., 2005). We found that genes with SDV expression in at least

one tissue showed significantly elevated phastCons scores compared to genes which did not exhibit SDV

expression in any tissues (p=4.1e-05, Mann-Whitney U test) (**Figure 3C**). Only genes expressed in all

tissues with at least 20 SDV genes were considered to remove bias, i.e., a gene is more likely to be SDV

if it's expressed in more tissues and genes expressed in many tissues tend to be more evolutionarily

constrained (Karczewski et al., 2020) .

Similarly, we also examined the LINSIGHT scores of SDV genes. LINSIGHT scores combine

information from intraspecific genetic polymorphism, interspecific genetic divergence, and functional

genomic data to provide a probability that a mutation at a given base-pair will have deleterious fitness

consequences (Huang et al., 2017). We found that genes with SDV expression in at least one tissue

showed significantly higher mean LINSIGHT scores within their proximal promoter regions than genes

without SDV expression (p=1.8e-02, Mann-Whitney U test).

Finally, we asked whether the increased constraint acting on the expression levels of SDV genes

would be reflected in other metrics of dosage sensitivity. To do this, we compared LOEUF scores in SDV

genes and non SDV genes. LOEUF (loss-of-function observed/expected upper bound fraction) is a score

that reflects the tolerance of a protein to loss-of-function mutations (Karczewski et al., 2020). The scores

range from 0 to 2, with higher scores indicating higher tolerance to loss-of-function mutations and

therefore reduced selective constraint. We only considered genes expressed in all tissues with at least 20

SDV genes and found that genes with SDV in at least one tissue had significantly reduced LOEUF scores

compared to genes without SDV expression in any tissue (p=4.2e-10, Mann-Whitney U test) (**Figure 3E**),

corroborating that SDV genes tend to be more constrained at expression level.

**SDV genes in breast tissue show evidence of sex-specific genetic regulation**

To determine if SDV genes showed evidence of sex-specific genetic regulation, we downloaded sex-

biased eQTL (sbQTL) data from GTEx (Oliva et al., 2020). We restricted our analysis to breast tissue

given that it was found to have both the largest number of SDV genes (in our study) and by far the largest

number of sbQTLs of any tissue (Oliva et al., 2020).  To test if genes with sbQTLs were enriched for

SDV genes we performed a gene set enrichment analysis (GSEA). By only considering genes tested in

both analyses and ranking genes according to their corrected -$\log_{10}$(p-values) in the sbQTL analysis we

found that SDV genes were significantly enriched at lower p-values (p=1.2e-2, permutation test)

(**Supplementary Figure 8**). Additionally, we found that all 13 SDV genes with higher variation in males

and significant sbQTLs (FDR<0.25, the threshold used in Oliva et al., 2020) had eQTLs with larger effect

sizes in males than females. This represents a significant enrichment of genes with larger eQTL effect

sizes in males relative to the set of all genes with significant sbQTLs (p=7.4e-4, hypergeometric test).

These results indicate that SDV genes are enriched for sex-specific genetic architectures regulating gene

expression and suggest that SDV expression may have an underlying genetic component.

**Discussion**

In this work, we examined a largely unappreciated aspect of transcriptomic sex differences: genes with sex differences in inter-individual expression variation. In particular, we focused on genes that showed sex differences in inter-individual expression overdispersion. We termed these genes sex-differentially variable (SDV) genes. Across 43 tested tissues, we found that roughly 18% of expressed genes showed SDV patterns of expression in at least one tissue (**Figure 1B**). Our results were undoubtedly affected by the unequal number of samples in different tissues – we had more power to detect SDV genes in some tissues than others. Nevertheless, we identified more SDV genes in breast tissue than in any other tissue. Issues of sample size aside, from a physiological perspective this seems logical since the breast is the only tissue tested known to have functional significance in only one sex: the production of milk is essentially female-specific. Other studies have found evidence that low inter-individual transcriptional variation in a gene is indicative of increased constraint acting on the gene (Fair et al., 2020). Since the vast majority (94%) of SDV genes identified in breast tissue showed higher expression variability in males than in females, this is consistent with the idea that expression levels should be more functionally constrained in female breast tissue than in male breast tissue.

To gain biological insight into the nature of SDV genes we performed a gene set overrepresentation analysis followed by meta-analysis. We found that SDV genes were enriched for genes from sets related to sex hormone response, immune response, and various signaling and cancer-associated pathways (**Figure 2A, B**). Additionally, we found that SDV genes were enriched for the targets of three transcription factors implicated in cancer (**Figure 2C**). The overrepresentation of SDV genes in gene sets related to various biological pathways and transcription factors indicates that SDV expression may be caused by SDV activity at the pathway level. In breast tissue, the overrepresentation of genes with high variability in males in sets related to estrogen response suggests that genes of particular biological importance to one sex may show higher variation in the opposite sex.

In order to further examine the biological significance of SDV genes in breast tissue we analyzed previously published single-cell RNA sequencing data of breast epithelial tissue. We found that SDV

genes (identified in the GTEx data) were expressed in a more cell-type specific manner (**Figure 2D**).

These results indicate that SDV genes may be more likely to be involved in cell-type specific functions.

Additionally, this suggests that sex differences in gene expression variability may be partially driven by

sex differences in cellular composition variability across individuals.

To examine a possible link between SDV expression and disease, we analyzed summary GWAS

data from a classically female-biased disease: Graves' disease. We found a significant enrichment of

Graves' associated genes in the set of SDV genes with higher variability in females in healthy thyroid

tissue (**Figure 2E**). In the future, the role of SDV expression in sex-biased diseases should be thoroughly

investigated by examining expression in both healthy and diseased tissue.

We found that SDV genes tended to be depleted for *cis*-regulatory variation overall (**Figure 3A.**

**B**). Additionally, we showed that SDV genes show significantly increased constraint in both proximal

promoter and protein-coding sequences. These results are consistent with increased constraint acting on

the expression levels of SDV genes and suggest that SDV genes may be evolving under increased sex-

specific constraint acting on gene expression.

We propose that SDV expression can arise through sex-specific increases in stabilizing selection

acting on gene expression. To test this, we performed a forward Monte Carlo simulation of the evolution

of a gene evolving under sex-specific stabilizing selection (**Figure 4**). Our model assumes that the

expression depends on the additive contributions of a shared genetic locus (this locus contributes equally

to both sexes), two sex-specific loci (each locus only contributes to expression in one sex), and noise (to

model environmental effects) (see methods). Therefore, an assumption of the model is some degree of

sex-specific genetic architecture for SDV genes, which is supported by empirical evidence

(**Supplementary Figure 8**). The three loci were in linkage equilibrium. Increased stabilizing selection in

one sex results in a difference in inter-individual phenotypic variances between the two sexes – with the

sex experiencing increased selection having relatively reduced variation (**Figure 4A, B**). Additionally,

increased selection in one sex results in reduced genetic variation in the set of shared genetic loci –

consistent with our observations of reduced *cis*-regulatory variation in SDV genes (**Figure 4C**). These

results are qualitatively consistent with predictions from quantitative genetics theory. Within a single

generation, stronger stabilizing selection in one sex – for example, females – would lead to reduced

additive genetic variance in females compared to males. Assuming that phenotypic variance is the sum of

additive genetic and environmental variances, and that environmental variances are the same in both

sexes, it implies that phenotypic variance will be lower in females than in males. Additionally, increased

stabilizing selection in females will decrease the additive genetic covariance between the sexes to a larger

extent than identical stabilizing selection in both sexes. This corresponds to our observation of reduction

in the variance of the shared genetic locus under conditions of increased selection in females (**Figure 4C**).

A more detailed description of the theoretical predictions is presented in the **Appendix**.

Although sex-specific selection resulting in decreased expression variance in the sex with

increased selection is one possible explanation for SDV expression, it is not the only possibility.

Alternatively, SDV expression, in particular genes with higher variability in females, could be a result of

hormonal fluctuations as part of the menstrual cycle. The overrepresentation of estrogen response genes

among female SDV genes supports this as a possible explanation (**Figure 2A**). These genes may also

evolve under increased selective constraint in females. Both explanations for SDV expression suggest

sex-specific or sex-biased functions for SDV genes, which could help explain the observation that a

majority of SDV genes also exhibit SBE expression in a few tissues (**Supplementary Figure 4**).

Overall, our results suggest that genes with sex differences in inter-individual gene expression

variability are possibly involved in sex-specific biological functions and may be evolving under sex-

specific constraint acting on gene expression. These results highlight the possible importance of SDV

genes in a functional and evolutionary sense, but future work needs to be done to properly understand the

role of SDV genes in human health.

**Methods**

**Power Simulations**

Random samples were generated from negative binomial distributions using the 'R' function 'rnegbin'

from the library 'MASS'. To simulate the effect of 'sex' on the overdispersion parameter, samples were

drawn from two negative binomial distributions with equal mean ($\mu$) parameters and different

overdispersion ($\sigma$, or equivalently $\frac{1}{\theta}$ in the 'MASS' documentation) parameters. Similarly, to determine

false positive rates samples were drawn from identical distributions. To restrict the search space, $\mu$ was

fixed at 1500 for all simulations and the size of both samples combined was fixed at 400. These

parameters were typical of the gene-expression levels and sample sizes found in GTEx data. Moreover,

varying the mean parameter only marginally affected power calculations. For the unbalanced group sizes

simulations, the size of one group was 320 while the size of the other was 80. Both the baseline

overdispersion parameter ($\sigma$) and the difference in overdispersion between groups ($\Delta\sigma$) were varied over

the course of the simulation experiments. The variances of the two samples were compared using the

Fligner-Killeen test ('fk.test' function from the 'R' library 'npsm'), Levene's test ('levene.test' function

from the 'R' library 'lawstat'), and a likelihood-ratio test implemented using negative binomial GAMLSS

models. The more complex GAMLSS model parametrized the mean ($\mu$) and the overdispersion ($\sigma$)

parameters as functions of sex, $\sigma \sim sex$ and $\mu \sim sex$, while the simplified model parametrized only the

mean in terms of sex with the overdispersion being modeled as constant. The p-value for the significance

of the sex overdispersion term was saved. At each parameter combination 1000 sample pairs were drawn

and p-values from each of the three tests were saved. Power was calculated as the fraction of 1000

simulations where the p-value fell below an alpha of 0.05. False positive rate was calculated as the faction

of 1000 simulations where the p-value fell below and alpha of 0.05 when the two underlying distributions

being sampled for a sample pair were identical. It was observed that power for Levene's test markedly

improved when log-transforming the sampled count data, while the Fligner-Killeen test showed largely

unaltered power: hence samples were transformed as ln(counts + 1) before testing with Levene's test and

the Fligner-Killeen test. Raw samples were used for the GAMLSS likelihood-ratio test.

**Identification of SDV genes**

For each tissue, SDV genes were identified separately using RNA-seq count data downloaded from

GTEx-v8 (dbGAP accession number # phs000424.v8.p2). To reduce confounding due to population

structure, only European Americans were considered. First, counts per million (CPM) values were

calculated using edgeR and genes with no expression (CPM=0) in at least one individual were removed.

To identify individuals that were global expression outliers, dimensionality reduction was then performed

on $\log_2$ transformed CPM values using principal component analysis (PCA) with the 'R' function

'prcomp'. Since outliers can greatly affect estimates of variation, we removed individuals whose global

expression profiles deviated significantly from other individuals in a given tissue. Individuals whose

global expression patterns were more than 1.5 interquartile ranges higher than the third quartile or lower

than the first quartile along any principal component explaining more than 5% of the total variance in the

dataset were considered outliers and removed from further analysis.

Following global outlier removal three negative binomial models were fit to the count data (one

gene at a time) using gamlss. The first and most complex model (model 1) parametrized the mean and

non-Poisson variation terms of the negative binomial distribution ($\mu$ and $\sigma$, respectively) as $\mu \sim sex +$

$age + ischemic\ time + RIN + PC1 + PC2 + PC3 + PC4 + PC5 + offset(offset)$ and $\sigma \sim sex +$

$age + ischemic\ time + RIN + PC1 + PC2 + PC3 + PC4 + PC5$, where $RIN$ represents RNA integrity

number and the 5 PCs represent genotype principal components obtained from GTEx . The offset term, to

account for library size was calculated as $\log(library\ size * normalization\ factors)$, where the

normalization factors were calculated using the 'calcNormFactors' function from edgeR using the

'upperquartile' method. Model 2 had the same form as model 1 for $\mu$, however for the formula for the

non-Poisson variation was $\sigma \sim age + ischemic\ time + RIN + 5PCs$. Model 3 had the same formula as

model 1 for $\sigma$ however the formula for the mean was $\mu \sim age + ischemic\ time + RIN + 5PCs +$

$offset(offset)$. To test for the significance of the effect of sex on non-Poisson variation ($p_{\sigma,sex}$) a

likelihood ratio test was performed comparing model 1 and model 2. To test for the significance of the

effect of sex on mean expression ($p_{\mu,sex}$) a likelihood ratio test was performed comparing model 1 and

model 3. The coefficients corresponding to the sex terms for both $\mu$ and $\sigma$ ($\beta_{\mu,sex}$ and $\beta_{\sigma,sex}$) from model

1 were recorded for each gene. After every gene was tested in a given tissue p-values for $\mu$ and $\sigma$ were

corrected independently using the Benjamini-Hochberg (BH) procedure using the 'R' function 'p.adjust'.

Unsurprisingly we noticed that individual expression outliers at the gene level dramatically

affected our estimates of $p_{\sigma,sex}$ and $\beta_{\sigma,sex}$. To account for this and to generally validate the robustness of

our models we performed randomization tests for all genes with a BH corrected $p_{\sigma,sex}$ <0.05. For each

gene the sex labels were randomly permuted, model 1 was fit to the randomized data, and $\beta_{\sigma,sex}$ values

were recorded. We performed 1000 permutations for every gene to generate an empirical null distribution

for $\beta_{\sigma,sex}$. An empirical p-value was calculated as the fraction of random permutations where

$abs(\beta_{\sigma,sex,perm}) > abs(\beta_{\sigma,sex,obs})$. These empirical p-values were then corrected using the BH

procedure. In the end genes were considered SDV if the original BH corrected p-values ($p_{\sigma,sex}$) were less

than 0.05 and the BH corrected empirical p-values were less than 0.05.

**Hallmark enrichment analysis**

Gene set annotations for collection 'H' (hallmark gene sets) were downloaded from MSigDB using the

'R' function 'msigdbr' from the package 'msigdbr'. Only tissues with at least 20 male or female high

variability genes were considered. Enrichment of gene sets for male high variability and female high

variability genes was calculated separately. Enrichment p-values for each gene set in each tissue were

calculated using a hypergeometric test (using the 'R' function 'phyper'). For each gene set a meta-

analysis was performed across all tested tissues using a variation of Fisher's method. This approach was

implemented as follows. For each gene set a test statistic was calculated by taking -2 times the sum of the

log p-values in each tested tissue. In an ordinary implementation of Fisher's method this test statistic

would have a chi-squared distribution under the null hypothesis. However, given that the underlying p-

values are based on a discrete test (the hypergeometric test) a core assumption of Fisher's method is

violated: namely that for an individual tissue p-values should be uniformly distributed under the null

hypothesis. To overcome this, we generated empirical null distributions for the test statistic described

above using a permutation approach: within each tissue we permuted the labels of each gene (not SDV,

female high variability, and male high variability) and calculated a p-value for the label of interest

(female high variability or male high variability) using a hypergeometric test. This permutation procedure

was done for every tissue tested – finally resulting in a null value for the sum of log p-values test statistic

described above. For each gene set 10,000 such permutations were performed to generate an empirical

null distribution for the test statistic. The empirical p-values for each gene set were calculated by finding

the fraction of permutations resulting in a test statistic larger than the one observed. These combined p-

values across gene sets were corrected using the BH procedure and gene sets with corrected p-

values<0.05 were considered significant. Gene sets that were highly tissue/developmentally specific were

not tested in the meta-analysis. The 7 removed hallmark gene sets were pancreas beta cells, adipogenesis,

myogenesis, spermatogenesis, bile acid metabolism, heme metabolism, and xenobiotic metabolism.

**Transcription factor enrichment analysis**

The analysis was performed in a fashion identical to the hallmark enrichment analysis except that the

gene sets from collection 'C3' subcollection 'GTRD' were used.

**Single-cell RNA sequencing analysis**

The count matrix for individual 4 was downloaded from GSE113197 (Nguyen et al., 2018). All

clustering and marker identification analysis was performed using the 'R' library 'Seurat' version 3.2.3.

Cells which had fewer than 500 and more than 6000 expressed features were removed. Cells with more

than 10% of their transcripts coming from mitochondrial genes were removed. The counts were

normalized using the function 'SCTransform'. Both dimensionality reduction (using the function

'RunUMAP') and clustering (function: 'FindClusters') were performed using the first 10 principal

components of the normalized counts. L1, L2, and basal cells were identified based on their expression of

markers "KRT18", "SPLI", and "KRT14". Clusters expressing "KRT18" and "SPLI" were deemed to be

L1 cells, clusters only expressing "KRT18" were deemed to be L2 cells, and clusters expressing

"KRT14" were taken to be basal cells.

To calculate the cell-type specificity of each gene raw counts were aggregated within each of the

three cell types, generated 3 pseudo-bulk count vectors (one for each cell type). For each cell-type CPMs

were calculated by multiplying the pseudo-bulk count vector by 1 million and dividing by the sum of the

vector. The CPM values were then $log_2$ transformed. Cell-type specificity was calculated using the

standard metric for tissue specificity $\tau = \frac{sum(1 - x_{hat})}{length(x_{hat}) - 1}$, where $x_{hat}$ is the vector representing a given

gene's expression across the 3 cell-types divided by the maximum expression of that gene in any of the

three cell-types.

**Graves' disease analysis**

Summary GWAS statistics were obtained from the NHGRI-EBI GWAS catalog for study EFO_0004237,

downloaded on 08/04/2022 (Buniello et al., 2019).

**eQTL analysis**

Summary statistics for single-tissue eQTL mapping performed in European-American individuals were

downloaded from the GTEx-v8 website. Genes with at least one eQTL with q-value<0.1 were considered

to be eGenes. A two-sided Fisher's exact test, as implemented in the 'fisher.test' function, was used to

test for the enrichment/depletion of eGenes among SDV genes in each tissue. Only tissues with at least 20

autosomal SDV genes were considered. Only autosomal genes with expression high enough to be tested

in both the original eQTL study and our study were considered. To test if the median fold enrichment of

eGenes across tissues was significant we randomly permuted the SDV labels in each tissue and calculated

a median fold enrichment across all tissues. In total 10,000 permutations were performed and for each

round of permutation the median fold enrichment of eGenes across the 20 tissues was recorded. The

empirical p-value was calculated as the fraction of all permutations where the fold enrichment was at least

as low as the observed fold enrichment.

**PhastCons analysis**

Genome-wide phastCons scores (in hg38 coordinates) generated from a 20 species alignment were

downloaded from http://hgdownload.cse.ucsc.edu/goldenPath/hg38/phastCons20way/. Transcription start

sites (TSS) for all annotated genes were downloaded from BioMart using the 'R' package biomaRt

(Durinck et al., 2005, 2009). The most distal TSS was used for each gene. Only genes expressed in all

tissues containing at least 20 SDV genes were tested (20 tissues overall). A gene was considered 'SDV' if

it displayed 'SDV' expression in at least 1 of the 20 tested tissues, otherwise it was considered to be 'not

SDV'.  Requiring genes to be expressed in all 20 tissues was done to remove bias, i.e., a gene is more

likely to be SDV if it's expressed in more tissues and genes expressed in many tissues tend to be more

evolutionarily constrained (Karczewski et al., 2020). Further increasing the number of tissues (by

decreasing the minimum number of SDV genes allowed in each tissue) reduces power by decreasing the

intersection of genes expressed across all tissues. Mean phastCons scores were obtained for each gene in

a region 100 bp upstream from their TSSs.

**LINSIGHT analysis**

Data was downloaded from the Siepel Laboratory website: https://siepellab.labsites.cshl.edu/software/ .

The data was processed and analyzed in the same manner as the phastCons data following conversion to

hg38 coordinates.

**Loss of function analysis**

Loss of function metrics were downloaded from https://gnomad.broadinstitute.org/downloads . In

particular the v2.1.1 loss of function metrics by gene were used. The analysis was similar to the analysis

for PhastCons score: only genes expressed across all tissues with at least 20 SDV genes were considered.

LOEUF scores were compared between 'SDV' and 'not SDV' genes.

**Sex-specific stabilizing selection simulation**

The simulation assumed a diploid, diecious, and randomly mating population of 500 individuals. In each

individual, the phenotype (expression) was modelled as the additive contribution of 3 genetic loci and

environmental noise. One of the genetic loci had equal effect in both sexes while each of the two other

genetic loci only had effect in one sex or the other. The allelic effects of each genetic locus in each

individual were initialized as draws from a standard normal distribution (mean=0 and standard

deviation=1). The environmental component was also taken to be a random number drawn from a

standard normal distribution. Similarly, mutation resulted in the addition of a number drawn from a

standard normal distribution to the allelic effect and the mutation rate was set to 5e-3 per generation at

each genetic locus in each individual. The fitness of each individual was taken to be $w = e^{-s(G+E)^2}$,

where '*s*' represented the strength of selection, '*G*' was the total sum of genetic allelic effects (from the

shared and sex-specific locus), and '*E*' was the environmental effect. For the simulations with equal

selection in both sexes '*s*' was 0.005, while in simulation with sex-specific selection '*s*' was 0.005 in

males and 0.05 in females. Selection only affected viability and had no impact on fecundity. An

individual survived to reproduce if a random draw from a uniform distribution on [0, 1] was less

than or equal to the individual's fitness; therefore, the fitness represents the probability of an

individual surviving to reproduce. In each generation 100 males and 100 females were chosen and

paired at random with replacement. Each pair produced 5 offspring and the sex of each offspring was

determined randomly. The simulations were run for a total of 2000 generations. Every 10 generations the

variance of the phenotype (across individuals in the population) was recorded along with the variance of

the allelic effect of the shared genetic component. For both the equal selection and sex-specific selection

models 30 independent simulations were run.

**GSEA analysis of sbQTL data**

Sex-biased QTL (sbQTL) data from breast tissue was downloaded from GTEx v8. Genes were only

considered if they were tested in our analysis and in the sbQTL analysis. Genes were ranked according to

corrected -$\log_{10}$(p-values) where the (Bonferroni) correction was made to account for multiple SNPs

tested per gene. GSEA analysis was performed using the 'R' function 'fgsea' from the library 'fgsea'

(Korotkevich et al., 2021).

**Appendix**

**Theory behind sex-specific stabilizing selection**

The evolution of a single quantitative trait in males and females can be predicted from the 2-by-2 additive

genetic variance-covariance matrix:

$$\boldsymbol{G} = \begin{bmatrix} G_m & B \\ B & G_f \end{bmatrix}. \qquad (1)$$

Where the diagonal terms represent the additive genetic variances in males and females, while the

identical off-diagonal terms represent the additive genetic covariance between the sexes (Barker et al.,

2010; Lande, 1980). In the absence of directional selection, the change in the genetic variance-covariance

matrix due to stabilizing selection over the course of a generation is (Barker et al., 2010; Phillips and

Arnold, 1989):

$$\Delta\boldsymbol{G} = \boldsymbol{G\gamma G} . \qquad (2)$$

Where $\boldsymbol{\gamma}$ is a matrix describing the strength of stabilizing selection within each sex:

$$\boldsymbol{\gamma} = \begin{bmatrix} \gamma_m & 0 \\ 0 & \gamma_f \end{bmatrix}. \qquad (3)$$

For stabilizing selection to occur the diagonal terms must be negative with decreasing values indicating

stronger stabilizing selection. Substituting (3) and (1) into (2) yields:

$$\Delta\boldsymbol{G} = \begin{bmatrix} \Delta G_m & \Delta B \\ \Delta B & \Delta G_f \end{bmatrix} = \begin{bmatrix} \gamma_m G_m^2 + \gamma_f B^2 & B(\gamma_m G_m + \gamma_f G_f) \\ B(\gamma_m G_m + \gamma_f G_f) & \gamma_f G_f^2 + \gamma_m B^2 \end{bmatrix}. \qquad (4)$$

589    By assuming that initially the trait has identical genetic variance in males and females (which is assumed

590    at the start of our simulation): we can use the simplification $G_m = G_f = G$. This leads to the following

591    equations for $\Delta G_m$ and $\Delta G_f$ :

$$\Delta G_m = \gamma_m G^2 + \gamma_f B^2 \qquad (5a)$$

$$\Delta G_f = \gamma_f G^2 + \gamma_m B^2 \; . \qquad (5b)$$

In the case of stronger stabilizing selection in females: $\gamma_f < \gamma_m < 0$. Additionally, theoretical and

empirical evidence suggests that generally the magnitude of intra-sex genetic variance is larger than the

magnitude of inter-sex genetic covariance: $G > B$ (Barker et al., 2010). Therefore, $(5a, b)$ yields:

$\Delta G_f < \Delta G_m < 0$. Following the assumptions made in our simulation (namely that the phenotypic

variance in each sex is just the sum of the environmental variances and the additive genetic variances $(G)$,

and that the environmental variance is the same in both sexes) leads to the result that phenotypic variance

will decrease to a larger extent in females than in males over the course of a generation. This is what we

observed (however, over multiple generations) in our simulation (**Figure 4A, B**).

To further relate our simulation results to the theory described above we can formulate the phenotypes

and their variances from the simulation as follows:

$$P_m = g_1 + g_2 + \varepsilon \qquad (6a)$$

$$P_f = g_1 + g_3 + \varepsilon \; . \qquad (6b)$$

607    $P_m$ and $P_f$ are the phenotypes in males and females respectively. $g_1$, $g_2$, and $g_3$ are the genetic

608    contributions of the shared genetic locus, the male specific genetic locus, and the female specific genetic

609    locus. The environmental contribution to the phenotypes is $\varepsilon$ . From $(6a, b)$, since all the phenotypic

610    components are independent (so their covariances are 0), the phenotypic variances are:

$$Var(P_m) = Var(g_1) + Var(g_2) + Var(\varepsilon) \qquad (7a)$$

$$Var(P_f) = Var(g_1) + Var(g_3) + Var(\varepsilon) \qquad (7b)$$

613 and the phenotypic covariances are:

$$Cov(P_m, P_f) = Var(g_1) + Var(\varepsilon). \qquad (7c)$$

From $(7a, b, c)$ it becomes apparent that $G_m$ corresponds to $var(g_1) + var(g_2)$, $G_f$ corresponds to

$var(g_1) + var(g_3)$, and $B$ corresponds to $var(g_1)$. From the off-diagonal terms in equation (4) it is

evident that increasing the strength of selection in females (decreasing $\gamma_f$) makes $\Delta B$ more negative –

which is consistent with the observed relative reduction in the genetic variance in the shared locus

$(var(g_1))$ (**Figure 4C**).

**Data access and code availability**

The code generated for this work has been deposited on GitHub: https://github.com/LiZhaoLab/SDV.

**Acknowledgments**

We thank Zhao lab members for discussions about the project, Jason Mezey and Christina Leslie for

helpful discussions during early stages of the project, Mike Love for statistical advice, and Andy Clark for

the suggestion of power analysis. We thank Matt Hahn and Ziyue Gao for their comments and

suggestions for a recent version of the manuscript. We thank Jason Banfelder and Bala Jayaraman from

The Rockefeller University High Performance Computing (HPC) Center for their dedicated work in these

extraordinary times.

**Funding**

This work was supported by National Institutes of Health (NIH) MIRA R35GM133780, the Robertson

Foundation, a Rita Allen Foundation Scholar Program, and a Vallee Scholar Program (VS-2020-35), a

Monique Weill-Caulier Career Scientist Award, and an Alfred P. Sloan Research Fellowship (FG-2018-

10627) to L.Z. C.S.J. was supported by UL1 TR001866 from the National Center for Advancing

Translational Sciences (NCATS), NIH Clinical and Translational Science Award (CTSA) program.

E.B.Z. was supported by a Medical Scientist Training Program grant from the National Institute of

General Medical Sciences of the NIH under award number: T32GM007739 to the Weill

Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program.

**Competing interest statement**

The authors declare that no competing interests exist.

**References**

Avery, J.T., Zhang, R., and Boohaker, R.J. (2021). GLI1: A Therapeutic Target for Cancer. Front. Oncol.
*11*.

Barker, B.S., Phillips, P.C., and Arnold, S.J. (2010). A test of the conjecture that G-matrices are more
stable than B-matrices. Evolution *64*, 2601–2613.

Beery, A.K., and Zucker, I. (2011). Sex bias in neuroscience and biomedical research. Neurosci.
Biobehav. Rev. *35*, 565–572.

Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C., McMahon, A.,
Morales, J., Mountjoy, E., Sollis, E., et al. (2019). The NHGRI-EBI GWAS Catalog of published
genome-wide association studies, targeted  arrays and summary statistics 2019. Nucleic Acids Res. *47*,
D1005–D1012.

Catalán, A., Hutter, S., and Parsch, J. (2012). Population and sex differences in Drosophila melanogaster
brain gene expression. BMC Genomics *13*, 654.

Chella Krishnan, K., Mehrabian, M., and Lusis, A.J. (2018). Sex differences in metabolism and
cardiometabolic disorders. Curr. Opin. Lipidol. *29*, 404–410.

Choleris, E., Galea, L.A.M., Sohrabji, F., and Frick, K.M. (2018). Sex differences in the brain:
Implications for behavioral and biomedical research. Neurosci. Biobehav. Rev. *85*, 126–145.

Chu, X., Pan, C.-M., Zhao, S.-X., Liang, J., Gao, G.-Q., Zhang, X.-M., Yuan, G.-Y., Li, C.-G., Xue, L.-
Q., Shen, M., et al. (2011). A genome-wide association study identifies two new risk loci for Graves'
disease. Nat. Genet. *43*, 897–901.

Clement, V., Sanchez, P., de Tribolet, N., Radovanovic, I., and Ruiz i Altaba, A. (2007). HEDGEHOG-

GLI1 Signaling Regulates Human Glioma Growth, Cancer Stem Cell Self-Renewal, and Tumorigenicity.

Curr. Biol. *17*, 165–172.

Cooper, J.D., Simmonds, M.J., Walker, N.M., Burren, O., Brand, O.J., Guo, H., Wallace, C., Stevens, H.,

Coleman, G., Franklyn, J.A., et al. (2012). Seven newly identified loci for autoimmune thyroid disease.

Hum. Mol. Genet. *21*, 5202–5208.

Dayan, C.M., Londei, M., Corcoran, A.E., Grubeck-Loebenstein, B., James, R.F., Rapoport, B., and

Feldmann, M. (1991). Autoantigen recognition by thyroid-infiltrating T cells in Graves disease. Proc.

Natl. Acad. Sci. *88*, 7415–7419.

Durinck, S., Moreau, Y., Kasprzyk, A., Davis, S., De Moor, B., Brazma, A., and Huber, W. (2005).

BioMart and Bioconductor: a powerful link between biological databases and  microarray data analysis.

Bioinformatics *21*, 3439–3440.

Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the integration of

genomic datasets with the R/Bioconductor  package biomaRt. Nat. Protoc. *4*, 1184–1191.

Fair, B.J., Blake, L.E., Sarkar, A., Pavlovic, B.J., Cuevas, C., and Gilad, Y. (2020). Gene expression

variability in human and chimpanzee populations share common determinants. Elife *9*, e59929.

Falconer, D.S. (1967). The inheritance of liability to diseases with variable age of onset, with  particular

reference to diabetes mellitus. Ann. Hum. Genet. *31*, 1–20.

Gershoni, M., and Pietrokovski, S. (2017). The landscape of sex-differential transcriptome and its

consequent selection in human adults. BMC Biol. *15*, 7.

Glassberg, E.C., Gao, Z., Harpak, A., Lan, X., and Pritchard, J.K. (2019). Evidence for Weak Selective

Constraint on Human Gene Expression. Genetics *211*, 757 LP – 772.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby,

C., Zagar, M., et al. (2020). Integrated analysis of multimodal single-cell data. BioRxiv

2020.10.12.335331.

Huang, Y.-F., Gulko, B., and Siepel, A. (2017). Fast, scalable prediction of deleterious noncoding

variants from functional and  population genomic data. Nat. Genet. *49*, 618–624.

Itoh, Y., and Arnold, A.P. (2015). Are females more variable than males in gene expression? Meta-

analysis of microarray datasets. Biol. Sex Differ. *6*, 1–9.

de Jong, T. V, Moshkin, Y.M., and Guryev, V. (2019). Gene expression variability: the other dimension

in transcriptome analysis. Physiol. Genomics *51*, 145–158.

Josephs, E.B., Lee, Y.W., Stinchcombe, J.R., and Wright, S.I. (2015). Association mapping reveals the

role of purifying selection in the maintenance of genomic variation in gene expression. Proc. Natl. Acad.

Sci. *112*, 15390 LP – 15395.

Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L.,

Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2020). The mutational constraint spectrum quantified

from variation in 141,456 humans. Nature *581*, 434–443.

Khodursky, S., Svetec, N., Durkin, S.M., and Zhao, L. (2020). The evolution of sex-biased gene

expression in the Drosophila brain. Genome Res. *30*, 874–884.

Klein, S.L., and Flanagan, K.L. (2016). Sex differences in immune responses. Nat. Rev. Immunol. *16*,

626–638.

Kolmykov, S., Yevshin, I., Kulyashov, M., Sharipov, R., Kondrakhin, Y., Makeev, V.J., Kulakovskiy, I.

V, Kel, A., and Kolpakov, F. (2021). GTRD: an integrated view of transcription regulation. Nucleic Acids

Res. *49*, D104–D111.

Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N., and Sergushichev, A. (2021). Fast

gene set enrichment analysis. BioRxiv 60012.

Lande, R. (1980). Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic Characters.

Evolution (N. Y). *34*, 292.

Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015). The

Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst. *1*, 417–425.

Lopes-Ramos, C.M., Chen, C.-Y., Kuijjer, M.L., Paulson, J.N., Sonawane, A.R., Fagny, M., Platig, J.,

Glass, K., Quackenbush, J., and DeMeo, D.L. (2020). Sex Differences in Gene Expression and

Regulatory Networks across 29 Human Tissues. Cell Rep. *31*, 107795.

Mauvais-Jarvis, F., Bairey Merz, N., Barnes, P.J., Brinton, R.D., Carrero, J.-J., DeMeo, D.L., De Vries,

G.J., Epperson, C.N., Govindan, R., Klein, S.L., et al. (2020). Sex and gender: modifiers of health,

disease, and medicine. Lancet *396*, 565–582.

McCarthy, D.J., Chen, Y., and Smyth, G.K. (2012). Differential expression analysis of multifactor RNA-

Seq experiments with respect to biological variation. Nucleic Acids Res. *40*, 4288–4297.

McIver, B., and Morris, J.C. (1998). The pathogenesis of graves' disease. Endocrinol. Metab. Clin. North

Am. *27*, 73–89.

Meisel, R.P., Malone, J.H., and Clark, A.G. (2012). Faster-X Evolution of Gene Expression in

Drosophila. PLOS Genet. *8*, e1003013.

Nakabayashi, K., Tajima, A., Yamamoto, K., Takahashi, A., Hata, K., Takashima, Y., Koyanagi, M.,

Nakaoka, H., Akamizu, T., Ishikawa, N., et al. (2011). Identification of independent risk loci for Graves'

disease within the MHC in the Japanese population. J. Hum. Genet. *56*, 772–778.

Naqvi, S., Godfrey, A.K., Hughes, J.F., Goodheart, M.L., Mitchell, R.N., and Page, D.C. (2019).

Conservation, acquisition, and functional impact of sex-biased gene expression in mammals. Science.

*365*, eaaw7317.

Nguyen, Q.H., Pervolarakis, N., Blake, K., Ma, D., Davis, R.T., James, N., Phung, A.T., Willey, E.,

Kumar, R., Jabart, E., et al. (2018). Profiling human breast epithelial cells using single cell RNA

sequencing identifies cell diversity. Nat. Commun. *9*, 2028.

Ober, C., Loisel, D.A., and Gilad, Y. (2008). Sex-specific genetic architecture of human disease. Nat.

Rev. Genet. *9*, 911–922.

Oliva, M., Muñoz-Aguirre, M., Kim-Hellmuth, S., Wucher, V., Gewirtz, A.D.H., Cotter, D.J., Parsana, P.,

Kasela, S., Balliu, B., Viñuela, A., et al. (2020). The impact of sex on gene expression across human

tissues. Science. *369*, eaba3066.

Parisi, M., Nuttall, R., Naiman, D., Bouffard, G., Malley, J., Andrews, J., Eastman, S., and Oliver, B.

(2003). Paucity of genes on the Drosophila X chromosome showing male-biased expression. Science *299*,

697–700.

Parsch, J., and Ellegren, H. (2013). The evolutionary causes and consequences of sex-biased gene

expression. Nat. Rev. Genet. *14*, 83–87.

Phillips, P.C., and Arnold, S.J. (1989). Visualizing Multivariate Selection. Evolution (N. Y). *43*, 1209–

1222.

Quaranta, R., Pelullo, M., Zema, S., Nardozza, F., Checquolo, S., Lauer, D.M., Bufalieri, F., Palermo, R.,

Felli, M.P., Vacca, A., et al. (2017). Maml1 acts cooperatively with Gli proteins to regulate sonic

hedgehog signaling pathway. Cell Death Dis. *8*, e2942–e2942.

Ranz, J.M., Castillo-Davis, C.I., Meiklejohn, C.D., and Hartl, D.L. (2003). Sex-dependent gene

expression and evolution of the Drosophila transcriptome. Science (80-. ). *300*, 1742–1745.

Rao, R.C., and Dou, Y. (2015). Hijacked in cancer: the KMT2 (MLL) family of methyltransferases. Nat.

Rev. Cancer *15*, 334–346.

Rigby, R.A., and Stasinopoulos, D.M. (2005). Generalized additive models for location, scale and shape.

J. R. Stat. Soc. Ser. C (Applied Stat. *54*, 507–554.

Robinson, M.D., and Smyth, G.K. (2007). Moderated statistical tests for assessing differences in tag

abundance. Bioinformatics *23*, 2881–2887.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140.

Sakaue, S., Kanai, M., Tanigawa, Y., Karjalainen, J., Kurki, M., Koshiba, S., Narita, A., Konuma, T.,

Yamamoto, K., Akiyama, M., et al. (2021). A cross-population atlas of genetic associations for 220

human phenotypes. Nat. Genet. *53*, 1415–1424.

Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson, H., Spieth,

J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved elements in vertebrate, insect,

worm, and yeast genomes. Genome Res. *15*, 1034–1050.

Simonovsky, E., Schuster, R., and Yeger-Lotem, E. (2019). Large-scale analysis of human gene

expression variability associates highly variable  drug targets with lower drug effectiveness and safety.

Bioinformatics *35*, 3028–3037.

Soldin, O.P., and Mattison, D.R. (2009). Sex differences in pharmacokinetics and pharmacodynamics.

Clin. Pharmacokinet. *48*, 143–157.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,

Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: A knowledge-

based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. U. S. A. *102*,

15545–15550.

Sun, Y., Wang, Y., Fan, C., Gao, P., Wang, X., Wei, G., and Wei, J. (2014). Estrogen promotes stemness

and invasiveness of ER-positive breast cancer cells through Gli1 activation. Mol. Cancer *13*, 137.

Totterman, T.H., Andersson, L.C., and Häyry, P. (1979). Evidence for thyroid antigen-reactive t

lymphocytes infiltrating the thyroid gland in graves'disease. Clin. Endocrinol. (Oxf). *11*, 59–68.

Williams, T.M., and Carroll, S.B. (2009). Genetic and molecular insights into the development and

evolution of sexual dimorphism. Nat. Rev. Genet. *10*, 797–804.

Wu, L., Aster, J.C., Blacklow, S.C., Lake, R., Artavanis-Tsakonas, S., and Griffin, J.D. (2000). MAML1,

a human homologue of Drosophila mastermind, is a transcriptional  co-activator for NOTCH receptors.

Nat. Genet. *26*, 484–489.

Zajitschek, S.R., Zajitschek, F., Bonduriansky, R., Brooks, R.C., Cornwell, W., Falster, D.S., Lagisz, M.,

Mason, J., Senior, A.M., Noble, D.W., et al. (2020). Sexual dimorphism in trait variability and its eco-

evolutionary and statistical  implications. Elife *9*.

Zhao, S.-X., Xue, L.-Q., Liu, W., Gu, Z.-H., Pan, C.-M., Yang, S.-Y., Zhan, M., Wang, H.-N., Liang, J.,

Gao, G.-Q., et al. (2013). Robust evidence for five new Graves' disease risk loci from a staged genome-

wide  association analysis. Hum. Mol. Genet. *22*, 3347–3362.

**Figure Legend**

792
793 **Figure 1. The identification of genes with sex-differentially-variable gene expression.** A) Outline of
794 pipeline to identify SDV genes. B) The number of genes with SDV expression identified in each tissue.
795 The X-axis has a pseudo-log scale. C) Two examples of SDV genes with higher variation in females
796 (WNT4 in the pituitary gland) and in males (CLEC7A in breast tissue), respectively.
797

798
**Figure 2. Biological properties of SDV genes.** A) Following meta-analysis SDV genes with higher
variation in males and females are enriched for terms related to various signaling pathways, immune
response, and sex-hormone response. B) Genes with higher variability in males are enriched for terms
related to estrogen and immune response in breast tissue. C) Following meta-analysis genes with higher
variability in females are enriched for targets of two transcription factors. D) SDV genes identified in
breast tissue show highly cell-type specific expression in breast epithelial tissue (p-value calculated using
the Mann-Whitney U test) . Higher τ indicates greater cell-type specificity. E) Genes with higher
variability in females in the thyroid are significantly enriched for genes associated with Graves' disease
(p-value calculated using a hypergeometric test).
808

809
**Figure 3. SDV genes are associated with increased constraint.** A) 14 out of 20 tested tissues with at
least 20 SDV genes show a depletion in eGenes among SDV genes. Significance was tested using
Fisher's exact test at FDR<0.05. B) Across the 20 tested tissues the median fold enrichment of eGenes is
significantly lower than expected by chance. The red line shows the observed median fold-enrichment,
while the blue distribution shows the empirical null distribution. C) Genes with SDV expression in at
least one tissue show significantly elevated mean phastCons scores in proximal promoters (100 bp
upstream of TSS) indicating increased conservation. D) Genes with SDV expression in at least one tissue
show significantly increased LINSIGHT scores, indicating greater functional importance. E) Genes with
SDV expression in at least one tissue show significantly reduced LOEUF scores, indicating more
intolerance to loss of function mutations. For C-E, SDV genes were considered to be SDV if they showed
SDV expression in at least one tissue out of the 20 tested tissues. Only genes expressed in all 20 tested
tissues were considered to remove bias. For C-E, p-values were calculated using the Mann-Whitney U
test.
823

824
**Figure 4**. **Simulations showing that SDV expression can arise through sex-specific constraint.**
Simulation results showing the output from 30 simulations in both the equal selection and sex-specific
selection conditions. A) Equal constraint in both sexes results in similar phenotypic variance in both
sexes. B) Increased sex-specific constraint acting in females reduces the phenotypic (expression) variation
across individuals. C) Increased constraint in females results in reduced genetic variance, consistent with
our observations of reduced *cis*-regulatory variation in SDV genes.
831
832