# Nuclear genetic diversity of head lice sheds light on human dispersal around the world

Marina S. Ascunce[1,2,*], Ariel C. Toloza[3], Angélica González-Oliver[4], and David L. Reed[5]

[1]Emerging Pathogens Institute, Department of Plant Pathology, University of Florida, Gainesville, Florida, United States of America

[2]USDA-ARS Center for Medical, Agricultural, and Veterinary Entomology, Gainesville, Florida, United States of America

[3]Centro de Investigaciones de Plagas e Insecticidas (CONICET-UNIDEF), Villa Martelli, Buenos Aires, Argentina

[4]Departamento de Biología Celular, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad de México, México

[5]Florida Museum of Natural History, University of Florida, Gainesville, Florida, United States of America

[*]Corresponding author E-mail: marina.ascunce@usda.gov (MSA)

# Abstract

The human louse, *Pediculus humanus*, is an obligate blood-sucking ectoparasite that has coevolved with humans for millennia. Because of the intimate relationship between this parasite and the human host, the study of human lice has the potential to shed light on aspects of the human evolution that are obscured or difficult to interpret using other biological evidence. In this study, we analyzed the genetic variation in 274 human lice from 25 geographic sites around the world by using nuclear microsatellite loci and female-inherited mitochondrial DNA sequences. Nuclear genetic diversity analysis revealed the presence of two distinct genetic clusters I and II, which are subdivided into subclusters: Ia-Ib and IIa-IIb, respectively. Among these samples, we observed the presence of the two most common louse mitochondrial haplogroups: A and B. Both mitochondrial haplotypes were found in nuclear Clusters I and II. Evidence of nuclear admixture was uncommon (33 lice) and was predominately found in the New World potentially mirroring the pattern of very recent host admixture between Native Americans and Europeans. These findings were supported by novel DIYABC-simulations that were built using both host and parasite data to define parameters and models. In addition to providing new evolutionary knowledge about this human parasite, our study could guide the development of new analyses in other host-parasite systems.

# Introduction

The human louse is a single insect species composed of two distinct morphological, behavioral, and ecological types: the human head louse (*Pediculus humanus capitis* De Geer), and the clothing or body louse (*Pediculus humanus humanus* L.) [1]. Human lice (Phthiraptera: Anoplura) are blood-sucking, wingless, host-specific ectoparasites of humans that are both obligate (cannot live off the host) and permanent (complete their life cycle on a single host). Thus, human lice are inextricably tied to their host in ecological and evolutionary time. The coevolution of these parasites and their primate hosts over the last 25 million years (MY) is very well documented [2,3]. Cophylogenetic analyses revealed that human and chimpanzee lice in the genus *Pediculus* diverged ca. 5.6 MY ago (MYA), which is contemporaneous with the known divergence of chimps and humans [4]. Human head lice have been found in archeological human remains, including a head louse nit obtained from human remains in northeastern Brazil dated 10,000 years old [5] and a nit from Israel dated as 9,000 years old [6] (Fig 1). In host-parasite systems, these long and intimate associations lead to reciprocal adaptations in the hosts and parasites (coevolution), which allows one to identify events of the same age in the host and parasite phylogenies, and provide an internal time calibration for comparative studies of rates of evolution in the two groups [7]. Thus, in cases where parasites coevolve with their hosts, parasite phylogenies will, to some extent, reflect host evolutionary history [8–10]. Parasites have some biological features that differ from their hosts, such as faster evolutionary rates and larger population sizes in parasites, which allows them to maintain greater levels of genetic diversity than their hosts. These attributes of human parasites have provided insights into aspects of human evolution that were unclear from studies of direct human evidence, such as fossil or molecular data [10–13]. Evolutionary studies of human lice have already provided insights about the ecology of early humans, such as when humans started

using clothes [14,15] and how humans travelled around the globe [16–18]. Recently, human DNA

was extracted from nits found on hairs from ancient mummies in South America revealing

ancestral migration routes [19]. Moreover, almost 20 years ago, Reed et al. [2] described highly

divergent human louse mitochondrial clades and hypothesized that each louse mitochondrial

lineage evolved on different hominin hosts. The authors suggested that as anatomically modern

humans (AMH) migrated out of Africa and interacted with other hominis, those different lineages

of lice were transferred among hominins through direct physical contact [2]. This hypothesis of

direct contact between modern and archaic hominins predicted using human louse data was later

confirmed through host genomic studies comparing ancient DNA genomes from hominid remains

with AMH genomes [20].

**Fig 1. Worldwide geographic distribution of collected modern lice (green dots).**

Archeological findings of human lice are shown with the figure of human lice on the map and

the corresponding estimated dates from: [3,5,6,22,100]. In addition, the map reflects the

proposed distribution of hominids: The approximate distribution and locations with dates of

known Neanderthal (Blue), Denisovan (Black), and Anatomical Modern Humans (Orange) fossil

remains based on: [101–116].

The genetic diversity of human lice has been widely studied based on the mitochondrial DNA.

However, we know little about the nuclear genetic diversity of this human parasite, particularly at

a worldwide scale. The first mitochondrial studies revealed the presence of several deeply

divergent mtDNA clades or haplogroups named A, B and C [2,3,14,17,21]. Haplogroup A is the

most common of the human louse haplogroups, has a worldwide distribution and has been found

in lice from human archeological remains in Israel from 2,000 years ago [22]. Haplogroup B has

been found in the Americas, Europe, Australia, North and South Africa [17,21,23,24], as well in

ancient human remains in Israel [22] and Peru [3]. Haplogroup C has a more geographically

restricted distribution and is only present in lice found in Nepal, Ethiopia, Republic of Congo,

Pakistan and Thailand [14,22,25]. Finally, recent studies have suggested the presence of three

more clades: D, E and F [24,26,27]. Clade D is sister to clade A found in Ethiopia, Zimbabwe and

Republic Democratic of Congo [24,26,28]. Clade E is sister to clade C and was detected in head

lice from Senegal and Mali [24]. Clade F appears to be specific to South America, as it has not

been reported elsewhere [27]. Head lice can be found in all the mitochondrial clades (A-F),

whereas clothing lice are only found in clades A and D [28]. Because of the haploid nature and

maternal inheritance of the mitochondrial DNA, events of hybridization cannot be detected with

this marker, thus new nuclear markers were developed [16] to provide a comprehensive

understanding of human louse genetic diversity and to assess admixture among human lice.

In our previous study using the same microsatellite markers with fewer samples [16], we

found that nuclear genetic diversity in human lice revealed continental differences and suggested

that the genetic pattern found in the Americas could be mirroring the (human host) colonization

of the New World represented by two major human settlements of the New World: the first

peopling of America and the European colonization after Columbus. In this new study we

expanded the sampling of lice to include more localities to deepen our understanding of human

louse nuclear genetic diversity while also assessing its correlation with mitochondrial diversity.

For this study, we examined 15 microsatellite loci and a mitochondrial gene (COX1) for nearly

300 individual head lice collected from 25 sites distributed throughout 10 main geographic

regions (Table 1, S1 Table, Fig 1). Furthermore, we included an new analysis by using novel

DIYABC-simulations that were built using both host and parasite data to define parameters and

models. In addition to providing new evolutionary knowledge about this human parasite, our

study could guide the development of new analyses in other host-parasite systems.

**Table 1. Microsatellite polymorphisms among human louse populations.** Groups reflects

their geographic origin and STRUCTURE clustering: Africa-Israel-Spain (AIS), South and

Southeast Asia (SAs), East Asia (EAs), Europe (E), North America (NA), Mesoamerica (MA),

Central America (CA), and South America (SAm).

| Geographic Region | Group | Country | Site | N | $Na$ | $H_O$ | $H_E$ | $F_{IS}$ | $P$-value |
|---|---|---|---|---|---|---|---|---|---|
| Africa | AIS | Rwanda | Rw | 1 | 1.533 | n/a | n/a | n/a | - |
| West Asia | AIS | Israel | Isr | 2 | 1.667 | n/a | n/a | n/a | - |
| Central Asia | SAs | Nepal | Np | 1 | 1.667 | n/a | n/a | n/a | - |
| South and | SAs | Thailand | Th | 1 | 1.333 | n/a | n/a | n/a | - |
| Southeast | SAs | Cambodia | Cam | 22 | 6.8 | 0.483 | 0.703 | 0.319 | *0.000* |
| Asia | SAs | Laos | Ls | 4 | 3.2 | n/a | n/a | n/a | - |
| East Asia | EAs | Mongolia | Mn | 10 | 5 | 0.524 | 0.706 | 0.267 | *0.005* |
| Europe | E | Turkey | Tur | 12 | 3 | 0.232 | 0.408 | 0.407 | *0.000* |
| | E | Croatia | Croa | 6 | 1.133 | n/a | n/a | n/a | - |
| | E | Norway | Nw | 1 | 1.067 | n/a | n/a | n/a | - |
| | E | Netherlands | Neth | 64 | 2.267 | 0.032 | 0.248 | 0.863 | *0.000* |
| | E | France | Fr | 3 | 1.133 | n/a | n/a | n/a | - |
| | E | United Kingdom | UK | 27 | 2.533 | 0.145 | 0.329 | 0.550 | *0.000* |
| | AIS | Spain | Spain | 3 | 2.067 | n/a | n/a | n/a | - |
| North | NA | USA | Oce | 25 | 2.8 | 0.083 | 0.352 | 0.793 | *0.000* |
| America | NA | USA | SF | 1 | 1 | n/a | n/a | n/a | - |
| | NA | USA | Martin | 1 | 1 | n/a | n/a | n/a | - |
| | NA | USA | WPB | 1 | 1.467 | n/a | n/a | n/a | - |
| | NA | USA | Nash | 18 | 2.467 | 0.037 | 0.311 | 0.864 | *0.000* |
| | NA | USA | MC | 22 | 3.867 | 0.173 | 0.452 | 0.587 | *0.000* |
| Meso | MA | Mexico | MX-1 | 19 | 3.13 | 0.208 | 0.396 | 0.481 | *0.000* |
| America | MA | Mexico | DF | 1 | 1 | n/a | n/a | n/a | - |
| Central America | CA | Honduras | Hon | 20 | 4.2 | 0.365 | 0.567 | 0.312 | *0.000* |
| South | SAm | Peru | Peru | 1 | 1.267 | n/a | n/a | n/a | - |
| America | SAm | Argentina | Arg | 9 | 3.2 | 0.349 | 0.482 | 0.231 | *0.012* |
| Total/Average | | | | 25 | 274 | 12 | 0.257 | 0.402 | 0.469 |

*(Total/Average row values: N=25, Na=274, then 12, 0.257, 0.402, 0.469, 0.000)*

107
108 $N$ = sampling size; $Na$ = Average number of alleles; $H_O$ = observed heterozygosity; $H_E$ =
109 expected heterozygosity; $F_{IS}$ = index of inbreeding of individuals resulting from the non-random
110 union of gametes within a subpopulation, positive value indicates heterozygote deficit; $P$
111 proportion of permutations leading to a $F_{IS}$ value larger or equal to the observed one. n/a
112 indicates values not estimated due to sample size smaller than 9 individuals.

# MATERIALS AND METHODS

## Ethics statement

The Institutional Review Board of the University of Florida exempted the study from review (Exemption of Protocol #2009-U-0422) and waived the need for written informed consent of the participants. This exemption is issued based on the United States Department of Health and Human Services (HHS) regulations. Specifically, HHS regulation 45 CFR part 46 applies to research activities involving human subjects. Because louse removal was voluntary and no information was recorded that would allow patients to be identified directly or through identifiers linked to them, the University of Florida waived the need for written informed consent from the participants.

## Sampling

Sampling details are provided in Fig 1, Table 1 and S1 Table. Briefly, 274 human head lice were collected from different individuals at 25 localities throughout the world covering 10 major geographic regions.

## DNA extraction

Lice were cut in half, placed into 1.5 ml Eppendorf tubes containing a cell lysis and proteinase K solution, and ground using a pestle. DNA was extracted from each individual louse using the Puregene Core Kit A (QIAGEN, Valencia, California) following manufacturer protocols with modifications. We used a NanoDrop 1000 spectrophotometer (Thermo Scientific) to determine the amount of DNA in each sample and then diluted them from their original concentrations to approximately 5-10 ng/µl.

## Multiple co-amplification: Multiplexes

A total of 15 microsatellite loci were amplified using four multiplexes as described in [16].

We used fluorescently labeled universal primers: M13 (5'-CACGACGTTGTAAAACGAC-3')

and CAG (5'-CAGTCGGGCGTCATCA-3'), each labeled with a unique fluorescent tag (e.g.

FAM, VIC, NED, PET; Applied Biosystems) to co-amplify multiple loci. Locus-specific primers

were modified by adding the matching 5′ universal primer sequence tails. The complete set of 15

loci was amplified in four separate 15 μL multiplex-PCR reactions containing 7.5 μL of 2X Master

Mix (Type-It Microsatellite PCR kit, Qiagen, Venlo, Netherlands), 0.015-0.15 μL of unlabelled

forward primer with tail (10 μM), 0.15-0.8 of unlabelled reverse primer (10 μM), 0.5 μL labelled

universal tail primer (10 μM), 1-2 μl of total genomic DNA (10-20 ng) and sdH$_2$0 added up to final

volume. Two thermal cycling profiles were used, both beginning with initial denaturation at 95°C

(5 min) and ending with a final extension of 72°C (40 min). The v1 (touch-down PCR) protocol

consisted of 10 cycles of 94°C (30 sec), 60°C/55°C (-0.5°C/cycle) for 45 sec, and 72°C (45 sec)

followed by 25 cycles of 94°C (30 sec), 55°C (45 sec), and 72°C (1 min). The v2 protocol

consisted of 35 cycles of 94°C (30 sec), 52°C (45 sec), and 72°C (45 sec). PCR products were

electrophoresed on 1.5% and 4% agarose gels stained with ethidium bromide and visualized under

ultraviolet light. Dilutions in dH$_2$0 from amplicons (1:100) were run on an ABI 3730xl 96-

capillary sequencer using GeneScan 600 LIZ as an internal size standard (Applied Biosystems) at

the University of Florida biotechnology facility (ICBR). Microsatellite genotypes were scored

using the GeneMarker version 1.60 software (SoftGenetics, LLC).

## Mitochondrial PCR amplification and sequencing

A polymerase chain reaction was performed to amplify the mitochondrial gene cytochrome *c*

oxidase 1 (COX1) using the primers H7005 and L6625 [9] as described in [2]. PCRs consisted of

25 μL total volume including 10 μL of MasterMix (5 PRIME), 1 μL of each primer, 2-4 μL of total

genomic DNA, and water.  The thermal cycling profile began with an initial denaturation at 94°C

(10 min) followed by 10 cycles of 94°C (1 min), 48°C (1 min), and 65°C (2 min) (decreased by

0.5°C per cycle).  This was followed by 35 cycles of 94°C (1 min), 52°C (1 min), and 65°C (2

min) and then a final extension of 65°C (10 min).  Amplified fragments were purified using

ExoSAP-IT (USB Corporation, Cleveland, Ohio).  Sequencing was performed at the University of

Florida DNA Sequencing Core Laboratory (ICBR, Gainesville, Florida) using standard fluorescent

cycle-sequencing PCR reactions (ABI Prism Big Dye terminator chemistry, Applied Biosystems).

***Sequence editing, alignment, and haplotype reconstruction***

The forward and reverse sequences were edited and aligned using Sequencher 4.5 (Gene Codes

Corporation, Ann Arbor, Michigan) with base calling confirmed by eye.  Consensus sequences

were generated for each sample using both forward and reverse sequences.  These new louse COX1

sequences were aligned along with previous sequences from Ascunce et al. (2013) [17].  A final

alignment of 379 bp was used to construct a matrix of pairwise differences using uncorrected *p*-

distances (proportion of nucleotide sites at which two sequences being compared are different)

among COX1 unique haplotypes was calculated with MEGA 4 [29].  The genetic relationships

were estimated by constructing neighbor-joining (NJ) trees [30] using PAUP* 4.0b10 [31].

Bootstrapping was performed using 100 pseudo-replications of the data set.

# Analyses of genetic diversity and population structure among collection sites

We used the software Arlequin 3.5.1.2 [32] to estimate population genetic diversity values.

First, for all sites regardless of the number of lice per site, we determined the number of alleles per

locus.  Then, for those 11 sites with sample sizes exceeding 8 lice per site, we also estimated

observed and expected heterozygosity ($H_O$ and $H_E$), and the mean $F_{IS}$ over loci per population with

significance tested using 1,023 random permutations.  For those same 11 sites with sample sizes

exceeding 8 lice per site, population pairwise $F_{ST}$ values were estimated using Arlequin 3.5.22

[33].  Significance values for pairwise $F_{ST}$ values were estimated using the permutation procedure

with 1,000 permutations.

To assess the distribution of the human louse genetic diversity among world populations

through clustering of similar genotypes, we used three different methods as they have different

assumptions.  First, we used STRUCTURE [34], a model-based Bayesian clustering method, that

assumes Hardy–Weinberg and linkage equilibrium among loci.  Second, we also used two

multivariate methods, which do not carry these assumptions and could visualize potential cryptic

structure, including PCoA (Principle Coordinates Analysis) in GenAlEx 6.5 [35,36] and

Discriminant Analysis of Principal Components (DAPC) [37].

**Bayesian clustering analyses**

The Bayesian clustering approach implemented in STRUCTURE 2.3.4 [34] uses the individual

multi-locus genotypic data to evaluate models assuming different numbers of genetic clusters ($K$)

based on the posterior probabilities given the data and model.  Based on the individual multilocus

genotypes and the allele frequencies estimated for the reconstructed clusters, each individual's

genome was probabilistically partitioned into membership fractions ($q$) (ancestry) in each cluster.

Admixed ancestry is modeled by assuming that an individual $i$ has inherited some fraction ($q$) of

its genome from ancestors in population $k$ [34].  Hybrids can then be inferred when observing an

individual being intermediate of some degree between two or more clusters in terms of the $q$-value.

For example, individuals with $q = 0.5$ would be likely first-generation hybrids.  We identify

possible hybrids based on $q$-values threshold of 0.2 as suggested in VäHä and Primmer (2005)

[38], where threshold $q$-value of 0.2 means that individuals with $q$-value between 0 and < 0.20 or >

0.8 and 1 are classified as non-admixed and any individual with $q$-value between 0.2 and 0.8 are

classified as a hybrids. All simulations used 50,000 Markov chain Monte Carlo (MCMC)

generations in the burn-in phase and 100,000 generations in the data collection phase. Ten

independent runs were performed using default parameters for each $K$, to ensure equilibration

during burn-in and consistency in estimation of the posterior probabilities. Selection of the number

of distinct clusters was based on the evaluation of the $\Delta K$ statistic [39] using the online tool

STRUCTURE HARVESTER [40]. The ten STRUCTURE runs at each $K$ produced nearly

identical individual membership coefficients. The run with the highest likelihood of the data given

the parameter values for the predominant clustering pattern (i.e. the mode) at each $K$ was used for

plotting with DISTRUCT [41].

**Multivariate analysis using Principal Coordinates Analysis (PCoA)**

This multivariate analysis was conducted using pairwise genetic difference between

individuals calculated in GenAlEx 6.5 [35,36] to validate and further define genetic clusters found

using STRUCTURE.

**Multivariate analysis using Discriminant Analysis of Principal Components**

**(DAPC) [37]**

This analysis was conducted using the '*adegenet*' package 2.1.3 [42], in the R software

environment version 3.6.3 [43]. For these analyses we only included 175 lice from which we were

able to obtain both microsatellite genotype and mitochondrial haplogroup sequence. An additional

DAPC included the same louse samples, however we considered four groups depending on nuclear

cluster and haplogroup membership.

## Allelic richness within each main nuclear STRUCTURE genetic cluster

We estimated the allelic richness withing each main nuclear STRUCTURE genetic cluster using the rarefaction method implemented in the computer program ADZE (Allelic Diversity Analyzer) version 1.0 [44]. Because allelic richness measurements considered sample size, the rarefaction method allows the estimation of allelic richness for different random subsamples of size "g" from the populations [45,46].

## Analyses based on STRUCTURE genetic clusters and mitochondrial clades

To gain a better understanding of the geographic distribution of the nuclear and mitochondrial diversity, we conducted a set of Kruskal–Wallis nonparametric analysis of variance tests [47]. We included the same 175 lice from which we were able to obtain both microsatellite genotype and mitochondrial haplogroup sequence data and the same eight geographic regions as described above. Calculated medians were compared and separated at $P < 0.05$ [48]. We compared the Observed vs. Expected Frequency $X^2$ test to evaluate the fit of the data (frequencies) to any arbitrary set of expected frequencies among the eight geographic regions. All statistical tests were performed with $\alpha=0.05$ for significance of statistical tests.

Analysis of molecular variance (AMOVA) considering the number of different microsatellite alleles ($F_{ST}$ based) was used to investigate the partitioning of genetic variation between the two nuclear clusters: cI and cII using Arlequin 3.5 [32]. Statistical significance of differentiation at each level was assessed by means of 1,023 permutations.

The historical migration patterns among the genetic clusters identified using the methods

describe above were estimated in Migrate-n 3.3.0 using the maximum likelihood approach [49,50].

Migrate employed estimates of gene genealogies to sample areas of the coalescence space with

highest likelihood.  The random number of seed and starting values of $\Theta$ and $4N_em$ (where $N_em$ is

the indirect measure of gene flow) were based on $F_{ST}$ as starting parameters and we used a

Brownian mutation model.  The data was analyzed using 10 short ($10^4$ MCMC steps) and 3 long

chains ($10^5$ MCMC steps), together with adaptive heating to assure convergence.

## Comparison of demographic models

We further inferred the demographic history of human lice, based on microsatellite data, using

the approximate Bayesian computation (ABC) approach [51], implemented in the program

DIYABC, v2.0 [52].  We tested different evolutionary scenarios in which each louse genetic

cluster evolves on different human host populations and diverge in tandem with their hosts.  A

total of seven different evolutionary scenarios were considered with population size estimates and

divergence times based on inferences done using data on human evolution and louse life history

traits references (Table 2, Fig 2).  The first three scenarios are directly testing Reed et al. (2004)

[2] hypothesis of ancient hybridization between modern and archaic humans.  The scenarios 1 and

2 were trying to unravel the retention of ancient polymorphisms on the hybrid lice, where the main

nuclear genetic clusters were derived from lice that parasitized the heads of Neandertals (scenario

1), or the heads of anatomical modern humans (AMH) (scenario 2).  In scenario 3, the model tested

hybridization events between lice parasitizing Neanderthals and AMH during periods of

contemporaneity between the two hosts.  The last four scenarios (scenario 4, 5, 6 and 7) do not

require ancient hominids, and are based solely on hybridization events within AMH during recent

time periods. In scenario 4, we tested the hypothesis that the hybrid lice were the product of

hybridization between lice from the Native Americans and the first Europeans during the European

colonization of the Americas 500 years ago. In scenario 5, we tested the hybridization occurring

in the last 100 to 40 years around the World Wars I and II. Scenario 6 reflects the beginning of

the globalization from 1980's to ten years ago, while scenario 7 refers to very recent events in the

last 10 years.

**Table 2. Prior distributions of historical and demographic parameters for the demographic scenarios.** Between [ ] are the host

values obtained from anthropological references.  For estimates of times, those values for louse represents louse generations (g), and

the values between [] for hominids are in years (y).  Louse values are estimate considering an infection of 20 lice in 30% of hominin

population.

| Details/Historic Events | Parameter | Distribution | Minimum-Maximum | References |
|---|---|---|---|---|
| *Effective population* | | | | |
| Louse Ancestral Population | $N_A$ | Uniform | 1,800-36,000 | |
| [Ancestral hominids] | | | [300-6,000] | |
| Louse cluster I | $N_1$ | Uniform | 1,800-36,000 | |
| [AMH] | | | [300-6,000] | |
| Louse cluster II | $N_2$ | Uniform | 1,800-36,000 | |
| [Neanderthals] | | | [300-6,000] | [113] |
| Louse Hybrids | $N_3$ | Uniform | 270-5,400 | 15% hybrids from current study |
| [Hybrids between AMH and Neanderthals] | | | [?] | |
| *Time Events\** | | | | |
| Time of split clusters I and II | $t_S$ | Uniform | 1,200,000 - 16,000,000g | |
| [Split between AMH and Neanderthals] | | | [600,000-800,000y] | [113] |
| T1 ancestral polymorphism from cII | T1 | Uniform | 2,400,000 - 2,800,000g | [114] |
| [Presence of Neanderthals in Mongolia] | | | [120,000 - 140,000y] | [116] |
| T2 ancestral polymorphism from cI | T2 | Uniform | 1,000,000 - 1,600,000g | |
| [Presence of AMH in Middle East] | | | [50,000-80,000y] | [105] |
| Time of ancestral admixture | $t_{aA}$ | Uniform | 1,000,000 - 1,600,000g | |
| [AMH-Neanderthals in sympatry] | | | [50,000-80,000y] | [20] |
| *Times of Recent Admixture* | | | | |
| Time of admixture E1 | $T_{aE1}$ | Uniform | 2,000 – 20,000g | |
| [European colonization] | | | [100 – 1,000y] | |
| Time of admixture E2 | $T_{aE2}$ | Uniform | 800 – 2000g | |
| [Migration after World Wars] | | | [40 – 100y] | |
| Time of admixture E1 | $T_{aG}$ | Uniform | 200 – 800g | |
| [Initial Globalization] | | | [10 – 40y] | |
| Time of admixture E1 | $T_{aC}$ | Uniform | 20 – 200g | |
| [Current time -t sampling] | | | [1 – 10y] | |

| | | | |
|---|---|---|---|
| *Admixture rate* | | | |
| Rate of admixture | ar | Uniform | 0.001-0.999 |
| *Mutation rate* | | | |
| Tetranucleotide microsatellite mutation rate | Um1 | Loguniform | $10^{-9}-10^{-3}$ |
| Trinucleotide microsatellite mutation rate | Um2 | Loguniform | $10^{-7}-10^{-3}$ |
| Dinucleotide microsatellite mutation rate | Um3 | Loguniform | $10^{-6}-10^{-2}$ |

*Time parameters for lice are expressed as the number of generations of lice back in time assuming a generation time of 18 days [15].

The values for hominids refer to years

**Fig 2. Evolutionary scenarios.** A total of seven different scenarios were compared using the

approach from the DIY-ABC program.

For each scenario, we considered three levels of parasitism: from a low level of parasitism

where each head human would have 10 lice and only 10% of the heads would be infected (S2

Table); an intermediate level, where each head would have 20 lice and only 30% of the population

is infested (Table 2); and a high level of lice, where there are 50 lice per head in 50% of the

population (S3 Table) [53]. The generation time of *P. humanus* was considered as 18 days [15].

All microsatellites used the default generalized stepwise mutation model and were assigned a

log uniform distribution with mean mutation rates between $10^{-9}$–$10^{-2}$. The standard mutation rate

for microsatellites is considered to be between $10^{-6}$–$10^{-2}$ per locus per generation [54]. However,

due to the possibility of different repeat lengths having different mutation rates [55] for

microsatellites, tetra-, tri- and di- nucleotide repeats were modeled separately. Thus, preliminary

runs were carried out in order to optimize mutation rates to account for other factors [56] following

Bertorelle et al. [57]. A range in allele size of 40 contiguous allelic states was used, as

recommended by the program developers (A. Estoup, personal communication).

Summary statistics calculated for microsatellite loci were as follows: Mean number of alleles,

Mean genetic diversity, and Mean size variance, and Mean $F_{ST}$. The selection of summary

statistics was empirical based on preliminary tests where these summary statistics were used and

able to recover the models and parameters following Bertorelle et al. [57]. Through these

preliminary tests the above summary statistics demonstrated their suitability for our data.

We performed about 714,000 simulations using prior parameters for each scenario, thereby

producing a reference table comprising 5,000,000 simulated data sets. First, we pre-evaluated the

fit of observed values to prior distributions of scenarios, by conducting a Principal Components

Analysis (PCA) using data sets simulated with the prior distributions of parameters and the

observed data as well as data sets from the posterior predictive distribution to check the suitability

of the model (scenarios and prior parameters). The posterior probability of each scenario was

assessed by performing direct approach and a polychotomous weighted logistic regression that

estimates the difference between simulated (from the reference table) and observed data sets

[58,59] after linear discriminant analysis on summary statistics [52]. Only the posterior

probabilities for the 1% (50,000) of simulated data sets closest to the observed data were retained.

Confidence in choice of a scenario was estimated as the proportion of times the selected scenario

did not possess the highest posterior probability compared to competing scenarios for the 500

simulated data sets (type I error). Confidence in scenario choice was further tested including the

estimation of the prior error rate, which is calculated as the probability of choosing an incorrect

model when drawing model index and parameter values into priors (the probability of selecting

the chosen scenario when it is not correct, a type-II error). After scenario choice, we proceeded to

parameter inference estimated from the modes and 95% confidence intervals (CI) of their posterior

distributions.

**RESULTS**

**Nuclear Genetic Diversity**

The 274 human louse specimens comprising 25 geographic sites through 10 worldwide

geographic regions revealed high microsatellite diversity with overall five to 20 alleles per locus

and an average of 12 alleles per locus (Table 1). In the 11 sites with more than 8 lice analyzed,

$H_O$ values ranged from 0.032 to 0.524 (average of 0.257) and $H_E$ values ranged from 0.248 to

0.706 (average of 0.402), reflecting a general patter of deficit of heterozygotes (Table 1).

Furthermore, in all 11 louse populations, genotype proportions deviated significantly from Hardy-

Weinberg expectations with values of $F_{IS}$ significantly positive and an overall $F_{IS}$ value of 0.469

($P < 0.00001$) (Table 1).  Overall, there were high values of genetic differentiation among the 11

sites, with $F_{ST}$ values of up to 0.663 between the sites: Mongolia (Mn) and Netherlands (Neth) (S4

Table).  The lowest $F_{ST}$ of 0.053 was found between sites in New York (Oce) and Tennessee

(Nash).

## Mitochondrial diversity

We also obtained the COX1 sequence data from 175 head lice spanning 18 sampling locations

throughout nine geographic regions (Table 3).  One hundred four (104) lice belonged to

haplogroup A and 71 to haplogroup B (Table 3).  As expected based on previous studies [17,21,23],

haplogroup A shows a worldwide distribution including Africa, Asia, Europe and the Americas.

For our sampling, haplogroup B has a narrower distribution and it is found only in lice from Europe,

North and Central America (Table 3).

**Table 3. Nuclear and mitochondrial louse genetic diversity.** Geographic distribution of 175 human lice including combined information of their mitochondrial DNA haplogroups based the COX1 gene and their assignment to each louse major nuclear genetic clusters described in this study.

| Geographic Region | Group | Country | Site | mtA+clusterI | mtB+clusterI | mtA+clusterII | mtB+clusterII |
|---|---|---|---|---|---|---|---|
| Africa | AIS | Rwuanda | Rw | 1 | | | |
| South Asia | SAs | Nepal | Np | 1 | | | |
| Southeast Asia | SAs | Cambodia | Cam | 21 | | | |
| | | Laos | Ls | 1 | | | |
| East Asia | EAs | Mongolia | Mn | 6 | | | |
| Europe | E | Turkey | Tur | | | | 7 |
| | E | Croatia | Croa | | | | 5 |
| | E | Norway | Nw | | | 1 | |
| | E | Netherlands | Neth | | | 49 | 5 |
| | E | France | Fr | | | 2 | |
| | E | U. Kingdom | UK | | | | 26 |
| North America | NA | USA | Oce | | | 12 | 4 |
| | NA | USA | SF | | | 1 | |
| | NA | USA | Nash | | | | 3 |
| | NA | USA | MC | | | 1 | 9 |
| Mesoamerica | MA | Mexico | MX | | | 6 | |
| Central America | CA | Honduras | Hon | | 12 | | |
| South America | SAm | Argentina | Arg | | | 2 | |
| Total | | | | 30 | 12 | 74 | 59 |

## Nuclear genetic clusters at the worldwide scale

All clustering methods: STRUCTURE, PCoA and DAPC were concordant in identifying the

distinct nuclear genetic clusters: Cluster I (cI) with subclusters Ia and Ib and Cluster II (cII)

including subclusters IIa and IIb (Fig 1; S1 Fig).  It is striking, the difference in their geographic

distribution, where cI defined by head lice from Africa, Israel, Asia and the Americas compass a

more worldwide distribution, while cII composed of head lice mainly Europe and the Americas

reflects a more restricted geographic distribution.  While most of the genetic diversity of human

louse populations from around the world seems to be highly structured geographically,

approximately 12% of the lice we sampled (33 of 274 lice) appear to be hybrids (lice with $q$-values

less than 0.8), with 75% (25 lice) of the hybrids found in the New World.

In the STRUCTURE analysis, after a signal for $K = 2$ ($\Delta K = 858.853$, S2 Fig), increasing $K$

values in the STRUCTURE plots revealed geographically localized structure.  At $K = 3$, a new

cluster formed within Europe from the original cII and it is composed of lice primarily from the

Netherlands forming the subcluster cIIb (light blue color).  When $K$ was increased to 4 a population

from Honduras and other parts of the Americas emerged from cluster I as the Ib subcluster, with

the other samples remaining in subcluster Ia.  This Honduras (Ib subcluster) might be indicative

of Native American louse lineage.

The Principal Coordinate Analysis (PCoA) supported the two main nuclear genetic clusters

and subclusters identified by STRUCTURE, with putative hybrid lice not assigned to either cluster,

rather were placed between them (Fig 3).  Finally, the eigenvalues obtained in the DAPC analysis

showed that the genetic structure was captured mainly by the first principal component

representing cI (to the right) and cII (to the left) (S1 Fig).

**Fig 3. Human louse worldwide nuclear genetic diversity**. A) Genetic clusters inferred from

STRUCTURE simulations at $K = 2$: Cluster I (cI, Orange) and Cluster II (cII, Blue), $K = 3$ cI and

cIIa-cIIb, and $K = 4$ cIa-cIb and cIIa-cIIb.  In the bar plot, each louse is represented by a single

vertical line and the length of each color segment represents the proportion of membership ($Q$) to

each cluster. B) Distributions of points in the first two dimensions resulting from Principal

Coordinates Analysis (PCoA) conducted using pairwise genetic distance comparisons.  Each site

was color-coded to match STRUCTURE results (e.g. sites in Asia were color-coded in orange,

and sites in Europe were color-coded in blue) codes correspond to geographic origin and

abbreviations as defined in Table 1.

Additionally, we conducted further analysis with all the dataset and excluding hybrids as

identified through STRUCTURE $q$-value.  Thus, when using the reduced dataset only non-

admixed individuals with shared ancestry $q$-values larger than 0.8 as suggested by [38], the

STRUCTURE analysis reflected similar results identifying cI and cII, with their corresponding

subclusters at $K = 4$ (S3 Fig).

## Worldwide cluster I

All lice that belong to the cI ($q$-values larger than 0.5) were included in a second STRUCTURE

simulations to assess the substructure within this cluster.  At $K = 2$, this figure strikingly showed

the presence of two clusters: one composed of lice from the Old World (Africa, Asia and Europe),

and the other of lice from the New World (Americas) (S4 Fig).  The further split of the genetic

clusters during the increase of defined $K$ in the STRUCTURE analysis showed at $K = 3$ the

emergence of an Asian cluster except for most of the lice from Mongolia that remains within the

Old Word cluster, however at $K = 4$, a subset of those lice from Mongolia formed a unique cluster

(S4 Fig).

## Geographically restricted cluster II

Considering our current sampling, cII has narrower geographic distribution than the cI and it is limited mostly to Europe and the Americas (Fig 3). This cluster showed a substructure at $K = 2$ where most of the lice from the Netherlands plus few others formed a distinct cluster (light blue in plot) (S5 Fig). Both these individual cluster analyses for cI and cII highlight the geographic structure observed among louse populations.

## Allelic richness analysis per main genetic cluster

We estimated the allelic richness per main non-admixed clusters I and II (without hybrids) using the rarefaction method implemented in the computer program ADZE (Allelic Diversity Analyzer) Version 1.0 [44]. Because allelic richness measurements considered sample size, the rarefaction method allows the estimation of allelic richness for different random subsamples of size "g" from the populations [45,46]. In addition, we conducted the same analysis for each cluster separately using the sub-clusters defined by STRUCTURE.

Allelic richness per non-admixed cluster was higher in cluster I than II (S6 Fig). This pattern reflects very different evolutionary histories for the two clusters. While high allelic richness is consistent with a large effective population size, it could also be concordant with greater genetic structure. It is noteworthy also that the clusters I and II differ in sample size (80 lice in the cI versus 174 in cII).

## Statistical analysis of genetic clusters and mitochondrial clades

The combined analysis of the nuclear cluster membership frequency and the mitochondrial haplogroup were plotted using a Correspondence Analysis (CA) and showed the presence of four well defined groups (S7 Fig). The dimensions 1 and 2 represented the largest amount of inertia or

variance representing a measure of deviation of independence. Each one of the dimensions possess

an eigenvalue that represents its relative importance and how much of the variance is explained.

In our study, one dimension represents 58.7 % of the inertia and two dimensions represents 99.6%

of the inertia. Of this, dimension 1 differentiated both nuclear genetic clusters: cI and cII; while

dimension 2 differentiated the mitochondrial haplogroups. One group corresponds to lice of

nuclear cluster II-haplogroup A and included lice from Europe, North America, Mesoamerica and

South America. The second group is also nuclear cluster II-haplogroup B from the same regions,

except South America. The third group is composed of individuals of cluster I-haplogroup A:

South Asia, Africa-Israel-Spain, some Europe, while the last group belonged to cluster I with

haplogroup B and were found in Central America (S7 Fig). Furthermore, multiple comparisons

(non-parametric Kruskal–Wallis test) indicated that the distribution of the individuals assigned to

the nuclear genetic clusters I and II were not distributed randomly among the eight geographic

regions (H = 103.55, $P < 0.0001$). The cluster membership median values were significantly

different among: South Asia (SAs), East Asia (EA), Africa-Israel-Spain (AIS), Central America

(CA), and Europe (E) ($P < 0.05$). On the other hand, cluster membership medians values for North

America (NA), South America (Sam), and Mesoamerica (MA) were not significantly different.

Both nuclear genetic assignment and haplogroup frequencies differed significantly among the

eight geographic regions (nuclear cluster assignment $X^2 = 175$, $P < 0.0001$; haplogroup frequencies

$X^2 = 42$, $P < 0.0001$).

## AMOVA by nuclear clusters and mitochondrial clades

431  Hierarchical analyses of molecular variance (AMOVA) employed to explore the distribution

432  of nuclear genetic variation considering the two nuclear genetic clusters: cI and cII revealed

433  significant genetic differentiation between them ($\Phi_{ST} = 0.33$; $P < 0.00$). When examining only the

434  non-admixed individuals (those with at least 0.8 of coefficient of ancestry values) the

435  differentiation between cI and cII increased ($\Phi_{ST} = 0.4$; $P < 0.00$).

## Migration patterns considering nuclear genetic clusters and mitochondrial clades

The historical migration pattern was estimated in Migrate-n 3.3.0 using maximum likelihood

inference and the Brownian motion mutation model [49,50]. Given that previous studies have

suggested a different evolutionary history between the mitochondrial haplogroups A and B, we

evaluated a full migration model with four populations defined by both the nuclear clusters inferred

from STRUCTURE and the mitochondrial haplogroups: cI − mtDNA A; cI − mtDNA B; cII −

mtDNA A and cII − mtDNA B. Migration rates between these four populations varied. In general,

migration rates within nuclear clusters were high, contrasting with low migration rates within

mitochondrial haplogroups (S8 Fig). For example, we observed the highest migration rates within

cluster II (cluster II − mtDNA A and cluster II − mtDNA B) and some of the lowest migration rates

between cluster I − mtDNA A and cluster II − mtDNA A. Because STRUCTURE analysis

indicated the effect of geography on the genetic structure of human lice, further studies including

more samples from each sampling site are needed to further understand migration patterns among

geographic human louse populations.

## Demographic history

We inferred relationships between sets of individuals based on the STRUCTURE genetic

clusters using approximate Bayesian computation (ABC) incorporated in the software DIYABC,

v2.0 [52]. Individuals were assigned to the genotype pool based on their $q$ values, where

individuals with $q$-value between 0 and < 0.20 or > 0.8 and 1 are classified as non-admixed and

any individual with *q*-value between 0.2 and 0.8 are classified as hybrids. We tested seven

different evolutionary scenarios in which each louse genetic cluster evolves on different human

host populations and diverges in tandem with their hosts. The scenarios reflect different

hypothesizes that followed human evolutionary demographic events from the admixture between

Neanderthals and AMH to current human movement around the globe during globalization (Table

2, Fig 2). We repeated the same 7 models using different levels of louse parasitism. Comparison

of the seven scenarios showed that the most strongly supported scenario was scenario 7 suggesting

a very recent admixture between the two nuclear clusters with a logistic posterior probability of

ca. 0.90 (Fig 4, Table 4). The three most strongly supported scenarios 5, 6 and 7, are topologically

similar each involving admixture between the two nuclear clusters at different time paralleling

different human host colonization and migration events from Europe into the New World.

**Fig 4. Posterior probabilities for each of the seven demographic scenarios.** Graphs of

posterior probabilities based on DIY-ABC analysis comparing the seven demographic scenarios

when considering intermedia louse infestation level where 30% of the hominin population have

lice with 20 lice per head. Each scenario was color-coded as follow Scenario 1 – green, Scenario

2 – red, Scenario 3 – light blue, Scenario 4 – pink, Scenario 5 – yellow, Scenario 6 – black,

Scenario 7 – gray.

**Table 4. Posterior probabilities for the seven demographic scenarios shown in Fig 2.** Probabilities obtained through the

logarithmic method (with 95% confidence intervals in brackets) for the seven scenarios inferred from DIY-ABC analysis.  For each

scenario three parasitism level were considered where 10% indicates that 10% of the hominin population have lice with 10 lice per

head, 30% and 50% indicate that 30% and 50% of the hominin population have 20 and 50 lice per head, respectively.  For each

comparison, the selected scenario (bold entry in shaded cell) was the one with the highest probability value. Confidence in scenario

choice, shown under probability values, was evaluated by estimating the type I error rate from 1000 randomly constructed simulated

data sets, for scenarios 5 and 7, respectively.

| Scenarios | Parasitism Level | | |
|---|---|---|---|
| | 10% | 30% | 50% |
| 1: Retain Polymorphisms from cI | 0.0000 [0.0000,0.0000] | 0.0000 [0.0000,0.0715] | 0.0000 [0.0000,0.1130] |
| 2: Retain Polymorphisms from cII | 0.0000 [0.0000,0.0000] | 0.0000 [0.0000,0.0715] | 0.0000 [0.0000,0.1130] |
| 3: Ancestral admixture cI and cII | 0.0000 [0.0000,0.0000] | 0.0000 [0.0000,0.0715] | 0.0000 [0.0000,0.1130] |
| 4: European colonization of the New World | 0.0000 [0.0000,0.0000] | 0.0001 [0.0000,0.0716] | 0.0560 [0.0000,0.1522] |
| 5: European migration to the New World after WWs | 0.0000 [0.0000,0.0000] | 0.0071 [0.0000,0.0757] | **0.5204 [0.4477,0.5930]** |
| 6: Globalization | 0.0026 [0.0011,0.0041] | 0.3792 [0.2622,0.4963] | 0.2754 [0.2042,0.3467] |
| 7: Current time - sampling | **0.9974 [0.9959,0.9989]** | **0.6136 [0.4959,0.7312]** | 0.1482 [0.0095,0.2869] |
| | 0.335 | 0.317 | 0.372 |

# DISCUSSION

In this study, we found that the nuclear microsatellite data from lice were grouped in two distinct genetic clusters: cI and cII. Cluster I had a worldwide distribution whereas cII was restricted to Europe and the Americas (Fig 5). Remarkably, the two nuclear clusters revealed a significant genetic differentiation between them ($\Phi_{ST} = 0.33$; $P < 0.00$) despite having geographic overlap in either Europe or the Americas. In the Structure analysis at $K = 4$ each of the two Clusters further revealed the subclusters: Ia-Ib and IIa-IIb, which were sorted based on their geographical origin (i.e., continents). These results are consistent with our previous work using fewer samples where we showed that human louse populations are genetically structured based on geography [16]. In that previous study, we assessed louse microsatellite genetic diversity of a small sampling from four main geographic regions (North America, Central America, Asia, and Europe) that showed that the head louse populations were strongly structured geographically, where one cluster included head lice from North America and Europe, a second cluster encompassed only head lice from Central America, with a third Asian cluster [16]. This current study with a larger dataset highlighted a similar pattern showing a close relationship between Central America (cIa) and Asia (cIb) subclusters suggesting that the Central America cluster is probably from Native American origin since East Asia has been suggested as a potential source population for the first people of the Americas [60–62] (Fig 5). Thus, microsatellite data support the idea that current populations of human lice in the Americas retain human louse genetic diversity brought by the first people [16]. This work adds to our previous studies and shows that the study of the genetic diversity of human lice provides insights to better understand patterns of human migrations worldwide.

**Fig 5. Proposed global co-migration human lice and humans.** Top: Map depicting the geographic distribution of the nuclear genetic diversity among the human louse populations

included in this study. Colored circles on map indicate collecting sites, with the color of each

circle corresponding to the majority nuclear genetic cluster to which sampled individuals were

assigned. Sites with admixed lice are indicated with patterned circles including colors of the two

major genetic clusters at that site. Thick grey arrows indicate proposed migrations of anatomically

modern humans out of Africa into Europe, Asia and the Americas, as well as the more recent

European colonization of the New World. Colored arrows represent hypothetical human louse co-

migrations. The bottom panel is the STRUCTURE plot from Fig 3A corresponding to the

assignment of 274 lice from 25 geographical sites at $K = 4$ (Table 1).

Other human parasites have also shed light into different aspects of human evolution [11–13].

For example, the evolutionary history of the bacterial parasite, *Helicobacter pylori*, which infects

the stomach of most humans, reflected major events in human migrations such as the out of Africa

expansion [63,64], the colonization of Austronesian speaking region [65], and the settlement of

the Americas by Asians across the Bering Strait [66]. However, microorganisms such as bacteria

and viruses often reproduce clonally, increasing the probability of observing a selective sweep

such that the parasite phylogeny is likely to reflect the selection history of some genes rather than

of the host tree. On the other hand, macroparasites (such as parasitic arthropods) often reproduce

sexually and may track host evolutionary history more accurately because selective sweeps affect

only closely linked loci in their genomes [67]. In cases where obligate parasites coevolve with

their hosts, parasite phylogenies will, to some extent, reflect host evolutionary history [8,9]. For

this reason, the use of macroparasites such as pinworms [68] and lice [2,3] has become increasingly

common in the study of human evolution. In fact, the human louse, *Pediculus humanus*, is one of

the most well studied human parasites [1–6,14–19,21–28].

A key feature of lice is that human lice can disperse from a host to another at any motile stage

(nymphs and adults), however adults are most likely to transfer from one head to another [69].

Lice can be transferred via direct body contact, for example when people fight, live in group and

their heads are in contact [70].  Additionally, lice could be transferred via indirect contact, like

when an infected person shares their hat and combs with other person, there could be a possibility

that live lice stayed on the personal staff and might be transferred to the other person head [71].

In addition, because parasites have larger population sizes and faster rates of molecular evolution

compared to their hosts, population genetic patterns of parasites can reveal life history traits of

their host such as host dispersal, interspecies interactions and population structure that might not

be evident from host data alone or not yet discernible in the host's genome [9,72,73].  Differences

between parasite and host intrinsic life-history traits may lead to incongruities in patterns of

population structure between hosts and their parasites [74–78].  Several types of parasites are

known to demonstrate this phenomenon, including flies from bats [78], helminths from pikas [79],

and feather lice from Galápagos hawks [80].  For example, the comparative evolutionary histories

between Arctic Galliformes and their parasitic lice suggest consistent movement of lice among

different host species and populations in Alaska [81].  While the bird hosts present some level of

population structure and/or isolation by distance, no structure was detected within either genus of

louse on these hosts [81].  These incongruent patterns in host and parasite genetic structure suggest

there is still contact or at least distributional overlaps in host populations which are not perceptible

from the host data alone.  In our current study, we show the presence of two louse distinct nuclear

genetic clusters: I and II.  These could be the result of the retention of ancient polymorphisms due

to incomplete lineage sorting (ILS) or admixture.  To test if louse current genetic diversity is the

result of ILS or admixture we used the approximate Bayesian computation (ABC) method as

implemented in the DIY-ABC package [52]. We performed ABC analyses for seven scenarios

and for three different level of parasitism (Table 2, 4, Figs 2, 4, S9 Fig). These analyses suggested

a scenario of recent admixture between nuclear Clusters I and as observed in the STRUCTURE

analysis, where most of the hybrids are being observed in the New World. While most of the

genetic diversity in human louse population from around the world seems to be highly

geographically structured, there are a total of 33 hybrids out of 274 lice analyzed reaching 12% of

our current sampling. However, 25 of those hybrids (75%) are found in the New World, which

may be mirroring human colonization history and reflecting a zone of secondary contact. All the

three scenarios considering recent admixture events from European migration after World Wars

(scenario 5), initial globalization (scenario 6) and current times at sampling (scenario 7) were the

most supported (Table 2, 4, Fig 4, S9 Fig).

A new and puzzling insight from this study was the finding of such low prevalence of hybrid

lice and that these hybrids are geographically concentrated in the New World. This pattern could

be reflecting the human colonization history from Europe to the New World as mentioned above.

However, if these colonization events have been occurring for many louse generations, why are

there so few hybrids? A tentative explanation could be associated to sampling issues as a large

number of samples came from the Americas (43%) (Table 1). Thus, future studies on genetic

diversity of human lice should consider larger sampling if possible. Despite that it is beyond the

scope of the present paper to investigate the reason behind the lack of hybrids, we hypothesize that

this pattern could be indicating some sort of genetic isolation between cI and cII such that the

hybrids are not able to persist for many generations within the human louse populations. Thus, it

is worthwhile to enumerate the potential causes for future researcher.

One potential mechanism of reproductive isolation may involve the effect of endosymbiont on

host biology.  Endosymbionts could also alter host biology in diverse ways, including the induction

of reproductive manipulations, such as feminization, parthenogenesis, male killing and sperm–egg

incompatibility, which could lead to reproductive incompatibility [82].  *Wolbachia* is a common

intracellular bacterial symbiont of arthropods and filarial nematodes and it has been shown to

infect human lice [83–85].  When populations of insect host that contain different strains of

*Wolbachia* interbreed, or when one population has no *Wolbachia* at all and the other population

does have *Wolbachia*, the interbreeding matings ("hybrids") result in reduced fitness or the

complete absence of progeny.  If this process were to occur in human lice, it could explain the low

number of hybrids; however, because *Wolbachia* endosymbiont is transmitted together with the

mitochondrial genome through the egg cytoplasm, endosymbionts and mtDNA should reflect a

similar history.   In our study, mitochondrial lineages were grouped in two mitochondrial

haplogroups. We found individuals of both the A and B haplogroups in both nuclear clusters (cI

and cII). This is consistent with our initial findings [16] in that there is no one-to-one correlation

between mitochondrial haplogroups (A and B) and nuclear genetic clusters.

Insect species with nutritionally incomplete diets (e.g., phloem or blood), such as human lice

often harbor mutualistic bacteria that synthesize missing nutrients [86–88].  As *Wolbachia*, these

endosymbionts are intracellular, and are transmitted vertically from mother to offspring.  The

human lice endosymbiont is *Candidatus* Riesia pediculicola [89], and based on genomic studies,

this endosymbiont carries a small plasmid needed for vitamin B5 synthesis [90] suggesting that

this bacterial symbiont supplies the lice with vitamin B-complex, which the louse cannot produce

by itself [91,92].  In Boyd et al. [90] they compared the genomic sequence of *Candidatus* Riesia

pediculicola found in lice from Cambodia (cI-mtDNA A) and the Netherlands (cII-mtDNA A)

with the lab strain, and they found no indels between these two genomes.  However, when they

included the genome obtained from louse from Honduras (cI-mtDNA B), the authors found

indels [90], reflecting that correlation between mtDNA clades and the endosymbiont.  Here

again, since there is no correlation between the mitochondrial DNA and the microsatellites,

*Candidatus* Riesia pediculicola cannot help to understand the low prevalence of louse hybrids.

An alternative explanation could be related to an unusual paternal genome elimination (PGE)

that has been reported in other members of Psocodea and has been suggested as the mode of sex

determination for the entire lineage that includes Liposcelididae and Phthiraptera [93].  Earlier

studies have shown that some, but not all, male of the clothing lice *P. humanus* only transmitted

their maternal copy of microsatellite markers [94].  Particularly, the allele transmission in

females followed Mendelian expectations but it was non-Mendelian in males: heterozygous male

parents systematically passed on one of their two alleles to their offspring, the one from the

mother [94].  This mode of inheritance is consistent with paternal genome elimination (PGE), a

type of pseudohaplodiploid reproduction found across several arthropod orders in which males

do not transmit paternally inherited alleles to their offspring [95].  A more recent study [96]

assessed this phenomenon in both clothing and head lice.  They found genome-wide male

transmission ratio distortion in both ecotypes of *P. humanus*: males exclusively (or, in some

cases, preferentially) transmit only one of their alleles to their offspring, in agreement with

previous findings from McMeniman and Barker (2006) (92).  In our study, out of the 33 hybrids,

19 were females, 10 males and 4 were juveniles.  Under the PGE scenario, all those 10 male

hybrids will only pass their maternally inherited alleles to the offspring, this will lessen

admixture in the populations.  However, as human louse sex ratios are female-biased, PGE

would have a lesser impact.  Future studies including more louse samples and genomic tools may

help to understand the low number of hybrid lice.

Another cause of genetic isolation may be related to epigenetic effects.  In the broad sense,

epigenetics refers to the differential expression of genes and phenotypes that cannot be explained

by changes in the DNA sequence.  Epigenetic mechanisms like DNA methylation and histone

modification are well known to influence gene regulation, phenotypic plasticity, development,

and the preservation of genome integrity.  However, a growing number of studies are showing

direct and indirect contributions of epigenetics to evolutionary processes since some epigenetic

states are transmitted intergenerationally and affect the phenotype of offspring [97].  One of the

modes for transgenerational epigenetic inheritance is through "genomic imprinting" for which a

gene is distinguished as patrigene or matrigene, depending on its paternal or maternal inheritance

[98].  This mode is often referred as parent-specific gene expression (PSGE) and has been well

study among social insects, where it is shown that epigenetic changes are functionally

responsible for caste formation [99].  Among lice epigenetics has not been studied and opens an

exciting area of new research.  There could also be a relationship between the mechanism

mentioned in the paragraph above related to paternal genome elimination (PGE) and parent-

specific gene expression (PSGE) that could be interfering with the viability and/or fecundity of

hybrid lice, a question that warrants epigenomic studies in human lice.

**Final remarks**

This work is part of the growing number of studies looking into human parasites as probes

for human migration around the world.  We focused on the human louse, which is one of the

oldest known human ectoparasites as indicated by archeological remains, the worldwide co-

distribution of both humans and lice, and coevolutionary studies.  Here, we used microsatellites

to assess human louse genetic diversity worldwide, which revealed the presence of two deep

divergent nuclear clusters, with very limited and recent admixture, mostly in the New World.

We suggest that this pattern is reflecting human migrations during the population of the New

World and the most recent European migration.  However, we hypothesize that there is an

undetermined mechanism that is preventing admixture between these two divergent nuclear

clusters, probably epigenetic.  We also provide a new approach to evaluate demographic

scenarios through DIY-ABC models by using data from the human host, which could guide the

development of new analyses in other host-parasite systems.  Due to the use of microsatellites

that are known to be fast-evolving markers, our analyses are more suited to recent events.  Other

slower evolving markers could provide insights into more ancient events.  Further studies in

human lice including more samples, whole genomic and epigenomic approaches could provide

new knowledge about louse evolution as well as their human host.

## Acknowledgments

We would like to thank former University of Florida undergraduate students: Gebreyes Kassu

and Jackie Fane for their help in the lab and former UF postdoc Marie de Gracia Coquerel,

currently at the Donald Danforth Plant Science Center for comments in early version of the

manuscript.  We greatly appreciate the contribution of Katie Shepherd (Founder & CEO of The

Shepherd Institute for Lice Solutions), Marieta A. H. Braks (Laboratory for Zoonoses and

Environmental, National Institute for Public Health and the Environment, Netherlands) and

collectors worldwide for providing lice used in this study.  We would like to thank Aida Miró-

Herrans and Niyomi House (University of Florida, USA) and Jan Štefka (Institute of Parasitology,

Biology Centre CAS, Czech Republic) for comments on earlier version on this manuscript.

## References

1.  Light JE, Toups MA, Reed DL. What's in a name: The taxonomic status of human head and
body lice. Mol Phylogenet Evol. 2008;47: 1203–1216. doi:10.1016/j.ympev.2008.03.014

2.  Reed DL, Smith VS, Hammond SL, Rogers AR. Genetic analysis of lice supports direct
contact between modern and archaic humans. PLoS Biol. 2004;2: e340.
doi:10.1371/journal.pbio.0020340

3.  Raoult D, Reed DL, Dittmar K, Kirchman JJ, Rolain J, Guillen S, et al. Molecular
Identification of Lice from Pre-Columbian Mummies. J Infect Dis. 2008;197: 535–543.
doi:10.1086/526520

4.  Reed DL, Light JE, Allen JM, Kirchman JJ. Pair of lice lost or parasites regained: the
evolutionary history of anthropoid primate lice. BMC Biol. 2007;5: 7. doi:10.1186/1741-
7007-5-7

5.  Araújo A, Ferreira LF, Guidon N, Maues da Serra Freire N, Reinhard KJ, Dittmar K. Ten
Thousand Years of Head Lice Infection. Parasitol Today. 2000;16: 269.
doi:10.1016/S0169-4758(00)01694-X

6.  Mumcuoglu KY. Human lice: Pediculus and Pthirus. Springer, Berlin, Germany; 2008.

7.  Page RDM, Holmes EC. Molecular evolution: a phylogenetic approach. Oxford ; Malden,
MA: Blackwell Science; 1998.

8.  Hafner MS, Nadler SA. Phylogenetic trees support the coevolution of parasites and their
hosts. Nature. 1988;332: 258–259. doi:10.1038/332258a0

9.  Hafner MS, Sudman PD, Villablanca FX, Spradling TA, Demastes JW, Nadler SA.
Disparate rates of molecular evolution in cospeciating hosts and parasites. Science.
1994;265: 1087–1090.

10. Perry GH. Parasites and human evolution. Evol Anthropol Issues News Rev. 2015;23: 218–
228. doi:10.1002/evan.21427

11. Nozais J-P. The origin and dispersion of human parasitic diseases in the old world (Africa,
Europe and Madagascar). Mem Inst Oswaldo Cruz. 2003;98 Suppl 1: 13–19.

12. Reed DL, Toups MA, Light JE, Allen JM, Flanagin S. Lice and other parasites as markers
of primate evolutionary history. Primate Parasite Ecology: The Dynamics and Study of
Host-Parasite Relationships. Cambridge University Press; 2009. p. 531.

13. Dittmar K, Araújo A, Reinhard KJ. The Study of Parasites Through Time:
Archaeoparasitology and Paleoparasitology. A companion to paleopathology. Chichester,
West Sussex; Malden, MA: Wiley-Blackwell; 2012.

14. Kittler R, Kayser M, Stoneking M. Molecular evolution of Pediculus humanus and the
origin of clothing. Curr Biol CB. 2003;13: 1414–1417.

15.  Toups MA, Kitchen A, Light JE, Reed DL. Origin of clothing lice indicates early clothing
use by anatomically modern humans in Africa. Mol Biol Evol. 2011;28: 29–32.
doi:10.1093/molbev/msq234

16.  Ascunce MS, Toups MA, Kassu G, Fane J, Scholl K, Reed DL. Nuclear Genetic Diversity
in Human Lice (*Pediculus humanus*) Reveals Continental Differences and High
Inbreeding among Worldwide Populations. PLOS ONE. 2013;8: e57619.
doi:10.1371/journal.pone.0057619

17.  Ascunce MS, Fane J, Kassu G, Toloza AC, Picollo MI, González-Oliver A, et al.
Mitochondrial diversity in human head louse populations across the Americas. Am J Phys
Anthropol. 2013;152: 118–129. doi:10.1002/ajpa.22336

18.  Reed DL, Allen JM, Toups MA, Boyd BM, Ascunce MS. The study of primate evolution
from a lousy perspective. In: Krasnov BR, Littlewood DTJ, Morand S, editors. Parasite
Diversity and Diversification: Evolutionary Ecology Meets Phylogenetics. Cambridge:
Cambridge University Press; 2015. pp. 202–214. doi:10.1017/CBO9781139794749.014

19.  Pedersen MW, Antunes C, De Cahsan B, Moreno-Mayar JV, Sikora M, Vinner L, et al.
Ancient human genomes and environmental DNA from the cement attaching 2,000 year-
old head lice nits. Mol Biol Evol. 2022; msab351. doi:10.1093/molbev/msab351

20.  Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A Draft Sequence
of the Neandertal Genome. Science. 2010;328: 710–722. doi:10.1126/science.1188021

21.  Light JE, Allen JM, Long LM, Carter TE, Barrow L, Suren G, et al. Geographic
Distributions and Origins of Human Head Lice (Pediculus humanus capitis) Based on
Mitochondrial Data. J Parasitol. 2008;94: 1275–1281. doi:10.1645/GE-1618.1

22.  Amanzougaghene N, Mumcuoglu KY, Fenollar F, Alfi S, Yesilyurt G, Raoult D, et al. High
Ancient Genetic Diversity of Human Lice, Pediculus humanus, from Israel Reveals New
Insights into the Origin of Clade B Lice. PLOS ONE. 2016;11: e0164659.
doi:10.1371/journal.pone.0164659

23.  Boutellis A, Abi-Rached L, Raoult D. The origin and distribution of human lice in the
world. Infect Genet Evol. 2014;23: 209–217. doi:10.1016/j.meegid.2014.01.017

24.  Ashfaq M, Prosser S, Nasir S, Masood M, Ratnasingham S, Hebert PDN. High diversity
and rapid diversification in the head louse, Pediculus humanus (Pediculidae: Phthiraptera).
Sci Rep. 2015;5. doi:10.1038/srep14188

25.  Sunantaraporn S, Sanprasert V, Pengsakul T, Phumee A, Boonserm R, Tawatsin A, et al.
Molecular survey of the head louse Pediculus humanus capitis in Thailand and its
potential role for transmitting Acinetobacter spp. Parasit Vectors. 2015;8: 127.
doi:10.1186/s13071-015-0742-4

26.   Drali R, Davoust B, Shako J-C, Raoult D, Diatta G. A New Clade of African Body and
Head Lice Infected by Bartonella quintana and Yersinia pestis—Democratic Republic of
the Congo. Am J Trop Med Hyg. 2015;93: 990–993. doi:10.4269/ajtmh.14-0686

27.   Amanzougaghene N, Fenollar F, Davoust B, Djossou F, Ashfaq M, Bitam I, et al.
Mitochondrial diversity and phylogeographic analysis of Pediculus humanus reveals a new
Amazonian clade "F." Infect Genet Evol. 2019;70: 1–8. doi:10.1016/j.meegid.2019.02.006

28.   Amanzougaghene N, Fenollar F, Raoult D, Mediannikov O. Where Are We With Human
Lice? A Review of the Current State of Knowledge. Front Cell Infect Microbiol. 2020;9:
474. doi:10.3389/fcimb.2019.00474

29.   Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis
(MEGA) software version 4.0. Mol Biol Evol. 2007;24: 1596–1599.
doi:10.1093/molbev/msm092

30.   Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing
phylogenetic trees. Mol Biol Evol. 1987;4: 406–425.

31.   Swofford DL. Phylogenetic Analysis Using Parsimony (*and Other Methods). Sunderland,
Massachusetts: Sinauer Associates; 2003.

32.   Excoffier L, Lischer HEL. Arlequin suite ver 3.5: a new series of programs to perform
population genetics analyses under Linux and Windows. Mol Ecol Resour. 2010;10: 564–
567. doi:10.1111/j.1755-0998.2010.02847.x

33.   Excoffier L, Laval G, Schneider S. Arlequin (ver. 3.0): An integrated software package for
population genetics data analysis. Evol Bioinforma Online. 2005;1: 47–50.

34.   Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus
genotype data. Genetics. 2000;155: 945–959.

35.   Peakall R, Smouse PE. genalex 6: genetic analysis in Excel. Population genetic software for
teaching and research. Mol Ecol Notes. 2006;6: 288–295. doi:10.1111/j.1471-
8286.2005.01155.x

36.   Peakall R, Smouse P. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software
for teaching and research - an update. Bioinforma Oxf Engl. 2012 [cited 3 Aug 2012].
doi:10.1093/bioinformatics/bts460

37.   Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a new
method for the analysis of genetically structured populations. BMC Genet. 2010;11: 94.
doi:10.1186/1471-2156-11-94

38.   VäHä J-P, Primmer CR. Efficiency of model-based Bayesian methods for detecting hybrid
individuals under different hybridization scenarios and with different numbers of loci:
GENETIC DETECTION OF HYBRIDS. Mol Ecol. 2005;15: 63–72. doi:10.1111/j.1365-
294X.2005.02773.x

39.  Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the
software STRUCTURE: a simulation study. Mol Ecol. 2005;14: 2611–2620.
doi:10.1111/j.1365-294X.2005.02553.x

40.  Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. Conserv Genet
Resour. 2012;4: 359–361. doi:10.1007/s12686-011-9548-7

41.  Rosenberg NA. distruct: a program for the graphical display of population structure. Mol
Ecol Notes. 2004;4: 137–138. doi:10.1046/j.1471-8286.2003.00566.x

42.  Jombart T. adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics. 2008;24: 1403–1405. doi:10.1093/bioinformatics/btn129

43.  R Core Team. R: A Language and Environment for Statistical Computing. R Foundation
For Statistical Computing, Vienna, Austria. 2015.

44.  Szpiech ZA, Jakobsson M, Rosenberg NA. ADZE: a rarefaction approach for counting
alleles private to combinations of populations. Bioinformatics. 2008;24: 2498–2504.
doi:10.1093/bioinformatics/btn478

45.  Hurlbert SH. The Nonconcept of Species Diversity: A Critique and Alternative Parameters.
Ecology. 1971;52: 577–586. doi:10.2307/1934145

46.  Petit RJ, El Mousadik A, Pons O. Identifying Populations for Conservation on the Basis of
Genetic Markers. Conserv Biol. 2008;12: 844–855. doi:10.1111/j.1523-
1739.1998.96489.x

47.  Kruskal WH, Wallis WA. Use of Ranks in One-Criterion Variance Analysis. J Am Stat
Assoc. 1952;47: 583. doi:10.2307/2280779

48.  StatSoft, Inc. STATISTICA (Data Analysis Software System). Tulsa, OK, USA; 2001.

49.  Beerli P, Felsenstein J. Maximum likelihood estimation of a migration matrix and effective
population sizes in n subpopulations by using a coalescent approach. Proc Natl Acad Sci.
2001;98: 4563–4568. doi:10.1073/pnas.081068098

50.  Beerli P, Palczewski M. Unified Framework to Evaluate Panmixia and Migration Direction
Among Multiple Sampling Locations. Genetics. 2010;185: 313–326.
doi:10.1534/genetics.109.112532

51.  Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population
genetics. Genetics. 2002;162: 2025–2035.

52.  Cornuet J-M, Pudlo P, Veyssier J, Dehne-Garcia A, Gautier M, Leblois R, et al. DIYABC
v2.0: a software to make approximate Bayesian computation inferences about population
history using single nucleotide polymorphism, DNA sequence and microsatellite data.
Bioinformatics. 2014;30: 1187–1189. doi:10.1093/bioinformatics/btt763

53.   Toloza AC, Laguna MF, Ortega-Insaurralde I, Vassena C, Risau-Gusman S. Insights About
Head Lice Transmission From Field Data and Mathematical Modeling. J Med Entomol.
2018;55: 929–937. doi:10.1093/jme/tjy026

54.   Schlötterer C. Evolutionary dynamics of microsatellite DNA. Chromosoma. 2000;109:
365–371. doi:10.1007/s004120000089

55.   Schug MD, Hutter CM, Wetterstrand KA, Gaudette MS, Mackay TF, Aquadro CF. The
mutation rates of di-, tri- and tetranucleotide repeats in Drosophila melanogaster. Mol Biol
Evol. 1998;15: 1751–1760.

56.   Stolle E, Kidner JH, Moritz RFA. Patterns of Evolutionary Conservation of Microsatellites
(SSRs) Suggest a Faster Rate of Genome Evolution in Hymenoptera Than in Diptera.
Genome Biol Evol. 2013;5: 151–162. doi:10.1093/gbe/evs133

57.   Bertorelle G, Benazzo A, Mona S. ABC as a flexible framework to estimate demography
over space and time: some cons, many pros. Mol Ecol. 2010;19: 2609–2625.
doi:10.1111/j.1365-294X.2010.04690.x

58.   Cornuet J-M, Santos F, Beaumont MA, Robert CP, Marin J-M, Balding DJ, et al. Inferring
population history with DIY ABC: a user-friendly approach to approximate Bayesian
computation. Bioinformatics. 2008;24: 2713–2719. doi:10.1093/bioinformatics/btn514

59.   Cornuet J-M, Ravignie V, Estoup A. Inference on population history and model checking
using DNA sequence and microsatellite data with the software DIYABC (v1.0). BMC
Bioinformatics. 2010;11: 401. doi:10.1186/1471-2105-11-401

60.   Torroni A, Sukernik RI, Schurr TG, Starikorskaya YB, Cabell MF, Crawford MH, et al.
mtDNA variation of aboriginal Siberians reveals distinct genetic affinities with Native
Americans. Am J Hum Genet. 1993;53: 591–608.

61.   Tamm E, Kivisild T, Reidla M, Metspalu M, Glenn Smith D, Mulligan CJ, et al. Beringian
standstill and spread of Native American founders. PLoS ONE. 2007;2: e829.

62.   Raghavan M, Steinrücken M, Harris K, Schiffels S, Rasmussen S, DeGiorgio M, et al.
Genomic evidence for the Pleistocene and recent population history of Native Americans.
Science. 2015;349: aab3884. doi:10.1126/science.aab3884

63.   Falush D. Traces of human migrations in Helicobacter pylori populations. Science.
2003;299: 1582–1585. doi:10.1126/science.1080857

64.   Moodley Y, Linz B, Bond RP, Nieuwoudt M, Soodyall H, Schlebusch CM, et al. Age of the
Association between Helicobacter pylori and Man. Ochman H, editor. PLoS Pathog.
2012;8: e1002693. doi:10.1371/journal.ppat.1002693

65.   Moodley Y, Linz B, Yamaoka Y, Windsor HM, Breurec S, Wu J-Y, et al. The peopling of
the Pacific from a bacterial perspective. Science. 2009;323: 527–530.
doi:10.1126/science.1166083

66.    Ghose C, Perez-Perez GI, Dominguez-Bello M-G, Pride DT, Bravi CM, Blaser MJ. East
839           Asian genotypes of Helicobacter pylori strains in Amerindians provide evidence for its
840           ancient human carriage. Proc Natl Acad Sci. 2002;99: 15107–15111.
841           doi:10.1073/pnas.242574599

842    67.    Grenfell BT, Pybus OG, Gog JR, Wood JLN, Daly JM, Mumford JA, et al. Unifying the
843           Epidemiological and Evolutionary Dynamics of Pathogens. Science. 2004;303: 327–332.
844           doi:10.1126/science.1090727

845    68.    Araújo A, Reinhard KJ, Ferreira LF, Gardner SL. Parasites as probes for prehistoric human
846           migrations? Trends Parasitol. 2008;24: 112–115. doi:10.1016/j.pt.2007.11.007

847    69.    Takano-Lee M, Edman JD, Mullens BA, Clark JM. Transmission potential of the human
848           head louse, Pediculus capitis (Anoplura: Pediculidae). Int J Dermatol. 2005;44: 811–816.
849           doi:10.1111/j.1365-4632.2005.02418.x

850    70.    Rózsa L, Apari P. Why infest the loved ones – inherent human behaviour indicates former
851           mutualism with head lice. Parasitology. 2012;139: 696–700.
852           doi:10.1017/S0031182012000017

853    71.    Mumcuoglu KY, Pollack RJ, Reed DL, Barker SC, Gordon S, Toloza AC, et al.
854           International recommendations for an effective control of head louse infestations. Int J
855           Dermatol. 2021;60: 272–280. doi:10.1111/ijd.15096

856    72.    Johnson KP, Allen JM, Olds BP, Mugisha L, Reed DL, Paige KN, et al. Rates of genomic
857           divergence in humans, chimpanzees and their lice. Proc R Soc B Biol Sci. 2014;281:
858           20132174. doi:10.1098/rspb.2013.2174

859    73.    Ricklefs RE, Outlaw DC. A Molecular Clock for Malaria Parasites. Science. 2010;329:
860           226–229. doi:10.1126/science.1188954

861    74.    Olival KJ, Dick CW, Simmons NB, Morales JC, Melnick DJ, Dittmar K, et al. Lack of
862           population genetic structure and host specificity in the bat fly, Cyclopodia horsfieldi,
863           across species of Pteropus bats in Southeast Asia. Parasit Vectors. 2013;6: 231.
864           doi:10.1186/1756-3305-6-231

865    75.    van Schaik J, Dekeukeleire D, Kerth G. Host and parasite life history interplay to yield
866           divergent population genetic structures in two ectoparasites living on the same bat species.
867           Mol Ecol. 2015;24: 2324–2335. doi:10.1111/mec.13171

868    76.    Mazé-Guilmo E, Blanchet S, McCoy KD, Loot G. Host dispersal as the driver of parasite
869           genetic structure: a paradigm lost? Thrall P, editor. Ecol Lett. 2016;19: 336–347.
870           doi:10.1111/ele.12564

871    77.    Talbot B, Vonhof MJ, Broders HG, Fenton B, Keyghobadi N. Comparative analysis of
872           landscape effects on spatial genetic structure of the big brown bat and one of its cimicid
873           ectoparasites. Ecol Evol. 2017;7: 8210–8219. doi:10.1002/ece3.3329

874    78.    Speer KA, Luetke E, Bush E, Sheth B, Gerace A, Quicksall Z, et al. A Fly on the Cave
875           Wall: Parasite Genetics Reveal Fine-Scale Dispersal Patterns of Bats. J Parasitol.
876           2019;105: 555. doi:10.1645/19-20

877    79.    Galbreath KE, Hoberg EP. Return to Beringia: parasites reveal cryptic biogeographic
878           history of North American pikas. Proc R Soc B Biol Sci. 2012;279: 371–378.
879           doi:10.1098/rspb.2011.0482

880    80.    Whiteman NK, Kimball RT, Parker PG. Co-phylogeography and comparative population
881           genetics of the threatened Galápagos hawk and three ectoparasite species: ecology shapes
882           population histories within parasite communities. Mol Ecol. 2007;16: 4759–4773.
883           doi:10.1111/j.1365-294X.2007.03512.x

884    81.    Sweet AD, Wilson RE, Sonsthagen SA, Johnson KP. Lousy grouse: Comparing
885           evolutionary patterns in Alaska galliform lice to understand host evolution and host–
886           parasite interactions. Ecol Evol. 2020;10: 8379–8393. doi:10.1002/ece3.6545

887    82.    Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology.
888           Nat Rev Microbiol. 2008;6: 741–751. doi:10.1038/nrmicro1969

889    83.    Perotti MA, Catalá SS, Ormeño A del V, Zelazowska M, Biliński SM, Braig HR. The sex
890           ratio distortion in the human head louse is conserved over time. BMC Genet. 2004;5: 10.
891           doi:10.1186/1471-2156-5-10

892    84.    Kyei-Poku GK, Colwell DD, Coghlin P, Benkel B, Floate KD. On the ubiquity and
893           phylogeny of Wolbachia in lice. Mol Ecol. 2005;14: 285–294. doi:10.1111/j.1365-
894           294X.2004.02409.x

895    85.    Covacin C, Barker SC. Supergroup F Wolbachia bacteria parasitise lice (Insecta:
896           Phthiraptera). Parasitol Res. 2007;100: 479–485. doi:10.1007/s00436-006-0309-6

897    86.    Baumann P. Biology of Bacteriocyte-Associated Endosymbionts of Plant Sap-Sucking
898           Insects. Annu Rev Microbiol. 2005;59: 155–189.
899           doi:10.1146/annurev.micro.59.030804.121041

900    87.    López-Sánchez MJ, Neef A, Peretó J, Patiño-Navarrete R, Pignatelli M, Latorre A, et al.
901           Evolutionary Convergence and Nitrogen Metabolism in Blattabacterium strain Bge,
902           Primary Endosymbiont of the Cockroach Blattella germanica. Bordenstein SR, editor.
903           PLoS Genet. 2009;5: e1000721. doi:10.1371/journal.pgen.1000721

904    88.    Manzano-Marín A, Oceguera-Figueroa A, Latorre A, Jiménez-García LF, Moya A. Solving
905           a Bloody Mess: B-Vitamin Independent Metabolic Convergence among
906           Gammaproteobacterial Obligate Endosymbionts from Blood-Feeding Arthropods and the
907           Leech *Haementeria officinalis*. Genome Biol Evol. 2015;7: 2871–2884.
908           doi:10.1093/gbe/evv188

89. Sasaki-Fukatsu K, Koga R, Nikoh N, Yoshizawa K, Kasai S, Mihara M, et al. Symbiotic
Bacteria Associated with Stomach Discs of Human Lice. Appl Environ Microbiol.
2006;72: 7349–7352. doi:10.1128/AEM.01429-06

90. Boyd BM, Allen JM, Nguyen N-P, Vachaspati P, Quicksall ZS, Warnow T, et al. Primates,
Lice and Bacteria: Speciation and Genome Evolution in the Symbionts of Hominid Lice.
Mol Biol Evol. 2017;34: 1743–1757. doi:10.1093/molbev/msx117

91. Puchta O. Experimentelle Untersuchungen uber die Bedeutung der Symbiose der
Kleiderlaus Pediculus Vestimenti Burm. Z Parasitenkd. 1955;17.
doi:10.1007/BF00260226

92. Perotti M, Kirkness E, Reed D, Braig H. Endosymbionts of lice. Insect Symbiosis. Bourtzis
K Mill T Ed. 2009;3.

93. Yoshizawa K, Johnson KP. How stable is the "Polyphyly of Lice" hypothesis (Insecta:
Psocodea)?: A comparison of phylogenetic signal in multiple genes. Mol Phylogenet Evol.
2010;55: 939–951. doi:10.1016/j.ympev.2010.02.026

94. McMeniman CJ, Barker SC. Transmission ratio distortion in the human body louse,
Pediculus humanus (Insecta: Phthiraptera). Heredity. 2006;96: 63–68.
doi:10.1038/sj.hdy.6800760

95. Normark BB. The Evolution of Alternative Genetic Systems in Insects. Annu Rev Entomol.
2003;48: 397–423. doi:10.1146/annurev.ento.48.091801.112703

96. De La Filia AG, Andrewes S, Clark JM, Ross L. The unusual reproductive system of head
and body lice ( Pediculus humanus ): Paternal genome loss in human lice males. Med Vet
Entomol. 2018;32: 226–234. doi:10.1111/mve.12287

97. Ashe A, Colot V, Oldroyd BP. How does epigenetics influence the course of evolution?
Philos Trans R Soc B Biol Sci. 2021;376: 20200111. doi:10.1098/rstb.2020.0111

98. Patten MM, Ross L, Curley JP, Queller DC, Bond3uriansky R, Wolf JB. The evolution of
genomic imprinting: theories, predictions and empirical tests. Heredity. 2014;113: 119–
128. doi:10.1038/hdy.2014.29

99. Oldroyd BP, Yagound B. The role of epigenetics, particularly DNA methylation, in the
evolution of caste in insect societies. Philos Trans R Soc B Biol Sci. 2021;376: 20200115.
doi:10.1098/rstb.2020.0115

100. Drali R, Mumcuoglu K, Raoult D. Human Lice in Paleoentomology and
Paleomicrobiology. Drancourt M, Raoult D, editors. Microbiol Spectr. 2016;4.
doi:10.1128/microbiolspec.PoH-0005-2014

101. Finlayson C. Biogeography and evolution of the genus Homo. Trends Ecol Evol. 2005;20:
457–463. doi:10.1016/j.tree.2005.05.019

102. McDougall I, Brown FH, Fleagle JG. Stratigraphic placement and age of modern humans
from Kibish, Ethiopia. Nature. 2005;433: 733–736. doi:10.1038/nature03258

103. Grün R, Stringer C, McDermott F, Nathan R, Porat N, Robertson S, et al. U-series and ESR
analyses of bones and teeth relating to the human burials from Skhul. J Hum Evol.
2005;49: 316–334. doi:10.1016/j.jhevol.2005.04.006

104. Krause J, Lalueza-Fox C, Orlando L, Enard W, Green RE, Burbano HA, et al. The Derived
FOXP2 Variant of Modern Humans Was Shared with Neandertals. Curr Biol. 2007;17:
1908–1912. doi:10.1016/j.cub.2007.10.008

105. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A Draft Sequence
of the Neandertal Genome. Science. 2010;328: 710–722. doi:10.1126/science.1188021

106. Higham T, Compton T, Stringer C, Jacobi R, Shapiro B, Trinkaus E, et al. The earliest
evidence for anatomically modern humans in northwestern Europe. Nature. 2011;479:
521–524. doi:10.1038/nature10484

107. Benazzi S, Douka K, Fornai C, Bauer CC, Kullmer O, Svoboda J, et al. Early dispersal of
modern humans in Europe and implications for Neanderthal behaviour. Nature. 2011;479:
525–528. doi:10.1038/nature10617

108. Arsuaga JL, Martínez I, Arnold LJ, Aranburu A, Gracia-Téllez A, Sharp WD, et al.
Neandertal roots: Cranial and chronological evidence from Sima de los Huesos. Science.
2014;344: 1358–1363. doi:10.1126/science.1253958

109. Devièse T, Karavanić I, Comeskey D, Kubiak C, Korlević P, Hajdinjak M, et al. Direct
dating of Neanderthal remains from the site of Vindija Cave and implications for the
Middle to Upper Paleolithic transition. Proc Natl Acad Sci. 2017;114: 10606–10611.
doi:10.1073/pnas.1709235114

110. Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E. Tracing the
peopling of the world through genomics. Nature. 2017;541: 302–310.
doi:10.1038/nature21347

111. Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, et al. A High-Coverage
Genome Sequence from an Archaic Denisovan Individual. Science. 2012;338: 222–226.
doi:10.1126/science.1224344

112. Wall JD, Yang MA, Jay F, Kim SK, Durand EY, Stevison LS, et al. Higher Levels of
Neanderthal Ancestry in East Asians than in Europeans. Genetics. 2013;194: 199–209.
doi:10.1534/genetics.112.148213

113. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, et al. The complete
genome sequence of a Neanderthal from the Altai Mountains. Nature. 2014;505: 43–49.
doi:10.1038/nature12886

114. Kuhlwilm M, Gronau I, Hubisz MJ, de Filippo C, Prado-Martinez J, Kircher M, et al.
Ancient gene flow from early modern humans into Eastern Neanderthals. Nature.
2016;530: 429–433. doi:10.1038/nature16544

115. Hershkovitz I, Weber GW, Quam R, Duval M, Grün R, Kinsley L, et al. The earliest
modern humans outside Africa. Science. 2018;359: 456–459.
doi:10.1126/science.aap8369

116. Douka K, Slon V, Jacobs Z, Ramsey CB, Shunkov MV, Derevianko AP, et al. Age
estimates for hominin fossils and the onset of the Upper Palaeolithic at Denisova Cave.
Nature. 2019;565: 640–644. doi:10.1038/s41586-018-0870-z

# Supporting information

**S1 Table. Detail of the human louse samples used in the current study.** Groups reflects their geographic origin and STRUCTURE clustering: Africa-Israel-Spain (AIS), South and Southeast Asia (SAs), East Asia (EAs), Europe (E), North America (NA), Mesoamerica (MA), Central America (CA), and South America (SAm).

**S2_S3 Tables. Prior distributions of population sizes parameters for the demographic scenarios described in Table 2 and Fig 2.** Between [ ] are the host values obtained from anthropological references. Louse values were estimated using anthropological references under different louse infection rate scenarios. S2 and S3 Tables values reflect an infection rate of 10% of hominin population having lice with 10 lice per head (S2 Table) and 50% of hominin population having lice with 50 lice per head (S3 Table). Times events are the same as in Table 2 of the manuscript.

**S4 Table. Genetic differentiation.** Louse pairwise genetic differentiation based on FST values from 15 microsatellite loci among sites where 8 or more lice were collected. Sample sizes are in parentheses following population name. Values below the diagonal indicate FST values, p-values are shown above the diagonal; * indicates significant at nominal alpha level of 0.05.

**S1 Fig. Discriminant analysis of principal components (DAPC).** Cluster membership of each louse is depicted by a different colour inside their 95% inertia ellipses. DA eigenvalues are shown in the inserted graphs at the bottom right, whereby the numbers of the discriminants plotted against each other are indicated by dark grey and the remaining discriminants retained for the analysis in light grey.

**S2 Fig. Number of clusters.** $\Delta K$ statistic of Evanno et al. [39] from STRUCTURE analysis from Fig. 3. K = 2 ($\Delta K$ = 858.853).

**S3 Fig. STRUCTURE plot including only non-admixed lice (no hybrids) from both clusters**

**I and II at K = 4.** An ancestry threshold value of q-values as suggested in [38] was used because

it is efficient and accurate at differentiating between non-interbreed and hybrids. Only individuals

with q-value between 0 and < 0.20 or > 0.8 and 1 are classified as non-admixed and were included

in this analysis.

**S4 Fig. Cluster I substructure.** STRUCTURE plot including only non-admixed lice (no hybrids).

**S5 Fig. Cluster II substructure.** STRUCTURE plot including only non-admixed lice (no hybrids).

**S6 Fig. Allelic richness.** Allelic richness per nuclear genetic cluster (cI, orange and cII, blue) using

the rarefaction method implemented in the computer program ADZE (Allelic Diversity Analyzer)

Version 1.0 [44]. Because allelic richness measurements considered sample size, the rarefaction

method allows the estimation of allelic richness for different random subsamples of size "g" from

the populations [45,46].

**S7 Fig. Correspondence analysis (CA).** CA of the nuclear cluster membership frequency and the

mitochondrial haplogroup of head lice.

**S8 Fig. Gene flow.** Full migration model with four units defined by both the nuclear clusters

inferred from STRUCTURE and the mitochondrial haplogroups: unit 1) cI − mtDNA A; unit 2)

cII − mtDNA A, unit 3) cI − mtDNA B; and unit 4) and cII − mtDNA B. Values above and below

the arrows indicate migration rates, arrows indicated directionality.

**S9 Fig. Posterior probabilities for each of the seven demographic scenarios.** Graphs of

posterior probabilities based on DIY-ABC analysis comparing the seven demographic scenarios

when considering low and high louse infestation levels where 10% and 50% indicate that 10% and

50% of the hominin population have 10 and 50 lice per head, respectively. Each scenario was

color-coded as follow Scenario 1 – green, Scenario 2 – red, Scenario 3 – light blue, Scenario 4 –

pink, Scenario 5 – yellow, Scenario 6 – black, Scenario 7 – gray.

Figure 2

A)

Africa
Middle East
Europe
Asia
North America
Mesoamerica
Central America
South America

*K* = 2

cI    cII

*K* = 3

Ia    IIaIIb

*K* = 4

Ia  Ib  IIaIIb

Rw | Isr | Tur | Croa | Nw | Neth | Fr | UK | Spain | Np | Ls Th | Cam | Mn | Oce | SF | Martin | WPB | Nash | MC | MX | Hon | Peru | Arg

B)

Principal Coordinates

Mongolia

Honduras

Europe
North America

Africa, Asia and
Central America

Figure 3

Figure 4

Figure 5

Figure 1