Nuclear genetic diversity of head lice sheds light on human dispersal around the world

Marina S. Ascunce[1,2,*], Ariel C. Toloza[3], Angélica González-Oliver[4], and David L. Reed[5]

[1]Emerging Pathogens Institute, Department of Plant Pathology, University of Florida, Gainesville, Florida, United States of America

[2]USDA-ARS Center for Medical, Agricultural, and Veterinary Entomology, Gainesville, Florida, United States of America

[3]Centro de Investigaciones de Plagas e Insecticidas (CONICET-UNIDEF), Villa Martelli, Buenos Aires, Argentina

[4]Departamento de Biología Celular, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad de México, México

[5]Florida Museum of Natural History, University of Florida, Gainesville, Florida, United States of America

[*]Corresponding author E-mail: marina.ascunce@usda.gov (MSA)

Abstract

The human louse, *Pediculus humanus*, is an obligate blood-sucking ectoparasite that has coevolved with humans for millennia. Because of the intimate relationship between this parasite and the human host, the study of human lice has the potential to shed light on aspects of human evolution that are difficult to interpret using other biological evidence. In this study, we analyzed the genetic variation in 274 human lice from 25 geographic sites around the world by using nuclear microsatellite loci and female-inherited mitochondrial DNA sequences. Nuclear genetic diversity analysis revealed the presence of two distinct genetic clusters I and II, which are subdivided into subclusters: Ia-Ib and IIa-IIb, respectively. Among these samples, we observed the presence of the two most common louse mitochondrial haplogroups: A and B that were found in both nuclear Clusters I and II. Evidence of nuclear admixture was uncommon (33 lice) and was predominately found in the New World potentially mirroring the history of colonization in the Americas. These findings were supported by novel DIYABC simulations that were built using both host and parasite data to define parameters and models suggesting that admixture between cI and cII was very recent. This pattern could also be the result of a reproductive barrier between these two nuclear genetic clusters. In addition to providing new evolutionary knowledge about this human parasite, our study could guide the development of new analyses in other host-parasite systems.

Introduction

The human louse (*Pediculus humanus*, Phthiraptera: Anoplura) is a blood-sucking, wingless, human-specific ectoparasite that is obligate (cannot live without the host) and permanent (complete their life cycle on a host) [1]. This host requirement led to this ectoparasite to be intimately tied to its host through evolutionary times making humans and lice an ideal system for coevolutionary studies [2,3]. For example, studies have shown that primate lice have cospeciated with their primate hosts over the last 25 million years (MY) [2,3]. Furthermore, cophylogenetic analyses revealed that human and chimpanzee lice in the genus *Pediculus* diverged ca. 5.6 MY ago (MYA), which is contemporaneous with the known divergence of chimps and humans [4]. Human head lice have been found in archeological sites, such as a head louse nit obtained from human remains in northeastern Brazil dated 10,000 years old [5] and a nit from Israel dated as 9,000 years old [6] (Fig 1). Multiple archeological findings of human lice around the world are shown in Fig 1 supporting the idea that lice are one of the oldest known human ectoparasites. In host-parasite systems, these long -term associations could lead to reciprocal adaptations in the hosts and parasites (i.e., coevolution), which allows one to build phylogenies for both the parasite and host. Thus obtaining a time estimation of events that co-occurred in both the host and parasite [7]. In cases where the parasite and host have an inextricable relationship, the parasite phylogeny could reflect host evolutionary history [8–10]. In some cases, there are parasite-specific life history traits such as relatively faster evolutionary rates and larger population sizes that allow parasites to maintain greater levels of genetic diversity over time than their hosts. This allows researchers to harness the parasites' genetic diversity in order to explore unclear aspects of human evolution that were based on direct human evidence, such as archeological remains or human molecular data [10–13]. Evolutionary studies

of human lice have already provided insights relative to the ecology of early humans, such as

when humans started using clothes [14,15] and how humans travelled globally [16–18].

Recently, human DNA was extracted from nits found on hairs from ancient mummies in South

America revealing ancestral migration routes [19]. Moreover, almost 20 years ago, Reed et al.

[2] described highly divergent human louse mitochondrial clades and hypothesized that each

louse mitochondrial lineage evolved on different hominin hosts. The authors suggested that as

anatomically modern humans (AMH) migrated out of Africa and interacted with other hominins,

direct physical contact among hominins allowed different lineages of lice to be transferred from

host to host [2]. The hypothesis of direct contact between modern and archaic hominins

predicted using human louse data was later confirmed through host genomic studies comparing

ancient DNA genomes from hominin remains with AMH genomes [20].

Fig 1. Humans and lice. The map represents the geographic distribution of the modern lice

using green dots. Archeological findings of human lice are shown with the figure of human lice

on the map and the corresponding estimated dates from: [3,5,6,21,22]. In addition, the map

reflects the approximate locations of hominin fossil remains and their proposed distribution

based on: [23–38]. Each hominin is color coded as follows: Neanderthal (Blue), Denisovan

(Black), and Anatomical Modern Humans (Orange).

The genetic diversity of human lice has been widely studied based on mitochondrial DNA.

However, little is known about the nuclear genetic diversity of this human parasite, particularly

at a worldwide scale. Early human louse mitochondrial studies revealed the presence of three

deeply divergent mtDNA clades or haplogroups named A, B and C [2,3,14,17,39]. Haplogroup

A is the most common of the human louse haplogroups, has a worldwide distribution and has

been found in lice from human archeological remains in Israel from 2,000 years ago [21].

Haplogroup B has been found in the Americas, Europe, Australia, North and South Africa

[17,39–41], as well in ancient human remains in Israel [21] and Peru [3]. Haplogroup C has a

more geographically restricted distribution and is only present in lice found in Nepal, Ethiopia,

Republic of Congo, Pakistan and Thailand [14,21,42]. Finally, recent studies have suggested the

presence of three more clades: D, E and F [41,43,44]. Clade D is sister to clade A found in

Ethiopia, Zimbabwe and the Democratic Republic of the Congo [41,43,45]. Clade E is sister to

clade C and was detected in head lice from Senegal and Mali [41]. Clade F appears to be

specific to South America, as it has not been reported elsewhere [44]. Head lice can be found in

all the mitochondrial clades (A-F), whereas clothing lice are only found in clades A and D [45].

Despite the extensive work that has been done by many researchers on louse mitochondrial

diversity, there are some aspects that this maternal transmitted molecular marker cannot resolve

such as hybridization events among different louse populations. To overcome this limitation

new nuclear microsatellite markers that have biparental inheritance were developed [16]. In this

study we used these 15 newly developed microsatellite markers in order to provide a

comprehensive understanding of human louse genetic diversity and to assess admixture among

human lice.

In our previous study of louse microsatellite diversity, we only included 91 lice from a

limited number of geographic sites (11 sites) [16]. We found that nuclear genetic diversity in

human lice revealed continental differences suggesting that the genetic pattern found in the

Americas could be mirroring the human host colonization of the New World. In this new study,

we expanded the sampling of lice to 275 and included 25 localities distributed throughout 10

main geographic regions (Table 1, S1 Table, Fig 1). This broader sampling allows us to improve

our understanding of human-louse nuclear genetic diversity while also assessing its correlation

with mitochondrial diversity at a more global scale. This new study also addresses some of the

questions covered in another work where we analyzed 450 mtDNA COX1 sequences from lice

collected in the Americas [17]. Besides enlarging the sampling size and widening the

geographic range in this study we tested seven different evolutionary models by conducting

DIYABC-simulations. For these new simulations, the parameters were developed with both host

and parasite information. We wanted to explore the origin of the hybrids between these louse

nuclear genetic clusters to test if the hybrid lice could be the result of the retention of ancient

polymorphisms due to incomplete lineage sorting (ILS) (Model 1 and 2) or admixture among

louse genetic clusters. For Model 1 and 2, we tested the retention of ancient polymorphisms due

to incomplete lineage sorting (ILS) on the hybrid lice, where the hybrid lice were derived from

lice that parasitized the heads of Neandertals (Model 1), or the heads of anatomical modern

humans (AMH) (Model 2). There were four scenarios considering different times for the

admixture. An ancient admixture model tested hybridization between lice parasitizing

Neanderthals and AMH during periods of contemporaneity between the two hosts (Model 3).

Admixture during European colonization of the Americas 500 years ago was analyzed in Model

4. In Model 5, we tested the hybridization occurring in the last 100 to 40 years around World

Wars I and II. Model 6 reflected the beginning of globalization from the 1980's to ten years ago,

while Model 7 referred to the last 10 years. In addition to providing new evolutionary

knowledge about this parasite, our study could guide the development of new analyses in other

host-parasite systems using this novel approach for the DIYABC-simulations.

Table 1. Microsatellite polymorphisms among human louse populations. Groups reflect

their geographic region and STRUCTURE clustering: Africa-Israel-Spain (AIS), South and

Southeast Asia (SAs), East Asia (EAs), Europe (E), North America (NA), Mesoamerica (MA),

Central America (CA), and South America (SAm).

Geographic Region	Group	Country	Site	N	*Na*	H_O	H_E	F_{IS}	*P-value*
Africa	AIS	Rwanda	Rw	1	1.533	n/a	n/a	n/a	-
West Asia	AIS	Israel	Isr	2	1.667	n/a	n/a	n/a	-
Central Asia	SAs	Nepal	Np	1	1.667	n/a	n/a	n/a	-
South and	SAs	Thailand	Th	1	1.333	n/a	n/a	n/a	-
Southeast	SAs	Cambodia	Cam	22	6.8	0.483	0.703	0.319	*0.000*
Asia	SAs	Laos	Ls	4	3.2	n/a	n/a	n/a	-
East Asia	EAs	Mongolia	Mn	10	5	0.524	0.706	0.267	*0.005*
Europe	E	Turkey	Tur	12	3	0.232	0.408	0.407	*0.000*
	E	Croatia	Croa	6	1.133	n/a	n/a	n/a	-
	E	Norway	Nw	1	1.067	n/a	n/a	n/a	-
	E	Netherlands	Neth	64	2.267	0.032	0.248	0.863	*0.000*
	E	France	Fr	3	1.133	n/a	n/a	n/a	-
	E	United Kingdom	UK	27	2.533	0.145	0.329	0.550	*0.000*
	AIS	Spain	Spain	3	2.067	n/a	n/a	n/a	-
North	NA	USA	Oce	25	2.8	0.083	0.352	0.793	*0.000*
America	NA	USA	SF	1	1	n/a	n/a	n/a	-
	NA	USA	Martin	1	1	n/a	n/a	n/a	-
	NA	USA	WPB	1	1.467	n/a	n/a	n/a	-
	NA	USA	Nash	18	2.467	0.037	0.311	0.864	*0.000*
	NA	USA	MC	22	3.867	0.173	0.452	0.587	*0.000*
Meso	MA	Mexico	MX-1	18	3.13	0.208	0.396	0.481	*0.000*
America	MA	Mexico	DF	1	1	n/a	n/a	n/a	-
Central America	CA	Honduras	Hon	20	4.2	0.365	0.567	0.312	*0.000*
South	SAm	Peru	Peru	1	1.267	n/a	n/a	n/a	-
America	SAm	Argentina	Arg	9	3.2	0.349	0.482	0.231	*0.012*
Total/Average				25	274	12	0.257	0.402	0.469	*0.000*

127
128 *N* = sampling size; *Na* = Average number of alleles; H_O = observed heterozygosity; H_E =
129 expected heterozygosity; F_{IS} = index of inbreeding of individuals resulting from the non-random
130 union of gametes within a subpopulation, positive value indicates heterozygote deficit; *P*
131 proportion of permutations leading to a F_{IS} value larger or equal to the observed one. n/a
132 indicates values not estimated due to sample size smaller than 9 individuals.

MATERIALS AND METHODS

Ethics statement

The Institutional Review Board of the University of Florida exempted the study from review (Exemption of Protocol #2009-U-0422) and waived the need for written informed consent of the participants. This exemption is issued based on the United States Department of Health and Human Services (HHS) regulations. Specifically, HHS regulation 45 CFR part 46 applies to research activities involving human subjects. Because louse removal was voluntary and no information was recorded that would allow patients to be identified directly or through identifiers linked to them, the University of Florida waived the need for written informed consent from the participants.

Sampling

Sampling details are provided in Fig 1, Table 1 and S1 Table. Briefly, 274 human head lice were collected from different individuals at 25 localities throughout the world covering 10 major geographic regions.

DNA extraction

Each individual louse was cut in half, one half was kept in 95% Ethanol for storage, the other half was placed into 1.5 ml Eppendorf tubes containing a cell lysis and proteinase K solution for DNA extraction. The other half of the louse was manually ground using a pestle and the DNA was extracted using the Puregene Core Kit A (QIAGEN, Valencia, California) following manufacturer protocols for tissues however only a third of the volume suggested in the protocol was used due to the small size of the louse material. DNA was quantified using a NanoDrop

154 1000 spectrophotometer (Thermo Scientific), and DNA samples were diluted to reach a

155 concentration between 5 to 10 ng/μL.

Multiple co-amplification: Multiplexes

A total of 15 microsatellite loci were amplified using four multiplexes as described in [16].

We used fluorescently labeled universal primers: M13 (5'-CACGACGTTGTAAAACGAC-3')

and CAG (5'-CAGTCGGGCGTCATCA-3'), each labeled with a unique fluorescent tag (e.g.,

FAM, VIC, NED, PET; Applied Biosystems) to co-amplify multiple loci. Locus-specific

primers were modified by adding the matching 5′ universal primer sequence tails. The complete

set of 15 loci were amplified in four separate 15 μL multiplex-PCR reactions containing 7.5 μL

of 2X Master Mix (Type-It Microsatellite PCR kit, Qiagen, Venlo, Netherlands), 0.015-0.15 μL

of unlabeled forward primer with tail (10 μM), 0.15-0.8 of unlabeled reverse primer (10 μM), 0.5

μL labelled universal tail primer (10 μM), 1-2 μL of total genomic DNA (10-20 ng) and sdH$_2$0

added up to the final volume of 15 μL. Two thermal cycling profiles were used, both beginning

with initial denaturation at 95°C (5 min) and ending with a final extension of 72°C (40 min).

The v1 (touch-down PCR) protocol consisted of 10 cycles of 94°C (30 sec), 60°C/55°C (-

0.5°C/cycle) for 45 sec, and 72°C (45 sec) followed by 25 cycles of 94°C (30 sec), 55°C (45 sec),

and 72°C (1 min). The v2 protocol consisted of 35 cycles of 94°C (30 sec), 52°C (45 sec), and

72°C (45 sec). PCR products were electrophoresed on 1.5% and 4% agarose gels stained with

ethidium bromide and visualized under ultraviolet light. Dilutions in dH$_2$0 from amplicons

(1:100) were run on an ABI 3730xl 96-capillary sequencer using GeneScan 600 LIZ as an

internal size standard (Applied Biosystems) at the University of Florida biotechnology facility

(ICBR). Microsatellite genotypes were scored using the GeneMarker version 1.60 software

(SoftGenetics, LLC).

Mitochondrial PCR amplification and sequencing

The mitochondrial gene cytochrome *c* oxidase 1 (COX1) was amplified through polymerase chain reaction (PCR) using the primers H7005 and L6625 [9] as described in [2]. Each PCR reaction consisted of 25 μL total volume, which included 10 μL of MasterMix (5 PRIME), 1 μL of each primer, 2-4 μL of total genomic DNA, and water. The PCR's cycling profile began with an initial denaturation at 94°C (10 min) followed by 10 cycles of 94°C (1 min), 48°C (1 min), and 65°C (2 min) (decreased by 0.5°C per cycle). Then, there were 35 cycles of 94°C (1 min), 52°C (1 min), and 65°C (2 min) and a final extension of 65°C (10 min). PCR products were run in agarose gel to verify amplification then purified using ExoSAP-IT (USB Corporation, Cleveland, Ohio). Purified PCR products were submitted for sequencing at the University of Florida DNA Sequencing Core Laboratory (ICBR, Gainesville, Florida) using standard fluorescent cycle-sequencing PCR reactions (ABI Prism Big Dye terminator chemistry, Applied Biosystems).

COX1 sequence editing, alignment, and haplotype reconstruction

For each louse sample, the forward and reverse sequences were edited manually and aligned using Sequencher 4.5 (Gene Codes Corporation, Ann Arbor, Michigan). Consensus sequences were generated for each sample using both forward and reverse sequences. These new louse COX1 sequences were aligned along with previous sequences of lice used in our previous work [17]. A final alignment of 379 bp was used to construct a matrix of pairwise differences using uncorrected *p*-distances (proportion of nucleotide sites at which two sequences being compared are different) with MEGA 4 [46]. The genetic relationships were estimated by constructing neighbor-joining (NJ) trees [47] using PAUP* 4.0b10 [48] following Light et. al. [1]. Bootstrapping was performed using 100 pseudo-replications of the data set.

Analyses of genetic diversity and population structure among collection sites using microsatellites

We used the software Arlequin 3.5.1.2 [49] to estimate population genetic diversity values based on the microsatellite data. For all sites regardless of the number of lice per site, we determined the number of alleles per locus. Then, for those 11 sites with sample sizes exceeding 8 lice per site, we also estimated observed and expected heterozygosity (H_O and H_E), and the mean F_{IS} over loci per population with significance tested using 1,023 random permutations. For those 11 sites with sample sizes exceeding 8 lice per site, population pairwise F_{ST} values were estimated using Arlequin 3.5.22 [50]. Significance values for pairwise F_{ST} values were estimated using the permutation procedure with 1,000 permutations.

To assess the distribution of the human louse genetic diversity among world populations through clustering of similar genotypes, we used three different methods utilizing different assumptions. First, we used STRUCTURE [51], a model-based Bayesian clustering method, that assumes Hardy–Weinberg and linkage equilibrium among loci. In addition, we used two multivariate methods, which do not carry these assumptions and could visualize potential cryptic structure, PCoA (Principal Coordinates Analysis) in GenAlEx 6.5 [52,53] and Discriminant Analysis of Principal Components (DAPC) [54].

Bayesian clustering analyses

The Bayesian clustering approach implemented in STRUCTURE 2.3.4 [51] uses the individual multi-locus genotypic data to evaluate models assuming different numbers of genetic clusters (K) based on the posterior probabilities given the data and model. Based on the individual multilocus genotypes and the allele frequencies estimated for the reconstructed clusters, each individual's genome was probabilistically partitioned into membership fractions (q)

(ancestry) in each cluster. Admixed ancestry is modeled by assuming that an individual i has

inherited some fraction (q) of its genome from ancestors in population k [51]. Hybrids can then

be inferred when observing an individual being intermediate of some degree between two or

more clusters in terms of the q-value. For example, individuals with $q = 0.5$ would likely be

first-generation hybrids. We identify possible hybrids based on a q-value threshold of 0.2 as

suggested in VäHä and Primmer (2005) [55], where a threshold q-value of 0.2 means that

individuals with q-value between 0 and < 0.20 or > 0.8 and 1 are classified as non-admixed and

any individual with q-value between 0.2 and 0.8 are classified as hybrids. All simulations used

50,000 Markov chain Monte Carlo (MCMC) generations in the burn-in phase and 100,000

generations in the data collection phase. Ten independent runs were performed using default

parameters for each K, to ensure equilibrium during burn-in and consistency in estimation of the

posterior probabilities. Selection of the number of distinct clusters was based on the evaluation

of the ΔK statistic [56] using the online tool STRUCTURE HARVESTER [57]. The ten

STRUCTURE runs at each K produced nearly identical individual membership coefficients. The

run with the highest likelihood of the data given the parameter values for the predominant

clustering pattern (i.e. the mode) at each K was used for plotting with DISTRUCT [58].

Multivariate analysis using Principal Coordinates Analysis (PCoA)

This multivariate analysis was conducted using pairwise genetic differences between

individuals calculated in GenAlEx 6.5 [52,53] to validate and further define genetic clusters

found using STRUCTURE.

Multivariate analysis using Discriminant Analysis of Principal Components

(DAPC)

This analysis was conducted using the '*adegenet*' package 2.1.3 [59], in the R software

environment version 3.6.3 [60]. For these analyses we only included 175 lice from which we

were able to obtain both microsatellite genotypes and mitochondrial haplogroup sequences. An

additional DAPC included the same louse samples, however we considered four groups

depending on nuclear clusters and haplogroup membership.

Allelic richness within each main nuclear STRUCTURE genetic cluster

251   We estimated the allelic richness within each main nuclear STRUCTURE genetic cluster

252   using the rarefaction method implemented in the computer program ADZE (Allelic Diversity

253   Analyzer) version 1.0 [61]. Because allelic richness measurements considered sample size, the

254   rarefaction method allows the estimation of allelic richness for different random subsamples of

255   size "g" from the populations [62,63].

Analyses based on STRUCTURE genetic clusters and mitochondrial clades

To gain a better understanding of the geographic distribution of the nuclear and

mitochondrial diversity, we conducted a set of Kruskal–Wallis nonparametric analyses of

variance tests [64]. We included the same 175 lice from which we were able to obtain both

microsatellite genotype and mitochondrial haplogroup sequence data and the same eight

geographic regions as described above. Calculated medians were compared and separated at $P <$

0.05 [65]. We compared the Observed vs. Expected Frequency X^2 test to evaluate the fit of the

data (frequencies) to any arbitrary set of expected frequencies among the eight geographic

regions. All statistical tests were performed with a significance value of $\alpha=0.05$.

Analysis of molecular variance (AMOVA) considering the number of different microsatellite

alleles (F_{ST} based) was used to investigate the partitioning of genetic variation between the two

nuclear clusters: cI and cII using Arlequin 3.5 [49]. Statistical significance of differentiation at

each level was assessed by means of 1,023 permutations.

The historical migration patterns among the genetic clusters identified using the methods

describe above were estimated in Migrate-n 3.3.0 using the maximum likelihood approach

[66,67]. Migrate employed estimates of genealogies to sample areas of the coalescence space

with the highest likelihood. The random number of seed and starting values of □ and $4N_em$

(where N_em is the indirect measure of gene flow) were based on F_{ST} as starting parameters and

we used a Brownian mutation model. The data was analyzed using 10 short (10^4 MCMC steps)

and 3 long chains (10^5 MCMC steps), together with adaptive heating to assure convergence.

Comparison of demographic models

We further inferred the demographic history of human lice, based on microsatellite data,

using the approximate Bayesian computation (ABC) approach [68], implemented in the program

DIYABC, v2.0 [69]. We tested different evolutionary Models in which each louse genetic

cluster evolves on different human host populations and diverges in tandem with their hosts. A

total of seven different evolutionary Models were considered with estimations of population size

and divergence times based on information of both human evolution and lice (Table 2, Fig 2).

The first three Models are directly testing Reed et al. (2004) [2] hypothesis of ancient

hybridization between modern and archaic humans. Models 1 and 2 aimed to unravel the

retention of ancient polymorphisms due to incomplete lineage sorting (ILS) on the hybrid lice,

where the hybrid lice were derived from lice that parasitized the heads of Neandertals (Model 1),

or the heads of anatomical modern humans (AMH) (Model 2). In Model 3, admixture events

between lice parasitizing Neanderthals and AMH during periods of host coexistence were tested.

The last four models (Model 4, 5, 6 and 7) do not require ancient hominins, and are based solely

on hybridization events within AMH during recent time periods. In Model 4, we tested the

hypothesis that the hybrid lice were the product of hybridization between lice from the Native

Americans and the first Europeans during the European colonization of the Americas 500 years

ago. In Model 5, we tested the hybridization occurring in the last 100 to 40 years around World

Wars I and II. Model 6 reflects the beginning of globalization in the 1980's to ten years ago,

while   Model   7   refers   to   very   recent   events   in   the   last   10   years.

Table 2. Prior distributions of historical and demographic parameters for the demographic scenarios. Between the brackets []

are the host values obtained from anthropological references. For estimates of times, those values for louse represents louse

generations (g), and the values between [] for hominins are in years (y). For this table, louse values were estimated considering an

infestation of 20 lice in 30% of hominin population and a generational time of 18 days.

Details/Historic Events	Parameter	Distribution	Minimum-Maximum	References
Effective population				
Louse Ancestral Population	N_A	Uniform	1,800-36,000	
[Ancestral hominins]			[300-6,000]	
Louse cluster I	N_1	Uniform	1,800-36,000	
[AMH]			[300-6,000]	
Louse cluster II	N_2	Uniform	1,800-36,000	
[Neanderthals]			[300-6,000]	[35]
Louse Hybrids	N_3	Uniform	270-5,400	15% hybrids from current study
[Hybrids between AMH and Neanderthals]			[?]	
Time Events*				
Time of split clusters I and II	t_S	Uniform	1,200,000 - 16,000,000g	
[Split between AMH and Neanderthals]			[600,000-800,000y]	[35]
T1 ancestral polymorphism from cII	T1	Uniform	2,400,000 - 2,800,000g	[36]
[Presence of Neanderthals in Mongolia]			[120,000 - 140,000y]	[38]
T2 ancestral polymorphism from cI	T2	Uniform	1,000,000 - 1,600,000g	
[Presence of AMH in Middle East]			[50,000-80,000y]	[27]
Time of ancestral admixture	t_{aA}	Uniform	1,000,000 - 1,600,000g	
[AMH-Neanderthals in sympatry]			[50,000-80,000y]	[20]
Times of Recent Admixture				
Time of admixture E1	T_{aE1}	Uniform	2,000 – 20,000g	
[European colonization]			[100 – 1,000y]	
Time of admixture E2	T_{aE2}	Uniform	800 – 2000g	
[Migration after World Wars]			[40 – 100y]	
Time of admixture E1	T_{aG}	Uniform	200 – 800g	
[Initial Globalization]			[10 – 40y]	
Time of admixture E1	T_{aC}	Uniform	20 – 200g	
[Current time -t sampling]			[1 – 10y]	

Admixture rate			
Rate of admixture	ar	Uniform	0.001-0.999
Mutation rate			
Tetranucleotide microsatellite mutation rate	Um1	Loguniform	10^{-9}–10^{-3}
Trinucleotide microsatellite mutation rate	Um2	Loguniform	10^{-7}–10^{-3}
Dinucleotide microsatellite mutation rate	Um3	Loguniform	10^{-6}–10^{-2}

301   *Time parameters for lice are expressed as the number of generations of lice back in time assuming a generation time of 18 days [73].

302   The values for hominins refer to years.

Fig 2. Evolutionary Models. We used seven different demographic historical models with three

different level of parasitism and three different generational times for the lice that lead to a total

of 63 different scenarios that were compared using the approach from the DIY-ABC program.

For each model, we considered three levels of parasitism: from a low level of parasitism

where each human head would have 10 lice and only 10% of the heads would be infested (S2

Table); an intermediate level, where each head would have 20 lice and only 30% of the

population is infested (Table 2); and a high level of lice, where there are 50 lice per head in 50%

of the population (S2 Table) [70].

The louse life cycle includes egg, three nymphal stages, and adult. Therefore, each

demographic model and each level of parasitism, was tested for three different generational times

to account for natural length variation that could occur as was reported in the literature. For this

study we considered the time of head lice maturation from egg to reproductive male adult and

egg-laying-adult female of 18 days, 27 days and 36 days (Table 2, S3 Table) [71–73]. All

microsatellites used the default generalized stepwise mutation model and were assigned a log

uniform distribution with mean mutation rates between 10^{-9}–10^{-2}. The standard mutation rate for

microsatellites is considered to be between 10^{-6}–10^{-2} per locus per generation [74]. However,

due to the possibility of different repeat lengths having different mutation rates [75] for

microsatellites, tetra-, tri- and di- nucleotide repeats were modeled separately. Thus, preliminary

runs were carried out in order to optimize mutation rates [76] following Bertorelle et al. [77]. A

range in allele size of 40 contiguous allelic states was used, as recommended by the program

developers (A. Estoup, personal communication).

Summary statistics calculated for microsatellite loci were as follows: mean number of alleles,

mean genetic diversity, mean size variance, and mean F_{ST}. The selection of summary statistics

was empirical based on preliminary tests where these summary statistics were used and able to

recover the models and parameters following Bertorelle et al. [77]. Through these preliminary

tests the above summary statistics demonstrated their suitability for our data.

We performed about 714,000 simulations using prior parameters for each scenario, thereby

producing a reference table comprising 5,000,000 simulated data sets. First, we pre-evaluated

the fit of observed values to prior distributions of scenarios, by conducting a Principal

Components Analysis (PCA) using data sets simulated with the prior distributions of parameters

and the observed data as well as data sets from the posterior predictive distribution to check the

suitability of the model (scenarios and prior parameters). The posterior probability of each

scenario was assessed by performing direct approach and a polychotomous weighted logistic

regression that estimates the difference between simulated (from the reference table) and

observed data sets [78,79] after linear discriminant analysis on summary statistics [69]. Only the

posterior probabilities for the 1% (50,000) of simulated data sets closest to the observed data

were retained. Confidence in choice of a scenario was estimated as the proportion of times the

selected scenario did not possess the highest posterior probability compared to competing

scenarios for the 500 simulated data sets (type I error). Confidence in scenario choice was

further tested including the estimation of the prior error rate, which is calculated as the

probability of choosing an incorrect model when drawing model index and parameter values into

priors (the probability of selecting the chosen scenario when it is not correct, a type-II error).

After scenario choice, we proceeded to parameter inference estimated from the modes and 95%

confidence intervals (CI) of their posterior distributions.

RESULTS

Nuclear Genetic Diversity

The 274 human louse specimens comprising 25 geographic sites through 10 worldwide

geographic regions revealed high microsatellite diversity with overall five to 20 alleles per locus

and an average of 12 alleles per locus (Table 1, S1 File). In the 11 sites with more than 8 lice

analyzed, H_O values ranged from 0.032 to 0.524 (average of 0.257) and H_E values ranged from

0.248 to 0.706 (average of 0.402), reflecting a general pattern of deficit of heterozygotes (Table

1). Furthermore, in all 11 louse populations, genotype proportions deviated significantly from

Hardy-Weinberg expectations with values of F_{IS} significantly positive and an overall F_{IS} value of

0.469 ($P < 0.00001$) (Table 1). Overall, there were high values of genetic differentiation among

the 11 sites, with F_{ST} values of up to 0.663 between the sites: Mongolia (Mn) and Netherlands

(Neth) (S4 Table). The lowest F_{ST} of 0.053 was found between sites in New York (Oce) and

Tennessee (Nash).

Mitochondrial diversity

We also obtained the COX1 sequence data from 175 head lice spanning 18 sampling

locations throughout nine geographic regions (Table 3). One hundred four (104) lice belonged to

haplogroup A and 71 to haplogroup B (Table 3). As expected based on previous studies

[17,39,40], haplogroup A shows a worldwide distribution including Africa, Asia, Europe and the

Americas. For our sampling, haplogroup B has a narrower distribution and it is found only in

lice from Europe, North and Central America (Table 3).

Table 3. Nuclear and mitochondrial louse genetic diversity. Geographic distribution of 175

human lice including combined information of their mitochondrial DNA haplogroups based the

COX1 gene and their assignment to each louse major nuclear genetic clusters described in this

study.

Geographic Region	Group	Country	Site	mtA+clusterI	mtB+clusterI	mtA+clusterII	mtB+clusterII
Africa	AIS	Rwanda	Rw	1			
South Asia	SAs	Nepal	Np	1			
Southeast Asia	SAs	Cambodia	Cam	21			
		Laos	Ls	1			
East Asia	EAs	Mongolia	Mn	6			
Europe	E	Turkey	Tur				7
	E	Croatia	Croa				5
	E	Norway	Nw			1	
	E	Netherlands	Neth			49	5
	E	France	Fr			2	
	E	United Kingdom	UK				26
North America	NA	USA	Oce			12	4
	NA	USA	SF			1	
	NA	USA	Nash				3
	NA	USA	MC			1	9
Mesoamerica	MA	Mexico	MX			6	
Central America	CA	Honduras	Hon		12		
South America	SAm	Argentina	Arg			2	
Total				30	12	74	59

Nuclear genetic clusters at the worldwide scale

All clustering methods: STRUCTURE, PCoA and DAPC were concordant in identifying the

distinct nuclear genetic clusters: Cluster I (cI) with subclusters Ia and Ib and Cluster II (cII) with

subclusters IIa and IIb (Fig 1; S1 Fig). The differences in their geographic distribution can be

seen where cI defined by head lice from Africa, Israel, Asia and the Americas comprised a more

worldwide distribution, while cII composed of head lice mainly from Europe and the Americas

reflects a more restricted geographic distribution. While most of the genetic diversity of human

louse populations from around the world seems to be highly structured geographically,

approximately 12% of the lice we sampled (33 of 274 lice) appear to be hybrids (lice with q-

values less than 0.8), with 75% (25 lice) of the hybrids found in the New World.

In the STRUCTURE analysis, after a signal for $K = 2$ ($\Delta K = 858.853$, S2 Fig), increasing K

values in the STRUCTURE plots revealed geographically localized structure. At $K = 3$, a new

cluster formed within Europe from the original cII, composed of lice primarily from the

Netherlands forming the subcluster cIIb (light blue color). When K was increased to 4 a

population from Honduras and other parts of the Americas emerged from cluster I as the Ib

subcluster, with the other samples remaining in subcluster Ia. This Honduras subcluster (Ib) may

be indicative of Native American louse lineage.

The Principal Coordinate Analysis (PCoA) supported the two main nuclear genetic clusters

and subclusters identified by STRUCTURE, with putative hybrid lice not assigned to either

cluster, rather placing hybrids between them (Fig 3). Finally, the eigenvalues obtained in the

DAPC analysis showed that the genetic structure was captured mainly by the first principal

component representing cI (to the right) and cII (to the left) (S1 Fig).

Fig 3. Human louse worldwide nuclear genetic diversity. A) Genetic clusters inferred from

STRUCTURE simulations at $K = 2$: Cluster I (cI, Orange) and Cluster II (cII, Blue), $K = 3$ cI and

cIIa-cIIb, and $K = 4$ cIa-cIb and cIIa-cIIb. In the bar plot, each louse is represented by a single

vertical line and the length of each color segment represents the proportion of membership (Q) to

each cluster. At the top of the STRUCTURE bar plot, an additional bar was added using different

colors for each continent and main geographic region where each louse sample was collected. At

the bottom of the STRUCTURE bar plot, the codes correspond to geographic origin from the lice

using abbreviations as defined in Table 1. B) Distributions of points in the first two dimensions

resulting from Principal Coordinates Analysis (PCoA) conducted using pairwise genetic distance

comparisons. Each site was color-coded to match STRUCTURE results (e.g., sites in Asia were

color-coded orange, and sites in Europe were color-coded blue). These codes correspond to

geographic origin and abbreviations as defined in Table 1. Map author: Maulucioni

(https://commons.wikimedia.org/wiki/File:World_map_with_the_Americas_on_the_right.png).

This file is licensed under the Creative Commons Attribution-Share Alike 4.0 International

license. https://creativecommons.org/licenses/by-sa/4.0/legalcode

Additionally, we conducted further analysis with all the dataset excluding hybrids as

identified through STRUCTURE q-values. Thus, when using the reduced dataset only non-

admixed individuals with shared ancestry q-values larger than 0.8 as suggested by [55],the

STRUCTURE analysis reflected similar results identifying cI and cII, with their corresponding

subclusters at $K = 4$ (S3 Fig).

Worldwide cluster I

All lice that belong to the cI (q-values larger than 0.5) were included in a second

STRUCTURE simulations to assess the substructure within this cluster. At $K = 2$, this figure

showed the presence of two clusters: one composed of lice from the Old World (Africa, Asia and

Europe), and the other of lice from the New World (Americas) (S4 Fig). The further split of the

genetic clusters during the increase of defined K in the STRUCTURE analysis showed at $K = 3$

the emergence of an Asian cluster except for most of the lice from Mongolia that remains within

the Old Word cluster (S4 Fig). At $K = 4$, a subset of those lice from Mongolia formed a unique

new cluster (S4 Fig).

Geographically restricted cluster II

424 According to our current sampling, cII has narrower geographic distribution than cI and is

425 limited mostly to Europe and the Americas (Fig 3). This cluster showed a substructure at $K = 2$

426 where most of the lice from the Netherlands plus a few others form a distinct cluster (light blue

427 in plot) (S5 Fig). Both these individual cluster analyses for cI and cII highlight the geographic

428 structure observed among louse populations.

Allelic richness analysis per main genetic cluster

We estimated the allelic richness per main non-admixed clusters I and II (without hybrids)

using the rarefaction method implemented in the computer program ADZE (Allelic Diversity

Analyzer) Version 1.0 [61]. Allelic richness per non-admixed cluster was higher in cluster I than

II (S6 Fig). This pattern reflects very different evolutionary histories for the two clusters. While

high allelic richness is consistent with a large effective population size, it could also be

concordant with greater genetic structure. It is also noteworthy that clusters I and II differ in

sample size (80 lice in the cI versus 174 in cII).

Statistical analysis of genetic clusters and mitochondrial clades

The combined analysis of the nuclear cluster membership frequency and the mitochondrial

haplogroup were plotted using a Correspondence Analysis (CA) and showed the presence of four

well defined groups (S7 Fig). The dimensions 1 and 2 represented the largest amount of inertia

or variance representing a measure of deviation of independence. Each one of the dimensions

possess an eigenvalue that represents its relative importance and how much of the variance is

explained. In our study, one dimension represents 58.7 % of the inertia and two dimensions

represents 99.6% of the inertia. Of this, dimension 1 differentiated both nuclear genetic clusters:

cI and cII, while dimension 2 differentiated the mitochondrial haplogroups. One group

corresponds to lice of nuclear cluster II-haplogroup A and included lice from Europe, North

America, Mesoamerica and South America. The second group is also nuclear cluster II-

haplogroup B from the same regions, except South America. The third group is composed of

individuals of cluster I-haplogroup A: South Asia, Africa-Israel-Spain, and some parts of Europe,

while the last group belonged to cluster I with haplogroup B and was found in Central America

(S7 Fig). Furthermore, multiple comparisons (non-parametric Kruskal–Wallis test) indicated

that the distribution of the individuals assigned to the nuclear genetic clusters I and II were not

distributed randomly among the eight geographic regions (H = 103.55, $P < 0.0001$). The cluster

membership median values were significantly different among: South Asia (SAs), East Asia

(EA), Africa-Israel-Spain (AIS), Central America (CA), and Europe (E) ($P < 0.05$). On the other

hand, cluster membership medians values for North America (NA), South America (Sam), and

Mesoamerica (MA) were not significantly different. Both nuclear genetic assignment and

haplogroup frequencies differed significantly among the eight geographic regions (nuclear

cluster assignment $X^2 = 175$, $P < 0.0001$; haplogroup frequencies $X^2 = 42$, $P < 0.0001$).

AMOVA by nuclear clusters and mitochondrial clades

Hierarchical analyses of molecular variance (AMOVA) employed to explore the distribution

of nuclear genetic variation considering the two nuclear genetic clusters: cI and cII revealed

significant genetic differentiation between them ($\Phi_{ST} = 0.33$; $P < 0.00$). When examining only

the non-admixed individuals (those with at least 0.8 of coefficient of ancestry values) the

differentiation between cI and cII increased ($\Phi_{ST} = 0.4$; $P < 0.00$).

Migration patterns considering nuclear genetic clusters and mitochondrial clades

The historical migration pattern was estimated in Migrate-n 3.3.0 using maximum likelihood

inference and the Brownian motion mutation model [66,67]. Given that previous studies have

suggested a different evolutionary history between the mitochondrial haplogroups A and B, we

evaluated a full migration model with four populations defined by both the nuclear clusters

inferred from STRUCTURE and the mitochondrial haplogroups: cI – mtDNA A; cI – mtDNA B;

cII – mtDNA A and cII – mtDNA B. Migration rates between these four populations varied. In

general, migration rates within nuclear clusters were high, contrasting with low migration rates

within mitochondrial haplogroups (S8 Fig). For example, we observed the highest migration

rates within cluster II (cluster II – mtDNA A and cluster II – mtDNA B) and some of the lowest

migration rates between cluster I – mtDNA A and cluster II – mtDNA A. Because

STRUCTURE analysis indicated the effect of geography on the genetic structure of human lice,

further studies including more samples from each sampling site are needed to further understand

migration patterns among geographic human louse populations.

Demographic history

The presence of two louse distinct nuclear genetic clusters (I and II) could be the result of the

retention of ancient polymorphisms due to incomplete lineage sorting (ILS) or admixture from

previous isolated lineages. We tested seven different demographic models in which each louse

genetic cluster and its hybrids evolve on different human host populations and diverge in tandem

with them using the Bayesian computation (ABC) method as implemented in the DIY-ABC

package [52]. The models reflect different hypotheses that followed human evolutionary events

from the admixture between Neanderthals and AMH (Models 1-3) to human movement around

the globe (Models 4-7) (Table 2, Fig 2). We tested these 7 demographic models using three

different levels of louse parasitism as well as three different generational times creating a total of

63 scenarios (S9 Fig). The three most strongly supported demographic models (5, 6 and 7) were

topologically similar with each involving admixture between the two nuclear clusters at different

times since migrations after World Wars (Model 5), initial globalization (Model 6) and current

times (Model 7) (Fig 4, Table 4, S9 Fig). Considering 10% and 30% levels of louse parasitism,

for any of the generational times (18 days, 27 days or 36 days), the comparison of the seven

demographic models showed Model 7 (current time) as the model with the highest logistic

posterior probability of ca. 0.90 (Fig 4, Table 4, S9 Fig). For the high level of parasitism at 50%,

the longer the generational time the older admixture models were shown with higher

probabilities (S9 Fig). For example, when the longest generational time of 36 days was used in

the scenario, then Model 4 appeared as the scenario with the highest probability. This model

corresponds to the European colonization of the New World. When 27 days were chosen as

generational time, Model 5, dated around the migrations after World Wars, presented the highest

probability. Finally, when the generational time was set at 18 days, Model 6 that it is associated

with Globalization shows the highest probability (S9 Fig).

Fig 4. Posterior probabilities for each of the seven demographic models. Graphs of posterior

probabilities based on DIY-ABC analysis comparing the seven demographic models when

considering intermediate louse infestation level where 30% of the hominin population have lice

with 20 lice per head. A) The posterior probabilities are estimated using the direct approach and

in B) they are inferred by a logistic approach [69]. Each scenario was color-coded as follows:

Model 1 – green, Model 2 – red, Model 3 – light blue, Model 4 – pink, Model 5 – yellow, Model

6 – black, Model 7 – gray.

Table 4. Posterior probabilities for the seven demographic models shown in Fig 2. Probabilities obtained through the logarithmic

method (with 95% confidence intervals in brackets) for the seven scenarios inferred from DIY-ABC analysis. For each model three

parasitism level were considered where 10% indicates that 10% of the hominin population have lice with 10 lice per head, 30% and 50%

indicate that 30% and 50% of the hominin population have 20 and 50 lice per head, respectively. Generational time was set to 18 days.

For each comparison, the selected scenario (bold entry in shaded cell) was the one with the highest probability value. Confidence in

scenario choice, shown under probability values, was evaluated by estimating the type I error rate from 1000 randomly constructed

simulated data sets, for Models 5 and 7, respectively.

Models	Parasitism Level		
	10%	30%	50%
1: Retain Polymorphisms from cI	0.0000 [0.0000,0.0000]	0.0000 [0.0000,0.0715]	0.0000 [0.0000,0.1130]
2: Retain Polymorphisms from cII	0.0000 [0.0000,0.0000]	0.0000 [0.0000,0.0715]	0.0000 [0.0000,0.1130]
3: Ancestral admixture cI and cII	0.0000 [0.0000,0.0000]	0.0000 [0.0000,0.0715]	0.0000 [0.0000,0.1130]
4: European colonization of the New World	0.0000 [0.0000,0.0000]	0.0001 [0.0000,0.0716]	0.0560 [0.0000,0.1522]
5: European migration to the New World after WWs	0.0000 [0.0000,0.0000]	0.0071 [0.0000,0.0757]	**0.5204 [0.4477,0.5930]**
6: Globalization	0.0026 [0.0011,0.0041]	0.3792 [0.2622,0.4963]	0.2754 [0.2042,0.3467]
7: Current time - sampling	**0.9974 [0.9959,0.9989]** 0.335	**0.6136 [0.4959,0.7312]** 0.317	0.1482 [0.0095,0.2869] 0.372

DISCUSSION

In this study, we found that the nuclear microsatellite data from lice were grouped in two distinct genetic clusters: cI and cII. Cluster I had a worldwide distribution whereas cII was restricted to Europe and the Americas (Fig 5). Remarkably, the two nuclear clusters revealed a significant genetic differentiation between them (Φ_{ST} = 0.33; P < 0.00) despite having geographic overlap in either Europe or the Americas. In the Structure analysis at $K = 4$ each of the two Clusters further split into subclusters: Ia-Ib and IIa-IIb, which were sorted based on their geographical origin (i.e., continents). These results are consistent with our previous work where we assessed louse microsatellite genetic diversity in fewer samples from only four geographic regions (North America, Central America, Asia, and Europe) [16]. Both this current study and our past work support the idea that head louse populations are strongly structured geographically. This current study highlights a genetic relationship between Asia (cIa) and Central America (cIb) subclusters supporting the idea that the Central America cluster (cIb) is probably of Native American origin since a potential source population for the first people of the Americas has been suggested to be East Asia [80–82] (Fig 5). Thus, microsatellite data suggest that current human louse populations in the Americas retain human louse genetic diversity from louse brought by the first people [16]. This work adds to our previous studies showing that human lice could be used to track aspects of human evolution.

Fig 5. Proposed global co-migration human lice and humans. Top: Map depicting the collection sites of the human lice included in this study. The color of each circle corresponds to the majority nuclear genetic cluster to which sampled individuals were assigned. Sites with admixed lice are indicated with patterned circles including colors of the two major genetic clusters at that site. The proposed migrations of anatomically modern humans out of Africa into

Europe, Asia and the Americas, as well as the more recent European colonization of the New

World are indicated with thick grey arrows. Hypothetical human louse co-migrations are

indicated with orange and blue arrows. At the bottom, the STRUCTURE plot from Fig 3A

corresponding to the assignment of 274 lice from 25 geographical sites at $K = 4$ (Table 1) is

shown.   Map   author: Maulucioni

(https://commons.wikimedia.org/wiki/File:World_map_with_the_Americas_on_the_right.png).

This file is licensed under the Creative Commons Attribution-Share Alike 4.0 International

license. https://creativecommons.org/licenses/by-sa/4.0/legalcode

Other human parasites have also shed light into different aspects of human evolution [11–13].

For example, the genetic diversity of a bacterial parasite, *Helicobacter pylori*, which infects the

stomach of humans, has been studied to provide insights into human migrations and strong

correlations between the parasite and human host have been found reflecting the out of Africa

expansion [83,84], the colonization of the Austronesian speaking region [85], and the settlement

of the Americas by Asians across the Bering Strait [86]. However, some microorganisms such

as bacteria and viruses that reproduce clonally could also be affected by selective sweeps that

could lead to the removal of entire lineages, or to gene specific as well as genome wide reduction

in genetic diversity. For those scenarios where parasites have clonal reproduction the effect of

selection could be strong resulting in parasite phylogenies reflecting the evolutionary history of

their genes under selection (selective sweep) rather than the demographic history of the host. On

the other hand, macroparasites (such as lice) often reproduce sexually, which lessens the effect of

selective sweeps since it would only affect closely linked loci in their genomes [87]. Thus,

sexually reproductive parasites may track host evolutionary history more accurately than clonal

parasites. For this reason, the use of macroparasites, such as pinworms [88] and lice [2,3], that

are also obligate parasites and co-evolved with their host, has become increasingly common in

the study of human evolution. In fact, the human louse, *Pediculus humanus*, is one of the most

well studied human parasites [1–6,14–19,21–28].

Human lice can disperse from a host to another at any motile stage (nymphs and adults),

however adults are more likely to transfer from one head to another [89]. Lice can be transferred

via direct body contact, for example during a fight, living with others, or through any other

means where their heads are in close proximity to each other [90]. Additionally, lice can be

transferred via indirect contact, such as through sharing clothes that cover the head or combs

between an infested person and a non-infested person. In this case the shared item may still have

living lice that can be transferred to the non-infested person [91]. Louse ability to transfer from

host to host directly or indirectly along with louse life history traits such as larger population

sizes and faster rates of molecular evolution compared to their hosts reveal cryptic aspects of

their hosts including how the host interact with other species, and host cryptic population

structure. These host aspects might not be evident using the host data alone or not yet

discernible in the host's genome [9,92,93]. Differences between parasite and host intrinsic life-

history traits may lead to incongruities in patterns of population structure between hosts and their

parasites [94–98]. This phenomenon has been observed amongst several parasites including flies

[98], helminths [99], and feather lice [100]. In one case, coevolutionary studies of the

relationship between Arctic Galliformes and their parasitic lice in Alaska suggest lice

consistently move among different host species and populations in that region [101]. While the

bird hosts present some level of population structure and/or isolation by distance, the parasitic

lice do not show any structure [101]. This lack of congruence between the genetic structures of

the parasite and host suggests that there are still some cryptic contact or distributional overlaps

among host bird populations. Although, these species interactions were not perceptible from the

host data alone, they could be useful for ecological and conservation efforts. In our current study,

we show the presence of two louse distinct nuclear genetic clusters: I and II, which are not

observable in the human host. These could be the result of the retention of ancient

polymorphisms due to incomplete lineage sorting (ILS) or admixture from previous isolated

lineages. To test if louse current genetic diversity is the result of ILS or admixture we used the

approximate Bayesian computation (ABC) method as implemented in the DIY-ABC package

[69].

We performed ABC analyses for seven demographic models, three different level of

parasitism and three different generational times (Table 2, 4, Figs 2, 4, S9 Fig). The

demographic model with higher probabilities under 10% and 30% parasitism level corresponded

to the model suggesting recent admixture between the two nuclear Clusters I and II in Model 7.

This model supports the STRUCTURE analysis, where most of the hybrids are being observed in

the New World. Although for cases of higher parasitism, older admixture models were

supported including models 4, 5, and 6. It could be possible that in the past both infestation rate

and generational time could have been different due to different climate and human

environmental conditions. In fact, several studies have shown that louse life history traits

changed due to temperature. For example, it was reported that under laboratory conditions

oviposition, egg hatchability, and fertility drastically varied with temperature suggesting that

louse population sizes might increase in the winter months and decrease in summer [102].

Another study reported that *P. h. humanus* ceased to lay eggs at temperatures beneath 20ºC and

lay rapidly at 37ºC. These conditions lead to a mean life span of 16 days and 27 days,

respectively [71]. Despite the variation in the life cycle in relation to temperature, head lice were

recorded from every group of man in every part of the world, even where it is known that lice are

rare [103]. This could be generally attributed to human habits rather than to the climate [104].

For the remainder of this discussion, we will focus on the 30% level of parasitism's scenarios as

it is the one that most likely reflects current louse population conditions [105–107].

While most of the genetic diversity in human louse populations around the world seem to be

highly geographically structured, there are a total of 33 hybrids out of 274 lice analyzed reaching

12% of our current sampling. However, 25 of those hybrids (75%) are found in the New World,

which could represent a zone of secondary contact as a consequence of human colonization

history. The three models that consider recent admixture events from European migration after

both World Wars (Model 5), initial globalization (Model 6) and current times at sampling

(Model 7) were the most supported (Table 2, 4, Fig 4, S9 Fig). Thus, a new and puzzling insight

from this study was the finding of such low prevalence of hybrid lice and that these hybrids are

geographically concentrated in the New World. This pattern could reflect human colonization of

the New World by Europe as mentioned above. However, if these colonization events have been

occurring for many louse generations, this does not explain the low occurrence of hybrids. It

could be possible that our sampling is spatially/geographically bias as many samples came from

the Americas (43%) (Table 1). If that would have been the case, it could be possible that while

hybrids are rare in the Americas, they could be more common on other regions of the world. To

answer this question, future studies on genetic diversity of human lice should consider sampling

other geographic areas. However, even if that would be the case, still the lack of hybrids in the

Americas deserves some further discussion. While this is beyond the scope of the present paper,

we hypothesize that this pattern of limited admixture could be indicating some sort of genetic

isolation between cI and cII such that the hybrids are not able to persist for many generations

within the human louse populations. Thus, it is worthwhile to enumerate the potential causes for

future researcher.

One potential mechanism of reproductive isolation may involve the effect of endosymbiont

on host biology. Endosymbionts could also alter host biology in diverse ways, including the

induction of reproductive manipulations, such as feminization, parthenogenesis, male killing and

sperm–egg incompatibility, which could lead to reproductive incompatibility [108]. *Wolbachia*

is a common intracellular bacterial symbiont of arthropods and filarial nematodes that has been

shown to infect human lice [109–111]. When populations of insect hosts that contain different

strains of *Wolbachia* interbreed, or when one population has *Wolbachia* and the other population

does not, the interbreeding matings ("hybrids") resulted in reduced fitness or the complete

absence of progeny. If this process were to occur in human lice, it could explain the low number

of hybrids; however, because *Wolbachia* endosymbiont is transmitted with the mitochondrial

genome through the egg cytoplasm, endosymbionts and mtDNA should reflect a similar history.

In our study, mitochondrial lineages were grouped in two mitochondrial haplogroups. We found

individuals of both the A and B haplogroups in both nuclear clusters (cI and cII). This is

consistent with our initial findings [16] in that there is no one-to-one correlation between

mitochondrial haplogroups (A and B) and nuclear genetic clusters.

The human louse, unlike most animals, which have single-chromosome mitochondrial (mt)

genomes [112], has fragmented mt genomes, each with 9-20 minichromosomes [113,114]. The

human head louse (*Pediculus humanus capitis*, family Pediculidae) and human body louse

(*Pediculus humanus corporis*) have the most fragmented mt genomes with 20 minichromosomes

with a conserved common pattern whereas each of the other protein-coding and rRNA genes has

its own minichromosome [113]. Furthermore, this pattern of gene distribution and the

fragmented mt genome apparently evolved in the MRCA of the human lice and have been

retained since [115]. Because stretches of identical sequences (26–127 bp long) have been found

to be shared between genes on different minichromosomes, this was thought to be evidence for

inter-minichromosome recombination [113] however the frequency of recombination has not

been clearly established. Although, there could be some concern about mt minichromosome

recombination since we only focused on one gene, the probability of that affecting our results is

expected to be minimal.

Insect species with nutritionally incomplete diets (e.g., phloem or blood), such as human lice

often harbor mutualistic bacteria that synthesize missing nutrients [116–118]. As *Wolbachia*,

these endosymbionts are intracellular, and are transmitted vertically from mother to offspring.

The human lice endosymbiont is *Candidatus* Riesia pediculicola [119], and based on genomic

studies, this endosymbiont carries a small plasmid needed for vitamin B5 synthesis [120]

suggesting that this bacterial symbiont supplies the lice with vitamin B-complex, which the louse

cannot produce by itself [121,122]. Studies have confirmed that this obligate human louse

endosymbiont is housed in the mycetome, which is localized on the ventral side of the louse

midgut. In females, this endosymbiont also migrates to the lateral oviducts at the beginning of

oogenesis, which is transovarially transmitted to the progeny confirming its maternal inheritance

[123,124]. In Boyd et al. [120] they compared the genomic sequence of *Candidatus* Riesia

pediculicola found in lice from Cambodia (cI-mtDNA A) and the Netherlands (cII-mtDNA A)

with the lab strain, and they found no indels between these two genomes. However, when they

included the genome obtained from louse from Honduras (cI-mtDNA B), the authors found

indels [120], reflecting that correlation between mtDNA clades and the endosymbiont. A recent

study using a larger number of lice has also shown the codivergence of the endosymbiont

Candidatus Riesia pediculicola and the mt clades of their *P. humanus* host [123,124]. In our

study, since there is no correlation between the mitochondrial DNA and the nuclear genetic

clusters and since *Candidatus* Riesia pediculicola co-diverge with the mtDNA lineages, this

endosymbiont cannot help to understand the low prevalence of louse hybrids.

An alternative explanation could be related to an unusual paternal genome elimination (PGE)

that has been reported in other members of Psocodea particularly Liposcelididae and Phthiraptera

where for these two lineages PGE has been suggested as the mode of sex determination [125].

Earlier studies have shown that some, but not all, male of the clothing lice, *P. h. humanus*, only

transmitted their maternal copy of microsatellite markers while females showed a Mendelian

gene transmission [126]. This mode of non-Mendelian inheritance by male lice is consistent a

type of pseudohaplodiploid reproduction known as paternal genome elimination (PGE), in which

males only transmit maternally inherited alleles to their offspring [127]. A more recent study

[128] assessed this phenomenon in both clothing and head lice. In both ecotypes of *P. humanus*

louse males exclusively (or, in some cases, preferentially) transmit only the maternally inherited

alleles to their offspring revealing a genome-wide male transmission ratio distortion for both

ecotypes in agreement with early findings from McMeniman and Barker (2006) [126]. In our

study, out of the 33 hybrids, 19 were female, 10 were male, and 4 were juvenile. Under the PGE

scenario, all 10 male hybrids would only pass their maternally inherited alleles to the offspring

thus lessening admixture in the populations. Since for our study lice were collected at a single

time point, there is no data of louse pedigree which would have been needed to address PGE in

detail. In addition, as human louse sex ratios are female-biased, PGE would have a lesser impact.

Future studies including louse family pedigree and the analysis of allele transmission patterns are

needed to understand if the low number of hybrid lice is a result of PGE.

If hybrids present some type of phenotypes that are not well adapted compared to the non-

admixed parental lice, this could lead to an alternative hypothesis. For example, epigenetic

mechanisms like DNA methylation and histone modification are well known to influence gene

regulation, phenotypic plasticity, development, and the preservation of genome integrity. When

most of the studies of epigenetics are related to intra-generational times, there is a growing

number of studies focusing on transgenerational epigenetics: the inheritance of a modified

phenotype from the parental generation without changes in genes or gene sequence. This

transgenerational epigenetics suggest that there are direct and indirect contributions of

epigenetics to evolutionary processes [129]. One of the modes for transgenerational epigenetic

inheritance is through "genomic imprinting" for which a gene is distinguished as patrigene or

matrigene, depending on its paternal or maternal inheritance [130]. This mode is often referred

as parent-specific gene expression (PSGE) and has been well study among social insects, where

it is shown that epigenetic changes are functionally responsible for caste formation [131].

Among lice, epigenetics has not been studied and opens an exciting area of new research. There

could also be a relationship between the mechanism mentioned in the paragraph above related to

paternal genome elimination (PGE) and parent-specific gene expression (PSGE) that could be

interfering with the viability and/or fecundity of hybrid lice, a question that warrants epigenomic

studies in human lice.

Final remarks

In this work, we assessed the use of human louse nuclear genetic diversity as a tool to

reconstruct human migration around the world, by analyzing microsatellite data from 274 lice.

In addition to well-known genetic analysis methods, we developed a new approach to evaluate

demographic scenarios through DIY-ABC models by using data from the human host. This

novel approach could also guide the development of new analyses in other host-parasite systems.

Our results showed the presence of two divergent nuclear louse genetic clusters, with very

limited and recent admixture, mostly in the New World. We suggested that this pattern is

reflective of human dispersal to the Americas, with an early wave of louse-human co-migration

during the population of the New World and the most recent European migration. We also

hypothesized that there is a potential mechanism that is preventing admixture between these two

divergent nuclear clusters, most likely epigenetic in origin. Due to the use of microsatellites that

are known to be fast-evolving markers, our analyses are more suited to recent events, and slower

evolving markers could provide insights into more ancient events. Further studies in human lice

including more samples, whole genomic, and epigenomic approaches could provide new

knowledge about louse evolution as well as their human host.

Acknowledgments

We would like to thank former University of Florida undergraduate students: Gebreyes

Kassu and Jackie Fane for their help in the lab, Lauren Justice (UF student) and former UF

postdoc Marie de Gracia Coquerel, currently at the Donald Danforth Plant Science Center for

comments in early version of the manuscript. We greatly appreciate the contribution of Katie

Shepherd (Founder & CEO of The Shepherd Institute for Lice Solutions), Marieta A. H. Braks

(Laboratory for Zoonoses and Environmental, National Institute for Public Health and the

Environment, Netherlands) and collectors worldwide for providing lice used in this study. We

would like to thank Aida Miró-Herrans and Niyomi House (University of Florida, USA) and Jan

Štefka (Institute of Parasitology, Biology Centre CAS, Czech Republic) for comments on earlier

version on this manuscript.

References

1. Light JE, Toups MA, Reed DL. What's in a name: The taxonomic status of human head and
body lice. Mol Phylogenet Evol. 2008;47: 1203–1216. doi:10.1016/j.ympev.2008.03.014

2. Reed DL, Smith VS, Hammond SL, Rogers AR. Genetic analysis of lice supports direct
contact between modern and archaic humans. PLoS Biol. 2004;2: e340.
doi:10.1371/journal.pbio.0020340

3. Raoult D, Reed DL, Dittmar K, Kirchman JJ, Rolain J, Guillen S, et al. Molecular
Identification of Lice from Pre-Columbian Mummies. J Infect Dis. 2008;197: 535–543.
doi:10.1086/526520

4. Reed DL, Light JE, Allen JM, Kirchman JJ. Pair of lice lost or parasites regained: the
evolutionary history of anthropoid primate lice. BMC Biol. 2007;5: 7. doi:10.1186/1741-
7007-5-7

5. Araújo A, Ferreira LF, Guidon N, Maues da Serra Freire N, Reinhard KJ, Dittmar K. Ten
Thousand Years of Head Lice Infection. Parasitol Today. 2000;16: 269.
doi:10.1016/S0169-4758(00)01694-X

6. Mumcuoglu KY. Human lice: *Pediculus* and *Pthirus*. Springer, Berlin, Germany; 2008.

7. Page RDM, Holmes EC. Molecular evolution: a phylogenetic approach. Oxford ; Malden,
MA: Blackwell Science; 1998.

8. Hafner MS, Nadler SA. Phylogenetic trees support the coevolution of parasites and their
hosts. Nature. 1988;332: 258–259. doi:10.1038/332258a0

9. Hafner MS, Sudman PD, Villablanca FX, Spradling TA, Demastes JW, Nadler SA.
Disparate rates of molecular evolution in cospeciating hosts and parasites. Science.
1994;265: 1087–1090. Available: http://www.ncbi.nlm.nih.gov/pubmed/8066445

10. Perry GH. Parasites and human evolution. Evol Anthropol Issues News Rev. 2015;23: 218–
228. doi:10.1002/evan.21427

11. Nozais J-P. The origin and dispersion of human parasitic diseases in the old world (Africa,
Europe and Madagascar). Mem Inst Oswaldo Cruz. 2003;98 Suppl 1: 13–19. Available:
http://www.ncbi.nlm.nih.gov/pubmed/12687757

12. Reed DL, Toups MA, Light JE, Allen JM, Flanagin S. Lice and other parasites as markers
of primate evolutionary history. Primate Parasite Ecology: The Dynamics and Study of
Host-Parasite Relationships. Cambridge University Press; 2009. p. 531.

13. Dittmar K, Araújo A, Reinhard KJ. The Study of Parasites Through Time:
Archaeoparasitology and Paleoparasitology. A companion to paleopathology. Chichester,
West Sussex; Malden, MA: Wiley-Blackwell; 2012.

14. Kittler R, Kayser M, Stoneking M. Molecular evolution of *Pediculus humanus* and the
origin of clothing. Curr Biol CB. 2003;13: 1414–1417. Available:
http://www.ncbi.nlm.nih.gov/pubmed/12932325

15. Toups MA, Kitchen A, Light JE, Reed DL. Origin of clothing lice indicates early clothing
use by anatomically modern humans in Africa. Mol Biol Evol. 2011;28: 29–32.
doi:10.1093/molbev/msq234

16. Ascunce MS, Toups MA, Kassu G, Fane J, Scholl K, Reed DL. Nuclear Genetic Diversity
in Human Lice (*Pediculus humanus*) Reveals Continental Differences and High
Inbreeding among Worldwide Populations. PLOS ONE. 2013;8: e57619.
doi:10.1371/journal.pone.0057619

17. Ascunce MS, Fane J, Kassu G, Toloza AC, Picollo MI, González-Oliver A, et al.
Mitochondrial diversity in human head louse populations across the Americas. Am J Phys
Anthropol. 2013;152: 118–129. doi:10.1002/ajpa.22336

18. Reed DL, Allen JM, Toups MA, Boyd BM, Ascunce MS. The study of primate evolution
from a lousy perspective. In: Krasnov BR, Littlewood DTJ, Morand S, editors. Parasite
Diversity and Diversification: Evolutionary Ecology Meets Phylogenetics. Cambridge:
Cambridge University Press; 2015. pp. 202–214. doi:10.1017/CBO9781139794749.014

19. Pedersen MW, Antunes C, De Cahsan B, Moreno-Mayar JV, Sikora M, Vinner L, et al.
Ancient human genomes and environmental DNA from the cement attaching 2,000 year-
old head lice nits. Mol Biol Evol. 2022; msab351. doi:10.1093/molbev/msab351

20. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A Draft Sequence
of the Neandertal Genome. Science. 2010;328: 710–722. doi:10.1126/science.1188021

21. Amanzougaghene N, Mumcuoglu KY, Fenollar F, Alfi S, Yesilyurt G, Raoult D, et al. High
Ancient Genetic Diversity of Human Lice, *Pediculus humanus*, from Israel Reveals New
Insights into the Origin of Clade B Lice. PLOS ONE. 2016;11: e0164659.
doi:10.1371/journal.pone.0164659

22. Drali R, Mumcuoglu K, Raoult D. Human Lice in Paleoentomology and
Paleomicrobiology. Drancourt M, Raoult D, editors. Microbiol Spectr. 2016;4.
doi:10.1128/microbiolspec.PoH-0005-2014

23. Finlayson C. Biogeography and evolution of the genus *Homo*. Trends Ecol Evol. 2005;20:
457–463. doi:10.1016/j.tree.2005.05.019

24. McDougall I, Brown FH, Fleagle JG. Stratigraphic placement and age of modern humans
from Kibish, Ethiopia. Nature. 2005;433: 733–736. doi:10.1038/nature03258

25. Grün R, Stringer C, McDermott F, Nathan R, Porat N, Robertson S, et al. U-series and ESR
analyses of bones and teeth relating to the human burials from Skhul. J Hum Evol.
2005;49: 316–334. doi:10.1016/j.jhevol.2005.04.006

26. Krause J, Lalueza-Fox C, Orlando L, Enard W, Green RE, Burbano HA, et al. The Derived
FOXP2 Variant of Modern Humans Was Shared with Neandertals. Curr Biol. 2007;17:
1908–1912. doi:10.1016/j.cub.2007.10.008

27. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A Draft Sequence
of the Neandertal Genome. Science. 2010;328: 710–722. doi:10.1126/science.1188021

28. Higham T, Compton T, Stringer C, Jacobi R, Shapiro B, Trinkaus E, et al. The earliest
evidence for anatomically modern humans in northwestern Europe. Nature. 2011;479:
521–524. doi:10.1038/nature10484

29. Benazzi S, Douka K, Fornai C, Bauer CC, Kullmer O, Svoboda J, et al. Early dispersal of
modern humans in Europe and implications for Neanderthal behaviour. Nature. 2011;479:
525–528. doi:10.1038/nature10617

30. Arsuaga JL, Martínez I, Arnold LJ, Aranburu A, Gracia-Téllez A, Sharp WD, et al.
Neandertal roots: Cranial and chronological evidence from Sima de los Huesos. Science.
2014;344: 1358–1363. doi:10.1126/science.1253958

31. Devièse T, Karavanić I, Comeskey D, Kubiak C, Korlević P, Hajdinjak M, et al. Direct
dating of Neanderthal remains from the site of Vindija Cave and implications for the
Middle to Upper Paleolithic transition. Proc Natl Acad Sci. 2017;114: 10606–10611.
doi:10.1073/pnas.1709235114

32. Nielsen R, Akey JM, Jakobsson M, Pritchard JK, Tishkoff S, Willerslev E. Tracing the
peopling of the world through genomics. Nature. 2017;541: 302–310.
doi:10.1038/nature21347

33. Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, et al. A High-Coverage
Genome Sequence from an Archaic Denisovan Individual. Science. 2012;338: 222–226.
doi:10.1126/science.1224344

34. Wall JD, Yang MA, Jay F, Kim SK, Durand EY, Stevison LS, et al. Higher Levels of
Neanderthal Ancestry in East Asians than in Europeans. Genetics. 2013;194: 199–209.
doi:10.1534/genetics.112.148213

35. Prüfer K, Racimo F, Patterson N, Jay F, Sankararaman S, Sawyer S, et al. The complete
genome sequence of a Neanderthal from the Altai Mountains. Nature. 2014;505: 43–49.
doi:10.1038/nature12886

36. Kuhlwilm M, Gronau I, Hubisz MJ, de Filippo C, Prado-Martinez J, Kircher M, et al.
Ancient gene flow from early modern humans into Eastern Neanderthals. Nature.
2016;530: 429–433. doi:10.1038/nature16544

37. Hershkovitz I, Weber GW, Quam R, Duval M, Grün R, Kinsley L, et al. The earliest
modern humans outside Africa. Science. 2018;359: 456–459.
doi:10.1126/science.aap8369

38. Douka K, Slon V, Jacobs Z, Ramsey CB, Shunkov MV, Derevianko AP, et al. Age
estimates for hominin fossils and the onset of the Upper Palaeolithic at Denisova Cave.
Nature. 2019;565: 640–644. doi:10.1038/s41586-018-0870-z

39. Light JE, Allen JM, Long LM, Carter TE, Barrow L, Suren G, et al. Geographic
Distributions and Origins of Human Head Lice (*Pediculus humanus capitis*) Based on
Mitochondrial Data. J Parasitol. 2008;94: 1275–1281. doi:10.1645/GE-1618.1

40. Boutellis A, Abi-Rached L, Raoult D. The origin and distribution of human lice in the
world. Infect Genet Evol. 2014;23: 209–217. doi:10.1016/j.meegid.2014.01.017

41. Ashfaq M, Prosser S, Nasir S, Masood M, Ratnasingham S, Hebert PDN. High diversity
and rapid diversification in the head louse, *Pediculus humanus* (Pediculidae: Phthiraptera).
Sci Rep. 2015;5. doi:10.1038/srep14188

42. Sunantaraporn S, Sanprasert V, Pengsakul T, Phumee A, Boonserm R, Tawatsin A, et al.
Molecular survey of the head louse *Pediculus humanus capitis* in Thailand and its
potential role for transmitting *Acinetobacter* spp. Parasit Vectors. 2015;8: 127.
doi:10.1186/s13071-015-0742-4

43. Drali R, Davoust B, Shako J-C, Raoult D, Diatta G. A New Clade of African Body and
Head Lice Infected by *Bartonella quintana* and *Yersinia pestis*—Democratic Republic of
the Congo. Am J Trop Med Hyg. 2015;93: 990–993. doi:10.4269/ajtmh.14-0686

44. Amanzougaghene N, Fenollar F, Davoust B, Djossou F, Ashfaq M, Bitam I, et al.
Mitochondrial diversity and phylogeographic analysis of *Pediculus humanus* reveals a
new Amazonian clade "F." Infect Genet Evol. 2019;70: 1–8.
doi:10.1016/j.meegid.2019.02.006

45. Amanzougaghene N, Fenollar F, Raoult D, Mediannikov O. Where Are We with Human
Lice? A Review of the Current State of Knowledge. Front Cell Infect Microbiol. 2020;9:
474. doi:10.3389/fcimb.2019.00474

46. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis
(MEGA) software version 4.0. Mol Biol Evol. 2007;24: 1596–1599.
doi:10.1093/molbev/msm092

47. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing
phylogenetic trees. Mol Biol Evol. 1987;4: 406–425.

48. Swofford DL. Phylogenetic Analysis Using Parsimony (*and Other Methods). Sunderland,
Massachusetts: Sinauer Associates; 2003.

49. Excoffier L, Lischer HEL. Arlequin suite ver 3.5: a new series of programs to perform
population genetics analyses under Linux and Windows. Mol Ecol Resour. 2010;10: 564–
567. doi:10.1111/j.1755-0998.2010.02847.x

50. Excoffier L, Laval G, Schneider S. Arlequin (ver. 3.0): An integrated software package for
population genetics data analysis. Evol Bioinforma Online. 2005;1: 47–50.

51. Pritchard JK, Stephens M, Donnelly P. Inference of population structure using multilocus
genotype data. Genetics. 2000;155: 945–959. Available:
http://www.ncbi.nlm.nih.gov/pubmed/10835412

52. Peakall R, Smouse PE. genalex 6: genetic analysis in Excel. Population genetic software for
teaching and research. Mol Ecol Notes. 2006;6: 288–295. doi:10.1111/j.1471-
8286.2005.01155.x

53. Peakall R, Smouse P. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software
for teaching and research - an update. Bioinforma Oxf Engl. 2012 [cited 3 Aug 2012].
doi:10.1093/bioinformatics/bts460

54. Jombart T, Devillard S, Balloux F. Discriminant analysis of principal components: a new
method for the analysis of genetically structured populations. BMC Genet. 2010;11: 94.
doi:10.1186/1471-2156-11-94

55. VäHä J-P, Primmer CR. Efficiency of model-based Bayesian methods for detecting hybrid
individuals under different hybridization scenarios and with different numbers of loci:
Genetic Detection of Hybrids. Mol Ecol. 2005;15: 63–72. doi:10.1111/j.1365-
294X.2005.02773.x

56. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the
software STRUCTURE: a simulation study. Mol Ecol. 2005;14: 2611–2620.
doi:10.1111/j.1365-294X.2005.02553.x

57. Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. Conserv Genet
Resour. 2012;4: 359–361. doi:10.1007/s12686-011-9548-7

58. Rosenberg NA. distruct: a program for the graphical display of population structure. Mol
Ecol Notes. 2004;4: 137–138. doi:10.1046/j.1471-8286.2003.00566.x

59. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics. 2008;24: 1403–1405. doi:10.1093/bioinformatics/btn129

60. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation
For Statistical Computing, Vienna, Austria. 2015.

61. Szpiech ZA, Jakobsson M, Rosenberg NA. ADZE: a rarefaction approach for counting
alleles private to combinations of populations. Bioinformatics. 2008;24: 2498–2504.
doi:10.1093/bioinformatics/btn478

62. Hurlbert SH. The Nonconcept of Species Diversity: A Critique and Alternative Parameters.
Ecology. 1971;52: 577–586. doi:10.2307/1934145

63. Petit RJ, El Mousadik A, Pons O. Identifying Populations for Conservation on the Basis of
Genetic Markers. Conserv Biol. 2008;12: 844–855. doi:10.1111/j.1523-
1739.1998.96489.x

64. Kruskal WH, Wallis WA. Use of Ranks in One-Criterion Variance Analysis. J Am Stat
Assoc. 1952;47: 583. doi:10.2307/2280779

65. StatSoft, Inc. STATISTICA (Data Analysis Software System). Tulsa, OK, USA; 2001.

66. Beerli P, Felsenstein J. Maximum likelihood estimation of a migration matrix and effective
population sizes in n subpopulations by using a coalescent approach. Proc Natl Acad Sci.
2001;98: 4563–4568. doi:10.1073/pnas.081068098

67. Beerli P, Palczewski M. Unified Framework to Evaluate Panmixia and Migration Direction
Among Multiple Sampling Locations. Genetics. 2010;185: 313–326.
doi:10.1534/genetics.109.112532

68. Beaumont MA, Zhang W, Balding DJ. Approximate Bayesian computation in population
genetics. Genetics. 2002;162: 2025–2035.

69. Cornuet J-M, Pudlo P, Veyssier J, Dehne-Garcia A, Gautier M, Leblois R, et al. DIYABC
v2.0: a software to make approximate Bayesian computation inferences about population
history using single nucleotide polymorphism, DNA sequence and microsatellite data.
Bioinformatics. 2014;30: 1187–1189. doi:10.1093/bioinformatics/btt763

70. Toloza AC, Laguna MF, Ortega-Insaurralde I, Vassena C, Risau-Gusman S. Insights About
Head Lice Transmission From Field Data and Mathematical Modeling. J Med Entomol.
2018;55: 929–937. doi:10.1093/jme/tjy026

71. Bacot A. A Contribution to the Bionomics of *Pediculus humanus* (*Vestimenti*) and
Pediculus capitis. Parasitology. 1917;9: 228–258. doi:10.1017/S0031182000006065

72. Meinking TL. Infestations. Curr Probl Dermatol. 1999;11: 73–118.

73. Takano-Lee M, Yoon KS, Edman JD, Mullens BA, Clark JM. In vivo and in vitro rearing
of *Pediculus humanus capitis* (Anoplura: Pediculidae). J Med Entomol. 2003;40: 628–
635.

74. Schlötterer C. Evolutionary dynamics of microsatellite DNA. Chromosoma. 2000;109:
365–371. doi:10.1007/s004120000089

75. Schug MD, Hutter CM, Wetterstrand KA, Gaudette MS, Mackay TF, Aquadro CF. The
mutation rates of di-, tri- and tetranucleotide repeats in *Drosophila melanogaster*. Mol
Biol Evol. 1998;15: 1751–1760.

76. Stolle E, Kidner JH, Moritz RFA. Patterns of Evolutionary Conservation of Microsatellites
(SSRs) Suggest a Faster Rate of Genome Evolution in Hymenoptera Than in Diptera.
Genome Biol Evol. 2013;5: 151–162. doi:10.1093/gbe/evs133

77.   Bertorelle G, Benazzo A, Mona S. ABC as a flexible framework to estimate demography
over space and time: some cons, many pros. Mol Ecol. 2010;19: 2609–2625.
doi:10.1111/j.1365-294X.2010.04690.x

78.   Cornuet J-M, Santos F, Beaumont MA, Robert CP, Marin J-M, Balding DJ, et al. Inferring
population history with DIY ABC: a user-friendly approach to approximate Bayesian
computation. Bioinformatics. 2008;24: 2713–2719. doi:10.1093/bioinformatics/btn514

79.   Cornuet J-M, Ravignie V, Estoup A. Inference on population history and model checking
using DNA sequence and microsatellite data with the software DIYABC (v1.0). BMC
Bioinformatics. 2010;11: 401. doi:10.1186/1471-2105-11-401

80.   Torroni A, Sukernik RI, Schurr TG, Starikorskaya YB, Cabell MF, Crawford MH, et al.
mtDNA variation of aboriginal Siberians reveals distinct genetic affinities with Native
Americans. Am J Hum Genet. 1993;53: 591–608. Available:
http://www.ncbi.nlm.nih.gov/pubmed/7688933

81.   Tamm E, Kivisild T, Reidla M, Metspalu M, Glenn Smith D, Mulligan CJ, et al. Beringian
standstill and spread of Native American founders. PLoS ONE. 2007;2: e829.

82.   Raghavan M, Steinrücken M, Harris K, Schiffels S, Rasmussen S, DeGiorgio M, et al.
Genomic evidence for the Pleistocene and recent population history of Native Americans.
Science. 2015;349: aab3884. doi:10.1126/science.aab3884

83.   Falush D. Traces of human migrations in Helicobacter pylori populations. Science.
2003;299: 1582–1585. doi:10.1126/science.1080857

84.   Moodley Y, Linz B, Bond RP, Nieuwoudt M, Soodyall H, Schlebusch CM, et al. Age of the
Association between *Helicobacter pylori* and Man. Ochman H, editor. PLoS Pathog.
2012;8: e1002693. doi:10.1371/journal.ppat.1002693

85.   Moodley Y, Linz B, Yamaoka Y, Windsor HM, Breurec S, Wu J-Y, et al. The peopling of
the Pacific from a bacterial perspective. Science. 2009;323: 527–530.
doi:10.1126/science.1166083

86.   Ghose C, Perez-Perez GI, Dominguez-Bello M-G, Pride DT, Bravi CM, Blaser MJ. East
Asian genotypes of *Helicobacter pylori* strains in Amerindians provide evidence for its
ancient human carriage. Proc Natl Acad Sci. 2002;99: 15107–15111.
doi:10.1073/pnas.242574599

87.   Grenfell BT, Pybus OG, Gog JR, Wood JLN, Daly JM, Mumford JA, et al. Unifying the
Epidemiological and Evolutionary Dynamics of Pathogens. Science. 2004;303: 327–332.
doi:10.1126/science.1090727

88.   Araújo A, Reinhard KJ, Ferreira LF, Gardner SL. Parasites as probes for prehistoric human
migrations? Trends Parasitol. 2008;24: 112–115. doi:10.1016/j.pt.2007.11.007

89.   Takano-Lee M, Edman JD, Mullens BA, Clark JM. Transmission potential of the human
head louse, *Pediculus capitis* (Anoplura: Pediculidae). Int J Dermatol. 2005;44: 811–816.
doi:10.1111/j.1365-4632.2005.02418.x

90.   Rózsa L, Apari P. Why infest the loved ones – inherent human behaviour indicates former
mutualism with head lice. Parasitology. 2012;139: 696–700.
doi:10.1017/S0031182012000017

91.   Mumcuoglu KY, Pollack RJ, Reed DL, Barker SC, Gordon S, Toloza AC, et al.
International recommendations for an effective control of head louse infestations. Int J
Dermatol. 2021;60: 272–280. doi:10.1111/ijd.15096

92.   Johnson KP, Allen JM, Olds BP, Mugisha L, Reed DL, Paige KN, et al. Rates of genomic
divergence in humans, chimpanzees and their lice. Proc R Soc B Biol Sci. 2014;281:
20132174. doi:10.1098/rspb.2013.2174

93.   Ricklefs RE, Outlaw DC. A Molecular Clock for Malaria Parasites. Science. 2010;329:
226–229. doi:10.1126/science.1188954

94.   Olival KJ, Dick CW, Simmons NB, Morales JC, Melnick DJ, Dittmar K, et al. Lack of
population genetic structure and host specificity in the bat fly, *Cyclopodia horsfieldi*,
across species of *Pteropus* bats in Southeast Asia. Parasit Vectors. 2013;6: 231.
doi:10.1186/1756-3305-6-231

95.   van Schaik J, Dekeukeleire D, Kerth G. Host and parasite life history interplay to yield
divergent population genetic structures in two ectoparasites living on the same bat species.
Mol Ecol. 2015;24: 2324–2335. doi:10.1111/mec.13171

96.   Mazé-Guilmo E, Blanchet S, McCoy KD, Loot G. Host dispersal as the driver of parasite
genetic structure: a paradigm lost? Thrall P, editor. Ecol Lett. 2016;19: 336–347.
doi:10.1111/ele.12564

97.   Talbot B, Vonhof MJ, Broders HG, Fenton B, Keyghobadi N. Comparative analysis of
landscape effects on spatial genetic structure of the big brown bat and one of its cimicid
ectoparasites. Ecol Evol. 2017;7: 8210–8219. doi:10.1002/ece3.3329

98.   Speer KA, Luetke E, Bush E, Sheth B, Gerace A, Quicksall Z, et al. A Fly on the Cave
Wall: Parasite Genetics Reveal Fine-Scale Dispersal Patterns of Bats. J Parasitol.
2019;105: 555. doi:10.1645/19-20

99.   Galbreath KE, Hoberg EP. Return to Beringia: parasites reveal cryptic biogeographic
history of North American pikas. Proc R Soc B Biol Sci. 2012;279: 371–378.
doi:10.1098/rspb.2011.0482

100.   Whiteman NK, Kimball RT, Parker PG. Co-phylogeography and comparative population
genetics of the threatened Galápagos hawk and three ectoparasite species: ecology shapes
population histories within parasite communities. Mol Ecol. 2007;16: 4759–4773.
doi:10.1111/j.1365-294X.2007.03512.x

101. Sweet AD, Wilson RE, Sonsthagen SA, Johnson KP. Lousy grouse: Comparing
1033   evolutionary patterns in Alaska galliform lice to understand host evolution and host–
1034   parasite interactions. Ecol Evol. 2020;10: 8379–8393. doi:10.1002/ece3.6545

1035   102. Nuttall GHF. The Biology of *Pediculus humanus*. Parasitology. 1917;10: 80–185.
1036   doi:10.1017/S0031182000003747

1037   103. Ferris GF. Contributions towards a Mono-graph of the Sucking Lice, viii. Stanford; 1935.

1038   104. Buxton PA. The Louse. 1st ed. Baltimore, MD.: Williams & Wilkins Co.; 1946.

1039   105. Gratz NG. Human lice: their prevalence, control and resistance to insecticides, A review
1040   1985-1997. Geneva, Switzerland: WHOPES, CTD, WHO; 1997.

1041   106. Falagas ME, Matthaiou DK, Rafailidis PI, Panos G, Pappas G. Worldwide Prevalence of
1042   Head Lice. Emerg Infect Dis. 2008;14: 1493–1494. doi:10.3201/eid1409.080368

1043   107. Takano-Lee M, Yoon KS, Edman JD, Mullens BA, Clark JM. In vivo and in vitro rearing
1044   of *Pediculus humanus capitis* (Anoplura: Pediculidae). J Med Entomol. 2003;40: 628–
1045   635. Available: http://www.bioone.org/doi/abs/10.1603/0022-2585-40.5.628

1046   108. Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology.
1047   Nat Rev Microbiol. 2008;6: 741–751. doi:10.1038/nrmicro1969

1048   109. Perotti MA, Catalá SS, Ormeño A del V, Zelazowska M, Biliński SM, Braig HR. The sex
1049   ratio distortion in the human head louse is conserved over time. BMC Genet. 2004;5: 10.
1050   doi:10.1186/1471-2156-5-10

1051   110. Kyei-Poku GK, Colwell DD, Coghlin P, Benkel B, Floate KD. On the ubiquity and
1052   phylogeny of Wolbachia in lice. Mol Ecol. 2005;14: 285–294. doi:10.1111/j.1365-
1053   294X.2004.02409.x

1054   111. Covacin C, Barker SC. Supergroup F Wolbachia bacteria parasitise lice (Insecta:
1055   Phthiraptera). Parasitol Res. 2007;100: 479–485. doi:10.1007/s00436-006-0309-6

1056   112. Boore JL. Animal mitochondrial genomes. Nucleic Acids Res. 1999;27: 1767–1780.

1057   113. Shao R, Kirkness EF, Barker SC. The single mitochondrial chromosome typical of animals
1058   has evolved into 18 minichromosomes in the human body louse, *Pediculus humanus*.
1059   Genome Res. 2009;19: 904–912. doi:10.1101/gr.083188.108

1060   114. Dong Y, Zhao M, Shao R. Fragmented mitochondrial genomes of seal lice (family
1061   Echinophthiriidae) and gorilla louse (family Pthiridae): frequent minichromosomal splits
1062   and a host switch of lice between seals. BMC Genomics. 2022;23: 283.
1063   doi:10.1186/s12864-022-08530-8

115. Shao R, Zhu X-Q, Barker SC, Herd K. Evolution of Extensively Fragmented Mitochondrial
Genomes in the Lice of Humans. Genome Biol Evol. 2012;4: 1088–1101.
doi:10.1093/gbe/evs088

116. Baumann P. Biology of Bacteriocyte-Associated Endosymbionts of Plant Sap-Sucking
Insects. Annu Rev Microbiol. 2005;59: 155–189.
doi:10.1146/annurev.micro.59.030804.121041

117. López-Sánchez MJ, Neef A, Peretó J, Patiño-Navarrete R, Pignatelli M, Latorre A, et al.
Evolutionary Convergence and Nitrogen Metabolism in *Blattabacterium* strain Bge,
Primary Endosymbiont of the Cockroach *Blattella germanica*. Bordenstein SR, editor.
PLoS Genet. 2009;5: e1000721. doi:10.1371/journal.pgen.1000721

118. Manzano-Marín A, Oceguera-Figueroa A, Latorre A, Jiménez-García LF, Moya A. Solving
a Bloody Mess: B-Vitamin Independent Metabolic Convergence among
Gammaproteobacterial Obligate Endosymbionts from Blood-Feeding Arthropods and the
Leech *Haementeria officinalis*. Genome Biol Evol. 2015;7: 2871–2884.
doi:10.1093/gbe/evv188

119. Sasaki-Fukatsu K, Koga R, Nikoh N, Yoshizawa K, Kasai S, Mihara M, et al. Symbiotic
Bacteria Associated with Stomach Discs of Human Lice. Appl Environ Microbiol.
2006;72: 7349–7352. doi:10.1128/AEM.01429-06

120. Boyd BM, Allen JM, Nguyen N-P, Vachaspati P, Quicksall ZS, Warnow T, et al. Primates,
Lice and Bacteria: Speciation and Genome Evolution in the Symbionts of Hominid Lice.
Mol Biol Evol. 2017;34: 1743–1757. doi:10.1093/molbev/msx117

121. Puchta O. Experimentelle Untersuchungen uber die Bedeutung der Symbiose der
Kleiderlaus Pediculus Vestimenti Burm. Z Parasitenkd. 1955;17.
doi:10.1007/BF00260226

122. Perotti M, Kirkness E, Reed D, Braig H. Endosymbionts of lice. Insect Symbiosis. Bourtzis
K Mill T Ed. 2009;3.

123. Eberle MW, McLean DL. Observation of symbiote migration in human body lice with
scanning and transmission electron microscopy. Can J Microbiol. 1983;29: 755–762.

124. Buchner P, editor. Endosymbiosis of Animals with Plant Microorganisms. New York:
Wiley; 1965. Available: https://books.google.com/books?id=lcZ3q4oiuYMC

125. Yoshizawa K, Johnson KP. How stable is the "Polyphyly of Lice" hypothesis (Insecta:
Psocodea)?: A comparison of phylogenetic signal in multiple genes. Mol Phylogenet Evol.
2010;55: 939–951. doi:10.1016/j.ympev.2010.02.026

126. McMeniman CJ, Barker SC. Transmission ratio distortion in the human body louse,
Pediculus humanus (Insecta: Phthiraptera). Heredity. 2006;96: 63–68.
doi:10.1038/sj.hdy.6800760

127. Normark BB. The Evolution of Alternative Genetic Systems in Insects. Annu Rev Entomol.
2003;48: 397–423. doi:10.1146/annurev.ento.48.091801.112703

128. De La Filia AG, Andrewes S, Clark JM, Ross L. The unusual reproductive system of head
and body lice (*Pediculus humanus*): Paternal genome loss in human lice males. Med Vet
Entomol. 2018;32: 226–234. doi:10.1111/mve.12287

129. Ashe A, Colot V, Oldroyd BP. How does epigenetics influence the course of evolution?
Philos Trans R Soc B Biol Sci. 2021;376: 20200111. doi:10.1098/rstb.2020.0111

130. Patten MM, Ross L, Curley JP, Queller DC, Bonduriansky R, Wolf JB. The evolution of
genomic imprinting: theories, predictions and empirical tests. Heredity. 2014;113: 119–
128. doi:10.1038/hdy.2014.29

131. Oldroyd BP, Yagound B. The role of epigenetics, particularly DNA methylation, in the
evolution of caste in insect societies. Philos Trans R Soc B Biol Sci. 2021;376: 20200115.
doi:10.1098/rstb.2020.0115

1113

Supporting information

S1 Table. Detail of the human louse samples used in the current study. Groups reflect their geographic distribution and STRUCTURE clustering: Africa-Israel-Spain (AIS), South and Southeast Asia (SAs), East Asia (EAs), Europe (E), North America (NA), Mesoamerica (MA), Central America (CA), and South America (SAm).

S2_S3 Tables. Prior distributions of population sizes parameters and times of events for testing for the demographic scenarios described in Table 2 and Fig 2. Between the brackets [] are the host values obtained from anthropological references as shown in Table 2 of the main manuscript. Louse values were estimated using anthropological references under different louse infestation levels as well as different generational times. **S2 Table.** Population size priors considering 10% and 50% louse infestation level. **S3 Table.** Louse time priors are expressed considering number of generations of lice back in time assuming a generation time of 27 and 36 days, respectively. The values for hominins refer to years.

S4 Table. Genetic differentiation. Louse pairwise genetic differentiation based on F_{ST} values from 15 microsatellite loci among sites where 8 or more lice were collected. Sample sizes are in parentheses following population name. Values below the diagonal indicate F_{ST} values, *p*-values are shown above the diagonal; * indicates significant at nominal alpha level of 0.05.

S1 Fig. Discriminant analysis of principal components (DAPC). Cluster membership of each louse is depicted by a different colour inside their 95% inertia ellipses. DA eigenvalues are shown in the inserted graphs at the bottom right, whereby the numbers of the discriminants plotted against each other are indicated by dark grey and the remaining discriminants retained for the analysis in light grey. A total of 10 axes were retained in this DAPC, this is referred to the n-

dim function in the adegenet package, and indicates the number of retained DAPC axes, which is

affected by both the number of PCA axes and DA axes retained.

S2 Fig. Number of clusters. ΔK statistic of Evanno et al. [39] from STRUCTURE analysis from

Fig. 3. K = 2 (ΔK = 858.853).

S3 Fig. STRUCTURE plot including only non-admixed lice (no hybrids) from both clusters

I and II at K = 4. An ancestry threshold value of q-values as suggested in [38] was used

because it is efficient and accurate at differentiating between non-interbreed and hybrids. Only

individuals with q-value between 0 and < 0.20 or > 0.8 and 1 are classified as non-admixed and

were included in this analysis.

S4 Fig. Cluster I substructure. STRUCTURE plot including only non-admixed lice (no

hybrids).

S5 Fig. Cluster II substructure. STRUCTURE plot including only non-admixed lice (no

hybrids).

S6 Fig. Allelic richness. Allelic richness per nuclear genetic cluster (cI, orange and cII, blue)

using the rarefaction method implemented in the computer program ADZE (Allelic Diversity

Analyzer) Version 1.0 [44]. Because allelic richness measurements considered sample size, the

rarefaction method allows the estimation of allelic richness for different random subsamples of

size "g" from the populations [45,46].

S7 Fig. Correspondence analysis (CA). CA of the nuclear cluster membership frequency and

the mitochondrial haplogroup of head lice.

S8 Fig. Gene flow. Full migration model with four units defined by both the nuclear clusters

inferred from STRUCTURE and the mitochondrial haplogroups: unit 1) cI – mtDNA A; unit 2)

cII – mtDNA A, unit 3) cI – mtDNA B; and unit 4) and cII – mtDNA B. Values above and

below the arrows indicate migration rates, arrows indicated directionality.

S9 Fig. Graphs of posterior probabilities for each of the 63 scenarios per level of parasitism

and generational time. Louse infestation included three levels: 10%, 30% and 50%, where 10%

indicates that 10% of the hominin population have 10 lice per head, the 30% level considers that

each head would have 20 lice and only 30% of the population is infested, while the 50% reflects

high level of parasitism with 50 lice per head in 50% of the population. For this study we

considered the time of head lice maturation from egg to reproductive male adult and egg-laying-

adult female of 18 days, 27 days and 36 days. Each demographic model was color-coded as

follow Scenario 1 – green, Scenario 2 – red, Scenario 3 – light blue, Scenario 4 – pink, Scenario

5 – yellow, Scenario 6 – black, Scenario 7 – grey.

Bering Land Bridge

European colonization