

Full title:

Assessing the intrageneric environmental boundaries of the  
extremophilic cyanobacterium *Chroococcidiopsis*  
(*Pleurocapsales*) and its implications for space exploration

Short title: Environmental boundaries of *Chroococcidiopsis* and  
its implications for space

John R. Cumbers<sup>1,2</sup> and Lynn J. Rothschild<sup>1,3</sup>

Mail Stop 239-20, NASA Ames Research Center,  
P.O. Box 1, Moffett Field, CA 94035

<sup>1</sup>Department of Molecular Biology, Cell Biology and Biochemistry  
Brown University, Box G-W. Providence,  
RI 02912, USA

<sup>2</sup>Current address: SynBioBeta, 818 Grayson Rd Suite 201, Pleasant Hill, CA 94523

<sup>3</sup>Space Science Division, NASA Ames Research Center, Moffett Field, CA 94035

Corresponding author: Lynn J. Rothschild

Office +1 (650) 604-6525, e-mail: [Lynn.J.Rothschild@nasa.gov](mailto:Lynn.J.Rothschild@nasa.gov)

## Abstract

The cyanobacterium *Chroococcidiopsis* contains species found in extreme environments, thus providing the opportunity to study adaptation to a range of environments within the confines of a single genus. Due to its extremophilic nature, it has attracted attention for space settlement as well as a model for life elsewhere. In this study, eight unialgal strains from diverse habitats, isolated in unialgal culture and grown in laboratory conditions, were characterized for their ability to survive a range of extreme environments including UVC (254 nm) radiation, oxidative damage, desiccation, and repeated freeze/thawing. The study revealed two previously uncharacterized saltwater isolates of *Chroococcidiopsis* that were more radiation resistant than most of the other isolates. Isolate CCMP 1991 from Hawaii survived up to  $1750 \text{ J}\cdot\text{m}^{-2}$ , and isolate CCMP 3184 from Samoa survived up to  $1000 \text{ J}\cdot\text{m}^{-2}$  (254 nm UVR) compared with  $250 \text{ J}\cdot\text{m}^{-2}$  for most other isolates tested. These two UV radiation-resistant isolates are closely related phylogenetically, but inhabit different environments. Each was further characterized for its ability to repair DNA damage as assessed by the repair of UV-induced thymine dimers and for oxidative damage tolerance via resistance to  $\text{H}_2\text{O}_2$ -induced (oxidative) damage. Both isolates repaired thymine dimers faster in the light than in the dark with the Hawaiian isolate repairing faster than the Samoan isolate in the light, suggesting repair by photoreactivation. The Hawaiian isolate was more tolerant to  $\text{H}_2\text{O}_2$  exposure than the Samoan isolate, indicating a possible role for antioxidants in the protection of the cell. Both isolates were more tolerant than the other isolates tested to freeze/thawing in liquid nitrogen, which is also known to cause DNA damage. Spectral absorbance scans were performed to detect pigments in each isolate. While all showed peaks likely to be chlorophyll a, carotenoids, phycocyanin, scytonemin and MAAs, the Hawaiian isolate contained a pigment that absorbed at around 325 nm that none of the other isolates contained. Although this pigment is outside the range of UVC absorbance, it is hypothesized that it may play a role in DNA protection as a UV sunscreen or as an antioxidant. The phenotypic similarities in radiation resistance and freeze/thawing resistance among the Hawaiian, Samoan and Negev isolates appear to be the

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

result of environmental adaptation rather than phylogenetic markers as the first of these have been previously shown to be part of a saltwater clade, while the Negev strain falls within a freshwater clade. As pigmentation, and likely resistance to other environmental conditions, can be induced, these data provide a baseline study of strains in an uninduced state. Thus, the true environmental limits to *Chroococcidiopsis* likely go beyond our current knowledge. The implications of this is discussed in relation to space exploration

**Keywords:** *Chroococcidiopsis*; DNA repair; Freezing tolerance; H<sub>2</sub>O<sub>2</sub> tolerance; oxidative damage; pigments; thymine dimers; UV tolerance, desiccation tolerance.

## Introduction

*Chroococcidiopsis* is a genus of photosynthetic cyanobacteria which has long had a reputation for the ability to survive various extreme environments. As such, the genus is of interest to understand the limits of life on Earth and the evolution of the ability to survive one or more extreme environments (Capece et al., 2013). With these abilities, such organisms make excellent model systems for the search for life elsewhere (e.g., Rothschild & Mancinelli, 2001). Imre Friedmann (1995) suggested that *Chroococcidiopsis* would be suitable for growth on Mars as we move towards human exploration and settlement. They have promise for use in making extraterrestrial bodies habitable (Verseux et al., 2015; Billi et al., 2016), for example, by aiding in the production of desert crusts to mitigate dust (Liu et al., 2008). The genetic capabilities of such organisms could provide these capabilities to other organisms of use that may be more space hardy or genetically tractable (Rothschild, 2010).

Members of the genus have been isolated from diverse environments (Fewer et al., 2002), ranging from the frigid Dry Valleys of Antarctica (Friedmann and Ocampo, 1976) to hot deserts such as the Atacama in Chile (Warren-Rhodes et al., 2006; 2007) and the Negev in Israel (Friedmann et al., 1967). These strains have acquired various traits that allow them to live in their respective extreme environments, such as efficient DNA repair mechanisms to survive in high UV radiation (UVR) environments (Billi et al., 2000a), and desiccation tolerance to survive in low water environments (Caiola et al., 1996a).

Solar radiation is a natural environmental variable for organisms that live on the surface of a planetary body, and exposure to radiation between 400-700 nm is obligatory for photosynthesis. Thus, organisms have evolved in direct response to solar radiation. In contrast, the UV radiation (UVR) portion of the spectrum (10-400 nm) is known to cause a range of different biological stresses (reviewed in Franklin and Forster, 1997). UVR damages biomolecules such as DNA and proteins, which have peak absorption at approximately 260 and 280 nm, respectively. Direct damage from UVR in the UVB (280-315 nm) and UVC (100-280 nm) induces thymine dimer formation, where two adjacent thymine residues are fused together on

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

a strand of DNA (Goodsell, 2001). Several DNA repair mechanisms have evolved to reverse the effects of direct UVR-induced damage, including nucleotide excision repair, base excision repair and photoreactivation (Ehling-Schulz and Scherer, 1999; Friedberg et al., 2005). For example, photolyase enzymes are light-activated DNA repair enzymes, known to work by recognizing cyclobutane-type pyrimidine dimers, and converting them back into monomers in the presence of blue/violet light (Ehling-Schulz and Scherer, 1999; Frohnmeier, 2003; Faraji and Dreuw, 2014).

UVR also damages other biological molecules in a variety of ways (rev in Rothschild and Cockell, 1999; Sinha and Häder, 2002). UVR can inhibit growth and enzymatic activity (Kumar et al., 1996; Wu et al., 2005), and is known to cause damage to cell membranes, pigments, and signal transduction pathways (Vincent and Roy, 1993; Häder et al., 2003; Cadoret, 2005; Portwich and Garcia-Pichel, 2007).

*Chroococcidiopsis* has been exposed to solar ultraviolet radiation (UVR) in the open environment (Cockell et al., 2008), and simulated Martian UVR environment in the laboratory (Cockell et al., 2005; Billi et al., 2011). Cockell et al. (2005) exposed dried monolayers of *Chroococcidiopsis* sp. 029 to a simulated Martian surface UVR flux as well as visible radiation, resulting in only 1% viability after 5 minutes, and no viability after 30 minutes. However, a 1 mm layer of Martian regolith simulant sieved to a grain size of <45  $\mu\text{M}$  provided protection, similar to that shown for UVR protection by evaporitic salt crusts (Rothschild, 1990). Further studies by Billi's lab in preparation for ESA's BioMEX (*BIOlogy and Mars EXperiment*) flight experiment aboard the International Space Station (ISS), confirmed that mineral analogs provide protection to the integrity of the DNA and photosynthetic pigments, even after exposure to 500  $\text{kJ}\cdot\text{m}^{-2}$  of polychromatic UVR and space vacuum ( $10^{-4}$  Pa), exposures that correspond to conditions during a one-year period in Low Earth Orbit (LEO) (Baqué et al., 2014).

Several studies have been conducted through controlled exposure of *Chroococcidiopsis* to UVC radiation at 254 nm. For example, Baqué et al. (2013a) showed that the DNA of dried biofilms

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

of *Chroococcidiopsis* sp. CCME 029 maintained DNA integrity virtually identical to unexposed liquid culture controls after exposures of up to  $1.000 \text{ J}\cdot\text{m}^{-2}$  at 254 nm.

Cellular aggregation is correlated with resistance to some environmental factors, including UVR. For example, a dried biofilm of CCME 029 exposed to  $10 \text{ kJ}\cdot\text{m}^{-2}$  of UVC radiation protects the lower cell layers by the top cell layers (Baqué et al., 2013a). The cells in the top layer had damaged photosynthetic pigments as suggested by yellowish autofluorescence, while the lower cell layers maintained a red autofluorescence characteristic of chlorophyll. Liquid cultures withstood exposure to  $13 \text{ kJ}\cdot\text{m}^{-2}$  from a germicidal lamp, which Baqué et al. (2013b) ascribe to multiple factors, including cell aggregation which protects some cells from the lethal effects of radiation.

Some strains of *Chroococcidiopsis* are remarkably resistant to x-ray irradiation (0.01-10 nm), even though unlike UVR, naturally occurring exposure is minimal on Earth. Billi and colleagues (2000) exposed ten different isolates from desert and hypersaline environments to x-rays; all showed between 35% and 80% survival at 2.5 kGy, where a Gy is defined as the absorption of one joule of radiation per kg of matter. Survival was reduced 1 to 2 orders of magnitude when the exposure was doubled, and no cyanobacteria survived exposure at 20 kGy. This is nearly as resistant as the most radiation-resistant organisms known can tolerate. For example, Rainey et al. (2005) were able to recover live *Deinococcus*, *Geodermatophilus*, and *Hymenobacter* from the arid soil of the Sonoran Desert after exposure to doses of 17 to 30 kGy. A subsequent study in Billi's lab was performed on two strains isolated by Roseli Ocampo-Friedmann:

*Chroococcidiopsis* spp. CCME 029, a cryptoendolith isolated from the sandstone of the Negev Desert (Israel), and CCME 057, isolated from chasmoendolithic growth in granite in the Sinai Desert (Egypt). In the desiccated state, both strains survived x-ray exposures up to 5 kGy without detectable damage to either their DNA or membrane integrity (Verseux et al., 2017). To put this in perspective, the average radiation dose from an abdominal X-ray is 0.7 mGy (0.0007 Gy). However, it should also be noted that the effect is correlated with rate of exposure: whereas 5 Gy is lethal to humans in a short period of time, it may be tolerated over a lifetime.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

Thus, comparisons between studies with different dose rates are not equivalent. With reference to potential exposure on Mars, an absorbed dose rate of 76 mGy year<sup>-1</sup> was measured by the Curiosity rover at the surface of Gale Crater (Hassler et al., 2013).

A byproduct of UVR, reactive oxygen species (ROS), can in itself cause oxidative damage to biomolecules (He and Häder, 2002; Pattanaik et al., 2007). The resulting damage often includes single-strand DNA breaks, and the production of 8-hydroxyguanosine residues (Cooke, 2003). Low levels of x-ray radiation also may cause oxidative damage indirectly (Puthran et al., 2009). Repair mechanisms, such as excision repair, can be used for both direct radiation and oxidative damage (rev. in Rastogi et al., 2010).

One mechanism for resistance to radiation and oxidative damage is the production of UVR-absorbing and antioxidant pigments. Mycosporine-like amino acids (MAAs) are some of the strongest natural compounds that absorb in the UVA and are found in a wide phylogenetic variety of organisms including cyanobacteria, red algae, dinoflagellates, corals, and many marine invertebrates. In addition to MAAs, the UVR-absorbing pigments scytonemin is widely distributed among the cyanobacteria, including *Chroococcidiopsis* where scytonemin is found extracellularly in the sheath (Sinha, 2000; Dillon et al., 2002; Řezanka et al., 2004). Some photopigments such as melanin, bacterioruberin and the carotenoids are also known to act as antioxidants (Geng et al., 2008; Karsten and Garcia-Pichel, 1996; Shahmohammadi et al., 1998; Stahl and Sies, 2003; He and Hader, 2002a; b).

A link between desiccation and radiation resistance was established in the exceptionally radiation-resistant bacterium *Deinococcus radiodurans* (Mattimore and Battista, 1996; Battista, 1997). *Chroococcidiopsis* is known to withstand desiccation, although there is strain-specific variation. Faglierone et al. (2017) revealed two orders of magnitude difference in the ability of 10 strains of *Chroococcidiopsis* to survive four years of air-dried storage. Differences even occurred among isolates collected from the same area.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

The ability of *Chroococcidiopsis* to withstand oxidative damage as assessed by protein oxidation was tested after four years of desiccation (Fagliarone et al., 2017). Strain CCME 029 from the Negev desert was resistant to 30 min exposure up to concentrations of 500 mM H<sub>2</sub>O<sub>2</sub>, whereas at the other end of the spectrum, CCME 584(M) from the Gobi Desert and *C. thermalis* PCC 7203, which was isolated from a soil sample near Greifswald, East Germany, showed significant protein carbonylation at only 10 mM H<sub>2</sub>O<sub>2</sub>. This tolerance to oxidative damage correlated with resistance to  $\gamma$ -ray exposure (Fagliarone et al., 2017).

Billi (2009b) did not find genomic fragmentation, nor loss of membrane integrity in CCME 029 (Negev) Cells that had survived desiccation for four years. Protection could be by the production of antioxidant enzymes such as superoxide dismutase (Caiola et al., 1996b, Billi, 2009b), an enzyme known to reduce oxidative stress, and carotenoids (Baqué et al., 2013b). Combined with the evidence of DNA repair, it seems that *Chroococcidiopsis* may survive in extreme conditions by both DNA protection by antioxidants and by DNA repair.

*Chroococcidiopsis* is thought to be able to withstand high levels of X-ray radiation and desiccation due to an efficient DNA repair mechanism (Billi et al., 2000a), possibly the same mechanism as that which is responsible for UVR resistance (Cockell et al., 2005; Billi, 2009a) as both forms of radiation are known to cause strand breakage in DNA. Double-strand breaks in the genomes of a number of *Chroococcidiopsis* isolates, including 029 (Negev) and 171 (Antarctica), were observed following X-ray irradiation by fragmented DNA on an agarose gel (Billi et al., 2000a). This damage was repaired within 24 hours, demonstrating that an efficient DNA repair system existed. Thus, as in *D. radiodurans*, a link between desiccation resistance and radiation resistance in *Chroococcidiopsis* seems likely, although the mechanism of resistance differs as *D. radiodurans* repairs DNA breaks, while *Chroococcidiopsis* prevents them.

Cyanobacteria were responsible for the rise in atmospheric oxygen levels, allowing for the formation of the stratospheric ozone layer. Because cyanobacteria had to access photosynthetically-active radiation at the time when UVR flux on the surface of the Earth should have been substantially higher, cyanobacteria evolved a variety of UVR-protection



Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

mechanisms. These include avoiding damage by physical barriers such as living under minerals (e.g., Baqué et al., 2014), or mitigating (e.g., Bebout and Garcia-Pichel, 1995) or repairing damage through physiological means. For example, planktonic *Chroococcidiopsis* are not as resistant to environmental extremes as those in a dried biofilm, including in low Earth Orbit through the EXPOSE R-2 experiment on ISS (Billi et al., 2019) due to both self-shielding and increased production of extracellular polymers.

With interest in *Chroococcidiopsis* high for its potential in space exploration and as a member of the toolkit for space synthetic biology, this study was undertaken to assess intra-generic phenotypic diversity, and to see if a correlation existed between radiation resistance, photolyase activity and pigmentation. The objective was to characterize previously unstudied isolates and compare them to well-studied isolates (e.g., CCMEE 029 Negev and PCC 7203, *C. thermalis*) in order to find novel mechanisms that allow them to survive in extreme environments on Earth in order to transfer these mechanisms to other organisms for the possible use in extreme environments of space (Rothschild, 2010). Although numerous isolates of *Chroococcidiopsis* have been identified and isolated, only a small number of isolates have been studied simultaneously for their abilities to survive in different extreme conditions, and few comparative studies have been conducted to determine the range of adaptive capabilities and mechanisms that have evolved within the genus.

This study was undertaken with the expectation that it would reveal useful phenotypes that could be exploited for space exploration, either by sending *Chroococcidiopsis* or the genes derived from it into space, or by using it as a model organism in the search for life elsewhere. Thus, the focus was on tolerance to UV radiation, desiccation, freeze/thawing and oxidative damage. The mechanism for resistance was approached by assessing the ability of select strains to repair UV-induced thymine dimers in the light and dark, and by obtaining absorbance spectra for each strain. Only strains that were in unialgal culture were selected as they would be the most likely to be used for future study.

## Results

Eight unialgal cultures of *Chroococcidiopsis*, representing a range of environmental sources (Table 1), were assessed for their environmental tolerances. The data indicate a broad range of environmental preferences that are species or strain specific.

### UVR tolerance

Radiation tolerance was assessed by irradiation with UVC radiation at 254 nm, a wavelength commonly used as a germicidal source. Dose response curves are plotted in Fig. 1. The cells were spotted on the surface of agar plates prior to irradiation to remove the possibility of attenuation by the culture medium, and to minimize self-shading. The figure shows that a range of UVR tolerances exist among the isolates tested. Two isolates survived similarly large doses of radiation: 1991 (Hawaii) which survived up to  $1750 \text{ J}\cdot\text{m}^{-2}$ , and 029 (Negev, Israel) which survived up to  $1500 \text{ J}\cdot\text{m}^{-2}$ . Isolate 3184 (Samoa) had moderate radiation resistance with survival up to a dosage of  $750 \text{ J}\cdot\text{m}^{-2}$ . Isolate 171 (Antarctica) show some survival at  $250 \text{ J}\cdot\text{m}^{-2}$ , but no survival by  $500 \text{ J}\cdot\text{m}^{-2}$ . The remaining four strains (2116, Madagascar; 1489, Galapagos; 7203, Germany; 6712, California) were killed by the lowest dosage of irradiation used in the experiment,  $250 \text{ J}\cdot\text{m}^{-2}$ . *Escherichia coli* (Strain TOP10) was irradiated for comparative purposes, and also did not survive the dosage of  $250 \text{ J}\cdot\text{m}^{-2}$ .

Previous work has shown that CCMP 1991 (Hawaii) and CCMP 3184 (Samoa) are phylogenetically similar, clustering close to each other in a saltwater clade of *Chroococcidiopsis* (Cumbers and Rothschild, 2014). Because these two isolates are phylogenetically similar but differ in radiation phenotypes, they were investigated in more detail. When triplicate samples of each isolate were exposed to  $250 \text{ J}\cdot\text{m}^{-2}$  of UVC radiation and immediately tested for accumulated damage, CCMP 1991 (Hawaii) contained  $73,697 \pm 1.7\%$  thymine dimers per Mb of DNA, 8.4% more than CCMP 3184 (Samoa) which contained  $67,958 \pm 2\%$ . The subsequent repair of the thymine dimers was investigated in both isolates over time as shown in Fig. 2. Thymine dimer repair occurred faster in in the light in isolate CCMP 1991 (Hawaii) than in

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

isolate CCMP 3184 (Samoa). After four hours, isolate CCMP 1991 (Hawaii) had repaired 51% of the thymine dimers in contrast to only 20% in CCMP 3184 (Samoa). For both isolates, thymine dimer repair takes place faster when cells are left to recover in the light than in the dark (Fig. 3).

### Oxidative damage tolerance

The ability of the strains to withstand oxidative damage was determined by mixing the cells with varying concentrations of H<sub>2</sub>O<sub>2</sub>, and then plating them on agar plates to determine subsequent colony-forming ability. Isolate 1991 (Hawaii) was more tolerant to H<sub>2</sub>O<sub>2</sub> than isolate 3184 (Samoa) as shown in Fig. 4. Isolate 1991 (Hawaii) survived up to 100 mM H<sub>2</sub>O<sub>2</sub>, an order of magnitude more than isolate 3184 (Samoa).

### Desiccation tolerance

Cell survival of air-dried cells after 9 months of storage at room temperature in the dark is shown in Table 2. When the dried cells were re-suspended, only isolate PCC 6712 (California) and CCMP 1489 *C. polansiana* (Galapagos) failed to produce colonies. Thus, under a simple air-drying protocol with storage in the dark, nearly all the strains are desiccation tolerant.

### Freeze/thaw tolerance

Repeated freeze/thawing of cells in liquid nitrogen (Fig. 5a) showed that 1991 (Hawaii) and 3184 (Samoa) both withstood multiple freeze/thaw cycles up to 16 rounds, while the majority of other isolates died after 12 rounds (data not shown.) Isolate 7203 *C. thermalis* died after a single round of freeze/thawing in liquid N<sub>2</sub>. The effect of temperature was assessed by conducting experiments where the cells were frozen in an ethanol/dry ice slurry, which has an approximate temperature of -68 °C, rather than liquid N<sub>2</sub>, which is -196 °C. Isolate 1991 (Hawaii) and 3184 (Samoa) were subjected to freeze/thawing in an ethanol/dry ice slurry (approximately -68 °C). In contrast to the results from freezing in liquid N<sub>2</sub>, neither isolate appeared to be affected by repeated freezing and thawing at the warmer temperature.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

## Pigment absorption

Pigments were extracted in methanol, per standard procedures. The absorbance spectra represent a sampling every 5 nm from 190 to 1000 nm. The spectra show several features expected of cyanobacteria, notably the characteristic peaks of chlorophyll *a* at around 432 and 665 nm (Rowan, 1989, Table 3.2), carotenoids around 490 nm (Erokhina et al., 2002), and phycocyanin at 620 (Sinha et al., 1999; Erokhina et al., 2002). Phycoerythrin (560 nm; Sinha et al., 1999) was not detectable.

Strong absorbance is detected in the UV. The varying strength peaks between 200 and 250 and 325 nm are likely to be the UV-screening pigments scytonemin and mycosporine-like amino acids, based on published studies where the absorption was confirmed by HPLC in the cyanobacterium *Lyngbya* (Sinha et al., 1999, Garcia-Pichel and Castenholz, 1993.) This is especially true of species 1991 (Hawaii) which has a pigment absorbing in the low 300s that is absent in the other species under the growth conditions of this experiment. The identification was not confirmed in this study.

## Discussion

The data presented in this study demonstrate that there is a diversity of phenotypes within the genus *Chroococcidiopsis*. The results show a range of UVR tolerance, indicating that two of the isolates are more tolerant to UVR than the rest. Isolate 1991 (Hawaii) and 029 (Negev) both show similar radiation resistance as seen in Fig. 1. Isolate 029 (Negev) is a well-studied isolate of *Chroococcidiopsis*, and has been shown to be also resistant to X-ray irradiation and desiccation (Billi et al., 2000a) via mechanisms that involve both DNA repair (Billi et al., 2000a) and anti-oxidant production (Caiola et al., 1996b). Isolate 029 (Negev) has also been shown to be UV radiation resistant (Cockell et al., 2005), so it was anticipated that this isolate would show high radiation resistance in this study too. Here two further isolates *Chroococcidiopsis* that were previously uncharacterized have been identified as UV radiation resistant. Isolate 1991 (Hawaii) is of a similar resistance to 029 (Negev) and isolate 3184 (Samoa) is about half as radiation resistant in comparison.

The results from the radiation exposure, repair and pigmentation experiments presented here, suggest that the two previously uncharacterized isolates survive UV radiation using a combination of DNA protection and repair mechanisms. Isolate 1991 (Hawaii) is more resistant to H<sub>2</sub>O<sub>2</sub>, indicating that it may contain enzymes such as superoxide dismutase (SOD) that play a role in reducing oxidative damage (Caiola et al., 1996b; He and Häder, 2002). Isolate 1991 (Hawaii) is also faster at repairing thymine dimers in its DNA than isolate 3184 (Somoa), indicating that it may have a better DNA repair mechanism.

The large error bars on the dose response curves are due to the clumping of the *Chroococcidiopsis* cells. In general, larger cellular aggregates were observed to contain more cells per aggregate which would likely decrease the chance of genome damage occurring per aggregate, and increase the chance of survivability of at least some members of the colony-forming unit. Isolate 1991 (Hawaii) has tighter packed cells than isolate 3184 (Samoa) which could contribute to the enhanced DNA repair ability observed. *D. radiodurans* is also known to form cellular aggregates which again indicate that this may be part of the DNA protection

and/or repair mechanism (Cox and Battista, 2005). So the very characteristic that makes *Chroococcidiopsis* cells difficult to disaggregate may also be the key to its radiation resistance. Billi et al. (2000a) suggested that part of the reason that the genus as a whole may be able to survive such extremes is linked to the morphology of the genus, where mother cells contain daughter cells (baeocytes) at various stages of active cell division. Thus, the fact that the cells were clumping and have baeocytes may contribute to their environmental resistance.

DNA repair was shown to take place faster in the light than in the dark in both 1991 (Hawaii) and 3184 (Somoa), although there was repair in both the light and the dark (Fig. 3). This suggests that a major repair is a light-dependent DNA repair mechanism. Photolyase genes such as the photorepair gene *phrA* (Hendrischk et al., 2007) are likely responsible for the light repair system, and other mechanisms, such as excision repair, for the dark repair. In particular, isolate CCMP 1991 (Hawaii) appears to contain a faster DNA repair mechanism occurring in the first two hours that is not as apparent in the first two hours of DNA repair in the Samoan (CCMP 3184) isolate. After irradiation, CCMP 1991 (Hawaii) contained more thymine dimers per Mb of DNA than the Samoan isolate CCMP 3184, but it was able to repair these thymine dimers faster, indicating that DNA repair may be more important than DNA protection in this isolate.

The linkage between radiation and desiccation tolerance is known to be through DNA repair (Mattimore and Battista, 1996), as discussed above. However, a link between freeze/thaw tolerance and DNA repair has not been established but could be an important future avenue of research. DNA repair may be a common link between many forms of environmental stresses faced by organisms due to the central role that the molecule plays as an information carrier in the cell. This study focused only on cell survival and DNA damage/repair, and this was limited to thymine dimer damage. Future studies could also investigate other forms of DNA damage that are known to take place (Friedberg, 2005) as well as protein oxidation, which has also been implicated in bacterial radiation resistance (Daly et al., 2007). Faglierone et al. (2017) demonstrated a clear correlation between desiccation and radiation resistance, and avoidance of protein oxidation. As they point out, mechanistic studies should be conducted examining the

possibility of an ROS-scavenging strategy based on a Mn-dependent system such as in *D. radiodurans* (Daly et al., 2004) and for other desiccation- and radiation-tolerant bacteria (Paulino-Lima et al., 2017). Desiccation-resistant organisms usually accumulate protective compounds such trehalose and/or saccharose. Trehalose forms hydrogen bonds with polar residues in proteins and phospholipids in cell membranes and may therefore substitute for water molecules that normally comprises the ‘hydration shell’ surrounding hydrophilic biomolecules, as well as forming a molecular ‘glass’ (vitrification) at biological temperatures (rev. in Morano, 2014). Future mechanistic studies will help explain the higher resistance of desiccated vs hydrated cells to ionizing radiation (Verseux et al. 2017), as well as their endurance when exposed in the dried state to ultraviolet doses simulating 1-year duration in low earth orbit (Billi et al. 2011; Baqué et al. 2013b).

A number of unexpected results exist in the data. Isolate 171 from Antarctica has been previously shown to be resistant to X-ray radiation (Billi et al., 2000a), but in this study, it died after a relatively low UVR dosage of 250 J·m<sup>2</sup> (Fig. 1). The isolate was also not particularly resistant to freeze/thawing in liquid nitrogen (Fig. 5) when the opposite might be expected from an isolate living in an environment such as Antarctica that undergoes regular freeze/thawing. However, both were resistant to the freeze/thaw regime in the dry ice/ethanol slurry, which is a far closer temperature to that found in the Antarctic (-68 °C, rather than liquid N<sub>2</sub>, which is -196 °C).

The pigment scans in Fig. 6 suggests a pigment in isolate 1991 (Hawaii) that has an absorption peak around 325 nm. This absorption is just in the range of UVA, but distant from 254 nm (UVC), which was the wavelength of the radiation source used in this study. This suggests that this pigment is unlikely to play a role in the protection of the organism from UVC shown here. However, the pigment may still play a role in protection of the organism from UVB. The pigment shows a similar absorbance profile to that of the carotene lycopene (Sicilia et al., 2005), the carotenoid 4-ketomyxol 2'-fucoside (Takaichi, 2005), or a mycosporine-like amino acid (MAA) or scytonemin, both well known from cyanobacteria.

A word of caution should be noted on the interpretation of these data, as well as prior studies with respect to the ability of isolates to withstand environmental extremes. Pigment production responds to the radiation environment, with humans producing more melanin in high UVR environments, and plants changing their pigmentation in response to light levels. Similarly, the production of MAAs and scytonemin are known to be inducible in cyanobacteria (e.g., Garcia-Pichel and Castenholz, 1993; Ehling-Schulz et al., 1997; Sinha et al., 2003), including in *Chroococcidiopsis* (e.g., Dillon et al., 2002). Thus, the pigmentation profiles in Fig. 6, and many prior studies, may reflect intrinsic pigment production after a number of years in culture in the absence of UVR, rather than the capabilities that these isolates would have shown when collected from their natural environments. The culturing of *Chroococcidiopsis* in the laboratory can, over time, make it less resistant to UV radiation and desiccation (Olsson-Francis and Cockell, 2010a). As the pigmentation is likely affecting the tolerance to UVB irradiation and other stresses, the data presented here should be taken as baseline data from organisms not recently exposed to UVR, and thus provide a pathway to further studies testing environmental resistance and pigmentation after induction.

Based on 16s rRNA and gltX sequence analysis, we suggested previously that the genus *Chroococcidiopsis* contained a saltwater clade within an ancestral freshwater taxon (Cumbers and Rothschild, 2014), and may be polyphyletic. Komárek et al. (2014) have suggested splitting the genus, with the type species, *C. thermalis*, being removed, but the other more extremophilic strains remaining in *Chroococcidiopsis sensu stricto*. The data presented in Fig. 1 might be interpreted as a phylogenetic correlation between UV resistance and the saltwater clade, but the high resistance of the strain from Negev desert (CCMEE 029), which is within the freshwater clade, suggests that this characteristic is an environmental adaptation, and thus polyphyletic. A similar situation exists for the freeze/thaw data presented in Fig. 5, which also shows the Negev phenotypically grouped with members of the saltwater clade, suggesting phenotypic plasticity. Fagliarone et al. (2017) also assessed the relationship between phylogeny and phenotypic characteristics in *Chroococcidiopsis*. These included representatives from



Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

deserts worldwide, including the Middle East (Israel, Egypt), Central Asia (Mongolia), South America (Chile) and Antarctica, and the same strain used here to represent the Negev, CCME 029, and showed multiple lineages but widespread resistance to desiccation. Even isolates from the same area showed differences in environmental resistance. As Fagliarone et al. (2017) focused on strains from deserts, none of the strains were examined that Cumbers and Rothschild (2014) that they had previously identified as the “saltwater clade” were included. With baseline data suggesting that 1991 (Hawaii) could be as environmentally robust as 029 (Negev), this strain should be included in future comparisons.

### Implications for space exploration

Cyanobacteria could play an important role in space exploration, from ecopoeisis (making Mars habitable to earthlings) to biological *in situ* resource utilization (ISRU) and human life support (Graham, 2004; Olsson-Francis and Cockell, 2010b; Verseux et al., 2015). Friedmann (1995) first proposed *Chroococcidiopsis* in particular as a suitable organism for terraforming Mars because of its resistance to harsh environmental conditions. While a few terrestrial organisms could probably survive the temperatures and pressures on the surface of Mars (Rothschild, 1990), it is unlikely that any terrestrial organism would thrive under those conditions. Even *Chroococcidiopsis* in desert rocks of the Negev need a minimum of –6.9 MPa and 90% RH at 20°C to photosynthesize (Palmer and Friedmann, 1990). Still, the use of polyextremophiles to enable human colonization should result in lower energy expenditures, and thus enhanced mission capabilities as the power is used for other activities.

To test the capabilities of *Chroococcidiopsis* to survive the space environment, the Biofilm Organisms Surfing Space (BOSS) experiment, part of the EXPOSE-R2 space mission exposed cells that were dried as biofilms and compared them to ones dried in the planktonic form (Billi et al., 2019.) Radiation fluences ranged from solar radiation and space vacuum to a Mars-like UVR flux and atmosphere. Upon return to Earth, cells were rehydrated and tested for damage to cell division, the genome and photosynthetic apparatus. Under all conditions, including dark, laboratory-based ground controls, the biofilms recovered better than the planktonic forms,

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

possibly due to the abundance of extracellular polymeric substances, and physiological changes induced by drying. This result confirmed the BOSS team's prior ground-based studies (Baqué et al., 2013a). While normally biotechnology uses bioreactors as production platforms for microbes, the BOSS results suggest that a biofilm configuration would be a better option for Mars as it would allow the system to better withstand more environmental exposure than planktonic cells, thus potentially lowering infrastructure costs. However, production may be lower in the biofilm than in the planktonic mode, so this would have to be taken into account.

*Chroococcidiopsis* was exposed to space on a second mission on EXPOSE-R2 facility on ISS, BIOMEX (BIOlogy and Mars EXperiment). For this mission, lichens, archaea, bacteria, cyanobacteria, snow/permafrost algae, meristematic black fungi, and bryophytes from alpine and polar habitats were embedded, grown, and cultured on a mixture of martian and lunar regolith analogs or other terrestrial minerals. The organisms and regolith analogs and terrestrial mineral mixtures were then exposed to space and to simulated Mars-like on the EXPOSE-R2 facility. *Chroococcidiopsis* showed survival and recovery on Mars analog minerals, and low DNA-damage particularly in the more protected endolithic dark control areas (de Vera et al., 2019). This result is not dissimilar to prior two-week exposure studies on the free-flying BioPan platform with *Bacillus subtilis* spores where the survival rate was increased by 5 orders of magnitude and more if the spores in the dry layer were directly mixed with powder of clay, rock or meteorites (Horneck et al., 2001). Up to 100% survival was reached in soil mixtures with spores comparable to the natural soil to spore ratio.

While extremophiles for the production of resources in space is of interest to the space community, the addition of synthetic biology greatly expands the potential for biology-based production (Rothschild, 2016). Towards that goal of making *Chroococcidiopsis* a tractable organism for genetic manipulation, Billi et al. (2001) demonstrated gene transformation in several strains of *Chroococcidiopsis* using electroporation and conjugative transfer and showed that CCME 029 was particularly efficient in conjugative transfer. Further, she created two GFP-based plasmids for *Chroococcidiopsis* CCME 029 and CCME 123. Both were maintained in the

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

host cells after 18 months of dry storage and rehydration. Equally exciting, Billi and colleagues (2000b) transformed *Escherichia coli* BL21DE3 with sucrose-6-phosphate synthase (SpsA) from the cyanobacterium *Synechocystis* sp. strain PCC 6803, conferring a 10,000-fold increase in survival following freeze-drying or air drying. *Synechocystis* sp. strain PCC 6803 is commonly used in industry and has some tolerance to short-term heat stress, but was isolated from a freshwater lake and has an optimal temperature for growth of 32-38°C (Červený et al., 2015). We propose the exciting possibility of mining the genome of the more extremophilic *Chroococcidiopsis* and using these genes to transform more commonly used synthetic biology chassis organisms such as *Bacillus subtilis* for space use.

This study provides a foundation for performing comparative genomics on isolates of *Chroococcidiopsis* that are phylogenetically similar, but phenotypically diverse, such as CCMP 1991 (Hawaii) and CCMP 3184 (Samoa). These two isolates showed high baseline levels of UV radiation resistance without induction and may prove interesting candidates for future studies in addition to the better studied 029 (Negev). Full genome sequencing or transcriptomics under environmental induction, including UVR, desiccation etc. should illuminate which mechanisms are at play and the relative contributions of DNA protection and repair. Both could lead to a fuller understanding of how *Chroococcidiopsis* survives in extreme environments and how other organisms might one day be engineered to do the same. A confirmation of the identification of UVR-absorbing peaks and their correlation to environmental resistance will allow a fuller understanding of the environmental resistance capabilities of this extremophile genus.

## Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

The goal of the study was to begin to understand the mechanisms responsible for survival in extreme environments and to use those as a toolkit to engineering other organisms for survival under such conditions. This was approached by characterizing the response of unialgal isolates from culture collections to the environmental stresses of UVR, oxidative damage desiccation and freeze/thawing. The results indicated that resistance to UV radiation in uninduced strains is not as widespread in the genus as may have been previously thought. Two isolates were examined in detail for their ability to repair thymine dimers in DNA following radiation and for their ability to survive exposure to H<sub>2</sub>O<sub>2</sub>. Although neither of the mechanisms for these traits has been elucidated, a number of isolates of interest were identified and preliminary work on dissecting the mechanisms of DNA protection and repair were demonstrated. It remains to be seen what the extremophilic capabilities of the genus are after environmental, particularly radiation, induction. The true environmental limits to *Chroococcidiopsis* likely go beyond our current knowledge, making the genus even more interesting for both astrobiology and human space exploration.

## Materials and methods

### Cyanobacterial cultures and culturing

The eight isolates used in this study are listed in Table 1. Each was obtained as a unialgal culture from the Provasoli-Guillard National Center for Culture Collection of Marine Phytoplankton (CCMP; since 2011 the NCMA, the National Center for Marine Algae and Microbiota), the Pasteur Culture Collection of Cyanobacteria, The University of Oregon's Culture Collection of Microorganisms from Extreme Environments (CCMEE) and Sammlung von Algenkulturen (SAG) at the University of Göttingen (Göttingen, Germany). We are grateful to the late R. Castenholz for the kind gift of isolates from the CCMEE, and to R. Anderson for the kind gift of isolates from the CCMP. The isolates were chosen to represent the geographical, environmental, and phylogenetic diversity of the genus *Chroococcidiopsis* while also being available in unialgal culture. Freshwater isolates were grown in BG11 growth medium (Stanier et al., 1971), made from a 50x concentrate (Sigma-Aldrich Inc., St. Louis, MO, USA, cat. # C3061-500ML). The pH of the BG11 was adjusted to 7.4 before filter sterilization. Saltwater isolates also were grown in BG11, with the addition of 4% (40 g·L<sup>-1</sup>) sea salt (Sigma-Aldrich Inc., St. Louis, MO, USA cat. # S9883).

Some strains of *Chroococcidiopsis* grow in cellular aggregates making plating of homogeneous monolayers difficult. In order to disaggregate cell aggregates, liquid cultures in mid-exponential phase were first sonicated gently (Ultrasonic processor XL, microtip size 1/16, repeated rounds of three seconds, Misonix Inc, Farmingdale, NY, USA) until the optical density at 730 nm had plateaued, suggesting that large clumps of cells had separated. Cells were then centrifuged at 1000 x g, the supernatant discarded, and the cells resuspended in fresh medium to a concentration of 1·10<sup>6</sup> colony-forming units mL<sup>-1</sup> as confirmed by visual inspection with a hemocytometer. Cells were plated on agar plates of the same medium used to grow the isolates, cultured overnight, and inspected the following day to confirm that the sonication had produced a homogeneous monolayer of viable cells. As with previous studies of *Chroococcidiopsis*, a single cell or cell aggregate was counted as a colony forming unit (Billi et al., 2000a). As isolate 1991 (Hawaii) had a morphology showing smaller, denser cell aggregates

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

than 3184 (Samoa), to compare the DNA repair ability of these two isolates, cells were normalized to  $6 \times 10^7$  cells·mL<sup>-1</sup> rather than normalizing to colony-forming units.

All growth and survival assays were performed by spotting replicate aliquots of 10 µL of cells onto agar plates, unless otherwise stated in the figure legends. Plates were sealed with parafilm to reduce evaporation before being placed in a growth incubator (Rexmed Inc., Taiwan) at 25°C under a 16h:8h light/dark regime and a mix of Philips TLD36W/840 (cool white) and Philips TLD36w (aqua) fluorescent light at  $54 \pm 2$  µmol photons·m<sup>-2</sup>·s<sup>-1</sup> as measured with a PAR detector (PMA2100/PMA2132, Solar Light Company, PA, USA). The Model PMA2132 Digital Quantum Light Sensor measures photon flux in the wavelength range 400 to 700 nm.

### Microscopy and image analysis

Agar plates were individually photographed on a light table using a digital camera. Micrographs were taken using a Zeiss Axio Imager Z1 with a DIC filter and a binocular microscope, Zeiss Stemi 2000-c (Carl Zeiss MicroImaging, Thornwood, NY, USA). ImageJ (Abramoff et al., 2004) was used to measure relative cell density. Each agar plate image was cropped to size and converted to a 32 bit grey scale image. The background was subtracted, and light density was measured using the Gel Analyzer tool in ImageJ (Abramoff et al., 2004).

### UVR tolerance

To determine the UVR tolerance of each isolate, cells were first plated onto agar plates in 10 µL spots as described above. The agar plates were then covered with sterile Petri dish lids and left for 24 hours before irradiation. After this time, lids were removed, and cells were placed in a sterile hood 44 cm beneath a UVC light source of 254 nm for different lengths of time to achieve various dosages of radiation. The intensity of the UVC source was 42 µWs·cm<sup>2</sup> at 254 nm, measured with a high-resolution spectrometer (HR4000, Ocean Optics, Dunedin, FL, USA). Dosages were also measured in real time using a radiometer (PMA2100, Solar Light Company, Glenside, PA, USA). For comparison, *Escherichia coli* (isolate TOP10, Invitrogen, Carlsbad, CA

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

USA) was irradiated after plating cells on LB medium in the same manner as the *Chroococcidiopsis*.

### Thymine dimer detection

Thymine dimers were measured using a dot blot assay according to Sinha et al. (2001) as modified by Leuko et al. (2011). DNA from irradiated cells was extracted using a DNeasy Plant Mini Kit (Qiagen, Gaithersburg, MD, USA) according to the manufacturer's instructions. The quality (OD 260/280 ratio) and quantity of the extracted DNA was measured on a Nanodrop (ThermoScientific, Wilmington, DE, USA). The DNA was diluted to 2 ng·μL<sup>-1</sup>, NaOH added to 1/10<sup>th</sup> volume, and incubated at 80° C for 30 min. The denatured DNA was then blotted onto a Hybond LFP membrane using a Minifold I dot blot system (both from GE Healthcare, Piscataway, NJ, USA) followed by a wash with 400 μL TE buffer per well. The membrane was dried for 30 min at 80° C in a drying oven before being blocked with 5% ECL advanced blocking reagent (GE Healthcare, Piscataway, NJ, USA) for one hour. The unwashed membrane was incubated in mouse anti-thymine dimer primary antibody (#MC-062, Kamiya Biomedical, Seattle, WA, USA) at a 1:3000 dilution in PBS-T buffer for two hours at room temperature. The membrane was then washed twice with PBS-T followed by a one hour incubation in a secondary antibody, sheep anti-mouse IgG with Cy3 labeling (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA, cat # 515-165-062) at a 1:1,000 dilution in PBS-T with 5% ECL advanced blocking reagent. The membrane was washed twice in PBS-T before being visualized on a Typhoon Trio Scanner (GE Healthcare, Piscataway, NJ, USA) and quantified with ImageJ (Abramoff et al., 2004).

### Oxidative damage tolerance

To assess the effect of an oxidant on survival, the cells were exposed to hydrogen peroxide. H<sub>2</sub>O<sub>2</sub> dilutions were freshly prepared from a 9.7 M, 30% v/v stock of H<sub>2</sub>O<sub>2</sub> (Sigma-Aldrich Inc., St. Louis, MO, USA) in fresh media. Cells were thoroughly mixed 1:1 for each dilution in PCR tubes by pipetting up and down, immediately plated and growth assayed as described above.

### Desiccation tolerance

Desiccation tolerance was assessed in air-dried cultures, kept in the dark at room temperature, according to Billi (2009b) and Caiola et al. (1993). After 9 months, cells were rehydrated by suspension in 50  $\mu$ L of sterile water rather than fresh medium in order to keep the solute concentration as it was before desiccation. The cells were re-plated, and their growth assayed via microscopy.

### Freeze/thaw tolerance

To measure tolerance to repeated cycles of freezing and thawing, 70  $\mu$ L of a liquid culture of each isolate was placed into a 0.2 mL PCR tube. The tube was then flash frozen by placing it for 30 sec into an ice bucket containing liquid nitrogen. The tube was then removed and allowed to thaw at room temperature. Cells were then repeatedly re-frozen or plated in 10  $\mu$ L replicates and growth was assayed as described above. The same protocol was followed for freeze/thawing in the dry ice/ethanol slurry.

### Pigment extraction and absorbance measurements

Pigments were extracted in the dark by first centrifuging 1.5 mL of  $1 \cdot 10^6$  cells·mL<sup>-1</sup> liquid cultures at 4000 x g for 3 min. The supernatant was removed, leaving behind ~50  $\mu$ L of liquid and ~25 mg of wet weight. Tubes were then placed in ice, and cells were ground with an electric pellet pestle motor (Kontes/Sigma-Aldrich Inc., St. Louis, MO, USA) in repetitions of 30 sec until cells were broken, as confirmed by light microscopy. An aliquot of 250  $\mu$ L of methanol was added, and the suspension mixed by hand. Cells were centrifuged at 10,000 x g for 10 min. The supernatant containing the extracted pigment was removed, and the absorbance measured from 190 to 1000 nm in 10 nm intervals on a Spectramax 384 Spectrophotometer (Molecular Devices, Sunnyvale, CA, USA).

### Acknowledgments



Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

The laboratory work was performed in partial fulfillment of the following Ph.D. thesis: Cumbers, J.R. Phylogenetic and Phenotypic Studies of the Extremophile *Chroococcidiopsis* (Cyanobacterium). A Foundation for Synthetic Biology in Space. Ph.D. Thesis, Brown University. 2012. <https://doi.org/10.7301/Z0VX0DTV>. We are enormously grateful to the other members of the thesis committee, Gary Wessel, Mark Tatar, Casey Dunn, and Daniel Weinreich for technical advice and support. We thank S. Pete Worden for funding this study while Center Director of NASA Ames Research Center, and for encouraging LJR to explore the potential for synthetic biology as an enabling technology for NASA's missions. This work was in part funded by NASA Cooperative Agreement Award Number, NNX08AZ52A.

## References

- Abramoff, M. D., Magelhaes, P. J., and Ram, S. J. 2004. Image Processing with ImageJ. *Biophotonics International* 11:36–42.
- Andersen, R., Morton, S., and Sexton, J. 1997. Provasoli-Guillard National Center for culture of marine phytoplankton 1997 list of strains. *J. Phycology* 33:1–75.
- Baqué, M., de Vera, J-P., Rettberg, P. and Billi, D. 2013a. The BOSS and BIOMEX space experiments on the EXPOSE-R2 mission: Endurance of the desert cyanobacterium *Chroococcidiopsis* under simulated space vacuum, Martian atmosphere, UVC radiation and temperature extremes. *Acta Astronautica* 91: 180–186.
- Baque, M., Viaggiu, E., Scalzi, G., and Billi, D. 2013b. Endurance of the endolithic desert cyanobacterium *Chroococcidiopsis* under UVC radiation. *Extremophiles* 17:161–169 DOI 10.1007/s00792-012-0505-5
- Baqué, M., Verseux, C., Rabbow, E., de Vera, J-P.P. and Billi, D. 2014. Detection of Macromolecules in Desert Cyanobacteria Mixed with a Lunar Mineral Analogue After Space Simulations. *Orig Life Evol Biosph.* DOI 10.1007/s11084-014-9367-4
- Battista, J. R. 1997. Against all odds: the survival strategies of *Deinococcus radiodurans*. *Ann. Rev. Microbiology*, 51:203–224.
- Bebout, B., and Garcia-Pichel, F. 1995. UV B-Induced Vertical Migrations of Cyanobacteria in a Microbial Mat. *Appl Environ Microbiol* 61:4215.
- Billi, D. 2009a. Loss of topological relationships in a Pleurocapsalean cyanobacterium (*Chroococcidiopsis* sp.) with partially inactivated ftsZ. *Annals of Microbiology* 59:235–238.
- Billi, D. 2009b. Subcellular integrities in *Chroococcidiopsis* sp. CCME029 survivors after prolonged desiccation revealed by molecular probes and genome stability assays. *Extremophiles* 13:49–57.
- Billi, D., 2012. Plasmid stability in dried cells of the desert cyanobacterium *Chroococcidiopsis* and its potential for GFP imaging of survivors on Earth and in space. *Orig. life Evol. Biosph.* 42, 235– 245.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

- Billi, D., Friedmann, E.I., Helm, R.F. and Potts, M. 2001. Gene Transfer to the Desiccation-Tolerant Cyanobacterium *Chroococcidiopsis*. *J. Bacteriol.* 183: 2298–305 DOI: 10.1128/JB.183.7.2298–2305.2001
- Billi, D., Friedmann, E. I., Hofer, K. G., Caiola, M. G., and Ocampo-Friedmann, R. 2000a. Ionizing-Radiation Resistance in the Desiccation-Tolerant Cyanobacterium *Chroococcidiopsis*. *Appl Environ Microbiol* 66:1489–1492.
- Billi, D., Wright, D., Helm, R., Prickett, T., Potts, M., and Crowe, J. 2000b. Engineering desiccation tolerance in *Escherichia coli*. *Appl Environ Microbiol* 66:1680–1684.
- Billi, D., Cockell, C. S., Rabbow, E., Horneck, G. and Onofri, S. 2011. Damage Escape and Repair in Dried *Chroococcidiopsis* spp. from Hot and Cold Deserts Exposed to Simulated Space and Martian Conditions. *Astrobiology* 11:65–73.
- Billi, D., Baqué, M., Verseux, C., Rothschild, L. & de Vera, J-P. 2016. “Desert Cyanobacteria - Potential for Space and Earth applications.” In: *Adaption of Microbial Life to Environmental Extremes* Second Edition, Springer, Eds. Helga Stan-Lotter and Sergiu Fendrihan. Pp. 133-46.
- Billi, D., Staibano, C., Verseux, C., Fagiarone, C., Mosca, C., Baqué, M., Rabbow, E. and Rettberg, P. 2019. Dried Biofilms of Desert Strains of *Chroococcidiopsis* Survived Prolonged Exposure to Space and Mars-like Conditions in Low Earth Orbit. *Astrobiology* 19: 1008-17. DOI: 10.1089/ast.2018.1900
- Cadoret, J. C. 2005. Cyclic Nucleotides, the Photosynthetic Apparatus and Response to a UV-B Stress in the Cyanobacterium *Synechocystis* sp. PCC 6803. *J. Biological Chemistry* 280:33935–33944.
- Caiola, M., Ocampo-Friedmann, R., and Friedmann, E. 1993. Cytology of long-term desiccation in the desert cyanobacterium *Chroococcidiopsis* (Chroococcales). *Phycologia* 32:315–322.
- Caiola, M. G., Billi, D., and Friedmann, E. I. (1996a. Effect of desiccation on envelopes of the cyanobacterium *Chroococcidiopsis* sp. (Chroococcales). *European Journal of Phycology* 31:97.
- Caiola, M. G., Canini, A., and Ocampo-Friedmann, R. 1996b. Iron superoxide dismutase (Fe-SOD) localization in *Chroococcidiopsis* sp. (Chroococcales, Cyanobacteria). *Phycologia* 35:90–94.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

- Capece, M., Clark, E., Saleh, J.K., Halford, D., Henl, N., Hoskins, S. & **Rothschild**, L.J. 2013. Polyextremophiles and the constraints for terrestrial habitability. In: *Polyextremophiles, Microorganisms and macroorganisms living under multiple forms of stress*. J. Seckback, Ed. Pp. 7-59.
- Červený, Jan; Sinetova, Maria A.; Zavřel, Tomáš; Los, Dmitry A. 2015. "Mechanisms of High Temperature Resistance of *Synechocystis* sp. PCC 6803: An Impact of Histidine Kinase 34. *Life* 5, no. 1: 676-699.
- Cockell, C.S., Schuerger, A.C., Billi, D., Friedmann, E.I., and Panitz, C. 2005. Effects of a simulated martian UV flux on the cyanobacterium, *Chroococcidiopsis* sp. 029. *Astrobiology* 5:127–140.
- Cockell, C. S., McKay, C. P., Warren-Rhodes, K., and Horneck, G. 2008. Ultraviolet radiation-induced limitation to epilithic microbial growth in arid deserts - dosimetric experiments in the hyperarid core of the Atacama Desert. *J Photochem Photobiol B, Biol* 90:79–87.
- Cooke, M. S. 2003. Oxidative DNA damage: mechanisms, mutation, and disease. *The FASEB Journal* 17:1195–1214.
- Cox, M. and Battista, J. 2005. *Deinococcus radiodurans* - the consummate survivor. *Nature Reviews Microbiology* 3:882–892.
- Cumbers, J. and Rothschild, L.J. (2014). Salt Tolerance and Polyphyly in the Cyanobacterium *Chroococcidiopsis* (Pleurocapsales). *J. Phycol.* 50:472–482.
- Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., Vasilenko, A., Zhai, M., Leapman, R. D., Lai, B., et al. 2007. Protein Oxidation Implicated as the Primary Determinant of Bacterial Radioresistance. *PLoS Biology*, 5(4), e92. Public Library of Science.
- Daly, M.J., Gaidamakova, E.K., Matrosova, V.Y., Vasilenko, A., Zhai, M., Venkateswaran, A., Hess, M., Omelchenko, M.V., Kostandarithes, H.M., Makarova, K.S., Wackett, L.P., Fredrickson, J.K., and Ghosal D. 2004. Accumulation of Mn(II) in *Deinococcus radiodurans* facilitates  $\gamma$ -radiation resistance. *Science* 306:1025–8.
- de Vera, J.-P., Alawi, M., Backhaus, T., Baqué, M., Billi, D., Böttger, U., Berger, T., Bohmeier, M., Cockell, C., Demets, R., de la Torre Noetzel, R., Edwards, H., Elsaesser, A., Fagiarone, C., Fiedler, A., Foing, B., Foucher, F., Fritz, J., Hanke, F., Herzog, T.,

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

- Horneck, G., Hübers, H-W., Huwe, B., Joshi, J., Kozyrovska, N., Kruchten, M., Lasch, P., Lee, N., Leuko, S., Leya, T., Lorek, A., Martínez-Frías, J., Meessen, J., Moritz, S., Moeller, R., Olsson-Francis, K., Onofri, S., Ott, S., Pacelli, C., Podolich, O., Rabbow, E., Reitz, G., Rettberg, P., Reva, O., Rothschild, L., Sancho, L.G., Schulze-Makuch, D., Selbmann, L., Serrano, P., Szewzyk, U., Verseux, C., Wadsworth, J., Wagner, D., Westall, F., Wolter, D., Zucconi, L. 2019. Limits of Life and the Habitability of Mars: The ESA Space Experiment BIOMEX on the ISS. *Astrobiology* 19:145-57. DOI: 10.1089/ast.2018.1897
- Dillon, J.G., Tatsumi, C.M., Tandingan, P.G. and Castenholz, R.W. 2002. Effect of environmental factors on the synthesis of scytonemin, a UV-screening pigment, in a cyanobacterium (*Chroococcidiopsis* sp.) *Arch Microbiol* 177 : 322–331 DOI 10.1007/s00203-001-0395-x
- Ehling-Schulz, M., Bilger, W., & Scherer, S. 1997. UV-B-induced synthesis of photoprotective pigments and extracellular polysaccharides in the terrestrial cyanobacterium *Nostoc commune*. *J. Bacteriology*, 179:1940-1945.
- Ehling-Schulz, M., and Scherer, S. 1999. UV protection in cyanobacteria. *European J. Phycology* 34:329–338.
- Erokhina, L.G., Shatilovich, A.V., Kaminskaya, O.P. and Gilichinskii, D.A. 2002. Chlorophylls, phycocyanin and carotenoids based on identification of: The Absorption and Fluorescence Spectra of the Cyanobacterial Phycobionts of Cryptoendolithic Lichens in the High-Polar Regions of Antarctica. *Microbiology* 71: 601–7. Translated from Mikrobiologiya, Vol. 71, No. 5, 2002, pp. 697–704
- Fagliarone, C., Mosca, C., Ubaldi, I., Verseux, C., Baqué, M., Wilmotte, A. and Billi, D. 2017. Avoidance of protein oxidation correlates with the desiccation and radiation resistance of hot and cold desert strains of the cyanobacterium *Chroococcidiopsis*. *Extremophiles* 21:981–991 DOI 10.1007/s00792-017-0957-8
- Faraji, S. and Dreuw, A. 2014. Physicochemical Mechanism of Light-Driven DNA Repair by (6-4) Photolyases. *Annu. Rev. Phys. Chem.* 65:275–92 doi: 10.1146/annurev-physchem-040513-103626

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

- Fewer, D., Friedl, T., and Büdel, B. 2002. *Chroococcidiopsis* and Heterocyst-Differentiating Cyanobacteria Are Each Other's Closest Living Relatives. *Molecular Phylogenetics and Evolution* 23:82–90.
- Franklin, L. A., and Forster, R. M. 1997. The changing irradiance environment: consequences for marine macrophyte physiology, productivity and ecology. *European J. Phycology* 32:207–232.
- Friedberg, E. C., Walker, G. C., Siede, W., Wood, R. D., Schultz, R. A., and Ellenberger, T. 2006. *DNA Repair and Mutagenesis*. (Washington D.C., USA: American Society for Microbiology Press)
- Friedmann, E. 1995. A primitive cyanobacterium as pioneer microorganism for terraforming Mars. *Advances in Space Research* 15:243–246.
- Friedmann, I., Lipkin Y, and Ocampo-Paus, R. 1967. Desert algae of the Negev (Israel). *Phycologia* 6:185–200
- Friedmann, E. I., and Ocampo, R. 1976. Endolithic blue-green algae in the dry valleys: primary producers in the Antarctic desert ecosystem. *Science* 193:1247–1249.
- Frohnmeier, H. 2003. Ultraviolet-B Radiation-Mediated Responses in Plants. Balancing Damage and Protection. *Plant Physiology* 133:1420–1428.
- Garcia-Pichel, F. and Castenholz, R. 1993. Occurrence of UV-absorbing, mycosporine-like compounds among cyanobacterial isolates and an estimate of their screening capacity. Identification of the peaks is discussed in the text. *Appl. Env. Microbiol.* 59: 163-169.
- Geng, J., Tang, W., Wan, X., Zhou, Q., Wang, X.-J., Shen, P., Lei, T.-C., and Chen, X.-D. (2008). Photoprotection of bacterial-derived melanin against ultraviolet A–induced cell death and its potential application as an active sunscreen. *J Eur Acad Dermatol Venerol* 22:852–858.
- Goodsell, D. S. 2001. The Molecular Perspective: Ultraviolet Light and Pyrimidine Dimers. *The Oncologist* 6:298–9.
- Graham, J.M. 2004. The Biological Terraforming of Mars: Planetary Ecosynthesis as Ecological Succession on a Global Scale. *Astrobiology* 4: 168-95.
- Hassler, D.M., Zeitlin, C., Wimmer-Schweingruber, R.F., Eh- resmann, B., Rafkin, S., Eigenbrode, J.L., Brinza, D.E., Weigle, G., Bö tcher, S., Bö hm, E., Burmeister, S., Guo, J., Kö hler, J.,

- Martin, C., Reitz, G., Cucinotta, F.A., Kim, M.H., Grinspoon, D., Bullock, M.A., Posner, A., Gomez-Elvira, J., Vasavada, A., Grotzinger, J.P.; MSL Science Team. 2013. Mars' surface radiation environment measured with the Mars Science Laboratory's Curiosity rover. *Science* 343, doi:10.1126/ science.1244797.
- He, Y., and Hader, D. 2002a. UV-B-induced formation of reactive oxygen species and oxidative damage of the cyanobacterium *Anabaena sp.*: protective effects of ascorbic acid and N-acetyl-L-cysteine. *J Photochem Photobiol B, Biol* 66:115–124.
- He, Y.-Y., and Häder, D.-P. 2002b. Reactive oxygen species and UV-B: effect on cyanobacteria. *Photochem. Photobiol. Sci.* 1:729–736.
- Hendrischk, A. K., Braatsch, S., Glaeser, J., and Klug, G. 2007. The phrA gene of *Rhodobacter sphaeroides* encodes a photolyase and is regulated by singlet oxygen and peroxide in a E-dependent manner. *Microbiology* 153:1842–1851.
- Horneck, G., Rettberg, P., Reitz, G., Wehner, J., Eschweiler, U., Strauch, K., Panitz, C., Starke, V., and Baumstark-Khan, C. 2001. Protection of bacterial spores in space, a contribution to the discussion on Panspermia. *Origins of Life and Evolution of the Biosphere* 31:527–547.
- Karsten, U., and Garcia-Pichel, F. 1996. Carotenoids and mycosporine-like amino acid compounds in members of the genus *Microcoleus* (Cyanobacteria): a chemosystematic study. *Systematic Applied Microbiology* 19:285–294.
- Komárek, J., Kaštovský, J., Mareš, J., and Johansen, J.R. 2014. Taxonomic classification of cyanoprokaryotes (cyanobacterial genera) 2014, using a polyphasic approach. *Preslia* 86: 295-335.
- Kumar, A., Sinha, R. P., and Hader, D. P. 1996. Effect of UV-B on enzymes of nitrogen metabolism in the cyanobacterium *Nostoc calcicola*. *J. Plant Physiology* 148:86–91.
- Leuko, S., Neilan, B. A., Burns, B. P., Walter, M. R., and Rothschild, L. J. 2011. Molecular assessment of UVC radiation-induced DNA damage repair in the stromatolitic halophilic archaeon, *Halococcus hamelinensis*. *J. Photochem Photobiol B, Biol* 102:140–145.
- Liu, Y., Cockell, C.S., Wang, G., Hu, C., Chen, L. and de Philippis, R. 2008. From Control of Lunar and Martian Dust—Experimental Insights from Artificial and Natural Cyanobacterial and

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

Algal Crusts in the Desert of Inner Mongolia, China. *Astrobiology* 8: DOI:  
10.1089/ast.2007.0122

Mattimore, V. and Battista, J. R. 1996. Radioresistance of *Deinococcus radiodurans*: functions necessary to survive ionizing radiation are also necessary to survive prolonged desiccation. *J. Bacteriol.* 178:633–37.

Morano K.A. 2014. Anhydrobiosis: drying out with sugar. *Current Biology* 24: R1121-R1123.  
[doi.org/10.1016/j.cub.2014.10.022](https://doi.org/10.1016/j.cub.2014.10.022)

Olsson-Francis, K., and Cockell, C. S. 2010a. Experimental methods for studying microbial survival in extraterrestrial environments. *J Microbiol Methods* 80:1–13.

Olsson-Francis, K. and Cockell, C.S. 2010b. Use of cyanobacteria for in-situ resource use in space applications. *Planetary And Space Science*, 58(10) pp. 1279–1285.

Palmer, R. J., and E. I. Friedmann. 1990. Water relations and photosynthesis in the cryptoendolithic microbial habitat of hot and cold deserts. *Microb. Ecol.* 19:111-118.

Pattanaik, B., Schumann, R., and Karsten, U. (2007). Effects of ultraviolet radiation on cyanobacteria and their protective mechanisms. In *Algae and Cyanobacteria in Extreme Environments*, J. Seckbach, ed. (New York, USA: Springer), pp. 31–45.

Paulino-Lima, I.G., Fujishima, K., Navarrete, J.U., Galante, D., Rodrigues, F., Azua-Bustos, A., Rothschild, L.J. 2017. Extremely high UV-C radiation resistant microorganisms from desert environments with different manganese concentrations. *J Photochem Photobiol B* 163: 327–36.

Portwich, A., and Garcia-Pichel, F. (2007). A Novel Prokaryotic UVB Photoreceptor in the Cyanobacterium *Chlorogloeopsis* PCC 6912. *Photochemistry and Photobiology* 71;493–498.

Puthran, S.S., Sudha, K., Rao, G.M. and Shetty, B.V. 2009. Oxidative stress and low dose ionizing radiation. *Indian J. Physiol. Pharmacol.* 53: 181-4.

Rainey, F.A., Ray, K., Ferreira, M., Gatz, B.Z., Nobre, M.F., Bagaley, D., Rash, B.A., Park, M.J., Earl, A.M., Shank, N.C., Small, A.M.*et al.* 2005. Extensive diversity of ionizing-radiation-resistant bacteria re- covered from Sonoran Desert soil and description of nine new



- species of the genus *Deinococcus* obtained from a single soil sample. *Appl Environ Microbiol* 71:5225–5235
- Rippka, R., Deruelles, J., and Waterbury, J. (1979). Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *J. General Microbiology* 111:1–61.
- Rastogi, R. P., Richa, Kumar, A., Tyagi, M. B., & Sinha, R. P. (2010). Molecular mechanisms of ultraviolet radiation-induced DNA damage and repair. *J. Nucleic Acids*, 2010, 592980. doi:10.4061/2010/592980
- Rothschild, L.J. 1990. Earth analogs for Martian life. Microbes in evaporites, a new model system for life on Mars. *Icarus* **88**: 246-260.
- Rothschild, L. J. 2010. A powerful toolkit for synthetic biology: Over 3.8 billion years of evolution. *BioEssays* 32:304–313.
- Rothschild, L.J. 2016. Synthetic biology as the enabling technology for humans on Mars. *Biochemical Society Transactions*, **44**(4): 1158-64. DOI: 10.1042/BST20160067;
- Rothschild, L. J., and Cockell, C. S. 1999. Radiation: microbial evolution, ecology, and relevance to mars missions. *Mutation Research* 430:281–291.
- Rothschild, L.J. & Mancinelli, R.L. 2001. Life in extreme environments. *Nature (London)* **409**: 1092-1101.
- Shahmohammadi, H. R., Asgarani, E., Terato, H., Saito, T., Ohyama, Y., Gekko, K., Yamamoto, O., and Ide, H. 1998. Protective roles of bacterioruberin and intracellular KCl in the resistance of *Halobacterium salinarum* against DNA-damaging agents. *J. Radiation Research* 39;251–262.
- Sicilia, T., Bub, A., Rechkemmer, G., Kraemer, K., Hoppe, P. P., & Kulling, S. E. 2005. Novel lycopene metabolites are detectable in plasma of preruminant calves after lycopene supplementation *The Journal of Nutrition*, 135:2616–2621.
- Sinha, R. 2000. Mycosporine-like amino acids in the marine red alga *Gracilaria cornea* - effects of UV and heat. *Environmental and Experimental Botany* 43:33–43.
- Sinha, R. P., Ambasht, N.K., Sinha, J.P., Klisch, M., a, Donat-D-P. 2003. UV-B-induced synthesis of mycosporine-like amino acids in three strains of *Nodularia* (cyanobacteria). *Journal of Photochemistry and Photobiology B: Biology*, 71: 51-8.

- Sinha, R. P., Dautz, M., and Häder, D.-P. 2001. A simple and efficient method for the quantitative analysis of thymine dimers in cyanobacteria, phytoplankton and macroalgae. *Acta Protozoologica* 40:187–195.
- Sinha, R. P., and Häder, D.-P. 2002. UV-induced DNA damage and repair: a review. *Photochemical & Photobiological Sciences* 1:225–236.
- Sinha, R. P., Dautz, M., and Häder, D.-P. 2001. A simple and efficient method for the quantitative analysis of thymine dimers in cyanobacteria, phytoplankton and macroalgae. *Acta Protozoologica* 40:187–195.
- Stahl, W. and Sies, H. 2003. Antioxidant activity of carotenoids. *Molecular Aspects of Medicine*, 24: 345-51. doi.org/10.1016/S0098-2997(03)00030-X
- Stanier, R., Kunisawa, R., and Mandel, M. 1971. Purification and properties of unicellular blue-green algae (order Chroococcales). *Bacteriological Reviews* 35:171–205.
- Takaichi, S. 2005. Myxol and 4-Ketomyxol 2'-Fucosides, not Rhamnosides, from *Anabaena* sp. PCC 7120 and *Nostoc punctiforme* PCC 73102, and Proposal for the Biosynthetic Pathway of Carotenoids. *Plant and Cell Physiology*, 46:497–504.
- Verseux, C., Baqué, M., Lehto, K., de Vera, J.-P., Rothschild, L.J. & Billi, D. 2015. Sustainable life support on Mars - the potential roles of cyanobacteria. *Internatl. J. Astrobiology*, 28 pp. doi:10.1017/S147355041500021X
- Verseux C, Baqué M, Cifariello R, Fagiarone C, Raguse M, Moeller R, Billi D 2017. Evaluation of the resistance of *Chroococcidiopsis* spp. to sparsely and densely ionizing irradiation. *Astrobiology* 17:118–125
- Vincent, W. F., and Roy, S. 1993. Solar ultraviolet-B radiation and aquatic primary production: damage, protection, and recovery. *Environmental Reviews* 1:1–12.
- Warren-Rhodes, K. A., Rhodes, K. L., Boyle, L. N., Pointing, S. B., Chen, Y., Liu, S., Zhuo, P., and McKay, C. P. 2007. Cyanobacterial ecology across environmental gradients and spatial scales in China's hot and cold deserts. *FEMS Microbiol Ecol* 61:470–482.
- Warren-Rhodes, K. A., Rhodes, K. L., Pointing, S. B., Ewing, S. A., Lacap, D. C., Gómez-Silva, B., Amundson, R., Friedmann, E. I., and McKay, C. P. (2006). Hypolithic cyanobacteria, dry

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

limit of photosynthesis, and microbial ecology in the hyperarid Atacama Desert. *Microb Ecol* 52:389–398.

Wu, H., Gao, K., Villafane, V. E., Watanabe, T., and Helbling, E. W. (2005). Effects of Solar UV Radiation on Morphology and Photosynthesis of Filamentous Cyanobacterium *Arthrospira platensis*. *Appl Environ Microbiol* 71:5004–5013.

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*


## Supplementary information

None


## Figures

**Fig. 1.** Dose response curves of species exposed to different dosages of UVC (254 nm). Species CCMP 1991 (Hawaii) and CCME 029 (Negev) have similar radiation resistance. Species CCMP 3184 (Samoa) has moderate radiation resistance. Cells were irradiated on agar plates and incubated as described above. Cell density was then measured and normalized to the density of un-irradiated cells. Values are means of two independent trials with six replicates per trial.

**Fig 1.**



**Fig. 2.** DNA repair of thymine dimers occurs faster in species 1991 (Hawaii) than in species 3184 (Samoa). Cells were irradiated with 250 J·m<sup>2</sup> of UV radiation and left to recover under laboratory light conditions. Samples were taken periodically during recovery and DNA was extracted. The number of thymine dimers was determined by immunoblotting. Mean values are from three analyses.



**Fig. 3.** DNA repair takes place faster in the light than in the dark in both 1991 (Hawaii) and 3184 (Samoa). As in Fig. 2, but cells were left to recover in the dark as well as the light. Mean values are from three analyses. A: 1991 (Hawaii); B 3184 (Samoa).





Fig. 4. Hydrogen peroxide tolerance in 1991 (Hawaii) and 3184 (Samoa) shows that 1991 (Hawaii) is more tolerant to  $H_2O_2$  than 3184 (Samoa). Cells were diluted 1:1 in  $H_2O_2$  and plated onto agar plates for growth. Cell density was then measured and normalized relative to the density of un-exposed cells. Mean values are shown from three independent trials with six replicates per trial.





**Fig. 5**





Fig. 5. Resistance of strains to repeated cycles of freezing and thawing. Cells were frozen for 30 seconds, and thawed at room temperature. A. Repeated freezing and thawing of cells in liquid nitrogen followed by plating and growth on agar plates. The results show that species 1991 (Hawaii) and 3184 (Samoa) are more resistant to freeze thawing in liquid N<sub>2</sub> than the other species. Density of cells was assayed and normalized relative to the cell density of control cells. Mean values are shown from two independent trials with six replicates per trial. Species 1489 (Galapagos) and 2116 (Madagascar) are from a single trial with six replicates. B. A comparison of freezing and thawing in liquid N<sub>2</sub> versus a slurry of dry ice and ethanol. The data are the average of five replicates (1991 dry ice) or six replicates (384 dry ice). Liquid N<sub>2</sub> data are as in A.

**Fig. 6.**



Spectral absorbance scans of pigments extracted in methanol normalized to the red peak of chlorophyll *a*. The spectra show several features expected of cyanobacteria, notably the characteristic peaks of chlorophyll *a* at around 432 and 665 nm, carotenoids around 490 nm, and phycocyanin at 620. Absorbance in the UV range is likely to be caused by the presence of scytonemin and mycosporine-like amino acids (MAAs). Species 1991 (Hawaii) has a pigment absorbing around 325 nm that is absent in the other species under the growth conditions of this experiment. Identification of the peaks is discussed in the text.

## Tables

Table 1: *Chroococcidiopsis* isolates used in this study

Species	strain	Growth medium	Location isolated
<i>unassigned</i>	CCMP 1991	BG11 + 4% sea salt	Hawaii
<i>unassigned</i>	CCMP 2116	BG11 + 4% sea salt	Nosy Kumba Island, Madagascar
<i>C. polansiana</i>	CCMP 1489	BG11 + 4% sea salt	Fernandina Island, Galapagos station 8
<i>unassigned</i>	CCMP 3184	BG11 + 4% sea salt	Island of Savaii, Western Samoa, Micronesia
<i>unassigned</i>	PCC 6712	BG11	Marin County, California
<i>C. thermalis</i>	PCC 7203	BG11	near Greifswald, East Germany.
<i>unassigned</i>	CCMEE 029	BG11	Makhtesh Ramon, Negev Desert, Israel
<i>unassigned</i>	CCMEE 171- A789-2	BG11	University Valley, McMurdo Dry Valleys, Antarctica

Abbreviations for culture collections: Provasoli-Guillard National Center for Culture Collection of Marine Phytoplankton (CCMP, Andersen et al., 1997; since 2011 the NCMA, the National Center for Marine Algae and Microbiota), the Pasteur Culture Collection of Cyanobacteria (Rippka et al., 1979), The University of Oregon's Culture Collection of Microorganisms from Extreme Environments (CCMEE) (<http://cultures.uoregon.edu>) and Sammlung von Algenkulturen (SAG) at the University of Göttingen (Göttingen, Germany).

Table 2: Colony forming ability after 9 months of desiccation

Isolate	Desiccation tolerant?
CCMEE 171 (Antarctica)	+
CCMEE 029 (Negev, Israel)	+
PCC 7203 ( <i>C. thermalis</i> )	+
PCC 6712 (Marin)	-
CCMP 1991 (Hawaii)	+

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

CCMP 1489, <i>C. polansiana</i> (Galapagos)	-
CCMP 3184 (Samoa)	+
CCMP 2116 (Madagascar)	+

Cumbers and Rothschild, Environmental boundaries of *Chroococcidiopsis*

## Funding Disclosure

The research was funded as part of a cooperative agreement between Brown University and NASA Award Number NNX08AZ52A to Prof. Marc Tatar, and funding from the NASA Ames Center Director's Discretionary Funds.

## Competing interests

None

## Related Manuscripts

Cumbers, J. & **Rothschild**, L.J., 2014. Salt Tolerance and Polyphyly in the Cyanobacterium *Chroococcidiopsis* (Pleurocapsales). *J. Phycol.* **50**: 472–482.

Ph.D. thesis: Cumbers, J.R. Phylogenetic and Phenotypic Studies of the Extremophile *Chroococcidiopsis* (Cyanobacterium). A Foundation for Synthetic Biology in Space. Ph.D. Thesis, Brown University. 2012. *Molecular Biology, Cell Biology, and Biochemistry Theses and Dissertations*. Brown Digital Repository. Brown University Library. <https://doi.org/10.7301/Z0VX0DTV>