Self-produced hydrogen sulfide improves ethanol fermentation by
***Saccharomyces cerevisiae* and other yeast species**

Emilio Espinoza-Simón [1#], Paola Moreno-Alvarez [1#], Elias Nieto-Zaragoza [1],
Carolina Ricardez-García [2], Emmanuel Ríos-Castro [3], Salvador Uribe-Carvajal [2]
and Francisco Torres-Quiroz [1,*]

[1]   División de Ciencia Básica, Departamento de Bioquímica y Biología
Estructural, Instituto de Fisiología Celular, Universidad Nacional Autónoma de
México (UNAM), Ciudad de México, México

[2]   División de Ciencia Básica, Departamento de Genética Molecular, Instituto
de Fisiología Celular, Universidad Nacional Autónoma de México (UNAM),
Ciudad de México, México

[3]   Unidad de Genómica, Proteómica y Metabolómica, LaNSE, Centro de
Investigación y de Estudios Avanzados del I.P.N., Ciudad de México, México

#   These authors contributed equally to this work

*   Correspondence: ftq@ifc.unam.mx

Abstract

Hydrogen sulfide (H_2S) is a gas produced endogenously in organisms from the
three domains of life. In mammals, it is involved in diverse physiological
processes, including the regulation of blood pressure, and its effects on memory.
In contrast, in unicellular organisms the physiological role of H_2S has not been
studied in detail. In yeast, for example, in the winemaking industry H_2S is an
undesirable byproduct because of its rotten egg smell; however, its biological

relevance during fermentation is not well understood. The effect of H_2S in cells is linked to a posttranslational modification in cysteine residues known as S-persulfidation. We evaluated S-persulfidation in the *Saccharomyces cerevisiae* proteome. We screened S-persulfidated proteins from cells growing in fermentable carbon sources and we identified several glycolytic enzymes as S-persulfidation targets. Pyruvate kinase, catalyzing the last irreversible step of glycolysis, increased its activity in the presence of a H_2S donor. Yeast cells treated with H_2S increased ethanol production; moreover, mutant cells that endogenously accumulated H_2S produced more ethanol and ATP during the exponential growth phase. This mechanism of the regulation of the metabolism seems to be evolutionarily conserved in other yeast species, because H_2S induces ethanol production in the pre-Whole Genome Duplication species *Kluyveromyces marxianus* and *Meyerozyma guilliermondii*. Our results suggest a new role of H_2S in the regulation of the metabolism during fermentation.

Keywords: H_2S; S-persulfidation; fermentation; yeast; metabolism; posttranslational modification; hydrogen sulfide.

Introduction

Hydrogen sulfide (H_2S) is a gasotransmitter produced endogenously in cells. It has been associated with diverse physiological processes such as vasodilation [1], pain [2] and longevity in animals [3], plant growth and development [4], bacterial antibiotic resistance [5], and as a byproduct of alcoholic fermentation in yeast [6]. Surprisingly, the biological function of H_2S in yeast is not fully understood [7]; the majority of reports describe how it is produced or how to prevent its production during fermentation [8–10]. In yeast, H_2S is involved in heavy metal detoxification [11], population synchrony [12], and chronological

aging [3]; however the molecular mechanisms behind these phenomena have not
been fully elucidated.

In yeast, the main metabolic pathway that produces H_2S is the sulfate assimilation
pathway [13], where inorganic sulfate is transformed to H_2S and used in the
synthesis of methionine and cysteine. This pathway is highly active in the
exponential growth phase, as the principal H_2S producer, sulfite reductase
(encoded by *MET5* and *MET10*) [6], is highly active. The synthesis of H_2S takes
place in the first hours of fermentation and decreases at the final stages when
cells reach the stationary phase [14]. The sulfur transferase Tum1p is another
protein involved in H_2S production during fermentation when high concentrations
of cysteine are present in the media [15]. H_2S is metabolized by Met17p a
sulfhydrylase that catalyze the incorporation of sulfide for the biosynthesis of
sulfur-containing amino acids [16].

The molecular effect of hydrogen sulfide depends on a posttranslational
modification named S-persulfidation (originally termed sulfhydration) [17]. S-
persulfidation involves addition of a thiol group to the cysteine residues (-S-SH)
in proteins. This posttranslational modification has been associated with the
activation and inhibition of protein activity [17,18].

In this work, for the first time, we evaluated the S-persulfidation of yeast proteins.
We report that hydrogen sulfide is a regulator of glycolysis that increases ethanol
production in *S. cerevisiae*. This was observed using an exogenous donor of
hydrogen sulfide or mutant strains that accumulate or produce less H_2S. This
mechanism of regulation was conserved in pre-Whole Genome Duplication
(WGD) species, such as the thermotolerant *Kluyveromyces marxianus* from the
KLE clade and the oleaginous yeast *Meyerozyma guilliermondii* from the CUG-
Ser1 clade. This work provides an insight into how H_2S regulates glucose
metabolism through an evolutionarily conserved mechanism, constituting an
important role of H_2S in fermentation.

Materials and Methods

Yeast strains, media, and growth conditions

Saccharomyces cerevisiae strains used in this study were S288C-derived laboratory strains BY4742 (*MATα his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*) referred as *wt*, BY4741 (*MATa his3Δ1 leu2Δ0 met17Δ0 ura3Δ0*). *Kluyveromyces marxianus* and *Meyerozyma guilliermondii* were isolated from mezcal producers in Michoacán, México [19]. Deletion strains derived from BY4742 were constructed by PCR-based gene replacement [20] using synthetic oligonucleotides and the kanMx and natMx disruption modules contained in plasmids pUG6 and pAG25. Gene deletions were confirmed by PCR using A and D oligos. Strains and oligonucleotides are listed in supplementary table 1. Strains were cultured at 30ºC in liquid YPD medium (1% yeast extract, 2% dextrose, 2% peptone) or YPG medium (1% yeast extract, 2% galactose, 2% peptone) until reaching the exponential growth phase (optical density at 660nm [OD660] =0.5-0.6) and cells were collected then for protein extraction.

Reagents

Sodium hydrosulfide (NaHS), Methyl methanethiosulfonate (MMTS), dithiothreitol (DTT), antibiotin antibody, neocuproine, deferoxamine and others chemicals were purchased from Sigma-Aldrich, St Louis MO, rabbit polyclonal anti-GAPDH (GTX100118, Genetex) were purchased from Genetex, Irvine, CA and N-(6-(biotinamido)hexyl)-3′-(2′-pyridyldithio)-propionamide (HPDP-biotin) (sc-207359), mouse monoclonal anti-enolase (sc-21738), goat polyclonal anti-CBS (sc-46830) and rabbit polyclonal anti-TIM (FL-249) were purchased from Santa Cruz Biotech, Dallas Tx.

Modified biotin switch assay

The modified biotin switch assay was performed as described previously [17,21]. Briefly, after yeast cultures reached exponential phase, cells were collected, and intracellular proteins were extracted with chilled glass-beads in HEN buffer (250 mM HEPES-NaOH pH 7.7, 1 mM EDTA) supplemented with 1% triton X-100, 0.1 mM neocuproine, 0.1 mM deferoxamine and 1X protease cocktail inhibitor (Roche, Switzerland). Cell lysates were centrifuged at 16900 x *g* for 1 hr at 4ºC, total extracts (1-2 mg) were blocked in HEN buffer with 2.5% SDS and 20 mM MMTS at 50ºC for 20 min. The MMTS was removed by acetone precipitation and the protein pellet was resuspended in HEN buffer with 1% SDS. Protein labeling was performed with 0.8 mM HPDP-biotin for 3 h at room temperature in the dark. The biotinylated proteins were separated by SDS-polyacrylamide gel electrophoresis (PAGE) and subjected to immunoblot analysis.

Purification of biotinylated proteins

After biotin switch assay, labeled extracts were subjected to streptavidin-based affinity precipitation with magnetic beads. Labeled extracts were incubated with 3X volumes of neutralization buffer (20 mM HEPES-NaOH pH 7.7, 100 mM NaCl, 1 mM EDTA, 0.5% triton) and 25 µl of streptavidin magnetic beads (Pierce) with agitation, overnight at 4ºC. Magnetic beads were collected and washed with wash buffer as indicated by manufacturer's instructions, biotinylated proteins were eluted with IP-MS elution buffer and analyzed using LS-MS or SDS-PAGE.

Immunoblot analysis

Protein extracts were separated by SDS-PAGE and transferred to polyvinylidene difluoride (PVDF) membranes (Millipore-Merck, Germany). Membranes were

blocked with 5% non-fat milk and incubated with a specific anti-biotin antibody
overnight at 4ºC. Proteins were detected with chemiluminescence using horse-
radish peroxidase conjugated secondary antibodies (Jackson ImmunoResearch,
West Grove, PA). Before to immunoblot, membranes were stained using
Ponceau red (Millipore-Merck, Germany) as protein loading control.

Quantification of intracellular ATP concentration

NaHS was added to cells cultures at specific timepoints, then cells were
centrifuged, and intracellular ATP was measured using the ATP Bioluminescent
Assay Kit HS II (Roche, Switzerland). Cell samples were prepared by diluting
treated cells to a final concentration of 3.7×10^9 cells·mL^{-1} in 500 µl with a buffer
containing 100 mM Tris-HCl pH 7.8 and 4 mM EDTA. After 2 min incubation,
samples were immersed in boiling water for 2 minutes, and the resulting cell
extracts were incubated for 5 minutes at 4ºC, cell debris were removed by
centrifugation at 16900 x *g* for 5 min and supernatants were used to measure the
amount of intracellular ATP using an ATP calibration curve prepared each time,
as indicated by the manufacturer. Bioluminescence was detected in a POLARstar
Omega luminometer (BGM LABTECH, Offenburg, Germany). Three independent
experiments with three replicas were performed, and values are represented as
mean ± standard error.

Detection of H$_2$S production

H$_2$S production by yeast strains colonies was detected through the generation of
a visible black precipitate indicating that the hydrogen sulfide gas has reacted
with lead nitrate [22]. Yeast strains were diluted, and cell density normalized to
3×10^7 cells·mL^{-1}. Cells were spotted in solid media (3.2% dextrose, 0.4% yeast
extract, 0.24% peptone, 0.016% ammonium sulfate, 0.08% lead nitrate, 1.6%

agar) and plates were kept at 30ºC for 5-7 days. Also, H_2S production was

measured as reported previously [23] with some modifications. BY4742 *wt* strain

were precultured at 30ºC with constant shaking for 2 days in fresh YPD media.

The assay was performed on a 96 well plate (COSTAR). Each well had 185 µL

of YPD media, 5 µL of methylene blue (1 mg·mL^{-1}) diluted in citrate buffer (100

mM, pH 4.5) and 10 µL of cells, for a final OD_{600} of 0.2. Growth was measured in

an Infinite 200 (TECAN, Life Sciences) at 600 nm and 663 nm during 15 hours

with intervals of 15 minutes between readings. During measures cells were

incubated at 30ºC with occasionally shaking. Three experimental replicates were

made, with six different biological replicates in each experiment. Data for the

hydrogen sulfide production were analyzed with the following formula:

$$\frac{((OD_{600nm\ t0} - OD_{663nm\ t0}) - (OD_{600nm\ tx} - OD_{663nm\ tx})}{OD_{600nm\ tx}\ from\ no\ reaction\ mix}$$

Fermentation assays

BY4742 *wt* and derived mutants were precultured in liquid YPD medium for 24 h

at 30ºC, under agitation at 90 rpm in an Excella E24 incubator Shaker (New

Brunswick Scientific, USA), then were inoculated in a 2L flask containing 500 mL

of fresh YPD with an initial OD_{600}=0.2 and incubated in the same conditions.

When cells reached DO_{660}=0.5, a pulse of NaHS was added. After 7h of NaHS

addition, aliquots of 2 ml were obtained, DO_{660} was measured, and cells

centrifuged at 16900 x *g* for 1 min. Supernatants were stored at -20ºC for

subsequent ethanol quantification. For mutant and *wt* strains, aliquots were taken

every hour after cells were inoculated, supernatants were stored at -20ºC.

Ethanol production was evaluated through enzymatic assay coupled to NAD^+

reduction. Briefly, supernatants were incubated in buffer (114 mM K_2HPO_4 pH

7.6), 1.8 mM NAD^+ and 39µg·mL^{-1} alcohol dehydrogenase (ADH) for 30 min at

30ºC with vigorous agitation [24]. Produced NADH was monitored by the increase

in absorbance at 340 nm. The results are reported as mM ethanol per 1x10^7 cells.

Three independent experiments with three replicas were performed, and values

are represented as mean $\pm$ standard error. *K. marxianus* fermentation assay was

performed as in *S. cerevisiae* strains, when cells reached $DO_{660}=0.5$, a pulse of

NaHS 0.1mM was added. After 7h of NaHS addition, aliquots of 2 ml were

obtained, and ethanol was qunatified. For *Meyerozyma guilliermondii* when cells

reached $DO_{660}=0.5$, a pulse of NaHS 0.1mM was added, 24h later another pulse

of same concentration was added and 7h later ethanol was quantified.

Enzymes activity assays

Glyceraldehyde 3 phosphate dehydrogenase (GAPDH) [17], pyruvate kinase

(PK) [25] and alcohol dehydrogenase (ADH) [26] activities were measured by

specific reaction assays and monitored spectrophotometrically at 340 nm,

recording the rate of NAD to NADH reduction. Cells cultures were exposed to

NaHS at different times, protein extracts were quantified, and 10 µg of protein

were incubated in assay buffer as follows: for GAPDH (20 mM Tris-HCl pH 7.8,

100 mM NaCl, 0.1 mg·mL^{-1} Bovine serum albumin, 2 mM NAD$^+$, 10 mM sodium

pyrophosphate, 20 mM sodium arsenate, 500 mM DTT buffer, phosphate

buffered saline (PBS) 1X, 27.3 mM glyceraldehyde 3-phosphate [G3P]). For PK

(50 mM Imidazole·HCl, 120 mM KCl, 62 mM $MgSO_4$ pH 7.6, 45 mM ADP, 6.6

mM NADH, 45 mM phosphoenolpyruvate [PEP], 1.3 KU·mL^{-1} lactate

dehydrogenase). For ADH (114 mM K_2HPO_4 pH 7.6), 1.8 mM NAD+, and 16.4

mM ethanol). Three independent experiments with three replicas were

performed, and values are represented as mean $\pm$ standard error.

Oxygen consumption rate assay

BY4742 *wt* and derived mutant cells were precultured in liquid YPD medium for

48 hrs at 30°C, then were cultured in YPD medium with an initial $DO_{600}=0.2$ under

agitation in an Excella E24 incubator (New Brunswick Scientific, USA) for 7h at
30°C. Basal oxygen consumption was measured in resting cells with a Clark
electrode (Oximeter model 782, Warner/Strathkelvin Instruments, North
Lanarkshire, Scotland) in a water-jacketed chamber. Temperature was kept at 30
°C using a water bath (PolyScience 7 L, IL). Oxygen consumption reaction
mixture was MES 10 mM pH 6 and 500 mg (wet weight) of cells were added at
chamber [24]. To evaluate role of NaHS addition, when BY4742 cells reached
OD_{600}=0.5, a pulse of 0.1 mM NaHS was added. Seven hours later, basal oxygen
consumption was measured as abovementioned. Results are reported as
natgO/wet weight g/ mn and values are represented as mean ± standard error.
et weight) of cells were added at chamber.

Sample preparation and LC-MALDI-MS/MS

Biotinylated proteins were digested with 250 ng of trypsin mass spectrometry
grade (Sigma-Aldrich, St. Louis, MO) in 50 mM of ammonium bicarbonate (ABC).
Resulting tryptic peptides were desalted using ZipTip C18 (Millipore) and
concentrated to an approximated volume of 10 µL. Afterward, 9 µL were loaded
into ChromXP Trap Column C18-CL precolumn (Eksigent, Redwood City CA);
350 µm X 0.5 mm, 120 A° pore size, 3 µm particle size and desalted with 0.1%
trifluoroacetic acid (TFA) in H2O at a flow of 5 µL min^{-1} for 10 min. Then, peptides
were loaded and separated on a 3C18-CL-120 column (Eksigent, Redwood City
CA); 75 µm X 150 mm, 120 A° pore size, 3 µm particle size, in a HPLC Ekspert
nanoLC 425 (Eksigent, Redwood City CA) using as a mobile phase A, 0.1% TFA
in H_2O and mobile phase B 0.1% TFA in acetonitrile (ACN) under the following
lineal gradient: 0-3 min 10% B, 60 min 60% B, 61-64 min 90 % B, 65 to 90 min
10% B at a flow of 250 nL min^{-1}. Eluted fractions were automatically mixed with a
solution of 2 mg·mL^{-1} of alfa-cyano-4-hydroxycinnamic acid (CHCA) in 0.1% TFA
and 50% ACN as a matrix, spotted in an Opti-TOF plate of 384 spots using a
MALDI Ekspot (Eksigent, Redwood City CA) with a spotting velocity of 20 s per
spot at a matrix flow of 1.6 µL min^{-1}. The generated spots were analyzed by a

MALDI-TOF/TOF 4800 Plus mass spectrometer (ABSciex, Framingham MA).

Each MS Spectrum was acquired by an accumulation of 1000 shots in a mass

range of 850-4000 Th with a laser intensity of 3800. The 100 more intense ions

with a minimum signal-noise (S/N) of 20 were programmed to fragmenting. The

MS/MS spectra were obtained by fragmentation of selected precursor ions using

Collision-Induced Dissociation (CID) and acquired by 3000 shots with a laser

intensity of 4300. Generated MS/MS spectrums were compared using Protein

Pilot software v. 2.0.1 (ABSciex, Framingham MA) against *Saccharomyces*

cerevisiae, strain ATCC 204508/S288c database (downloaded from Uniprot,

6049 protein sequences) using Paragon algorithm. Search parameters were: Not

constant modifications in cysteines, trypsin as a cutter enzyme, all the biological

modifications and amino acids substitution set by the algorithm (including

carbamidomethylated cysteine as a variable modification); as well as

phosphorylation emphasis and Gel-based ID as special factors. The detection

threshold was considered in 1.3 to acquire 95% of confidence; additionally, the

identified proteins showed a local FDR of 5% or less. Since a peptide derived

from a given fragmentation spectra may be shared among redundant proteins

during database search, it is necessary group all competing proteins and report

only the protein with more spectrometric evidence; for this reason, identified

proteins were grouped by ProGroup algorithm contained in the software Protein

Pilot to minimize redundancy.

Results

S-Persulfidation of yeast proteins growing on a fermentable carbon source

Protein S-persulfidation was detected using the modified biotin switch method

[17]. In order to validate the method in yeast, we performed the assay in either a

poor producer (*met5Δmet10Δ*) or an accumulator (*met17Δ*) strain of H_2S and

compared them to the *wt* (BY4742) (Figure 1A). Cells were grown using glucose

as the carbon source, and at the exponential phase, when H_2S is produced [6],

the protein was extracted. S-Persulfidated proteins were accumulated in *met17Δ*

in comparison to the *met5Δmet10Δ* strain and wt as expected (Figure 1B).

In yeast, H_2S is produced during fermentation, however, the S-persulfidation

target proteins are not known. We used mass spectrometry to analyze the S-

persulfidated proteins in cells growing at the exponential phase in two different

fermentable carbon sources: glucose and galactose. Glucose is the preferred

fermentable carbon source of yeast, while galactose needs to be isomerized to

enter the glycolytic pathway. We found 42 S-persulfidated proteins; 21 were

specific to glucose-grown cells, 4 were specific to galactose-grown cells, and 17

proteins were found in both conditions (Supplementary Table 2). Among the

generally expressed 17 proteins, 15 were reported before as proteins with a

redox-regulated cysteine [27], which is a feature of cysteines susceptible to

posttrans-lational modifications [28]. Cytoplasmic translation (seven proteins)

and glycolysis (seven proteins) where the most represented biological processes

in the cells growing in either condition. Interestingly, pyruvate decarboxylase 1

(Pdc1), a key enzyme in alcoholic fermentation, and Adh1, the major enzyme

responsible for ethanol synthesis, were also found, suggesting a possible role of

S-persulfidation in fermentation. The identities of some glycolytic enzymes,

glyceraldehyde 3-phosphate dehydrogenase (GAPDH), eno-lase, and

triosephosphate isomerase (Tdh3, Eno2, and Tpi1, respectively) were confirmed

using specific antibodies (Figure 2). We also tested cystathionine beta-synthase

(Cys4), which was described before as a possible S-persulfidated protein [17,

29], although it must be considered that in our mass spectrometry analysis it did

not pass the threshold (unused score of 1.04, coverage of 43.98%), we did find

that Cys4 was a target of S-persulfidation.

H_2S production during yeast growth

In order to evaluate H_2S production during yeast growth, we determined H_2S in

the wt. The H_2S reached a maximal amount during the log phase and dropped its

production (Figure 3). Suggesting that the H_2S concentration is not constant and
drops when the culture stops growing.

H_2S increases glycolytic enzymes activities

Among the first effects of S-persulfidation described was the increase in GAPDH
activity [17]. Considering that Tdh3 (GAPDH) was one of the glycolytic enzymes
targeted for S-persulfidation, we decided to test the effect of NaHS (a donor of
H_2S) on GAPDH activity. Cells were stimulated with 0.1 mM or 0.25 mM NaHS
for two and seven hours. Then, the protein was extracted, and the GAPDH activity
was measured. We found that after two hours of NaHS stimulation, both at 0.1
mM and at 0.25 mM, increased GAPDH activity 1.4 times (Figure 4A), and the
effect was lost at seven hours (Figure 4B). Another protein identified by mass
spectrometry was pyruvate kinase (PK) that catalyzes the last irreversible step of
glycolysis. We measured the activity of the pyruvate kinase at two hours of
treatment with 0.1 mM or 0.25 mM NaHS finding that NaHS increased PK activity
2.39 times (Figure 4C, Supplementary Table 3), and lost its effect at seven hours
(Figure 4D). Finally, we subjected alcohol dehydrogenase (ADH) to the same
treatment, and we did not find any significant difference between the treated and
untreated cells, i.e., at these concentrations NaHS did not affect ADH activity
(Supplemental Figure 1).

H_2S stimulates fermentation

The glycolytic enzymes GAPDH and PK increased their activity in response to
H2S. In addition, mass spectrometry data indicated that these and other enzymes
from gly-colysis were S-persulfidated. Thus, we decided to test whether H_2S
influenced the syn-thesis of ethanol. Exponential-phase-grown cells were treated
with 0.1 mM NaHS and after seven hours, the supernatant was collected. It was

observed that the treated cells had increased ethanol production as compared to
the control (Figure 5A).

In order to test the effect of endogenous H_2S in fermentation, we decided to
compare ethanol production in the two isogenic lab strains BY4741 and BY4742.
The only difference between these two strains is that BY4741 has a deletion of
MET17 and strain BY4742 has a deletion of LYS2. As mentioned before, the
met17Δ strain endogenously accumulates H_2S, because *MET17* codifies for the
enzyme using H_2S and O-acetyl homoserine to synthetize homocysteine. An
overnight preculture of each strain was diluted to an OD600=0.2, the
supernatants were collected after 24 h, and the ethanol was quantified. The strain
ac-cumulating H_2S endogenously, BY4741 *met17Δ*, produced more ethanol than
the strain BY4742 *MET17* (Figure 5B).

Ethanol is the main product of fermentation. Additionally, during glycolytic
fermentation two ATP molecules are synthesized. Thus, we decided to quantify
the ATP after treatment with NaHS. Exponential-phase cells were treated with
the same quantities of NaHS used before, and the ATP was quantified two and
four hours after treatment. After two hours, the treated cells produced more ATP
than the untreated cells; this effect was lost four hours after treatment (Figure
5C). These results suggest that H_2S stimulates glycolysis, and that fermentation
is enhanced to produce both ATP and ethanol.

Based on these results we decided to compare whether the endogenous H_2S had
an influence on the onset of ethanol synthesis. We measured the ethanol
production of the poor H_2S producer strain *met5Δmet10Δ*, the H_2S accumulator
met17Δ strain, and the wt. The cells from 48 hours preculture were resuspended
in fresh media and aliquots from the supernatant were collected every hour. The
met17Δ strain initiated ethanol production at five hours, the wt strain initiated
production at six hours, and the *met5Δmet10Δ* initiated production after seven
hours (Figure 6A). This result showed that the cells with high endogenously
accumulated H_2S began ethanol production before the cells with a lower H_2S
concentration.

Finally, in each of these strains we quantified the ATP at the exponential or
stationary phase. We found that the endogenously H_2S accumulator strain
produced the most ATP during the exponential phase, while there were no
differences in the ATP concentration at the stationary phase between the wt and
mutant strains (Figure 6B). These results support our proposal that H_2S
stimulates ethanol and ATP production.

Endogenous H_2S accumulation promotes basal oxygen consumption

ATP could be synthesized as product of the glycolysis and the oxidative phos-
phorylation. In order to elucidate if the ATP produced by H_2S stimulation was from
oxidative phosphorylation, we measured oxygen consumption from wt strain and
mu-tants. A 48 hours preculture of each strain was diluted to an OD600= 0.2, and
oxygen consumption was measured. Then, after seven hours (when cells were
at exponential phase) oxygen was measured again (Figure 7). We found, in the
diluted cells, that the *met5Δmet10Δ*, and the *met17Δ* consumed more oxygen
than the *wt* strain. On the other hand, after seven hours of growing, we found that
the *met17Δ* strain maintained the el-evated rate of oxygen consumption. This
result suggest that endogenously accumulated H_2S promotes oxygen
consumption.

H_2S stimulates ethanol production in Meyerozyma guilliermondii and
Kluyveromyces marxianus

Ethanol synthesis is more robust in Crabtree positive yeast species; this
phenomenon is associated with the WGD [30]. Evidence suggests that the WGD
event arose from an interspecies hybridization between a strain from the KLE
clade (genera Kluyveromyces, Lachancea and Eremothecium) and a strain from
the ZT clade (Zygosaccharomyces and Torulaspora) [31]. The CUG-Ser1 clade

first appeared approximately 117 million years before the WGD event; the CUG-
Ser1 clade is characterized by a change in codon usage [32]. Considering this,
we decided to test the effect of H_2S during ethanol synthesis on the KLE clade
strain, *K. marxianus* and in *M. guilliermondii* from the CUG-Ser1 clade. *K.*
marxianus exponential-phase cells were stimulated with NaHS. Treatment with
the H_2S donor in *K. marxianus* increased ethanol synthesis as in *S. cerevisiae*
(Figure 8a). In *M. guilliermondii*, it was noted that ethanol synthesis took longer
than 24 h when glucose was the carbon source [33]; for this reason, the
exponential-phase cells were treated with NaHS and treated again 24 h later. As
observed in *K. marxianus*, ethanol synthesis increased after the H_2S donor
treatment on *M. guilliermondii* (Figure 8b) confirming that there is an effect of H_2S
on the fermentation activity of these two species of yeast.

Discussion

Hydrogen sulfide is produced endogenously in yeast, and it is considered a fer-
mentation byproduct; however its biological role is unknown. The biological
effects of H_2S are linked to a cysteine posttranslational modification termed S-
persulfidation [17]. In this work, we analyzed S-persulfidated proteins on the yeast
proteome. We identified several glycolytic enzymes as S-persulfidation targets,
as reported previously in tissues such as the brain, heart and liver [29], in
hepatocytes [17], a pancreatic beta cell line [34], HEK293 cells [29], plant [35,36]
and bacteria [37]. Interestingly, Fu and collaborators reported six S-persulfidated
glycolytic enzymes (ALDOA, GAPDH, PGK1, ENO1, PKM and LDHA) when they
evaluated S-persulfidated proteins in cells overexpressing the H2S-producer
enzyme cystathionine gamma-lyase (CSE) [29]. In a previous report, in
pancreatic beta cells, metabolites were measured, and H_2S was associated with
an increased glycolytic metabolic flux of cells under chronic stress. All these
reports agreed that several glycolytic enzymes were S-persulfidated, even when
the activity of GAPDH was the only one measured [17]. Cysteine's
posttranslational modifications of glycolytic enzymes regulate the subcellular

localization and oligomerization, which can impact its activity [38–40]. In a cell culture, H_2S production is not constant, it starts to decline when the cells are at the middle of the logarithmic phase, suggesting that H_2S synthesis could be regulated by the metabolic conditions. In *S. cerevisiae*, we found that GAPDH increased its activity with the H_2S donor NaHS two hours after the treatment, and the effect was lost at seven hours. It is important to note that H_2S is released from NaHS just a few seconds after the sodium salt is dissolved [41]; hence, the effect of NaHS two hours after treatment may be attributed to a chemical modification of the enzymes, such as S-persulfidation. The thioredoxin system eliminates this posttranslational modification [42,43], which is consistent with the idea that cellular mechanisms maintain protein S-persulfidation homeostasis [44]. Seven hours after stimulation, there was no effect on GAPDH activity probably due to the loss of S-persulfidation by the protein. Furthermore, the cells were no longer at an exponential phase seven hours after the OD600 reached 0.5, and the yeast metabolism changed to aerobic at the diauxic shift. The enzymes catalyzing the irreversible steps regulate the glycolytic pathway [45]; and we found that the enzyme involved in the last irreversible step, pyruvate kinase, increased its activity when stimulated with NaHS. The increase in pyruvate kinase activity may have at least two important consequences: feeding the Krebs cycle and/or stimulating the synthesis of ethanol. Considering that yeast synthe-tizes H_2S during fermentation, we decided to test whether NaHS increased ethanol production. We found that cells treated with the H_2S donor produced more ethanol. In order to confirm our observations, we measured the ethanol production in the isogenic strains BY4741 and BY4742. These strains have almost the same selection markers, and they differ only in one of them, BY4741 accumulates H_2S because it is *met17∆*; BY4742 is *lys2∆*, and thus, it does not accumulate H_2S. We observed that BY4741 produced more ethanol than BY4742, suggesting that endogenous H_2S levels increase ethanol synthesis. Previously, it was reported that BY4742 fermenting activity was slower than in BY4741; it would be interesting to analyze the role of H_2S in this system [46]. Considering these results, we proposed that if H_2S stimulates fermentation, then mutants accumulating H_2S would begin ethanol production before strains producing less H_2S. We tested this hy-pothesis by comparing ethanol production between a lower producer of H_2S, the strain *met5∆met10∆*, the accumulator

strain *met17Δ*, and the *wt*. We found that the *met17Δ* strain started to produce

ethanol before the wt strain; in turn, the *met5Δmet10Δ* strain production of ethanol

was delayed even longer. The results confirmed that endogenous concentra-

tions of H_2S affects ethanol synthesis. Finally, we measured ATP production in

all strains, and we found that at the logarithmic phase the *met17Δ* strain produced

more ATP than the others. This result supports the idea that in addition to

increasing an early synthesis of ethanol, H_2S and also enhances ATP production

at the exponential phase of growth.

The fermentation and the oxidative phosphorylation could yield ATP. We found

that at exponential phase the *met17Δ* strain produced more ATP than the wt and

met5Δmet10Δ strains. In order to test if the ATP was produced from the oxidative

phosphorylation, we measured basal oxygen consumption on these strains. We

found that the *met17Δ* strain has an elevated rate of oxygen consumption, and

this is sustained when cells were at ex-ponential phase of growth; suggesting that

endogenously accumulated H_2S induces oxygen consumption. This would be

contradictory to a report where described that ex-ogenous H_2S inhibit respiration

[47], however, at physiological concentrations H_2S could induce the S-

persulfidation of the ATP synthase from mammals and increases its activity [48].

The S-persulfidation takes place at cysteines 244 and 294 of human ATP

synthase. The yeast ATP synthase (Atp1) conserved the cysteine 244

(Supplementary figure 2) in lineal sequence and has similar orientation on protein

structure; therefore, the S-persulfidation could also be carried out in Atp1. Our

results suggest that endogenous H_2S has an effect on glycolysis and oxygen

consumption. The effect of endogenous H_2S on metabolism may explain the

advantage of the *met17Δ* strain growing on fermentable carbon sources (glucose

and galactose), over the *wt* and *met5Δmet10Δ* strains (Supple-mentary figure 3).

It is accepted that the origin of *S. cerevisiae* comes from a WGD event, probably

by the interspecies hybridization between a strain from the KLE clade and a strain

from the ZT clade [31]. WGD species have a more pronounced Crabtree effect

than non-WGD species [30], so we decided to test whether H2S influenced yeast

from the parental KLE clade that originated S. cerevisiae and a Crabtree-negative

species from the CUG-Ser1 clade. The origin of this clade is estimated to be

between 178 and 248 million years ago (mya), and this event occurred before

WGD, estimated to be between 82 and 105 mya [32]. We found that both species

increase ethanol production after the NaHS treatment suggesting that i) H2S is a

positive regulator of fermentation and ii) this effect is evolutionarily conserved.

Overall, H2S is considered as a fermentation byproduct on yeast even when its

biological effect is unknown. Here, we proposed a very different picture that will

change our vision of how H2S regulates cell metabolism.

490   5. Conclusions

In conclusion, our data demonstrated that H_2S is a regulator of energetic

metabolism. These results fill a major gap in the understanding of H_2S and its

control of ethanol production, which is evolutionarily conserved among yeast

species. Finally, our work provides the foundation for a mechanistic

understanding of the effects of H_2S.

Supplementary Materials: Figure S1: Alcohol dehydrogenase activity; Figure S2:

ATP synthase alignment; Figure S3: wt, met5Δmet10Δ, and met17Δ strains

growth curves; Table S1: Strains and oligonucleotides primers; Table S2: Mass

spectrometry results; Table S3: Protein activity data

501   Funding

This work was supported by the Programa de Apoyo a Proyectos de

Investigación e Innovación Tecnológica from the Dirección General de Asuntos

del Personal Academico of the Universidad Nacional Autonoma de Mexico (FTQ:

UNAM-DGAPA-PAPIIT IA200315, IA202217 and IN209219; SUC: IN208821)
and from the Consejo Nacional de Ciencia y Tecnología, Convocatoria de Ciencia
Básica to FTQ (CONACyT-CB-238681).

Acknowledgments

Authors acknowledge Dr. Antonio Peña and Dr. Norma Silvia Sanchez for the
Kluyveromyces marxianus and Meyerozyma guilliermondii strains provided, Dr.
Natalia Chi-quete-Félix, Dr. Gabriel del Rio and Dr. Teresa Lara for technical
assistance.

References

1.   Yang, G.; Wu, L.; Jiang, B.; Yang, W.; Qi, J.; Cao, K.; Meng, Q.; Mustafa,
A.K.; Mu, W.; Zhang, S.; et al. H2S as a Physiologic Vasorelaxant: Hypertension
in Mice with Deletion of Cystathionine Gamma-Lyase. Science 2008, 322, 587–
590, doi:10.1126/science.1162667.

2.   Kawabata, A.; Ishiki, T.; Nagasawa, K.; Yoshida, S.; Maeda, Y.;
Takahashi, T.; Sekiguchi, F.; Wada, T.; Ichida, S.; Nishikawa, H. Hydrogen
Sulfide as a Novel Nociceptive Messenger. Pain 2007, 132, 74–81,
doi:10.1016/j.pain.2007.01.026.

3.   Hine, C.; Harputlugil, E.; Zhang, Y.; Ruckenstuhl, C.; Lee, B.C.; Brace, L.;
Longchamp, A.; Treviño-Villarreal, J.H.; Mejia, P.; Ozaki, C.K.; et al. Endogenous
Hydrogen Sulfide Production Is Essential for Dietary Restriction Benefits. Cell
2015, 160, 132–144, doi:10.1016/j.cell.2014.11.048.

4.   Dooley, F.D.; Nair, S.P.; Ward, P.D. Increased Growth and Germination
Success in Plants Following Hydrogen Sulfide Administration. PLoS One 2013,
8, e62048, doi:10.1371/journal.pone.0062048.

5.   Shatalin, K.; Shatalina, E.; Mironov, A.; Nudler, E. H2S: A Universal
Defense against Antibiotics in Bacteria. Science 2011, 334, 986–990,
doi:10.1126/science.1209855.

6.   Jiranek, V.; Langridge, P.; Henschke, P.A. Regulation of Hydrogen Sulfide
Liberation in Wine-Producing Saccharomyces Cerevisiae Strains by Assimilable
Nitrogen. Appl Environ Microbiol 1995, 61, 461–467, doi:10.1128/aem.61.2.461-
467.1995.

7.   Huang, C.-W.; Walker, M.E.; Fedrizzi, B.; Gardner, R.C.; Jiranek, V.
Hydrogen Sulfide and Its Roles in Saccharomyces Cerevisiae in a Winemaking
Context. FEMS Yeast Res 2017, 17, doi:10.1093/femsyr/fox058.

8.   Huang, C.; Roncoroni, M.; Gardner, R.C. MET2 Affects Production of
Hydrogen Sulfide during Wine Fermentation. Appl Microbiol Biotechnol 2014, 98,
7125–7135, doi:10.1007/s00253-014-5789-1.

9.   Boudreau, T.F.; Peck, G.M.; O'Keefe, S.F.; Stewart, A.C. The Interactive
Effect of Fungicide Residues and Yeast As-similable Nitrogen on Fermentation
Kinetics and Hydrogen Sulfide Production during Cider Fermentation. J Sci Food
Agric 2017, 97, 693–704, doi:10.1002/jsfa.8096.

10.   Wang, C.; Liu, M.; Li, Y.; Zhang, Y.; Yao, M.; Qin, Y.; Liu, Y. Hydrogen
Sulfide Synthesis in Native Saccharomyces Cerevisiae Strains during Alcoholic
Fermentations.   Food   Microbiol   2018,   70,   206–213,
doi:10.1016/j.fm.2017.10.006.

11.   Sun, G.L.; Reynolds, Erin.E.; Belcher, A.M. Using Yeast to Sustainably
Remediate and Extract Heavy Metals from Waste Waters. Nat Sustain 2020, 3,
303–311, doi:10.1038/s41893-020-0478-9.

12.   Sohn, H.Y.; Murray, D.B.; Kuriyama, H. Ultradian Oscillation of
Saccharomyces Cerevisiae during Aerobic Continuous Culture: Hydrogen
Sulphide Mediates Population Synchrony. Yeast 2000, 16, 1185–1190,
doi:10.1002/1097-0061(20000930)16:13<1185::AID-YEA619>3.0.CO;2-W.

13.   Mendoza-Cózatl, D.; Loza-Tavera, H.; Hernández-Navarro, A.; Moreno-
Sánchez, R. Sulfur Assimilation and Glutathione Metabolism under Cadmium
Stress in Yeast, Protists and Plants. FEMS Microbiol Rev 2005, 29, 653–671,
doi:10.1016/j.femsre.2004.09.004.

14.   Oka, K.; Hayashi, T.; Matsumoto, N.; Yanase, H. Decrease in Hydrogen
Sulfide Content during the Final Stage of Beer Fermentation Due to Involvement
of Yeast and Not Carbon Dioxide Gas Purging. J Biosci Bioeng 2008, 106, 253–
257, doi:10.1263/jbb.106.253.

15.   Huang, C.-W.; Walker, M.E.; Fedrizzi, B.; Roncoroni, M.; Gardner, R.C.;
Jiranek, V. The Yeast TUM1 Affects Pro-duction of Hydrogen Sulfide from
Cysteine Treatment during Fermentation. FEMS Yeast Res 2016, 16, fow100,
doi:10.1093/femsyr/fow100.

16.   D'Andrea, R.; Surdin-Kerjan, Y.; Pure, G.; Cherest, H. Molecular Genetics
of Met 17 and Met 25 Mutants of Saccharomyces Cerevisiae: Intragenic
Complementation between Mutations of a Single Structural Gene. Mol Gen
Genet 1987, 207, 165–170, doi:10.1007/BF00331505.

17.   Mustafa, A.K.; Gadalla, M.M.; Sen, N.; Kim, S.; Mu, W.; Gazi, S.K.; Barrow,
R.K.; Yang, G.; Wang, R.; Snyder, S.H. H2S Signals through Protein S-
Sulfhydration. Sci Signal 2009, 2, ra72, doi:10.1126/scisignal.2000464.

18.   Krishnan, N.; Fu, C.; Pappin, D.J.; Tonks, N.K. H2S-Induced Sulfhydration
of the Phosphatase PTP1B and Its Role in the Endoplasmic Reticulum Stress
Response. Sci Signal 2011, 4, ra86, doi:10.1126/scisignal.2002329.

19.   Estrada-Ávila, A.K.; González-Hernández, J.C.; Calahorra, M.; Sánchez,
N.S.; Peña, A. Xylose and Yeasts: A Story beyond Xylitol Production. Biochim
Biophys Acta Gen Subj 2022, 1866, 130154, doi:10.1016/j.bbagen.2022.130154.

20.   Bergkessel, M.; Guthrie, C.; Abelson, J. Yeast-Gene Replacement Using
PCR Products. Methods Enzymol 2013, 533, 43–55, doi:10.1016/B978-0-12-
420067-8.00005-2.

21.   Paul, B.D.; Snyder, S.H. Protein Sulfhydration. Methods Enzymol 2015,
555, 79–90, doi:10.1016/bs.mie.2014.11.021.

22.   Cost, G.J.; Boeke, J.D. A Useful Colony Colour Phenotype Associated with
the Yeast Selectable/Counter-Selectable Marker MET15. Yeast 1996, 12, 939–
941,   doi:10.1002/(SICI)1097-0061(199608)12:10%3C939::AID-
YEA988%3E3.0.CO;2-L.

23.   Choi, K.-M.; Kim, S.; Kim, S.; Lee, H.M.; Kaya, A.; Chun, B.-H.; Lee, Y.K.;
Park, T.-S.; Lee, C.-K.; Eyun, S.-I.; et al. Sulfate Assimilation Regulates Hydrogen
Sulfide Production Independent of Lifespan and Reactive Oxygen Species under
Methionine Restriction Condition in Yeast. Aging (Albany NY) 2019, 11, 4254–
4273, doi:10.18632/aging.102050.

24.   Peña, A.; Sánchez, N.S.; González-López, O.; Calahorra, M. Mechanisms
Involved in the Inhibition of Glycolysis by Cyanide and Antimycin A in Candida
Albicans and Its Reversal by Hydrogen Peroxide. A Common Feature in Candida
Species. FEMS Yeast Res 2015, 15, fov083, doi:10.1093/femsyr/fov083.

25.   Sakai, H.; Suzuki, K.; Imahori, K. Purification and Properties of Pyruvate
Kinase from Bacillus Stearothermophilus. J Biochem 1986, 99, 1157–1167,
doi:10.1093/oxfordjournals.jbchem.a135579.

26.   Bergmeyer, H.V.; Gawhn, K.; Grassel, M. Alcohol Dehydrogenase. In
Methods of Enzymatic Analysis; Academic Press: New York, 1974; Vol. 1, pp.
428–429.

27.   Marino, S.M.; Li, Y.; Fomenko, D.E.; Agisheva, N.; Cerny, R.L.; Gladyshev,
V.N. Characterization of Surface-Exposed Reactive Cysteine Residues in
Saccharomyces Cerevisiae. Biochemistry 2010, 49, 7709–7721,
doi:10.1021/bi100677a.

28.   Alcock, L.J.; Perkins, M.V.; Chalker, J.M. Chemical Methods for Mapping
Cysteine Oxidation. Chem Soc Rev 2018, 47, 231–268,
doi:10.1039/c7cs00607a.

29.   Fu, L.; Liu, K.; He, J.; Tian, C.; Yu, X.; Yang, J. Direct Proteomic Mapping
of Cysteine Persulfidation. Antioxid Redox Signal 2020, 33, 1061–1076,
doi:10.1089/ars.2019.7777.

30.   Dashko, S.; Zhou, N.; Compagno, C.; Piškur, J. Why, When, and How Did
Yeast Evolve Alcoholic Fermentation? FEMS Yeast Res 2014, 14, 826–832,
doi:10.1111/1567-1364.12161.

31.   Marcet-Houben, M.; Gabaldón, T. Beyond the Whole-Genome
Duplication: Phylogenetic Evidence for an Ancient In-terspecies Hybridization in
the Baker's Yeast Lineage. PLoS Biol 2015, 13, e1002220,
doi:10.1371/journal.pbio.1002220.

32.   Shen, X.-X.; Opulente, D.A.; Kominek, J.; Zhou, X.; Steenwyk, J.L.; Buh,
K.V.; Haase, M.A.B.; Wisecaver, J.H.; Wang, M.; Doering, D.T.; et al. Tempo and

Mode of Genome Evolution in the Budding Yeast Subphylum. Cell 2018, 175, 1533-1545.e20, doi:10.1016/j.cell.2018.10.023.

33. Fabricio, M.F.; Valente, P.; Záchia Ayub, M.A. Oleaginous Yeast Meyerozyma Guilliermondii Shows Fermentative Metabolism of Sugars in the Biosynthesis of Ethanol and Converts Raw Glycerol and Cheese Whey Permeate into Polyunsaturated Fatty Acids. Biotechnol Prog 2019, 35, e2895, doi:10.1002/btpr.2895.

34. Gao, X.-H.; Krokowski, D.; Guan, B.-J.; Bederman, I.; Majumder, M.; Parisien, M.; Diatchenko, L.; Kabil, O.; Willard, B.; Banerjee, R.; et al. Quantitative H2S-Mediated Protein Sulfhydration Reveals Metabolic Reprogramming during the Integrated Stress Response. Elife 2015, 4, e10067, doi:10.7554/eLife.10067.

35. Aroca, Á.; Serna, A.; Gotor, C.; Romero, L.C. S-Sulfhydration: A Cysteine Posttranslational Modification in Plant Systems. Plant Physiol 2015, 168, 334–342, doi:10.1104/pp.15.00009.

36. Aroca, A.; Benito, J.M.; Gotor, C.; Romero, L.C. Persulfidation Proteome Reveals the Regulation of Protein Function by Hydrogen Sulfide in Diverse Biological Processes in Arabidopsis. J Exp Bot 2017, 68, 4915–4927, doi:10.1093/jxb/erx294.

37. Peng, H.; Zhang, Y.; Palmer, L.D.; Kehl-Fie, T.E.; Skaar, E.P.; Trinidad, J.C.; Giedroc, D.P. Hydrogen Sulfide and Reactive Sulfur Species Impact Proteome S-Sulfhydration and Global Virulence Regulation in Staphylococcus Aureus. ACS Infect Dis 2017, 3, 744–755, doi:10.1021/acsinfecdis.7b00090.

38. Aroca, A.; Schneider, M.; Scheibe, R.; Gotor, C.; Romero, L.C. Hydrogen Sulfide Regulates the Cytosolic/Nuclear Partitioning of Glyceraldehyde-3-Phosphate Dehydrogenase by Enhancing Its Nuclear Localization. Plant Cell Physiol 2017, 58, 983–992, doi:10.1093/pcp/pcx056.

39. Mitchell, A.R.; Yuan, M.; Morgan, H.P.; McNae, I.W.; Blackburn, E.A.; Le
Bihan, T.; Homem, R.A.; Yu, M.; Loake, G.J.; Michels, P.A.; et al. Redox
Regulation of Pyruvate Kinase M2 by Cysteine Oxidation and S-Nitrosation.
Biochem J 2018, 475, 3275–3291, doi:10.1042/BCJ20180556.

40. Ford, A.E.; Denicourt, C.; Morano, K.A. Thiol Stress-Dependent
Aggregation of the Glycolytic Enzyme Triose Phosphate Isomerase in Yeast and
Human Cells. Mol Biol Cell 2019, 30, 554–565, doi:10.1091/mbc.E18-10-0616.

41. Li, L.; Whiteman, M.; Guan, Y.Y.; Neo, K.L.; Cheng, Y.; Lee, S.W.; Zhao,
Y.; Baskar, R.; Tan, C.-H.; Moore, P.K. Characterization of a Novel, Water-
Soluble Hydrogen Sulfide-Releasing Molecule (GYY4137): New Insights into the
Biology of Hydrogen Sulfide. Circulation 2008, 117, 2351–2360,
doi:10.1161/CIRCULATIONAHA.107.753467.

42. Ju, Y.; Wu, L.; Yang, G. Thioredoxin 1 Regulation of Protein S-
Desulfhydration. Biochem Biophys Rep 2016, 5, 27–34,
doi:10.1016/j.bbrep.2015.11.012.

43. Wedmann, R.; Onderka, C.; Wei, S.; Szijártó, I.A.; Miljkovic, J.L.; Mitrovic,
A.; Lange, M.; Savitsky, S.; Yadav, P.K.; Torregrossa, R.; et al. Improved Tag-
Switch Method Reveals That Thioredoxin Acts as Depersulfidase and Controls
the Intracellular Levels of Protein Persulfidation. Chem Sci 2016, 7, 3414–3426,
doi:10.1039/c5sc04818d.

44. Paul, B.D.; Filipovic, M.R. Editorial: Molecular Mechanisms of Thiol-Based
Redox Homeostasis and Signaling in the Brain. Front Aging Neurosci 2021, 13,
771877, doi:10.3389/fnagi.2021.771877.

45. Campbell-Burk, S.L.; Shulman, R.G. High-Resolution NMR Studies of
Saccharomyces Cerevisiae. Annu Rev Microbiol 1987, 41, 595–616,
doi:10.1146/annurev.mi.41.100187.003115.

46.   Harsch, M.J.; Lee, S.A.; Goddard, M.R.; Gardner, R.C. Optimized Fermentation of Grape Juice by Laboratory Strains of Saccharomyces Cerevisiae. FEMS Yeast Res 2010, 10, 72–82, doi:10.1111/j.1567-1364.2009.00580.x.

47.   Wang, T.; Yang, Y.; Liu, M.; Liu, H.; Liu, H.; Xia, Y.; Xun, L. Elemental Sulfur Inhibits Yeast Growth via Producing Toxic Sulfide and Causing Disulfide Stress. Antioxidants (Basel) 2022, 11, 576, doi:10.3390/antiox11030576.

48.   Módis, K.; Ju, Y.; Ahmad, A.; Untereiner, A.A.; Altaany, Z.; Wu, L.; Szabo, C.; Wang, R. S-Sulfhydration of ATP Synthase by Hydrogen Sulfide Stimulates Mitochondrial Bioenergetics. Pharmacol Res 2016, 113, 116–124, doi:10.1016/j.phrs.2016.08.023.

Figure 1. H_2S productivity correlates with S-persulfidated proteins levels. (A) H_2S

productivity by *wt*, *met5Δmet10Δ* and *met17Δ*. Cells were incubated at 30°C on

YPDL plates for 4 days. (B) S-persulfidated proteins in *wt*, *met5Δmet10Δ* and

met17Δ strains. Whole cell extracts from exponential phase cultures were

subjected to the modified biotin switch assay with antibody against biotin (α-Biotin

Ab) to detect S-persulfidation.

Figure 2. Confirmation of S-persulfidated proteins by streptavidin beads

precipitation. A. Whole cell extracts from exponential phase cultures were subject

to the modified biotin switch assay, precipitated with streptavidin beads and

detected with antibodies specific to each protein; enolase (Eno2), glyceraldehyde

3 phosphate dehydrogenase (Tdh3), cystathionine beta synthase (Cys4) and

triose phosphate isomerase (Tpi1). HPDP-B: (N-[6-(biotinamido)hexyl]-3'-(2'-

pyridyldithio)propionamide). B. Whole cell extract from exponential phase

cultures after modified biotin switch assay was used as input control. (-) line

shows the proteins that reacts with anti-biotin antibody. (+) line shows biotinylated

proteins after modified biotin switch assay. TCL: Total cell lysate, PP STREP:

Streptavidin precipitation.

Figure 3. H_2S production reach a maximal during log phase. Yeast cell were

cultured in 96 wells microplate at 30°C and growth was measured at 600 nm. H_2S

was detected measuring methylene blue reduction at 663 nm.

Figure 4. H2S increase the activity of GAPDH and Pyruvate Kinase two hours

after stimu-lation. (A) Yeast cell cultures at exponential phase were treated with

NaHS 0.1 and 0.25 mM. Two hours later whole cell extracts were used to

measure GAPDH activity in vitro at 37°C. One-way ANOVA ** P<0.0001. (B)

Yeast cell cultures at exponential phase were treated with NaHS 0.1 and 0.25

mM. Seven hours later whole cell extracts were used to measure GAPDH ac-

tivity in vitro at 37°C. (C) Yeast cell cultures at exponential phase were treated

with NaHS 0.1 and 0.25 mM. Two hours later whole cell extracts were used to

measure Pyruvate Kinase activity in vitro at 37°C. (D) Yeast cell cultures at

exponential phase were treated with NaHS 0.1 and 0.25 mM. Seven hours later

whole cell extracts were used to measure Pyruvate Kinase activity in vitro at 37°C. One-way ANOVA * P<0.01. Closed circles, untreated cells; closed squares, NaHS 0.1 mM; closed triangles, NaHS 0.25 .

Figure 5. Exogenous and endogenous H_2S on yeast cells induce ethanol production and ATP synthesis. (A) BY4742 yeast cell cultures were treated with NaHS 0.1 mM and seven hour later supernatants were collected. Ethanol production was measured in vitro at 37°C. Closed circles, untreated cells; closed squares, NaHS 0.1 mM. Unpaired t * P=0.04. (B) Yeast cell cultures of the strains BY4741 and BY4742 supernatants were collected at 24 h. Ethanol production was measured in vitro at 37°C. Closed squares, BY4741; open squares, BY4742. Unpaired t * P=0.02 (C) Yeast cell cultures at exponential phase were treated with NaHS 0.1 and 0.25 mM. Two and four hours later whole cell extracts were lysated and ATP was quantified. ATP production was measure in vitro at 37°C. Closed circles, untreated cells; closed squares, NaHS 0.1 mM; closed triangles, NaHS 0.25 mM. One-way ANOVA ** P<0.001.

Figure 6. Yeast mutants that accumulate H_2S synthesize ethanol faster and more ATP than lower endogenously accumulated H_2S. (A) 48 hours precultures of yeast cell of the strains *wt*, *met5∆met10∆* and *met17∆* were resuspended in fresh media and supernatants were collected every hour. Ethanol production was measured in vitro at 37°C. (B) Yeast cell cultures at exponential and stationary phase of the strains *wt*, *met5∆met10∆* and *met17∆* were lysated and ATP was quantified. ATP production was measure in vitro at 37°C. One-way ANOVA * P<0.05, ** P<0.01. Open circles, *wt*; closed squares, *met5∆met10∆*; closed diamonds, *met17∆*.

Figure 7. Endogenous H_2S promotes basal oxygen consumption. 48 hours precultures of the *wt* and the mutants was diluted to an $OD_{600}= 0.2$, and oxygen consumption was measured. The basal oxygen consumption was measured in resting cells in a Clark electrode at 30°C. Seven hours after dilution oxygen consumption was measured again. One-way ANOVA * P<0.05, ** P<0.01. Open circles, *wt*; closed squares, *met5∆met10∆*; closed diamonds, *met17∆*.

Figure 8. Exogenous H_2S induce ethanol production in *K. marxianus* and *M.*
guilliermondi. (A) *K. marxianus* yeast cell cultures were treated with NaHS 0.1
mM and seven hour later supernatants were collected. Ethanol production was
measured in vitro at 37°C. Unpaired t ** P=0.002. (B) *M. guilliermondi* yeast cell
cultures were treated with NaHS 0.1 mM and 24 h later cells were treated again
with same concentration of NaHS. Seven hour later supernatants were collected.
Ethanol production was measured in vitro at 37°C. Unpaired t * P=0.04. Closed
circles, untreated cells; closed squares, NaHS 0.1 mM

A GAPDH activity 2h

B GAPDH activity 7h

C PK activity 2h

D PK activity 7h

Basal oxygen consumption rate

A *K. marxianus*

B *M. guilliermondi*