Platelet-derived transcription factors license human monocyte inflammation

Ibrahim Hawwari[1], Lukas Rossnagel[1], Nathalia Sofia Rosero Reyes[1], Salie Maasewerd[1], Marius Jentzsch[2], Agnieszka Demczuk[1], Lino L Teichmann[3], Lisa Meffert[3], Damien Bertheloot[1], Lucas S. Ribeiro[1], Sebastian Kallabis[1], Felix Meissner[1], Moshe Arditi[4,5], Magali Noval Rivas[4,5], and Bernardo S. Franklin[1*]

Affiliations:

1) Institute of Innate Immunity, University Hospitals, Medical Faculty, University of Bonn, Bonn, Germany

2) Institute of Clinical Chemistry and Clinical Pharmacology, University and University Hospital Bonn, Bonn, Germany

3) Department of Medicine III, University Hospital Bonn, Bonn, Germany.

4) Department of Pediatrics, Division of Pediatric Infectious Diseases, Guerin Children's, Cedars Sinai Medical Center, Los Angeles, CA USA.

5) Infectious and Immunologic Diseases Research Center (IIDRC), Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, California, USA.

*Correspondence: franklin@uni-bonn.de

KEYWORDS: Cytokines, Immunoparalysis, Immune Thrombocytopenia, Toll-like Receptors, Inflammasomes.

Running Title: Platelets supply monocyte inflammation with Transcription Factors

Key Points

- Platelets are essential to TLR and NLR cytokine responses of human monocytes,
- Immune thrombocytopenia leads to monocyte immunoparalysis;
- Platelet supplementation reverses monocyte immunoparalysis;
- Platelets transfer NFκB that reactivates cytokine production in genetically deficient monocytes.

ABSTRACT

CD14+ monocytes, the predominant population in human blood, are primarily engaged in host defense and pro-
inflammatory cytokine responses. Aberrant monocyte activity causes life-threatening cytokine storms, while
dysfunctional monocytes lead to 'immunoparalysis.' Understanding the mechanisms controlling monocyte
functions is therefore paramount. Here, we reveal platelets' vital role in human monocytes' pro-inflammatory
responses. Low platelet counts in immune thrombocytopenia (ITP) patients, or platelet depletion in healthy
monocytes result in monocyte immunoparalysis, characterized by reduced pro-inflammatory gene expression
and weakened cytokine responses to immune challenge. Remarkably, adding fresh platelets reverses monocyte
immunoparalysis. In mice, thrombocytopenia results in down-regulation of myeloid innate immune genes, and
compromised host defense transcriptional programs in monocytes despite normal responses to LPS. Platelets
control monocyte cytokines independently of traditional cross-talk pathways, acting as reservoirs of transcription
factors like NFκB and MAPK p38. We pinpointed megakaryocyte-derived NFκB2 transfer to human monocytes
by mass spectrometry-based proteomics. Functionally, platelets proportionally restored impaired cytokine
secretion in human monocytes lacking p38a and NFκB. We unveil the intercellular transfer of inflammatory
regulators, positioning platelets as central checkpoints in monocyte-mediated inflammation.

175 **words**

INTRODUCTION

Classical $CD14^+CD16^-$ monocytes constitute about 85% of the human circulating monocyte pool. Compared to other subsets, these monocytes are adeptly prepared for host defense programs, excel in innate immune detection of pathogens phagocytosis, and display an increased capability to release cytokines (Wong *et al*, 2011). Monocyte-driven innate immune responses can yield extreme outcomes. While heightened monocyte activity triggers hyper-inflammation and cytokine storms (Bonnet *et al*, 2021; Fajgenbaum & June, 2020; Ferreira *et al*, 2021; Jafarzadeh *et al*, 2020; Junqueira *et al*, 2022; Schulte-Schrepping *et al*, 2020; Vanderbeke *et al*, 2021), dysfunctional monocytes contribute to a potentially life-threatening state of hypo-responsiveness, known as 'immunoparalysis,' predisposing patients to opportunistic infections. Monocyte immunoparalysis is often observed after sepsis (Arens et al, 2016; Frazier & Hall, 2008; Roquilly et al, 2020), major visceral surgery (Frazier & Hall, 2008), and has recently been associated with the severity of SARS-CoV-2 infections (Agrati *et al*, 2020; Arunachalam *et al*, 2020). Monocytes are also vital cells mediating "trained immunity," a series of long-lasting epigenetic and metabolic adaptations that enhance innate immune responsiveness upon subsequent encounters with pathogen molecules (Bekkering *et al*, 2014; Netea *et al*, 2016). Hence, understanding the mechanisms regulating monocyte functions is exceptionally relevant in diverse clinical settings, as inappropriate monocyte activities can have long-lasting immunological consequences (Agrati *et al.*, 2020; Arunachalam *et al.*, 2020; Bonnet *et al.*, 2021; Ferreira *et al.*, 2021; Jafarzadeh *et al.*, 2020; Junqueira *et al.*, 2022; Schulte-Schrepping *et al.*, 2020; Vanderbeke *et al.*, 2021).

Here, we reveal a crucial role for platelets in the effector inflammatory functions of human monocytes. In the bloodstream, monocytes continuously interact with platelets, forming monocyte-platelet aggregates (MPAs) under physiological conditions (Rinder *et al*, 1991). MPAs increased in numerous inflammatory and thrombotic disorders and are usually associated with poor outcomes (Allen *et al*, 2019; Liang *et al*, 2015; Manne *et al*, 2020; Stephen *et al*, 2013). Interaction with platelets amplifies several effector functions of monocytes in numerous diseases (D'Mello *et al*, 2017; Fu *et al*, 2021; Rong *et al*, 2014; Singhal *et al*, 2017). Nevertheless, the physiological functions of platelets or MPAs to monocyte innate functions remain undefined. Moreover, studies in the human system are lacking and were primarily done in conditions where platelets are added to monocytes or describing pathological conditions where MPAs are enhanced.

In this study, we revealed a previously unrecognized critical dependency on platelets for the cytokine production of human monocytes. Using different human monocyte isolation kits and untouched monocytes from patients with primary immune thrombocytopenia (ITP), an autoimmune disease characterized by low blood platelet counts, we demonstrate that platelet numbers directly impact the cytokine output of $CD14^+$ monocytes towards Toll-like receptor (TLR) and Nod-like receptor (NLR) stimulation. Removal of platelets from healthy human monocytes caused monocyte immunoparalysis, characterized by transcriptional silencing of pro-inflammatory genes and impaired capacity to secrete pro-inflammatory cytokines. Notably, monocyte

immunoparalysis is dynamic and can be reversed by replenishing monocytes with fresh platelets. Moreover,

monocytes from ITP patients were inherently impaired in their capacity to produce cytokines in response to TLR

and NLR stimulation. Remarkably, the supplementation of ITP monocytes with healthy platelets reactivated

monocytes, reinvigorating their cytokine responses.

Mechanistically, we show that platelets are abundant cellular sources of transcription factors (TFs) and

signaling molecules, including numerous NFκB and MAPK p38 subunits, which are potent cytokine synthesis

regulators. Furthermore, we used Stable Isotope Labeling with Amino acids in Cell culture (SILAC) combined

with high-resolution mass spectrometry (MS)-based proteomics to demonstrate the transfer of platelet-derived

transcription factors (TFs), namely p65 (RelA), p52/p100 (NFκB2) and p38 MAPK into human monocytes.

Finally, we reveal the capacity of fresh platelets to bypass pharmacological inhibition or genetic ablation of p38,

RelA, and NFκB2 in human monocytes and restore their impaired cytokine secretion in a platelet:monocyte

ratio-dependent manner. Our findings elucidate the pivotal role of platelet-derived TFs in regulating monocyte

pro-inflammatory activity, suggesting the potential therapeutic application of platelet supplementation against

monocyte immunoparalysis. Our research challenges the earlier notion of monocytes being immune and self-

sufficient, unveiling their reliance on platelets for optimal pro-inflammatory cytokine production.

RESULTS

Platelets are required for cytokine secretion by human monocytes.

Hyper-inflammation, resulting from inflammasome activation in monocytes, leads to excessive production of

pro-inflammatory cytokines (Junqueira *et al.*, 2022; Rodrigues *et al*, 2021). To thoroughly analyze the

inflammasome-mediated cytokine response in human monocytes, we isolated classical monocytes (CD14$^+$CD16$^-$

) via immune-magnetic isolation kits (**Fig S1**, and **Materials and Methods**). Consistent with previous studies

(Bhattacharjee *et al*, 2017; Han *et al*, 2020; Rolfes *et al*, 2020), monocytes purified through standard magnetic

isolation methods, termed "standard monocytes" (StdMo), comprised a mixture of platelet-free monocytes,

monocyte-platelet aggregates (MPAs), and free platelets (Fig S1**A-D**). Employing a kit-provided platelet-

depleting antibody to remove platelets from StdMo allowed us to obtain platelet-depleted monocytes (PdMo)

while maintaining stable monocyte counts (Fig S1**A-C**). PdMo remained viable and were phenotypically similar

to StdMo (Fig S1**A**). However, upon exposure to lipopolysaccharide (LPS) or LPS + nigericin (LPS + Nig), the

secretion of IL-1β, TNFα, and IL-6 in PdMo drastically declined compared to StdMo, highlighting an essential

role of platelets in monocyte cytokine production (Fig **1A-B** and S1**D-E**). Notably, reintroducing autologous

platelets to PdMo (PdMo + Plts) revitalized PdMo's impaired cytokine responses. Importantly, platelets alone

(Plts) did not secrete any of the measured cytokines, confirming monocytes as their primary source. Consistent

with pyroptosis downstream of inflammasome activation, we also observed reduced viability in nigericin-treated

StdMo (Fig **1C**). However, platelet depletion (PdMo) led to decreased caspase-1 activity (Fig **1D**) and

intracellular maturation of IL-1β (Fig **1B**) while increasing cell viability (Fig **1C**). Platelet supplementation (PdMo

+ Plts) restored caspase-1 activity, IL-1β maturation, and release and re-sensitized platelet-depleted monocytes

to nigericin-induced pyroptosis (**Fig 1B-D**). These findings reveal the dependency on platelets for the monocyte

cytokine production. To validate our findings using alternative methods, we juxtaposed two techniques for

isolating CD14⁺ monocytes: positive and negative selection. We found different degrees of platelet

contamination between positive vs. negative selection of CD14⁺ monocytes (Fig S**2**). Positively selected StdMo

contained fewer platelets (Fig S**2B-C**) but responded poorly to TLR and inflammasome stimulation than

negatively isolated StdMo, which had significantly more platelets (Fig S**2D**). The impairment of CD14-positively-

selected monocytes was not due to pre-engagement of this receptor by the magnetic beads (Bhattacharjee *et*

al., 2017), as these cells also displayed reduced cytokine response to Pam3CysK4, a TLR1/2 agonist. Notably,

supplementing positively selected monocytes with platelets enhanced their cytokine responses (Fig S**2D**). These

findings conclusively demonstrate that platelets directly impact the cytokine output of monocytes and exclude

that the effects we observed were due to a specific isolation method. These findings support the requirement for

platelets for inflammasome functions of human monocytes.

In human monocyte-derived macrophages (hMDMs), platelets specifically license NLRP3 mRNA and protein

expression, exclusively boosting the activation of the NLRP3 inflammasome (Rolfes *et al.*, 2020). However, in

human monocytes, we found that platelet removal additionally modulated the activity of the AIM2 and NLRC4

inflammasomes, indicating a broader influence of platelets on monocyte inflammasome function (Fig **1E-F**) and

suggesting that platelets may influence additional innate immune sensors in monocytes. Supporting this

conclusion, platelet removal impacted the monocyte production of inflammasome-independent cytokines IL-6

and TNFα upon exposure to LPS (Fig **1A** and S**1E**), Pam3CysK4, a TLR2 ligand, and R848, which activates

TLR7 and TLR8 (Fig S**1F**). Platelet depletion further affected the production of numerous cytokines,

chemokines, and growth factors upon TLR4 and NLRP3 inflammasome activation (Fig **1G**) or TLR1/2 and

TLR7/8 activation (Fig S**1G**). While StdMo released copious amounts of IL-1β, IL-6, CCL2, CCL4, IL-8, PDGF,

and IL-1RA (> 10 ng ml⁻¹), the overall cytokine response of PdMo was blunted. However, the PdMo's impaired

cytokine response was reverted by reintroducing autologous platelets (50:1 platelet:monocytes ratio) to PdMo

(Fig **1G** and S**1G**). Hence, the disruption of cytokine responses of PdMo was not limited to a specific TLR or the

NLRP3 inflammasome but portrayed a widespread impairment of monocyte inflammation towards pattern

recognition receptors (PRRs). Together, our findings underscore the paramount role of platelets in the PRR-

induced cytokine response of human monocytes.

Platelet Supplementation reverts monocyte immunoparalysis in Immune Thrombocytopenia

To assess the impact of platelets on the effector functions of monocytes in a clinical setting where blood
platelet counts are inherently low, we isolated monocytes from patients with ITP. ITP is an autoimmune disorder
characterized by the destruction of platelets and impaired platelet production in the absence of infections or
other causes of thrombocytopenia (Cooper & Ghanima, 2019). Patients in our ITP cohort had low platelet counts
despite showing average blood leukocyte counts and plasma C-reactive protein concentrations (Fig **2A-D**).
Importantly, ITP patients were asymptomatic, free of infections, and were not treated with glucocorticoids (Fig
2A). We found that untouched monocytes taken from ITP patients (referred to as "ITPMo") were inherently
impaired in their capacity to produce cytokines in response to LPS and LPS + Nig stimulation compared to
monocytes from healthy volunteers (StdMo) (Fig **2E-G**). Remarkably, when healthy fresh platelets were
supplemented to ITPMo, their cytokine production was revitalized to levels equivalent to stimulated healthy
monocytes (StdMo), demonstrating that platelets can revert monocyte immunoparalysis. These findings confirm
the relevance of platelets as checkpoints for monocyte-driven immune responses in a clinical setting and
highlight the potential of platelet supplementation in counteracting monocyte immunoparalysis in ITP.

Platelet depletion induces transcriptional silencing of inflammatory genes in human monocytes.

To gain insights into the transcriptional changes in human monocytes upon platelet
depletion/supplementation, we analyzed the expression of 770 genes comprising the myeloid innate immune
response. We compared untouched StdMo, PdMo, and PdMo replenished with autologous platelets (PdMo +
Plts) under resting conditions or upon *ex vivo* stimulation with LPS (**Fig 3**).

Firstly, we examined the transcriptional effects of LPS or LPS + Nig stimulation in StdMo, PdMo, PdMo +
Plts, and platelets alone (Plts). Principal component analysis (PCA) revealed that platelet transcripts formed
distinct clusters separated from monocytes, indicating a negligible contribution of platelet transcripts to the
overall mRNA pool. The transcriptional response of stimulated monocytes was clearly distinguished from that of
unstimulated StdMo (**Fig 3A**), consistent with a typical LPS-induced gene expression (with 100 up-regulated and
54 down-regulated genes) compared to unstimulated conditions (**Fig 3B**, left panel). Strikingly, stimulated PdMo
showed a less distinct transcriptional profile, clustering closer to unstimulated StdMo (**Fig 3A**), indicating a loss
of transcriptional response to stimulation. Indeed, only four genes were significantly induced by LPS in PdMo,
indicating a general suppression of inflammatory gene expression. Notably, the reintroduction of autologous
platelets to PdMo (50:1 Platelet:Monocyte ratio) re-approximated their PCA-clustering towards StdMo (**Fig 3A**)
and restored the expression of 86% of LPS-induced genes in PdMo, resembling the profile of StdMo (**Fig 3B**,
right panel). These results were consistent with the multiplex cytokine analysis (**Fig 1G** and **S1F**).

Next, to specifically address the effects of platelet depletion/replenishment on monocytes, we separately
compared StdMo vs. PdMo in unstimulated and stimulated conditions (**Fig 3E-F**). Platelet depletion alone

modified 87 genes in steady-state monocytes, notably suppressing genes like MAPK14 (p38α) and BTK

involved in pro-inflammatory signaling. Meanwhile, numerous transcription factors (TFs) related to monocyte

differentiation and other processes were induced (Fig S**3A-C**). We identified several monocyte cytokine function

regulators among the most highly differentially expressed genes (DEGs). For example, genes involved in

sensing chemokines or external stimuli (e.g., *CCR2*, *FCGR3A*, and *CD14*) were down-regulated in PdMo. In

contrast, several TFs (e.g., *EGR2*, *PPARG*) and ERK-MAPK repressor genes (e.g., *ATF3* and *SPRY2*) were up-

regulated in PdMo (**Fig 3E-F** and S**3**). These findings demonstrate that transcriptional reprogramming underlies

the functional effects of platelet depletion in primary human monocytes. Furthermore, in line with the cytokine

secretion (**Fig 1G** and S**1F**), the transcriptional shutdown of pro-inflammatory gene expression in PdMos can be

reversed by their replenishment with fresh platelets.

Monocytes from thrombopenic mice display altered host immune defense transcriptional programs.

Our results in ITP patients demonstrate that monocytes from thrombocytopenic individuals display naturally

impaired cytokine responses to TLR and NLR activation. To recapitulate these findings *in vivo* and investigate

the underlying mechanisms, we induced thrombocytopenia in mice by i.v. injection of 2 mg/kg of body weight of

a rat anti-glycoprotein Ibα (GPIbα) mAb or, as control, with equal amounts of a rat IgG isotype (Sreeramkumar

et al, 2014; Xiang *et al*, 2013) (Fig **4A**). We then challenged platelet-depleted mice with i.v. injection of LPS (2

mg/kg). Antibody-dependent platelet clearance efficiently lowered blood platelet counts in treated mice (Fig **4B**).

Contrary to what our experiments with human monocytes indicate, but in line with earlier findings in mice

injected with LPS or in models of bacterial infection (Carestia *et al*, 2019; Claushuis *et al*, 2016; de Stoppelaar

et al, 2014; Xiang *et al*., 2013), platelet depletion did not significantly influence the plasma levels of IL-1β,

TNFα, and IL-6 in challenged mice (Fig S**4A**).

To investigate the effects of platelet depletion specifically on blood monocytes, we performed RNASeq

analysis on FACS-sorted (Ly6G⁻ IA/IE⁻ CD45⁺ CD11b⁺ CD115⁺) murine PBMCs. Consistent with the removal of

platelets by anti-GPIbα mAbs, the expression of several platelet transcripts (e.g., *Pf4*; *Gp9*; *Itga2b*, *Ppbp*,

Tubb1, *Treml1*, and *Clu*) was down-regulated in monocytes from platelet-depleted compared to IgG-treated mice

in the absence of immune stimulation (PBS group) (Fig S**4B**). Nevertheless, platelet depletion increased the

expression of genes involved in the complement cascade (e.g., *C1qa*, *C1qb*, and C1qc) and cell activation

processes (e.g. *Htra3* and *Mertk*) (Fig S**4B**), correlating with enrichment in pathways involved in blood

coagulation, hemostasis, platelet activation, complement, and wound healing processes (Fig S**4C**), and

suggesting that anti-GPIbα-mediated platelet depletion resulted in platelet activation.

We next compared the effects of platelets on the monocyte transcriptional response to LPS challenge *in*

vivo. Expression profiling analysis (Fig **4C**) and gene ontology (GO) (Fig **4E-F**) confirmed that monocytes

isolated from IgG-treated mice displayed a typical LPS-induced transcription profile (Alasoo *et al*, 2015) with the enrichment of genes involved in immune defense among the highest DEGs (e.g. *Cxcl10, Ccl12, Il1a, Ly6c*, interferon-stimulated genes (ISGs), and *Il12a*) and the concomitant and expected down-regulation of *Card10 and Cx3cr1* (Pachot *et al*, 2008) (Fig **4C**). Overall, LPS stimulation altered the expression of a higher number of transcripts in monocytes from platelet-depleted (anti-GPIbα-treated) compared to IgG-treated mice (4088 vs. 1282 transcripts, Fig S**4D**). However, the expression of several LPS-induced genes was comparable between anti-GPIbα and IgG-treated mice (Fig S**3E**). More than 80% of the LPS-induced gene signature was shared between monocytes from IgG-treated and platelet-depleted mice. While 3008 DEGs were exclusively expressed in monocytes from platelet-depleted mice, 202 DEGs were exclusively regulated in monocytes from IgG-treated mice (Fig S**4D**). Next, we focused the analysis to encompass the 770 myeloid innate immunity genes used in the human monocyte NanoString (**Fig 3**). Akin to the transcription profile of human PdMos, monocytes from thrombopenic mice had a significantly reduced expression of genes related to the myeloid innate immune response in response to i.v. LPS challenge (Fig **4D**). Consistent with the importance of the myeloid compartment in innate immunity, the LPS-induced expression profiles identified higher enrichment scores for pathways involved in host immune defense programs in IgG-treated mice (e.g., immune response to virus, immune system processes, cellular responses to IFNβ, and inflammation (Fig **4E-F**). These pathways were also represented but with lower enrichment scores in the gene profile of platelet-depleted mice, which was skewed towards apoptosis and cell cycle regulation (Fig **4E-F**). From these findings, we conclude that, although monocytes in thrombopenic mice maintain their proinflammatory cytokine capacity, platelet depletion induces transcriptional reprogramming towards cell cycle and division at the expense of immune defense programs. These findings align with previous observations of compromised immunity (Carestia *et al.*, 2019; Claushuis *et al.*, 2016; van den Boogaard *et al*, 2015; Xiang *et al.*, 2013), and higher susceptibility to sepsis in thrombopenic mice, despite normal inflammatory responses.

Platelets regulate cytokine secretion in human monocytes in trans and independently of classical platelet-monocyte crosstalk mechanisms.

Evidence highlights that innate immune cells can engulf platelets (Lang *et al*, 2002; Maugeri *et al*, 2009; Rolfes *et al.*, 2020; Senzel & Chang, 2013). Our observations did not reveal distinguishable internalization of platelets by human monocytes (Fig **5A**). However, blocking actin polymerization in platelet-depleted monocytes (PdMo) via Cytochalasin D (CytoD) (Fig **5B)** or Latrunculin B (Lat-B) (Fig S**5A**) ablated the ability of platelets to restore their faulty cytokine secretion. The potential role of endocytosis in this mechanism was also assessed, but the treatment of PdMo with Dynasore (DS), an inhibitor of dynamin 1/2 endocytosis, did not prevent the platelets' ability to reconstitute the faulty cytokine response of PdMo (Fig **5B**). These data underpins the significance of monocyte's phagocytic machinery for the influence exerted by platelets, albeit without directly

visible platelet phagocytosis, supporting a scenario where platelet-derived vesicles (PMPs), although too small
for imaging, are internalized by monocytes (Vajen Mause & Koenen, 2015). Supporting this conclusion, platelet
released molecules (Plts Sups) efficiently restored cytokine production in LPS-stimulated human PdMo (Fig **5C**).

Lastly, we examined whether ligand-receptor interactions mediate platelet-monocyte crosstalk. A well-
documented mechanism of monocyte-platelet or monocyte-PMP binding hinges on the interaction between
platelet P-selectin (P-sel or CD62P) and P-selectin glycoprotein ligand-1 (PSGL-1 or CD162) present on
monocytes (Frenette *et al*, 2000; Han *et al.*, 2020). To delineate the importance of the Psel-PSGL-1 axis, we
employed KPL-1, an antibody targeting P-sel. Although KPL-1 markedly reduced monocyte-platelet binding (**Fig
5D-E** and S**5B**), it neither inhibited pro-inflammatory cytokine production in stimulated StdMo nor impeded
platelets in restoring cytokine dysfunction in PdMos (**Fig 5F**). Although both soluble and immobilized P-sel can
influence innate immune functions in stimulated monocytes (Weyrich *et al*, 1995), and platelets can release P-
sel in vesicles (Forlow McEver & Nollert, 2000), introducing soluble recombinant human P-sel to PdMo didn't
rectify their impaired cytokine secretion, compared to the re-addition of autologous platelets (**Fig S5C**). These
insights suggest the P-sel-PSGL-1 interaction is not central to revitalizing the faulty cytokine response of PdMo.
Despite KPL-1 causing a physical separation between platelets and StdMo, it did not hinder cytokine secretion,
supporting the involvement of vesicles that may still be present.

As the platelet-monocyte crosstalk is additionally facilitated by platelet glycoprotein-Ib (GPIIb) and monocyte
CD11b (Carestia *et al.*, 2019; Malehmir *et al*, 2019), we blocked this ligand-receptor axis with anti-CD11b, or the
GPIIb/IIIa inhibitor trihydrochloride (GR-144053) (Fig S**5D**). Notably, neither targeting the GPIIb-CD11b complex
nor inhibiting GPIIb/IIIa prevented the platelet's ability to rescue the faulty cytokine production in platelet-
depleted monocytes (Fig S**5D**). Next, we blocked integrins, additional players in platelet adhesion, and
interactions with leukocytes (Bennett Berger & Billings, 2009; Nieswandt *et al*, 2000). We used RGD, an
antagonist of platelet glycoprotein IIb/IIIa receptors and other integrins (Haskel & Abendschein, 1989). We pre-
incubated platelets with RGD before their addition to PdMo. Irrespective of RGD, platelets similarly rescued
cytokine secretion of PdMo (Fig S**5E**), indicating that this class of integrins is not involved in the platelet
regulation of monocyte cytokine responses. Using similar inhibition strategies with mAbs, and simulation of
ligands with recombinant human proteins, we further excluded a role for RANTES (CCL5) (Alard *et al*, 2015),
CXCL12(Chatterjee *et al*, 2015), sialic acids (SAs)(Kullaya *et al*, 2018), Siglec-7(Varchetta *et al*, 2016), and the
CD40-CD40L axis(Henn *et al*; Inwald *et al*, 2003) in the effects described in our study (Fig S**5F-H**). Together,
our findings indicate that the classical and well-described mechanisms of platelet-monocyte crosstalk (i.e., co-
stimulatory molecules, integrins, CD40-CD40L axis, CCL5, CXCL12, and SAs) are not involved in the effects
described in our study.

276
Platelets are cellular sources of transcription factors (TFs)

Our transcriptomic approach revealed that platelets influence the expression of genes controlled by key
transcription regulators orchestrating pro-inflammatory cytokine production in monocytes, such as NFκB, BTK,
and MAPKs. We, therefore, profiled the activation of serine/threonine (STK) and protein tyrosine (PTK) kinase
networks on StdMo, PdMo, or PdMo + Plts as well as on platelets alone (Plts) and observed that platelet
depletion markedly impacted the kinome network of LPS-stimulated monocytes (Fig **6A-D** and S**6A**). With few
exceptions, the LPS-induced phosphorylation of numerous PTK and STK subtracts was significantly lower in
PdMos compared to StdMo (Fig **6C-D**). Substantiating the results of mRNA expression (**Fig 3**), platelet depletion
decreased the phosphorylation of BTK, p38 MAPK, MAPK-activated protein kinases (MAPKAPK), IKK kinases,
erythropoietin-producing human hepatocellular receptors (Eph), Cyclin-depended kinases (CDK), protein kinase
A, (PKA) and protein kinase C (PKC). In contrast, platelet-depleted monocytes displayed increased activation of
Janus kinases (JAKs) and IKKs, which are known modulators of NFκB signaling (Solt & May, 2008) (Fig **6A**). In
line with the altered function of these groups of kinases, pathway analysis revealed that platelet depletion
predominantly changed the activity of kinases involved in signal transduction associated with NFκB activation,
immune responses via CD40 signaling, MAPK, and JAK/STAT signaling pathways (Fig **6E**). Combined, the
transcriptome (Fig **3**) and kinome (Fig **6**) of human monocytes subjected to platelet depletion/re-addition
revealed dysregulation of key cytokine-regulatory pathways in PdMo.

Confirming the dynamic changes in NFκB activation caused by platelet depletion/reconstitution in human
monocytes, the LPS-induced phosphorylation of the NFκB subunits p55 (RelB) and p65 (RelA) was impaired in
PdMo, as measured by HTRF (Fig S**6B**) and immunoblotting (Fig **6F**), and it was restored by the re-addition of
platelets. Supporting that platelets regulate the NFκB activity on monocytes, increasing concentrations of
platelets enhanced TLR2-triggered NFκB activity in THP-1 monocytes encoding an NFκB-inducible promoter
(Fig **6G**). Together, these findings highlight the involvement of the NFκB and MAPK signaling pathways as
candidate mechanisms by which platelets regulate the cytokine production of human monocytes.

Interestingly, throughout our kinase and immunoblotting assays, platelets consistently showed high basal
kinase activity (Fig **6C-D**), and protein levels of NFκB subunits (RelA, p65) (Fig **6F** and S**6C**), and p38 MAPK
(Fig S**6D**), supporting prior findings that platelets are cellular sources of transcription factors (TFs) and signaling
molecules (Lannan *et al*, 2015). Indeed, by assessing the native and phosphorylated form of these proteins in
the lysates or supernatants of human platelets, we demonstrated that platelets express and release NFκB and
confirmed previous observations that they release p38α (Fig S**6C-D**), the major isoform expressed on human
and mouse platelets (Shi *et al*, 2017). Although phosphorylated p65 (P-p65) and P-p38 were predominantly
enriched in platelets, these proteins were additionally detected in platelet supernatants (Fig S**6D**) (Lannan *et al.,*
2015).

Platelets transfer TFs to monocytes.

Given the abundance of these critical transcriptional regulators in non-nucleated platelets (Beaulieu &
Freedman, 2009; Ezumi Takayama & Okuma, 1995; Lannan *et al.*, 2015; Poli *et al*, 2021) and the corresponding
dynamic activation of these pathways on platelet-depleted/replenished monocytes (**Figs 3** and **6**), we
speculated that platelets transfer TFs to monocytes. Cell-to-cell propagation of TFs and other signaling
molecules has been reported to occur through gap junctions between connecting cells, with biological functions
demonstrated in recipient cells (Ablasser *et al*, 2013; Kasper *et al*, 2010). However, little is known about the
trans-cellular propagation of signaling molecules when circulating cells adhere to one another.

To precisely delineate the repertoire of platelet proteins transferred to monocytes, we employed an unbiased
proteomic approach combined with Stable Isotope Labeling by Amino acids in Cell culture (SILAC). For this, we
cultured human megakaryocytes (MEG-01 cells) for several days in a medium containing stable isotope labeled
amino acids $C^{13}L^{15}$] L-lysine and [$C^{13}L^{15}$] L-Arginine (hereafter referred to as heavy AAs). Under these
conditions, newly translated proteins in MEG01 incorporate heavy AAs and can be distinguished from pre-
existing proteins on monocytes by a mass shift (Wolf *et al*, 2020). Of note, cell-free supernatants from MEG-01
(termed MK-Sups), akin to platelets, were equally capable of restoring the impaired cytokine responses
observed in PdMos (Fig **7A**). Subsequently, PdMos were incubated with MK-Sups, followed by a stringent
washing protocol to eliminate non-specific associations before subjecting the monocyte lysates to MS analysis
(Fig **7B-C**). We detected 33 proteins harboring heavy AAs enrichment in PdMos exposed to MK-Sups under
LPS-stimulated conditions (Fig **7D**) and numerous TFs in unstimulated monocytes (Fig S**7A**). Our data revealed
notable entities among these proteins, including the transcriptional activator NFκB2 (p100/p52), transcriptional
regulators (e.g., ANP32B, THOC1), enzymes involved in mRNA translation and stability (ZFP36, EOF2A, and
PAIP1), and proteins implicated in the MAPK/ERK signaling cascade (Fig S**7**). Furthermore, 5 out of the 33
heavily labeled proteins enriched in PdMo after LPS were involved in transmembrane functions, pointing
towards a mechanism for the internalization of MK-derived proteins. We confirmed the expression and release
of NFκB2 on human platelets by immunoblotting (Fig **7E**). These findings substantiate the transfer of platelet-
originated transcriptional and signaling molecules to monocytes, underscoring a potentially crucial intercellular
communication mechanism.

Platelet-derived TFs license monocyte innate immune responses.

Next, to specifically address whether platelet-derived NFκB or p38 functions in monocytes, we tested
whether pharmacological inhibition of NFκB and p38 MAPKs on platelets interfere with their capacity to
reactivate cytokine production in PdMos (**Fig 8A**). Indeed, pre-treatment of platelets with the irreversible
inhibitor of IKKβ phosphorylation BAY 11-7082 (BAY) or the p38 inhibitor SB203580 (Yamashita Hishinuma &
Shimada, 2009) extinguished their ability to recover cytokine response of PdMo (**Fig 8B**). The addition of intact
platelets, but not NFκB- or p38-inhibited platelets, restored IL-6 and TNFα secretion of LPS-stimulated PdMo as

well as LPS + nigericin-induced IL-1β production (**Fig 8B**). To account for inhibitor carryover, we separately

treated monocytes with BAY or SB203580 (**Fig 8B**, last two right bars). BAY blocked the thrombin-induced P-

selectin expression on treated platelets (Fig S**8A-B**). While BAY completely abrogated monocyte cytokine

responses to TLR and NLRP3 activation, direct incubation of monocytes with SB203580 partially inhibited their

response. These findings support that p38 and NFκB or their downstream signaling in platelets are essential for

the platelet regulation of cytokine responses of human monocytes.

Next, to address whether platelet-derived TFs function in recipient monocytes, we examined whether

supplementation with fresh platelets could bypass NFκB or p38 inhibition in primary human monocytes. To

demonstrate this, we pre-incubated monocytes with high doses of BAY (50 μM) and SB203580 (20 μM). After

washing the inhibitors away, we supplemented BAY- or SB203580-treated monocytes with growing

concentrations of intact platelets freshly isolated from the same donors (Fig **8C**). In line with previous

observations of toxicity associated with BAY-11-7082 in human cells (Rauert-Wunderlich *et al*, 2013; White &

Burchill, 2008), BAY-treated monocytes displayed decreased viability (Fig **8D**), precluding us from addressing

the importance of platelet-derived NFκB in these experiments. Consistent with dysfunctional p38 signaling (Lee

et al, 1994), SB203580-treated monocytes were incapable of secreting cytokines in response to LPS or LPS +

Nig (Fig **8E**). Strikingly, the addition of intact platelets dose-dependently bypassed p38 inhibition and restored

cytokine secretion in SB203580-treated monocytes (Fig **8E**). Hence, these findings support that platelet

supplementation restores dysfunctional p38 activity in p38-inhibited monocytes in a dose-dependent manner.

Supporting this conclusion, SB203580-treated platelets failed to convert cytokine responses in SB203580-

treated monocytes (Fig S**8C**).

Finally, to specifically address the function of platelet-derived NFκB on monocytes and to simulate a

scenario in which platelets would be the only cellular source of functional NFκB, we genetically ablated RelA or

NFκB2 in THP-1 monocytes using CRISPR-Cas9 (Fig **8F**). Again, and consistent with a genetic deficiency in

these molecules, confirmed by immunoblotting (Fig **8F**), RelA[−/−] or NFκB2[−/−] monocytes were unable to secrete

IL-1β in response to Pam3Cysk4 + Nig stimulation (Fig **8G**). As previously reported (Rolfes *et al.*, 2020),

platelets boosted IL-1β secretion in wild-type monocytes (Fig **8G**). Notably, adding platelets to RelA[−/−] or

NFκB2[−/−] monocytes dose-dependently boosted their production of IL-1β, demonstrating the reactivation of

these pathways in otherwise knockout recipient cells. These findings demonstrate that platelets supply human

monocytes with functional signaling TFs, which license their optimal cytokine responses and reveal a novel layer

of cell-to-cell communication in innate immunity.

DISCUSSION

Monocytes play a pivotal role in the innate immune system, acting as both frontline defenders against pathogens, being progenitors of macrophages and dendritic cells, and releasing pro-inflammatory cytokines like IL-1β, IL-6, and TNFα crucial in initiating the inflammatory response (Guilliams Mildner & Yona, 2018). Unrestricted monocyte activity can lead to detrimental hyperinflammation, manifesting as cytokine storms, tissue damage, and even death. Regulating monocyte activity is, therefore, essential for maintaining immune homeostasis, with platelets playing a notable role.

In the blood, platelets surround monocytes, with whom they engage in complex interactions, forming MPAs. These aggregates are common in homeostasis but intensify during inflammation, serving as indicators of inflammatory diseases (Allen *et al.*, 2019; Barrett *et al*, 2019; Manne *et al.*, 2020). Despite their prevalence, the specific roles of MPAs in both health and disease remain to be fully elucidated.

Our research has uncovered that platelets are integral to sustaining the inflammatory responses of monocytes. Depletion of platelets from human monocytes results in an anergic state, with reduced cytokine production and downregulation of pro-inflammatory genes following stimulation of pattern recognition receptors (PRRs). This hyporesponsive state is also observed in monocytes from ITP patients, who have an impaired ability to produce cytokines in response to PRR activation. Remarkably, reintroducing autologous platelets to platelet-depleted monocytes reversed monocyte immunoparalysis, underscoring the crucial role of platelets in modulating monocyte function. We extended these investigations to the clinical domain by demonstrating the ability of healthy platelets to revert the immunoparalysis of ITP monocytes.

Immunoparalysis is commonly observed in occasions of increased blood loss, such as trauma, major organ surgery (Haupt *et al*, 1998), hemorrhagic fevers (Vangeti *et al*, 2021), sepsis (Cao Yu & Chai, 2019; Weisheit *et al*, 2020), liver failure (Lin *et al*, 2007; Wasmuth *et al*, 2005), and more recently in SARS-CoV-2 infections (Agrati *et al.*, 2020; Arunachalam *et al.*, 2020). Thrombocytopenia is a common feature among these disparate conditions. However, additional confounding factors, such as co-infections and lower leukocyte counts, render it difficult to assess the actual contribution of platelets to monocyte immunoparalysis. Our study provided evidence of monocyte immunoparalysis in ITP patients without other confounding factors such as infections or glucocorticoids and with average leukocyte counts and C-reactive protein levels.

Our findings reinforce the longstanding belief that normal platelet counts are essential for maintaining immune homeostasis. Low platelet count is a significant risk factor for infections (Birnie *et al*, 2019; de Stoppelaar *et al.*, 2014), and ITP patients experience compromised host defense, which is associated with a higher risk of infection, poorer treatment response, and more extended hospital stays (Qu *et al*, 2018). In spite of that, systemic levels of IL-1β, IL-6, and TNFα in ITP patients are usually comparable to healthy individuals (Andreescu, 2023). Our findings *in vivo* recapitulate the human scenario and suggest that platelet depletion

skews the immune response, prioritizing cell survival over host defense. Platelet-depleted mice exhibited no
significant differences in plasma levels of pro-inflammatory cytokines or in their monocyte transcriptional
response to intraperitoneal LPS. However, monocytes from these mice displayed weakened transcription
programs associated with host defense. Although contradictory to the transcriptional profile of platelet-depleted
human monocytes, our findings *in vivo* are fully corroborated by mouse models of sepsis, where
thrombocytopenia is associated with disturbed immune defense programs and increased mortality independently
of disease severity but without influencing local or systemic inflammation, plasma cytokine levels, and neutrophil
recruitment (Carestia *et al.*, 2019; Claushuis *et al.*, 2016; van den Boogaard *et al.*, 2015; Xiang *et al.*, 2013).
Furthermore, and in agreement with our findings, platelet depletion resulted in enhanced coagulation,
endothelial cell activation, and even increased plasma levels of some pro-inflammatory cytokines (Carestia *et*
al., 2019; Claushuis *et al.*, 2016; van den Boogaard *et al.*, 2015; Xiang *et al.*, 2013). Overall, our findings
highlight platelets as essential checkpoints determining the magnitude of monocyte innate immune programs
and may have consequences for the treatment of immunosuppression or other life-threatening conditions
associated with unrestrained monocyte responses (Ferreira et al., 2021; Junqueira et al., 2022).

The interaction between platelets and monocytes enhances various monocyte functions, such as their ability
to migrate through vessel walls, produce reactive oxygen species, and differentiate into dendritic cells,
ultimately aiding in the activation of CD8+ T cells (D'Mello *et al.*, 2017; Fu *et al.*, 2021; Han *et al.*, 2020; Rong
et al., 2014; Singhal *et al.*, 2017). However, the molecular mechanisms of how platelets influence these
monocyte functions remain partially understood (Kral *et al*, 2016). Our study introduces a novel mechanism of
platelet regulation on monocyte immunity, highlighting platelet-derived NFκB as a key trans-cellular regulator of
cytokine production in monocytes. Adding fresh platelets counteracted genetic ablation of NFκB signaling in
human monocytes. Furthermore, while pharmacological NFκB inhibition was toxic to primary human monocytes,
we found that platelets bypass SB203580 inhibition of p38 MAPK. These findings align with existing literature
that specific ablation of p38 MAPK in platelets confers protection against myocardial infarction (Shi *et al.*, 2017)
and platelet-activating factor (PAF)-induced lethality (Abhilasha *et al*, 2019). Indeed, MAPK p38 inhibitors have
emerged as pertinent targets for cytokine-suppressive anti-inflammatory drugs (Anand *et al*, 2011; Christie *et al*,
2015; Yong Koh & Moon, 2009).

We also corroborated previous findings indicating a rich stock of NFκB subunits in platelets and MKs
(Beaulieu & Freedman, 2009; Lannan *et al.*, 2015; Spinelli *et al*, 2010). Transcriptional and kinome analyses
have provided insight into the molecular underpinnings of this interplay, revealing that platelets contribute to key
cytokine-regulating signaling pathways in monocytes, notably NFκB and MAPK. Interestingly, it has been
previously demonstrated that interaction with platelets enhances the nuclear translocation of p65 within
monocytes (Ueda *et al*, 1994; Weyrich *et al*, 1996; Weyrich *et al.*, 1995). However, the possibility that a
considerable portion of p65 arises from platelets was neither raised nor experimentally demonstrated. Our study

provides the first evidence for the translocation of NFκB from platelets to monocytes, with maintained functions
in monocytes. We have shown that platelets can reactivate TLR-induced cytokine secretion in RelA$^{-/-}$ and
NFκB2$^{-/-}$ THP1 monocytes, pinpointing platelets as the only viable source of these molecules.

Although platelets could also reconstitute cytokines in SB203580-treated monocytes, our mass speck
proteomics approach did not identify platelet-derived p38a MAPK on human monocytes. However, as MS is an
abundance-based method, we cannot exclude the transfer of low anounts of p38a contributing to pathway
activity. As p38 MAPK signaling is required for the transcriptional NFκB activity (Anderson, 2010; Gottschalk *et*
al, 2016; Guma *et al*, 2011; Saha Jana & Pahan, 2007), platelets may circumvent SB203580 inhibition in
monocytes by supplying NFκB. Supporting this hypothesis, SB203580 reduced the phosphorylation of p65
(RelA) in treated monocytes (Fig. **EV7E**).

Although the transfer of TFs from platelets to monocytes has been anticipated (Lannan *et al.*, 2015), it has
never been experimentally demonstrated. In this regard, our study establishes platelet-monocyte TF transfer as
a sophisticated mechanism of intercellular communication. This finding is particularly intriguing given the unique
presence of platelets in mammals, as opposed to invertebrates, which rely on a single-cell type for host defense
and wound healing.

Although we ruled out the participation of numerous well-known ligand-receptor interactions in the platelet-
monocyte crosstalk, the mechanisms by which platelets supply TFs to monocytes remain unidentified. Though
we have not visualized phagocytosis of platelets by monocytes by confocal microscopy, using phagocytosis
inhibitors, and transfer of platelet supernatants, we ruled out the involvement of a humoral platelet-derived
factor, supporting that vesicles restore cytokine activity in platelet-depleted monocytes, a known mechanism for
the transfer of cellular receptors (Rozmyslowicz *et al*, 2003), bioactive lipids (Barry *et al*, 1997; Rossaint *et al*,
2016) and intact organelles (Levoux *et al*, 2021) that are transferred from platelets, or platelet vesicles, to
leukocytes, with preserved activity in the cytosol of recipient cells. We also observed numerous platelets bound
to monocytes, in line with previous observations that a significant fraction of blood monocytes (up to 45%) are
physiologically associated with platelets in steady-state conditions (Rinder *et al.*, 1991). Furthermore, breaking
MPAs via interfering with P-sel-PSGL-1 interactions did not prevent the platelet effect on monocytes. However,
based on our experiments, we cannot exclude that cell-surface receptor-activated signaling cascades involving
NFκB contribute to platelet-induced monocyte activation. In a randomized clinical trial, two anti-platelet
medications reduced MPA formation and the systemic levels of CCL2, IL-6, IL-8, and TNFα in humans
challenged with intravenous LPS injection (Thomas *et al*, 2015). However, depending on the disease setting,
platelets attenuate monocyte inflammatory responses by triggering the production of IL-10 or sequestering IL-6
and TNF-α released by monocytes (Carestia *et al.*, 2019; Xiang *et al.*, 2013). Hence, platelet-monocyte
interactions can have different and context-dependent outcomes (Kral *et al.*, 2016). Our findings further indicate
that other classical and well-described mechanisms of platelet-monocyte crosstalk (i.e., co-stimulatory

molecules, integrins, CD40-CD40L axis, CCL5, CXCL12, and SAs) are not involved in the effects described.

The exclusion of these mechanisms highlights a previously unrecognized mechanism of immune regulation,

where platelets license the pro-inflammatory cytokine responses of human monocytes. Our findings indicate that

the complex association between monocytes and platelets may be part of a homeostatic mechanism to equip

monocyte immune functions when needed.

In summary, our study offers a novel perspective on the regulatory role of platelets in monocyte-mediated

immunity. By identifying platelet-derived NFκB and MAPK signaling as crucial for monocyte cytokine production,

we open the door to potential therapeutic strategies that could modulate this platelet-monocyte interaction to

manage inflammatory conditions and improve outcomes in diseases characterized by monocyte dysfunction.

Our findings may also help to pave the way for new clinical approaches for improving platelet transfusions,

which are usually safe but can occasionally have deleterious inflammatory consequences.

METHODS

Reagents

Cell culture reagents (e.g., PBS, Fetal calf serum, GlutaMax, and RPMI) were from Gibco, Thermo Fisher Scientific. Stimuli were as follows: LPS (Ultrapure from E.coli O111:B4), Pam4CysK4, and R848 (Resiquimod) were from Invivogen. Nigericin acid was from Thermo Fisher Scientific. Inhibitors used were Cytochalasin D, BAY 11-7082 (Sigma-Aldrich). The CellTiter-BlueTM Cell Viability Assay and Caspase-Glo. 1 Inflammasome Assay was from Promega. HTRF kits to detect IL-1b, IL-6, TNFa, and NFκB were from Cisbio. ProcartaPlexTM kits for detecting other Cytokine/Chemokine/Growth Factors were from Invitrogen. Antibodies were anti-IL-1b antibody (BAF401, R&D Systems, 1:1,000). DRAQ5 was from Thermo Fisher Scientific. For FACS, we used FcR Blocking Reagent, human and mouse (Miltenyi Biotec). Anti-human CD41 directly conjugated to Alexa Fluor 488 (Clone: HIP8), Anti-human CD14-A647 (Clone: HCD14), and isotype Mouse IgG1, k Isotype Ctrl (Clone: MOPC-21) Antibody were all from BioLegend.

Study Subjects

Peripheral blood was obtained by venipuncture of healthy volunteers after the signature of informed consent and approval of the study by the Ethics Committee of the University of Bonn (Protocol #282/17), following the Declaration of Helsinki. Patients with immuno-thrombocytopenia were recruited from a tertiary hematology referral hospital. We recruited five patients (3 females and two males, average age of 46 ± 9) from December 2019 to May 2020. All patients were diagnosed with ITP, and comorbidities/infections were ruled out. Written informed consent was obtained from all subjects according to the Declaration of Helsinki and approval by the Institutional Review Board of the University of Bonn (313/21). All participants were instructed about the study's objectives and signed an informed consent following guidelines for human research.

Primary cells:

Primary human monocytes were isolated from peripheral blood collected from healthy volunteers in S-Monovette K3EDTA tubes. Blood was diluted 1:3 in PBS, and peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll® Paque PLUS density gradient centrifugation at 700 x g, 20 min. Primary human CD14+ monocytes were isolated from PBMCs using positive magnetic separation with the Classical Monocyte Isolation Kit (Miltenyi Biotech) or negative selection with the EasySep™ Human Monocyte Isolation Kit (STEMCELL™ Technologies) following manufacturer instructions. Platelet-depleted monocytes were generated by adding a Platelet Removal Component (50 μl ml⁻¹) supplied with the isolation kit (STEMCELL™ Technologies). The purity of monocyte populations was assessed by flow cytometry. Cells were incubated with

FcR Blocking Reagent for 15 min at 37 °C, followed by staining with fluorochrome-conjugated monoclonal antibodies against human CD14 APC (eBioscience), anti-CD45 PE (eBioscience), and anti-CD41a FITC (eBioscience) for 30 min at RT. Cells were washed with PBS, and fluorescence was measured on a MACSQuant® Analyzer 10.

Platelet isolation

Human platelets were isolated from venous blood from healthy volunteers collected as previously described[22, 87]. Briefly, whole blood was centrifuged at 330 x g for 15 min, and platelet-rich plasma (PRP) was collected. To prevent platelet aggregation during isolation, PRP was treated with 200 nM Prostaglandin E1 (PGE1), diluted 1:1 with PBS, and centrifuged at 240 x g for 10 min to pellet leukocytes. Platelet suspensions were collected and washed by centrifugation at 430 xg for 15 min in the presence of PBS PGE1. Washed platelets were pellet and resuspended in pre-warmed RPMI. Human platelets were stimulated with 1 U ml−1 Thrombin from human plasma (Sigma-Aldrich) or left unstimulated for 30 min to assess the purity and activation of isolated cells. Platelets were incubated with FcR Blocking Reagent followed by staining with CD45 (leukocyte marker), and CD41a FITC (eBioscience) and anti-CD62P APC (eBioscience).

Generation of platelet supernatants

Platelet supernatants were generated from suspensions of 1×10^8 human platelets incubated in RPMI and left untreated (PltSups) or stimulated with 2 ng ml−1 LPS (PltsLPS-Sups) for 4.5 h at 37ºC. Cell suspensions were centrifuged at 3,000 x g for 10 min. The platelet supernatants were harvested and used to stimulate human monocytes.

Cell lines

The human monocytic THP-1 cell line (ATCC TIB-202) and the THP-1 DualTM reporter cell (InvivoGen, Thpd-nfis) were culture in RPMI supplemented with 10% heat-inactivated fetal calf serum, 1% Penicillin/Streptomycin and 1x GlutaMax. For THP-1 DualTM cells, the growth medium contained additionally 25 mM HEPES. Cells were either studied in a monocyte-like phenotype (without PMA-differentiation) or after their differentiation into macrophages by PMA (50 µM) treatment for 24 h.

Stimulation Assays

Monocytes (1 x 10^5/well) were cultured alone or co-cultured with 50 (5 x 10^6/well) to 100 platelets (1 x 10^7/well).

For stimulation, cells were treated with agonists of TLR1/2: Pam3CysK4 (1 µg ml^{-1}); TLR4: LPS (2 ng ml^{-1}); or

TLR7/8: Resiquimod R848 (3.5 µg ml^{-1}); for for 4.5h. For inflammasome activation, cells were primed with a

TLR agonist for 3 h (indicated in Figs), followed by activation with nigericin (10 µM, 90 min) Prgl from *Bacillus*

anthracis (100 ng ml^{-1}) and protective antigen (PA, 1 µg ml^{-1}), for 90 min) or transfection of poly(dA:dT, 0.5 µg

ml^{-1}, 3 h). After stimulation, CFS was harvested and used for cytokine detection.

Cytokine measurement

558   According to the manufacturer's instructions, cytokines were detected in cell-free supernatants or whole cell

559   lysates by homogeneous time-resolved fluorescence technology (HTRF, Cisbio). A Human ProcartaPlexTM

560   (Invitrogen) immunoassay was additionally used to detect 45 human cytokines, chemokines, and growth factors,

561   according to the manufacturer's instructions, using a MAGPIX® System.

Cell Viability Assays

Cell viability was determined by measuring the release of Lactate Dehydrogenase Assay (LDH) in CFS or the

CellTiter-Blue Assay (CTB) in stimulated cells. For LDH measurement, CFS from stimulated cells were

incubated with LDH buffer for 30 min at 37 °C without CO2, and absorbance was measured at 490 nm using a

Spectramax i3 plate reader (ThermoFisher). Cells were incubated with CTB buffer at 37°C, and the

fluorescence was read at different time points in both assays, cells treated with Triton X-100 served as a 100%

cell death control. Fluorescence was detected with a SpectraMax.

Assessment of Caspase-1 Activity

572   The specific activity of Caspase-1 was assessed in cell-free supernatants using Caspase-Glo® 1 Inflammasome

573   Assay (Promega). Briefly, CFS was mixed 1:1 with Glo Buffer and incubated for 30 min at RT. Luminescence

574   was measured with SpectraMax i3 (Molecular devices).

Confocal Imaging

Platelet interactions with human monocytes were imaged in a Leica TCS SP5 SMD confocal system (Leica

Microsystem). Cells were seeded in microslide 8-Well IBIDI chamber, fixed with 4% paraformaldehyde (PFA) for

30 min, washed and incubated with staining mAbs in permeabilization/blocking buffer (10% goat serum, 1%

FBS, and 0.5% Triton X-100 in PBS) supplemented with human FcR Blocking Reagent for 30 min. Cells were

stained overnight at 4ºC with anti-human CD41Alexa Fluor 488, anti-human CD14-A647, or Hoechst 34580
(ThermoFisher).

Immunoblotting

Cells were lysed with RIPA complete lysis buffer supplemented with EDTA-free protease and phosphatase
inhibitors (Roche) and 25 U of Benzonase Nuclease. Samples were loaded onto NuPAGETM Novex 4-12% Bis-
Tris gels and transferred to Immobilon-FL Polyvinylidene fluoride (PVDF) membranes. After blocking with 3%
BSA [w/v] in Tris-buffered saline (TBS), membranes were incubated overnight with primary antibodies diluted in
TBS containing 3% BSA [w/v] and 0.1% Tween. After washing steps, membranes were incubated with
secondary antibodies conjugated to IRDye680 or IRDye800 (1:25000) for 2 h at RT. The membrane was
scanned at Odyssey Imager (Li-Cor Biosciences). The following primary antibodies were used for: β-Actin rabbit
mAb (1:5000), Phospho-NFκB p65 (Ser536) (93H1) Rabbit mAb #3033 (1:1000), NFκB p65 (C22B4) Rabbit
mAb #4764 (1:1000) and IkBa (L35A5) Mouse mAb (Amino-terminal Antigen) #4814 (1:1000).

Platelet depletion *in vivo*

In vivo, platelet depletion was performed in female C57BL/6J mice (12 weeks) by intravenousl (i.v.) injection of 2
mg/kg of polyclonal rat anti-mouse GPIbα (R300) or same amounts of a polyclonal rat IgG none-immune (C301)
antibody as control (Emfret Analytics). Mice were then challenged 12 hours later with i.v. injection of 2 mg/kg
LPS or PBS as control. Blood was collected after 3 h, and PBMCs were isolated by density gradient
centrifugation in Ficoll Paque PLUS. Mouse monocytes were separated from PBMCs by FACS-sorting gating on
Ly6G⁻, IA/IE⁻, CD45⁺, CD11b⁺, and CD115⁺ cells and sorted directly into QIAzol lysis reagent. Lysates were
frozen in liquid nitrogen and sent to GENEWIZ Azenta Life Sciences for bulk RNA sequencing. All animal
experimentation was approved by the local ethical committee (LANUV-NRW #84-02.04.2016.A487). Mice were
purchased from Charles River and housed at the House for Experimental Therapy (HET) of the University
Hospital of Bonn.

Fluorescence-activated cell sorting was used to obtain mouse monocytes. Mouse blood monocytes were
isolated from pooled blood of 4 mice per experimental group in 3 different experiments. For this, mouse PBMCs
were isolated through Ficoll gradient and stained with anti-Ly6G BV785 (Biolegend, dilution: 1:40), anti-I-A/I-E
BV785 (Biolegend, dilution: 1:80), anti CD45 FITC (Biolegend, dilution: 1:200), anti-CD11 b PE (eBioscience,
dilution: 1:166), anti-CD115 APC (Biolegend, dilution: 1:80) and Hoechst 33258 (Abcam, 1 µM) as viability
staining. Finally, viable Ly6G-I-A/I-ECD45+ CD11b+CD115+ monocytes were sorted directly into QIAzol lysis

reagent using the BD FACS Aria Fusion (387333827) and BD FACS Aria III (216372545) at the Flow Cytometry
Core Facility of the Medical Faculty of the University of Bonn.

Bulk RNA sequencing of isolated murine monocytes

For bulk RNA sequencing, samples were processed according to GENEWIZ (Azenta Life Sciences) pipeline.
RNA extraction was performed, followed by a library preparation. Ultra-low input RNA-Seq was performed at
Illumina HiSeq PE 2x150 bp with ~350M reads. Reads were then mapped to the Mus musculus GRCm38
reference genome (ENSEMBL) using the STAR aligner v.2.5.2b, and unique gene hit counts were generated
with the featureCounts from the Subread package v.1.5.2. Finally, standard analysis was performed by
GENEWIZ. Differential gene expression analysis was performed using DESeq2, p-values, and log2 fold changes
were calculated by applying the Wald test, and genes with adjusted p-value < 0.05 and absolute fold change > 2
were considered significantly altered differentially expressed genes (DEGs). Gene ontology (GO) analysis of the
significant DEGs was applied by GENEWIZ (Azenta Life Sciences) using the Fisher exact test. GO terms were
assigned using the GeneSCF v.1-p2 software. Data visualization was created using the R package ggplot2.

Transcriptional profiling of human monocytes and platelets with NanoString

According to the manufacturer protocol, the nCounter® Human Myeloid Innate Immunity Panel v2 (NanoString®
Technologies) was used to assess the mRNA expression of 770 human transcripts. Briefly, single or co-cultures
of monocytes and platelets were lysed in RLT Buffer (Qiagen) containing β-Meracptoethanol (1 x 10^4 cells/µl).
RNA was isolated with and homogenized with CodeSets and left for hybridization overnight. RNA/CodeSet
complexes were immobilized on nCounter cartridges at the nCounter® Prep Station 5s, and data collection
(RCC files) and quality check was performed in the nCounter® Digital Analyzer 5s. The nSolver™ Analysis and
Partek® Genomics Suite® software was used for analysis.

Kinase activity profiling microarray. Primary monocytes and platelets were isolated from 6 healthy donors.
Monocytes were stimulated *ex vitro* before or after the removal of platelets or after the re-addition of platelets to
PdMo (biological replicates with n = 6). As before, platelets alone (n = 6) were assessed. Stimulated cultures
were lysed in M-PER mammalian protein extraction reagent supplemented with PhosphoSTOP (Roche) and
cOmplete Tablets (Roche) incubated on ice for 15 minutes. After centrifugation at 20.000x g for 15 minutes at
4 C, the protein concentration in the supernatant is evaluated via Bradford assay. The lysate is aliquoted, snap-
frozen, and stored at -80 C. Only unthawed aliquots are used for the kinase activity assay.

Ser/Thr Kinase (STK) activity profiling assays based on measuring peptide phosphorylation by protein

kinases (PamGene International BV, The Netherlands) were performed as instructed by the manufacturer. In

summary, samples with 1 µg protein were applied on PamChip 4 arrays containing 144 (STK) or 196 (PTK)

peptides immobilized on a porous aluminum oxide membrane. The peptide sequences (13 amino acids long)

harbor phosphorylation sites and are correlated with one or multiple upstream kinases. Fluorescently labeled

anti-phospho antibodies detect phosphorylation activity of kinases present in the sample. Instrument operation

and imaging are controlled by the EVOLVE 2.0 software and quantified using BioNavigator 6.3 (BN6; PamGene

International BV, The Netherlands). Signal intensities at multiple exposure times were integrated by linear

regression (S100), Log2-transformed, and normalized using a Combat correction model for batch correction

where the scaling parameters (mean / sd) are estimated using an empirical Bayes approach, which effectively

reduces the noise associated with applying the correction.

The normalized values were used to perform statistics comparing groups or the upstream kinase analysis

(UKA) tool (BN6; PamGene International BV). The following statistical test is used to generate a list of

differentially significant phosphorylated peptides: Paired T-test for LPS effect/ comparisons and ANOVA-

PostHoc Test for multiple treatments versus control for comparing each group to StdMo unstimulated in both

unstimulated and LPS-15 min conditions. For Kinase interpretation (differential): To generate a ranked list of

putative kinases responsible for differences in the peptide phosphorylation, PamGene's in-house method called

Upstream Kinase Analysis (UKA) was used. For Pathway interpretation (differential): To generate a ranked list

of possible canonical pathways (and networks) responsible for differences in the peptide phosphorylation, there

are many open-source tools to perform this.

The phylogenetic kinome tree, applicable to group the kinases into sequence families, is plotted using the

online portal CORAL: http://phanstiel-lab.med.unc.edu/CORAL/ (12). The upstream kinase analysis functional

scoring tool (PamGene International) rank-orders the top kinases differential between the two groups, the

ranking factor being the final (median) kinase score (represented by node size). This score is based on a

combined sensitivity score (difference between treatment and control groups, expressed as node color) and

specificity score for a set of peptides to kinase relationships derived from existing databases. An arbitrary

threshold of a final score 1.2 was applied based on the specificity scores. Significant peptides (t-tests, p-value <

0.05) or kinases (UKA, final scores > 1.2) were imported to the MetaCore pathway analysis tool (Clarivate

Analytics), where an enrichment analysis was performed for pathways and networks. It matches the kinases or

substrates in the kinome array data with functional ontologies in MetaCore. The probability of a random

intersection between a set of IDs in the target list with ontology entities is estimated in the p value of

hypergeometric intersection. The lower p-value means higher relevance of the entity to the dataset, which

shows a higher rating for the entity. Direct interaction network algorithms were used to build interconnected

networks within each comparison. The "Add interactions" feature was used to add the interaction between
RIPK1 and the data in the MetaCore™ database after it was built.

Statistical Analysis

Statistical analyses were performed with GraphPad Prism Version 9.0f. Unless indicated otherwise, all graphs
are built from pooled data from a minimum of two independent experiments (biological replicates) performed in
triplicates (technical replicates). All statistical analyses were preceded by normality and lognormality tests,
followed by the recommended parametric or nonparametric tests. For most experiments with several groups
(StdMo, PdMo, PdMo + Ptls, and Plts, stimulated vs. unstimulated), P values were determined by two-way
ANOVA with Tukey's or Sidak's or multiple comparison tests. No outliers were detected or removed from the
analysis. Additional statistical details are given in the respective Fig legends when appropriate.

Data sharing availability

All datasets that support this manuscript will be made fully available upon publication. Raw data for RNA-seq
datasets are being deposited into public databases, and we will share with reviewers the assessment number
upon request. Additional requests to assess raw data and transcriptomic datasets before publication should be
made via e-mail to the corresponding author, along with an analysis proposal.

Data Presentation

Unless indicated otherwise (in Fig legends), all graphs are represented as Floating Bars (with mean and
minimum to maximum values) and are built from pooled data from a minimum of two independent experiments
(biological replicates) performed in triplicates (technical replicates) with monocytes or platelets from different
donors. Each symbol represents the average from 3 technical replicates per donor or experiment in the case of
cell lines. Symbols are coded (●, ▼, □□, ●, etc.) to indicate donors so readers can track the internal variability
between different donors or experiments. Dots are semi-transparent, with darker symbols showing overlapping
points.

FUNDING

This study was funded by the European Research Council (EC | H2020 | H2020 Priority Excellent Science |
H2020 European Research Council (ERC) PLAT-IL-1, 714175). Bernardo S Franklin is further supported by the
HORIZON Lump Sum Grant (ERC PoC UNBIAS, 101123144), (BEATSep, 101137484), as well as Germany's

Excellence Strategy (EXC 2151 – 390873048, SFB TRR259) from the Deutsche Forschungsgemeinschaft (DFG,
German Research Foundation).

ACKNOWLEDGMENTS

We thank Marco A. Ataide, Florian I Schmidt, and Dagmar Wachten for scientific discussions and input and
for proofreading the manuscript. We thank Cornelia Rohland and Maximilian Rothe for technical help and
organization and maintenance of mouse lines, respectively. We thank Jonathan Schmid-Burgk for their advice
and support in generating THP1 KO cells using CRISPR-Cas9. We thank Dr. Savithri Rangarajan and Dr. Rik de
Wijn (PamGene, Diagnostic Assay Services, 's-Hertogenbosch, The Netherlands) for performing the peptide
quality control and bioinformatics analysis of the Protein Kinase Assay. We thank the Microscopy and Flow
Cytometry Core Facility of the Medical Faculty at the University of Bonn for providing help, services, and
devices funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, project number
388158066, 216372545 and 388159768). Finally, we thank Andrea Schlichting and Leonie Verwohlt for
administrative support.

AUTHOR CONTRIBUTIONS

Ibrahim Hawwari: Investigation; Formal analysis; Data curation; Visualization; Methodology; Writing—review
and editing. **Lukas Rosnagel:** Investigation; Formal analysis; Data curation; Visualization; Methodology;
Writing—review and editing. **Nathalia Sofia Rosero Reyes:** Investigation; Technical Support. **Lino L
Teichmann** and **Lisa Meffert**: ITP patient samples. **Agnieszka Demczuk:** Investigation. **Lucas S. Ribeiro:**
Technical Support. **Damien Bertheloot:** Supervision; Investigation. **Magali Noval Rivas**: Investigation. **Moshe
Arditi:** Formal analysis; Data curation. **Sebastian Kallabis:** Investigation; Formal analysis; Data curation. **Felix
Meissner:** Formal analysis; Data curation. **Bernardo S Franklin:** Conceptualization; Funding acquisition;
Project administration; Resources; Formal analysis; Supervision; Validation; Investigation; Visualization;
Writing—original draft.

DISCLOSURE AND COMPETING INTERESTS STATEMENT

All authors declare that they have no conflict of interest.

736
REFERENCES

Abhilasha KV, Sumanth MS, Chaithra VH, Jacob SP, Thyagarajan A, Sahu RP, Rajaiah R, Prabhu KS, Kemparaju K, Travers JB
et al (2019) p38 MAP-kinase inhibitor protects against platelet-activating factor-induced death in mice. Free Radic Biol
Med 143: 275-287

Ablasser A, Schmid-Burgk JL, Hemmerling I, Horvath GL, Schmidt T, Latz E, Hornung V (2013) Cell intrinsic immunity spreads to
bystander cells via the intercellular transfer of cGAMP. Nature 503: 530-534

Agrati C, Sacchi A, Bordoni V, Cimini E, Notari S, Grassi G, Casetti R, Tartaglia E, Lalle E, D'Abramo A et al (2020) Expansion of
myeloid-derived suppressor cells in patients with severe coronavirus disease (COVID-19). Cell Death Differ 27: 3196-
3207

Alard J-E, Ortega-Gomez A, Wichapong K, Bongiovanni D, Horckmans M, Megens RTA, Leoni G, Ferraro B, Rossaint J, Paulin N
et al (2015) Recruitment of classical monocytes can be inhibited by disturbing heteromers of neutrophil HNP1 and
platelet CCL5. Sci Transl Med 7: 317ra196-317ra196

Alasoo K, Martinez FO, Hale C, Gordon S, Powrie F, Dougan G, Mukhopadhyay S, Gaffney DJ (2015) Transcriptional profiling of
macrophages derived from monocytes and iPS cells identifies a conserved response to LPS and novel alternative
transcription. Sci Rep 5: 12524

Allen N, Barrett TJ, Guo Y, Nardi M, Ramkhelawon B, Rockman CB, Hochman JS, Berger JS (2019) Circulating monocyte-
platelet aggregates are a robust marker of platelet activity in cardiovascular disease. Atherosclerosis 282: 11-18

Anand P, Shenoy R, Palmer JE, Baines AJ, Lai RY, Robertson J, Bird N, Ostenfeld T, Chizh BA (2011) Clinical trial of the p38
MAP kinase inhibitor dilmapimod in neuropathic pain following nerve injury. Eur J Pain 15: 1040-1048

Anderson P (2010) Post-transcriptional regulons coordinate the initiation and resolution of inflammation. Nat Rev Immunol 10: 24-
35

Andreescu M (2023) The link between immune thrombocytopenia and the cytokine profile: a bridge to new therapeutical targets.
Frontiers in Hematology 2

Arens C, Bajwa SA, Koch C, Siegler BH, Schneck E, Hecker A, Weiterer S, Lichtenstern C, Weigand MA, Uhle F (2016) Sepsis-
induced long-term immune paralysis--results of a descriptive, explorative study. Crit Care 20: 93

Arunachalam PS, Wimmers F, Mok CKP, Perera R, Scott M, Hagan T, Sigal N, Feng Y, Bristow L, Tak-Yin Tsang O et al (2020)
Systems biological assessment of immunity to mild versus severe COVID-19 infection in humans. Science 369: 1210-
1220

Barrett TJ, Schlegel M, Zhou F, Gorenchtein M, Bolstorff J, Moore KJ, Fisher EA, Berger JS (2019) Platelet regulation of myeloid
suppressor of cytokine signaling 3 accelerates atherosclerosis. Sci Transl Med 11

Barry OP, Pratico D, Lawson JA, FitzGerald GA (1997) Transcellular activation of platelets and endothelial cells by bioactive
lipids in platelet microparticles. J Clin Invest 99: 2118-2127

Beaulieu LM, Freedman JE (2009) NFkappaB regulation of platelet function: no nucleus, no genes, no problem? J Thromb
Haemost 7: 1329-1332

Bekkering S, Quintin J, Joosten LA, van der Meer JW, Netea MG, Riksen NP (2014) Oxidized low-density lipoprotein induces
long-term proinflammatory cytokine production and foam cell formation via epigenetic reprogramming of monocytes.
Arterioscler Thromb Vasc Biol 34: 1731-1738

Bennett JS, Berger BW, Billings PC (2009) The structure and function of platelet integrins. J Thromb Haemost 7 Suppl 1: 200-205

Bhattacharjee J, Das B, Mishra A, Sahay P, Upadhyay P (2017) Monocytes isolated by positive and negative magnetic sorting
techniques show different molecular characteristics and immunophenotypic behaviour. F1000Res 6: 2045

Birnie E, Claushuis TAM, Koh G, Limmathurotsakul D, Day NPJ, Roelofs J, Ware J, Hou B, de Vos AF, van der Poll T et al (2019)
Thrombocytopenia Impairs Host Defense Against Burkholderia pseudomallei (Melioidosis). J Infect Dis 219: 648-659

Bonnet B, Cosme J, Dupuis C, Coupez E, Adda M, Calvet L, Fabre L, Saint-Sardos P, Bereiziat M, Vidal M et al (2021) Severe
COVID-19 is characterized by the co-occurrence of moderate cytokine inflammation and severe monocyte
dysregulation. EBioMedicine 73: 103622

Cao C, Yu M, Chai Y (2019) Pathological alteration and therapeutic implications of sepsis-induced immune cell apoptosis. Cell
Death Dis 10: 782

Carestia A, Mena HA, Olexen CM, Ortiz Wilczyñski JM, Negrotto S, Errasti AE, Gómez RM, Jenne CN, Carrera Silva EA,
Schattner M (2019) Platelets Promote Macrophage Polarization toward Pro-inflammatory Phenotype and Increase
Survival of Septic Mice. CellReports 28: 896-908.e895

Chatterjee M, von Ungern-Sternberg SNI, Seizer P, Schlegel F, Büttcher M, Sindhu NA, Müller S, Mack A, Gawaz M (2015)
Platelet-derived CXCL12 regulates monocyte function, survival, differentiation into macrophages and foam cells
through differential involvement of CXCR4–CXCR7. Cell Death and Disease 6: e1989-e1989

Christie JD, Vaslef S, Chang PK, May AK, Gunn SR, Yang S, Hardes K, Kahl L, Powley WM, Lipson DA et al (2015) A
Randomized Dose-Escalation Study of the Safety and Anti-Inflammatory Activity of the p38 Mitogen-Activated Protein
Kinase Inhibitor Dilmapimod in Severe Trauma Subjects at Risk for Acute Respiratory Distress Syndrome. Crit Care
Med 43: 1859-1869

Claushuis TAM, van Vught LA, Scicluna BP, Wiewel MA, Klouwenberg PMCK, Hoogendijk AJ, Ong DSY, Cremer OL, Horn J,
Franitza M et al (2016) Thrombocytopenia is associated with a dysregulated host response in critically ill sepsis
patients. Blood 127: 3062-3072

Cooper N, Ghanima W (2019) Immune Thrombocytopenia. N Engl J Med 381: 945-955

D'Mello C, Almishri W, Liu H, Swain MG (2017) Interactions Between Platelets and Inflammatory Monocytes Affect Sickness
Behavior in Mice With Liver Inflammation. Gastroenterology 153: 1416-1428.e1412

de Stoppelaar SF, van 't Veer C, Claushuis TA, Albersen BJ, Roelofs JJ, van der Poll T (2014) Thrombocytopenia impairs host
defense in gram-negative pneumonia-derived sepsis in mice. Blood 124: 3781-3790

Ezumi Y, Takayama H, Okuma M (1995) Thrombopoietin, c-Mpl ligand, induces tyrosine phosphorylation of Tyk2, JAK2, and
STAT3, and enhances agonists-induced aggregation in platelets in vitro. FEBS Lett 374: 48-52

Fajgenbaum DC, June CH (2020) Cytokine Storm. N Engl J Med 383: 2255-2273

Ferreira AC, Soares VC, de Azevedo-Quintanilha IG, Dias SdSG, Fintelman-Rodrigues N, Sacramento CQ, Mattos M, de Freitas
CS, Temerozo JR, Teixeira L et al (2021) SARS-CoV-2 engages inflammasome and pyroptosis in human primary
monocytes. Cell Death Discov 7: 43

Forlow SB, McEver RP, Nollert MU (2000) Leukocyte-leukocyte interactions mediated by platelet microparticles under flow. Blood
95: 1317-1323

Frazier WJ, Hall MW (2008) Immunoparalysis and adverse outcomes from critical illness. Pediatr Clin North Am 55: 647-668, xi

Frenette PS, Denis CV, Weiss L, Jurk K, Subbarao S, Kehrel B, Hartwig JH, Vestweber D, Wagner DD (2000) P-Selectin
glycoprotein ligand 1 (PSGL-1) is expressed on platelets and can mediate platelet-endothelial interactions in vivo. J
Exp Med 191: 1413-1422

Fu G, Deng M, Neal MD, Billiar TR, Scott MJ (2021) Platelet-Monocyte Aggregates: Understanding Mechanisms and Functions in
Sepsis. Shock 55: 156-166

Gottschalk RA, Martins AJ, Angermann BR, Dutta B, Ng CE, Uderhardt S, Tsang JS, Fraser ID, Meier-Schellersheim M, Germain
RN (2016) Distinct NF-kappaB and MAPK Activation Thresholds Uncouple Steady-State Microbe Sensing from Anti-
pathogen Inflammatory Responses. Cell Syst 2: 378-390

Guilliams M, Mildner A, Yona S (2018) Developmental and Functional Heterogeneity of Monocytes. Immunity 49: 595-613

Guma M, Stepniak D, Shaked H, Spehlmann ME, Shenouda S, Cheroutre H, Vicente-Suarez I, Eckmann L, Kagnoff MF, Karin M
(2011) Constitutive intestinal NF-kappaB does not trigger destructive inflammation unless accompanied by MAPK
activation. J Exp Med 208: 1889-1900

Han P, Hanlon D, Arshad N, Lee JS, Tatsuno K, Robinson E, Filler R, Sobolev O, Cote C, Rivera-Molina F et al (2020) Platelet P-
selectin initiates cross-presentation and dendritic cell differentiation in blood monocytes. Sci Adv 6: eaaz1580

Haskel EJ, Abendschein DR (1989) Deaggregation of human platelets in vitro by an RGD analog antagonist of platelet
glycoprotein IIb/IIIa receptors. Thromb Res 56: 687-695

Haupt W, Riese J, Mehler C, Weber K, Zowe M, Hohenberger W (1998) Monocyte function before and after surgical trauma. Dig
Surg 15: 102-104

Henn V, Slupsky JR, Gräfe M, Nature IA, 1998 CD40 ligand on activated platelets triggers an inflammatory reaction of endothelial
cells. naturecom

Inwald DP, McDowall A, Peters MJ, Callard RE, Klein NJ (2003) CD40 is constitutively expressed on platelets and provides a
novel mechanism for platelet activation. Circ Res 92: 1041-1048

Jafarzadeh A, Chauhan P, Saha B, Jafarzadeh S, Nemati M (2020) Contribution of monocytes and macrophages to the local
tissue inflammation and cytokine storm in COVID-19: Lessons from SARS and MERS, and potential therapeutic
interventions. Life Sci 257: 118102

Junqueira C, Crespo Â, Ranjbar S, de Lacerda LB, Lewandrowski M, Ingber J, Parry B, Ravid S, Clark S, Schrimpf MR et al
(2022) FcγR-mediated SARS-CoV-2 infection of monocytes activates inflammation. Nature

Kasper CA, Sorg I, Schmutz C, Tschon T, Wischnewski H, Kim ML, Arrieumerlou C (2010) Cell-Cell Propagation of NF-κB
Transcription Factor and MAP Kinase Activation Amplifies Innate Immunity against Bacterial Infection. Immunity 33:
804-816

Kral JB, Schrottmaier WC, Salzmann M, Assinger A (2016) Platelet Interaction with Innate Immune Cells. Transfus Med
Hemother 43: 78-88

Kullaya V, de Jonge MI, Langereis JD, van der Gaast-de Jongh CE, Bull C, Adema GJ, Lefeber D, Cremers AJ, Mmbaga BT, de
Groot PG et al (2018) Desialylation of Platelets by Pneumococcal Neuraminidase A Induces ADP-Dependent Platelet
Hyperreactivity. Infect Immun 86

Lang D, Dohle F, Terstesse M, Bangen P, August C, Pauels HG, Heidenreich S (2002) Down-regulation of monocyte apoptosis
by phagocytosis of platelets: involvement of a caspase-9, caspase-3, and heat shock protein 70-dependent pathway.
J Immunol 168: 6152-6158

Lannan KL, Sahler J, Kim N, Spinelli SL, Maggirwar SB, Garraud O, Cognasse F, Blumberg N, Phipps RP (2015) Breaking the
mold: transcription factors in the anucleate platelet and platelet-derived microparticles. Front Immunol 6: 48

Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Keys JR, Land vatter SW et al
(1994) A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. Nature 372: 739-746

Levoux J, Prola A, Lafuste P, Gervais M, Chevallier N, Koumaiha Z, Kefi K, Braud L, Schmitt A, Yacia A et al (2021) Platelets
Facilitate the Wound-Healing Capability of Mesenchymal Stem Cells by Mitochondrial Transfer and Metabolic
Reprogramming. Cell Metab 33: 283-299 e289

Liang H, Duan Z, Li D, Li D, Wang Z, Ren L, Shen T, Shao Y (2015) Higher levels of circulating monocyte-platelet aggregates are
correlated with viremia and increased sCD163 levels in HIV-1 infection. Cell Mol Immunol 12: 435-443

Lin CY, Tsai IF, Ho YP, Huang CT, Lin YC, Lin CJ, Tseng SC, Lin WP, Chen WT, Sheen IS (2007) Endotoxemia contributes to
the immune paralysis in patients with cirrhosis. J Hepatol 46: 816-826

Malehmir M, Pfister D, Gallage S, Szydlowska M, Inverso D, Kotsiliti E, Leone V, Peiseler M, Surewaard BGJ, Rath D et al (2019)
Platelet GPIbalpha is a mediator and potential interventional target for NASH and subsequent liver cancer. Nat Med
25: 641-655

Manne BK, Denorme F, Middleton EA, Portier I, Rowley JW, Stubben C, Petrey AC, Tolley ND, Guo L, Cody M et al (2020)
Platelet gene expression and function in patients with COVID-19. Blood 136: 1317-1329

Maugeri N, Rovere-Querini P, Evangelista V, Covino C, Capobianco A, Bertilaccio MT, Piccoli A, Totani L, Cianflone D, Maseri A,
Manfredi AA (2009) Neutrophils phagocytose activated platelets in vivo: a phosphatidylserine, P-selectin, and {beta}2
integrin-dependent cell clearance program. Blood 113: 5254-5265

Netea MG, Joosten LA, Latz E, Mills KH, Natoli G, Stunnenberg HG, O'Neill LA, Xavier RJ (2016) Trained immunity: A program of
innate immune memory in health and disease. Science 352: aaf1098

Nieswandt B, Bergmeier W, Rackebrandt K, Gessner JE, Zirngibl H (2000) Identification of critical antigen-specific mechanisms in
the development of immune thrombocytopenic purpura in mice. Blood 96: 2520-2527

Pachot A, Cazalis MA, Venet F, Turrel F, Faudot C, Voirin N, Diasparra J, Bourgoin N, Poitevin F, Mougin B et al (2008)
Decreased expression of the fractalkine receptor CX3CR1 on circulating monocytes as new feature of sepsis-induced
immunosuppression. J Immunol 180: 6421-6429

Poli V, Di Gioia M, Sola-Visner M, Granucci F, Frelinger AL, 3rd, Michelson AD, Zanoni I (2021) Inhibition of transcription factor
NFAT activity in activated platelets enhances their aggregation and exacerbates gram-negative bacterial septicemia.
Immunity

Qu M, Liu Q, Zhao HG, Peng J, Ni H, Hou M, Jansen AJG (2018) Low platelet count as risk factor for infections in patients with
primary immune thrombocytopenia: a retrospective evaluation. Ann Hematol 97: 1701-1706

Rauert-Wunderlich H, Siegmund D, Maier E, Giner T, Bargou RC, Wajant H, Stuhmer T (2013) The IKK inhibitor Bay 11-7082
induces cell death independent from inhibition of activation of NFkappaB transcription factors. PLoS ONE 8: e59292

Rinder HM, Bonan JL, Rinder CS, Ault KA, Smith BR (1991) Dynamics of leukocyte-platelet adhesion in whole blood. Blood 78:
1730-1737

Rodrigues TS, de Sa KSG, Ishimoto AY, Becerra A, Oliveira S, Almeida L, Goncalves AV, Perucello DB, Andrade WA, Castro R
et al (2021) Inflammasomes are activated in response to SARS-CoV-2 infection and are associated with COVID-19
severity in patients. J Exp Med 218

Rolfes V, Ribeiro LS, Hawwari I, Bottcher L, Rosero N, Maasewerd S, Santos MLS, Prochnicki T, Silva CMS, Wanderley CWS et
al (2020) Platelets Fuel the Inflammasome Activation of Innate Immune Cells. Cell Rep 31: 107615

Rong MY, Wang CH, Wu ZB, Zeng W, Zheng ZH, Han Q, Jia JF, Li XY, Zhu P (2014) Platelets induce a proinflammatory
phenotype in monocytes via the CD147 pathway in rheumatoid arthritis. Arthritis Res Ther 16: 478

Roquilly A, Jacqueline C, Davieau M, Molle A, Sadek A, Fourgeux C, Rooze P, Broquet A, Misme-Aucouturier B, Chaumette T et
al (2020) Alveolar macrophages are epigenetically altered after inflammation, leading to long-term lung
immunoparalysis. Nat Immunol 21: 636-648

Rossaint J, Kühne K, Skupski J, Van Aken H, Looney MR, Hidalgo A, Zarbock A (2016) Directed transport of neutrophil-derived
extracellular vesicles enables platelet-mediated innate immune response. Nature Communications 7: 13464

Rozmyslowicz T, Majka M, Kijowski J, Murphy SL, Conover DO, Poncz M, Ratajczak J, Gaulton GN, Ratajczak MZ (2003)
897   Platelet-and megakaryocyte-derived microparticles transfer CXCR4 receptor to CXCR4-null cells and make them
898   susceptible to infection by X4-HIV. AIDS 17: 33-42

899 Saha RN, Jana M, Pahan K (2007) MAPK p38 regulates transcriptional activity of NF-kappaB in primary human astrocytes via
900   acetylation of p65. J Immunol 179: 7101-7109

901 Schulte-Schrepping J, Reusch N, Paclik D, Bassler K, Schlickeiser S, Zhang B, Kramer B, Krammer T, Brumhard S, Bonaguro L
902   et al (2020) Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell Compartment. Cell 182: 1419-1440 e1423

903 Senzel L, Chang C (2013) Platelet phagocytosis by neutrophils. Blood 122: 1543

904 Shi P, Zhang L, Zhang M, Yang W, Wang K, Zhang J, Otsu K, Huang G, Fan X, Liu J (2017) Platelet-Specific p38alpha
905   Deficiency Improved Cardiac Function After Myocardial Infarction in Mice. Arterioscler Thromb Vasc Biol 37: e185-
906   e196

907 Singhal R, Chawla S, Rathore DK, Bhasym A, Annarapu GK, Sharma V, Seth T, Guchhait P (2017) Development of pro-
908   inflammatory phenotype in monocytes after engulfing Hb-activated platelets in hemolytic disorders. Clinical
909   Immunology 175: 133-142

910 Solt LA, May MJ (2008) The IkappaB kinase complex: master regulator of NF-kappaB signaling. Immunol Res 42: 3-18

911 Spinelli SL, Casey AE, Pollock SJ, Gertz JM, McMillan DH, Narasipura SD, Mody NA, King MR, Maggirwar SB, Francis CW et al
912   (2010) Platelets and megakaryocytes contain functional nuclear factor-kappaB. Arterioscler Thromb Vasc Biol 30:
913   591-598

914 Sreeramkumar V, Adrover JM, Ballesteros I, Cuartero MI, Rossaint J, Bilbao I, Nacher M, Pitaval C, Radovanovic I, Fukui Y et al
915   (2014) Neutrophils scan for activated platelets to initiate inflammation. Science 346: 1234-1238

916 Stephen J, Emerson B, Fox KA, Dransfield I (2013) The uncoupling of monocyte-platelet interactions from the induction of
917   proinflammatory signaling in monocytes. J Immunol 191: 5677-5683

918 Thomas MR, Outteridge SN, Ajjan RA, Phoenix F, Sangha GK, Faulkner RE, Ecob R, Judge HM, Khan H, West LE et al (2015)
919   Platelet P2Y12 Inhibitors Reduce Systemic Inflammation and Its Prothrombotic Effects in an Experimental Human
920   Model. Arterioscler Thromb Vasc Biol 35: 2562-2570

921 Ueda A, Okuda K, Ohno S, Shirai A, Igarashi T, Matsunaga K, Fukushima J, Kawamoto S, Ishigatsubo Y, Okubo T (1994) NF-
922   kappa B and Sp1 regulate transcription of the human monocyte chemoattractant protein-1 gene. J Immunol 153:
923   2052-2063

924 Vajen T, Mause SF, Koenen RR (2015) Microvesicles from platelets: novel drivers of vascular inflammation. Thromb Haemost
925   114: 228-236

926 van den Boogaard FE, Schouten M, de Stoppelaar SF, Roelofs JJ, Brands X, Schultz MJ, van't Veer C, van der Poll T (2015)
927   Thrombocytopenia impairs host defense during murine Streptococcus pneumoniae pneumonia. Crit Care Med 43:
928   e75-83

929 Vanderbeke L, Van Mol P, Van Herck Y, De Smet F, Humblet-Baron S, Martinod K, Antoranz A, Arijs I, Boeckx B, Bosisio FM et
930   al (2021) Monocyte-driven atypical cytokine storm and aberrant neutrophil activation as key mediators of COVID-19
931   disease severity. Nat Commun 12: 4117

932 Vangeti S, Strandin T, Liu S, Tauriainen J, Raisanen-Sokolowski A, Cabrera L, Hassinen A, Makela S, Mustonen J, Vaheri A et al
933   (2021) Monocyte subset redistribution from blood to kidneys in patients with Puumala virus caused hemorrhagic fever
934   with renal syndrome. PLoS Pathog 17: e1009400

Varchetta S, Mele D, Lombardi A, Oliviero B, Mantovani S, Tinelli C, Spreafico M, Prati D, Ludovisi S, Ferraioli G et al (2016)
Lack of Siglec-7 expression identifies a dysfunctional natural killer cell subset associated with liver inflammation and
fibrosis in chronic HCV infection. Gut 65: 1998-2006

Wasmuth HE, Kunz D, Yagmur E, Timmer-Stranghoner A, Vidacek D, Siewert E, Bach J, Geier A, Purucker EA, Gressner AM et
al (2005) Patients with acute on chronic liver failure display "sepsis-like" immune paralysis. J Hepatol 42: 195-201

Weisheit CK, Kluners A, Wild L, Casalter A, Heilmann-Heimbach S, Sivalingam S, Kleiner JL, Ehrentraut SF, Hoeft A, Frede S,
Ehrentraut H (2020) Sustained Immunoparalysis in Endotoxin-Tolerized Monocytic Cells. Mediators Inflamm 2020:
8294342

Weyrich AS, Elstad MR, McEver RP, McIntyre TM, Moore KL, Morrissey JH, Prescott SM, Zimmerman GA (1996) Activated
platelets signal chemokine synthesis by human monocytes. J Clin Invest 97: 1525-1534

Weyrich AS, McIntyre TM, McEver RP, Prescott SM, Zimmerman GA (1995) Monocyte tethering by P-selectin regulates
monocyte chemotactic protein-1 and tumor necrosis factor-alpha secretion. Signal integration and NF-kappa B
translocation. J Clin Invest 95: 2297-2303

White DE, Burchill SA (2008) BAY 11-7082 induces cell death through NF-kappaB-independent mechanisms in the Ewing's
sarcoma family of tumours. Cancer Lett 268: 212-224

Wolf T, Jin W, Zoppi G, Vogel IA, Akhmedov M, Bleck CKE, Beltraminelli T, Rieckmann JC, Ramirez NJ, Benevento M et al
(2020) Dynamics in protein translation sustaining T cell preparedness. Nat Immunol 21: 927-937

Wong KL, Tai JJ, Wong WC, Han H, Sem X, Yeap WH, Kourilsky P, Wong SC (2011) Gene expression profiling reveals the
defining features of the classical, intermediate, and nonclassical human monocyte subsets. Blood 118: e16-31

Xiang B, Zhang G, Guo L, Li X-A, Morris AJ, Daugherty A, Whiteheart SW, Smyth SS, Li Z (2013) Platelets protect from septic
shock by inhibiting macrophage-dependent inflammation via the cyclooxygenase 1 signalling pathway. Nature
Communications 4: 264

Yamashita Y, Hishinuma M, Shimada M (2009) Activation of PKA, p38 MAPK and ERK1/2 by gonadotropins in cumulus cells is
critical for induction of EGF-like factor and TACE/ADAM17 gene expression during in vitro maturation of porcine
COCs. J Ovarian Res 2: 20

Yong HY, Koh MS, Moon A (2009) The p38 MAPK inhibitors for the treatment of inflammatory diseases and cancer. Expert Opin
Investig Drugs 18: 1893-1905

Figure Legends

Fig 1. Platelets regulate the cytokine secretion of human monocytes.

(A) Concentrations of IL-1β, TNFα, and IL-6 released into the cell-free supernatants (CFS) by untouched (StdMo, light blue), platelet-depleted (PdMo, blue), or PdMo that were supplemented with autologous platelets (PdMo + Plts, dark blue) at a 100:1 platelet:monocyte ratio. Cytokine levels secreted by platelets alone (Plts, purple) were measured as control. Cells were stimulated with LPS (2 ng ml⁻¹ for 3h) followed by activation with nigericin (10 µM for 1.5 h).

(B) Intracellular maturation and secretion of IL-1β assessed by WES capillary electrophoresis coupled with Ab-based detection in whole cell lysates (WCL) or CFS of primary human monocytes stimulated as in **A**.

(C) Cell viability assessed every 15 min in StdMo, PdMo, PdMo + Plts, and Plts treated as in **A**.

(D) Fluorometric assessment of caspase-1 activity in the CFS of primary human monocytes stimulated as in **A**.

(E - F) IL-1β concentrations in CFS of LPS-primed StdMo, PdMo, or PdMo + Plts that were transfected with Lipofectamin (LF) containing (E) double-strand DNA (dsDNA, 0.5 µg ml−1) for the activation of the AIM2 inflammasome. In (F), cells were treated with PrgI (100 ng ml⁻¹) and protective antigen (PA, 1 µg ml⁻¹) for 90 min, for the activation of the NLRC4 inflammasome. Floating bars display the max/min values with indication for the mean (white bands). Each symbol represents one independent experiment or blood donor.

(G) Radar plots displaying the 19 significantly (p < 0.05, 2-Way Anova, Tukey's multiple comparison test) altered cytokines detected by Cytokine Luminex in the CFS of human monocytes and platelets stimulated as in **A**. Protein concentrations are represented by the spread from inner (0 ng ml⁻¹) to outer circles (≥ 10 ng ml⁻¹). Colors represent stimuli (Grey: Unstim, Red: LPS, dark blue: LPS + Nig). Each symbol represents one independent experiment or blood donor (n = 5 for monocytes, and n = 2 for platelets alone). For better visualisation, cytokine concentrations (y axis) are displayed with a cut off at 8 ng ml⁻¹. Levels of IL-1RA and CCL4 reached 20 ng ml⁻¹. **See also Fig S1 and EV2.**

Fig 2. Platelet supplementation reverts monocyte immunoparalysis in Immune Thrombocytopenia (ITP).

(A) Demographic and clinical characteristics of ITP patients (n = 5).

(B) Representative flow cytometry percentages and (C) CASY automated cell quantification of monocytes and platelets in preparations of untouched or platelet-depleted monocytes from healthy donors (± platelet-depletion) or untouched monocytes from ITP patients (ITPMo).

(D) Clinical laboratory quantification of platelets in the peripheral blood of ITP patients. Blue shaded area display the healthy reference ranges.

(E-G) Concentrations of IL-1β, IL-6 and TNFα released into the CFS by untouched (StdMo), and PdMo isolated from healthy volunteers (n = 4), or untouched monocytes from ITP patients (ITPMo, n = 5). Healthy platelet-depleted monocytes (PdMo + Plts), or ITPMo were supplemented with platelets (ITPMo + Plts). Cells were stimulated with LPS (2 ng ml⁻¹) for IL-6 and TNFα measurements, or LPS + Nig (10 µM) for IL-1β measurements. Data is displayed as floating bars with the max/min values and mean (white bands). P values were calculated with 2-Way Anova, Tukey's multiple comparison test, and are displayed in the Fig. Each symbol represents one independent experiment or blood donor.

Fig 3. Platelet depletion causes transcriptional shutdown of myeloid innate immune genes in human monocytes.

(A) Principal Component Analyses (PCA) of gene expression in StdMo (green), platelet-depleted monocytes (PdMo, red), platelet-depleted monocytes reconstituted with platelets (PdMo + Plts, blue), and platelets alone (Plts, pink). Each symbol represents one donor. Stimulations are represented with shapes (elypse = unstimulated, square = LPS).

(B) Volcano plots showing log2 fold change (x-axis) and significance (−log10 *p-value; y-axis) of genes differentially expressed
comparing LPS-stimulated vs. unstimulated (Unstim) StdMo, PdMo, or PdMo + Plts.

(C) Pie charts indicate the number of up- and downregulated DEGs in each group upon LPS stimulation.

(D) Quantitative expression (Log2 Fold Change) of the LPS-induced (>= 2 Fold Change) genes in StdMo (green), PdMo (red)
or PdMo + Plts (blue).

(E) Scatter plots displaying the LSMean (x-axis) of StdMo vs PdMo (y-axis) from DEGs comparing the effects of platelet
removal in Unstim and LPS-stimulated conditions. DEGs with 2-Fold change are highlighted in red.

(F) Pie charts representing the DEGs of the comparisons in **D**, representing DEGs induced by platelet depletion in
unstimulated or LPS-stimulated monocytes. See also Fig S**3**.

1012

Fig 4. Dysregulation of host defense transcriptional programs in monocytes from thrombopenic mice.

(A) Schematic representation of the experimental setting for platelet depletion in vivo depletion followed by intravenous
challenge with LPS, FACS-sorting and bulk-RNASeq of blood monocytes. Monocytes were isolated from PBMCs from
pooled blood of 4 mice per group (IgG- or anti-GPIbα-treated) each challenged either with LPS or PBS in 3 different
experiments.

(B) Flow cytometric quantification of platelets in IgG-treated and anti-GPIba treated mice (n = 24 per condition).

(C) Volcano plots (log2 fold change vs. −log$_{10}$ adjusted P-value) of the significant DEGs (adjusted P-value < 0.05, Fold
Change ≤ -2 or ≥ 2) comparing PBS vs. LPS-challenge in IgG (left) or anti-GPIba-treated mice (right).

(D) Fold change of the DEGs that compose the myeloid innate immunity that were modulated by LPS in monocytes from IgG
vs. antiGPIbα-treated mice. P values from t.test are shown.

(E) Pathway enrichment analysis of the LPS-induced DEGs comparing monocytes from IgG (dark blue) or anti-GPIba-treated
mice (light blue).

(F) Main represented categories of pathways (GO analysis) from the DEGs induced by LPS on monocytes from IgG-treated
vs. platelet-depleted mice (See also Fig S**4**).

1027

Fig 5: Platelets modulate monocyte cytokine responses in trans.

(A) Confocal imaging of StdMo and platelets. Cells were stained with CD41a AF488 (platelets: green) and CD14 AF647
(monocytes: red). Nuclei were stained with Hoechst 34580 (blue). Scale: 24 μm (top panel); 4.8 μm (bottom panel). White
arrows indicate points of contacts between platelets and monocytes. Images are from one representative of four
independent experiments.

(B) IL-1β and TNFα levels in CFS of StdMo, PdMo, and PdMo that were supplemented with autologous platelets (PdMo + Plts,
100:1 platelets/ monocytes). Monocytes were pre-treated with Cytochalasin D (CytoD, 50 μM) or Dynasore (DS, 30 μM)
before being supplied with Plts. Cells were stimulated with LPS (2 ng ml-1), followed by activation with nigericin (10 μM).

(C) TNFα and IL-6 levels in CFS of LPS stimulated StdMo, PdMo, PdMo, or PdMo reconstituted with platelet releasates (Plt-
Sups).

(D) Representative flow cytometry assessment and gating strategy of StdMo that were incubated with growing concentrations
of KPL-1 (5, 10 or 25 μg ml-), or IgG (25 μg ml-). Gates indicate the frequencies of free platelets (CD45- CD41a+), MPAs
(CD45+CD41a+) and platelet-free monocytes (CD45+ CD41a-). Data is representative of two independent experiments
with several donors.

(E)   Cumulative flow cytometry frequency of free monocytes (CD45- CD41a-) in StdMo treated with growing concentrations of KPL-1 (5, 10 or 25 μg ml-1), a specific mAb against P-selectin, or an unrelated isotype IgG control (25 μg ml-1). See **Fig S5B** for gating strategy.

(F)   IL-1β, TNFα and IL-6 levels released by stimulated StdMo that were treated with increasing concentrations of KPL-1 (5, 10 or 25 μg ml-1), or IgG (25 μg ml-1).

Graphs display floating bars with the max/min values and mean (white bands) combined from multiple experiments. P values were calculated with 2-Way Anova, Tukey's multiple comparison test, and are displayed in the Fig. Each symbol represents one independent experiment or blood donor.

Fig 6. Platelets are cellular sources of active kinases and transcription factors.

(A-B)   Coral trees display the activity of Protein Tyrosine (upper branches) and Serine/Threonine kinases (lower branches) in StdMo, PdMo, PdMo + Plts and platelets from 5 donors, upon LPS stimulation for 15 min. (A) Impact of platelet depletion on the activity of PTK and STK in unstimulated monocytes. (B) Effects of LPS stimulation on the activation of PTKs and STKs in StdMo, PdMo, PdMo + Plts and platelets. Red: increased kinase activity, Blue: reduced kinase activity.

(C-D)   Scatter plots and **D** box plot showing the signal intensity for the phosphorylation status of phosphosites (x.axis) that are substracts of PTK (top) and STK (bottom) comparing StdMo, PdMo, PdMo + Plts, and platelets alone (Plts). Each symbol shows one donor (n = 5) and only include peptides which passed the QC. batch correction with a 2-step Combat correction. P values in D were calculated by Anova multiple comparison.

(E)   Pathway analysis of the most represented signaling pathways dynamically changed upon platelet depletion/re-addition to primary human monocytes. See also **Fig S6**.

(F)   Immunoblot of total RelA (p65), and Phosphorylated-p65 (P-p65) in MoStd, MoPD and Plts that were left untreated (−) or stimulated with LPS for 5, 15, 30 or 60 min. Representative of three independent experiments.

(G)   NFκB activity assay in CFS of unstimulated (Unstim), or Pam3CysK4-stimulated (100 ng ml-1) NFκB-SEAP and IRF-Lucia luciferase Reporter Monocytes cultured alone (−) or co-cultured with increasing concentrations of human platelets (1:10, 1:20, 1:50 and 1:100). SEAP-Activity was measured with QuantiBlue Buffer. Graphs show floating bars display max/min values with indication to the mean (white bands). Each symbol represents one donor, or independent experiment.

Fig 7: Enrichment of megakaryocyte-derived NFκB2 in human monocytes.

(A)   IL-1β released from LPS-primed StdMo, and PdMo that were reconstituted with platelets (PdMo + Plts) or culture supernatants from MEG-01 cells that were cultured in SILAC medium. Floating bars display the max/min values with indication for the mean (white bands). Each symbol represents one independent blood donor (n = 10).

(B - D)   Analysis of Mass speck proteomics combined with Stable isotope labeling with amino acids in cell culture (SILAC) in human PdMo exposed to cell-free supernatants of MEG-01 cells saturated with heavy AAs. Boxplots of quantitative distributions and (B) Principal component analysis (PCA) of SILAC proteomics samples from PdMo reconstituted with cell-free supernatants of MEG-01 cells.

(D)   Volcano plots of proteins with heavy-AAs detected in PdMo exposed to cell-free supernatants of MEG-01 cells through Mass speck proteomics combined with Stable isotope labeling with amino acids in cell culture (SILAC). Cells were stimulated with LPS. Unstimulated conditions are shown in **Fig S7**.

1082 **(E)**   WES capillary electrophoresis and immunoblotting of NFkB2 on whole cell lysates (Lysates) or cell-free supernatants
1083   (Sups) from human platelets. Representative of 2 independent experiments.

1084

Fig 8: Platelets restore cytokine function in NFκB2-deficient monocytes.

(A)   Schematic representation of the supplementation of platelet-depleted (PdMo) primary human monocytes with autologous
platelets that were left untreated (+ Fresh Plts), or pre-incubated for 20 min with 50 μM of BAY 11-7082 to inhibit NFκB (+
Plts$_{NFκB}$) or 20 μM of SB203580 to inhibit p38 MAPKs (+ Plts$_{p38}$) and washed before being added to PdMo.

(B)   IL-1β, TNFα, and IL-6 concentrations in CFS of StdMo, PdMo, or PdMo that were supplemented with platelets pretreated
with BAY11-7082 (+ Plts$_{BAY}$), SB203580 (+ Plts$_{SB}$), or left untreated (+ Plts$_{Intact}$). Co-cultures were stimulated with LPS (2 ng/ml)
followed by nigericin stimulation (10 μM). Side bars show the effects of direct exposure of monocytes to the inhibitors
(StdMo + Inhibitors).

(C)   Schematic representation of the inhibition of NFkB or MAPK p38 on primary human monocytes (StdMo) followed by their
replenishment with fresh autologous platelets.

(D)   Cell viability assay in human monocytes treated with BAY 11-7082 (50 μM, 30 min) or SB203580 (20 μM, 30 min) followed
by stimulation with LPS or LPS + Nig, as depicted in **C**.

(E)   IL-1β, TNFα, and IL-6 concentrations in the CFS of StdMo that were treated with SB203580 before been added with
increasing ratios of freshly isolated platelets (1:5, 1:50, and 1:100). Cells were stimulated with LPS or LPS and nigericin
(LPS + Nig). Floating bars display max/min values with indication to the mean (white bands). Each symbol represents one
donor. See also Fig S**8**.

(F)   Schematic representation of the genetic ablation of p65 (RelA) or p52/100 (NFκB2) on human THP-1 monocytes, and
knockout confirmation at the protein level by immunoblotting for each protein. Two clones are shown. Data is representative
of 3 independent experiments.

(G)   IL-1β concentrations in the CFS of stimulated RelA$^{-/-}$ or NFκB2$^{-/-}$ THP-1 monocytes that were cultured alone or added with
increasing ratios of freshly isolated platelets (1:20, 1:50, and 1:100). Cells were stimulated with Pam3CysK4 or Pam3CysK4
and nigericin (LPS + Nig). Floating bars display max/min values with indication to the mean (white bands). Each symbol
represents one donor. Data is pooled from 2 independent experiments.

1108

Fig 1. Platelets regulate the cytokine secretion of human monocytes.

(**A**)   Concentrations of IL-1β, TNFα, and IL-6 released into the cell-free supernatants (CFS) by untouched (StdMo, light blue), platelet-depleted (PdMo, blue), or PdMo that were supplemented with autologous platelets (PdMo + Plts, dark blue) at a 100:1 platelet:monocyte ratio. Cytokine levels secreted by platelets alone (Plts, purple) were measured as control. Cells were stimulated with LPS (2 ng ml^{-1} for 3h) followed by activation with nigericin (10 μM for 1.5 h).

(**B**)   Intracellular maturation and secretion of IL-1β assessed by WES capillary electrophoresis coupled with Ab-based detection in whole cell lysates (WCL) or CFS of primary human monocytes stimulated as in **A**.

(**C**)   Cell viability assessed every 15 min in StdMo, PdMo, PdMo + Plts, and Plts treated as in **A**.

(**D**)   Fluorometric assessment of caspase-1 activity in the CFS of primary human monocytes stimulated as in **A**.

(**E - F**) IL-1β concentrations in CFS of LPS-primed StdMo, PdMo, or PdMo + Plts that were transfected with Lipofectamin (LF) containing (**E**) double-strand DNA (dsDNA, 0.5 μg ml^{-1}) for the activation of the AIM2 inflammasome. In (**F**), cells were treated with PrgI (100 ng ml^{-1}) and protective antigen (PA, 1 μg ml^{-1}) for 90 min, for the activation of the NLRC4 inflammasome. Floating bars display the max/min values with indication for the mean (white bands). Each symbol represents one independent experiment or blood donor.

(**G**)   Radar plots displaying the top 19 highest cytokines detected by Cytokine Luminex in the CFS of human monocytes and platelets stimulated as in **A**. Protein concentrations are represented by the spread from inner (0 ng ml^{-1}) to outer circles (≥ 10 ng ml^{-1}). Colors represent stimuli (Grey: Unstim, Red: LPS, dark blue: LPS + Nig). Each symbol represents one independent experiment or blood donor (n = 5 for monocytes, and n = 2 for platelets alone). For better visualisation, cytokine concentrations (y axis) are displayed with a cut off at 8 ng ml^{-1}. Levels of IL-1RA and CCL4 reached 20 ng ml^{-1}. **See also Fig S1 and S2**.

A

ITP (n = 5)	
Demographic data	
Age (mean ± SD)	46 ± 9
Gender (Female/Male)	3/2
Confirmed ITP Diagnosis (%)	100
Duration since diagnosis (years mean ± SD)	6.5 ± 5.9
Laboratory results at baseline	
Platelet counts (G/L mean ± SD). Ref. Ranges: (150 - 370)	**53 ± 26**
Leukocyte count (G/L mean ± SD). Ref. Ranges: (3.9 - 10.2)	**6.8 ± 4.2**
CRP - mg/L (mean ± SD). Ref. Ranges: (0 - 3)	**1.9 ± 1.8**
Therapy at baseline	
Glucocorticoids (# / %)	0 / 0%
Thrombopoietin receptor agonist (# / %)	2 / 40%

Fig 2. Platelet supplementation reverts monocyte immunoparalysis in Immune Thrombocytopenia (ITP).

(**A**) Demographic and clinical characteristics of ITP patients (n = 5).

(**B**) Representative Flow Cytometry percentages and (**C**) CASY automated cell quantification of monocytes and platelets in preparations of untouched or platelet-depleted monocytes from healthy donors (± platelet-depletion) or untouched monocytes from ITP patients (ITPMo).

(**D**) Clinical laboratory quantification of platelets in the peripheral blood of ITP patients. Blue shaded area display the healthy reference ranges.

(**E-G**) Concentrations of IL-1β, IL-6 and TNFα released into the CFS by untouched (StdMo), and PdMo isolated from healthy volunteers (n = 4), or untouched monocytes from ITP patients (ITPMo, n = 5). Healthy platelet-depleted monocytes (PdMo + Plts), or ITPMo were supplemented with platelets (ITPMo + Plts). Cells were stimulated with LPS (2 ng ml^{-1}) for IL-6 and TNFα measurements, or LPS + Nig (10 μM) for IL-1β measurements. Data is displayed as floating bars with the max/min values and mean (white bands). P values were calculated with 2-Way Anova, Tukey's multiple comparison test, and are displayed in the Fig. Each symbol represents one independent experiment or blood donor.

Fig 3. Platelet depletion/supplementation dynamically governs gene expression of human monocytes.

(**A**) Principal Component Analyses (PCA) of gene expression in StdMo (green), platelet-depleted monocytes (PdMo, red), platelet-depleted monocytes reconstituted with platelets (PdMo + Plts, blue), and platelets alone (Plts, pink). Each symbol represents one donor. Stimulations are represented with shapes (elypse = unstimulated, square = LPS, triangle = LPS + Nig).

(**B**) Volcano plots showing log2 fold change (x-axis) and significance (−log10 *p-value; y-axis) of genes differentially expressed comparing LPS-stimulated vs. unstimulated (Unstim) StdMo, PdMo, or PdMo + Plts.

(**C**) Pie charts indicate the number of up- and downregulated DEGs in each group upon LPS stimulation.

(**D**) Quantitative expression (Log2 Fold Change) of the LPS-induced (>= 2 Fold Change) genes in StdMo (green), PdMo (red) or PdMo + Plts (blue).

(**E**) Scatter plots displaying the LSMean (x-axis) of StdMo vs PdMo (y-axis) from DEGs comparing the effects of platelet removal in Unstim and LPS-stimulated conditions. DEGs with 2-Fold change are highlighted in red.

(**F**) Pie charts representing the DEGs of the comparisons in **D**, representing DEGs induced by platelet depletion in unstimulated or LPS-stimulated monocytes.

Fig 4: The platelet effect depends on the phagocytic machinery of monocytes.

(**A**) Confocal imaging of StdMo and platelets. Cells were stained with CD41a AF488 (platelets: green) and CD14 AF647 (monocytes: red). Nuclei were stained with Hoechst 34580 (blue). Scale: 24 µm (top panel); 4.8 µm (bottom panel). White arrows indicate points of contacts between platelets and monocytes. Images are from one representative of four independent experiments.

(**B**) IL-1β and TNFα levels in CFS of StdMo, PdMo, and PdMo that were supplemented with autologous platelets (PdMo + Plts, 100:1 platelets/ monocytes). Monocytes were pre-treated with Cytochalasin D (CytoD, 50 µM) or Dynasore (DS, 30 µM) before being supplied with Plts. Cells were stimulated with LPS (2 ng·ml-1), followed by activation with nigericin (10 µM).

(**C**) TNFα and IL-6 levels in CFS of LPS stimulated StdMo, PdMo, PdMo, or PdMo reconstituted with platelet releasates (Plt-Sups).

(**D**) Representative flow cytometry assessment and gating strategy of StdMo that were incubated with growing concentrations of KPL-1 (5, 10 or 25 µg ml-1), or IgG (25 µg ml-1). Gates indicate the frequencies of free platelets (CD45- CD41a+), MPAs (CD45+CD41a+) and platelet-free monocytes (CD45+ CD41a-). Data is representative of two independent experiments with several donors.

(**E**) Cumulative flow cytometry frequency of free monocytes (CD45+ CD41a-) in StdMo treated with growing concentrations of KPL-1 (5, 10 or 25 µg ml-1), a specific mAb against P-selectin, or an unrelated isotype IgG control (25 µg ml-1). See **Fig S4B** for gating strategy.

(**F**) IL-1β, TNFα and IL-6 levels released by stimulated StdMo that were treated with increasing concentrations of KPL-1 (5, 10 or 25 µg ml-1), or IgG (25 µg ml-1).

Graphs display floating bars with the max/min values and mean (white bands) combined from multiple experiments. P values were calculated with 2-Way Anova, Tukey's multiple comparison test, and are displayed in the Fig. Each symbol represents one independent experiment or blood donor.

Fig 5. Platelets are cellular sources of MAPKs and transcription factors.

(**A-B**) Coral trees display the activity of Protein Tyrosine (upper branches) and Serine/Threonine kinases (lower branches) in StdMo, PdMo, PdMo + Plts and platelets from 5 donors, upon LPS stimulation for 15 min. (**A**) Impact of platelet depletion on the activity of PTK and STK in unstimulated monocytes. (**B**) Effects of LPS stimulation on the activation of PTKs and STKs in StdMo, PdMo, PdMo + Plts and platelets. Red: increased kinase activity, Blue: reduced kinase activity.

(**C-D**) Scatter plots and **D** box plot showing the signal intensity for the phosphorylation status of phosphosites (x.axis) that are substracts of PTK (top) and STK (bottom) comparing StdMo, PdMo, PdMo + Plts, and platelets alone (Plts). Each symbol shows one donor (n = 5) and only include peptides which passed the QC. batch correction with a 2-step Combat correction. P values in D were calculated by Anova multiple comparison.

(**E**) Pathway analysis of the most represented signaling pathways dynamically changed upon platelet depletion/re-addition to primary human monocytes. See also **Fig S5**.

(**F**) NFκB activity assay in CFS of unstimulated (Unstim), or Pam3CSK4-stimulated (100 ng·ml₋₁) NFκB-SEAP and IRF-Lucia luciferase Reporter Monocytes cultured alone (−) or co-cultured with increasing concentrations of human platelets (1:10, 1:20, 1:50 and 1:100). SEAP-Activity was measured with QuantiBlue Buffer. Graphs show floating bars display max/min values with indication to the mean (white bands). Each symbol represents one donor, or independent experiment.

(**G**) Immunoblot of RelA (p65), Phosphorylated-p65 (P-p65), and IκBα in MoStd, MoPD and Plts that were left untreated (−) or stimulated with LPS for 5, 15, 30 or 60 min. Representative of three independent experiments.

Fig 6: Enrichment of platelet-derived NFκB2 in human monocytes.

(**A**) IL-1β released from LPS-primed StdMo, and PdMo that were reconstituted with platelets (PdMo + Plts) or culture supernatants from MEG-01 cells that were cultured in SILAC medium. Floating bars display the max/min values with indication for the mean (white bands). Each symbol represents one independent blood donor (n = 10).

(**B - D**) Analysis of Mass speck proteomics combined with Stable isotope labeling with amino acids in cell culture (SILAC) in human PdMo exposed to cell-free supernatants of MEG-01 cells saturated with heavy AAs. Boxplots of quantitative distributions and (**B**) Principal component analysis (PCA) of SILAC proteomics samples from PdMo reconstituted with cell-free supernatants of MEG-01 cells.

(**D**) Volcano plots of proteins with heavy-AAs detected in PdMo exposed to cell-free supernatants of MEG-01 cells through Mass speck proteomics combined with Stable isotope labeling with amino acids in cell culture (SILAC). Cells were stimulated with LPS. Unstimulated conditions are shown in **Fig S6**.

(**E**) WES capillary electrophoresis and immunoblotting of NFkB2 on whole cell lysates (Lysates) or cell-free supernatants (Sups) from human platelets. Representative of 2 independent experiments.

Fig 7: Platelet-derived NFκB2 restore cytokine function in NFκB2-deficient monocytes.

(**A**) Schematic representation of the supplementation of platelet-depleted (PdMo) primary human monocytes with autologous platelets that were left untreated (+ Fresh Plts), or pre-incubated for 20 min with 50 μM of BAY 11-7082 to inhibit NFκB (+ Plts^{NFκB}) or 20 μM of SB203580 to inhibit p38 MAPKs (+ Plts^{p38}) and washed before being added to PdMo.

(**B**) IL-1β, TNFα, and IL-6 concentrations in CFS of StdMo, PdMo, or PdMo that were supplemented with platelets pretreated with BAY11-7082 (+ Plts^{BAY}), SB203580 (+ Plts^{SB}), or left untreated (+ Plts^{Intact}). Co-cultures were stimulated with LPS (2 ng/ml) followed by nigericin stimulation (10 μM). Side bars show the effects of direct exposure of monocytes to the inhibitors (StdMo + Inhibitors).

(**C**) Schematic representation of the inhibition of NFκB or MAPK p38 on primary human monocytes (StdMo) followed by their replenishment with fresh autologous platelets.

(**D**) Cell viability assay in human monocytes treated with BAY 11-7082 (50 μM) or SB203580 (20 μM) followed by stimulation with LPS or LPS + Nig.

(**E**) IL-1β, TNFα, and IL-6 concentrations in the CFS of StdMo that were treated with SB203580 before been added with increasing ratios of freshly isolated platelets (1:5, 1:50, and 1:100). Cells were stimulated with LPS or LPS and nigericin (LPS + Nig). Floating bars display max/min values with indication to the mean (white bands). Each symbol represents one donor. See also Fig S**7**.

(**F**) Schematic representation of the genetic ablation of p65 (RelA) or p52/100 (NFκB2) on Cas9-expressing human THP-1 monocytes, and knockout confirmation at the protein level by immunoblotting for each protein. Two clones are shown. Data is representative of 3 independent experiments.

(**G**) IL-1β concentrations in the CFS of stimulated RelA$^{-/-}$ or NFκB2$^{-/-}$ THP-1 monocytes that were cultured alone, or added with increasing ratios of freshly isolated platelets (1:20, 1:50, and 1:100). Cells were stimulated with Pam3cysK4 or Pam3cysK4 and nigericin (LPS + Nig). Floating bars display max/min values with indication to the mean (white bands). Each symbol represents one donor. Data is pooled from 2 independent experiments.