Ageing in a collective: The impact of ageing individuals on social network structure

Erin R. Siracusa[a*], André S. Pereira[a,b], Josefine Bohr Brask[c], Josué E. Negron-Del Valle[d], Daniel

Phillips[d], Cayo Biobank Research Unit, Michael L. Platt[e,f,g], James P. Higham[h], Noah Snyder-

Mackler[d,i,j], Lauren J. N. Brent[a]

Affiliations:

[a]School of Psychology, Centre for Research in Animal Behaviour, University of Exeter, Exeter,

UK

[b]Research Centre for Anthropology and Health, Department of Life Sciences, University of

Coimbra, Portugal

[c]Department of Applied Mathematics and Computer Science, Technical University of Denmark,

Kongens Lyngby, Denmark

[d]Center for Evolution and Medicine, Arizona State University, Arizona, USA

[e]Department of Neuroscience, University of Pennsylvania, PA, USA

[f]Department of Psychology, University of Pennsylvania, PA, USA

[g]Department of Marketing, University of Pennsylvania, PA, USA

[h]Department of Anthropology, New York University, New York, USA

[i]School of Life Sciences, Arizona State University, Arizona, USA

[j]School for Human Evolution and Social Change, Arizona State University, Arizona, USA

*corresponding author; email: erinsiracusa@gmail.com

Abstract

Ageing affects many phenotypic traits, but its consequences for social behaviour have

only recently become apparent. Social networks emerge from associations between individuals.

The changes in sociality that occur as individuals get older are thus likely to impact network

structure, yet this remains unstudied. Here we use empirical data from free-ranging rhesus

macaques and an agent-based model to test two things: (1) how age-based changes in social

behaviour influence an individual's level of indirect connectedness in their network, and (2) how

social ageing feeds up to affect overall network structure. Our empirical analyses revealed that

female macaques became less indirectly connected as they aged for some, but not all measures

examined, suggesting that ageing animals can remain well integrated in some social contexts.

Our agent-based model showed that aged-based changes in social behaviour can scale up to the

network level. Networks with a higher proportion of old individuals were less connected and less

cohesive. We did not find similar effects of age distribution on the structure of empirical

networks, which may reflect limited age variation in our data. Overall, our results suggest a

potentially important and underappreciated role of age in the structure and function of animal

collectives.

41
42   **Introduction**
43
44   The costs and benefits of an animal's behaviour toward members of their collectives can

shape lifespans, life history evolution, and the pace of senescence. For example, the support that

cooperative breeding species receive from non-breeding helpers may lead to longer lifespans than

those of related solitary species [1]. The transfer of enhanced ecological knowledge to close

relatives during collective foraging may have contributed to the evolution of reproductive

senescence and prolonged post-reproductive lifespan (i.e., menopause) in female killer whales [2–

4]. Differentiated relationships formed between groupmates have also been linked to enhanced

longevity in a range of group-living taxa [5]. It is therefore clear that living in collectives can alter

the adult ageing process. However, an important question is whether ageing, in turn, influences

the behaviour of adults in collectives, and ultimately the structure of collectives themselves.

54   Growing evidence suggests that older adults differ in their social behaviours from young

ones [6–11]. One pattern that seems to be emerging across taxa is that older adults interact with

fewer individuals than do younger adults, concentrating social relationships on close associates

and kin [6,12–15]. Given that social networks are an emergent feature of association rules between

individuals [16], shifts in patterns of social behaviour with age could also affect high-order

network structure. Age-based changes in social behaviour could therefore scale up to alter an

individual's connections to the partners of their social partners (their indirect connectedness [17])

as well as the overall structure ("topology") of the social network, both of which can have

consequences for disease transmission [18–23], information transfer [24–26], the cohesive

movement of groups [27,28], and many other eco-evolutionary dynamics [29,30]. Yet little

attention has been given to understanding the impact of social ageing for the polyadic social world

or the structure of the collective.

Recent work has offered some glimpses into how measures of indirect connectedness can

differ between young and old adults. In some species, older adults have measures of indirect

connectedness that suggest they have partners who are themselves not well connected [9,31],

(but see [32,33]), and that they are less effective at reaching disparate nodes in the network

compared to younger adults [32–34]. In contrast, older adults are more strongly embedded in

cliques or clusters in their networks compared to younger adults in some species [9], but not in

others [31,33]. However, most research to date has compared differences in measures of indirect

connectedness among adults of different age classes (e.g., old versus young), but has lacked the

longitudinal data required to quantify how the social positioning of individuals changes across

their lifetimes. Such longitudinal analyses have the potential to reveal important patterns that

might otherwise be masked by differences between individuals or cohorts [35]. Tracking within-

individual changes in measures of indirect connectedness is essential for more firmly placing

changes in sociality across the lifespan within the larger ageing phenotype and therefore

understanding the causes and consequences of these patterns of social senescence [36].

Populations composed of a greater proportion of older (or younger) adults may also be

structured in meaningfully different ways, and this could affect important processes such as

communication and cooperation. For example, the loss of old individuals through age-related

disease or trophy hunting can disrupt intergenerational flow of accumulated social and ecological

knowledge, impeding collective movement and the ability to locate critical resources [4,37–40].

The age structure of a group can also regulate the behaviour of younger individuals [41,42],

influencing aggression rates and social cohesion [43]. The impact of diminished cohorts of

younger individuals on higher-order network structure is less well understood, but likely to have

repercussions for network connectivity and cohesiveness given that younger adults are more

socially active in many populations [8,10,12–14]. Despite the established ecological and

evolutionary importance of network structure [29,30] the underlying drivers of variation in

network topology remain understudied [44–47]. Ageing, as an important process underlying

patterns of individual-level variation in sociality, might therefore provide a window into how

simple processes can generate complex network structures [48,49].

Using both empirical data and a theoretical model, we explore how social ageing of

individuals relates to measures of indirect connectedness and overall network structure in a

group-living primate, the rhesus macaque (*Macaca mulatta*), which is an emerging model in

social ageing research [36,50]. Rhesus macaques are highly social cercopithecine primates that

live in matrilineal kin-groups and exhibit clearly differentiated social relationships with kin-

biased affiliation [51–53]. Patterns of social interactions and social attention vary from adulthood

into old age [6,32,54]. Critically, as female rhesus macaques grow older, they reduce the size of

their social networks, and focus their social effort on a few important partners including close kin

[6]. Despite this, females do not reduce the rate at which they engage in social interactions as

they age, indicating that although their ego-networks get smaller, older females continue to

invest the same amount of time into fewer relationships [6].

Here, we set out to test if age-based social changes relate to measures of an individual's

level of indirect connectedness [17,55], and if they scale up to influence network structure as a

whole. Individual-level social network metrics, including measures of direct associations and

indirect connectedness to the wider social world, relate to underlying, putatively simple, social

differences or processes, such as individual-level variation in general sociability or reassociation

tendency [48]. For example, an individual's general tendency to be sociable can be intuitively

quantified using a direct network metric (e.g., strength, or the sum of the weights of an individual's

ties to their partners), or using an indirect metric (e.g., weighted eigenvector centrality, which

measures how well-connected an individual is to its partners and how well connected those

partners are to others [48]. Drawing on this idea that underlying processes can predict an

individual's position in the network, and on the age-based changes in social behaviour we have

documented in rhesus macaque females, we made predictions for four common measures of

indirect connectedness: eigenvector centrality, betweenness, closeness, and clustering coefficient.

Adult female rhesus macaques maintain their strength of grooming ties to others as they age [6].

We therefore predicted that weighted eigenvector centrality would also remain stable with age.

We also previously found that as female rhesus macaques age, they increase their likelihood of

interacting with certain partners, especially their kin. That is, ageing females increase both their

tendency to reassociate with others, and their tendency to interact with someone from their own

sub-group. Individuals with higher within-group association tendencies mix less widely in their

networks and those with greater reassociation tendency are less likely to associate with new

individuals and connect disparate parts of the network [48]. Therefore, we predicted that as females

aged, they would have lower measures of betweenness and closeness, but a higher clustering

coefficient. Betweenness calculates the number of shortest paths between others in a network that

pass through an individual, thus individuals with low betweenness have a low capacity for linking

discrete clusters in a network [56]. Closeness measures the inverse distance to all other individuals

in the network, and so the lower an individual's closeness score, the more difficult it is for them

to reach others or to be reached [56]. Clustering coefficient measures how densely (or sparsely)

the network is clustered around a focal individual, with a high clustering coefficient indicating that

an individual's partners are highly connected to each other [56].

To determine if age-based changes at the individual level can facilitate changes in overall

network structure we built an agent-based model to simulate how changing proportions of old

individuals in a network affects network structure. We simulated two simple changes in

individuals' interaction patterns with age based on our previous findings in female rhesus

macaques. We then used empirical data from 19 networks of female rhesus macaques to explore

whether the simulated results were borne out under natural conditions. Using both the simulated

networks and empirical networks we determined how changing proportions of old individuals in

the network was related to three common measures of global network structure: mean degree,

network diameter and transitivity. Mean degree represents the average number of ties that each

individual has to others in the network. Diameter measures the longest path length in the network

and provides a measure of overall connectedness. Transitivity captures the level of clustering in

the network [56]. We predicted that networks with a greater proportion of old individuals would

be more sparsely connected due to older animals having fewer social partners (i.e., have lower

mean degree and network diameter). Given greater kin clustering with age, we expected

networks with more old individuals would be more clustered (i.e., have higher transitivity). Our

results provide a first step to understanding how individual changes in social tendencies with age

can scale up to global network structure, offering important insight into the consequences of

shifting demography for the structure and function of collectives.

152
153 **Methods**
154
155 **Study population and data**
156
Data used in this study were collected on a well-studied population of rhesus macaques

on the island of Cayo Santiago, off the southeastern coast of Puerto Rico. The current population

is maintained by the Caribbean Primate Research Center (CPRC) and is descended from 409

macaques that were introduced to the island from India in 1938. The animals are food

supplemented and provided with ad libitum access to water. There is no medical intervention,

and so the major causes of death at this predator-free site are illness and injury [52,57]. The

CPRC staff collect demographic data five days per week and thus track dates of birth and death

of all individuals with a high degree of accuracy.

Subjects were mature adult females 6 years and older [58] from 6 naturally formed

mixed-sex social groups. We used data collected between 2010-2017, a time period for which we

had detailed behavioural data from which to estimate social networks. We collected behavioural

data between 07:30 and 14:00, which are the working hours of the field station, using 10-min

focal animal samples and recording all behaviours continuously [59]. We balanced data

collection to ensure equal sampling of individuals throughout the day and over the course of the

year, resulting in approximately the same number of focal samples per individual per year.

Female rhesus macaques have a strict dominance hierarchy with maternal rank inheritance and

youngest ascendency [60]. We therefore established dominance ranks for all females in a given

year using observed win-loss interactions (as per [61,62]). Rank was assigned as "high" ($\geq 80\%$

of other females dominated), "medium" (50%-79% of other females dominated) or "low" ($\leq$

49% of other females dominated).

Because individuals showed the clearest age-based changes in their grooming

associations [6], we used grooming interactions to build social networks. Grooming behaviour

was collected by recording the duration of a grooming bout along with the identities of the

interactants and the direction of grooming. We focused only on interactions between adult

females and did not include interactions with males as we wanted to avoid capturing changes in

socio-sexual behaviour with age. Given that the behaviour of juveniles' is commonly influenced

by their lack of independence from their mother we also did not include interactions with

juveniles of either sex. We built 19 grooming networks from the following group-years (group F

2010-2017; group HH 2014; group KK 2015; group R 2015-2016; group S 2011; group V 2015-

2016), with data availability based on the focus of projects over time given limited person power.

The average size of our networks was 50.7 ($\pm$ SE = 3.9) adult females. All network metrics were

calculated in R (version 4.2.0; [63]) using the igraph package (version 1.3.1; [64]).

189
190   **Investigating the relationship between age and indirect connectedness**
191
To assess age-based changes in indirect connectedness we used weighted network

metrics, as these are more robust and provide higher resolution than binary measures [67]. In

these weighted networks, edges represented the undirected rate of grooming between a pair of

individuals (seconds of grooming/total number of hours that both individuals were observed in

focal animal samples). For these analyses we set out to test how individuals' level of indirect

connectedness to their social network changed from prime adulthood into later life. At six years

old, females are deemed adults [58] and previous research on the macaque population of Cayo

Santiago has shown that, for females that survive to reproductive age, the median lifespan is 18

years with a maximum lifespan of about 30 years [50,68]. In our dataset, the median age of adult

females was 10-years. To capture later-life changes in indirect connectedness we thus focused

our analyses on individuals aged 10 and older (in line with previous ageing studies in this system

[6]. Females ranged between 10 and 28 years old. We had 563 macaque years of data over 204

unique females, with an average of 2.8 years of data per individual (range: 1-8 years; Fig. S1).

Because there was variation in the age-ranges over which individuals were sampled, we used a

within-individual centering approach to capture changes in indirect connectedness across

individuals' lifespans [69]. Briefly, following the methodology of van de Pol & Wright (2009)

we split our age term into a between-individual effect (calculated as the mean age of an

individual across all observations; hereafter called average age) and a within-individual effect

(calculated as the deviation of an individual's age from their mean age; hereafter called within-

individual age). This within-age term was our primary variable of interest and reflects how an

individual's deviation from its age affects its indirect connectedness in the network.

We fit four models with our four response variables of interest: eigenvector centrality,

betweenness, closeness, and clustering coefficient. All statistical models included average age

and within-individual age as continuous fixed effects. Given the strict dominance hierarchy

exhibited by female rhesus macaques [60], we assessed whether social status affected the change

in an individual's measures of indirect connectedness with age [6] by fitting an interaction

between rank and within-individual age in all models. We removed the interaction when not

significant. We included individual ID, group, and year, as random effects to account for

repeated observations and to capture any variation in indirect connectedness measures that might

be due to differences between individuals, groups, or years. We also fitted within-individual age

as a random slope over individual ID to capture any among-individual variation in the change in

indirect connectedness with age. We had no biological reason to expect non-linearities in the

relationship between age and measures of indirect connectedness given that we were looking at

changes from prime adulthood to old age and we've previously found a linear relationship

between direct measures of connectedness and age [6]. Nevertheless, we fit a model with

quadratic terms for within-individual age and average age and compared that model to the model

with only linear age terms using leave-one-out cross-validation in the brms package (version

2.17.0) [66]. The quadratic terms never improved the model fit and so were not considered

further. For the eigenvector centrality model, we logit-transformed the response variable to

improve model fit and fitted a model with a Gaussian error distribution. For betweenness we

fitted a model with a zero-inflated Poisson distribution. For closeness and clustering coefficient

we fitted models with a Beta error distribution and a zero-one-inflated Beta error distribution,

respectively.

235
236   **Investigating the relationship between age distribution and network structure**
237
Agent-based model

The purpose of the model was to gain insights into how age-based changes in sociality

may affect global network structure – specifically how the age distribution of the population

impacts overall network structure. Our general approach was therefore to model artificial

populations where social behaviour depends on age, and use these to investigate the relationship

between age distribution and global network metrics.

We were particularly interested in the potential effects of the age-dependencies observed

in the rhesus macaque system on Cayo Santiago [6]. As mentioned above, female rhesus

macaques have been observed to change two aspects of their sociality with age: their number of

social partners, and the proportion of partners that are kin [6]. We therefore included kinship as

an attribute of individuals in the model, and parameterised the model based on our empirical

249 data.

250   More specifically, the model simulates social networks of varying age distributions, in

251 which the probability that two nodes (i.e., individuals) have an edge (i.e., a grooming link)

252 depends on age and kinship of the individuals. Individuals belonged to two age categories (old

253 adults and young adults, hereafter "old" and "young") and two kin categories (kin and non-kin).

254 This gives 6 dyad types, which we denote by the age categories of the two individuals and their

255 kinship status (e.g. old/old kin). We established the linking probability (i.e., the chance of having

256 an edge in the social network) for each dyad type based on the mean proportion of dyads of each

257 type that had an edge across the 19 empirical grooming networks. The linking probability for

258 dyads that were constituted of old individuals who were related to each other (i.e., old/old kin)

259 was 0.33; for old/old non-kin dyads it was 0.02; for old/young kin dyads it was 0.37; for

260 old/young non-kin dyads it was 0.05; for young/young kin dyads it was 0.27 and for

261 young/young non-kin dyads it was 0.08. We fixed group size at 50 individuals, which

262 approximates the mean number of adult females in real groups on Cayo Santiago (mean $\pm$ SE =

263 50.7 $\pm$ 3.9). Each simulated network had 10 clusters of individuals who were related to each

264 other (i.e., kin groups) with 5 individuals in each. Kinship was determined by kin group

265 membership: individuals in the same kin group were kin and individuals in different kin groups

266 were non-kin. The number of individuals within each kin group mirrors the mean number of

267 close adult female kin (relatedness coefficient $\geq$ 0.125) that each adult female rhesus macaque

268 had in our empirical data (mean $\pm$ SE = 5.2 $\pm$ 0.87).

269   Each simulation round (i.e., construction and quantification of one network) proceeded as

270 follows: We first randomly drew the number of old individuals in the group (n_{old}) from a uniform

271 distribution with the range [0, 50]. We then randomly assigned all 50 group members to age-

groups (n_{old} old individuals and $n_{young} = 50-n_{old}$ young individuals) and kin groups. This allowed

the age structure of the groups to vary from only young individuals to only old individuals. We

then constructed the social network by drawing links, where the chance of each dyad getting a

link depended on their dyad type (link probabilities given above). To determine whether a dyad

was given a link, we extracted a random value from a binomial distribution with a sample of 1

and probability equal to the linking probability of the type of dyad. If the extracted value was 1,

the dyad was given a link, if it was 0, the dyad was not given a link.

To confirm that the model was working as intended, we ran 10000 simulations (i.e., we

generated 10000 networks), from which we calculated the proportion of dyads of each type that

had a link. We confirmed that the means of these proportions corresponded well to their

respective empirical linking probability (Fig. S2). Furthermore, to ensure that the conclusions

from the model were not highly sensitive to the linking probability estimates, we investigated

how the shape of the relationships between the proportion of old individuals and the network

metrics changed when the difference between linking probabilities of old and young individuals

became smaller. Our endpoint was the case where there was no difference between the social

behaviour of old and young individuals (i.e., their respective linking probabilities converged to

the same values). We ran the model for this extreme case, for the original data-based values, and

for five other sets of linking probabilities, which were at regular distances between the original

values and the endpoint values (Table S1). For each of these cases, we ran 10000 simulations.

Under high sensitivity, any observed correlations would disappear quickly (e.g., at the first

reduction of difference in linking probabilities between age groups), whereas a more gradual

reduction would indicate robustness to the input. Given that our input values are only estimates

of the real values, robustness to the input values is important for drawing general conclusions

about the potential effects in the real system.

To investigate the relationship between age distribution and network structure, we ran

100000 replicates of the simulation (i.e., generated 100000 networks) and visualised the

relationship between the proportion of old individuals and network mean degree, diameter, and

transitivity. Additionally, we used the agent-based model to investigate if we should expect to

find relationships between the proportion of old individuals and network metrics in the empirical

analysis. To do this, we limited the number of old individuals in each network to between 2 and

10 (i.e., 4-20% of the group, mirroring the empirical data), and we ran 19 simulations (mirroring

the 19 empirical networks). We repeated this 50 times (i.e., 50 sets of 19 networks) to gauge the

consistency of the results.

Empirical analyses

To assess the relationship between age distribution and network structure, we built

undirected, unweighted networks (where edges simply represented the presence of a grooming

relationship or not) to align with our agent-based model. Our networks included all females aged

6 and older. We had three response variables of interest and fitted a separate model for each:

mean degree, diameter, and transitivity. To ask if age distribution relates to network structure, we

included the proportion of old individuals as a continuous predictor in all our models. Since 18 is

the median age of death in this population and maximum lifespan is about 30 years, we

considered individuals above 18 to be "old" [50,68]. The proportion of old females in our 19

networks ranged from 0.03 to 0.19 (Fig. 1). Given that rhesus macaques live in matrilineal

groups with kin-biased behaviour [51,70], we included the average relatedness of the network as

a continuous covariate to account for differences between groups in general relatedness. We also

included density as a continuous fixed effect in the models of diameter and transitivity to account

for the fact that density can drive variation in other global network metrics. We included year as

a random effect to account for yearly variation that might affect network structure. We fit all

models with a Gaussian error distribution. To account for a potential non-linear relationship

between age distribution and network structure we fit all models with a quadratic term for

proportion of old individuals and compared that to the model with only linear terms using leave-

one-out cross-validation in the brms package (version 2.17.0; [66]). The quadratic terms never

improved model fit and so were not considered further.

Statistical analyses

All empirical models were fitted in a Bayesian framework with different error structures

and random effects dependent on the data analysed (see above). We conducted all analyses using

R (version 4.2.0; [63]) and fitted all models in the Bayesian software STAN [65] using the brms

package (version 2.17.0; [66]). All fixed effects were given weakly informative priors (see

Supplementary Information for more details). We ran all models for 10,000 iterations across two

chains with a warm-up period of 2,000 iterations. We assessed model convergence by examining

trace plots to assess sampling mixing and by ensuring Rhat = 1. We considered estimates of

fixed effects to be significantly different from zero when the 95% credible intervals of the

posterior distribution did not overlap zero.

337
338   **Results**
339
340   **Relationship between age and indirect measures of network connectedness**
341

Female macaques did not show any change in eigenvector centrality with age (within-

individual age: β = -0.04; 95% CI = -0.30, 0.24; Fig. 2A-B, Table S2). That is, as females aged,

the strength of their relationships to their partners, and to their partners' partners, remained stable.

We did find evidence of a within-individual decline in betweenness with age, suggesting older

individuals were less effective at reaching disparate nodes in the network. This effect was rank

dependent and seems to be driven primarily by a decline in betweenness as high-ranking

individuals got older (within-individual age:rankH: β = -0.35; 95% CI = -0.41, -0.29; Fig. 2C-D,

Table S3), although mid-ranking individuals also showed greater declines in betweenness with age

than did low-ranking females (within-individual age:rankM: β = -0.14; 95% CI = -0.18, -0.09; Fig.

2C-D, Table S3). Individuals showed a reduction in their closeness with age (within-individual

age: β = -0.18; 95% CI = -0.35, -0.02; Fig. 2E-F, Table S4), meaning they were harder to reach

and be reached in the network. There was no within-individual effect of age on clustering

coefficient, that is, how cliquish individuals were (within-individual age: β = 0.04; 95% CI = -

0.06, 0.14; Fig. 2G-H, Table S5).

Relationship between age distribution and network structure

Agent-based model results

360   Shifting group demographics in age appeared to have important consequences for

network structure (Fig. 3; Fig. S3). Specifically, our simulations showed that as the proportion of

old individuals in the population increased, mean degree (i.e., the mean number of partners that

each individual associated with) decreased (Fig. 3D), and the diameter of the network (i.e., the

longest path length in the network) increased (Fig. 3E), as expected. Network transitivity (i.e.,

the degree of clustering in the network; Fig. 3F) showed little variation in relation to age-based

distribution. Overall, these results suggest that compared to networks with more young

individuals, networks with more old individuals are sparser and less cohesive but are not

substantially different in how clustered they are. The patterns described above were only

gradually reduced when the difference in social behaviour between young and old individuals

was reduced, indicating that conclusions are robust to input parameter values (Fig. S4). When we

limited the sample size and variation in age distribution to those of the empirical data (i.e., 19

networks constituted by 4-20% of old individuals) we found no clear effects of age distribution

on network structure (Fig. 4C-D). These results hint that we might not expect to find a strong

effect of age-based changes in sociality on the structure of 19 empirical networks.

Empirical results

We found little evidence that groups with a greater number of old individuals were

structured differently from groups with fewer old individuals. Networks were not more sparsely

connected as the proportion of older animals increased. We found no overall effect of the

proportion of old individuals in the group on mean degree (β = -2.76; 95% CI = -10.24, 5.25; n =

19; Fig. 4A, Table S6), or network diameter (β = 2.02; 95% CI = -6.21, 10.00; n = 19; Fig. 4B,

Table S7). Older networks were also not more clustered or cliquish, as measured by transitivity

(β = 0.17; 95% CI = -0.31, 0.65; n = 19; Fig. 4C, Table S8).

383
384 **Discussion**
385
Age has begun to emerge as an important attribute shaping the social decisions of

individuals [6,7,14]. This implies that it is a potentially significant feature underlying the

behaviour of animals in collectives. Here we have shown that age-based changes in social

behaviour are likely to have important effects on higher-order network structure. As individuals

aged, they showed changes in some, but not all of their indirect measures of connectedness.

Furthermore, our agent-based model provides evidence that older populations can have

substantially different network structures than younger populations. While this suggests that age

plays a role in the overall structuring of rhesus macaque networks, this effect was not detected,

and may not have been detectable, in our empirical data.

In line with our predictions, we found no evidence for changes in eigenvector centrality

with age. This was expected because female rhesus macaques maintain the amount of time they

engage in social interactions as they get older [6], allowing them to continue to have strong

connections to some of their partners, and (indirectly) to the partners of their partners. A similar

pattern has been observed in a population of wild rhesus macaques in China whereby older

individuals had fewer partners but exhibited similar weighted eigenvector centrality to those of

younger individuals [32]. Unexpectedly, we did not find any evidence that clustering coefficient

increased as individuals aged in this population. Females increase their preference for grooming

kin as they age [6], suggesting that within-group (e.g., homophilic) tendencies increase with age,

which should lead to greater clustering as females from the same matriline would cluster together

in the network. However, although kin-biases increase with age, they exist for females of all ages

[53]. It is therefore possible that ties with kin are the main driver of clustering coefficient scores

regardless of age. Pruning a few non-kin relationships as an individual ages could thus have a

relatively minor impact on their clustering coefficient, leading to stable values for this metric

across the adult lifespan. In general, the stability of weighted eigenvector centrality and

clustering coefficient across the adult lifespan suggest that older animals can remain socially

central and well-integrated in some respects, despite other aspects of their social life changing

(e.g., fewer partners overall, and the patterns observed below). Old individuals may therefore

continue to reap some of the advantages of social relationships [5].

As predicted, we found declines in betweenness and closeness with age. As individuals

got older, they had less influence on and became less well-connected to the wider network.

These metrics are associated with an individual's level of reassociation tendency and within-

group association [48]. In other words, the more likely individuals are to reassociate with the

same partners, and to select partners of similar characteristics to themselves, the less they mix

with the wider network and the less likely they are to connect distinct subgroups (i.e., have high

betweenness) or be easily reached by all others in the network (i.e., have high closeness). We've

previously shown that ageing rhesus macaque females have fewer partners, but engage with

those partners more often, and increase their preference for close relatives. In other words, while

mean dyadic association strength to associates increases with age, mixing with the wider network

should decline [6]. Our betweenness and closeness results seem to reflect these underlying

changes in behavioural patterns. The declines in betweenness with age were rank dependent,

with mid-ranking and high-ranking rhesus macaque females showing a greater decline in

betweenness with age than low-ranking individuals. This may simply be due to a floor effect –

because low ranking individuals have such low betweenness to begin with, there is little capacity

for further decline with age.

Generally, our findings demonstrate that how indirectly connected an individual is to

their social world can change across their lifespan. How an individual is positioned in the wider

social environment can modulate their exposure to information [26,71], parasites [18,72], and

pathogens [21–23]. Individuals with high betweenness and closeness occupy a critical position in

the acquisition and transfer of 'goods' within a network [17,55,73]. Decreases in both of these

measures of indirect connectedness with age may benefit aged individuals who may experience

greater susceptibility to disease or illness as a result of immunosenescence [74]. Such benefits

need not imply that individuals actively change complex network positions with age. By

changing simple behavioural rules or processes with age, changes in polyadic ties with age are

likely to emerge [48]. Measures of indirect connectedness have been linked to fitness proxies

including future social status [75,76], survival [77,78], and reproductive success [61,76,79].

Indirect network metrics can be an even stronger predictor of fitness proxies [17,79] than

measures of direct dyadic associations and may more strongly reflect the underlying behavioural

rules that give rise to individual differences in sociality [48]. As such, documenting how ageing

shapes the polyadic social world may be particularly relevant for understanding how changes in

sociality across the lifespan influence patterns of senescence and fitness in later life.

In addition to age being associated with changes in the behaviour and social

connectedness of animals living in groups, our simulations provide evidence that age can also

impact the overall structure of those groups. By modelling age-based differences in two simple

interaction patterns, we found that the age composition of a group can have important

consequences for its cohesiveness and connectedness. As predicted, mean degree declined and

diameter increased as the proportion of old individuals increased, while transitivity exhibited no

strong relationship to age-distribution. Contrary to the results of the agent-based model, we

found no relationship between age distribution and network metrics in the observed macaque

data (Fig. 4). The impact of age distribution on network structure may only be detectable at the

extreme ends of the age distribution. That is, populations made up of very large numbers of

young, or old, individuals may diverge in structure at detectable levels, whereas populations

within a smaller range of age distributions may not. Although our empirical dataset ranged from

3-19% of individuals in the group classed as 'old', this may not have been wide enough to detect

differences in network structure. Results from our simulation were, as would be expected, the

most extreme at the more extreme ends of the age distributions examined, and when we

restricted our simulated networks to approximately 3-19% of individuals classed as 'old', we

found no relationship between the proportion of old individuals and network structure. We may

additionally have been unable to detect an effect of age distribution on network structure given

the amount of data available. Nineteen rhesus macaque networks from 6 groups were collected

across 8 years, which represents a considerable investment of research time and effort. But when

we restricted our agent-based model to a similar sample size, age distribution did not predict

network structure. However, it is also possible that our empirical results reflect a true null result

if there are processes occurring that our model has not accounted for. Our model, like all models,

is a simplistic version of the real world and does not fully replicate the complete suite of changes

that might occur with age.

The aim of our agent-based model was to capture two simple changes in individual-level

behaviours with age that have been identified across a wide range of taxa. Although the model

has the potential to be generalizable to other systems, it was parameterized based on data from

the macaque system and so its outputs may not hold in species that do not have a matrilineal kin

structure or low levels of modularity. Future work in other systems, conditions, and contexts

could reveal consequences for network structure that are even more stark. For example, more

dramatic differences between age classes in the overall probability of forming a social tie may be

expected in societies where the probability of connecting decreases with age but there is no

compensatory increase in the probability of forming ties with kin. This could lead to stronger

absolute effects on global network metrics, including substantial changes in transitivity, and thus

would likely have further consequences for network connectivity and spreading processes.

However, a greater proportion of older individuals in a population may not always lead to

changes in network structure. This can occur, of course, if individuals do not change their social

behaviour with age, but also in scenarios where age-based changes in sociality do exist. Within-

individual changes in social behaviour with age do not necessarily lead to between-individual

differences [69]. A cohort may start out very sociable due to, say, their environmental conditions

at birth. Even though members of this cohort become less sociable as they age, they could exhibit

similar average levels of sociality to a younger, less social, cohort. This would result in no

differences between old and young individuals in their social tendencies and therefore no effects

of age distribution on network structure. However, while cohort effects might occasionally mask

ageing effects, it seems unlikely that they would do so consistently. Future research may

therefore seek to assess at what point between individual differences can negate the effects of

within-individual changes with age.

Our model provides a glimpse into the powerful effects that age-based changes in

behaviour can have on network structure, offering a deeper understanding of the potential

importance of demographic changes for the structure and function of collectives. Much attention

has been given to the potential for the social structure of populations to be affected by the

removal of older individuals, for example through practices like trophy hunting or fishing

[80,81]. This is relevant given that older individuals can play a critical role in their social groups

through leadership [4,37], intergenerational transfers [38], and the stabilisation of social

relationships [82]. Despite the importance of the loss of older individuals from networks, our

results suggest that asking what happens when populations contain large numbers of older

individuals is an equally salient question.

In humans, population ageing is poised to be one of the most significant social

transformations of the 21st century, facilitated by falling fertility rates and increasing longevity

[83,84]. Meanwhile, in other group-living species, anthropogenic challenges, including climate

change, poaching, disease outbreaks, and pollution, among others, are likely to cause major

population declines and demographic shifts [85]. The consequences of declining fertility and/or

increasing mortality rates may be further exacerbated by the concomitant shifts in age structure

and resulting implications for network structure and function. For instance, declines in

cohesiveness and structural connectedness in older networks may limit or slow the transfer of

information [24,86], restrict the potential for cooperation [87] or reduce the stability of

populations [30], feeding back to further limit population growth. Alternatively, such changes

could reduce the vulnerability of populations to infectious diseases or parasites [30,88] or

enhance cooperation if network modularity is increased [89]. Consequently, through its effects

on network structure, shifting age demography may have broader implications than previously

appreciated for group dynamics and persistence and warrants further research.

Acknowledgments: Thank you to Kate and Christos for organising a stimulating meeting and to

the Royal Society for hosting. We thank the Caribbean Primate Research Center for maintaining

the Cayo Santiago population and for access to the study site, and all the field technicians who

have contributed to the long-term behavioural database over the years. We are grateful to

members of the Centre for Research in Animal Behaviour at the University of Exeter, as well as

Social Network Club, for thoughtful discussion during the manuscript's development. This work

was supported by the following grants from the National Institute of Health (NIH): grant nos

R01-AG060931, R00-AG051764, R01-MH096875, R37-MH109728, R01-MH108627, R01-

MH118203, U01MH121260, R01-NS123054 and the Kaufman Foundation: grant no KA2019-

105548. The Cayo Santiago Field Station is supported by the Office of Research Infrastructure

Programs of the NIH (2P40OD012217).

Data availability: All data and code associated with the analyses will be made publicly available

on the Figshare Repository following acceptance of the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Figure 1. Distribution of ages in each of the 19 empirical macaque networks. The dotted black line at age 18 indicates the median age of death in this population.

Figure 2. Relationship between within-individual changes in age and indirect measures of connectedness including (A & B) eigenvector centrality, (C & D) betweenness, (E & F) closeness, and (G & H) clustering coefficient in female rhesus macaques. In **A, C, E, G** the points represent raw data. Shaded grey bars indicate 95% confidence intervals around the predicted values. In **B, D, F, H** effect sizes and 95% credible intervals for all fixed effects and interaction terms are shown. Instances where the 95% CI overlaps zero are coloured in purple.

Figure 3. (**A-C**) Example networks from the agent-based model illustrating differences in network structure with the proportions of old individuals in the network equal to 0 (**A**), 0.5 (**B**) and 1 (**C**). Node colour represents kin groups where nodes of the same colour belong to the same kin group. (**D-F**) Results from the agent-based model showing relationship between the proportion of old individuals and (**D**) mean degree, (**E**) diameter, and (**F**) transitivity. Data points from the 100K simulations are shown with a small amount of jitter introduced to show overlapping points and are fitted with a smoothing term.

Figure 4. (A-C) The relationship between the proportion of old individuals in female rhesus
macaque networks and: **(A)** mean degree, **(B)** network diameter, and **(C)** network transitivity.
Points represent raw data and shaded grey bars indicate 95% confidence intervals around the
predicted values from models. **(D-F)** Relationship between the proportion of old individuals and
(D) mean degree, **(E)** network diameter, and **(F)** transitivity based on 19 simulations from the
agent-based model where the proportion of old individuals in the population was restricted
between 0.04 and 0.20. We re-simulated these 19 networks with limited age variation 50 times to
help ensure our results were robust. Each colour represents a different 'bout' of 19 networks.
Data points are shown fitted with a linear smoothing term.
577
578

579
580
581

582
583
584
585

References

1. Lucas ER, Keller L. 2020 The coevolution of longevity and social life. *Funct. Ecol.* **34**, 76–87.

2. Johnstone RA, Cant MA. 2010 The evolution of menopause in cetaceans and humans: the role of demography. *Proc. Biol. Sci.* **277**, 3765–3771.

3. Croft DP, Brent LJN, Franks DW, Cant MA. 2015 The evolution of prolonged life after reproduction. *Trends Ecol. Evol.* **30**, 407–416.

4. Brent LJN, Franks DW, Foster EA, Balcomb KC, Cant MA, Croft DP. 2015 Ecological knowledge, leadership, and the evolution of menopause in killer whales. *Curr. Biol.* **25**, 746–750.

5. Snyder-Mackler N *et al.* 2020 Social determinants of health and survival in humans and other animals. *Science* **368**. (doi:10.1126/science.aax9553)

6. Siracusa ER, Negron-Del Valle JE, Phillips D, Platt ML, Higham JP, Snyder-Mackler N, Brent LJN. 2022 Within-individual changes reveal increasing social selectivity with age in rhesus macaques. *bioRxiv.* , 2022.05.31.494118. (doi:10.1101/2022.05.31.494118)

7. Machanda ZP, Rosati AG. 2020 Shifting sociality during primate ageing. *Philosophical Transactions of the Royal Society B: Biological Sciences*. **375**, 20190620. (doi:10.1098/rstb.2019.0620)

8. Weiss MN *et al.* 2021 Age and sex influence social interactions, but not associations, within a killer whale pod. *Proceedings of the Royal Society B: Biological Sciences* **288**, 20210617.

9. Kroeger SB, Blumstein DT, Martin JGA. 2021 How social behaviour and life-history traits change with age and in the year prior to death in female yellow-bellied marmots. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **376**, 20190745.

10. Albery GF, Clutton-Brock TH, Morris A, Morris S. 2021 Ageing red deer alter their spatial behaviour and become less social. *bioRxiv*

11. Verhulst S, Geerdink M, Salomons HM, Boonekamp JJ. 2014 Social life histories: jackdaw dominance increases with age, terminally declines and shortens lifespan. *Proc. Biol. Sci.* **281**, 20141045.

12. Wrzus C, Hänel M, Wagner J, Neyer FJ. 2013 Social network changes and life events across the life span: a meta-analysis. *Psychol. Bull.* **139**, 53–80.

13. Rosati AG, Hagberg L, Enigk DK, Otali E, Emery Thompson M, Muller MN, Wrangham RW, Machanda ZP. 2020 Social selectivity in aging wild chimpanzees. *Science* **370**, 473–476.

14. Almeling L, Hammerschmidt K, Sennhenn-Reulen H, Freund AM, Fischer J. 2016

Motivational Shifts in Aging Monkeys and the Origins of Social Selectivity. *Curr. Biol.* **26**,
1744–1749.

15. Schino G, Pinzaglia M. 2018 Age-related changes in the social behavior of tufted capuchin
monkeys. *Am. J. Primatol.* **80**, e22746.

16. Hinde RA. 1976 Interactions, Relationships and Social Structure. *Man* **11**, 1–17.

17. Brent LJN. 2015 Friends of friends: are indirect connections in social networks important to
animal behaviour? *Anim. Behav.* **103**, 211–222.

18. White LA, Forester JD, Craft ME. 2017 Using contact networks to explore mechanisms of
parasite transmission in wildlife. *Biol. Rev. Camb. Philos. Soc.* **92**, 389–409.

19. Firth JA. 2020 Considering Complexity: Animal Social Networks and Behavioural
Contagions. *Trends Ecol. Evol.* **35**, 100–104.

20. Godfrey SS, Sih A, Bull CM. 2013 The response of a sleepy lizard social network to altered
ecological conditions. *Anim. Behav.* **86**, 763–772.

21. Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in wild
meerkats. *Proc. Biol. Sci.* **277**, 633–642.

22. Morrison RE, Mushimiyimana Y, Stoinski TS, Eckardt W. 2021 Rapid transmission of
respiratory infections within but not between mountain gorilla groups. *Sci. Rep.* **11**, 19622.

23. Weber N, Carter SP, Dall SRX, Delahay RJ, McDonald JL, Bearhop S, McDonald RA.
2013 Badger social networks correlate with tuberculosis infection. *Curr. Biol.* **23**, R915–6.

24. Aplin LM, Farine DR, Morand-Ferron J, Cockburn A, Thornton A, Sheldon BC. 2015
Experimentally induced innovations lead to persistent culture via conformity in wild birds.
Nature **518**, 538–541.

25. Evans JC, Silk MJ, Boogert NJ, Hodgson DJ. 2020 Infected or informed? Social structure
and the simultaneous transmission of information and infectious disease. *Oikos* **129**, 1271–
1288.

26. Claidière N, Messer EJE, Hoppitt W, Whiten A. 2013 Diffusion Dynamics of Socially
Learned Foraging Techniques in Squirrel Monkeys. *Curr. Biol.* **23**, 1251–1255.

27. King AJ, Sueur C, Huchard E, Cowlishaw G. 2011 A rule-of-thumb based on social
affiliation explains collective movements in desert baboons. *Anim. Behav.* **82**, 1337–1345.

28. Bode NWF, Wood AJ, Franks DW. 2011 The impact of social networks on animal
collective motion. *Anim. Behav.* **82**, 29–38.

29. Cantor M *et al.* 2021 The importance of individual-to-society feedbacks in animal ecology
and evolution. *J. Anim. Ecol.* **90**, 27–44.

30. Kurvers RHJM, Krause J, Croft DP, Wilson ADM, Wolf M. 2014 The evolutionary and

ecological consequences of animal social networks: emerging issues. *Trends Ecol. Evol.* **29**,
326–335.

31. Rathke E-M, Fischer J. 2021 Social aging in male and female Barbary macaques. *Am. J.*
Primatol. , e23272.

32. Liao Z, Sosa S, Wu C, Zhang P. 2018 The influence of age on wild rhesus macaques'
affiliative social interactions. *Am. J. Primatol.* **80**. (doi:10.1002/ajp.22733)

33. Brent LJN, Ruiz-Lambides A, Platt ML. 2017 Persistent social isolation reflects identity and
social context but not maternal effects or early environment. *Scientific Reports*. **7**.
(doi:10.1038/s41598-017-18104-4)

34. Wey TW, Blumstein DT. 2010 Social cohesion in yellow-bellied marmots is established
through age and kin structuring. *Anim. Behav.* **79**, 1343–1352.

35. Nussey DH, Coulson T, Festa-Bianchet M, Gaillard J-M. 2008 Measuring Senescence in
Wild Animal Populations: Towards a Longitudinal Approach. *Funct. Ecol.* **22**, 393–406.

36. Siracusa ER, Higham JP, Snyder-Mackler N, Brent LJN. 2022 Social ageing: exploring the
drivers of late-life changes in social behaviour in mammals. *Biol. Lett.* **18**, 20210643.

37. Allen CRB, Brent LJN, Motsentwa T, Weiss MN, Croft DP. 2020 Importance of old bulls:
leaders and followers in collective movements of all-male groups in African savannah
elephants (Loxodonta africana). *Scientific Reports*. **10**. (doi:10.1038/s41598-020-70682-y)

38. McComb K, Moss C, Durant SM, Baker L, Sayialel S. 2001 Matriarchs as repositories of
social knowledge in African elephants. *Science* **292**, 491–494.

39. Mueller T, O'Hara RB, Converse SJ, Urbanek RP, Fagan WF. 2013 Social learning of
migratory performance. *Science* **341**, 999–1002.

40. Lee HC, Teichroeb JA. 2016 Partially shared consensus decision making and distributed
leadership in vervet monkeys: older females lead the group to forage. *Am. J. Phys.*
Anthropol. **161**, 580–590.

41. Allen CRB, Croft DP, Brent LJN. 2021 Reduced older male presence linked to increased
rates of aggression to non-conspecific targets in male elephants. *Proc. Biol. Sci.* **288**,
20211374.

42. Slotow R, van Dyk G, Poole J, Page B, Klocke A. 2000 Older bull elephants control young
males. *Nature* **408**, 425–426.

43. Bourjade M, des Roches A de B, Hausberger M. 2009 Adult-Young Ratio, a Major Factor
Regulating Social Behaviour of Young: A Horse Study. *PLoS ONE*. **4**, e4888.
(doi:10.1371/journal.pone.0004888)

44. Cook PA, Baker OM, Costello RA, Formica VA, Brodie ED 3rd. 2022 Group composition
of individual personalities alters social network structure in experimental populations of

forked fungus beetles. *Biol. Lett.* **18**, 20210509.

45. Shizuka D, Johnson AE. 2019 How demographic processes shape animal social networks.
Behav. Ecol. **31**, 1–11.

46. Pinter-Wollman N. 2015 Persistent variation in spatial behavior affects the structure and
function of interaction networks. *Curr. Zool.* **61**, 98–106.

47. Ilany A, Akçay E. 2016 Social inheritance can explain the structure of animal social
networks. *Nat. Commun.* **7**, 12084.

48. Firth JA, Sheldon BC, Brent LJN. 2017 Indirectly connected: simple social differences can
explain the causes and apparent consequences of complex social network positions. *Proc.*
Biol. Sci. **284**. (doi:10.1098/rspb.2017.1939)

49. Ilany A. 2019 Complex societies, simple processes: a comment on Shizuka and Johnson.
Behav. Ecol. **31**, 13–13.

50. Chiou KL *et al.* 2020 Rhesus macaques as a tractable physiological model of human ageing.
Philos. Trans. R. Soc. Lond. B Biol. Sci. **375**, 20190612.

51. Bernstein IS, Judge PG, Ruehlmann TE. 1993 Kinship, association, and social relationships
in rhesus monkeys (Macaca mulatta). *Am. J. Primatol.* **31**, 41–53.

52. Widdig A *et al.* 2016 Genetic studies on the Cayo Santiago rhesus macaques: A review of
40 years of research. *Am. J. Primatol.* **78**, 44–62.

53. Berman CM. 2016 Primate Kinship: Contributions from Cayo Santiago. *Am. J. Primatol.* **78**,
63–77.

54. Rosati AG, Santos LR. 2017 Tolerant Barbary macaques maintain juvenile levels of social
attention in old age, but despotic rhesus macaques do not. *Anim. Behav.* **130**, 199–207.

55. Wey T, Blumstein DT, Shen W, Jordán F. 2008 Social network analysis of animal
behaviour: a promising tool for the study of sociality. *Anim. Behav.* **75**, 333–344.

56. Wasserman S, Faust K, Stanley (University of Illinois Wasserman, Urbana-Champaign).
1994 *Social Network Analysis: Methods and Applications*. Cambridge University Press.

57. Pavez-Fox MA *et al.* 2022 Reduced injury risk links sociality to survival in a group-living
primate. *bioRxiv.* , 2022.04.05.487140. (doi:10.1101/2022.04.05.487140)

58. Blomquist GE, Sade DS, Berard JD. 2011 Rank-Related Fitness Differences and Their
Demographic Pathways in Semi-Free-Ranging Rhesus Macaques (Macaca mulatta). *Int. J.*
Primatol. **32**, 193–208.

59. Altmann J. 1974 Observational study of behavior: sampling methods. *Behaviour* **49**, 227–
267.

60. Datta S. 1988 The acquisition of dominance among free-ranging rhesus monkey siblings.

Anim. Behav. **36**, 754–772.

61. Brent LJN, Heilbronner SR, Horvath JE, Gonzalez-Martinez J, Ruiz-Lambides A, Robinson
AG, Skene JHP, Platt ML. 2013 Genetic origins of social networks in rhesus macaques. *Sci.*
Rep. **3**, 1042.

62. Ellis S, Snyder-Mackler N, Ruiz-Lambides A, Platt ML, Brent LJN. 2019 Deconstructing
sociality: the types of social connections that predict longevity in a group-living primate.
Proc. Biol. Sci. **286**, 20191991.

63. R Core Team. 2021 R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

64. Csardi G, Nepusz T. 2006 The igraph software package for complex network research.
InterJournal, **Complex Systems**, 1695. https://igraph.org.

65. Stan Development Team. 2021 Stan Modeling Language Users Guide and Reference
Manual, 2.29. https://mc-stan.org.

66. Bürkner PC. 2017 brms: An R package for Bayesian multilevel models using Stan. *Journal*
of Statistical Software 80. doi:10.18637/jss.v080.i01.

67. Whitehead H. 2008 *Analyzing Animal Societies*. University of Chicago Press.

68. Brent LJN, Ruiz-Lambides A, Platt ML. 2017 Family network size and survival across the
lifespan of female macaques. *Proc. Biol. Sci.* **284**. (doi:10.1098/rspb.2017.0515)

69. Van de Pol M, Wright J. 2009 A simple method for distinguishing within-versus between-
subject effects using mixed models. *Anim. Behav.* **77**, 753.

70. Widdig A, Nürnberg P, Krawczak M, Streich WJ, Bercovitch FB. 2001 Paternal relatedness
and age proximity regulate social relationships among adult female rhesus macaques. *Proc.*
Natl. Acad. Sci. U. S. A. **98**, 13769–13773.

71. Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. 2012 Social networks predict patch
discovery in a wild population of songbirds. *Proc. Biol. Sci.* **279**, 4199–4205.

72. Rimbach R, Bisanzio D, Galvis N, Link A, Di Fiore A, Gillespie TR. 2015 Brown spider
monkeys (Ateles hybridus): a model for differentiating the role of social networks and
physical contact on parasite transmission dynamics. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*
370. (doi:10.1098/rstb.2014.0110)

73. Freeman LC. 1977 A Set of Measures of Centrality Based on Betweenness. *Sociometry* **40**,
35–41.

74. Peters A, Delhey K, Nakagawa S, Aulsebrook A, Verhulst S. 2019 Immunosenescence in
wild animals: meta-analysis and outlook. *Ecol. Lett.* **22**, 1709–1722.

75. McDonald DB. 2007 Predicting fate from early connectivity in a social network. *Proc. Natl.*
Acad. Sci. U. S. A. **104**, 10910–10914.

76. Gilby IC, Brent LJN, Wroblewski EE, Rudicell RS, Hahn BH, Goodall J, Pusey AE. 2013
FITNESS BENEFITS OF COALITIONARY AGGRESSION IN MALE CHIMPANZEES.
Behav. Ecol. Sociobiol. **67**, 373–381.

77. Stanton MA, Mann J. 2012 Early social networks predict survival in wild bottlenose
dolphins. *PLoS One* **7**, e47508.

78. Lehmann J, Majolo B, McFarland R. 2016 The effects of social network position on the
survival of wild Barbary macaques, *Macaca sylvanus*. *Behavioral Ecology*. **27**, 20–28.
(doi:10.1093/beheco/arv169)

79. Cheney DL, Silk JB, Seyfarth RM. 2016 Network connections, dyadic bonds and fitness in
wild female baboons. *R Soc Open Sci* **3**, 160255.

80. Birkeland C, Dayton PK. 2005 The importance in fishery management of leaving the big
ones. *Trends Ecol. Evol.* **20**, 356–358.

81. Roach DA, Carey JR. 2014 Population biology of aging in the wild. *Annu. Rev. Ecol. Evol.*
Syst. **45**, 421–443.

82. Archie EA, Chiyo PI. 2012 Elephant behaviour and conservation: social relationships, the
effects of poaching, and genetic tools for management. *Mol. Ecol.* **21**, 765–778.

83. Oeppen J, Vaupel JW. 2002 Demography. Broken limits to life expectancy. *Science* **296**,
1029–1031.

84. Harper S. 2014 Economic and social implications of aging societies. *Science* **346**, 587–591.

85. Selwood KE, McGeoch MA, Mac Nally R. 2015 The effects of climate change and land-use
change on demographic rates and population viability. *Biol. Rev. Camb. Philos. Soc.* **90**,
837–853.

86. Cantor M, Whitehead H. 2013 The interplay between social networks and culture:
theoretically and among whales and dolphins. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368**,
20120340.

87. Croft DP, James R, Thomas POR, Hathaway C, Mawdsley D, Laland KN, Krause J. 2006
Social structure and co-operative interactions in a wild population of guppies (Poecilia
reticulata). *Behav. Ecol. Sociobiol.* **59**, 644–650.

88. VanderWaal KL, Atwill ER, Hooper S, Buckle K, McCowan B. 2013 Network structure and
prevalence of Cryptosporidium in Belding's ground squirrels. *Behav. Ecol. Sociobiol.* **67**,
1951–1959.

89. Marcoux M, Lusseau D. 2013 Network modularity promotes cooperation. *J. Theor. Biol.*
324, 103–108.