Loss of Bcl6 transcriptionally alters classical dendritic cells and diminishes T follicular helper

cell and Th17-inducing cDC2 in mice

Hongkui Xiao[1,*], Isabel Ulmert[1,*], Johanna Huber[2], Hamsa Narasimhan[3,4], Luisa Bach[5], Ilia

Kurochkin[6,7], Urs Mörbe[1], Lili Zhang[8], Andreas Schlitzer[8], Filipe Pereira[6,7,9], Barbara U. Schraml[3,4],

Dirk Baumjohann[2,4], Katharina Lahl[1,10]

Affiliations:

[1]Section for Experimental and Translational Immunology, Institute for Health Technology,

Technical University of Denmark (DTU), 2800 Kongens Lyngby, Denmark

[2]Institute for Immunology, Biomedical Center, Faculty of Medicine, LMU Munich, Planegg-

Martinsried, Germany.

[3]Biomedical Center, Institute of Cardiovascular Physiology and Pathophysiology, Faculty of

Medicine, Ludwig-Maximillians-Universität München, Planegg-Martinsried, Germany.

[4]Walter-Brendel-Centre of Experimental Medicine, University Hospital, LMU Munich, Planegg-

Martinsried, Germany

[5]Medical Clinic III for Oncology, Hematology, Immuno-Oncology and Rheumatology, University

Hospital Bonn, University of Bonn, Bonn, Germany.

[6]Cell Reprogramming in Hematopoiesis and Immunity Laboratory, Lund Stem Cell Center,

Molecular Medicine and Gene Therapy, Lund University, Lund, Sweden.

[7]Wallenberg Center for Molecular Medicine, Lund University, Lund, Sweden.

[8]Quantitative Systems Biology, Life and Medical Sciences (LIMES) Institute, University of Bonn,

Germany.

[9]Center for Neuroscience and Cell Biology, University of Coimbra, Coimbra, Portugal.

[10]Immunology Section, Lund University, Lund 221 84, Sweden

*these authors contributed equally to this work

Corresponding authors: Katharina Lahl (1atharina.lahl@med.lu.se) and Dirk Baumjohann

(dirk.baumjohann@uni-bonn.de)

Keywords: dendritic cells, transcriptional regulation, Bcl6, cDC1, cDC2, Tfh

ABSTRACT

Conventional dendritic cells (cDC) play key roles in immune induction, but what drives their

heterogeneity and functional specialization is still ill-defined. Here we show that deletion of the

transcriptional repressor Bcl6 in murine cDC alters the phenotype and transcriptome of cDC1 and

cDC2, while their lineage identity is preserved. Bcl6-deficient cDC1 were diminished in the

periphery but maintained their ability to cross-present antigen to CD8[+] T cells, confirming general

maintenance of this subset. Surprisingly, the absence of Bcl6 in cDC caused a complete loss of

Notch2-dependent cDC2 in the spleen and intestinal lamina propria. DC-targeted Bcl6-deficient

mice induced fewer T follicular helper cells in response to the model antigen ovalbumin and

mounted diminished Th17 immunity to *Citrobacter rodentium* in the colon. Our findings establish

Bcl6 as an essential transcription factor for subsets of cDC and add to our understanding of the

transcriptional landscape underlying cDC heterogeneity.

INTRODUCTION

Conventional dendritic cells (cDC) have the unique ability to present antigen to naïve T cells

within lymphoid organs[1–3]. Upon encounter of antigen, migratory cDC scanning the peripheral

environment mature and relocate to the draining lymph nodes (LNs), where they collaborate

with LN-resident cDC to initiate immunity[4]. The spleen is the secondary lymphoid organ

responsible for induction of immune responses to blood-derived antigen[5] and maturation of

splenic cDC leads to similar relocation from the marginal zone and red pulp into the white pulp,

where naïve B and T cells reside[6,7]. "Resident" resting DC are also found in the spleen, where they

are located in the T cell zones in the absence of stimulation[8].

Conventional DC are broadly divided into two main subsets, cDC1 and cDC2[1]. cDC1 are

characterized by their expression of the cDC1-exclusive chemokine receptor XCR1 and the

absence of CD11b and Sirpα. Their development and functions are dependent on the

transcription factors (TFs) IRF8[9,10], BATF3[11], Nfil3[12,13] and Id2[14] and they excel at cross-presenting

antigen to CD8[+] T cells, which is at least in part attributable to their ability to engulf apoptotic

material[15–19]. cDC2 express high levels of CD11b and Sirpα, but not XCR1, and the transcriptional

network governing their development is less well defined. cDC2 show a higher level of

heterogeneity than cDC1[20] and recent attempts to subset cDC2 further have suggested

separation based on Notch2 versus Klf4-dependency[21]. IRF4-deficiency leads to a partial loss of

cDC2 and is thought to impact primarily on their survival and ability to migrate[22–24]. Other TFs

reported to influence cDC2 development or function include ZEB2[25], IRF-2[26] and Rel-B[27]. In the

spleen, the majority of cDC2 express endothelial cell specific adhesion marker (ESAM), a marker

defining cDC2 located in the bridging channels of the marginal zone, where they fulfill critical

functions in activating CD4[+] T cells and germinal center (GC) formation[28,29]. The development of

ESAM[hi] cDC2 depends on Notch2-signaling induced by delta ligand 1 (DLL1) expressed by stromal

cells[22,28,30,31] and on the chemokine receptor Ebi2 (GPR183)[32].

The TF Bcl6 plays an essential role in GC reactions, being required for the differentiation of both

GC B cells and T follicular helper (Tfh) cells in a cell intrinsic manner[33]. Bcl6 is expressed in cDC1,

and to a lower extent in cDC2[20] and we have shown previously that complete Bcl6-deficiency

causes a loss of CD11b[−] DC, presumed to be cDC1[20]. Other work has confirmed that Bcl6-deficient

mice lack splenic CD8-expressing cDC1, but also reported a reduction in splenic CD4[+] cDC2[34].

However, a more recent study has disputed these findings, by showing normal cDC1 numbers in

mice with conditional *Bcl6* deletion driven by cre recombinase under the control of the *Csf1*

promoter. Importantly, these authors suggested that down-regulation of CD11c in the absence

of Bcl6 may have compromised the detection of cDC1 in the earlier studies[35].

Here we reconcile these conflicting data by performing in-depth analysis of the cDC compartment

in mice lacking Bcl6 in all cDC or specifically in cDC1, using CD11c- and Clec9a- or XCR1-driven cre

drivers, respectively. Our results show that while cDC1 showed substantial alterations in gene

expression including deregulation of genes involved in cytokine signaling and cell adhesion, their

ability to cross-present antigen was largely conserved in the absence of Bcl6. Bcl6-deficiency in

DC led to a complete loss of the Notch2-dependent ESAM[hi] cDC2 subset in the spleen and

consequently to a defect in splenic Tfh cell induction to ovalbumin as a model antigen. Likewise,

Notch2-dependent cDC2 in the intestinal lamina propria (LP) were reduced in the absence of Bcl6

in DC, leading to a decrease in Th17-responses and delayed healing upon infection with the

enteric pathogen *Citrobacter rodentium*. Our findings suggest that the core subset identities of

cDC1 and cDC2 are maintained in the absence of Bcl6, but that its deficiency in DC causes cell-

intrinsic, subset-specific alterations in DC abundance and function, which translates into

corresponding defects in adaptive immune responses.

RESULTS

Bcl6 deficiency alters DC subset phenotypes and numbers across tissues

We have previously shown that mice completely lacking Bcl6 are largely devoid of CD11b⁻ cDC,

presumed to be cDC1[20]. To better define the impact of Bcl6-deficiency on cDC1, we generated

mice lacking Bcl6 specifically in cDC1 by crossing XCR1.cre mice to mice carrying a floxed Bcl6

gene (*XCR1.Bcl6^KO*). Additionally, we deleted Bcl6 in all CD11c-expressing cells by crossing

CD11c.cre mice to Bcl6^flox mice (*CD11c.Bcl6^KO*). As controls, we used the respective cre negative

Bcl6^flox littermates (designated as control throughout). As expected, flow cytometry revealed

mutually exclusive populations of CD11c⁺MHCII⁺ cDC in control spleens based on the expression

of XCR1 or CD11b to identify cDC1 and cDC2, respectively. However, there was a clear population

of XCR1⁺CD11b⁺ cDC in the absence of Bcl6 in the spleens of both Bcl6-deficient mouse lines (Fig.

1A). XCR1⁺CD11b⁺ cDC were also abundant within the resident populations of peripheral and

mesenteric lymph node (pLN and mLN) cDC in the absence of Bcl6 (Fig. 1B). The absolute numbers

of total XCR1⁺ cDC (independent of CD11b expression) were only mildly reduced in *XCR1.Bcl6^KO*

and normal in *CD11c.Bcl6^KO* mice, while XCR1⁻CD11b⁺ cDC were significantly diminished in

CD11c.Bcl6^KO mice, but normal in *XCR1.Bcl6^KO* mice (Fig. 1A). Resident cDC numbers were not

substantially altered in either pLNs or mLNs (Fig. 1B).

We next analyzed migratory DC populations in different organs. Mice lacking Bcl6 in cDC1 or in

all DC had significantly fewer migratory XCR1⁺ cDC in pLNs and mLNs and these also expressed

CD11b (Fig. 2A). Interestingly, we also detected a loss of XCR1⁻CD103⁺CD11b⁺ cDC in the

migratory compartment of the mLNs of *CD11c.Bcl6*^KO mice, whereas the numbers of XCR1⁻CD103⁻

CD11b⁺ cDC were normal (Fig. 2A). As the majority of migratory cDC in the mLNs derives from the

small intestinal lamina propria (SILP), we next analyzed this compartment. There were fewer

XCR1$^+$ cDC in the SILP of both *XCR1.Bcl6^{KO}* and *CD11c.Bcl6^{KO}* mice, all of which again expressed

CD11b (Fig. 2B). In addition, the numbers of XCR1$^-$CD103$^+$CD11b$^+$ cDC were reduced in

CD11c.Bcl6^{KO} mice, while the numbers of XCR1$^-$CD103$^-$CD11b$^+$ cDC were less affected (Fig. 2B).

Together these results suggest that the decreases in migratory XCR1$^+$ and XCR1$^-$CD103$^+$CD11b$^+$

cDC seen in mLNs are caused by absence of these populations, rather than a disability to migrate.

Similar changes in cDC were seen in the lung in the absence of Bcl6, with reduced numbers and

upregulation of CD11b in XCR1$^+$ cDC in both *XCR1.Bcl6^{KO}* and *CD11c.Bcl6^{KO}* mice, together with

reduced numbers of XCR1$^-$CD11b$^+$ cDC in *CD11c.Bcl6^{KO}* mice (Suppl. Fig. 1A). Thus, Bcl6 deficiency

affects cDC in several locations.

124   cDC1 and cDC2 signatures are preserved in the absence of Bcl6 in DC

One of the most surprising findings from our studies was the expression of the cDC2 marker

CD11b by the XCR1$^+$ cDC that was preserved in Bcl6-deficient mice. To determine whether this

reflects acquisition of CD11b by cDC1, or cDC2 with aberrant XCR1 expression, we sorted splenic

XCR1$^+$ cDC (independent of CD11b expression) and XCR1$^-$CD11b$^+$ cDC from control, *XCR1.Bcl6^{KO}*

and *CD11c.Bcl6^{KO}* mice for bulk mRNA-sequencing (see Fig. 1A for gating). Principal component

(PC) analysis showed that XCR1$^+$ cDC segregated from XCR1$^-$CD11b$^+$ cDC in all mouse models

along PC axis 1, which accounted for 69% of the variance. This strongly supports the idea that

Bcl6-deficient cDC1 and cDC2 maintained overall lineage identity (Fig. 3A). Nevertheless, PC2

separated the populations on the basis of Bcl6 presence or absence, accounting for 13% of the

variance and suggesting that Bcl6 has a significant impact on the gene transcription profile of

both cDC1 and cDC2. Interestingly, cDC1 from *XCR1.Bcl6^{KO}* and *CD11c.Bcl6^{KO}* mice clustered

closely and were separated from control cDC1, while cDC2 in *CD11c.Bcl6^{KO}* mice segregated from

those in *XCR1.Bcl6^{KO}* and control mice, consistent with XCR1-cre being specific to cDC1 and

indicating that Bcl6 regulates gene expression in both cDC subsets in a cell-intrinsic manner. PC3

accounted for 4% of the variance and showed opposite directional bias in Bcl6-deficient cDC1

and cDC2 (Fig. 3A). This suggests that Bcl6-mediated regulation of gene transcription may affect

some targets in cDC1 and cDC2 differently.

To explore further how Bcl6 might regulate cDC lineage identity, we generated cDC1 and cDC2

signatures by identifying the top 50 genes that were differentially expressed between our control

cDC1 and cDC2 and interrogated how these signatures were affected by a loss of Bcl6 expression

(for gene lists and expression levels see Suppl. Table 1). Supporting their maintenance of lineage

identity, XCR1$^+$ cDC from both *XCR1.Bcl6^{KO}* and *CD11c.Bcl6^{KO}* mice retained the expression of

cDC1 markers, including *Irf8*, *Tlr3*, *Clec9a*, and *Cadm1* (Fig. 3B). However, *Cd8a* was

downregulated in Bcl6-deficient XCR1$^+$ cDC and several genes associated with cDC2 identity were

upregulated by XCR1$^+$ cDC in the absence of Bcl6, including *Sirpa*, *Zeb2*, *Csf1r* and *Itgam* (encoding

CD11b) (foldchange > 2, padjust < 0.01). Although Bcl6-deficient CD11b$^+$ cDC2 did not express

cDC1 specific markers, cDC2 from *CD11c.Bcl6^{KO}* mice showed altered expression of several

classical cDC2 markers, of which some were further upregulated such as *Csf1*, *Zeb2* and *Itgam*,

while others were downregulated, including *Clec4a4* and *CD4* (Fig. 3B). Despite maintenance of

core lineage expression patterns, the overall transcriptional profiles of XCR1$^+$ and XCR1$^-$CD11b$^+$

cDC were significantly altered in the absence of Bcl6 (Fig. 3A, C). In total, 607 genes were

differentially regulated in XCR1$^+$ cDC from *XCR1.Bcl6^{KO}* mice compared with control cDC1, with

473 being upregulated and 134 downregulated. In contrast, of the 452 differentially regulated

genes in cDC2 from *CD11c.Bcl6^{KO}* mice, only 206 were upregulated, compared with 246

downregulated genes (Fig. 3D, for list of contributing genes see Suppl. Table 2). Interestingly, only

79 genes were coordinately overexpressed in Bcl6-deficient cDC1 and cDC2, and 32 genes were

coordinately downregulated in the absence of Bcl6 in both subsets. 12 genes were reciprocally

regulated in the two subsets, of which 11 genes were upregulated in cDC1 and downregulated in

cDC2 (including receptor encoding genes *Ffar1* and *Lair1,* enzyme-encoding genes *Tgm2, Pigv*

and *Malt1* and genes encoding proteins implicated in cellular and intracellular adhesion and

transport *Cdh17, Ctnnd2, Ston2 and Rin2*). The gene for the actin-binding protein Plectin was the

only gene upregulated in cDC2 and downregulated in cDC1. Taken together, although the lineage

identity of splenic cDC1 and cDC2 is generally maintained in the absence of Bcl6, this does result

in significant changes in subset-specific gene expression in both cDC subsets.

Bcl6-deficiency in cDC1 causes alterations in immunologically relevant pathways

To better characterize the impact of Bcl6-deficiency on cDC1, we next compared the expression

of cDC1-related markers using flow cytometry. This confirmed the maintenance of Tlr3, Xcr1, Irf8,

CD103 and Clec9a expression by XCR1$^+$ cDC from the spleen of *XCR1.Bcl6^{KO}* mice, as well as

upregulation of CD11b and Sirpα and loss of CD8α (Fig. 4A). In addition, we detected higher

expression of the costimulatory ligand CD24 on Bcl6-deficient cDC1. Next, we analyzed how the

absence of Bcl6 might influence the immunological functions of cDC1 by performing gene set

enrichment analysis of the RNA-Seq data from XCR1$^+$ cDC in *XCR1.Bcl6^{KO}* and control mice. This

showed that several immunologically relevant gene ontology pathways were enriched within the

differentially expressed gene sets, including *leukocyte adhesion*, *proliferation* and *activation*, as

well as *production of IL-1* (Fig. 4B). In parallel, KEGG pathway analysis highlighted differences in

cytokine-cytokine receptor interactions, NOD-like receptor signaling pathway, MERK signaling

pathway, hematopoiesis cell lineage, Tuberculosis and *C-type lectin receptor signaling pathway*

(Fig. 4C, D). These data suggest that Bcl6-mediated transcriptional repression in cDC1 may tune

the function of this subset.

186   Bcl6-deficient cDC1 maintain the ability to cross-present antigen

The most characteristic property of cDC1 is to cross-present exogenous antigen to class I MHC-

restricted CD8$^+$ T cells, a function which involves cell-surface receptors such as Clec9A (Dngr1)

that mediate the uptake of dead cells[18,19]. Our gene expression analysis suggested that XCR1$^+$

cDC1 maintained the expression of *Clec9a* in the absence of Bcl6 and this was confirmed by flow

cytometry (Fig. 4A). We therefore examined whether cross-presentation was intact in

XCR1.Bcl6^{KO} animals by injecting heat-killed ovalbumin-expressing mouse embryonic fibroblasts

(OVA-MEFs) into recipient mice adoptively transferred with OVA-specific TCR-transgenic CD8$^+$ T

cells from OT-I mice. OT-I proliferation strictly depends on cDC1-mediated cross-presentation in

this model[17]. OT-I cells proliferated to a similar extent in the spleens of OVA-MEF injected

XCR1.Bcl6^{KO} and control mice (Fig. 5A, B), suggesting that Bcl6-deficient cDC1 retain their ability

to cross-present. OTI proliferation was also observed in the mLN. In the mLN of *XCR1.Bcl6^{KO}* mice

OTI cells showed somewhat diminished proliferation compared to control mice, however, this

most likely reflects the decreased number of migratory cDC1 in the mLN of *XCR1.Bcl6*[KO] mice (Fig.

2A).

Taken together, although cDC1-specific Bcl6 deficiency alters the phenotype and transcriptional

profile of cDC1, the hallmark function of cDC1 to cross-present antigen is largely unaffected by

the loss of Bcl6.

Bcl6-deficiency results in a selective loss of Notch2-dependent cDC2

In addition to its role in cDC1, our initial results also revealed a significant role for Bcl6 in the gene

expression signature of cDC2. Furthermore, the numbers of cDC2 were reduced in the spleen of

CD11c.Bcl6[KO] mice, as were the numbers of cDC2 expressing CD103 in the intestinal LP and mLNs.

Diminished cDC2 numbers in the blood of CD11c.Bcl6[KO] mice suggest that this was not due to a

release of cDC2 into the bloodstream (Suppl. Fig. 1B). The remaining cDC in the spleen of

CD11c.Bcl6[KO] mice also showed somewhat lower expression of MHCII and higher expression of

CD11b than control cDC2 (Fig. 6A). Further phenotyping of spleen cDC2 in *CD11c.Bcl6*[KO] mice

revealed a striking loss of the ESAM[hi] cDC2 subset (Fig 6B and Suppl. Fig. 2). We verified our

findings an independent mouse model in which Bcl6 deletion was induced by cre recombinase

under the control of the *Clec9a* promoter, which targets cDC precursors and their progeny and

some pDCs but no other leukocytes[36]. These mice additionally carry a YFP reporter following a

floxed stop cassette in the ubiquitous ROSA26 locus to monitor targeting efficiency. As in

CD11c.Bcl6[KO] mice, spleen cDC in *Clec9a.Bcl6*[KO] mice showed altered CD11b expression and loss

of ESAM[hi] cDC2 (Fig. 6C, D and Suppl. Fig. 2). This was apparent in the analysis of bulk cDC2 in

Clec9a.Bcl6[KO] mice and was even more pronounced when examining the YFP[+] cDC2, which are

enriched for cDC2 with a history of *Clec9a*-promoter activity (Fig. 6E).

As expected[28], the ESAM[lo] subset of cDC2 expressed slightly lower levels of CD11c compared to

their ESAM[hi] counterparts and a significant proportion lacked expression of 33D1 and CD4, which

were expressed uniformly by ESAM[hi] cDC2 (Fig. 6F). The total population of cDC2 remaining in

CD11c.Bcl6[KO] mice resembled the ESAM[lo] population in terms of these markers but expressed

higher levels of CD11b and IRF4 than either ESAM[hi] or ESAM[lo] control cDC2 subset, suggesting

these markers may be specific targets of Bcl6 repression.

To explore the nature of the cDC2 remaining in the spleen of *CD11c.Bcl6[KO]* mice further, we

performed gene set enrichment analysis (GSEA) of differentially expressed genes between

CD11c.Bcl6[KO] and control cDC2 in gene sets derived from publicly available bulk RNA-Seq data

from ESAM[hi] and ESAM[lo] cDC2 from control spleen[37]. As expected from their usual predominance

in control spleen, control cDC2 gene signatures were highly enriched in ESAM[hi] vs ESAM[lo] gene

sets, while *CD11c.Bcl6[KO]* cDC2 gene signatures were highly enriched in ESAM[lo] vs ESAM[hi] gene

sets (Fig. 7A). We next directly compared the set of genes that was differentially expressed in

ESAM[hi] vs ESAM[lo] cDC2 with the gene set differentially expressed in *CD11c.Bcl6[KO]* vs control cDC2.

In accordance with the GSEA data, the majority of the top 50 genes expressed more prominently

by ESAM[hi] cDC2 were also significantly over-represented in the population of total control cDC2,

which expressed much fewer of the top 50 genes from ESAM[lo] cDC2 (Fig. 7B). Conversely, the

genes over-expressed by total cDC2 from *CD11c.Bcl6[KO]* spleen showed much more overlap with

the top 50 genes expressed by ESAM[lo] cDC2, while the genes over-expressed by ESAM[hi] cDC2

were mostly downregulated in Bcl6-deficient cDC2 (Fig. 7B, for a complete gene list containing

expression levels see Suppl. Table 3). These data confirm that Bcl6 deficiency in DC causes the

loss of the ESAM[hi] lineage of splenic cDC2.

DC-specific loss of Bcl6 causes defects in T follicular helper cell priming in the spleen

ESAM[hi] spleen DC reside in the marginal zone bridging channels, where they play a prominent

role in the induction of germinal centers[29]. Confocal microscopy using CD11c as a pan-DC marker

and TLR3 and Sirpα to define cDC1 and cDC2, respectively, confirmed a substantial loss of DC

from this region in the *CD11c.Bcl6[KO]* spleen (Fig. 7C). To test the functional relevance of this

defect, we adoptively transferred OVA-specific TCR-transgenic CD4[+] T cells (OT-II) into control

and *CD11c.Bcl6[KO]* mice, immunized the mice with OVA + poly(I:C) intraperitoneally and assessed

the generation of antigen-specific Tfh cells three days later (Fig. 7D). We chose this timepoint

because this early Tfh cell induction is independent of GC B cells[38,39], which are known to express

CD11c and are required for the maintenance of Tfh cells at later times[40]. Although OT-II cells

expanded normally in the *CD11c.Bcl6^KO* mice in response to immunization, the early induction of

antigen-specific Tfh cells was markedly reduced in immunized *CD11c.Bcl6^KO* mice compared to

control mice (Fig. 7E), suggesting that Bcl6-dependent cDC indeed drive early Tfh cell induction

in this model. To enable analysis of later timepoints of the GC reaction and to confirm findings in

mice in which Bcl6 deletion is confined to cDC, we adoptively transferred naïve OT-II cells into

control or *Clec9a.Bcl6^KO* mice, followed by i.p. immunization with OVA in alum adjuvants and

analysis of the spleen seven days later (Fig. 7D,F). The frequency of CXCR5$^+$PD-1$^+$ Tfh cells was

significantly reduced in OT-II cells recovered from *Clec9a.Bcl6^KO* hosts as compared to control

animals (Fig. 7F) and this was also the case for the endogenous Tfh cell pool in these mice. Thus,

Bcl6 deficiency in cDC causes the loss of ESAM^{hi} cDC2 and significantly impairs Tfh responses in

the spleen.

DC-specific loss of Bcl6 causes defects in Th17 cell priming in the colon of mice infected with *C.

rodentium*

In addition to their role in driving GC reactions in the spleen, Notch2-dependent cDC2 have been

shown to support Th17-dependent immunity towards *C. rodentium* in the colon[22]. The colonic LP

(cLP) of *CD11c.Bcl6^KO* mice contained significantly fewer cDC than that of control mice, with the

most pronounced loss detected in the CD103$^+$ cDC2 population (Fig. 8A, B). To analyze whether

Bcl6-deficiency recapitulated the functional consequences of Notch2-deficiency, we infected

CD11c.Bcl6^KO and control littermates with *C. rodentium. CD11c.Bcl6^KO* mice lost comparable

weight to control littermates early after infection, with delayed recovery as evidenced by

significantly lower body weight from day 11 onwards (Fig. 8C). Consistent with a lack in weight

difference between genotypes at day nine post infection, colon length and bacterial load were

comparable at this time point, while there was a tendency towards shorter colons and elevated

bacterial load in *CD11c.Bcl6^KO* mice compared to control littermates 21 days post infection (Fig.

8C). At this late stage during infection, colons from *CD11c.Bcl6^KO* mice continued to show mild

pathology, while control littermates appeared fully healed (Fig. 8D). In parallel, there were

blunted Th17 responses in the cLP of *CD11c.Bcl6^KO* mice 9 days after *C. rodentium* infection (Fig.

8E), while Th1 and Treg numbers were not affected. This defect in Th17 priming was no longer

evident 21 days after infection (Fig. 8F).

These data show that Bcl6-deficiency in DC causes the loss of a prominent colonic cDC2 subset,

with a consequent defect in Th17 priming in response to *C. rodentium* infection, leading to

delayed healing and recovery.

DISCUSSION

We show here that Bcl6-deficiency in the DC compartment causes major transcriptional

alterations with subset-specific consequences. Specifically, Bcl6-deficient cDC1 acquire a

XCR1/CD11b double-positive phenotype not seen in control mice, where XCR1 and CD11b are

mutually exclusive markers of cDC1 and cDC2, respectively. However, Bcl6-deficient cDC1

retained a cDC1-like transcriptome, cDC1 lineage markers and the ability to cross-present. The

lack of Bcl6 in DC also revealed a critical role for Bcl6 in the development of the ESAM^hi subset of

cDC2 in the spleen and cDC2 in the intestines and there was a parallel defect in splenic Tfh cell

induction in response to protein antigen immunization and in intestinal Th17 induction in

response to *C. rodentium* infection. Together these findings extend the role of Bcl6 to the

regulation of cDC2 behavior in GC function and the periphery.

We previously described a selective loss of putative cDC1 in the spleen of Bcl6-deficient mice

based on the absence of CD8α⁺CD11b⁻ cDC[20]. Here we refine this interpretation by showing that

the numbers of resident cDC1 are in fact normal when Bcl6 is lacking in CD11c⁺ cells or cDC1, but

they upregulate CD11b and down-regulate CD8α, causing them to merge into the classical cDC2

gate. In contrast, the numbers of migratory CD103⁺CD11b⁻ cDC1 are decreased in the mLN and

intestinal mucosa of mice lacking Bcl6 within the DC compartment, a finding supported by

defective cross presentation activity in the mLNs. We currently cannot explain the differential

effects of Bcl6 deficiency on resident versus migratory cDC1 populations. However, a similar

dichotomy has been described recently for dependence on DC-SCRIPT, whose absence

predominantly diminished resident cDC1[41]. As Bcl6 primarily acts as a transcriptional repressor,

it is possible that DC-SCRIPT and Bcl6 integrate in a feedback loop to regulate cDC1 migration or

survival upon maturation and this idea requires further study.

As Bcl6 is not expressed by DC progenitors in the bone marrow and their frequency is not affected

by the lack of Bcl6[35,42], Bcl6 seems to act downstream of other cDC1 lineage specifying TFs such

as IRF8, Nfil3, BATF3, DC-SCRIPT, and Id2. In models deficient of those TFs, the cDC1 lineage is

affected from the precursor level onwards, preventing their expression of all lineage markers, as

well as their ability to cross-present antigen[10,12–14,41,43–46], hallmarks that were preserved in Bcl6-

deficient cDC1. Rather, it appears that Bcl6 may fine-tune cDC1 identity by suppressing selected

cDC2 genes at the mature state and as a result, Bcl6-deficient cDC1 maintain key cDC1 genes such

as XCR1, TLR3, and CLEC9A as well as cross-presentation function. However, they show a gene

expression profile that suggests a cDC2-related "gain-of-function", with enhanced expression of

genes of the cytokine-cytokine receptor pathway, cell adhesion and migration pathways and

pathways associated with inflammasome activity. Interestingly, cDC1 with upregulated CD11b

and Sirpα have also been described when the TF ZEB2 is over-expressed from the pre-cDC stage

onwards[25]. Exploring how Bcl6 interacts with this and other lineage determining factors to

regulate cDC1 gene expression and functions will be an important topic for future research.

CD11c.Bcl6^{KO} and *Clec9a.Bcl6^{KO}* mice lacked the dominant subset of ESAM[hi] cDC2 in the spleen

and comparison of gene expression profiles between ESAM[hi] and ESAM[lo] control cDC2 subsets

and cDC2 from *CD11c.Bcl6^{KO}* mice showed that Bcl6-deficient cDC2 closely aligned with control

ESAM[lo] cDC2. However, this overlap was not absolute, suggesting that Bcl6-deficiency causes

transcriptional changes in all cDC2. For example, the extent of dysregulation of IRF4 and CD11b

expression in the absence of Bcl6 cannot solely be explained by the loss of ESAM[hi] cDC2. In

addition, both markers were also upregulated in Bcl6-deficient cDC1, again indicating a wider

role for Bcl6 in control of mature cDC differentiation. Interestingly, the consequences of Bcl6

deficiency on cDC overlap with those previously shown for Notch2 deficiency[20,28,34]. Specifically,

both mice show deficiencies in cDC1 and ESAM[hi] cDC2, but it remains to be determined whether

the apparent loss of cDC1 in Notch2-deficient animals reflects genuine absence of these cells, or

is because their phenotype has simply changed, as we show in Bcl6-deficient mice. Of note, the

cDC1 marker CD24 is upregulated in both Bcl6- and Notch2-targeted mice (our study and [22,28]),

arguing against a developmental defect within the cDC1 spleen compartment in Notch2-targeting

models. It will be of interest to explore how the Notch2- and Bcl6-pathways interact in DC biology.

Consistent with the findings that cDC2 equivalent to the ESAM^{hi} population locate preferentially

in marginal zone bridging channels in the spleen[47], the dysregulation in splenic cDC populations

in both Bcl6- and Notch2-deficient mice was associated with loss of cDC from this region of the

spleen ([29] and current results). Localization of cDC2 to the bridging channels is dependent on the

homing receptor GPR183[48,49] and Bcl6 regulates GPR183 expression in B cells[50]. However,

GPR183 expression levels were not altered in Bcl6-deficient cDC2, suggesting either that

additional factors are required for the correct positioning of cDC2 in the spleen, or that Bcl6-

deficiency causes a development block or diminished survival within ESAM^{hi} cDC2 irrespective of

their location. Along those lines, GPR183 deficiency in cDC2 specifically diminished antibody

response generation to particulate antigen, while responses to soluble antigen were

unaffected[49]. The defect in ESAM^{hi} cDC2 we found in *CD11c.Bcl6^{KO}* and *Clec9a.Bcl6^{KO}* mice was

accompanied by reduced induction of Tfh cells in response to parenteral immunization with

soluble OVA, suggesting a general deficiency in supporting GC reactions in these mice. While it

has been suggested that Notch2 is needed to ensure correct positioning of cDC2 for access to

antigen in the bloodstream[28], it has to our knowledge not been addressed whether response

deficits were limited to particulate antigen in these mice. Further investigation is needed to

explore the importance of correct localization on development, survival, and antigen capture by

cDC2 and how this might be controlled by a Bcl6-Notch2 axis.

XCR1$^-$CD103$^+$CD11b$^+$ intestinal LP DC were also reduced in the absence of Bcl6 in DC, again

mirroring the phenotype observed in Notch2-deficient mice[22,28]. cDC2 are potent drivers of Th17

responses in the intestines[23,51], but recent findings suggest that DC-independent Th17 cell

induction can also occur[52]. As *CD11c.Notch2^{KO}* mice succumb early to *C. rodentium* infection due

to deficient IL-23 driven innate immunity, the adaptive immune response has not been

investigated in this model[22]. In contrast, the early stages of infection were normal in *CD11c.Bcl6^{KO}*

mice, but these mice showed a delay in recovery at later stages and altered kinetics of Th17

induction, with significantly lower Th17 numbers at day nine, which had normalized by day 21.

We cannot currently explain the differences in the early innate immune response between

CD11c.Notch2^KO and *CD11c.Bcl6^KO* mice, but our data are consistent with the idea that DC play

the dominant role in coordinating the adaptive immunity required to terminate infection.

However, while Bcl6-dependent cDC2 support optimal Th17 induction to *C. rodentium*, they are

not essential for this response, as Th17 numbers normalized over time.

Our results show that Bcl6 deficiency does not alter cDC1 and cDC2 lineage identity, but it alters

their transcriptional profile and relative abundance of both lineages and their subsets in different

tissues. Specifically, although cDC1 in all tissues examined acquired several cDC2 markers in the

absence of Bcl6, they retained their transcriptional and functional identity as cDC1. In contrast,

Bcl6 deficiency had differing effects on cDC2 subsets that depended on their location and

phenotype, with selective depletion of ESAM^hi cDC2 in spleen and CD103^+CD11b^+ cDC2 in the

intestine. These effects were similar to those described for a deficiency in Notch2 and had

significant functional consequences for adaptive immunity in the spleen and intestine. These data

add to the emerging and poorly understood heterogeneity within the cDC2 compartment by

establishing that certain cDC2 subsets require Bcl6 in addition to Notch2- and lymphotoxin β-

signaling for their development. Finally, complementary to its established roles in controlling B

and T cell-intrinsic cell fate decisions[53], our findings establish Bcl6 as a subset-specific regulator

of DC development and function with critical consequences for the induction of adaptive immune

responses.

MATERIALS AND METHODS

Mice

Xcr1.cre (*B6-Xcr1^tm2Ciphe*)[54], *CD11c.cre*[31] and *Clec9A.cre*[36] mice were crossed to *Bcl6^fl/fl* mice[55]

(from Alexander Dent or JAX.org stock number #023727) to obtain DC-specific Bcl6-knockout

models. *Clec9A.Bcl6^KO* mice and *Clec9A.Bcl6^CONTROL* control mice were both maintained

homozygous for cre and additionally contained homozygous a *Rosa26^fl-Stop-fl-YFP* allele (JAX.org

stock number 006148). *CD11c.Bcl6* and *XCR1.Bcl6* models were maintained heterozygous for cre

and negative littermates were used as control controls. Additional strains used in this study were

B6.SJL-Ptprc[a] Pepc[b]/BoyJ (congenic C57BL/6.CD45.1 mice from the Jackson Laboratory), C57BL/6-

Tg (Tcra Tcrb)1100Mjb/J and B6.Cg-Tg(TcraTcrb)425Cbn/J (OT-I and OT-II mice from Jackson

Laboratory). In some experiments, OT-II (Thy1.1) mice were used as donors. Mice were group-

housed in individually ventilated cages and maintained under specific pathogen-free conditions.

Experimental groups were sex and age-matched. Male and female littermates were used

between 8 and 15 weeks of age, RNA-Seq was performed on female mice only. All animal

experiments were performed in accordance with European regulation and federal law of

Denmark and Germany, either approved by the Danish Animal Experiments Inspectorate, the

Regierung von Oberbayern or the Landesamt für Natur-, Umwelt und Verbraucherschutz NRW.

Cell preparation

Spleens, mLNs and pLN (axillary, brachial and inguinal) were digested in RPMI containing 5% FCS,

0.5 mg/ml Collagenase IV and 15 µg/ml DNase I for 45 min at room temperature. Red blood cells

in spleen and blood samples were subsequently lysed using red blood cell lysis buffer (ammonium

chloride, potassium bicarbonate, EDTA, and MiliQ water) for 3 minutes at room temperature. For

the OT-II Thy1.1 adoptive transfer experiments, splenic single cell suspensions were prepared

and analyzed as previously described[56].

Single cell suspensions of the intestinal LP were prepared as described previously[57]. Briefly, the

intestines were flushed with HBSS supplemented with FCS (10 %), Peyer's patches were removed

and epithelial cells and mucus were removed by incubating tissue in HBSS supplemented with

FCS (10%) and EDTA (2 mM) 3 times each for 15 min at 37 °C with continual shaking at 450 rpm.

Remaining tissue pieces were digested in R10 media (RPMI 1640 supplemented with FCS (10 %)

containing DNase I grade II (30 µg/ml) and Liberase TM (58 µg/ml, Roche) for 20 min at 37 °C with

a magnetic stirrer.

Lungs were perfused with PBS prior to resection. All lung lobes were cut into small pieces and

digested in R5 media (RPMI 1640 supplemented with FCS (5 %) containing DNase I grade II (30

µg/ml) and Liberase TM (50 µg/ml, Roche) for 45 min at 37 °C with a magnetic stirrer.

Intestinal LP and lung cell suspensions were filtered through 100 µm cell strainers (Fisher

Scientific) and leukocytes were enriched by density gradient centrifugation with 40 % / 70 %

Percoll (GE Healthcare) prior to further analysis.

Flow cytometry

Single-cell suspensions were stained for multicolor analysis with the indicated fluorochrome- or

biotin-conjugated antibodies after blocking with Fc block (purified anti-CD16/32). Antibodies

were diluted in flow cytometry buffer (DPBS, 2% FCS) and staining was performed for 30 min on

ice in the dark. LIVE/DEAD™ Fixable Near-IR Dead Cell Stain Kit (Invitrogen), Fixable Viability Dye

eFluor™-780 (Invitrogen) or DRAQ7 (BioLegend) were used for determining the viability of cells.

Lymphocytes from *C. rodentium* infection experiments were treated with rmIL-23 (40 ng/ml,

R&D) for 3 hours at 37°C with 5% CO_2, washed and further incubated with rmIL-23 (40 ng/ml),

PMA (50 ng/ml, Sigma Aldrich), Ionomycin (750 ng/ml, Sigma Aldrich) and BD GolgiStop™

(containing monensin at 1:1000 dilution, BD) for an additional 3.5 hours at 37 °C with 5% CO_2.

For intracellular staining, cells were fixed with the Foxp3 Fixation/Permeabilization kit from

eBioscience. Samples were acquired on a Fortessa X20, a LSRFortessa or a Symphony flow

cytometer (BD Biosciences) using FACSDiva software (BD Biosciences) and analyzed with

FlowJo software (Tree Star). The following antibodies were used: PE-Cy7/APC-Cy7 anti-mouse

NK1.1 (clone PK136), PE-Cy7/BUV737/APC-Cy7 anti-mouse CD3 (clone 145-2C11 or 17A2), PE-

Cy7/APC-Cy7 anti-mouse B220 (clone RA3-6B2), PE-Cy7/APC-Cy7 anti-mouse CD19 (clone

eBio1D3 or 6D5), PE-Cy7 anti-mouse CD64 (clone X54-5/7.1), APC-Cy7 anti-mouse TER119 (clone

TER-119), APC/BV711/PerCP-Cy5.5 anti-mouse CD11c (clone N418/HL3), FITC/BUV661 anti-

mouse MHC-II (clone M5/114.15.2), APC-Cy7/BUV737/FITC/APC-R700 anti-mouse CD11b (clone

M1/70), PE/BV421 anti-mouse XCR1 (clone ZET), BV421/BV786 anti-mouse CD103 (clone M290),

APC-Cy7 anti-mouse CD24 (clone M1/69), BV605 anti-mouse CD8α (clone 53-6.7), APC/PE anti-

73LR3 (clone 11F8), BUV395/V500 anti-mouse CD45 (clone 30-F11), PE-CF594 anti-mouse

CD172a (SIRPα) (clone P84), APC/PE-Cy7 anti-mouse ESAM (clone 1G8/ESAM), APC/PE anti-

mouse IRF4 (clone 3E4), PerCP eF710 anti-mouse IRF8 (clone V3GYWCH), PE anti-mouse Clec9a

(clone 7H11), FITC anti-mouse Vβ 5.1,5.2 (clone MR9-4), PerCp-Cy5.5 anti-mouse CD44 (clone

IM7), PE anti-mouse CD45.2 (clone 104), Alexa Fluor® 647 anti-human/mouse Bcl-6 (clone K112-

91), APC-Cy7 anti-mouse CD45.1 (clone A20), BUV395 anti-mouse CD4 (clone GK1.5), BV786 anti-

mouse IL-17A (clone TC11-18H10), FITC anti-mouse CD90.2 (clone 53-21), PerCP-Cy5.5 anti-

mouse IFNγ (clone XMG1.2), PE-eF610 anti-mouse RORγt (clone BD2), PE anti-mouse IL-22 (clone

1H8BWSR) , PE-Cy7 anti-mouse T-bet (clone 4B10), APC anti-mouse Foxp3 (clone FJK-16S), APC-

eF780 anti-mouse TCRβ (clone H57-597), biotin anti-mouse CXCR5 (clone L138D7), BV785 anti-

mouse CD4 (clone RM4-5), Pacific Blue™ anti-rat CD90/mouse CD90.1 (Thy-1.1) (clone OX-7),

PerCP/Cyanine5.5 anti-mouse CD19 (clone 6D5), PE-eFluor™ 610 anti-mouse CD279 (PD-1) (clone

J43), and streptavidin conjugated to BUV737 or APC.

Splenic dendritic cell sorting and cDNA library preparation for RNA-seq

Splenic DC were enriched using the Dynabeads Mouse DC Enrichment kit (Invitrogen) and sorted

on a BD FACS Melody to obtain cDC1 (B220⁻CD64⁻CD3⁻CD19⁻NK1.1⁻CD11c^{hi}MHCII⁺XCR1⁺) and

cDC2 (B220⁻CD64⁻CD3⁻CD19⁻NK1.1⁻CD11c^{hi}MHCII⁺CD11b⁺XCR1⁻). Sorted cells (2-4x10^4 per

sample) were resuspended in 350 μl RNeasy Lysis Buffer (QIAGEN), and RNA was extracted using

the QIAGEN RNeasy Mini kit (QIAGEN). RNA was washed, treated with DNase I (QIAGEN), and

eluted per the manufacturer's instructions. RNA-seq libraries were prepared by

combining the Nugen Ovation RNA-seq system with NuGEN's Ovation RNA-Seq System V2

(NuGEN Technologies). The amplified libraries were purified using AMPure beads, quantified by

qPCR, and visualized on an Agilent Bioanalyzer using BioA DNA High sensitivity (Agilent). The

libraries were pooled equimolarly and run on a HiSeq 2500 as single end reads of

50 nucleotide length.

RNA-sequencing data analysis

Sequencing reads were mapped to the mouse reference genome (GRCm38.85/mm10) using the

STAR aligner (v2.7.6a)[58]. Alignments were guided by a Gene Transfer Format (Ensembl GTF

GRCm38.101). Read count tables were normalized based on their library size factors using

DESeq2 (v1.30.1)[59], and differential expression analysis was performed with a filter of

log2foldChange >1 and a p-adjust value < 0.01. For selected gene signatures (cDC1 and cDC2 or

ESAM^hi and ESAM^lo DC2s) the 100 DEGs with highest normalized count values were selected, out

of which then the top 50 genes with highest absolute log2FoldChange were depicted in

heatmaps. RNA-seq reads aligned to genes from the GENCODE gene annotation were counted

using Ensembl's BiomaRt-v2.46.3. Heatmap visualizations were done using pheatmap v1.0.12.

Downstream statistical analyses and plot generation were performed in the R environment

(v4.0.5).[60] The online tool Venny 2.1 was used for the drawing of the Venn plots.[61]

Package PlotMA was used for visualizing of gene expression changes from two different

conditions (parameters: alpha = 0.01, type ="ashr").[62] Principal Component Analysis (PCA) plots

were drawn using PCAtools 2.2.0, with removeVar=0.01.[63] Volcano plots were drawn using the

function of ggpubr 0.4.0.[64] Gene Set Enrichment Analysis was performed using Gene Ontology R

package clusterProfiler 3.18.1 (category biological process) on DEGs obtained from DESeq2. The

p-values were adjusted using "BH" (Benjamini-Hochberg) correction and the cutoff was

0.01.[65] GO biological process terms were ranked by GeneRatio. The gene ratio is defined as the

number of genes associated with the term in DEGs divided by the number of DEGs. For KEGG

enrichment, DESeq2 derived DEGs were used.[66] Enrichments with a p-value from Fisher's exact

test ≤ 0.05 were considered significant.

Confocal Microscopy

Spleen and mLNs were harvested, washed with cold DPBS and placed in 4% PFA for 12 hours,

followed by an overnight wash with PBS-XG (PBS + 5% FCS + 0.2% Triton-X100 + 0.01%

Thimerosal). Samples were embedded in warm 4% low melting agarose. 70-100μm sections were

cut using a vibratome and stored until use. Samples were incubated with primary antibodies in

500ul PBS-XG in a 24 well plate overnight at 4°C under constant agitation (120rpm), followed by

at least 6 hours of washing in 5ml PBS-XG. The following antibodies were used: BV421 anti-mouse

B220/CD45R (clone RA3-6B2), APC anti-mouse TLR3 (clone 11F8), Alexa Fluor® 488 anti-mouse

CD11c (clone N418) and PE-CF594 anti-mouse CD172a (SIRPα) (clone P84). Sections were

mounted using ProLong Gold and images were acquired on a Zeiss LSM710 confocal microscope

equipped with 488-, 561-, and 633-nm lasers and a tunable two-photon laser. Digital pictures at

1,024 × 1,024-pixel density and 8-bit depth were acquired with ZEN software (Zeiss).

Cross-presentation assay

Naïve CD8 T cells from OT-I donors were enriched using the Dynabeads® Untouched™ Mouse T

Cells kit (Invitrogen) and stained in 1ml DPBS with CellTrace Violet dye (Invitrogen) for 10 min

at 37°C. 1×10^6 OT-I cells were transferred into recipients one day before i.p. injection of 5 x

10^6 cells/mouse of H-$2^{\beta m1}$ MEFs expressing truncated non-secreted OVA (OVA-MEF).[17] One day

after immunization, recipients were injected with FTY720 (20 µg/mouse; Cayman) in saline i.p..

Mice were sacrificed three days after immunization.

Induction of Tfh cells *in vivo*

For adoptive transfer experiments into *CD11c.Bcl6^{KO}* or control hosts, naïve CD4$^+$ T cells were

enriched from homozygous TCR-transgenic OT-II (CD45.1/.2) donors using the Dynabeads

Untouched™ Mouse CD4 Cells kit (Invitrogen). 1×10^5 OT-II cells were transferred intravenously

(i.v.) and one day later, recipients were immunized i.p. with ovalbumin grade V (OVA, 0.5 mg,

Sigma Aldrich) together with poly(I:C) (100 µg, Sigma-Aldrich). Controls received PBS alone and

mice were sacrificed three days after immunization. For adoptive transfer experiments into

Clec9a.Bcl6^{CONTROL} or *Clec9a.Bcl6^{KO}* hosts, naïve CD4$^+$ T cells were enriched from heterozygous

OT-II (Thy1.1) donors using the EasySep™ Mouse Naïve CD4$^+$ T Cell Isolation Kit (Stemcell

Technologies) according to the manufacturer's instructions. 2×10^5 OT-II cells were injected i.v.,

and one day later, recipient mice were immunized i.p. with a 1:1 mixture of 50 µg OVA (Sigma)

in PBS and Imject Alum (Thermo Fisher). Seven days later, spleens were dissected and analyzed

by flow cytometry.

***Citrobacter rodentium* infection**

The *C. rodentium* strain DBS100 (ATCC 51459; American Type Culture Collection) was used as

described[67] with some amendments. Briefly, *C. rodentium* cultures were grown in Luria-Bertani

(LB) medium to an OD_{600} of 1-1.5. Mice were orally inoculated with LB broth or $2x10^9$ CFU *C.*

rodentium, and their weight was monitored daily. Fecal samples were collected on indicated days

post infection, serially diluted in PBS and plated on Brilliance™ E. coli/coliform Agar (Thermo

Fisher). *C. rodentium* colonies, from at least three dilutions per sample, were enumerated and

normalized to weight of feces.

For histopathology, 0.5 cm of the distal colon was immediately placed in 10% formalin and

processed by the DTU Histology Core for hematoxylin and eosin (H&E) staining. Images were

acquired on a Pannoramic MIDI II scanner (3DHISTECH) and visualized with CaseViewer software

2.4 (3DHISTECH). The severity of the infection was assessed in a blinded manner.

Statistical Analysis

Statistical analyses were performed with Graphpad Prism 9.0 (Graphpad Software).

Supplementary Materials

Supplementary Figure 1: Phenotype and abundance of lung and blood cDC lacking Bcl6

Supplementary Figure 2: Quantification of spleen cDC in *Clec9a.Bcl6^{KO}* mice

Supplementary Figure 3: Intracellular Foxp3 and cytokine staining of colonic CD4+ T cells from

Citrobacter rodentium infected mice

Supplementary Table 1: Expression levels of top 50 DEGs between control cDC1 and control cDC2

Supplementary Table 2: DEGs between control and *XCR.Bcl6^{KO}* cDC1 and control and

CD11c.Bcl6^{KO} cDC2

Supplementary Table 3: Expression levels of top 50 DEGs between ESAM^{hi} and ESAM^{lo} cDC2

References

1.   Guilliams, M. *et al.* Dendritic cells, monocytes and macrophages: a unified nomenclature

based on ontogeny. *Nat. Rev. Immunol.* **14**, 571–578 (2014).

2.   Steinman, R. M. the Dendritic Cell System. *Annu. Rev. Immunol.* **9**, 271–296 (1991).

3.   Durai, V. & Murphy, K. M. Functions of Murine Dendritic Cells. *Immunity* **45**, 719–736

(2016).

4.   Allenspach, E. J., Lemos, M. P., Porrett, P. M., Turka, L. A. & Laufer, T. M. Migratory and

Lymphoid-Resident Dendritic Cells Cooperate to Efficiently Prime Naive CD4 T cells.

Immunity **29**, 795–806 (2008).

5.   Lewis, S. M., Williams, A. & Eisenbarth, S. C. Structure and function of the immune

system in the spleen. *Sci. Immunol.* **4**, (2019).

6.   Eisenbarth, S. C. Dendritic cell subsets in T cell programming: location dictates function.

Nat. Rev. Immunol. **19**, (2018).

7.   Calabro, S. *et al.* Differential Intrasplenic Migration of Dendritic Cell Subsets Tailors

Adaptive Immunity. *Cell Rep.* **16**, 2472–2485 (2016).

8.   Idoyaga, J., Suda, N., Suda, K., Park, C. G. & Steinman, R. M. Antibody to Langerin/CD207

localizes large numbers of CD8α + dendritic cells to the marginal zone of mouse spleen.

Proc. Natl. Acad. Sci. U. S. A. **106**, 1524–1529 (2009).

9.   Sichien, D. *et al.* IRF8 Transcription Factor Controls Survival and Function of Terminally

Differentiated Conventional and Plasmacytoid Dendritic Cells, Respectively. *Immunity* **45**,

626–640 (2016).

10.   Schiavoni, G. *et al.* ICSBP is essential for the development of mouse type I interferon-

producing cells and for the generation and activation of CD8α+ dendritic cells. *J. Exp.*

Med. **196**, 1415–1425 (2002).

11.   Hildner, K. *et al.* Batf3 deficiency reveals a critical role for CD8alpha+ dendritic cells in

cytotoxic T cell immunity. *Science* **322**, 1097–1100 (2008).

12.   Kashiwada, M., Pham, N. L. L., Pewe, L. L., Harty, J. T. & Rothman, P. B. NFIL3/E4BP4 is a

key transcription factor for CD8α+ dendritic cell development. *Blood* **117**, 6193–6197

(2011).

13.   Bagadia, P. *et al.* An Nfil3–Zeb2–Id2 pathway imposes Irf8 enhancer switching during

cDC1 development. *Nat. Immunol.* **20**, 1174–1185 (2019).

14.   Hacker, C. *et al.* Transcriptional profiling identifies Id2 function in dendritic cell

development. *Nat. Immunol.* **4**, 380–386 (2003).

15.   Cerovic, V. *et al.* Lymph-borne CD8α+ dendritic cells are uniquely able to cross-prime

CD8+ T cells with antigen acquired from intestinal epithelial cells. *Mucosal Immunol.* **8**,

38–48 (2015).

16.   Joeris, T. *et al.* Intestinal cDC1 drive cross-tolerance to epithelial-derived antigen via

induction of FoxP3+CD8+ Tregs. *Sci. Immunol.* **6**, (2021).

17.   Sancho, D. *et al.* Identification of a dendritic cell receptor that couples sensing of necrosis

to immunity. *Nature* **458**, 899–903 (2009).

18.   Ahrens, S. *et al.* F-actin is an evolutionarily conserved damage-associated molecular

pattern recognized by DNGR-1, a receptor for dead cells. *Immunity* **36**, 635–45 (2012).

19.   Zhang, J.-G. *et al.* The dendritic cell receptor Clec9A binds damaged cells via exposed
616   actin filaments. *Immunity* **36**, 646–57 (2012).

617   20.   Watchmaker, P. B. *et al.* Comparative transcriptional and functional profiling defines
618   conserved programs of intestinal DC differentiation in humans and mice. *Nat. Immunol.*
619   **15**, 98–108 (2014).

620   21.   Tussiwand, R. *et al.* Klf4 Expression in Conventional Dendritic Cells Is Required for T
621   Helper 2 Cell Responses. *Immunity* **42**, 916–928 (2015).

622   22.   Satpathy, A. T. *et al.* Notch2-dependent classical dendritic cells orchestrate intestinal
623   immunity to attaching-and-effacing bacterial pathogens. *Nat Immunol* **14**, 937–948
624   (2013).

625   23.   Persson, E. K. *et al.* IRF4 transcription-factor-dependent CD103(+)CD11b(+) dendritic cells
626   drive mucosal T helper 17 cell differentiation. *Immunity* **38**, 958–69 (2013).

627   24.   Williams, J. W. *et al.* Transcription factor IRF4 drives dendritic cells to promote Th2
628   differentiation. *Nat Commun* **4**, 2990 (2013).

629   25.   Scott, C. L. *et al.* The transcription factor Zeb2 regulates development of conventional
630   and plasmacytoid DC by repressing Id2. *J. Exp. Med.* jem.20151715 (2016).
631   doi:10.1084/jem.20151715

632   26.   Ichikawa, E. *et al.* Defective development of splenic and epidermal CD4+ dendritic cells in
633   mice deficient for IFN regulatory factor-2. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 3909–14
634   (2004).

635   27.   Wu, L. *et al.* RelB is essential for the development of myeloid-related CD8alpha- dendritic
636   cells but not of lymphoid-related CD8alpha+ dendritic cells. *Immunity* **9**, 839–47 (1998).

637   28.   Lewis, K. L. *et al.* Notch2 receptor signaling controls functional differentiation of dendritic
638   cells in the spleen and intestine. *Immunity* **35**, 780–791 (2011).

639   29.   Briseño, C. G. *et al.* Notch2-dependent DC2s mediate splenic germinal center responses.
640   *Proc. Natl. Acad. Sci. U. S. A.* **115**, 10726–10731 (2018).

30.　Fasnacht, N. *et al.* Specific fibroblastic niches in secondary lymphoid organs orchestrate

distinct Notch-regulated immune responses. *J. Exp. Med.* **211**, 2265–2279 (2014).

31.　Caton, M. L., Smith-Raska, M. R. & Reizis, B. Notch–RBP-J signaling controls the

homeostasis of CD8 − dendritic cells in the spleen . *J. Exp. Med.* **204**, 1653–1664 (2007).

32.　Lu, E., Dang, E. V., McDonald, J. G. & Cyster, J. G. Distinct oxysterol requirements for

positioning naïve and activated dendritic cells in the spleen. *Sci. Immunol.* **2**, eaal5237

(2017).

33.　Choi, J. & Crotty, S. Bcl6-Mediated Transcriptional Regulation of Follicular Helper T cells

(TFH). *Trends Immunol.* **42**, 336–349 (2021).

34.　Ohtsuka, H. *et al.* Bcl6 Is Required for the Development of Mouse CD4+ and CD8 +

Dendritic Cells. *J. Immunol.* **186**, 255–263 (2011).

35.　Bagadia, P. *et al.* Bcl6 -Independent In Vivo Development of Functional Type 1 Classical

Dendritic Cells Supporting Tumor Rejection . *J. Immunol.* ji1901010 (2021).

doi:10.4049/jimmunol.1901010

36.　Schraml, B. U. *et al.* Genetic tracing via DNGR-1 expression history defines dendritic cells

as a hematopoietic lineage. *Cell* **154**, 843–858 (2013).

37.　Lau, C. M. *et al.* Leukemia-associated activating mutation of Flt3 expands dendritic cells

and alters T cell responses. *J. Exp. Med.* **213**, 415–431 (2016).

38.　Baumjohann, D., Okada, T. & Ansel, K. M. Cutting Edge: Distinct Waves of BCL6

Expression during T Follicular Helper Cell Development. *J. Immunol.* **187**, 2089–2092

(2011).

39.　Goenka, R. *et al.* Cutting Edge: Dendritic Cell-Restricted Antigen Presentation Initiates the

Follicular Helper T Cell Program but Cannot Complete Ultimate Effector Differentiation. *J.*

Immunol. **187**, 1091–1095 (2011).

40.　Baumjohann, D. *et al.* Persistent Antigen and Germinal Center B Cells Sustain T Follicular

Helper Cell Responses and Phenotype. *Immunity* **38**, 596–605 (2013).

41.   Zhang, S. *et al.* Type 1 conventional dendritic cell fate and function are controlled by DC-
668   SCRIPT. *Sci. Immunol.* **6**, (2021).

669   42.   Zhang, T. T. *et al.* Dynamic expression of BCl6 in murine conventional dendritic cells
670   during in vivo development and activation. *PLoS One* **9**, 1–12 (2014).

671   43.   Durai, V. *et al.* Cryptic activation of an Irf8 enhancer governs cDC1 fate specification. *Nat.*
672   *Immunol.* **20**, 1161–1173 (2019).

673   44.   Grajales-Reyes, G. E. *et al.* Batf3 maintains autoactivation of Irf8 for commitment of a
674   CD8α+ conventional DC clonogenic progenitor. *Nat. Immunol.* **16**, 708–717 (2015).

675   45.   Seillet, C. *et al.* Nfil3 is required for the development of all innate lymphoid cell subsets.
676   *J. Exp. Med.* **211**, 1733–40 (2014).

677   46.   Kusunoki, T. *et al.* TH2 dominance and defective development of a CD8+ dendritic cell
678   subset in Id2-deficient mice. *J. Allergy Clin. Immunol.* **111**, 136–142 (2003).

679   47.   Dudziak, D. *et al.* Differential antigen processing by dendritic cell subsets in vivo. *Science*
680   **315**, 107–11 (2007).

681   48.   Gatto, D. *et al.* The chemotactic receptor EBI2 regulates the homeostasis, localization
682   and immunological function of splenic dendritic cells. *Nat. Immunol.* **14**, 446–453 (2013).

683   49.   Yi, T. & Cyster, J. G. EBI2-mediated bridging channel positioning supports splenic
684   dendritic cell homeostasis and particulate antigen capture. *Elife* **2**, e00757 (2013).

685   50.   Huang, C. *et al.* The BCL6 RD2 Domain Governs Commitment of Activated B Cells to Form
686   Germinal Centers. *Cell Rep.* **8**, 1497–1508 (2014).

687   51.   Schlitzer, A. *et al.* IRF4 transcription factor-dependent CD11b+ dendritic cells in human
688   and mouse control mucosal IL-17 cytokine responses. *Immunity* **38**, 970–83 (2013).

689   52.   Kedmi, R. *et al.* A RORγt+ cell instructs gut microbiota-specific Treg cell differentiation.
690   *Nature* (2022). doi:10.1038/s41586-022-05089-y

53.   Crotty, S., Johnston, R. J. & Schoenberger, S. P. Effectors and memories: Bcl-6 and Blimp-

1 in T and B lymphocyte differentiation. *Nat Immunol* **11**, 114–120 (2010).

54.   Wohn, C. *et al.* Absence of MHC class II on cDC1 dendritic cells triggers fatal

autoimmunity to a cross-presented self-antigen. *Sci. Immunol.* **5**, (2020).

55.   Hollister, K. *et al.* Insights into the Role of Bcl6 in Follicular Th Cells Using a New

Conditional Mutant Mouse Model. *J. Immunol.* **191**, 3705–3711 (2013).

56.   Baumjohann, D. & Ansel, K. M. Tracking Early T Follicular Helper Cell Differentiation In

Vivo. in *Methods in molecular biology* **1291**, 27–38 (2015).

57.   Johansson-Lindbom, B. *et al.* Functional specialization of gut CD103 + dendritic cells in

the regulation of tissue-selective T cell homing . *J. Exp. Med.* **202**, 1063–1073 (2005).

58.   Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21

(2013).

59.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).

60.   R Core Team (2020). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. (2020).

61.   Oliveros, J. C. Venny. An Interactive Tool for Comparing Lists with Venn's Diagrams.

62.   Stephens, M. False discovery rates: A new deal. *Biostatistics* **18**, 275–294 (2017).

63.   Blighe, K. and A. L. PCAtools: everything Principal Components

Analysis.https://github.com/kevinblighe/PCAtools. (2019).

64.   Alboukadel Kassambara. ggpubr:'ggplot2' Based Publication Ready Plots. (2020).

65.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for Comparing

Biological Themes Among Gene Clusters. *Omi. A J. Integr. Biol.* **16**, (2012).

66.   Qiu, Y.-Q. KEGG Pathway Database. in *Encyclopedia of Systems Biology* (eds. Dubitzky,

W., Wolkenhauer, O., Cho, K.-H. & Yokota, H.) 1068–1069 (Springer New York, 2013).

doi:10.1007/978-1-4419-9863-7_472

67.   Rizk, J. *et al.* The cIAP ubiquitin ligases sustain type 3 γδ T and innate lymphoid cells

718   during aging to allow normal cutaneous and mucosal responses. *bioRxiv* (2022).

719   doi:10.1101/2020.05.14.094334

Acknowledgements

We thank Katrine Fog Starup and Julien Vandamme for excellent technical support, the Lahl, Agace and Bekiaris laboratories for fruitful discussions, and Prof Allan Mowat for intellectual input and manuscript editing. *Bcl6^{fl/fl}* mice were kindly gifted by Alexander Dent and the OVA-MEF line was from Caetano Reis e Sousa. H&E sections were processed by Susanne Primdahl and histology data was generated using research infrastructure at DTU National Food Institute, including FOODHAY (Food and Health Open Innovation Laboratory, Danish Roadmap for Research Infrastructure).

Funding

This study was funded by the China Scholarship Council fellowship (HX), a Lundbeck Foundation Fellowship R215-2015-4100, the Ragnar Söderberg Foundation Fellowship in Medicine, Vetenskapsrådet 2020-01977, Cancerfonden 21 1826 Pj, the Novo Nordisk Foundation 18038, and the Crafoord Foundation (all KL), the European Research Council ERC-2016-STG-715182, Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Emmy Noether grant Schr 1444/1-1, Project-ID 360372040 – SFB 1335 (project 8) and 322359157-FOR2599-A03 (all BS), the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Emmy Noether Programme BA 5132/1-2 (252623821), an SFB 1054 Project B12 (210592381) (DB) and Germany's Excellence Strategy EXC2151 (390873048) (all to DB) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC2151 – 390873048 (AS), an Emmy Noether research grant (SCHL2116/1-1 to AS), the CRC1454 - INST 217/1034-1(Subproject P05) and a DFG Research Grant (SCHL2116/6-1) to AS.

Author Contributions:

Conceptualization: KL

Methodology: HX, IU, JH, HN, LB, IK, UM, LZ, AS, FP, BS, DB, KL

Investigation: HX, IU, JH, HN, LB, IK, UM, LZ, BS, DB, KL

Visualization: HX, IU, UM, KL

Funding acquisition: HX, DB, BS, KL

Project administration: KL

Supervision: FP, AS, BS, DB, KL

Writing – original draft: KL

Writing – review & editing: IU, BS, DB, KL

Competing Interests

The authors declare that they have no competing interests.

Data Availability

Raw RNA-sequencing data files are accessible through the NCBI SRA database under the

accession number PRJNA834905.

Figures:

Fig. 1: Phenotype and abundance of lymphoid tissue resident cDC lacking Bcl6

A. Left: Splenic XCR1$^+$ and XCR1$^-$CD11b$^+$ DC subsets in control (black), *XCR1.Bcl6^{KO}* (orange), and

766   *CD11c.Bcl6^{KO}* (red) mice. Plots show representative staining profiles of gated live, single lineage$^-$

767   (CD3, CD19, CD64, B220, NK1.1), MHCII$^+$, CD11c^{hi} splenocytes. Right: Absolute numbers of splenic

768   XCR1$^+$ and XCR1$^-$CD11b$^+$ DC from control, *XCR1.Bcl6^{KO}*, and *CD11c.Bcl6^{KO}* mice. Statistical analysis

769   by one-way ANOVA. * $P<0.01$ and ** $P<0.001$. Data points represent values from individual mice

770   pooled from 8-9 experiments and lines represent the medians.

771   B. Left: Flow cytometry plots showing resident XCR1$^+$ and XCR1$^-$CD11b$^+$ DC subsets in the pLNs and

772   mLNs of control, *XCR1.Bcl6^{KO}*, and *CD11c.Bcl6^{KO}* mice. Plots show representative staining profiles

773   of gated single live, CD45$^+$, lineage$^-$ (CD3, CD19, CD64, B220, NK1.1), CD11c^{hi}MHCII^{int} resident DC.

774   Right: Absolute numbers of resident cDC subsets in the pLN and mLN of control, *XCR1.Bcl6^{KO}*, and

775   *CD11c.Bcl6*[KO] mice. Statistical analysis by one-way ANOVA. * P<0.05. Data points represent values

776   from individual mice pooled from at least four experiments and lines represent the medians.

778   **Fig. 2: Phenotype and abundance of migratory cDC subsets lacking Bcl6**

779   A. Left: Flow cytometry plots showing migratory XCR1[+] and XCR1[−]CD11b[+] DC subsets in the pLN (top)

780   and mLN (middle) of control (black), *XCR1.Bcl6*[KO] (orange), and *CD11c.Bcl6*[KO] (red) mice. mLN

781   XCR1[−] DC are further sub-divided into CD103[+] and CD103[−] (bottom). Plots show representative

782   staining profiles of live single, CD45[+], lineage[−] (CD3, CD19, CD64, B220, NK1.1), CD11c[int]MHCII[hi]

783   migratory DC. Right: Absolute numbers of migratory cDC subsets from pLN and mLN of control,

784   *XCR1.Bcl6*[KO], and *CD11c.Bcl6*[KO] mice. Statistical analysis by one-way ANOVA. * P<0.05, ** P<0.01,

785   *** P<0.001 and **** P<0.0001. Data points represent values from individual mice pooled from

786   at least four experiments and lines represent the medians.

787   B. Left: Flow cytometry plots showing SILP XCR1[+] and XCR1[−]CD11b[+] DC subsets from control,

788   *XCR1.Bcl6*[KO], and *CD11c.Bcl6*[KO] mice. Plots show representative staining profiles of live, CD45[+],

789   lineage[−] (CD3, CD19, CD64, B220, NK1.1), MHCII[+], CD11c[hi] gated single cells.

790   Right: Absolute numbers of XCR1[+] and XCR1[−]CD11b[+] DC subsets from control, *XCR1.Bcl6*[KO], and

791   *CD11c.Bcl6*[KO] mice. Statistical analysis was performed by one-way ANOVA.* P<0.05 and ** P<0.01.

792   Data points represent values from individual mice pooled from three experiments and lines

793   represent the medians.

795   **Fig. 3: Lineage identity of splenic Bcl6-deficient cDC subsets**

796   A. Principal component analysis (PCA) plot of gene expression data obtained from bulk RNA-seq

797   using DESeq2. Color coded icons show cDC1 and cDC2 from the spleen of control, *XCR1.Bcl6*[KO],

798   and *XCR1.Bcl6*[KO] mice. Three replicates were included in each group.

799   B. Heatmap of selected highly expressed cDC1 or cDC2 associated genes in all RNA-Seq samples (50

800   most differentially expressed genes between cDC1 and cDC2 chosen from the 100 genes with

801   highest expression level among total differentially expressed genes (fold change $\geqslant$ 2, padj: 0.01)

802   in control samples). The color scale represents the row Z score.

803   C. Volcano plots showing the global transcriptional differences between splenic cDC2 and cDC1 in

804   control (left), *XCR1.Bcl6*[KO] (middle) and *CD11c.Bcl6*[KO] mice (right). Each circle represents one gene.

The log2FoldChange (LFC) is represented on the x-axis. The y-axis shows the –(log10 of the p adjust

value). A padj value of 0.01 and an absolute LFC of 1 are indicated by dashed lines. The

upregulated or downregulated genes are shown in red or blue dots respectively, and the top

ranked gene symbols with the lowest padj value were annotated in the plot.

D. VENN diagram showing significantly elevated differentially expressed genes (DEGs)

(|Log2foldchange|⩾1 and padj<0.01) between *XCR1.Bcl6^{KO}* cDC1 (red) and control cDC1 (blue),

and between *CD11c.Bcl6^{KO}* cDC2 (orange) and control cDC2 (green).

Fig. 4: Phenotype and gene expression profiles of Bcl6-deficient splenic cDC1

A. Flow cytometric analysis of cDC1 in spleen of control and *XCR1.Bcl6^{KO}* mice. Representative

histograms show indicated protein expression by control (black) and *XCR1.Bcl6^{KO}* (red) cDC1, with

FMO staining shown in grey. Pre-gating on cDC1 was performed as in 1A. Data are representative

of two independent experiments with 2-3 mice each.

B. Gene ontology (GO) term enrichment analysis of differentially expressed genes (DEGs) in

XCR1.Bcl6^{KO} and control cDC1. The p value calculated from GO term enrichment analysis is

represented on the x-axis and the y-axis shows the GO Biological Process (BP) terms.

C. KEGG (Kyoto Encyclopedia of Genes and Genomes) term enrichment analysis of DEGs in

XCR1.Bcl6^{KO} vs control cDC1. The GeneRatio value calculated from KEGG term enrichment analysis

is represented on the x-axis and the y-axis shows the KEGG enriched terms.

D. Heatmap of all genes contributing to the enriched KEGG pathways depicted in B. The color scale

represents the row Z score.

Fig. 5: Cross-presentation by cDC1 deficient in Bcl6

A. Representative flow cytometry plots showing CTV dilution and CD44 expression on OT-I cells

(gated on live, CD3^{+}CD8^{+}CD45.1^{+} single cells) in the spleen (top) and mLN (bottom) of control and

XCR1.Bcl6^{KO} recipient mice three days after i.p. immunization with heat-shocked OVA-MEFs. Data

are representative of three independent experiments with 2-5 mice each.

B. Total number of OT I cells in the spleen (right) and mLN (left) of immunized control and

XCR1.Bcl6^{KO} mice. Data are pooled from three independent experiments. Each dot represents one

mouse and lines represent the medians. Statistical analysis using Mann-Whitney test, *p<0.05.

Fig. 6: Phenotype and gene expression profiles of splenic Bcl6-deficient cDC2

A. Representative flow cytometric analysis of cDC2 in the spleen of control and *CD11c.Bcl6^{KO}* mice,

showing expression of CD11c vs MHCII (top panels) and XCR1 vs CD11b (lower panels) by live,

lineage$^−$ (CD3, CD19, CD64, B220, NK1.1) single cell-gated splenocytes. Data are representative of

at least 8 independent experiments with 1 – 3 mice each.

B. Representative flow cytometric analysis of ESAM^{hi} and ESAM^{lo} cDC2 in the spleen of control and

CD11c.Bcl6^{KO} mice. Cells were pre-gated as XCR1$^-$CD11b$^+$ cDC2 from live, lineage$^−$ (CD3, CD19,

CD64, B220, NK1.1) single cells. Data are representative of three independent experiments with

1 – 3 mice each.

C. Representative flow cytometric analysis of cDC2 in the spleen of control and *Clec9a.Bcl6^{KO}* mice,

showing expression of CD11c vs MHCII (top panels) and XCR1 vs CD11b (lower panels) by live,

lineage$^−$ (CD3, CD19, CD64, B220, NK1.1) single cell-gated splenocytes. Data are representative of

two independent experiments with 3 mice each.

D. Representative flow cytometric analysis of ESAM^{hi} and ESAM^{lo} cDC2 in the spleen of control and

Clec9a.Bcl6^{KO} mice. Cells were pre-gated as XCR1$^-$CD11b$^+$ cDC2 from live, lineage$^−$ (CD3, CD19,

CD64, B220, NK1.1) single cells (top). Data are representative of two independent experiments

with 3 mice each.

E. Same data as in D, but pre-gated on YFP$^+$ cells to enrich for cDC2 with a history of Clec9A-promoter

activity.

F. Representative flow cytometric analysis of expression of CD11c, 33D1, CD4, CD11b and IRF4 by

splenic XCR1$^−$CD11b$^+$ cDC2 from control (ESAM^{hi}: black, ESAM^{lo}: blue) or *CD11c.Bcl6^{KO}* mice

(orange). FMO controls are depicted in gray. Gates correspond to those depicted in Fig. 6B. Results

are representative of at least two (CD11c) or three (all others) independent experiments with 3

mice each.

Fig. 7: Consequences for germinal center reactions caused by Bcl6 deficiency in splenic cDC

A. Gene set enrichment plots following *CD11c.Bcl6^{KO}* vs control cDC2 in an ESAM^{hi} cDC2 related gene

set (top) vs an ESAM^{lo} cDC2 related gene set (bottom) (ESAM^{hi} and ESAM^{lo} gene sets derived from

GSE76132). NES = normalized enrichment score. FDR q-value: 0.0.

B.   The top 50 genes associated with spleen ESAM[hi] cDC2 (left) or ESAM[lo] cDC2 (right) were derived

from GSE76132. Heatmaps show the expression of these genes in the RNA-Seq analysis of cDC2

from control and *CD11c.Bcl6^{KO}* spleen. The color scale represents the row Z score (Expression Z-

scale).

C.   Confocal microscopy of spleens from *CD11c.Bcl6^{KO}* and control mice showing expression of TLR3

(grey: upper left quadrant), CD11c (green: upper right quadrant), Sirpα (red: lower left quandrant)

and B220 (blue: lower right quadrant). Scale bar=50μm. Orange arrows indicate marginal zone

bridging channels. The images shown are from one out of four (*CD11c.Bcl6^{KO}*) or three (control)

replicates derived from individual mice.

D.   Schematic depiction of immunization schedules.

E.   Top: Representative flow cytometric analysis of early Tfh cell induction amongst OT-II cells in the

spleen of control and *CD11c.Bcl6^{KO}* mice 3 days after i.p. immunization with OVA and poly(I:C) or

with PBS as a control. Tfh were gated as live, CD4^+Vβ5.2^+CD45.1^+CD45.2^+CD44^+Bcl6^+CXCR5^+ and

the results are representative of three experiments with 3-5 mice each. Each experiment

contained at least one unimmunized (PBS) control per genotype. Bottom: Numbers of total CD44^+

OT-II T cells (left panel), together with absolute numbers (middle panel) and proportions (right

panel) of OVA-specific Tfh cells in the spleens of control and *CD11c.Bcl6^{KO}* mice 3 days after i.p.

immunization with OVA and poly(I:C) or with PBS as a control. Statistical analysis was performed

by one-way ANOVA. * P<0.05, ** P<0.01, *** P<0.001 and **** P<0.0001. Data points represent

values from individual mice pooled from three experiments with 3-5 mice each in OVA/poly(I:C)

immunized groups. Each experiment contained at least one untreated control per genotype; lines

represent the medians.

F.   Representative flow cytometric analysis of splenic OT-II cells transferred into *Clec9a.Bcl6^{CONTROL}*

and *Clec9a.Bcl6^{KO}* recipients 7 days after i.p. immunization with OVA/alum. Tfh cells were gated

as CXCR5^+PD-1^+ cells among live CD4^+Thy1.2^+CD19^− lymphocytes. Statistical analysis was

performed by Mann-Whitney test, ** P<0.01. Each symbol represents one biological replicate

(n=5), lines represent the median. Data are representative of two independent experiments.

Fig. 8: *Citrobacter rodentium* infection in mice with Bcl6 deletion in cDC

A.   Left: Flow cytometry plots showing cLP XCR1$^+$ and XCR1$^-$CD11b$^+$ DC subsets gated on all DC (live,

CD45$^+$, lineage$^-$ (CD3, CD19, CD64, B220, NK1.1), MHCII$^+$, CD11c^{hi} single cells. Right: XCR1$^-$ DC are

further sub-divided into CD103$^+$ and CD103$^-$ cDC2s from control and *CD11c.Bcl6^{KO}* mice.

B.   Absolute numbers (top) and proportions (bottom) of cLP XCR1$^+$, XCR1$^-$CD103$^+$CD11b$^+$ and XCR1$^-$

CD103$^-$CD11b$^+$ DC subsets from control and *CD11c.Bcl6^{KO}* mice. Data points represent values from

individual mice pooled from 3 experiments and lines represent the medians. Statistical analysis

using Mann-Whitney test, ** P<0.01 and *** P<0.001.

C.   Percentage of weight loss of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice. Each

symbol represents the mean weight loss of all mice within the group at the indicated time point

post infection versus their starting weight and error bars represent SD; statistical analysis using

two-way ANOVA (and Tukey's multiple comparison test) with Geisser-Greenhouse correction, *P

< 0.05 and **P < 0.01 (top left). Colon lengths in mock and infected mice at 9 or 21 days post

inoculation; statistical analysis by one-way ANOVA, ** P<0.01, **** P<0.0001 (bottom left). *C.*

rodentium titers in feces measured as colony-forming units (CFU) of infected control and

CD11c.Bcl6^{KO} mice at indicated days post infection; statistical analysis using Mann-Whitney test,

not significant (right). All data points represent values from individual mice, lines represent the

median and data is pooled from 2-3 experiments with 3-5 mice per mock/*C. rodentium* infected

group.

D.   H&E-stained sections of the distal colon of mock and *C. rodentium* infected control and

CD11c.Bcl6^{KO} mice 21 days post inoculation. Scale bar, 100 μm. Symbols: # epithelial lining; ¤

leukocyte infiltration; % goblet cell destruction; & villus length. Representative of 4-5 mice per

group from one experiment.

E.   Numbers of total CD4$^+$ T cells, IL-17A$^+$IL-22$^-$, IL-17A$^+$IL-22$^+$, IL-17A$^+$IFNγ$^-$, IL-17A$^+$IFNγ$^+$ and Foxp3$^+$

CD4$^+$ T cells in the cLP of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice at day 9

post inoculation. Statistical analysis performed by one-way ANOVA, * P<0.05, ** P<0.01, ***

P<0.001 and **** P<0.0001.

F.   Numbers of total CD4$^+$ T cells, IL-17A$^+$IL-22$^-$, IL-17A$^+$IL-22$^+$, IL-17A$^+$IFNγ$^-$, IL-17A$^+$IFNγ$^+$ and Foxp3$^+$

T cells in the cLP of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice at day 21 post

inoculation. * P<0.05, ** P<0.01 and **** P<0.0001, by one-way ANOVA.

Supplementary Figure 1

A. Left: Flow cytometry plots show XCR1$^+$ and XCR1$^-$CD11b$^+$ DC subsets in the lungs of control,

XCR1.Bcl6^{KO} and *CD11c.Bcl6^{KO}* mice. The plots show representative staining profiles of live, CD45$^+$,

lineage$^-$ (CD3, CD19, CD64, B220, NK1.1), MHCII$^+$, CD11c^{hi} gated single cells. Right: Absolute

numbers of cDC subsets in the lungs of control, *XCR1.Bcl6^{KO}*, and *CD11c.Bcl6^{KO}* mice. Statistical

analysis by one-way ANOVA.** P<0.01, **** P<0.0001. Data points represent values from

individual mice pooled from 2-3 experiments and lines represent the medians.

B. Left: Representative flow cytometry plots of blood XCR1$^+$ and XCR1$^-$CD11b$^+$ DC subsets from

control, XCR1.Bcl6^{KO} and CD11c.Bcl6^{KO}. Shown are representative staining profiles of live, CD45$^+$,

lineage$^-$ (CD3, CD19, CD64, B220, NK1.1), MHCII$^+$, CD11c^{hi} gated single cells. Right: Absolute

numbers of cDC1 and cDC2 in blood of control, XCR1.Bcl6^{KO}, and CD11c.Bcl6^{KO} mice. Statistical

analysis by one-way ANOVA, significance at * P<0.05. Data points are normalized to volume of

blood drawn and represent values from individual mice pooled from 4 experiments; lines

represent the median cell numbers from each genotype.

Supplementary Figure 2

A. Relative proportions of XCR1$^+$, XCR1$^-$CD11b$^+$ESAM^{hi} and XCR1$^-$CD11b$^+$ESAM^{lo} subsets in the spleen

of Clec9a.*Bcl6^{KO}* (top) and CD11c.*Bcl6^{KO}* (bottom) mice. Cells were pre-gated on single live, CD45$^+$,

lineage$^-$ (CD3, CD19, TER119, B220, NK1.1), MHCII$^+$, CD11c^{hi}. Data points represent values from

individual mice pooled from 2-3 experiments and lines represent the medians. Statistical analysis

using Mann-Whitney test, *** P<0.001 and **** P<0.0001.

B. Numbers of XCR1$^+$, XCR1$^-$CD11b$^+$ESAM^{hi} and XCR1$^-$CD11b$^+$ESAM^{lo} subsets in the spleen of control,

Clec9a.*Bcl6^{KO}* normalized to volume of suspension (top) and CD11c.*Bcl6^{KO}* shown as absolute

numbers (bottom). Data points represent values from individual mice pooled from 2-3

experiments and lines represent the medians. * P<0.05, *** P<0.001 and **** P<0.0001, by

Mann-Whitney test.

Supplementary Figure 3

A. Representative flow cytometry plots showing Foxp3$^+$ CD4$^+$ T cells (gated on live, CD45$^+$,

CD3$^+$TCRβ$^+$ singlets) in the cLP of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice

at day 9 (left) or day 21 (right) post inoculation.

B. Representative flow cytometry analysis showing IL-17A$^+$IL-22$^+$ CD4$^+$ T cells (gated on live, CD45$^+$,

CD3$^+$TCRβ$^+$ singlets) in the cLP of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice

at day 9 (left) or day 21 (right) post inoculation.

C. Representative flow cytometry plots showing IL-17A$^+$IFNγ$^+$ CD4$^+$ T cells (gated on live, CD45$^+$,

CD3$^+$TCRβ$^+$ singlets) in the cLP of mock and *C. rodentium* infected control and *CD11c.Bcl6^{KO}* mice

at day 9 (left) or day 21 (right) post inoculation.

Figure 1

988   **Figure 2**

990 **Figure 3**

994   **Figure 4**

996   **Figure 5**

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012   **Figure 6**

1014   **Figure 7**

1016   **Figure 8**

1017

1018

Supplementary Figure 1

1019

1020

1021

1022

Supplementary Figure 2

A   Relative abundance of cDC subsets in the spleen in Clec9a.Bcl6^(KO) and CD11c.Bcl6^(KO) mice

B   Absolute count of cDC subsets in the spleen in Clec9a.Bcl6^(KO) and CD11c.Bcl6^(KO) mice

1023

Supplementary Figure 3