Statistical integration of multi-omics and drug screening data from cell lines

Said el Bouhaddani[1,*†], Matthias Höllerhage[2†], Hae-Won Uh[1], Claudia Moebius[3], Marc Bickle[4], Günter Höglinger[2,5,6‡], Jeanine Houwing-Duistermaat[1,7‡]

1 Dept. Data science & Biostatistics, UMC Utrecht, Utrecht, Netherlands
2 Department of Neurology, Hannover Medical School, Hannover, Germany
3 Roche Institute for Translational Bioengineering, Basel, Switzerland
4 Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany
5 Department of Neurology, Ludwig-Maximilians-Universität, Munich, Germany
6 German Center for Neurodegenerative Diseases, Munich, Germany
7 Dept. of Mathematics, Radboud University, Nijmegen, Netherlands

†These authors contributed equally to this work.
‡These authors also contributed equally to this work.
* sbouhad2@umcutrecht.nl

Abstract

Data integration methods are used to obtain a unified summary of multiple datasets. For multi-modal data, we propose a computational workflow to jointly analyze datasets from cell lines. The workflow comprises a novel probabilistic data integration method, named POPLS-DA, for omics data.

The workflow is motivated by a study on synucleinopathies where transcriptomics, proteomics, and drug screening data are measured in affected LUHMES cell lines and controls. The aim is to highlight potentially druggable pathways and genes involved in synucleinopathies. First, POPLS-DA is used to prioritize genes and proteins that best distinguish cases and controls. For these genes an integrated interaction network is constructed, where the drug screen data is incorporated to highlight druggable genes and pathways in the network. Finally functional enrichment analyses are performed to identify clusters of synaptic and lysosome-related genes and proteins targeted by the protective drugs.

We found that *HSPA5*, a member of the heat shock protein 70 family, was one of the most targeted genes by the validated drugs, in particular by AT1-blockers. *HSPA5* and AT1-blockers have been previously linked to α-synuclein pathology and Parkinson's disease, showing the relevance of our findings.

Our computational workflow identified new directions for therapeutic targets for synucleinopathies. An implementation based on R and markdown is freely available online.

Author summary

We present a computational workflow that combines the analysis of different types of data measured in cell line studies with non-overlapping samples. We apply the workflow to measurements of gene expression, protein abundances, and a screening of a wide range of FDA-approved drugs. These different types of data are obtained from

LUHMES brain cells and jointly analyzed to discover new treatment options in synucleinopathies, such as Parkinson's disease. Our workflow includes a new probabilistic method, named POPLS-DA. POPLS-DA combines the analysis of the genes and proteins to pinpoint a set of relevant genes and proteins that can distinguish affected and non-affected cells. Furthermore, we constructed a network that connects together the relevant genes and proteins that interact with each other. We incorporate the drug screening data to highlights which part of the network is relevant to the disease and druggable. Through additional analysis of the functionality, we discovered that the genes and proteins that are targeted by protective drugs share relevant properties, namely they are synaptic and lysosome related genes. Notably, we found that a specific type of drugs, namely AT1-blockers such as Telmisartan, are protective and target the network of relevant genes and proteins. These drugs are approved by the FDA and readily available to further investigate its potential in treating synucleinopathies. We further found that a gene named HSPA5, a member of the heat shock protein 70 family, is highly targeted by the protective drugs. This gene has been linked to Parkinson's disease in previous scientific literature. Our computational workflow and the implementation in R and markdown are freely available online.

Introduction

Nowadays, studies often include measurements of several omics datasets as well as other types of data. Particularly in cell line studies, datasets such as genomics, transcriptomics, and proteomics can be generated in a reproducible and standardized experimental system. Experiments using cell lines have also been set up to identify new drug targets and measure their efficacy. How to summarize these different datasets is an ongoing research topic. A joint analysis of all data is most efficient but also challenging since it might need a model describing the relationship between the datasets. Another way is to integrate datasets by using functional information about the top features of the datasets. This paper is motivated by a cell line study with proteomics, transcriptomics, and drug compound screening data. These data have been analyzed separately, but this did not yield an overarching overview of involved genes. New computational tools are needed to obtain an overview of the relationship between the omics data, and their interplay with drugs in the context of MSA. Here, we present a workflow that analyze the omics data jointly and integrate these with the drug compounds screening data using functional information.

For the analysis of multiple homogeneous omics data from cell lines, the multivariate integrative (MINT) method, which is an extension of partial least squares discriminant analysis (PLS-DA) [1] for multiple omics levels, is available. This method estimates linear combinations of features that maximally discriminate between experimental groups across all omics datasets. However, MINT does not include omics-specific variation, while it has been shown that modeling omics-specific variation yields better results [2,3] in the case of heterogeneous datasets. Since transcriptomics and proteomics data are different (e.g. technical platform and scale), including data-specific components in the model can be beneficial. For clinical and epidemiological studies, these approaches for heterogeneous multi-omics are available, namely the algorithmic and the probabilistic two-way orthogonal partial least squares (O2PLS [4] resp. PO2PLS [5]). For small datasets, a probabilistic approach is preferred since algorithmic approaches tend to overfit [3]. However, these methods are suited for dataset measured on the same samples and cannot directly be applied to data obtained from non-overlapping samples such as cell lines. We propose a new approach, probabilistic OPLS-DA (POPLS-DA) which is suited for heterogeneous multi-omics data from non-overlapping cell lines.

In our motivating study, transcriptomic and proteomic data are measured in

LUHMES cells. LUHMES cells are derived from human embryonic dopaminergic cells [6] and are used to investigate the neurobiological processes underlying neurodegeneration, specifically synucleinopathies. Synucleinopathies are neurodegenerative disorders where a pathological aggregation of α-synuclein is present. The most common synucleinopathy is Parkinson's disease (PD) and multiple system atrophy (MSA). It is assumed that aggregated forms of α-synuclein are toxic, leading to cell death and neurodegeneration. Integrative multi-omics approaches can provide insights into the biological processes, molecular functions, and interactions underlying the diseases. The transcriptomics and proteomics data were measured in cells overexpressing α-synuclein or GFP as a control [7]. Furthermore, to identify drugs that protect against α-synuclein-induced cell death, LUHMES cells were used to screen 1600 FDA-approved drugs. The primary screening was performed in triplicates in two different concentrations (3 μM and 10 μM) and revealed 53 drugs that reduced cell death in LUHMES cells overexpressing α-synuclein in at least one of the screening runs [8]. Here, we propose to integrate the drug compound screening data with the top features from the omics data analysis by using functional information on the genes targeted by the drugs and the interactome. Thus we aim to perform an integrated analysis of the three datasets to obtain a holistic overview of gene deregulation in α-synuclein overexpressing cells and to obtain a unified list of the most relevant features (genes and drug compounds) [9]. Note that such a list can comprise features that were not found with single dataset analyses, because of a lack of statistical evidence when analyzed separately.

Our contribution is firstly a novel data integration workflow comprising a joint analysis of the omics data and an integration approach using functional information to combine the top omic features and the drug compound screening results. Secondly a new probabilistic method named POPLS-DA for joint analysis of omics datasets in cell lines is developed. Thirdly application of the workflow including POPLS-DA to transciptomics, proteomics and drug compound screening datasets in LUMES cell lines. We perform dose-response testing of all 53 compounds to validate the protective efficacy of the drug compounds. Then we analyse the three datasets using our novel integrated computational workflow. We apply POPLS-DA to integrate the two omics datasets and identify relevant genes and proteins that distinguished the two experimental conditions (α-synuclein versus control protein overexpression). Based on these genes and proteins, a protein-protein interaction (PPI) network was constructed and the genes/proteins that are targeted by the drugs were identified. This approach highlighted genes in the network that are associated with α-synuclein overexpression and targeted by validated drugs, and therefore are potential targets for a novel therapy for synucleinopathies.

Results

We first present the results for the drug screen validation. Next, we show the results of our computational procedure applied to the multi-omics and drug screening data from the LUHMES cells. Finally enrichment analyses are presented.

Validation of the protective efficacy of the FDA-approved drugs

Previously, we identified 53 FDA-approved drugs as being protective against αSyn-induced toxicity in a primary drug screening [8]. Now, using a dose-response analysis, 41 of these 53 compounds were validated as being protective. The maximum effective concentrations appeared to range from 2.5 μM to 80 μM. Fig 1 shows a schematic overview and results of the dose-response testing.

Figure 1. Workflow of FDA-approved drug screening and dose-response validation. 1,600 FDA-approved drugs were screened in two different concentrations (3 μM, 10 μM) in three runs each in an αSyn toxicity cell model. 53 compounds were identified as being protective in at least one run (A). Timeline of dose-response testing (B). In the first round, concentrations from 0.6 nM to 20 μM of each compound were investigated. 36 compounds could be confirmed as being protective. Of these, 20 already reached their maximal protective potential and were selected not to be tested in a higher concentration. The remaining 16 compounds were tested again in concentrations between 2.4 and 80 μM. All of these were confirmed as being protective in at least one concentration. Of the 17 compounds that were not protective in the first round dose-response testing, 8 were already toxic and considered not being protective. The remaining 9 were tested again in concentration from 2.4 to 80 μM. 5 of these compounds were protective in higher concentrations. In total, 41 could be confirmed as protective against αSyn induced toxicity (C).

POPLS-DA integration of transcriptome and proteome data

For the integrated analysis of the pre-preocessed transcriptomics and proteomics datasets, we used POPLS-DA. For each dataset we used two joint components representing the relationship between the two datasets and two data specific

components. The number of components was determined by a scree plot of the eigenvalues of the row-wise concatenated datasets (Supplementary material). Based on a scree plot of the squared effect sizes per gene (given by the squared elements of $W\beta$), presented in Fig 2, 200 genes/proteins were retained for further analysis. A list of all genes/proteins and their weights are shown in the supplementary material.

Of the total variance in the transcriptomics and proteomics data, 21 and 18 percent, respectively, were explained by the integrative parts. Omics-specific variation accounted for around 27 percent of the total variation. The selected relevant genes/proteins appeared to be able to distinguish between the two experimental groups (Fig 2 and Supplementary material). We performed 400 permutations (see supplementary material) of the experimental group labels to test the null-hypothesis of no relation between the experimental groups and the genes/proteins. We found that, under the null hypothesis, the probability of achieving the same discrimination (100% accuracy) as observed in the data was 0.0025 (one in 400 permutations).

Figure 2. Selection of relevant genes/proteins and their ability to discriminate. The *left panel* shows the sorted squared effect size per gene given by the squared elements of $W\beta$. The 200 relevant genes/proteins correspond to the 'elbow', which is visually determined and approximately where the curve crosses the green vertical line. The *right panel* shows boxplots of score predictions based on the selected 200 relevant genes/proteins. The dots, representing individual samples, are added with a horizontal 'jitter' to reduce overlap. The transcriptomics and proteomics samples are colored orange and blue, respectively. Abbreviations: aSyn: α-synuclein, mRNA: transcriptome, Prot: proteome.

Integrated interaction network based on multi-omics and drug data

Fig 3 shows the integrated interaction network constructed using String-DB. The network had significantly more interactions than expected from an arbitrary subset of 200 genes. The network enrichment p-value was smaller than 10^{-15}. From the 41 drugs that were validated to reduce α-synuclein-induced toxicity [8], we found 27 targeting at least one gene in the integrated interaction network. The efficacy of the maximal effective dose of each of these compounds is shown in Table 1.

The *HMGCR*, *NQO1*, *HSPA5*, and *YBX1* genes were most targeted by the validated drugs, namely by 20, 18, 17, and 17 drugs, respectively.

Table 1 shows for each compound the number of direct neighbors it targets in the network. Dexibuprofen, a non-steroidal anti-inflammatory drug (NSAID), targets the highest number of direct neighbors (N=60), followed by Imatinib (N=47), which is acting on the tyrosine kinase enzyme and is involved in apoptosis. The most effective compounds were Telmisartan with an improvement of the cell survival of α-synuclein-overexpressing cells by 33.16% at a concentration of 5 μM, followed by Risperidone which improved cell survival by 28.58% at 10 μM.

Figure 3. String-DB and clustering analyses of the top 200 genes/proteins.
Figures are numbered from left to right, from top to bottom. In panel (a), a network of
interactions between the top 200 genes/proteins (estimated with POPLS-DA) was
constructed using String-DB. Each node is a gene, and a connection between genes
indicates evidence for a biologically plausible link. Text-mining was excluded as an
evidence source, and a medium confidence threshold was used. For genes that were
(indirectly) targeted by a drug compound, a green 'halo' is drawn. The intensity of the
green color is proportional to the number of drug compounds for which the gene was an
(indirect) target. In panel (b), the interaction network from (a) was clustered using the
MCL clustering algorithm from the String-DB website. The edges between the clusters
are removed for visual aid. In panel (c), an interaction network is shown for a druggable
subset of the top 200 genes/proteins, consisting of 116 genes that were (indirectly)
targeted by an FDA-approved drug compound. In panel (d), an interaction network of
top genes in the "Parkinson's disease" DisGeNet term was constructed using String-DB.
Text mining as evidence was included here.

Drug name	Direct neighbors	Most effective concentration	Cell survival	P-value
Dexibuprofen	60	80 μM	118.29%	<0.0001
Imatinib	47	20 μM	126.04%	<0.0001
Amiodarone	44	80 mM	108.05%	0.002
Risperidone	41	10 μM	128.58%	<0.0001
Bepridil	36	80 μM	115.89%	<0.0001
Astemizole	26	1.25 μM	106.49%	0.02
Telmisartan	24	5 μM	133.16%	<0.0001
Amlodipine	21	10 μM	116.42%	<0.0001
Benazepril	18	80 μM	117.96%	<0.0001
Repserpine	18	80 μM	107.64%	0.01
Dipyridamole	16	40 μM	118.09%	<0.0001
Trifluopromazine	16	40 μM	114.30%	<0.0001
Nefazodone	15	40 μM	115.08%	<0.0001
Trimipramine	15	20 μM	125.54%	<0.0001
Flunarizine	14	20 μM	104.38%	0.03
Quinacrine	13	5 μM	119.07%	0.008
Pentoxyverine	10	1.25 μM	109.69%	0.002
Dicyclomine	8	20 μM	120.15%	<0.0001
Doxazosin	8	40 μM	116.12%	<0.0001
Ajmaline	7	20 μM	108.94%	<0.0001
Ifenprodil	7	2.5 μM	116.74%	0.0003
Lomerizine	7	80 μM	110.35%	<0.0001
Pentamidine	6	2.5 μM	119.86%	0.003
Guanfacine	5	5 μM	117.01%	<0.0001
Dyclonine	4	5 μM	125.69%	<0.0001
Tropisetron	3	20 μM	131.63%	<0.0001
Clemastine	1	10 μM	117.52%	0.0004

Table 1. FDA-approved drugs protective against α-synuclein-induced toxicity with direct neighbors of the top 200 genes. For each drug, the number of direct interactors is shown. The most effective concentration is the concentration at which each compound had the strongest protective effect. Cell survival is expressed as percentage compared to αSyn-overexpressing cells that were treated with DMSO as control at the maximum effective concentration. P-Values of testing the dose-reponse were calculated with an ANOVA with Dunnett's post-hoc test for multiple comparisons.

The network was split into sub-clusters using the MCL clustering algorithm on the integrated interaction network, see Fig 3. Several clusters consist of genes that are involved in neurogenerative disorders. The largest cluster is a highly connected hub of 22 genes, primarily ribosomal. Ribosomal proteins have been reported to mediate neurodegeneration in PD [10]. Another cluster of five genes includes the top two genes, *MTHFD2* and *CBS*. Overexpression of the *CBS* (cystathionine-beta-synthase) gene was protective in a PD toxin model in rats [11]. The network also contains clusters with genes involved in neurological development, e.g. *TFRC* (transferrin receptor), and neuronal adhesion and growth, e.g. *TNC* (tenascin C). Further, four genes encoding intermediate filaments, such as *NES* (nestin), are clustered together. They are primarily responsible for cell adhesion [12]. Furthermore, a cluster of three genes from the solute carrier family was found (*SLC3A2*, *SLC7A5*, and *SLC7A6*). These genes may play a role in the transport across the blood-brain barrier.

Enrichment analyses of relevant genes and proteins ₁₂₈

In Table 2, the most significant pathways enriched in the top 200 genes are shown. The ₁₂₉
GO terms that contained "extracellular" and "cytoplasm" were the most significant. ₁₃₀
Genes with these annotations are involved in vesicular transport and signaling [13]. ₁₃₁
Nine of the ten terms were from the "cellular compound" GO category. ₁₃₂

Term	Ontology	Genes in set	FDR
cytoplasm	CC	178	1.28e-19
extracellular exosome	CC	71	4.67e-17
extracellular vesicle	CC	71	5.51e-16
extracellular organelle	CC	71	5.51e-16
intracellular anatomical structure	CC	190	7.62e-12
cell junction	CC	61	7.62e-12
vesicle	CC	88	2.30e-11
extracellular space	CC	81	7.09e-11
protein binding	CC	179	7.65e-11
cytosol	MF	100	1.48e-10

Table 2. Gene ontology enrichment analysis of the top 200 genes. Out of 1732
genes/proteins, the 200 genes/proteins with the highest weight (estimated using
POPLS-DA) were used to perform GO enrichment analysis. All three GO ontologies
were considered: cellular compound (CC), biological process (BP), and molecular
function (MF). For each term, a p-value was computed using Fisher's exact test with
the whole genome as background and corrected for multiple testing. The ten most
significant terms are shown, ranked by their GO enrichment false discovery rate (FDR)
adjusted p-value. The column "Genes in set" shows how many of the top 200
genes/proteins were annotated with the respective term.

Table 3 shows the five diseases for which the most significant enrichment was found ₁₃₃
in the top 200 genes. The second term was "Parkinson's disease" and contained 48 ₁₃₄
genes. Fig 3 shows the PPI network for genes/proteins in this disease term. Note that ₁₃₅
the *HSPA5* gene, which was one of the most targeted by the FDA drug compounds, is ₁₃₆
within this network. ₁₃₇

Term	Genes in set	FDR
Neoplasms	142	2.84e-07
Parkinson's Disease	48	1.94e-06
Carcinogenesis	100	3.36e-06
Malignant neoplasm of prostate	75	2.54e-05
Neoplasm Metastasis	98	2.89e-05

Table 3. Top five disease enrichment clusters based on DisGeNet. The top
200 relevant genes/proteins were analyzed using DisGeNet, a disease-gene database.
The BEFREE database was used. For each term, a p-value was calculated using a
Fisher exact test and FDR adjusted. The terms are ranked by this adjusted p-value.

Finally, enrichment analyses were performed for the subset of 116 genes/proteins ₁₃₈
that were direct neighbors of genes targeted by at least one of the drugs. Fig 3 shows ₁₃₉
the protein-protein interaction network for this subset. The corresponding network ₁₄₀
enrichment p-value was smaller than 10^{-10}. Also, the most significant GO terms agreed ₁₄₁
with the GO analysis of the total set of 200 genes/proteins (Supplementary material). ₁₄₂

Discussion

In this study we proposed a computational workflow to integrate omics and drug screen data from cell lines. A novel method, PO2PLS-DA, for analysis of multiple omics datasets in non-overlapping cell lines was developed. The workflow is illustrated by jointly analyzing multi-omics and drug screening data from LUHMES cells overexpressing α-synuclein and those overexpressing GFP as a control protein. Transcriptomics and proteomics data were first integrated with POPLS-DA. We found 200 relevant genes/proteins based on an elbow plot of the squared effect size of each gene (Fig 2). These genes/proteins appeared to perfectly discriminate between the two experimental groups. Using a dose-response approach, 41 FDA-approved drugs were validated to be protective against α-synuclein-induced toxicity. An integrated interaction network was constructed based on the 200 relevant genes/proteins by identifying direct neighbors of the 41 drugs in the network. Interactome analyses revealed several genes and pathways linked to PD, the most common synucleinopathy.

The drugs that targeted the highest number of direct neighbors were Dexibuprofen and Imatinib. Both drugs were previously found to lower the risk of PD in epidemiological studies [14, 15]. Interestingly, independent of the screening, we previously investigated the neuroprotective potential of imatinib as a macroautophagy-stimulating drug. Despite the protective effect, imatinib in our model led to a reduction of the expression of the dopaminergic markers tyrosine hydroxylase (TH) and dopamine transport (DAT) [16]. Therefore, further investigation would be necessary before using this drug in PD patients. However, other drugs we identified could be readily investigated in PD. Telmisartan, an AT1-blocker, which is used for arterial hypertension, was the most effective drug with an improvement of cell survival >30% at a comparably low concentration of 5μM. Since there are no major restrictions in PD other than a blood pressure lowering effect, Telmisartan could be a promising candidate for a neuroprotective therapy. Similarly, also Benazepril, an ACE-inhibitor and also blood-pressure lowering drug could be investigated with similar caution. Interestingly, AT1-blockers and ACE inhibitors have been shown to reduce the risk of falls in PD, and this association was not related to blood pressure [17]. Furthermore, the use of AT1-blockers and ACE inhibitors as potential neuroprotective drugs was previously discussed, because these drugs were protective in PD animal models, most likely due to an anti-oxidative effect [18]. In line with that, our data further emphasize that it would be promising to further investigate AT1-blockers and ACE-inhibitors in PD.

Other drugs, including Risperidone and Trifluopromazine, are antipsychotic drugs with an anti-dopaminergic effect. Though validated to be protective, they can lead to drug-induced parkinsonism [19] and are contraindicated in PD. Also, Flunarizine, a drug used in migraine prophylaxis can lead to drug-induced parkinsonism [20, 21].

The genes that were targeted by the majority of the drugs were *HMGCR*, *NQO1*, *HSPA5*, and *YBX1*. Note that *HMGCR* (3-Hydroxy-3-Methylglutaryl-CoA reductase) is the main target of statins. *NQO1* (NAD(P)H quinone dehydrogenase 1) has multiple functions and has been linked to Alzheimer's disease [22]. The *HSPA5* gene (heat shock protein family A (Hsp70) member 5) encodes a heat shock protein of the HSP70 family. It is found to be involved in maintaining the correct folding behavior of proteins [23]. HSPA5, also known as glucose-regulated protein 78 (GRP78), is located on the membrane of the endoplasmic reticulum and is essential for the unfolded protein response [24]. Furthermore, it was previously shown that overexpression of this protein reduced α-synuclein toxicity to dopaminergic cells in a rat PD model [25]. Interestingly, a previous study showed that GRP78 levels were reduced in the temporal cortex and cingulate gyrus of PD patients compared to healthy controls but markedly increased in PD patients with dementia and patients with dementia with Lewy bodies [26]. In line

with that, our data that show an interaction of drugs that protect against α-Syn with this protein support that GRP78 plays an important role in α-Syn toxicity and deliver further evidence that GRP78 could be an promising target for neuroprotective therapies of PD. The *YBX1* gene (Y-box binding protein 1) is a cold shock domain protein, predominantly localized to neurons and essential for brain development [27]. Furthermore, several clusters were observed in the network involving ribosomal proteins and intermediate filaments. Enrichment analyses identified several terms involving "extracellular" and "vesicles".

Using DisGeNet, the second most significantly enriched disease term was "Parkinson's disease", with 48 genes from our list of relevant genes/proteins. The other disease terms involved neoplasms and carcinogenesis. Note that previous studies suggested shared molecular mechanisms between PD and cancer [28–30].

In a previous study on these data [7], the authors identified significant deregulation of synaptic proteins, in particular, *STXBP1*, *STX1B*, *SYT1*, and *FYN*. In our POPLS-DA integration, we found three of these genes: *STXBP1*, *SYT1*, and *FYN*, ranked 16, 18, and 72, respectively. We identified two additional synaptic genes in the top 200, namely *STX12*, encoding syntaxin-12, as well as *SYT11*, encoding synaptotagmin-11. Single nucleotide polymorphisms in *SYT11* have been linked to PD in a previously performed GWAS [31]. Furthermore, a previous study showed that synaptotagmin-11 plays a role in Parkin-related parkinsonism. Additionally, it was previously reported that lysosome related proteins, *SCARB2*, *CTSB*, *SMPD1*, were differentially regulated upon overexpression of α-synuclein in LUHMES cells [7]. From our top genes/proteins, 22 were annotated with the lysosome GO term. Among these 22 genes are the previously found *SCARB2* at position 7 in our top genes/proteins, as well as *SLC7A5* and *SLC3A2* at position 5 and 6, respectively. The authors pointed to lysosome membrane protein 2 expression (encoded by *SCARB2*) as a therapeutic target to increase lysosomal glucocerebrosidase activity to promote α-synuclein clearance. Note that one of our clusters in Fig 3 consists of genes encoding lysosome related proteins (i.e., *SCARB2*, *AP2A2*, *ATP6V1C1*, *IGF2R*, and *TFRC*) that are also direct neighbors of the FDA-approved drugs.

The data used in this study were acquired from a human (LUHMES) cell line, where each dataset was measured in a different sample. Most data integration methods require multiple datasets measured in the same samples to calculate correlations across omics datasets from which joint components can be derived. These methods are sometimes referred to as horizontal integration approaches, where the data are laid out next to each other with the samples across the rows. In the present paper, we proposed a framework for vertical integration, where different groups of samples (representing different omics data) are laid down below each other with the same (type of) variables across the columns. Currently, POPLS-DA is restricted to the case that no missing variables are present across each group of samples.

Several databases are available with information on the interaction between genes and other molecular variables (interactomes) [32]. We chose String-DB since it was found to perform well with regard to disease prediction [33]. However, they also showed that other databases might comprise complementary information. Currently, to the best of our knowledge, no method can utilize all information across different interactomes. Information bias is another unsolved problem when using these databases: less studied diseases have less coverage in these databases [34]. Since synucleinopathies are less common and less studied (compared to, e.g., neoplasms), the bioinformatics interpretation of our results are likely to be biased towards well-studied disease. Indeed, in DisGeNet, PD has only 2078 annotated genes, while neoplasms have more than 10000 genes. This could explain that our enrichment analysis based on DisGeNet (Table 3) showed many genes annotated with "neoplasms".

The POPLS-DA method can integrate more than two datasets to distinguish between experimental conditions in cell lines if the variables in these datasets can be mapped to the same unit, i.e. genes. For example, SNPs from a GWAS study can be integrated by mapping them to genes. Here the distance of a SNP to a gene or its transcription factor binding site [35] can be used. For methylation, CpG sites are commonly mapped to genes based on their distance to a transcription start site [36]. Since, typically, multiple SNPs and CpG sites are mapped to the same gene, an aggregation step is needed to obtain one measurement per gene. For example, the first principal component of the SNPs or CpG sites corresponding to a gene can be taken as the final aggregated measurement.

The data from the FDA-approved drugs were integrated and projected onto the interaction network of the 200 relevant genes obtained from the transcriptomics and proteomics data analyses. An alternative approach is to simultaneously analyze the transcriptomics and proteomics data together with data on the validated compounds, i.e. drug-augmented data integration. It has been shown that including such information via penalties into data integration analysis yields robust results [37]. Such a joint model can pinpoint genes that can discriminate the two groups while targeted by the compounds, improving the clinical relevance of the list of top genes.

Conclusion   We validated 41 FDA-approved drugs to be protective against α-synuclein-induced toxicity in dose-response analyses. We applied a novel data integration approach to combine these drug data with experimental multi-omics data. We identified a set of 200 relevant genes/proteins that discriminated between samples overexpressing α-synuclein and controls, as well their drug targets. Some of the drugs (e.g. an ACE-inhibitor and an AT1-blocker) could be readily investigated in PD patients. These findings can potentially be used to develop therapeutic targets for Parkinson's disease, multiple system atrophy, and other synucleinopathies.

Materials and methods

Processing samples and measuring data

We describe the screening of 1600 FDA-approved drugs and the acquisition of proteome and transcriptome data in brief. Details are available elsewhere [7,8].

FDA-approved drug compound validation   For the dose-response testing, LUHMES cells were plated in flasks coated with poly-L-lysine (0.1 mg/ml) and fibronectin (5 μg/ml) in growth medium (DMEM/F12 with 1% N2 supplement and 0.04 μl/ml basic fibroblast growth factor). After 24h, the medium was changed to differentiation medium (DMEM/F12 with 1% N2-supplement, 1 μg/ml tetracycline, 0.49 μg/ml N6,2'-O-dibutyryladenosine 3',5'-cyclic monophosphate, 2 ng/ml glial derived neurotrophic factor). After another 24 hours, the cells were transduced with adenoviral vectors leading to overexpression of human wild-type α-synuclein. On the day after transduction, viral vectors were washed away, and the cells were replated on 384-well multi-well plates and treated with the different drugs solved in dimethyl sulfoxide. For validation, compounds were tested in concentrations between 0.6 nM to 20 μM. Compounds that were not protective or did not reach a protective plateau were further tested in higher concentrations up to 80 μM. The dose-response data were analyzed using an ANOVA with Dunnett's post hoc test to correct for multiple comparison with GraphPad Prism version 9.4 for Windows 64-bit (GraphPad Software, San Diego, CA, USA).

Figure 4. Workflow of the multi-omics integration approach. (1) To link the two datasets, the transcripts and proteins were mapped to their Entrez identifier, referred to as genes/proteins. (2) To identify the relevant genes/proteins that discriminate between the two experimental groups (α-synuclein overexpression vs control), POPLS-DA was applied. (3) The prioritized genes/proteins were integrated with the FDA-approved drug screening data using a direct neighbor approach. As an illustration, in the figure under "Direct Neighbor approach" are two drug target genes (green circles), which are neighbors of genes identified in (2) (blue circles). (4) Bioinformatics analyses: interactome, GO and DisGeNet enrichment analyses analyses. Protein-protein interaction networks were built using String-DB where each node is a gene. The genes that were direct neighbors of a drug target were highlighted with a green halo, where its color intensity is proportional to the number of direct neighbors. Abbreviations: aSyn: α-synuclein, GO: gene ontology , FDA: Food and Drug Administration.

Transcriptome and proteome data acquisition and processing For the transcriptome and proteome data, cells were plated directly in differentiation medium and transduced with adenoviral vectors leading to overexpression of either human wild-type α-synuclein or GFP, respectively. After 24 hours, the remaining viral vectors were removed. Samples were collected four days after transduction. Transcriptome data

were measured with Illumina HumanHT-12_V3 bead chips (Illumina, San Diego, CA, USA). Proteome data were measured by liquid chromatography-mass spectrometry (LC-MS/MS). The transcription probes were filtered based on the Illumina detection p-value, where probes were removed when all samples had a p-value above 0.05. A variance filter was applied where probes were removed if the inter-quantile range (IQR) of a probe did not exceed the 0.5-th quantile of all IQRs. Probes that could not be mapped to an Entrez identifier were removed.

After pre-processing and filtering, the transcriptomics and proteomics datasets contained 15660 transcription probes and 2577 proteins that could be mapped to Entrez identifiers. 2292 of these identifiers were present in both datasets. We excluded the *SNCA* measurements from the data analysis.

For the transcriptomics data, we analyzed three samples from LUHMES cells overexpressing α-synuclein, and three samples from LUHMES cells expressing GFP as a control protein. For proteomics, the numbers were equal to nine. The correlation of the transcripts with the α-synuclein overexpressing cells versus the control cells was taken as a reference; if this correlation was negative, the corresponding protein feature was multiplied by minus one. The measurements for both datasets were scaled to have zero mean and unit variance.

Computational workflow for multi-modal data

Our workflow for integrated analysis is depicted in Fig. 4. To summarize, the workflow begins with preprocessing the multi-omics data into a data matrix where variables are mapped to the same nomenclature. We then propose a joint model, POPLS-DA, to analyze the multi-omics data in terms of the experimental groups.

Next, we prioritize the most relevant genes based on POPLS-DA and integrate them with validated drugs using a 'direct neighbor' approach. This step involves identifying drugs that significantly reduce the toxicity of α-synuclein aggregation and retrieving the list of genes targeted by these drugs from DrugBank. We then add their direct neighbors in the String-DB database to the gene lists for each compound.

Finally, we construct an integrated gene-gene interaction network based on the relevant genes and perform functional enrichment analysis. The network is built with String-DB, and the nodes are colored according to the number of compounds targeting each gene. We use the Markov cluster algorithm to identify sub-clusters of genes/proteins. Gene ontology pathway and disease enrichment analyses are performed using the `goana` function from the `limma` package in R and DisGeNet, respectively.

POPLS-DA model for multi-omics cell line data For the joint analysis of the transcriptomics and proteomics datasets, a novel statistical data integration method, probabilistic orthogonal partial least squares discriminant analysis (POPLS-DA) is developed. Details can be found in the supplementary materials.

Let x_k be a random vector of size p for $k = 1, \ldots, K$. In our data example, $k = 1, 2$ represents the 2292 transcriptomics and proteomics data, respectively. Similarly, let y_k be a univariate random variable underlying the outcome. In our data example, y_k are the two experimental conditions (one for cells overexpressing α-synuclein, and zero for the controls) for $k = 1, 2$. In the POPLS-DA model, the group status and omics data are linked via latent variables u_k of size r, with r much smaller than the data dimensions. These latent variables represent joint parts of the two omics datasets that are correlated with the group status. To model omics variables that contribute to the variation in the omics dataset but do not play a role in distinguishing between the two experimental groups, specific components v_k were added to the model. Residual variation is modeled by noise terms e_k, and ϵ_k. All random variables are assumed to be zero-mean normally

distributed. The mathematical model for POPLS-DA can be written as 347

$$x_k = u_k W^{\mathrm{T}} + v_k P_t^{\mathrm{T}} + e_k, \qquad y_k = u_k \beta + \epsilon_k, \qquad k = 1, \ldots, K. \qquad (1)$$

The matrix W contains the joint loadings, and the matrices P_k contain the specific 348
loadings for each gene in each component. The vector β contains the regression 349
coefficients of y_k on the joint components u_k. The vector $W\beta$ represents effect sizes for 350
the association of each variable with the outcome. 351

The observed datasets are assumed to consist of N_k i.i.d. samples from the 352
POPLS-DA model (1), with $k = 1, \ldots, K$. Note that the sample size per k is allowed to 353
differ. Based on these data, the parameters of POPLS-DA are estimated with 354
memory-efficient maximum likelihood. Technical details are given in the supplementary 355
materials. 356

The numbers of components here are denoted by r_k, respectively. The number of 357
joint and specific components, r and r_k, have to be specified and can be based on scree 358
plots [38] of the eigenvalues of the datasets. To select a set of relevant genes/proteins, 359
the obtained effect sizes $W\beta$ are ranked and plotted. The elbow criterion [38] is used to 360
determine the threshold between relevant and irrelevant genes/proteins. 361

Computational workflow for integrating multi-omics and drug screening 362
data   Based on the POPLS-DA model (1) for multiple omics datasets, we propose a 363
computational procedure for integration and analysis of cell line data. First, the 364
multi-omics data are analyzed with POPLS-DA to prioritize genes that discriminate 365
cases from controls. The relevant genes/proteins identified by POPLS-DA are further 366
analyzed using several databases on protein-protein interactions (PPI), drug targets, 367
gene ontology, and gene-disease association. 368

To construct a integrated interaction network of the relevant genes/proteins, we 369
employed the String-DB [39] website and the `STRINGdb` package in R [40]. The network 370
was based on curated, computational, and experimental data sources, text mining was 371
excluded. A medium evidence threshold (0.4) was used to define an interaction (edge in 372
the network). 373

To integrate the FDA-approved drug screening data with the obtained integrated 374
interaction network, we first identified drugs that significantly reduced the toxicity of 375
α-synuclein aggregation. We implemented a automated R procedure using 376
DrugBank [41] to retrieve the list of genes targeted by these drugs. Using the 377
`string_db$get_neighbors` function, the direct neighbors of these targeted genes in the 378
String-DB database were added to the gene lists for each compound. Then, the number 379
of compounds targeting each gene in the α-Syn-PPI network was determined. Nodes 380
(genes and proteins) in the network were colored according to this number using the 381
"payload" function on the String-DB website. 382

To identify sub-clusters of genes/proteins in the integrated interaction network, the 383
Markov cluster algorithm (MCL) [42] on the String-DB website was used. The inflation 384
parameter was set to two. 385

Finally, gene ontology (GO) pathway and disease enrichment analyses were 386
performed. For GO, the `goana` function from the `limma` package [43] in R was used, 387
with the whole human genome as background. For disease enrichment analysis, 388
DisGeNet [44], a disease-gene association database implemented in the `disgenet2r` R 389
package, was used. Here, the "BEFREE" text mining database was used for the 390
disease-gene enrichment score as it contains more annotations regarding MSA and PD. 391

Our computational workflow, including POPLS-DA, is available as an R markdown 392
file automating most of the tasks detailed above. After the datasets are loaded into 393
memory, the file can be executed and an html file is generated with results and 394

visualizations. The workflow is available on
github.com/selbouhaddani/MultiOmicsCompWorkflow.

Supporting information

**S1 Supporting information. Supplementary materials for POPLS-DA and
data analysis.** The mathematical details and proofs for POPLS-DA are given, as well
as additional results and figures for the data analysis.

Acknowledgments

All authors were funded by ERA-Net E-Rare JTC 2018 (MSA-omics)
[40-44000-98-2006/ 90030376507]. JD was supported by European Union's Horizon 2020
(IMforFUTURE) [721815] and the EU funded Cost action DYNALIFE [CA21169]. GH
was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) under Germany's Excellence Strategy within the framework of the Munich
Cluster for Systems Neurology (EXC 2145 SyNergy) [390857198] and within the
Hannover Cluster RESIST (EXC 2155) [39087428], DFG [HO2402/18-1] MSAomics, the
Bavarian Ministry for Education, Culture, Science and Art (ForIPS) [8810001412],
Niedersächsisches Ministerium für Wissenschaft und Kunst [ZN3440.TP]: REBIRTH –
Forschungszentrum für translationale regenerative Medizin, VolkswagenStiftung
(Niedersächsisches Vorab), Petermax-Müller Foundation (Etiology and Therapy of
Synucleinopathies and Tauopathies). MH and GH received funding from
ParkinsonFonds Deutschland (Hypothesis-free compound screening in a new human
neuronal model of wild type alpha-synuclein-induced cell death).

References

1. Ståhle L, Wold S. Partial least squares analysis with cross-validation for the
   two-class problem: A Monte Carlo study. J Chemom. 1987;1(3):185–196.
   doi:10.1002/cem.1180010306.

2. Trygg J, Wold S. O2-PLS, a two-block (X-Y) latent variable regression (LVR)
   method with an integral OSC filter. J Chemom. 2003;17(1):53–64.
   doi:10.1002/cem.775.

3. el Bouhaddani S, Uh HW, Jongbloed G, Hayward C, Klarić L, Kiełbasa SM, et al.
   Integrating omics datasets with the OmicsPLS package. BMC Bioinformatics.
   2018;19(1):371. doi:10.1186/s12859-018-2371-3.

4. el Bouhaddani S, Houwing-Duistermaat J, Salo P, Perola M, Jongbloed G, Uh
   HW. Evaluation of O2PLS in Omics data integration. BMC Bioinformatics.
   2016;17(S2):S11. doi:10.1186/s12859-015-0854-z.

5. el Bouhaddani S, Uh HW, Jongbloed G, Houwing-Duistermaat J. Statistical
   integration of heterogeneous omics data: Probabilistic two-way partial least
   squares (PO2PLS). J R Stat Soc Ser C (Applied Stat.
   2022;doi:10.1111/rssc.12583.

6. Lotharius J. Progressive Degeneration of Human Mesencephalic Neuron-Derived
   Cells Triggered by Dopamine-Dependent Oxidative Stress Is Dependent on the
   Mixed-Lineage Kinase Pathway. J Neurosci. 2005;25(27):6329–6342.
   doi:10.1523/JNEUROSCI.1746-05.2005.

7. Höllerhage M, Stepath M, Kohl M, Pfeiffer K, ho Chua OW, Duan L, et al. Transcriptome and Proteome Analysis in LUHMES Cells Overexpressing Alpha-Synuclein. Front Neurol. 2022;13(April):1–19. doi:10.3389/fneur.2022.787059.

8. Höllerhage M, Moebius C, Melms J, Chiu WH, Goebel JN, Chakroun T, et al. Protective efficacy of phosphodiesterase-1 inhibition against alpha-synuclein toxicity revealed by compound screening in LUHMES cells. Sci Rep. 2017;7(1):11469. doi:10.1038/s41598-017-11664-5.

9. Ritchie MD, Holzinger ER, Li R, Pendergrass SA, Kim D. Methods of integrating data to uncover genotype–phenotype interactions. Nat Rev Genet. 2015;16(2):85–97. doi:10.1038/nrg3868.

10. Martin I, Kim JW, Lee BD, Kang HC, Xu JC, Jia H, et al. Ribosomal Protein s15 Phosphorylation Mediates LRRK2 Neurodegeneration in Parkinson's Disease. Cell. 2014;157(2):472–485. doi:10.1016/j.cell.2014.01.064.

11. Yin Wl, Yin Wg, Huang Bs, Wu Lx. Neuroprotective effects of lentivirus-mediated cystathionine-beta-synthase overexpression against 6-OHDA-induced parkinson's disease rats. Neurosci Lett. 2017;657:45–52. doi:10.1016/j.neulet.2017.07.019.

12. Perrot R, Eyer J. Intermediate Filaments in Neurodegenerative Diseases. InTech; 2013. Available from: `https://www.intechopen.com/chapters/44544http://www.intechopen.com/books/neurodegenerative-diseases/intermediate-filaments-in-neurodegenerative-diseases`.

13. Kao CY, Papoutsakis ET. Extracellular vesicles: exosomes, microparticles, their parts, and their targets to enable their biomanufacturing and clinical applications. Curr Opin Biotechnol. 2019;60:89–98. doi:10.1016/j.copbio.2019.01.005.

14. Rees K, Stowe R, Patel S, Ives N, Breen K, Clarke CE, et al. Non-steroidal anti-inflammatory drugs as disease-modifying agents for Parkinson's disease: evidence from observational studies. Cochrane Database Syst Rev. 2011;(11). doi:10.1002/14651858.CD008454.pub2.

15. Kumar M, Kulshrestha R, Singh N, Jaggi AS. Expanding spectrum of anticancer drug, imatinib, in the disorders affecting brain and spinal cord. Pharmacol Res. 2019;143:86–96. doi:10.1016/j.phrs.2019.03.014.

16. Höllerhage M, Fussi N, Rösler TW, Wurst W, Behrends C, Höglinger GU. Multiple molecular pathways stimulating macroautophagy protect from alpha-synuclein-induced toxicity in human neurons. Neuropharmacology. 2019;149:13–26. doi:10.1016/j.neuropharm.2019.01.023.

17. Laudisio A, Lo Monaco MR, Silveri MC, Bentivoglio AR, Vetrano DL, Pisciotta MS, et al. Use of ACE-inhibitors and falls in patients with Parkinson's disease. Gait Posture. 2017;54:39–44. doi:10.1016/j.gaitpost.2017.02.007.

18. Perez-Lloret S, Otero-Losada M, Toblli JE, Capani F. Renin-angiotensin system as a potential target for new therapeutic approaches in Parkinson's disease. Expert Opin Investig Drugs. 2017;26(10):1163–1173. doi:10.1080/13543784.2017.1371133.

19. López-Sendón JL, Mena MA, de Yébenes JG. Drug-induced parkinsonism in the elderly: incidence, management and prevention. Drugs Aging. 2012;29(2):105–18. doi:10.2165/11598540-000000000-00000.

20. Teive HAG, Troiano AR, Germiniani FMB, Werneck LC. Flunarizine and cinnarizine-induced parkinsonism: a historical and clinical analysis. Parkinsonism Relat Disord. 2004;10(4):243–5. doi:10.1016/j.parkreldis.2003.12.004.

21. Lin W, Lin CL, Hsu CY, Wei CY. Flunarizine Induced Parkinsonism in Migraine Group: A Nationwide Population-Based Study. Front Pharmacol. 2019;10:1495. doi:10.3389/fphar.2019.01495.

22. Simidjievski N, Bodnar C, Tariq I, Scherer P, Andres Terre H, Shams Z, et al. Variational Autoencoders for Cancer Data Integration: Design Principles and Computational Practice. Front Genet. 2019;10:1205. doi:10.3389/fgene.2019.01205.

23. Lyon MS, Milligan C. Extracellular heat shock proteins in neurodegenerative diseases: New perspectives. Neurosci Lett. 2019;711:134462. doi:10.1016/j.neulet.2019.134462.

24. Ibrahim IM, Abdelmalek DH, Elfiky AA. GRP78: A cell's response to stress. Life Sci. 2019;226:156–163. doi:10.1016/j.lfs.2019.04.022.

25. Gorbatyuk MS, Shabashvili A, Chen W, Meyers C, Sullivan LF, Salganik M, et al. Glucose regulated protein 78 diminishes α-synuclein neurotoxicity in a rat model of Parkinson disease. Mol Ther. 2012;20(7):1327–37. doi:10.1038/mt.2012.28.

26. Baek JH, Mamula D, Tingstam B, Pereira M, He Y, Svenningsson P. GRP78 Level Is Altered in the Brain, but Not in Plasma or Cerebrospinal Fluid in Parkinson's Disease Patients. Front Neurosci. 2019;13:697. doi:10.3389/fnins.2019.00697.

27. Bobkova NV, Lyabin DN, Medvinskaya NI, Samokhin AN, Nekrasov PV, Nesterova IV, et al. The Y-Box Binding Protein 1 Suppresses Alzheimer's Disease Progression in Two Animal Models. PLoS One. 2015;10(9):e0138867. doi:10.1371/journal.pone.0138867.

28. Tacik P, Curry S, Fujioka S, Strongosky A, Uitti RJ, van Gerpen JA, et al. Cancer in Parkinson's disease. Parkinsonism Relat Disord. 2016;31:28–33. doi:10.1016/j.parkreldis.2016.06.014.

29. Ejma M, Madetko N, Brzecka A, Guranski K, Alster P, Misiuk-Hojło M, et al. The Links between Parkinson's Disease and Cancer. Biomedicines. 2020;8(10):416. doi:10.3390/biomedicines8100416.

30. Filippou PS, Outeiro TF. Cancer and Parkinson's Disease: Common Targets, Emerging Hopes. Mov Disord. 2021;36(2):340–346. doi:10.1002/mds.28425.

31. International Parkinson Disease Genomics Consortium, Nalls MA, Plagnol V, Hernandez DG, Sharma M, Sheerin UM, et al. Imputation of sequence variants for identification of genetic risks for Parkinson's disease: a meta-analysis of genome-wide association studies. Lancet (London, England). 2011;377(9766):641–9. doi:10.1016/S0140-6736(10)62345-8.

32. Rigden DJ, Fernández XM. The 2022 Nucleic Acids Research database issue and the online molecular biology database collection. Nucleic Acids Res. 2022;50(D1):D1–D10. doi:10.1093/nar/gkab1195.

33. Mosca E, Bersanelli M, Matteuzzi T, Di Nanni N, Castellani G, Milanesi L, et al. Characterization and comparison of gene-centered human interactomes. Brief Bioinform. 2021;00(October 2020):1–16. doi:10.1093/bib/bbab153.

34. Haynes WA, Tomczak A, Khatri P. Gene annotation bias impedes biomedical research. Sci Rep. 2018;8(1):1362. doi:10.1038/s41598-018-19333-x.

35. Hu X, Kim H, Stahl E, Plenge R, Daly M, Raychaudhuri S. Integrating Autoimmune Risk Loci with Gene-Expression Data Identifies Specific Pathogenic Immune Cell Subsets. Am J Hum Genet. 2011;89(4):496–506. doi:10.1016/j.ajhg.2011.09.002.

36. Bibikova M, Barnes B, Tsan C, Ho V, Klotzle B, Le JM, et al. High density DNA methylation array with single CpG site resolution. Genomics. 2011;98(4):288–295. doi:10.1016/j.ygeno.2011.07.007.

37. Gu Z, el Bouhaddani S, Pei J, Houwing-Duistermaat J, Uh HW. Statistical integration of two omics datasets using GO2PLS. BMC Bioinformatics. 2021;22(1):131. doi:10.1186/s12859-021-03958-3.

38. Mardia KV, Kent JT, Bibby JM. Multivariate analysis. Academic Press; 1979.

39. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res. 2019;47(D1):D607–D613. doi:10.1093/nar/gky1131.

40. R Core Team. R: A Language and Environment for Statistical Computing; 2014. Available from: http://www.r-project.org/.

41. Wishart DS, Knox C, Guo AC, Shrivastava S, Hassanali M, Stothard P, et al. DrugBank: a comprehensive resource for in silico drug discovery and exploration. Nucleic Acids Res. 2006;34(Database issue):D668–D672. doi:10.1093/nar/gkj067.

42. Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection of protein families. Nucleic Acids Res. 2002;30(7):1575–84. doi:10.1093/nar/30.7.1575.

43. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 2015;43(7):e47–e47. doi:10.1093/nar/gkv007.

44. Piñero J, Ramírez-Anguita JM, Saüch-Pitarch J, Ronzano F, Centeno E, Sanz F, et al. The DisGeNET knowledge platform for disease genomics: 2019 update. Nucleic Acids Res. 2020;48(D1):D845–D855. doi:10.1093/nar/gkz1021.