Negative interactions and virulence differences drive the dynamics in multispecies bacterial infections

Désirée A. Schmitz[a], Richard C. Allen[a] & Rolf Kümmerli [a]

[a] Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

Bacterial infections are often polymicrobial, leading to intricate pathogen-pathogen and pathogen-host interactions. There is increasing interest in studying the molecular basis of pathogen interactions and how such mechanisms impact host morbidity. However, much less is known about the ecological dynamics between pathogens and how they affect virulence and host survival. Here we address these open issues by co-infecting larvae of the insect model host *Galleria mellonella* with one, two, three or four bacterial species, all of which are opportunistic human pathogens. We found that host mortality was always determined by the most virulent species regardless of the number of species and pathogen combinations injected. In certain combinations, the more virulent pathogen simply outgrew the less virulent pathogen. In other combinations, we found evidence for negative interactions between pathogens inside the host, whereby the more virulent pathogen typically won a competition. Taken together, our findings reveal positive associations between a pathogen's growth inside the host, its competitiveness towards other pathogens, and its virulence. Beyond being generalizable across species combinations, our findings predict that treatments against polymicrobial infections should target the most virulent species to reduce host morbidity, a prediction we validated experimentally.

multispecies bacterial infections | microbial and microbe-host interactions | virulence

Correspondence to: *desiree.schmitz@uzh.ch* or *rolf.kuemmerli@uzh.ch*

Importance
There is increasing evidence that polymicrobial infections are common and that host morbidity is impacted by interactions between the co-infecting pathogens. In this study, we infected the larvae of the insect host *Galleria mellonella* with four opportunistic human pathogens as mono or multispecies bacterial infections in all possible combinations and followed host survival and pathogen interactions over time. We discovered that species followed the exact same rank order in terms of their virulence, their growth inside the host, and their competitive ability against co-infecting pathogens. As consequence of this consistent rank order, we found that the most virulent species determines host survival dynamics in mixed infections. Beyond revealing a generalizable predictive pattern for virulence outcomes, our findings also predict that treatments should target the most virulent species in polymicrobial infections, a prediction we validated experimentally.

Introduction

Research over the past decades has revealed that many bacterial infections are polymicrobial (1–4) . This can lead to complicated pathogen-pathogen and pathogen-host interactions. The ecological interactions between pathogens can range from competition through commensalism to cooperation, and such interactions can have consequences for the host (5). For example, mutually beneficial cross-feeding between a human commensal bacterium and a pathogen was shown to increase the virulence of the latter, thus exacerbating host morbidity (6). Competition can also enhance virulence as revealed for the pathogen *Pseudomonas aeruginosa* that increases the production of the toxin pyocyanin thereby harming both its competitor *Staphylococcus aureus* and host tissue (7). Interaction patterns can become even more complex in chronic infections, where (co-)evolution can occur between pathogens and between pathogens and the host (8–13). There is increasing awareness that such eco-evolutionary dynamics affect host health and are important to consider in the context of treatment options (5).

For this reason, research on interactions between pathogenic bacteria has flourished in the past years. For example, there is a wealth of work on interactions between *P. aeruginosa and S. aureus*, two of the most troublesome nosocomial pathogens (14–16). This research, often carried out *in vitro*, has successfully identified molecular mechanisms of pathogen interactions and evolutionary patterns of how pathogens adapt to one another. However, we know much less about pathogen-pathogen interactions within hosts and how interactions drive virulence. Moreover, it is often unclear whether insights from a particular pathogen pair are generalizable and hold for other pathogen and strain combinations (17).

Here, we aim to tackle these questions by studying bacterial interactions between four different opportunistic human pathogens in an insect host, the larvae of the greater wax moth *Galleria mellonella*. This model host is suitable for our purpose because it allows for relatively high-throughput experiments where many different pathogen combinations can be tested, and well-defined doses of pathogens can be injected into a larva. Moreover, *G. mellonella* has an innate immune system that resembles the vertebrate innate immune response and lives approximately at human body temperature, which makes it an excellent model organism to study human bacterial pathogenesis (18–21). Regarding the pathogens, our aim was to choose four gram-negative bacterial species that exhibit different levels of virulence in order to test whether generalizable patterns of host survival and pathogen interactions arise across a spectrum of pathogens. We picked the following four opportunistic human pathogens. *Pseudomonas aeruginosa* (P) is a species of high clinical relevance that causes both community- and hospital-acquired infections including skin and wound infections, urinary tract infections, bloodstream infections, and pneumonias (22, 23). *Burkholderia cenocepacia* (B) causes chronic lung infections in immunocompromised patients, e.g. suffering from cystic fibrosis (24). *Klebsiella michiganensis* (K) belongs to the *K. oxytoca* complex (25), which are human pathogens that lead to health care-associated infections as well as to a variety of infections such as diarrhea, bacteremia, and meningitis (26–28). *Cronobacter sakazakii* (C) can also cause bacteremia and meningitis as well as necrotizing enterocolitis and brain abscess/lesions (29). While some of these pathogens can be found together – P and B in lung infections of cystic fibrosis patients (4, 30, 31) and P and K in burn wound infections (32, 33) – co-occurrence for other combinations is rarer. Thus, our model reflects a scenario where different opportunistic pathogens co-incidentally infect the same host without any prior interaction history or coevolution between the co-infecting pathogens. Such a scenario reflects the starting point of many acute and chronic polymicrobial infections.

In a first set of experiments, we aimed to understand the demographics of pathogen-host interactions in single species (mono) infections. For this purpose, we manipulated the injection dose and host age to understand how these factors affect each pathogen's virulence and mortality for the host. Second, we conducted mixed species infection experiments (double, triple, and quadruple) to assess how polymicrobial infections affect virulence patterns and host survival. Third, we quantified pathogen growth within the host at two time points for all mono and pairwise infections to test whether pathogen load links to virulence. Fourth, we followed changes in pathogen frequencies in mixed infections to derive pathogen interaction patterns. While this approach does not reveal specific molecular mechanisms of pathogen interactions, it allows us to distinguish between negative, neutral, and positive interactions between co-infecting pathogens and how such interactions link to virulence. One key insight from our experiments was that host mortality was always driven by the most virulent species, with the virulence of a pathogen correlating with its growth inside the host and its competitiveness towards the co-infecting species. Based on these insights, we predicted that the most virulent species should be targeted in an infection to reduce host morbidity and mortality. In a follow-up experiment, we validated this prediction by specifically targeting the more virulent *P. aeruginosa* co-infecting the host together with the less virulent *B. cenocepacia*.

Results

Injection dose, host age, and species identity determine virulence
To assess how the four different pathogens affect host survival, we exposed *G. mellonella* larvae (last instar larvae, relative age: 5, 10, or 15 days after arrival in our laboratory) to a range of pathogen injection doses (100 to 1 million CFU) for each of the four pathogens (B, C, K, P) and tracked their survival over 48 hours (Fig. 1A and 1B and Fig. S1).

We found that the hazard to die differed between the four species (Cox proportional hazard: χ_3^2 = 308.6, p < 0.0001) and was influenced by significant interactions between species and larval age (χ_3^2 = 67.28, p < 0.0001), and species and injection dose (χ_3^2 = 10.96, p = 0.0119). When examining these interactions more closely, we observed that the hazard to die grew significantly with increasing larval age for C (z = 2.69, p = 0.0071) and K (z = 6.43, p < 0.0001), but only marginally for B (z = 1.90, p = 0.0573), and not at all for P (z = -0.72, p = 0.471). For the injection dose, we found that higher CFU significantly increased the hazard to die in all four species, but the hazard increase was more pronounced for B and P than for K and C (see Table S1 for the full statistical analysis). The reason for the species-specific relationship between virulence and infection dose stems from the fact that the survival decreased in a log-linear fashion for B and P with higher pathogen doses, whereas C and K only became virulent above a certain threshold injection dose of about 10^4 CFU for K and 10^5 CFU for C.

Based on the insights from this experiment, we decided to use an injection dose of 10^5 CFU for all subsequent experiments. At this injection dose, all four pathogens are virulent with the following order of virulence: *P. aeruginosa* (P) > *B. cenocepacia* (B) > *K. michiganensis* (K) > *C. sakazakii* (C), and with significant host age effects for C and K (Fig. 1C).

of *G. mellonella* larval age (5, 10, 15 days), injection dose (CFU), and pathogen species (blue: *B. cenocepacia*, green: *C. sakazakii*, yellow: *K. michiganensis*, red: *P. aeruginosa*). The lines and the shaded area (95% CI) depict the relationship between mean survival time (across the different larval ages and replicates) and infection dose for each species. The black box highlights the injection dose chosen for all subsequent experiments. (B) Kaplan Meier survival curves of *G. mellonella* larvae for an injection dose of 10^5 CFU for each of the four pathogen species. Relative larval age – 5, 10, and 15 days after the arrival in our laboratory – is indicated by increasing line opaqueness. (C) The arrow chart shows the rank order of virulence among the four bacterial pathogens, from lowest to highest based on our experimental data. Data are from three independent experiments, each featuring 10-12 larvae per treatment, resulting in a total of 30-36 larvae per treatment.

Host mortality in multispecies infections is determined by the most virulent species

Next, we compared host survival between mono and mixed infections. For this, we injected pairwise combinations of our four bacterial species into larvae of *G. mellonella* and observed their survival over 48 h (Fig. 2A). The injection dose was always 10^5 CFU in total, with equal amounts of each co-injected species. An intuitive expectation is that the virulence in co-infections should be intermediate between the mono infections of the two species. In contrast to this expectation, we found that any mixed infection followed the dynamics of its most virulent species (Fig. 2B). A statistical examination confirmed that larval survival was not different between the mixed infection and the mono infection of the more virulent species in three out of six cases (log-rank test B+K vs. B: p = 0.2459; C+K vs. K: p = 0.9770; C+P vs. P: p = 0.0548). In the remaining three cases, survival in the mixed infection was significantly lower than the mono infection of the more virulent species (B+C vs. B: p = 0.0033; B+P vs. P: p = 0.0230; K+P vs. P: p = 0.0041), but the actual biological differences observed are extremely small (Fig. 2B).

We then tested whether the same patterns arise in the four triple and the quadruple infections. As above, we kept the total number of CFU constant at 10^5 CFU and mixed equal amounts of each pathogen. As for the paired infections, we found that the survival trajectory of larvae infected with three and four pathogens followed the

trajectories of the mono infection of the most virulent species in the mix (Fig. 2C). For example, in the triple infection with B, C, and K, larval survival followed the one of the B mono-infection, which is the most virulent of the three species. In the remaining four mixes, the most virulent species was P and in all these cases larval survival followed the one of P mono infections. Statistical analyses revealed subtle but significant differences: mixed infections with P were always slightly less virulent than the mono infections with P (see Table S2 for the full statistical analysis). We can explain this pattern by a density effect: fewer P cells were injected in higher order infections, and thus bacteria needed more time to replicate and reach sufficiently high numbers to kill the larval host (Fig. S2).

curves of mono versus pairwise (B), triple, and quadruple (C) infections with the shaded area indicating the 95% CI. The denotation ns (non-significance) marks cases for which survival does not significantly differ between one of the mono infections compared to the mixed

infection in the same panel. Data are from five independent experiments, each with 10-12 larvae per treatment, resulting in a total of 50-60 larvae per treatment.

The most virulent species is also more abundant both in mono and mixed infections

We then asked what the underlying reason could be for our observation that the most virulent species drives host survival dynamics. One explanation would be that the more virulent species grows better in the host, making it more abundant and thereby exerting a stronger effect on the host. We tested this hypothesis in mono infections first and found that bacterial load (CFU per larva) of the four pathogens followed the exact order of their virulence at 12 hours post infection (hpi, see Fig. S3 in the supplemental information, and Table S3 for the full statistical analysis). At the earlier timepoint (6 hpi), the same pattern holds for B, C, and P, whereas K had much higher CFU inside the host than expected from its virulence. The data imply that K initially grows well in larvae, while it is compromised later during the infection. Overall, the mono infection data suggest that pathogen load in a host positively links to its level of virulence (i.e.,

the percentage of larvae in which either coexistence (grey) of pathogens occurred or only a single species (color) was left. Data are shown for all pairwise infections both at 6 hpi and 12 hpi. Data are from 4-5 individual experiments with 2-3 larvae per treatment, resulting in a total of 8-12 larvae per treatment.

Next, we explored whether the same is true for mixed infections, i.e., whether the dominant effect of the more virulent species might be caused by its higher abundance in a co-infection. Indeed, in all cases the more virulent species was also the more abundant species at 12 hpi (Fig. 3). For example, P as the most virulent of our pathogens dominated in terms of abundance in all larvae in any of its mixes at 12 hpi. However, we also found evidence for more complex dynamics, where relative species abundance changed over time. Especially in the infection pairs K+P and K+B, we observed that K dominated at 6 hpi despite being the less virulent species, a pattern that disappeared at 12 hpi. Important to note is that in five out of six pairings both pathogen species coexisted in the host during the course of the infection in a large proportion of larvae.

Negative interactions dominate in pairwise infections
We used the CFU data from mixed infections to test whether the growth of any of the four species was influenced by the presence of a co-infecting species across two time points (at 6 and 12 hpi). We found that the co-infecting species had either no effect or a negative effect on a focal species in a pathogen-pair specific way (Fig. 4, see Table S4 for the full statistical analysis). The negative effect was either independent of how prevalent the co-infecting species was (main effect) or stronger with increasing CFU of the co-infecting species (CFU effect).

B. cenocepacia (B): The growth of this pathogen increased from 6 hpi to 12 hpi (t_{77} = 2.66, p = 0.0095), and was not affected by C (t_{77} = -0.10, p = 0.9171). In contrast, B growth was compromised by increasing numbers of K at the earlier timepoint and P at both timepoints (CFU effect of second species, for K: t_{77} = -4.73, p < 0.0001; for P: t_{77} = -4.17, p < 0.0001). The presence of P at 12 hpi resulted in a CFU drop of B even below the inoculum size in many larvae.

C. sakazakii (C): In mono infections, this pathogen's CFU dropped below the inoculum size at both timepoints, suggesting that it cannot replicate within the host. While the presence of B did not affect C at any given time, both K and P significantly compromised C abundance (main effect, for K: t_{66} = -3.29, p = 0.0016; for P: t_{66}= -2.20, p = 0.0314).

K. michiganensis (K): In line with the host survival data (Fig. 1A and B), we found more CFU of this pathogen the older the larval host was (t_{71}= 4.27, p < 0.0001). The growth of K significantly decreased with higher CFU of the co-infecting B at both timepoints (CFU effect of B: t_{71} = -2.48, p = 0.0156). Interestingly, our model predicts that K reaches higher CFU if C is present at sufficiently high numbers (CFU effect of C: t_{71} = 2.90, p = 0.0050), but because C never reached high CFU itself, the overall effect of C on K was negative in our experiments (main effect of C: t_{71} = -2.92, p < 0.0047).

P. aeruginosa (P): The CFU of P significantly increased from 6 to 12 hpi (t_{73} = 9.89, p < 0.0001). P growth was only affected by K, which had a strong negative impact at 6 hpi (CFU effect of K: t_{73} = -4.40, p < 0.0001).

While the number of interactions decreased over time (58% at 6 hpi; 42% at 12 hpi), all interactions were negative at both timepoints. Our four bacterial species spanned from being inhibited by one (P) or two other species (B, C, K) and inhibiting

Coexistence ○ *Yes* △ *No*   DominantSpecies ● *B* ● *C* ● *K* ● *P*

| *P. aeruginosa* | *P. aeruginosa* | *P. aeruginosa* | *B. cenocepacia* | *B. cenocepacia* | *K. michiganensis* |
| *P. aeruginosa* | *P. aeruginosa* | *P. aeruginosa* | *B. cenocepacia* | *B. cenocepacia* | *K. michiganensis* |

Fig. 4B). At 12 hpi, the competitive
of virulence and growth, whereby P
C mutually repress each other with

P

| *cenocepacia* | *K. michiganensis* |
| *cenocepacia* | *K. michiganensis* |

B+P

P

| *cenocepacia* | *K. michiganensis* |
| *cenocepacia* | *K. michiganensis* |

P

| *cenocepacia* |
| *cenocepacia* |

Hours post infection [hpi]

Hours post infection [h]

pathogens 6 and
for the mono inf
measured. Dots
injection dose of
was 100 000 CF
infections, we ha
the median (line
interquartile rang
interquartile range. Data are from 4-5
treatment, resulting in a total of 8-12
bacterial consortium based on the stat
indicated otherwise. Significant negativ
stop arrow. The label "independent of
the amount of the co-infecting pathogen

Rel

Hours post infection [h]

Targeting the more virulent species in mixed infections increases host survival
The main conclusion of the above results is that there are positive associations between a pathogen's growth inside the host, its competitiveness towards other pathogens, and its virulence. Consequently, we predicted that treating the most virulent species in a multispecies bacterial infection would be an effective strategy to increase host survival. To test this prediction, we co-infected *G. mellonella* larvae with B+P and compared host survival of the mixed infection to the mono infections of B and P, with all conditions receiving either no treatment or a single dose of gentamicin after two hours. Given that B is inherently resistant to gentamicin, we predicted that the more virulent pathogen P should be selectively killed and the survival dynamics of the mixed infection should follow the one of the B mono infection.

In the absence of gentamicin, we recovered our previous findings (Fig. 2B) that mixed B+P infections followed the virulence trajectory of P mono infections (Fig. 5). In stark contrast and in support of our prediction, we observed that host survival in B+P mixed infection more closely followed the trajectory of the B mono infection under gentamicin treatment (Fig. 5). This finding suggests that P was selectively removed by gentamicin, which is supported by our mono infection data showing that larval survival is very high for the P infection treated with gentamicin. Overall, B+P infections treated with gentamicin were significantly less virulent than B+P infections not receiving gentamicin treatment (log-rank test: B+P+gentamicin vs. B+P: p < 0.0001). These results support the view that treating the most virulent species in a polymicrobial infection is a promising strategy to reduce host mortality.

Figure 5. Treating the more virulent pathogen reduces host mortality in mixed infections to the level of the less virulent pathogen. Kaplan Meier survival curves of *G. mellonella* larvae injected with either NaCl (control, grey), B (less virulent pathogen, blue), P (more virulent pathogen, red), or B+P mixes (black). Larvae received either a gentamicin treatment (dashed lines) two hours after the bacterial/NaCl injection or no antibiotic treatment (solid lines). Survival curves are based on a total of 48-54 larvae per treatment from three independent experiments.

Discussion
There is increasing evidence that polymicrobial infections are common (1–4) and that ecological and evolutionary interactions between co-infecting pathogens can affect host morbidity (5–7). In our study, we examined whether there are generalizable patterns that characterize interaction dynamics between pathogens and a common host in multispecies infections. For our experiments, we used the larvae of *G. mellonella* as the host and infected it with four opportunistic human bacterial pathogens alone and in all possible combinations of mixed infections. We tested the rank order of virulence for the four pathogens and found that it was identical to the rank order of growth in the host. In addition, the more virulent species in mono infections were also better at outcompeting other species in mixed infections. A consequence of this was

that the most virulent species determined host survival dynamics in mixed infections regardless of the number and type of pathogens mixed. Our findings, which held for all pathogen combinations tested, reveal an infection dynamic that is not covered by any of the current models for pathogen virulence in mixed infections.

While we found that the virulence of a specific pathogen is positively linked to its growth and competitiveness in the host, our co-occurrence analysis could give us an idea of the relative importance of pathogen growth versus competitiveness and how it varies across pathogen species combinations (Fig. 4). In the pathogen interaction network (Fig. 4B), the absence of an interaction suggests that differences in growth dominate a co-infection, meaning that the faster growing species simply outperformed the slower growing one. This was the case for five (6 hpi) and seven (12 hpi) out of the total twelve interactions. Conversely, the remaining seven (6 hpi) and five (12 hpi) interactions that were negative would imply that competitiveness played a more prominent role. At least two types of competition could be involved. First, the faster-growing species limits resource availability for the slow-growing species, reducing its growth and survival due to starvation. Second, the more competitive species secretes a toxin or deploys contact-dependent mechanisms to target and kill the less competitive species in interference competition. While our results do not allow to differentiate between resource and interference competition (34), it is likely that both mechanisms matter. For example, in our mixed infections with K+B, and K+P, we observed that K inhibits its competitors early on during the infections (6 hpi) but is outcompeted at a later stage (12 hpi). One option is that the inhibitory effects early during the infection could be explained by K secreting a toxin, while the dominance of B and P later in the infection could be due to resource competition advantages. Alternatively, it could be that K grows quickly but inefficiently while its competitors (B and P) grow more slowly but efficiently and thus outpace K over time.

Important to note is also that all pathogen interactions were either negative or neutral, but never positive. This finding supports the view that competition is much more prevalent between bacterial species than positive interactions, where one species unilaterally or mutually benefits another species (35, 36). The prevalence of competitive interactions is perhaps expected given that the pathogens interacted in a closed environment (i.e., the larva), where both resource availability and host longevity are limited (37).

Competition between pathogens is also a major component of mathematical models predicting virulence levels (38). A traditional set of models assumes that genetically diverse pathogens engage in increased levels of resource competition in mixed infections, which is predicted to exacerbate virulence (37). Other models examined the effect of infighting between pathogens through toxin secretion (39) or competition for publicly shared virulence factors (40). These models predict that increased competition between pathogens should decrease virulence in mixed infections. Empirical support for these models vary, lending support to both types of predictions (41–44). A key insight from these studies is that the biological details of pathogen interactions matter. While our study was not designed to test specific model predictions, our results put forth a third virulence scenario, namely that pathogen interactions in mixed infections neither increase nor decrease virulence, but the virulence trajectory simply follows that of the more harmful species. A first example of this scenario was reported by Massey et al. (43) and here we reveal its generality. As discussed above, this pattern can arise when pathogen traits that are relevant for infections (virulence, growth, and competitiveness) are positively connected with one another. This virulence model could be particularly relevant for infections with

opportunistic pathogens that co-incidentally end up in the same host with no prior interaction and coevolution history (3).

Our data further indicate that host factors also influence pathogen virulence patterns. For example, we found that younger larvae had longer survival times than older larvae when infected with B, C, and K. Given that *G. mellonella* has an innate immune system similar to the one of vertebrates, our results indicate that the immune response works well against weaker pathogens (like C and K) but deteriorates with age (i.e. number of days after the arrival of the larvae in our laboratory). Another host effect we observed is that *G. mellonella* managed to control infections of C and K at low injection doses, again highlighting the potency of the insect's immune system. A further indication that pathogen performance is modulated by the host stems from an additional experiment, which we conducted with a medium that mimics the hemolymph of the insect order Lepidoptera (Fig. S4). Although this medium nutritionally matches insect conditions, we found the rank order of pathogen species growth to be markedly different compared to growth in live hosts (insect condition *in vitro*: C > P > K > B vs. *in vivo*: P > B > K > C). Clearly, host effects are important and likely feedback on pathogen-pathogen interactions, which is why they should be considered in future work on polymicrobial infections.

In conclusion, we can draw two generalities from our four-bacteria-infection system. First, no matter what pathogen combination we used, the most virulent pathogen dictated host survival. This led us to predict that targeting the most virulent pathogen seems a promising strategy to reduce host morbidity/mortality in polymicrobial infections. We found support for this prediction in a proof-of-principle experiment, where we selectively targeted the more virulent species in a mixed infection with antibiotics. Second, more virulent pathogens grow better in the host and are better in competition with less virulent pathogens. These observations suggest that the same traits (or co-regulated traits) responsible for attacking the host (e.g., virulence factors) could promote pathogen growth and help in competition with other pathogens. Identifying these traits could give rise to promising strategies to control polymicrobial infections.

Materials and Methods

Bacterial strains & host
We used the following four opportunistic human pathogens: *Pseudomonas aeruginosa* PAO1 (45), *Burkholderia cenocepacia* K56-2 (46), *Klebsiella michiganensis*, and *Cronobacter sakazakii* (ATCC29004). We purchased the larvae of the greater wax moth *G. mellonella* in their last instar stage from a local vendor (Bait Express GmbH, Basel, Switzerland). We contacted the vendor who confirmed that their larvae had not been treated with any antibiotics. We can therefore rule out the possibility that pathogen interactions in our experiments are affected by residual antibiotic concentrations in the larvae. Upon arrival, we stored them at 4-8° C without food. We considered larval age as the number of days after arrival in our laboratory, since the exact age of larvae was unknown to us. Hence, we only investigated larval age as a relative measure. When examining larval age, we compared larvae that were ordered at the same time (i.e., same batch), but were infected at different time points. To put host age into a wider context, the *G. mellonella* life cycle comprises four developmental phases: egg (3-30 d), larva/caterpillar (6-7 weeks), pupa (6-55 d), and adult (47). We only worked with the second phase, i.e., the larva, at its last out of seven larval stages before turning into a pupa (last instar stage) (48).

Culturing conditions of bacteria

Bacterial stocks were kept in 25% glycerol and stored at -80° C. For all experiments, we grew bacteria overnight until stationary phase in 5 mL lysogeny broth (LB) in 50 mL Falcon tubes at 37° C and at 170 rpm with aeration (Infors HT, Multitron Standard Shaker). For all species, we centrifuged 1 mL overnight-cell culture in a 1.5 mL Eppendorf tube at 7'500 rcf for 5 min (Eppendorf, tabletop centrifuge MiniSpin plus with rotor F-45-12-11) and washed cultures three times with a 0.8% NaCl solution to remove all original media. Next, we measured the optical density at 600 nm (OD_{600}) of the washed cell culture (Amersham Biosciences, Ultrospec 2100 pro spectrophotometer), adjusted it to OD_{600}=1, and diluted each species individually to obtain similar cell numbers per milliliter for all species.

While all our main experiments were carried out in the *G. mellonella* larvae, we conducted one additional *in vitro* experiment. Specifically, we grew all four pathogens as monocultures in a medium that mimics the hemolymph of Lepidoptera – the insect order to which *G. mellonella* belongs – called Grace's insect medium (GIM, Fig. S4). The aim of this experiment was to show that our model host directly affects pathogen growth and is not just serving as a sack to conduct growth assays. Thus, we expected pathogen growth to be different inside the host compared to the *in vitro* GIM environment. We prepared bacterial overnight cultures as stated above to obtain washed and OD_{600}=1 adjusted cell cultures. A 96-well cell culture plate (Eppendorf, non-treated, flat bottom) was then filled with 190 µL GIM (Gibco, GIM 1X, supplemented) and 10 µL cells to reach a starting OD_{600}=0.001 per well. To reduce evaporation the inter-well spaces and the outer moat space of the microplate were filled with sterile water (13 mL in total). We incubated the microplate at 37° C in a microplate reader (Tecan, Infinite MPlex, monochromator optics) and measured OD_{600} every 15 min for 24 h, shaking for 60 sec before every read. The plate layout was randomized between experiments. All chemicals were purchased from Sigma Aldrich, Switzerland, unless indicated otherwise.

Infection experiments
Prior to infections, we sorted larvae and evenly distributed them across treatments according to size. Their weight ranged from 278-731 mg with an average weight of 499 mg. We put larvae on ice in a petri dish to immobilize them. Next, we surface-sterilized larvae with 70% ethanol and injected them between the posterior prolegs. We used a sterile hypodermic needle (Braun, 0.45 x 12 mm BI/LB, 26G), a sterile syringe (Braun, 0.01-1 mL Injekt-F Luer Solo), and a programmable syringe pump (New Era Pump Systems Inc, model NE-300) to standardize injection speed. With a flow rate of 2 ml/h, we either injected 10 µL control or bacterial solution (both in 0.8 % NaCl) into larvae within 18 s. We had a second control treatment, where larvae were not injected. The injection dose ranged from 10^2 to 10^6 CFU for mono infections and was kept at 10^5 CFU for experiments that included multispecies infections. In multispecies infections, we mixed equal amounts of each species. The order of treatments was randomized between experiments. For all experiments, we confirmed the injection dose in triplicate by CFU on 1.5% LB agar plates. We distributed injected larvae to individual wells of a 12-well plate for incubation at 37° C in the dark without food. We monitored survival of every larva in regular intervals (hourly during 12-24 hours post infection (hpi) and every two hours during 36-48 hpi). Larvae were considered dead when they did not move upon touch with a pipette tip.

We used gentamicin to selectively target P in co-infections with B. First, we determined the gentamicin concentration that killed P but had no effect on B *in vitro*, since B is inherently resistant to gentamicin (up to a certain concentration). We found that 8 µg/mL gentamicin fulfilled the condition. Thus, we injected 10 µL of a 400 µg/mL gentamicin solution into the larvae, which should result in an internal gentamicin concentration of 8 µg/mL based on the average larval weight (assuming 499 mg ~ 500 mL according to (49)). The gentamicin treatment was applied two hours after the bacteria had been injected so that they had time to establish an infection in the host. To control for any impact caused by the second injection, half of the NaCl sterile control larvae also received the gentamicin treatment.

Bacterial load in the hemolymph
We determined the bacterial load in all larvae that received mono-, mixed-, NaCl, and no infection at two time points, at 6 and 12 hpi. Since larvae had to be sacrificed during this

process, we had two independent batches of larvae that were used for the respective timepoints. First, we placed larvae individually in 2 mL Eppendorf tubes and submerged those in ice until larval movement halted. Then, we opened each larva with a surgical blade (Aesculap AG, Germany) behind the posterior prolegs, with the cut length spanning half of the body width. This procedure is similar to the one used by Harding et al. (50), McCloskey et al. (51), and Admella & Torrents (52). We sterilized the blade with 70% ethanol between different larvae. Next, we drained 10-20 µL hemolymph into a fresh 1.5 mL Eppendorf tube by gently squeezing the larva. The collected hemolymph was mixed and stored on ice. To enumerate bacterial CFU, we serially diluted 10 µL of the collected hemolymph in 0.8% NaCl up to 10^{-6} and plated the appropriate dilutions on LB-agar plates (1.5% agar) in duplicates. The appropriate dilutions varied between 10^{0} (undiluted) and 10^{-6} and depended on the bacterial species. We incubated the plates at 37° C overnight and then manually counted CFU. Because *B. cenocepacia* grew slower than the other species, plates were incubated for two days. Plates with *C. sakazakii* were kept at room temperature for an additional day following overnight incubation for colonies to develop their characteristic yellow color. All four species could be distinguished from each other based on their different colony morphologies (see Fig. S5). We considered plates with a minimum of 25 CFU up to a number of CFU for which we could still distinguish single colonies with confidence. Plates with less than 25 CFU were only considered if none of the plated dilutions from the same individual adhered to this threshold. For mixes where one species occurred at very low frequency (mostly mixes with P), we plated the undiluted hemolymph to see whether the rare species was present at all. While both C and K were visible even in a lawn of P, this was not true for B. For this species combination, we cannot conclude that B was completely absent in certain larvae but rather that P was highly dominant. We calculated the CFU/larva by multiplying the volume of plated hemolymph with the average larval weight since larval weight is almost identical to larval liquid volume according to Andrea et al. (49).

In approximately 20% of the extracted larval hemolymph, we detected bacteria different from our four focal species. We determined the identity of ten representative isolates by 16S rRNA sequencing (colony PCR using standard primers 1492r and 27f). We identified *Enterococcus casseliflavus*, and *E. gallinarum*, two common insect gut bacteria (53, 54), as the closest relatives of our isolates with sequence identities of 89-99%. These results suggest that we had drained certain larvae too extensively, so that we not only collected bacteria from the hemolymph but also from the gut. For this reason, we excluded these larvae from all further analysis.

Statistical analysis
All statistical analyses were performed with R (version 4.1.1) and the interface RStudio (version 2021.09.0) (55). For the dose-response curves in Fig. 1A, we measured mean survival time of the host as the area under the full survival curve using the function survmean from the survival package (56). To build a log-logistic model we used the drm function from the drc package (57). To compare host survival in mono versus mixed infections, we built a separate model for each panel shown in Fig. 2, comparing host survival of a particular pathogen combination. The statistical results were robust across three different methods comparing host survival: the non-parametric log-rank test, the semi-parametric Cox proportional hazards model, and the parametric Weibull regression. See Table S2 for a model comparison. Non-parametric log-rank tests were also used to compare host survival between mono and mixed infections with or without gentamicin treatment (Fig. 5, Table S2).

To determine whether the pathogens significantly vary in their bacterial load in mono infections, we compared the CFU extracted from the hemolymph of host individuals at 6 hpi and 12 hpi. We chose a linear model using generalized least squares to account for differences in variance between the species, which was mainly caused by *K. michiganensis* at 12 hpi (see Table S3 for the full statistical analysis). Next, we performed a post-hoc analysis using the Tukey honest significant difference test with p-value adjustment to compare the bacterial load between species.

To compare bacterial load in the hemolymph in mono- versus mixed infections, we built a separate linear model for each pathogen to test whether its growth was influenced by the presence of another species, the CFU of this other species and time (at 6 hpi versus 12 hpi). Since the injection dose was always the same independent of how many species were injected, we compared for each species half the CFU found in mono infections to the total CFU (for the focal species) observed in pairwise infections. To obtain normally distributed residuals, we transformed all CFU values in our models. We used the function transformTukey from the rcompanion package to find the best transformation (58). Non-significant interaction terms and main effects were removed from models until a minimal model was obtained. Most of the minimal models included both a main effect of the co-infecting species and its CFU effect. We often found that these two explanatory variables had opposing signs, which means that the influence of the co-infecting species on the focal species depended on the actual CFU of the co-infecting species. For this reason, we assessed the threshold CFU of the co-infecting pathogen at which its effect on the focal pathogen switches from being positive to being negative. This was done for each pathogen combination. We could then compare this CFU threshold to our experimental CFU values to check whether a co-infecting species stimulated or inhibited a focal species. To calculate the standard error for confidence intervals of this threshold and the p-value (one sample t-test versus zero) we used the Taylor expansion in the R package propagate (59).

Data availability
All raw data sets will be deposited in the figshare repository (LINK).

Supplementary information
Supplementary information will be made available online: Supplemental file, XLSX file, XX MB.

Acknowledgements
We thank Kayla King, Anna-Liisa Laine, Alex Hall, and Roland Regoes for their scientific inputs. We also thank Nadine Koch for showing us how to conduct injections with the larvae of *G. mellonella*.

Funding
This project has received funding from the Swiss National Science Foundation (grant no. 31003A_182499 to RK) and from the Novartis Foundation for Medical-Biological Research (to RK).

Compliance with ethical standards
Conflict of interest: The authors declare that they have no conflict of interest.

Bibliography
1. Peters BM, Jabra-rizk MA, Costerton JW, Shirtliff ME. 2012. Polymicrobial Interactions: Impact on Pathogenesis and Human Disease. Clin Microbiol Rev 25:193–213.
2. Brogden KA, Guthmiller JM, Taylor CE. 2005. Human polymicrobial infections. Lancet 365:253–255.
3. Short FL, Murdoch SL, Ryan RP. 2014. Polybacterial human disease: the ills of social networking. Trends Microbiol 22:508–516.
4. Folkesson A, Jelsbak L, Yang L, Johansen HK, Ciofu O, Hoiby N, Molin S. 2012. Adaptation of Pseudomonas aeruginosa to the cystic fibrosis airway: An evolutionary perspective. Nat Rev Microbiol 10:841–851.
5. Rezzoagli C, Granato ET, Kümmerli R. 2019. Harnessing bacterial interactions to manage infections: a review on the opportunistic pathogen Pseudomonas aeruginosa as a case example. J Med Microbiol 68:1–15.
6. Stacy A, Everett J, Jorth P, Trivedi U, Rumbaugh KP, Whiteley M. 2014. Bacterial fight-and-flight responses enhance virulence in a polymicrobial infection. Proc Natl Acad Sci U S A 111:7819–7824.
7. Korgaonkar A, Trivedi U, Rumbaugh KP, Whiteley M. 2013. Community surveillance

enhances Pseudomonas aeruginosa virulence during polymicrobial infection. Proc Natl Acad Sci U S A 110:1059–1064.

8. Frank SA. 1992. Models of plant-pathogen coevolution. Trends Genet 8:213–219.

9. Frank SA. 1993. Coevolutionary genetics of plants and pathogens. Evol Ecol 7:45–75.

10. Van Baalen M. 1998. Coevolution of recovery ability and virulence. Proc R Soc London B 265:317–325.

11. Buckling A, Rainey PB. 2002. Antagonistic coevolution between a bacterium and a bacteriophage. Proc R Soc London B 269:931–936.

12. Nourmohammad A, Otwinowski J, Plotkin JB. 2016. Host-Pathogen Coevolution and the Emergence of Broadly Neutralizing Antibodies in Chronic Infections. PLoS Genet 12:1–23.

13. Papkou A, Guzella T, Yang W, Koepper S, Pees B, Schalkowski R, Barg MC, Rosenstiel PC, Teotónio H, Schulenburg H. 2019. The genomic basis of red queen dynamics during rapid reciprocal host–pathogen coevolution. Proc Natl Acad Sci U S A 116:923–928.

14. Limoli DH, Hoffman LR. 2019. Help, hinder, hide and harm: What can we learn from the interactions between Pseudomonas aeruginosa and Staphylococcus aureus during respiratory infections? BMJ https://doi.org/10.1136/thoraxjnl-2018-212616.

15. Orazi G, Ruoff KL, O'Toole GA. 2019. Pseudomonas aeruginosa Increases the Sensitivity of Biofilm-Grown Staphylococcus aureus to Membrane-Targeting Antiseptics and Antibiotics. MBio 10:1–15.

16. Tognon M, Köhler T, Luscher A, Van Delden C. 2019. Transcriptional profiling of Pseudomonas aeruginosa and Staphylococcus aureus during in vitro co-culture. BMC Genomics 20:1–15.

17. Niggli S, Kümmerli R. 2020. Strain Background, Species Frequency, and Environmental Conditions Are Important in Determining Pseudomonas aeruginosa and Staphylococcus aureus Population Dynamics and Species Coexistence. Appl Environ Microbiol 86:1–14.

18. Sheehan G, Garvey A, Croke M, Kavanagh K. 2018. Innate humoral immune defences in mammals and insects: The same, with differences? Virulence 9:1625–1639.

19. Tsai CJY, Loh JMS, Proft T. 2016. Galleria mellonella infection models for the study of bacterial diseases and for antimicrobial drug testing. Virulence 7:214–229.

20. Ramarao N, Nielsen-Leroux C, Lereclus D. 2012. The insect Galleria mellonella as a powerful infection model to investigate bacterial pathogenesis. J Vis Exp 1–7.

21. Junqueira JC. 2012. Galleria mellonella as a model host for human pathogens: Recent studies and new perspectives. Virulence 3:474–476.

22. Revuz J, Penso D, Roujeau J, Guillaume J-C, Payne CR, Wechsler J, Touraine R. 1987. Toxic Epidermal Necrolysis. Arch Dermatol 123:1160–1165.

23. Gaynes R, Edwards JR. 2005. Overview of nosocomial infections caused by gram-negative bacilli. Clin Infect Dis 41:848–854.

24. Vandamme P, Holmes B, Vancanneyt M, Coenye T, Hoste B, Coopman R, Revets H, Lauwers S, Gillis M, Kersters K, Govan JRW. 1997. Occurrence of Multiple Genomovars of Burkholderia cepacia in Cystic Fibrosis Patients and Proposal of Burkholderia multivorans sp. nov. Int J Syst Bacteriol 47:1188–1200.

25. Yang J, Long H, Hu Y, Feng Y, McNally A, Zong Z. 2022. Klebsiella oxytoca Complex: Update on Taxonomy, Antimicrobial Resistance, and Virulence. Clin Microbiol Rev 35:1–39.

26. Podschun R, Ullmann U. 1998. Klebsiella spp. as nosocomial pathogens: Epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev 11:589–603.

27. Högenauer C, Langner C, Beubler E, Lippe IT, Schicho R, Gorkiewicz G, Krause R, Gerstgrasser N, Krejs GJ, Hinterleitner TA. 2006. Klebsiella oxytoca as a Causative Organism of Antibiotic-Associated Hemorrhagic Colitis. N Engl J Med 355:2418–26.

28. Neog N, Phukan U, Puzari M, Sharma M, Chetia P. 2021. Klebsiella oxytoca and Emerging Nosocomial Infections. Curr Microbiol 78:1115–1123.

29. Iversen C, Lehner A, Mullane N, Bidlas E, Cleenwerck I, Marugg J, Fanning S, Stephan R, Joosten H. 2007. The taxonomy of Enterobacter sakazakii: Proposal of a new genus Cronobacter gen. nov. and descriptions of Cronobacter sakazakii comb. nov. Cronobacter sakazakii subsp. sakazakii, comb. nov., Cronobacter sakazakii subsp. malonaticus subsp. nov., ... BMC Evol Biol 7:1–11.

30. Govan JRW, Deretic V. 1996. Microbial pathogenesis in cystic fibrosis: Mucoid Pseudomonas aeruginosa and Burkholderia cepacia. Microbiol Rev 60:539–574.

31. Eberl L, Tümmler B. 2004. Pseudomonas aeruginosa and Burkholderia cepacia in cystic fibrosis: Genome evolution, interactions and adaptation. Int J Med Microbiol 294:123–131.

32. Abbas HA, El-Masry EM, Shaker GH. 2013. Bacterial etiology and antimicrobial resistance of burn wound infections in a burn unit in Hehia General Hospital in Egypt. Int J Biol Pharm Res 4:1251–1255.

33. Forson OA, Ayanka E, Olu-Taiwo M, Pappoe-Ashong PJ, Ayeh-Kumi PJ. 2017. Bacterial infections in burn wound patients at a tertiary teaching hospital in Accra, Ghana. Ann Burns Fire Disasters 30:116–120.

34. Ghoul M, Mitri S. 2016. The Ecology and Evolution of Microbial Competition. Trends Microbiol 24:833–845.

35. Oliveira NM, Niehus R, Foster KR. 2014. Evolutionary limits to cooperation in microbial communities. Proc Natl Acad Sci U S A 111:17941–17946.

36. Foster KR, Bell T. 2012. Competition, not cooperation, dominates interactions among culturable microbial species. Curr Biol 22:1845–1850.

37. Frank SA. 1996. Models of Parasite Virulence. Q Rev Biol 71:37–78.

38. Buckling A, Brockhurst MA. 2008. Kin selection and the evolution of virulence. Heredity 100:484–488.

39. Inglis RF, Gardner A, Cornelis P, Buckling A. 2009. Spite and virulence in the bacterium Pseudomonas aeruginosa. Proc Natl Acad Sci U S A 106:5703–5707.

40. West SA, Buckling A. 2003. Cooperation, virulence and siderophore production in bacterial parasites. Proc R Soc London B 270:37–44.

41. Read AF, Taylor LH. 2001. The Ecology of Genetically Diverse Infections. Science 292:1099–1103.

42. Granato ET, Ziegenhain C, Marvig RL, Kümmerli R. 2018. Low spatial structure and selection against secreted virulence factors attenuates pathogenicity in Pseudomonas aeruginosa. ISME J 12:2907–2918.

43. Massey RC, Buckling A, Ffrench-Constant R. 2004. Interference competition and parasite virulence. Proc R Soc London B 271:785–788.

44. Boots M, Mealor M. 2007. Local Interactions Select for Lower Pathogen Infectivity. Science (80-) 315:1284–1286.

45. Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ, Brinkman FSL, Hufnagle WO, Kowallk DJ, Lagrou M, Garber RL, Goltry L, Tolentino E, Westbrock-Wadman S, Yuan Y, Brody LL, Coulter SN, Folger KR, Kas A, Larbig K, Lim R, Smith K, Spencer D, Wong GKS, Wu Z, Paulsen IT, Relzer J, Saler MH, Hancock REW, Lory S, Olson M V. 2000. Complete genome sequence of Pseudomonas aeruginosa PAO1, an opportunistic pathogen. Nature 406:959–964.

46. Mahenthiralingam E, Coenye T, Chung JW, Speert DP, Govan JRW, Taylor P, Vandamme P. 2000. Diagnostically and experimentally useful panel of strains from the Burkholderia cepacia complex. J Clin Microbiol 38:910–913.

47. Ellis JD, Graham JR, Mortensen A. 2013. Standard methods for wax moth research. J Apic Res 52:1–17.

48. Wojda I, Staniec B, Sułek M, Kordaczuk J. 2020. The greater wax moth Galleria mellonella: Biology and use in immune studies. Pathog Dis 78:1–15.

49. Andrea A, Krogfelt KA, Jenssen H. 2019. Methods and challenges of using the greater wax moth (Galleria mellonella) as a model organism in antimicrobial compound discovery. Microorganisms 7:1–9.

50. Harding CR, Schroeder GN, Collins JW, Frankel G. 2013. Use of Galleria mellonella as a model organism to study Legionella pneumophila infection. J Vis Exp 81:1–10.

51. McCloskey AP, Lee M, Megaw J, McEvoy J, Coulter SM, Pentlavalli S, Laverty G. 2019. Investigating the In Vivo Antimicrobial Activity of a Self-Assembling Peptide Hydrogel Using a Galleria mellonella Infection Model. ACS Omega 4:2584–2589.

52. Admella J, Torrents E. 2022. A Straightforward Method for the Isolation and Cultivation of Galleria mellonella Hemocytes. Int J Mol Sci 23:1–14.

53. Peng BY, Su Y, Chen Z, Chen J, Zhou X, Benbow ME, Criddle CS, Wu WM, Zhang Y. 2019. Biodegradation of Polystyrene by Dark (Tenebrio obscurus) and Yellow (Tenebrio molitor) Mealworms (Coleoptera: Tenebrionidae). Environ Sci Technol 53:5256–5265.

54. Allonsius CN, Van Beeck W, De Boeck I, Wittouck S, Lebeer S. 2019. The microbiome of the invertebrate model host Galleria mellonella is dominated by Enterococcus. Anim

Microbiome 1:1–7.

55. R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
56. Therneau TM, Grambsch PM. 2000. Modeling Survival Data: Extending the Cox Model. Springer, New York.
57. Ritz C, Baty F, Streibig JC, Gerhard D. 2015. Dose-Response Analysis Using R. PLOS One 10:e0146021.
58. Mangiafico S. 2021. rcompanion: Functions to Support Extension Education Program Evaluation. R package version 2.4.1.
59. Spiess A-N. 2018. propagate: Propagation of Uncertainty. R package version 1.0-6.