Proteome capacity constraints favor respiratory ATP generation

Authors

Yihui Shen [1,2], Hoang V. Dinh [5], Edward Cruz [2,4], Catherine M. Call [1,2], Heide Baron [2], Rolf-Peter Ryseck [1,2], Jimmy Pratas [1,2], Arjuna Subramanian [2], Zia Fatma [6], Daniel Weilandt [1,2], Sudharsan Dwaraknath [8], Tianxia Xiao [1,2], John I. Hendry [5], Vinh Tran [6], Lifeng Yang [1,2], Yasuo Yoshikuni [8], Huimin Zhao [6,7], Costas D. Maranas [5], Martin Wühr [2,4,*], Joshua D. Rabinowitz [1,2,3,*]

Author affiliations

Department of Chemistry, Princeton University, Princeton, NJ, USA

Lewis Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ, USA

Ludwig Institute for Cancer Research, Princeton Branch, Princeton, NJ, USA

Department of Molecular Biology, Princeton University, Princeton, New Jersey 08544, USA

Department of Chemical Engineering, The Pennsylvania State University, University Park, PA, USA

Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, Urbana, IL, 61801, USA

Department of Chemical and Biomolecular Engineering, University of Illinois at Urbana-Champagne, Urbana, IL, USA

US Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

* Corresponding authors

Proteome capacity constraints favor respiratory ATP generation

Abstract

Cells face competing metabolic demands. These include efficient use of both limited substrates
and limited proteome capacity, as well as flexibility to deal with different environments. Flexibility
requires spare enzyme capacity, which is proteome inefficient. ATP generation can occur via
fermentation or respiration. Fermentation is much less substrate-efficient, but often assumed to be
more proteome efficient [1–3], thereby favoring fast-growing cells engaging in aerobic glycolysis [4–
8]. Here, however, we show that mitochondrial respiration is actually more proteome-efficient than
aerobic glycolysis. Instead, aerobic glycolysis arises from cells maintaining the flexibility to grow
also anaerobically. These conclusions emerged from an unbiased assessment of metabolic
regulatory mechanisms, integrating quantitative metabolomics, proteomics, and fluxomics, of two
budding yeasts, *Saccharomyces cerevisiae* and *Issatchenkia orientalis*, the former more
fermentative and the latter respiratory. Their energy pathway usage is largely explained by
differences in proteome allocation. Each organism's proteome allocation is remarkably stable
across environmental conditions, with metabolic fluxes predominantly regulated at the level of
metabolite concentrations. This leaves extensive spare biosynthetic capacity during slow growth
and spare capacity of their preferred bioenergetic machinery when it is not essential. The greater
proteome-efficiency of respiration is also observed in mammals, with aerobic glycolysis occurring
in yeast or mammalian cells that maintain a fermentation-capable proteome conducive to both
aerobic and anaerobic growth.

Introduction

Metabolism is subject to physical constraints. Given the law of conservation of matter, to maintain
homeostasis, limited metabolic inputs must balance with outputs ('flux balance'). These outputs
include high energy cofactors (most importantly ATP), building blocks for cell replication, and
waste. The resources, including physical space and protein synthesis capacity, to sustain these
fluxes are also limited. Thus, cells are under pressure to produce their required metabolic fluxes
efficiently. As proteins catalyze most metabolic reactions and comprise the majority of biomass in

many cell types, the challenges of limited biosynthetic machinery and physical space can be
viewed largely as constraints on proteome capacity [2,3,9–17].

To manage proteome capacity, cells tailor protein expression to their conditions [18–21]. For example,
rapidly growing cells express copious ribosomes [18,22,23]. This requires less expression of other
protein types, such as anabolic and catabolic enzymes (e.g. those required for assimilating limiting
quantities of nitrogen [24], or breaking down non-preferred carbon sources [21,25]). Conversely, under
less favorable conditions, ribosome expression falls to make room for other proteins. Perfect
proteome tailoring, however, is not necessarily feasible or desirable [26–30], as proteome remodeling
is expensive and spare enzyme capacity can allow cells to quickly ramp up fluxes to deal with
changing conditions.

An important case of metabolic tailoring involves energy production from fermentation versus
respiration. Respiration is by far more energy-efficient, producing roughly 10-fold more ATP per
glucose [31]. Nevertheless, many organisms ferment, producing organic waste, even when oxygen is
available ('aerobic glycolysis'). Aerobic glycolysis is associated with fast growing cells including
bacteria, yeast and cancer cells [4–8]. Indeed, as their growth accelerates, both *Escherichia coli* and
S. cerevisiae switch from respiration to fermentation [2,32].

Why do cells engage in aerobic glycolysis when it is so much less energy-efficient than respiration?
One possibility involves a 'rate-yield tradeoff' [1–3,33]. More specifically, it is often believed that
fermentation is capable of producing ATP faster per unit enzyme expression, i.e. more 'proteome
efficient' [9–17]. The proteome efficiency of glycolysis versus respiration, however, has not been
carefully experimentally tested in eukaryotes.

Here we examine this question, building from an extensive systems-level analysis of metabolism
in two evolutionarily distant budding yeasts (separated by 200 million years [34]): *S. cerevisiae*
(Baker's yeast) and *I. orientalis* (also known as *Candida krusei* and *Pichia kudriavzevii*, a species
abundant in fermented food, fruit, and soil with favorable properties for bioengineering) [35–40]. We
find that, across diverse environmental conditions spanning a broad range of growth rates, the
proteome of each yeast varies only modestly, with metabolic flux explained primarily by

metabolite levels. Across the yeasts, however, metabolic flux differences are explained mainly by

proteome allocation, with *I. orientalis* expressing more respiratory enzymes and outcompeting *S.*

cerevisiae across diverse aerobic contexts. Both in *I. orientalis* and in respiring *S. cerevisiae*,

quantitative measurements show that respiration is actually several-fold more proteome efficient

than glycolysis. Similar results are attained in mammalian tissues and cancer cells. The origin of

aerobic glycolysis accordingly does not lie in proteome efficiency, but rather in a proteome

hedging strategy where cells maintain spare glycolytic capacity in preparation for potential future

hypoxia.

Results

Metabolic fluxes in *S. cerevisiae* and *I. orientalis*

We first characterized aerobic growth and metabolism of *S. cerevisiae* and *I. orientalis* in glucose

minimal medium (Fig. S1a). *I. orientalis* grows faster than *S. cerevisiae* (μ = 0.52 vs. 0.39 h^{-1}),

respires much more, consumes less glucose, and excretes less ethanol (Fig. S1a). We then resolved

flux through the entire metabolic network with ^{13}C metabolic flux analysis (Fig. 1). Specifically,

we developed genome-scale metabolic models of both yeasts including complete atom mapping

[41]. The models were then constrained by flux balance and experimentally derived extracellular

fluxes, biomass fluxes, and isotope labeling (from two distinct ^{13}C-tracer strategies: [1,2-

$^{13}C_2$]glucose and [U-$^{13}C_6$]glucose, each at 1:1 molar ratio with unlabeled glucose). This enabled

comprehensive yeast metabolic flux analysis, at a rigor previously achieved only in prokaryotes

[41,42].

^{13}C-tracing resolved key internal flux branch points. For example, [1,2-$^{13}C_2$]glucose revealed

markedly higher [M+1] pyruvate labeling in *I. orientalis* (Fig. S1b), reflecting greater oxidative

pentose phosphate pathway (PPP) flux in this yeast species (Fig. 1). The same tracer also reveals

greater [M+1] tricarboxylic acid (TCA) cycle intermediates in *I. orientalis* (Fig. S1c), consistent

with higher oxidative TCA cycle flux relative to *S. cerevisiae*, where clockwise TCA flux was

truncated at α-ketoglutarate (Fig. 1). The greatest difference between the two yeasts is in the way

glucose is catabolized. Namely, *S. cerevisiae* prefers carbon-inefficient fermentation, and

correspondingly makes most ATP from glycolysis and consumes most NADH via ethanol

fermentation (Fig. 1, Fig. S1d). In contrast, *I. orientalis* prefers respiration for ATP production,

and re-oxidizes NADH by a blend of complex I and the quinone oxidoreductase Nde1, the

knockout of which impairs *I. orientalis* but not *S. cerevisiae* growth [43] (Fig. 1, Fig. S1e).

Flux control across environmental conditions

When nutrients become scarce, cells adjust metabolic fluxes and growth. Such fluxes can be

controlled through enzyme concentration ($k_{cat}[E]$ in the Michaelis-Menten kinetics), active site

occupancy ($[S]/([S]+K_m)$), or allosteric regulation (Fig. 2a). To assess flux control mechanisms in

S. cerevisiae and *I. orientalis*, we grew each yeast in aerobic chemostats at diverse growth rates

controlled by limiting glucose, ammonia, or phosphate availability (Fig. 2b). In each nutrient

environment, we measured enzyme concentrations via quantitative proteomics, metabolite

concentrations via metabolomics, and metabolic fluxes via ^{13}C-informed fluxomics (Fig. 2c).

Fluxes aligned remarkably closely with growth rate in both yeasts (Fig. S2a), with the exception

of metabolic switching to respiration and the PPP (relative to glycolysis) in glucose-limited *S.*

cerevisiae, which renders glucose-limited *S. cerevisiae* metabolically similar to *I. orientalis* (Fig.

S2, b and c). On average, 53% of flux variation in *S. cerevisiae* and 71% in *I. orientalis* was

explained by growth rate alone.

The corresponding metabolomics and proteomics data provide a valuable resource for

understanding the biochemical basis by which these fluxes are achieved, especially in the non-

model yeast. For example, they can be assessed on a reaction-by-reaction basis to identify

physiologically meaningful metabolic regulators [44] (Fig. S3a showing glyceraldehyde-3-phosphate

dehydrogenase, or GAPD, as an example). We were able to identify allosteric regulation in 19 out

of 51 examined reactions in *I. orientalis* (Supplementary Table). Seven of these regulations have

also been reported in *S. cerevisiae*, including classical ones such as citrate inhibition of

phosphofructokinase [31] and fructose-1,6-bisphosphate activation of pyruvate kinase [45]. Our

analysis also revealed multiple previously unappreciated regulatory interactions. For example,

ATP inhibits *I. orientalis* GAPD, a novel interaction that we biochemically verified (Fig. S3, a-c).

Overall, the integration of *in vivo* enzyme and metabolite concentrations via Michaelis-Menten

kinetics explained the vast majority of flux variation across physiological conditions (Fig. 2d).

This reflects enzyme concentration, active site occupancy, and allosteric regulation by metabolites

collectively accounting for most yeast flux control, without the need to invoke other mechanisms

like enzyme covalent modification or localization.

Flux control by enzyme concentration

Cells contain extensive programs for regulating protein levels. Across nutrient conditions, however,

we observed remarkably stable enzyme concentrations. In contrast, both metabolic fluxes and

metabolite concentrations varied much more than enzymes (Fig. 2c, Fig. S2a). For some reactions,

enzyme levels even show negative correlation with flux (negative Pearson's R in Fig. 2d with

enzyme only). We assessed the extent of physiological flux control residing in enzymes and

metabolites based on their metabolic leverage, the product of their physiological concentration

variation across conditions and their flux control coefficient based on the best-supported kinetic

model from the above quantitative analysis of physiological metabolic regulation. Across 51

evaluable reactions, we found that, across nutrient conditions, metabolites exert much more

metabolic leverage than enzymes (Fig. 2e, Fig. S3d).

Indeed, within both yeast species, flux changes across physiological conditions correlate better

with pathway substrate concentration changes than pathway enzyme concentration changes (based

on median of fold change across pathway components) (Fig. 2, f-i). The maintenance of enzyme

concentrations with reduced growth and metabolic flux suggests substantial spare enzyme capacity,

which may facilitate rapid growth acceleration when nutrient conditions improve [26,27,46,47].

In contrast to metabolite-dominant flux control within each yeast in response to changing nutrient

environment, flux differences between *S. cerevisiae* and *I. orientalis* strongly aligned with enzyme

concentrations (Fig. 2, j-k). Given the 200 million years of evolution separating these two species

[34], we expected that there might be substantial differences in enzyme properties that change

metabolic flux between the two organisms. Highly expressed proteins, including central metabolic

enzymes, however remained strongly conserved between the two yeasts at the protein sequence

level (Fig. S4). Correspondingly, enzyme abundances account for a large fraction of flux variation,

including the greater glycolysis flux in *S. cerevisiae* and faster TCA turning and oxidative

phosphorylation in *I. orientalis* (Fig. 2k). Thus, within the tested yeast species, flux is

predominately regulated at the level of metabolite concentrations and active site occupancy. In

contrast, flux differences between these yeast species is predominately explained by enzyme
concentrations.
181
Proteome-efficiency of ATP generation
We next examined overall proteome allocation of both yeasts with absolute proteomic
quantification calibrated by UPS2 standard, and found that metabolic genes (enzymes, transporters,
and mitochondrial proteins) together accounted for just over half of the proteome in both species,
with the other major proteome sectors being translation (mostly ribosomes) and transcriptional
machinery (Fig. 3a). The fractional proteome allocation to these three major sectors was nearly
identical across the two yeasts. The major difference was within the metabolic sector, which had
three major components: anabolic, glycolytic, and respiratory (the latter being composed of
mitochondrial, TCA, and oxidative phosphorylation proteins). The anabolic enzymes constituted
a similar proteome fraction in both species, but there was a major reallocation between the two
other sectors: *S. cerevisiae* expresses more glycolytic proteins, and *I. orientalis* more respiratory
proteins.
194
Together, our absolute proteome quantitation and flux analyses enabled quantification of ATP
production per protein mass (i.e. proteome efficiency) of glycolysis and respiration in both species.
We obtained the ATP flux from the genome-scale model, which included a mechanistic ratio of 3
ATP produced for every 10 protons translocated by ATP synthase [48]. Including both TCA and
oxidative phosphorylation proteins (but not other mitochondrial proteins) as the proteome cost of
respiration, we found that, in batch culture, respiration is more proteome-efficient than glycolysis
in both yeasts (Fig. 3b).
202
Despite comparing favorably to glycolysis, respiration in glucose-rich batch-cultured *S. cerevisiae*
was much less proteome-efficient than in *I.orientalis.* Besides the absence of proton-pumping
complex I in *S. cerevisiae*, we hypothesized that this also reflects spare respiratory capacity when
glucose is abundant. Consistent with this, the proteome-efficiency of *S. cerevisiae* respiration
increased (and of glycolysis fell) under glucose limitation (Fig. 3c, Fig. S5). In contrast, since *I.*
orientalis defaults to respiration even when glucose is abundant, proteome-efficiency was
unaffected by glucose availability.

Part of glycolytic and TCA flux is diverted to biosynthesis, and glycolytic flux is needed to generate respiratory substrate. We further assessed proteome efficiency of ATP generation of fermentation (turning glucose to ethanol) and respiration (turning glucose to CO_2) by calculating a flux-partitioned proteome cost [2], which counts glycolytic proteins in the cost of respiration and discounts flux diverted to other pathways. In both yeasts, this flux-partitioned analysis identified respiration as the more proteome-efficient ATP production pathway (Fig. S6, a-b). Similarly, even if counting all mitochondrial proteins into respiration's proteome cost, respiration remains more proteome-efficient than glycolysis in *I. orientalis* and glucose-limited *S. cerevisiae* (Fig. S6, c-d).

Benefit of aerobic glycolysis

In *S. cerevisiae*, rate-yield tradeoff was believed to underlie the switching to carbon-inefficient fermentation at faster growth [14–17]. Our data reveals, however, that respiration is both more energy- and proteome-efficient than glycolysis. Such efficiency would be expected to lead to greater fitness.

Consistently, *I. orientalis* outcompetes *S. cerevisiae* in co-culture under conditions requiring respiratory ATP production (ethanol, glucose limitation). Importantly, however, it also outcompetes under conditions where fermentation is a viable strategy (abundant glucose, nitrogen limitation, phosphorus limitation, and even sucrose as the sole carbon source, which *S. cerevisiae* can ferment on while *I. orientalis* alone cannot metabolize and presumably takes in glucose and fructose liberated by *S. cerevisiae*) (Fig. 4a, Fig. S7).

If respiration is both more energy- and proteome-efficient than glycolysis, why does aerobic glycolysis occur? One possibility is the production of a toxic product that impairs competitors: ethanol [49,50]. Another is carbon resource competition, essentially quick uptake of glucose [1,33,51]. These might help *S. cerevisiae* compete with bacteria, but did not against *I. orientalis* (Fig. 4a).

We wondered whether the benefit of aerobic glycolysis might instead not be during aerobic growth, but rather in hedging for oxygen limitation (hypoxia) [13]. *S. cerevisiae* was repeatedly exposed to oxygen limitation during human baking and winemaking [52]. But oxygen limitation can also occur readily naturally, due to oxygen's limited solubility ($c_{O2} \approx 230\,\mu$mol/L) and slow diffusion in water

(D ≈ 2.3×10^{-9} m^2/s). We measured the dissolved oxygen at the bottom of unstirred *S. cerevisiae*
and *I. orientalis* cultures, and found that oxygen depletion occurred at relatively low culture
density (Fig. S8). Notably, *S. cerevisiae* outcompeted *I. orientalis* in fully or cyclically oxygen-
depleted co-cultures (Fig. 4a, Fig. S7). Consistently, we observed about 60% growth rate reduction
in *I. orientalis* but not *S. cerevisiae* upon oxygen depletion or pharmacological inhibition of
electron transport chain complex III by antimycin (Fig. 4b). Both oxygen depletion and antimycin
increased glucose uptake rate by more than two-fold in *I. orientalis* (Fig. 4c), which is mediated
by about 3-fold higher expression of glycolytic proteins (Fig. 4d). Notably, this glycolytic protein
expression came at the expense of ribosomal proteins, consistent with the growth defect in
anaerobic *I. orientalis* (Fig. 4d, Fig. S9).
251

Proteomic hedging and aerobic glycolysis

253   Both *I. orientalis* and *S. cerevisiae* can tailor their respiratory versus glycolytic enzyme expression
254   to environmental conditions. But this tailoring is incomplete: *I. orientalis* partially retains
255   respiratory enzyme and mitochondrial protein expression in hypoxia (Fig. 4d), while *S. cerevisiae*
256   retains high glycolytic enzymes in aerobic conditions (Fig. 2g).
257

To explore the consequences of incomplete proteome tailoring, we assembled a coarse-grained
quantitative model of yeast growth and metabolism, where growth is limited both by ATP
generation (fermentative or respiratory) and by translational machinery, jointly constrained by
proteome capacity (Fig. 4e and Extended Data Note). Growth optimization is performed to find
the optimal respiratory and glycolytic proteome allocation (f_R and f_G, respectively). This minimal
model captures the proteome tradeoff between optimal aerobic and anaerobic growth (Fig. 4e).
While optimal aerobic growth is achieved via respiratory ATP production, when the proteome is
constrained to always contain enough glycolytic enzyme for rapid anaerobic growth, aerobic
glycolysis emerges as an optimal strategy (Fig. 4f).
267

Mammalian ATP generation

We were curious if the greater proteome efficiency of respiration generalizes from yeast to
mammals. To investigate this, we quantified the proteome efficiency of glycolysis and respiration
in cultured cancer cells and mouse tissues (Fig. 5). ATP flux was from previous reports or

estimated based on reported oxygen consumption rates [54,55] (Fig. 5a). The proteome fraction
allocated to glycolysis and respiration was computed from published proteomics data [56,57], which
shows that mouse tissues in general have greater respiratory proteome capacity like in *I. orientalis*,
whereas cancer cell lines have more glycolytic proteome like in *S. cerevisiae* (Fig. 5b). Overall,
the proteome efficiency of both glycolysis and respiration was lower in mammals than in yeast,
consistent with mammals being under less stringent selection for proteome efficiency.
Nevertheless, in both cultured cancer cells and *in vivo* tissues, respiration was the more proteome-
efficient ATP generation pathway (Fig. 5c). Thus, in both yeast and mammals, mitochondrial
respiration is more proteome-efficient than glycolysis.

Discussion

Here we report in-depth proteomic, metabolomic, and metabolic flux characterization of two
divergent budding yeasts across various environmental conditions. The resulting data reveal
principles of yeast metabolism and its regulation. Prior integrative 'omic analysis of *E. coli* and *S.*
cerevisiae concluded that metabolism is substantially 'self-regulated', i.e. that changes in
metabolic flux are caused more by metabolites themselves than transcriptional and translational
reprogramming of enzyme levels [44,58–60]. This conclusion is reinforced by analogous analysis of *I.*
orientalis here, which shows even less proteome variation across most environmental conditions
than *S. cerevisiae,* and yet greater dominance of metabolic flux control by metabolites themselves.

In contrast to the limited impact of the proteome on flux control within each species, across the
two species, metabolic differences are mainly encoded by protein abundances. Given that these
two yeasts diverged roughly 200 million years ago, the ability to explain most of their metabolic
differences through enzyme concentrations – rather than changes in the properties of enzymes
themselves – is notable, and speaks to the importance of proteome allocation in driving metabolic
divergence even across long timescales [61].

The most striking metabolic difference between *I. orientalis* and *S. cerevisiae* is that, in the
presence of abundant glucose, the former respires while the latter engages in aerobic glycolysis.
We show that, across a wide range of aerobic conditions, the more respiratory yeast grows faster
and has superior competitive fitness. This aligns with respiration requiring less of the cell's

precious proteome capacity to achieve the same growth-required ATP flux. Quantitative analysis
of mammalian cancer cells and tissues demonstrates that respiration is also more proteome-
efficient than glycolysis in mammals.
306
Prior careful evaluation of proteome efficiency in *E. coli* reached a seemingly opposite conclusion,
finding that acetate fermentation is favored for its proteome efficiency [2]. Acetate overflow
metabolism in *E. coli*, however, involves a blend of glycolytic and respiratory ATP generation
with 4 NADH feeding into the electron transport chain for each glucose. This provides an ATP
yield of about 12 per glucose, the majority of which is made via the oxidative phosphorylation
(compared to 2 ATP per glucose in yeast or mammalian aerobic glycolysis). Thus, aerobic
'fermentation' in *E. coli* is proteome efficient only because it generates substantial respiratory ATP.
314
Overall, supported by our data in *S. cerevisiae* and *I. orientalis*, we propose that cells of a given
type tend to have a characteristic metabolic proteome that varies only modestly across conditions.
In this nearly fixed enzyme network, changing substrate levels induce different fluxes, providing
metabolic flexibility without the need for extensive proteome remodeling. A benefit of such
proteome constancy is that cells are prepared in advance for changing metabolic environments.
One of the most important metabolic fluctuations cells face is shifting oxygen availability [62]. We
thus posit that aerobic glycolysis occurs not because it is beneficial per se, but as a side effect of
maintaining a fermentative proteome that effectively supports both aerobic and anaerobic growth.
323

Reference

1. Pfeiffer, T., Schuster, S. & Bonhoeffer, S. Cooperation and Competition in the Evolution of

ATP-Producing Pathways. *Science* **292**, 504–507 (2001).

2. Basan, M. *et al.* Overflow metabolism in Escherichia coli results from efficient proteome

allocation. *Nature* **528**, 99–104 (2015).

3. Molenaar, D., van Berlo, R., de Ridder, D. & Teusink, B. Shifts in growth strategies reflect

tradeoffs in cellular economics. *Mol. Syst. Biol.* **5**, 323 (2009).

4. Crabtree, H. G. Observations on the carbohydrate metabolism of tumours. *Biochem. J.* **23**,

536–545 (1929).

5. De Deken, R. H. The Crabtree Effect: A Regulatory System in Yeast. *J. Gen. Microbiol.* **44**,

149–156 (1966).

6. Vander Heiden, M. G., Cantley, L. C. & Thompson, C. B. Understanding the Warburg Effect:

The Metabolic Requirements of Cell Proliferation. *Science* **324**, 1029–1033 (2009).

7. DeBerardinis, R. J. & Chandel, N. S. We need to talk about the Warburg effect. *Nat. Metab.*

2, 127–129 (2020).

8. Wolfe, A. J. The Acetate Switch. *Microbiol. Mol. Biol. Rev.* **69**, 12–50 (2005).

9. Adadi, R., Volkmer, B., Milo, R., Heinemann, M. & Shlomi, T. Prediction of Microbial

Growth Rate versus Biomass Yield by a Metabolic Network with Kinetic Parameters. *PLoS*

Comput. Biol. **8**, e1002575 (2012).

10. O'Brien, E. J., Lerman, J. A., Chang, R. L., Hyduke, D. R. & Palsson, B. Ø. Genome- scale

models of metabolism and gene expression extend and refine growth phenotype prediction.

Mol. Syst. Biol. **9**, 693 (2013).

11. Mori, M., Hwa, T., Martin, O. C., De Martino, A. & Marinari, E. Constrained Allocation Flux

Balance Analysis. *PLOS Comput. Biol.* **12**, e1004913 (2016).

12. Salvy, P. & Hatzimanikatis, V. The ETFL formulation allows multi-omics integration in

thermodynamics-compliant metabolism and expression models. *Nat. Commun.* **11**, 30 (2020).

13. Wortel, M. T., Noor, E., Ferris, M., Bruggeman, F. J. & Liebermeister, W. Metabolic enzyme

cost explains variable trade-offs between microbial growth rate and yield. *PLOS Comput. Biol.*

14, e1006010 (2018).

14. Sánchez, B. J. *et al.* Improving the phenotype predictions of a yeast genome-scale metabolic

model by incorporating enzymatic constraints. *Mol. Syst. Biol.* **13**, 935 (2017).

15. Oftadeh, O. *et al.* A genome-scale metabolic model of Saccharomyces cerevisiae that

integrates expression constraints and reaction thermodynamics. *Nat. Commun.* **12**, 4790

(2021).

16. Elsemman, I. E. *et al.* Whole-cell modeling in yeast predicts compartment-specific proteome

constraints that drive metabolic strategies. *Nat. Commun.* **13**, 801 (2022).

17. Chen, Y. & Nielsen, J. Energy metabolism controls phenotypes by protein efficiency and

allocation. *Proc. Natl. Acad. Sci.* **116**, 17592–17597 (2019).

18. You, C. *et al.* Coordination of bacterial proteome with metabolism by cyclic AMP signalling.

Nature **500**, 301–306 (2013).

19. Li, G.-W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying Absolute Protein Synthesis

Rates Reveals Principles Underlying Allocation of Cellular Resources. *Cell* **157**, 624–635

(2014).

20. Dekel, E. & Alon, U. Optimality and evolutionary tuning of the expression level of a protein.

Nature **436**, 588–592 (2005).

21. Keren, L. *et al.* Massively Parallel Interrogation of the Effects of Gene Expression Levels on

Fitness. *Cell* **166**, 1282-1294.e18 (2016).

22. Bosdriesz, E., Molenaar, D., Teusink, B. & Bruggeman, F. J. How fast-growing bacteria
robustly tune their ribosome concentration to approximate growth-rate maximization. *FEBS J.*
282, 2029–2044 (2015).

23. Peebo, K. *et al.* Proteome reallocation in Escherichia coli with increasing specific growth rate.
Mol. Biosyst. **11**, 1184–1193 (2015).

24. Hui, S. *et al.* Quantitative proteomic analysis reveals a simple strategy of global resource
allocation in bacteria. *Mol. Syst. Biol.* **11**, 784 (2015).

25. Gancedo, J. M. Yeast Carbon Catabolite Repression. *Microbiol. Mol. Biol. Rev.* **62**, 334–361
(1998).

26. Metzl-Raz, E. *et al.* Principles of cellular resource allocation revealed by condition-dependent
proteome profiling. *eLife* **6**, e28034 (2017).

27. O'Brien, E. J., Utrilla, J. & Palsson, B. O. Quantification and Classification of E. coli Proteome
Utilization and Unused Protein Costs across Environments. *PLOS Comput. Biol.* **12**, e1004998
(2016).

28. Yu, R. *et al.* Nitrogen limitation reveals large reserves in metabolic and translational capacities
of yeast. *Nat. Commun.* **11**, 1881 (2020).

29. Postma, E., Verduyn, C., Scheffers, W. A. & Van Dijken, J. P. Enzymic analysis of the crabtree
effect in glucose-limited chemostat cultures of Saccharomyces cerevisiae. *Appl. Environ.*
Microbiol. **55**, 468–477 (1989).

30. Christodoulou, D. *et al.* Reserve Flux Capacity in the Pentose Phosphate Pathway Enables
Escherichia coli's Rapid Response to Oxidative Stress. *Cell Syst.* **6**, 569-578.e7 (2018).

31. Lehninger, A. L., Nelson, D. L. & Cox, M. M. *Lehninger principles of biochemistry.* (W.H.
Freeman, 2005).

32. Van Hoek, P., Van Dijken, J. P. & Pronk, J. T. Effect of Specific Growth Rate on Fermentative

Capacity of Baker's Yeast. *Appl. Environ. Microbiol.* **64**, 4226–4233 (1998).

33. Bachmann, H. *et al.* Availability of public goods shapes the evolution of competing metabolic

strategies. *Proc. Natl. Acad. Sci.* **110**, 14302–14307 (2013).

34. Shen, X.-X. *et al.* Tempo and Mode of Genome Evolution in the Budding Yeast Subphylum.

Cell **175**, 1533-1545.e20 (2018).

35. Radecka, D. *et al.* Looking beyond *Saccharomyces* : the potential of non-conventional yeast

species for desirable traits in bioethanol fermentation. *FEMS Yeast Res.* **15**, fov053 (2015).

36. Fatma, Z., Schultz, J. C. & Zhao, H. Recent advances in domesticating non-model

microorganisms. *Biotechnol. Prog.* **36**, (2020).

37. Xiao, H., Shao, Z., Jiang, Y., Dole, S. & Zhao, H. Exploiting Issatchenkia orientalis SD108

for succinic acid production. *Microb. Cell Factories* **13**, 121 (2014).

38. Suthers, P. F. *et al.* Genome-scale metabolic reconstruction of the non-model yeast

Issatchenkia orientalis SD108 and its application to organic acids production. *Metab. Eng.*

Commun. **11**, e00148 (2020).

39. Cao, M. *et al.* A genetic toolbox for metabolic engineering of Issatchenkia orientalis. *Metab.*

Eng. **59**, 87–97 (2020).

40. Douglass, A. P. *et al.* Population genomics shows no distinction between pathogenic Candida

krusei and environmental Pichia kudriavzevii: One species, four names. *PLOS Pathog.* **14**,

e1007138 (2018).

41. Gopalakrishnan, S. & Maranas, C. D. 13C metabolic flux analysis at a genome-scale. *Metab.*

Eng. **32**, 12–22 (2015).

42. McCloskey, D., Young, J. D., Xu, S., Palsson, B. O. & Feist, A. M. Modeling Method for
Increased Precision and Scope of Directly Measurable Fluxes at a Genome-Scale. *Anal. Chem.*
88, 3844–3852 (2016).

43. Luttik, M. A. H. *et al.* The Saccharomyces cerevisiae NDE1 and NDE2 Genes Encode
Separate Mitochondrial NADH Dehydrogenases Catalyzing the Oxidation of Cytosolic
NADH. *J. Biol. Chem.* **273**, 24529–24534 (1998).

44. Hackett, S. R. *et al.* Systems-level analysis of mechanisms regulating yeast metabolic flux.
Science **354**, aaf2786–aaf2786 (2016).

45. Schomburg, D., Schomburg, I. & Chang, A. *Springer handbook of enzymes*. (Springer, 2007).

46. Reznik, E. *et al.* Genome-Scale Architecture of Small Molecule Regulatory Networks and the
Fundamental Trade-Off between Regulation and Enzymatic Activity. *Cell Rep.* **20**, 2666–2677
(2017).

47. Jansen, M. L. A. *et al.* Prolonged selection in aerobic, glucose-limited chemostat cultures of
Saccharomyces cerevisiae causes a partial loss of glycolytic capacity. *Microbiology* **151**,
1657–1669 (2005).

48. Symersky, J. *et al.* Structure of the c10 ring of the yeast mitochondrial ATP synthase in the
open conformation. *Nat. Struct. Mol. Biol.* **19**, 485–491 (2012).

49. Zhou, N. *et al.* Coevolution with bacteria drives the evolution of aerobic fermentation in
Lachancea kluyveri. *PLOS ONE* **12**, e0173318 (2017).

50. Dashko, S., Zhou, N., Compagno, C. & Piškur, J. Why, when, and how did yeast evolve
alcoholic fermentation? *FEMS Yeast Res.* **14**, 826–832 (2014).

51. MacLean, R. C. & Gudelj, I. Resource competition and social conflict in experimental
populations of yeast. *Nature* **441**, 498–501 (2006).

52. Dekker, W. J. C., Wiersma, S. J., Bouwknegt, J., Mooiman, C. & Pronk, J. T. Anaerobic growth of *Saccharomyces cerevisiae* CEN.PK113-7D does not depend on synthesis or supplementation of unsaturated fatty acids. *FEMS Yeast Res.* **19**, foz060 (2019).

53. Martin, A. W. & Fuhrman, F. A. The Relationship between Summated Tissue Respiration and Metabolic Rate in the Mouse and Dog. *Physiol. Zool.* **28**, 18–34 (1955).

54. Zielinski, D. C. *et al.* Systems biology analysis of drivers underlying hallmarks of cancer cell metabolism. *Sci. Rep.* **7**, 41241 (2017).

55. Bartman, C. R. *et al. Slow TCA flux implies low ATP production in tumors.* http://biorxiv.org/lookup/doi/10.1101/2021.10.04.463108 (2021) doi:10.1101/2021.10.04.463108.

56. Wang, M., Herrmann, C. J., Simonovic, M., Szklarczyk, D. & Mering, C. Version 4.0 of PaxDb: Protein abundance data, integrated across model organisms, tissues, and cell-lines. *PROTEOMICS* **15**, 3163–3168 (2015).

57. Gholami, A. M. *et al.* Global Proteome Analysis of the NCI-60 Cell Line Panel. *Cell Rep.* **4**, 609–620 (2013).

58. Fendt, S. *et al.* Unraveling condition-dependent networks of transcription factors that control metabolic pathway activity in yeast. *Mol. Syst. Biol.* **6**, 432 (2010).

59. Millard, P., Smallbone, K. & Mendes, P. Metabolic regulation is sufficient for global and robust coordination of glucose uptake, catabolism, energy production and growth in Escherichia coli. *PLOS Comput. Biol.* **13**, e1005396 (2017).

60. Gerosa, L. *et al.* Pseudo-transition Analysis Identifies the Key Regulators of Dynamic Metabolic Adaptations from Steady-State Data. *Cell Syst.* **1**, 270–282 (2015).

61. Kito, K. *et al.* Yeast Interspecies Comparative Proteomics Reveals Divergence in Expression

Profiles and Provides Insights into Proteome Resource Allocation and Evolutionary Roles of

Gene Duplication. *Mol. Cell. Proteomics* **15**, 218–235 (2016).

62. Dewhirst, M. W., Cao, Y. & Moeller, B. Cycling hypoxia and free radicals regulate

angiogenesis and radiotherapy response. *Nat. Rev. Cancer* **8**, 425–437 (2008).

Acknowledgement

We thank the members of Rabinowitz lab for discussion on experiments and the manuscript; S.

Silverman and J. Avalos for yeast strains, L. Ryazanova for help with proteomics experiment, P.

F. Suthers for discussion on the genome-scale model, M. Gupta for discussion on protein

regulation, N. Piyush and Z. Zhang for advice on competitive fitness, and Z.-Y. Wu for

experimental support. This work was funded by DOE grant DE-SC0018260 and the DOE Center

for Advanced Bioenergy and Bioproducts Innovation (Award Number DE-SC0018420). Any

opinions, findings, conclusions or recommendations expressed in this publication are those of the

author(s) and do not necessarily reflect the views of the U.S. Department of Energy.

Author contributions

Y.S., M.W. and J.D.R. designed this study. Y.S. performed most of the experiments and data

analysis. H.V.D. designed and performed genome-scale metabolic flux analysis with the input

from Y.S., J.I.H., and C.D.M. E.C., H.B., and A.S. performed proteomics measurement. C.M.C.

performed nutrient limited culture and measurement. R.P.R. designed and performed enzyme

purification and qPCR. J.P. performed enzyme purification and competitive growth experiments.

Z.F. created mutant yeast strains with the input from H.Z., and contributed to enzyme purification.

S.D. and Y.Y. contributed to yeast growth measurement. V.T. contributed to enzyme purification.

T.X. contributed to metabolomics measurements. D.W. contributed to enzyme constrained

modeling. L.Y. contributed to oxygen consumption measurement. Y.S. and J.D.R. wrote the

manuscript with input from all co-authors.

491
492

493
494 **Figure 1. Genome-scale flux analysis shows more active respiratory metabolism in *I.***
495 ***orientalis*.**

496
Metabolic flux (in mmol/gDW/h) of *S. cerevisiae* (CEN.PK) and *I. orientalis* (SD108) in aerobic
exponential growth in YNB with 20 g/L glucose. Fluxes are best estimate from genome-scale [13]C-
informed metabolic flux analysis (MFA), with the input data including metabolite [13]C labeling
from two [13]C-glucose tracers (each with n = 3 or 4 biological replicates) and consumption and
excretion fluxes (at least n = 3 biological replicates). Color represents metabolic pathways:
glycolysis in red, oxidative phosphorylation (ox phos) in blue, TCA in green, and PPP in orange.
Numbers represents flux in mmol/h/gDW.

Figure 2. Flux change is explained by metabolites within yeast species, and by enzymes across

yeasts. (Legend continues on next page.)

Figure 2. Flux change is explained by metabolites within yeast species, and by enzymes across

yeasts.

(a) Flux change can be achieved through change in either enzyme or substrate level via Michaelis-

Menten kinetics.

(b) Multi-omics data were obtained in steady-state yeast grown in nutrient-limited continuous

culture or nutrient-replete batch culture.

(c) Genome-wide metabolic flux, protein abundance, and metabolite concentrations in *I. orientalis*

across growth conditions. Values are normalized to the geometric mean across all the conditions.

Metabolomics, mean of n = 3 technical replicates (independent sampling from continuous culture).

Proteomics, n = 1 biological replicate. Fluxomics, best estimates from ^{13}C-MFA similar to Fig. 1.

(d) Distribution of Pearson's R for 51 reactions between measured flux and flux predicted from

Michaelis-Menten kinetics accounting for different variables: concentration of enzyme, reactant,

and best data-supported allosteric regulator (if any).

(e) Partition of metabolic control among enzyme, reactants, and regulator for 51 reactions in *I.*

orientalis.

(f-k) Correlation between flux and metabolite concentration (f-h) or between flux and enzyme

concentration (i-k). Data within an organism (f,g,i,j) compares nutrient limited to batch conditions.

Data across organisms (h,k) compares *S. cerevisiae* to *I. orientalis*. Each point represents median

flux and concentration fold change for the pathway. Pearson's R and p value are shown. Symbols

in (h, k) are diamonds, CEN.PK in batch culture; circle, FY4 in batch culture; triangle, FY4 in

nutrient limitation at 0.22 h^{-1}. Black line shows slope = 1. Other pathways (folate, sugar, nucleic

acid, lipid, amino acid) are plotted in different shades of grey.

Figure 3. Proteome efficiency across nutrient conditions in *S.cerevisiae* and *I. orientalis*.

(a) Proteome allocation of *S. cerevisiae* (CEN.PK) and *I. orientalis* in exponentially growing batch culture. Mean ± SE, n = 4 biological replicates.

(b) ATP fluxes (from [13]C-informed MFA), proteome mass fraction (of whole cell dry weight), and proteome efficiency for glycolysis and respiration in exponentially growing aerobic glucose-fed batch culture. ATP fluxes are shown as mean ± SE based on [13]C-MFA confidence interval. Proteome efficiency is shown as mean ± SE, with error propagated from flux and proteome fraction measurements.

(c) Proteome efficiency of respiration and glycolysis across different nutrient conditions in *S. cerevisiae* (left) and *I. orientalis* (right). For raw data, see Fig. S5. Solid line shows linear regression in glucose replete conditions (Batch, N-limit, and P-limit). Dashed line is glucose-depleted conditions (C-limit). P values are from ANOVA of linear model.

Figure 4. Aerobic glycolysis emerges from anaerobically primed proteome.

(a) Fitness of *I. orientalis* and *S. cerevisiae* measured in competitive co-culture. Relative abundance of the two yeasts was measured by qPCR at 4 to 6 time points and used to obtain fitness. See method for details. Mean ± SE, n = 3 or 4 biological replicates.

(b-c) Specific growth rates (b) and glucose consumption rates (c) for batch-cultured *S. cerevisiae* and *I. orientalis* with oxygen (blue), without oxygen (light grey), and with 10 µM antimycin (dark grey). Mean ± SE, n = 6 or 7 biological replicates.

(d) Proteome allocation in *I. orientalis* in above conditions. Mean ± SE, n=3 biological replicates. Arrows show fold change in translational and glycolytic proteome compared to aerobic condition.

(e) Respiro-fermentative growth rate (µ) was predicted from a proteome-constrained coarse-grained model parameterized with proteome efficiency measured from *S. cerevisiae*. Glycolytic (f_G) and respiratory proteome fraction (f_R), are mass fractions of whole cell dry weight. Optimal proteome fractions in aerobic and anaerobic condition were indicated as stars. Measured proteome

fractions in glucose-fed batch cultures of *I. orientalis* (aerobic, $+O_2$; or anaerobic, $-O_2$) and *S.*

cerevisiae (aerobic, $+O_2$) are shown in circles.

(f) Experimental glucose consumption (J_{GLC}) and ethanol excretion (J_{ETOH}) rates (symbols, in

mmol/h/gDW) and prediction from proteome-constrained model (lines) under high (*S. cerevisiae*)

or low (*I. orientalis*) glycolytic capacity (f_G relative to its anaerobic optimum, $f_{G,anae}$). Literature

data was obtained from Van Hoek 1998 [32].

Figure 5. Respiratory ATP production is more proteome efficient than glycolysis in mammals.

(a) Ratio between glycolytic and respiratory ATP production in *I. orientalis*, *S. cerevisiae*, NCI60 cancer cell lines, and mouse tissues. For yeasts, each data point represents a nutrient condition. For cancer cell lines and mouse tissues, each point represents an individual cell line or tissue.

(b) As in (a), for proteome allocation.

(c) Corresponding glycolytic versus respiratory proteome efficiency.