Tup1 is Required for Transcriptional Repression Necessary in Quiescence in *S. cerevisiae*

Thomas B. Bailey[1], Phaedra A. Whitty[1], Eric U. Selker[1], Jeffrey. N. McKnight[1], Laura E. McKnight[1,*]

[1] Institute of Molecular Biology, University of Oregon, Eugene OR 97403, USA
[*] Correspondence: lmcknigh@uoregon.edu

Abstract

Upon glucose starvation, *S. cerevisiae* shows a dramatic alteration in transcription, resulting in wide-scale repression of most genes and activation of some others. This coincides with an arrest of cellular proliferation. A subset of such cells enters quiescence, a reversible non-dividing state. Here, we demonstrate that the conserved transcriptional corepressor Tup1 is critical for transcriptional repression after glucose depletion. We show that Tup1-Ssn6 binds new targets upon glucose depletion, where it remains as the cells enter the G0 phase of the cell cycle. In addition, we show that Tup1 represses a variety of glucose metabolism and transport genes. We explored how Tup1 mediated repression is accomplished and demonstrated that Tup1 coordinates with the Rpd3L complex to deacetylate H3K23. We found that Tup1 coordinates with Isw2 to affect nucleosome positions at glucose transporter HXT family genes during G0. Finally, microscopy revealed that a quarter of cells with a Tup1 deletion contain multiple DAPI puncta. Taken together, these findings demonstrate the role of Tup1 in transcriptional reprogramming in response to environmental cues leading to the quiescent state.

Keywords: Groucho/Gro, Tle, Grg , Diauxic Shift, Metabolism, Histone Acetylation, Nucleosome Positioning, Rpd3, Xbp1, Isw2

Introduction

Cellular quiescence is a non-proliferative cell state that appears conserved across all life (Marescal & Cheeseman, 2020; Rittershaus et al., 2013; van Velthoven & Rando, 2019). Quiescent cells are distinct from other non-dividing cells, such as senescent and terminally differentiated cells, in that quiescent cells can return to a proliferative state if the right conditions are met. Quiescence is necessary for the survival of unicellular organisms in environments that are unfavorable for proliferation; in fact, most microbes in nature are quiescent (O'Farrell, 2011; Valcourt et al., 2012). In multicellular eukaryotes, quiescence plays a critical role in various cell types including stem cells, fibroblasts, and lymphocytes (Tümpel & Rudolph, 2019; van Velthoven & Rando, 2019; Yao, 2014). The inability to properly regulate quiescence can lead to cancer; additionally, cancer stem cells can become quiescent, leading to drug resistance and the ability to initiate relapse at a later time (Vallette et al., 2019). All quiescent cells have conserved characteristics including transcriptional repression, increased cell density, translational repression, and decreased cell metabolism (de Virgilio, 2012; Gray et al., 2004; Miles et al., 2021; Sun & Gresham, 2021). Many of the components that regulate quiescence are found throughout life and likely have conserved functions; however, some of the fundamental ways cells regulate entry, exit, and maintenance of quiescence are not fully understood (Yanagida, 2009).

Saccharomyces cerevisiae is a well-established model organism for studying cellular quiescence because it can reliably produce a population of quiescent cells (Gray et al., 2004; Sun & Gresham, 2021). Quiescence is notoriously difficult to study in multicellular organisms due to the lack of quiescence-specific biomarkers and because it is difficult to determine which cells are quiescent until they begin dividing again. *S. cerevisiae* can produce a large population of quiescent cells that can be isolated by density centrifugation (Allen et al., 2006). Additionally, budding yeast is capable of surviving mutations that would be lethal in higher order organisms, allowing study of critically important factors in the process of establishing quiescence (Miles et al., 2021; Sun & Gresham, 2021).

Because many of the fundamental components that regulate the cell cycle are conserved between budding yeast and higher eukaryotes, we suspect that many players involved in cell cycle arrest are highly conserved.

Tup1-Ssn6 is a transcriptional corepressor complex that globally regulates many cell processes in *S. cerevisiae* (Malavé & Dent, 2006). Tup1 serves mostly as a repressor, although it can also induce gene expression in some contexts (Conlan et al., 1999). Tup1 forms a complex with Ssn6 in which 3-4 Tup1 molecules associate with each Ssn6 molecule (Varanasi et al., 1996). The structure of Tup1 contains an N-terminal helix, which facilitates oligomerization and interacts with histone tails, and a beta-propeller domain composed of seven WD40 domains, which interact with other proteins such as histone deacetylases and DNA binding factors (Sprague et al., 2000). While the detailed mechanism of repression by Tup1 is unknown, some relevant information is available. Tup1 is recruited by sequence-specific transcription factors to DNA (Hanlon et al., 2011), where it recruits histone deacetylases (HDACs) to generate hypoacetylated regions of chromatin, typically associated with transcriptional repression (Davie et al., 2003; Fleming et al., 2014; Watson et al., 2000; Wu et al., 2001). In addition to HDAC recruitment, Tup1 interacts directly with nucleosomes by binding histone H3 and H4 tails, preferentially to hypoacetylated tails, to stabilize +1 nucleosomes that have been repositioned by Isw2 (Chen et al., 2013; Li & Reese, 2001; Rizzo et al., 2011; Zhang & Reese, 2004b). Tup1 also interacts directly with the Mediator complex through its subunit Hrs1, potentially inhibiting it from transcribing into gene bodies (Papamichos-Chronakis et al., 2000). Lastly, Tup1 is thought to influence transcription by blocking the binding of transcription factors to DNA, which may explain why it can appear to simultaneously activate and repress targets (Parnell & Stillman, 2011; Wong & Struhl, 2011).

Homologues of Tup1-Ssn6 found in higher eukaryotes include groucho in drosophila, grg in mice, and TLE1 in humans (Jennings & Ish-Horowicz, 2008). While the beta-propeller domain is well conserved in both sequence and structure, the sequence of the N-terminus is poorly conserved; however, it appears to be structurally conserved in simulations (Jumper et al., 2021; Varadi et al., 2022). Cross-species interactions have been established, where TLE family proteins can pull down yeast Ssn6 from cellular lysate, suggesting strong conservation of structure and function (Flores-Saaib & Courey, 2000; Grbavec et al., 1999). Mice and flies exhibit severe developmental phenotypes in Groucho/TLE null or hypomorphic mutations (Agarwal et al., 2015). Additionally, in humans TLE1 is currently being studied as both a cancer biomarker and a potential cancer therapeutic drug target (Ogawa et al., 2019; Yuan et al., 2017).

During quiescence in yeast, it has been observed that nucleosomes are repositioned and histone tails are deacetylated (McKnight, Boerma, et al., 2015; McKnight, Breeden, et al., 2015). Tup1 plays a role in histone deacetylation and chromatin remodeling in other contexts (Davie et al., 2003; Z. Zhang & Reese, 2004b); however, it is unclear whether Tup1 contributes to these processes in quiescence. Tup1 has been identified as a gene that is essential for viability of cells in G0 and is implicated in glucose repression in yeast, but there is no molecular mechanism to explain its role in these processes (Lin et al., 2017; Papamichos-Chronakis et al., 2004; Reimand et al., 2012; Williams et al., 1991; Williams & Trumbly, 1990). Here, we demonstrate that Tup1 is essential for repressing target genes in stationary phase by interacting with Rpd3 and Isw2 to generate regions of repressed chromatin. We also show that cells lacking Tup1 display morphological defects in stationary phase, suggestive of a possible role for Tup1 in mitosis.

Results

Tup1 Relocalizes After Glucose Starvation

Because Tup1 is a master regulator of many processes in *S. cerevisiae*, we sought to investigate the consequences of glucose starvation on Tup1 localization across the genome. We performed ChIP-Seq analysis on a Myc-tagged Tup1 strain during log phase, diauxic shift, and stationary phase (**Figure 1A**). Ideally, we would work with purified quiescent cells, but due to the flocculent nature of Tup1 deletion strains it is impossible to isolate pure populations of quiescent cells. However, we do anticipate that the findings in stationary phase translate to quiescence because we observed that a fraction of stationary cells in Tup1 knockout strains are viable and can re-enter the cell cycle when plated on glucose rich media. Additionally, we know that Tup1 knockouts have reduced viability in G0, indicating that Tup1 plays a critical role in quiescence.

In log phase, we observed Tup1 binding to the promoters of roughly 923 genes, including genes for ribosomal protein subunits, tRNAs, and cell wall organization (**Figure 1B-C**). During diauxic shift, we observed Tup1 at 174 new targets across the genome, where it persisted into stationary phase. Gene ontology analysis revealed that during diauxic shift, Tup1 binds to the promoters of sugar transmembrane transporters, carbohydrate kinases, ATP from ADP factors, zinc transporters, and DNA binding transcription factors (**Figure 1D**). Motif analysis revealed significant enrichment for binding motifs of 78 transcription factors, including Mig1, which is known to recruit Tup1 to sites of repression (**Supplementary Table 1**) (Bailey & Grant, 2021; Papamichos-Chronakis et al., 2004; Treitel & Carlson, 1995).

Figure 1. Tup1 Relocalizes After Glucose Starvation.
(A) Illustration of a yeast growth curve indicating diauxic shift (DS) and stationary phase (3d). **(B)** Integrated Genome Browser (IGB) tracks representative of Tup1-myc ChIP-Seq during log phase, diauxic shift, and stationary phase. **(C)** ChIP-Seq heatmap of Tup1 binding during log phase, diauxic shift, and stationary phase. Heatmap was clustered using deeptools and sorted by binding intensity. **(D)** Gene ontology analysis from GOrilla of Tup1 binding locations with relative enrichment (e) and p-value (p) indicated for each term.

A)

B)

C)

Repressed by Tup1
Carbohydrate Transport (e = 2.29, p = 6.99E-06)
Ion Transport (e = 1.43, p = 7.58E-07)
rRNA and Ribosome Processing (e = 1.69, p = 5.64E-08)
Translation (e = 2.16, p = 6.51E-21)

Activated by Tup1
Fatty Acid Metabolism (e = 2.73, p = 2.63E-06)
Glyoxylate Cycle(e = 5.42, p = 2.63E-04)
Synaptonemal Complex (e = 3.19, p = 2.68E-05)
Golgi Transport (e = 1.60, p = 9.08E-06)

Figure 2. Tup1 Activates and Represses Key Targets for Quiescence Initiation. (A) Volcano plots of RNA-Seq differential expression data comparing a Tup1 deletion to wild type. Expression changes are based on two biological replicates. **(B)** Venn diagram comparing genes that are repressed by Tup1 in diauxic shift to those repressed in stationary phase. **(C)** Gene ontology analysis from GOrilla of transcripts affected by Tup1 in both diauxic shift and stationary phase with relative enrichment (e) and p-value (p) indicated for each term.

Tup1 Both Activates and Represses Key Targets for Quiescence Initiation

Next, we sought to understand how Tup1 binding correlates with transcriptional changes upon glucose repression. We performed differential expression analysis on RNA-Seq datasets, comparing transcription between wild type and Tup1 knockout strains during diauxic shift and stationary phase (**Figure 2A**). We were able to detect differential expression of 2102 genes in diauxic shift and 2999 genes during stationary phase in the Tup1 deletion, suggestive of Tup1's role in altering transcriptional programming during glucose exhaustion. Of the genes that were significantly differentially expressed in diauxic shift, nearly 60% had a 2-fold increase in expression, while the remaining 40% showed significant reduction in expression (**Figure 2A**). During stationary phase, only slightly more genes showed increased expression than showed decreased expression (approximately 52% versus 48%). Among genes that are repressed by Tup1, 787 were repressed in both diauxic shift and stationary phase (**Figure 2B**).

A previous screen for genes required for quiescence in yeast found 137 deletion mutants that were unable to form a quiescent cell population (L. Li et al., 2015). In our experiments, Tup1 was found to regulate expression of 57% of these genes, indicating that Tup1 plays a key role in transcriptional reprogramming during quiescence. Gene ontology analysis revealed that during diauxic shift, Tup1 represses genes associated with transmembrane transport and metabolism of carbohydrates (**Figure 2C**). We also observed Tup1 repressing genes implicated in transportation of amino acids, metal ions (Zn^{2+}, Mg^{2+}, Cu^{2+}, and Fe^{2+}), vitamins, and fatty acids. While transportation of fatty acids appears repressed, metabolism of fatty acids appears to be activated by Tup1. In addition to repressing transportation of many components, Tup1 apparently stalls translation by repressing expression of ribosomal protein subunits, ribosome biogenesis genes, and tRNA ligases. During stationary phase, Tup1 represses many of the same genes found in diauxic shift, and also represses additional gene sets including rRNAs and DNA transcription factors associated with gene activation.

By comparing the results from RNA-Seq and ChIP-Seq, we found that Tup1 occupancy does not correlate solely with repression or activation of genes. For example, out of the 981 genes Tup1 releases from in DS, 35% have higher levels of expression in the Tup1 deletion and 25% have lower levels, while 40% are not significantly affected. Perhaps Tup1 release frees up binding for additional factors, which may be repressive or may induce transcription. Of the 176 genes where we detect Tup1 binding during DS, we find that most (60%) are repressed by Tup1 and less than 15% are activated, demonstrating that Tup1 recruitment to a site during diauxic shift is usually indicative of repression. Because the number of sites bound by Tup1 in the ChIP dataset does not match the number of repressed genes, it is likely that we are not detecting all Tup1 binding that occurs during diauxic shift; however, we do not believe that the inability to detect all instances of Tup1 binding fully accounts for this difference. We observed that Tup1 binds to and represses many DNA binding transcription factors including Hap1, Nrg1, and Mig1 (**Supplementary Table 1**) and downstream effects of the repression of these transcription factors may account for why there are more repressed targets in the RNA-Seq dataset compared to the ChIP-Seq dataset.

Figure 3. Sds3 and Xbp1 Repress Targets in Common with Tup1. (A) Volcano plots of RNA-Seq differential expression data comparing Xbp1 and Sds3 deletions to wild type in both diauxic shift and stationary phase. Expression changes are based on two biological replicates. **(B)** Venn diagram of transcripts repressed by Tup1, Sds3, and Xbp1 in diauxic shift. **(C)** Venn diagram of transcripts repressed by Tup1, Sds3, and Xbp1 in stationary phase.

Sds3 and Xbp1 Repress Targets in Common With Tup1

Tup1 recruits the Rpd3L complex to sites of repression through DNA-bound transcription factors (Davie et al., 2003; Hanlon et al., 2011; Treitel & Carlson, 1995; Watson et al., 2000). We know that histone deacetylation during quiescence is largely Rpd3L-dependent and that Tup1 interacts with Rpd3L (McKnight, Boerma, et al., 2015). To explore this connection further, we first compared our Tup1 ChIP data to an existing Rpd3 ChIP dataset in quiescent cells (**Supplementary Figure 1**) (McKnight, Boerma, et al., 2015). We found substantial overlap in Rpd3 and Tup1 peaks, indicating that they localize to similar targets, if not interact directly. We also performed ChIP-Seq for the stress response factor Xbp1, which is required to recruit Rpd3 at Xbp1 motifs, and compared these data to Tup1 ChIP and found that Xbp1 and Tup1 also share binding sites (**Supplementary Figure 1**).

To elucidate the relationship between Tup1 and Rpd3 in quiescence, we knocked out Sds3, an Rpd3L-specific subunit required for deacetylation, and compared RNA-Seq datasets to that of the Tup1 knockout (**Figure 3A**). Upon deletion of Sds3 we observed that 1680 genes are differentially expressed in diauxic shift (57% repressed by Sds3 and 43% activated) and 3212 are differentially expressed during stationary phase (52% were repressed by Sds3 and 48% were activated). 734 genes that were upregulated upon Sds3 deletion in diauxic shift were also found to be upregulated during the stationary

phase. A gene ontology analysis revealed that Sds3 represses many genes in common with Tup1, including those implicated in carbohydrate transport and ribosome biogenesis. Sds3 is different from Tup1 in that it represses cell cycle associated genes, which did not appear in Tup1's gene ontology analysis.

When comparing the two RNA-Seq datasets we found that during diauxic shift, 37% of genes repressed by Sds3 were also repressed by Tup1 (p < 5.5e-42) (**Figure 3B**). During stationary phase, Tup1 represses 41% of genes repressed by Sds3 (p < 1.6e-65). This suggests widespread collaboration between Tup1 and Rpd3L in repressing target genes upon glucose starvation.

The transcriptional repressor Xbp1 has been shown to recruit Rpd3L to its binding sites (McKnight, Boerma, et al., 2015). MACS2 (Y. Zhang et al., 2008) peak calling from our ChIP dataset on Xbp1 and Tup1 binding suggests that Tup1 and Xbp1 localize to similar genes (Supplementary Figure 1). We performed ChIP-Seq experiments in strains lacking Xbp1 and found loss of Xbp1 has no effect on Tup1 binding, demonstrating that Xbp1 does not directly recruit Tup1, even though they colocalize (**Supplementary Figure 2**). To measure the effect of Xbp1 deletion on transcriptional silencing, we deleted Xbp1 and performed RNA-Seq experiments during diauxic shift and log phase. We observed that in an Xbp1 deletion 528 genes are differentially expressed during diauxic shift, and during stationary phase 1437 genes were differentially expressed (**Figure 3A**).

Figure 4. Tup1, Sds3, and Xbp1 are Required for H3K23 Deacetylation at Repressed Genes. (A) Heatmap showing the change in H3K23ac levels in deletion mutants compared to wild-type across all yeast genes. **(B)** H3K23ac profile levels across gene promoters, integrated across all yeast genes. **(C)** Venn diagram comparing the 1000 genes with the largest change in H3K23ac to genes repressed by Tup1 and Sds3 in stationary phase.

Of these genes, 86% were repressed by Xbp1, and 14% were activated. During stationary phase, 51% of differentially expressed genes were repressed by Xbp1 and 49% activated. A gene ontology analysis revealed that Xbp1 is responsible for repressing genes associated with carbohydrate metabolism during diauxic shift, while repressing carbohydrate transport and cell cycle genes during stationary phase. Xbp1 and Tup1 repress 150 (p < 3.8e-28) shared genes during diauxic shift and 379 (p < 2.8e-60) genes during stationary phase (**Figure 3C**). Xbp1 represses 153 (p < 4.8e-45) genes in common with Sds3 during diauxic shift and 455 (p < 1.3e-101) during stationary phase. Xbp1 also regulates cell cycle genes along with Sds3, while Tup1 has no effect on these genes.

Tup1, Sds3, and Xbp1 Are Required for H3K23 Deacetylation at Repressed Genes

Since Tup1 facilitates H3K23 deacetylation in other contexts (Malavé & Dent, 2006), we next wanted to see if Tup1 contributes to the H3K23 deacetylation we observe during quiescence. To determine if Tup1 is required for deacetylation, we performed H3K23ac ChIP-Seq during stationary phase where Tup1, Xbp1, or Sds3 had been deleted. Cells with the Sds3 deletion showed global hyperacetylation in the promoter region during the stationary phase (**Figure 4A-B**). Additionally, Δtup1 cells also showed global hyperacetylation, though not as dramatically as with the Δsds3 strain. It was difficult to see a global effect of Xbp1 deletion on acetylation levels; however, when we filtered genes with an Xbp1 motif, it appears that deletion of Xbp1 influences acetylation levels as well, although not nearly as dramatically as with loss of Sds3 or Tup1.

To determine the effect of acetylation of transcription, we took the 1000 genes with the greatest changes in acetylation levels across all three mutants and compared them to genes repressed by Tup1 or Sds3 in stationary phase (**Figure 4C**). We observed a strong relationship between genes that are deacetylated in stationary phase and genes that are repressed.

Tup1 and Isw2 Regulate Nucleosome Positions at HXT Family Genes

During quiescence, cells reposition their +1 nucleosomes, either to shrink the nucleosome depleted region (NDR) or to expand it. Shrinking of the NDR is generally regarded as repressive, because it presumably blocks recruitment of DNA transcription factors and RNA polymerase II. Expansion of the nucleosome NDR, on the other hand, presumably facilitates recruitment of RNA polymerase and is associated with active transcription. The vast majority of NDRs shrink during quiescence in yeast, while a few are expanded (McKnight, Breeden, et al., 2015). The chromatin remodeler Isw2 is known to interact with Tup1, and both proteins are required to position nucleosomes in some contexts (Rizzo et al., 2011; Z. Zhang & Reese, 2004b, 2004a). Because Tup1 is involved in +1 nucleosome stabilization in other contexts, we performed MNase-Seq to determine if Tup1 affects the position of nucleosomes in stationary phase. We found that specifically, Tup1 and Isw2 are required to position nucleosomes at the HXT family of glucose transporter genes HXT1, HXT3, HXT8, HXT9, HXT11, HXT13, HXT15, and HXT16 (**Figure 5A**).

Figure 5. Tup1 and Isw2 Regulate Nucleosome Positions at HXT Family Genes. (A) Integrated Genome Browser (IGB) tracks of MNase-Seq in wild type, Δisw2, and Δtup1 in stationary phase. Dashed gray lines indicate wild type nucleosome positions. **(B)** Volcano plot of Δisw2 deletion in stationary phase demonstrating shifted nucleosomes at the HXT1 locus. **(C)** MNase-Seq dyad signal across 217 genes with shifted nucleosomes in Δisw2 during stationary phase. "-1" indicates the first nucleosome upstream of the TSS, while "+1" indicates the first nucleosome downstream of the TSS. **(D)** Gene ontology terms for genes with shifted nucleosome positions in Δisw2 yeast at stationary phase, with relative enrichment (e) and p-value (p) indicated for each term.

RNA-Seq data revealed that Isw2 is required to repress 106 genes during diauxic shift and 172 genes in stationary phase (**Figure 5B**). Isw2 represses many shared targets with Tup1: 33% of genes repressed by Isw2 are also repressed by Tup1 in diauxic shift, whereas almost half of the genes repressed by Isw2 in stationary phase are repressed by Tup1. In addition, we found a greater number of mispositioned nucleosomes in the Δisw2 strain, which had mispositioned nucleosomes in the -2, -1, +1, and +2 positions relative to the transcription start site (TSS) (**Figure 5C**). Our gene ontology analysis of the genes with mis-positioned nucleosomes in the Δisw2 deletion showed that Isw2 regulates nucleosomes in the promoters of genes involved in glucose transport, cell cycle, and DNA repair pathways in stationary phase (**Figure 5D**).

Deletion of Tup1 or Sds3 Leads to Morphological Defects in Stationary Phase

Due to the multimeric nature of the Tup1-Ssn6 complex, it has been hypothesized that Tup1 may act as a scaffold for long-distance chromatin interactions. We know that during quiescence there are large scale chromatin folding events in yeast (Swygert et al., 2019). We have biochemical evidence that Tup1 may regulate chromatin folding in vitro; however, it is unknown whether this occurs in vivo. We initially wondered if deleting Tup1 would lead to less compaction of chromatin and sought to determine if Δtup1 cells had larger nuclei using fluorescence microscopy and DAPI staining. Although we failed to detect statistically significant changes in nuclei size, we found that Δtup1 cells showed multiple DAPI puncta not observed in wild type cells (**Figure 6A-B**).

Deletion of Sds3 did not cause this phenotype; however, Δsds3 cells were often misshapen, which we termed "peanut shaped cells" (**Figure 6C**). Peanut shaped cells have been observed in other mutants, particularly mutants involved in schmoo formation such as AFR1, PEA2, and SPA2 (Chenevert et al., 1994). While Sds3 does not appear to significantly regulate the expression of these genes, Sds3 is involved in regulating the cell cycle, bud site selection, and cell wall organization. While the phenotype between Δsds3 cells and Afr1 mutant appear similar, it is likely caused by different types of misregulation. The morphological defects observed in both Δtup1 and Δsds3 yeast suggest a potential mitotic defect, leading to aberrant distribution of DNA within nuclei in Δtup1 cells and abnormal cell division in the Δsds3 mutant.

Discussion

Tup1 plays a key role in regulating large scale transcriptional responses to environmental conditions in yeast (Hanlon et al., 2011). Our ChIP-Seq studies with epitope-tagged Tup1 revealed that when glucose is depleted during the diauxic shift, Tup1 releases from numerous chromatin sites where it is bound in log phase and binds at new positions that persist into the stationary phase. Tup1 binds some other sites in both log phase and during diauxic shift, typically associated with genes involved in cell wall organization. During log phase, Tup1 is bound to the promoters of ribosomal subunit proteins and tRNAs, which are actively transcribed at this stage but become repressed during diauxic shift; it seems that Tup1 is involved in both gene activation and repression. During diauxic shift, Tup1 relocalizes to genes associated with carbohydrate metabolism and transport, as well as factors that convert ADP to ATP, zinc

A)

WT   Δtup1   Δsds3

B)

C)

Figure 6. Deletion of Tup1 or Sds3 Leads to Morphology Defects in Stationary Phase. **(A)** Fluorescence microscopy of DAPI stained nuclei across the genome at 67x magnification in wild type, Δtup1, and Δsds3 yeast. **(B)** Quantification of Δtup1 cells with multiple DAPI puncta. **(C)** Quantification of peanut-shaped cells in Δsds3 yeast.

transporters, and DNA-binding transcription factors; Tup1 relocalization to these genes represses their expression. The sites occupied by Tup1 regardless of glucose availability all become derepressed in a Tup1 knockout strain, indicating that Tup1 represses these targets. We suspect that repression of transcription factors by Tup1 may play a role in its ability to affect transcription of more factors in addition to those directly associated with its binding sites. Additionally, because Tup1 does not directly interact with DNA, we suspect that ChIP fails to reveal every instance of Tup1 binding.

Tup1 null mutants fail to repress many of the genes required for diauxic shift and establishment of quiescence. Many of the targets that are repressed/activated by Tup1 account for characteristics of quiescent cells. For example, common hallmarks of quiescent cells are fewer ribosomes, stalled translation, cell wall reorganization, and repression of metabolism. Our RNA-Seq data indicate that Tup1 is required to repress genes associated with all of these processes, demonstrating its role in regulating the transition to quiescence. Li et. al performed a genetic screen for factors that are required for quiescence in yeast. We found that Tup1 regulates more than 56% of these factors, further demonstrating the importance of Tup1 in regulating quiescence.

Tup1 regulates metabolism by simultaneously repressing genes involved in RNA, lipid, amino acid, and carbohydrate biosynthetic processes and activating the glyoxylate cycle and fatty acid metabolism. Transmembrane transporters of amino acids, carbohydrates, fatty acids, and metals (specifically magnesium, zinc, copper, and iron) are repressed by Tup1. We also find that two of the three yeast aquaporins are repressed by Tup1. It has been observed in other organisms that transmembrane transport is related to cellular proliferation (Peters-Regehr et al., 1999; Shyh-Chang & Daley, 2015). These findings indicate that yeast cells repress the vast majority of transmembrane transporters, which likely allows the cells to maintain a controlled intracellular environment in a context where metabolism is dramatically slowed.

The importance of the Rpd3 complex in establishing a quiescent state has been well demonstrated, and one factor that recruits Rpd3 is the transcription factor Xbp1 (McKnight, Boerma, et al., 2015). Xbp1 and Rpd3 share many common targets of repression with Tup1, while simultaneously having targets independent of Tup1. For example, Tup1 and Sds3 both strongly repress tRNA processing factors, ribosomal subunits, transmembrane transport, sugar catabolism, and cell wall components. Xbp1 and Tup1 overlap in their repression of transmembrane transport genes and sugar metabolism. However, Xbp1 and Sds3 strongly repress cell cycle genes, which are unaffected by Tup1.

Histone deacetylation is strongly associated with transcriptional repression in quiescence (McKnight, Breeden, et al., 2015) and we demonstrated the importance of Tup1, Xbp1, and Sds3 to facilitate histone deacetylation. Whether or not histone deacetylation is the main driving force of transcriptional repression is up for debate. We are only able to detect changes in histone deacetylation three days into stationary phase, which raises the possibility that histone deacetylation is not the main driving force in repressing transcription. However, recent findings demonstrate that the acetylome can change within 5 minutes of glucose depletion, so perhaps changes in the acetylome during diauxic shift are too subtle for us to detect (Hsieh et al., 2022; Cai, et al., 2011). We know that Tup1 and Rpd3 have repressive activities outside of histone deacetylation Malavé, & Dent, 2006), but it is difficult to decipher which mechanism is employed to repress targets. Overall, there is a strong correlation between genes that are repressed and those that lose their acetyl histone marks.

Tup1 and Isw2 both coordinately regulate nucleosome positions in the promoter regions of HXT family genes. Our RNA-Seq dataset revealed that these genes are also repressed by Tup1 and Isw2, suggesting that repositioning of nucleosomes by Isw2 and Tup1 at these genes leads to repression. Interestingly, these are the only genes in which both Tup1 and Isw2 affect nucleosome positions; deletion of Tup1 has little effect on the positions of nucleosomes at any other genes. We see a greater number of nucleosomes that are affected by Isw2 in genes involved in DNA repair pathways, glucose transport, and cell cycle process. We found that in a Sds3 knockout, cells form peanut shapes, indicative of a cell attempting to divide but ultimately failing to complete the process. Sds3 regulates many cell cycle genes, and it seems likely that failure to repress key cell cycle genes leads to the incomplete cell divisions we observed.

The multiple DAPI puncta observed in Tup1 knockout cells warrant further investigation. One possibility is that these cells have multiple nuclei or fragments of nuclei, which could be the result of a failure to properly segregate chromosomes during mitosis. Another possibility is these puncta are fragments of a broken-down nucleus, which is the

result of cell cycle dysregulation. In any case, the observation highlights the importance of Tup1 in regulating the exit from the cell cycle. Taken together these results indicate that Tup1 is critical for gene repression in the exit from the cell cycle and establishment of quiescence.

It has been demonstrated that the Tup1 homologue GRG5 is critical for embryonic stem cell fate decisions in mice (Chanoumidou et al., 2018). Additionally, TLE1 is associated with many cancers in humans, including synovial carcinoma, lung cancer, breast cancer, glioblastoma, gastric cancer, and pancreatic cancer (Yuan et al., 2017). This indicates a role of Tup1/Groucho/TLE family proteins in determining cell fate in higher eukaryotes. Perhaps a conserved role of Tup1 family proteins is to regulate entry and exit from the cell cycle, which may explain why dysregulation of Groucho of TLE in higher eukaryotes leads to cancer.

Materials and Methods

Generation of Deletion Strains

All yeast strains were generated from the parent strain *S. cerevisiae* W303 RAD5+ (ATCC 208352). Null mutants were generated by replacing the gene of interest with antibiotic resistance markers amplified from pAG vectors (Longtine et al., 1998). FLAG- and Myc-tagged strains were made from a pFA6a vector containing their respective sequences. We then inserted the tags through homologous recombination of PCR products using selectable drug markers (Longtine et al., 1998).

Growth Conditions

S. cerevisiae W303 cultures were streaked from glycerol stocks onto yeast extract peptone dextrose (YPD) plates and grown at 30° C. Individual colonies were picked and used to inoculate 25 mL YPD with shaking at 30° C. These cultures were grown to an O.D. of 0.6 to 0.8 for log phase cultures. Diauxic shift cultures were obtained by monitoring glucose levels with glucose test strips (Precision Laboratories). Cells were harvested two hours after there was no detectable glucose. Stationary phase cultures were prepared by growing yeast for three days in YPD medium.

Chromatin Immunoprecipitation

Chromatin immunoprecipitation was performed as previously described (Rodriguez et al.; 2014). 100 OD-mL of cells were crosslinked in 1% formaldehyde for 20 minutes at 30° C. Cells were then quenched with a final concentration of 125 mM glycine, pelleted, and frozen at -80° C. Cells were lysed in a Bead Beater (Biospec) for two minutes with acid-washed glass beads, diameter 425-600 m (Sigma Aldrich) in the presence of protease inhibitors (Proteoloc Protease Inhibitor Cocktail, Expedeon) and then sheared in a Bioruptor sonicator for a total of 30 min (high output 30 sec on 30 sec off for 10 minutes 3x) and then centrifuged. The sheared chromatin supernatant was incubated with antibodies bound to G protein beads (Invitrogen). The crosslinks were reversed, overnight at 65° C and the sample was treated with 20 g RNase A for two hours at 45° C and 100 g Proteinase K for 4 hours at 55° C, and then purified using Qiagen MinElute columns. DNA samples were quantified using a Qubit fluorescence assay and libraries were prepared using a NuGEN Ovation Ultralow kit. Sequencing was performed by the University of Oregon Genomics and Cell Characterization Core Facility on an Illumina NextSeq500 on the 37 cycle, paired-end setting, yielding approximately 10-20 million paired-end reads per sample.

RNA Extraction

Cells were removed from -80° C and ground with a mortar and pestle in liquid nitrogen. RNA was extracted using hot acid-phenol (Uppuluri et al., 2007) and cleaned up using the RNeasy kit (Qiagen). Libraries were prepared using the NuGEN Universal Plus mRNA kit. Sequencing was performed by University of Oregon Genomics and Cell Characterization Core Facility on an Illumina NextSeq500 with 37 cycles of paired-end setting. Paired end reads were filtered and aligned to the *S. cerevisiae* sacSer3 genome using bowtie2 (Langmead & Salzberg, 2012). Differential expression analysis was performed using DESeq2 (Love et al., 2014).

Micrococcal Nuclease Digestion

We utilized our rapid MNase protocol (McKnight et al., 2021). In short, cross-linked cells were permeabilized by zymolyase digestion then treated with MNase and Exonuclease III. Next, the samples were treated with RNaseA and Proteinase K and purified using the Qiagen MinElute PCR Cleanup kit.

DAPI Staining and Fluorescence Microscopy

Staining and microscopy was performed as described by Swygert et al. (2019). Cells were fixed in 3.7% formaldehyde for 20 min at 4° C in 0.1 M KPO4 pH 6.4 buffer. Cells were washed twice in sorbitol citrate (1.2 M sorbitol, 100 mM K2HPO4, 36.4 mM citric acid) and incubated with 2.5 µg of Zymolyase for 30 minutes. Cells were added to polytetrafluoroethylene (PFTE) slides with 0.1% polylysine then washed with ice cold methanol and acetone. Next 5 µL of DAPI mount (0.1 µg/ml DAPI, 9.25 mM p-phenylenediamine, dissolved in PBS, and 90% glycerol) was added to the well. Images were taken on a Zeiss LSM 880 confocal microscope with Airyscan. Images were analyzed and processed using Imaris software (Oxford Instruments).

Computational Analysis

All alignments were performed using Bowtie2 (Langmead & Salzberg, 2012). Further analysis was performed using deepTools including generation of heatmaps (Ramírez et al., 2016). Gene tracks were visualized using Integrated Genome Browser (Freese et al., 2016). Venn diagrams were generated using euler.com (Larson, 2020). Motif analysis was performed using SEA in MEME-suite (Bailey & Grant, 2021).

Author Contributions

Conceptualization, T.BB., E.U.S., J.N.M.; Methodology, T.B.B, L.E.M., J.N.M.; Investigation, T.B.B., P.A.A., L.E.M., J.N.M.; Additional Resources, E.U.S.; Writing - Original Draft, T.B.B.; Writing - Reviewing and Editing, T.B.B., E.U.S.; L.E.M.; Visualization, T.B.B., L.E.M.; Supervision, E.U.S., J.N.M., L.E.M.; Project Administration, L.E.M., J.N.M.; Funding Acquisition, J.N.M.

Declaration of Interest

The authors have no competing interests to declare.

Data Availability

The datasets generated during this study are available at GEO under accession code GSE210934.

Acknowledgments

The authors thank the Genomics Core (GC3F) at the University of Oregon for high throughput sequencing services. The authors also thank Christine Cucinotta (Fred Hutchinson Cancer Research Center) and Hideki Tanizawa, Osamu Iwasaki, Ken-Ichi Noma for helpful discussions related to this work. This work was supported by a National Institutes of Health training grant T32 GM007759 (to T.B.B), and by NIGMS R01 GM129242 (to J.N.M. and L.E.M).

Works Cited

Agarwal, M., Kumar, P., & Mathew, S. J. (2015). The Groucho/Transducin-like enhancer of split protein family in animal development. *IUBMB Life*, 67(7), 472–481. https://doi.org/10.1002/iub.1395

Allen, C., Büttner, S., Aragon, A. D., Thomas, J. A., Meirelles, O., Jaetao, J. E., Benn, D., Ruby, S. W., Veenhuis, M., Madeo, F., & Werner-Washburne, M. (2006). Isolation of quiescent and nonquiescent cells from yeast stationary-phase cultures. *Journal of Cell Biology*, 174(1), 89–100. https://doi.org/10.1083/jcb.200604072

Bailey, T. L., & Grant, C. E. (2021). SEA: Simple Enrichment Analysis of motifs. *BioRxiv*, 2021.08.23.457422. https://www.biorxiv.org/content/10.1101/2021.08.23.457422v1

Cai, L.; Sutter, B. M.; Li, B.; Tu, B. P. (2011). Acetyl-CoA Induces Cell Growth and Proliferation by Promoting the Acetylation of Histones at Growth Genes. *Molecular Cell*, 42(4), 426–437. https://doi.org/10.1016/j.molcel.2011.05.004.Acetyl-CoA

Chanoumidou, K., Hadjimichael, C., Athanasouli, P., Ahlenius, H., Klonizakis, A., Nikolaou, C., Drakos, E., Kostouros, A., Stratidaki, I., Grigoriou, M., & Kretsovali, A. (2018). Groucho related gene 5 (GRG5) is involved in embryonic and neural stem cell state decisions. *Scientific Reports*, 8(1), 1–16. https://doi.org/10.1038/s41598-018-31696-9

Chen, K., Wilson, M. A., Hirsch, C., Watson, A., Liang, S., Lu, Y., Li, W., & Dent, S. Y. R. (2013). Stabilization of the promoter nucleosomes in nucleosome-free regions by the yeast Cyc8-Tup1 corepressor. *Genome Research*, 23(2), 312–322. https://doi.org/10.1101/gr.141952.112

Chenevert, J., Valtz, N., & Herskowitz, I. (1994). Identification of genes required for normal pheromone-induced cell polarization in Saccharomyces cerevisiae. *Genetics*, 136(4), 1287–1296. https://doi.org/10.1093/genetics/136.4.1287

Conlan, R. S., Gounalaki, N., Hatzis, P., & Tzamarias, D. (1999). The Tup1-Cyc8 Protein Complex Can Shift from a Transcriptional Co-repressor to a Transcriptional Co-activator. *Journal of Biological Chemistry*, 274(1), 205–210. https://doi.org/10.1074/jbc.274.1.205

Davie, J. K., Edmondson, D. G., Coco, C. B., & Dent, S. Y. R. (2003). Tup1-Ssn6 Interacts with Multiple Class I Histone Deacetylases in Vivo. *Journal of Biological Chemistry*, 278(50), 50158–50162. https://doi.org/10.1074/jbc.M309753200

de Virgilio, C. (2012). The essence of yeast quiescence. *FEMS Microbiology Reviews*, 36(2), 306–339. https://doi.org/10.1111/j.1574-6976.2011.00287.x

Fleming, A. B., Beggs, S., Church, M., Tsukihashi, Y., & Pennings, S. (2014). The yeast Cyc8-Tup1 complex cooperates with Hda1p and Rpd3p histone deacetylases to robustly repress transcription of the subtelomeric FLO1 gene. *Biochimica et Biophysica Acta - Gene Regulatory Mechanisms*, 1839(11), 1242–1255. https://doi.org/10.1016/j.bbagrm.2014.07.022

Flores-Saaib, R. D., & Courey, A. J. (2000). Analysis of Groucho-histone interactions suggests mechanistic similarities between Groucho- and Tup1-mediated repression. Nucleic Acids Research, 28(21), 4189–4196. https://doi.org/10.1093/nar/28.21.4189

Freese, N. H., Norris, D. C., & Loraine, A. E. (2016). Integrated genome browser: Visual analytics platform for genomics. *Bioinformatics*, 32(14), 2089–2095. https://doi.org/10.1093/bioinformatics/btw069

Gray, J. V, Petsko, G. A., Johnston, G. C., Ringe, D., Singer, R. A., & Werner-washburne, M. (2004). "Sleeping Beauty": Quiescence in Saccharomyces cerevisiae. *Society*, 68(2), 187–206. https://doi.org/10.1128/MMBR.68.2.187

Grbavec, D., Lo, R., Liu, Y., Greenfield, A., & Stifani, S. (1999). Groucho/transducin-like Enhancer of split (TLE) family members interact with the yeast transcriptional co-repressor SSN6 and mammalian SSN6-related proteins: Implications for evolutionary conservation of transcription repression mechanisms. *Biochemical Journal*, 337(1), 13–17. https://doi.org/10.1042/0264-6021:3370013

Hanlon, S. E., Rizzo, J. M., Tatomer, D. C., Lieb, J. D., & Buck, M. J. (2011). The stress response factors Yap6, Cin5, Phd1, and Skn7 direct targeting of the conserved co-repressor Tup1-Ssn6 in S. cerevisiae. *PLoS ONE*, 6(4). https://doi.org/10.1371/journal.pone.0019060

Hsieh, W. C., Sutter, B. M., Ruess, H., Barnes, S. D., Malladi, V. S., & Tu, B. P. (2022). Glucose starvation induces a switch in the histone acetylome for activation of gluconeogenic and fat metabolism genes. *Molecular Cell*, 82(1), 60-74.e5. https://doi.org/10.1016/j.molcel.2021.12.015

Jennings, B. H., & Ish-Horowicz, D. (2008). The Groucho/TLE/Grg family of transcriptional co-repressors. *Genome Biology*, 9(1), 205. https://doi.org/10.1186/gb-2008-9-1-205

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4), 357–359. https://doi.org/10.1038/nmeth.1923

Larsson J (2020). eulerr: Area-Proportional Euler and Venn Diagrams with Ellipses. R package version 6.1.0, https://cran.r-project.org/package=eulerr.

Li, B., & Reese, J. C. (2001). Ssn6-Tup1 Regulates RNR3 by Positioning Nucleosomes and Affecting the Chromatin Structure at the Upstream Repression Sequence. *Journal of Biological Chemistry*, 276(36), 33788–33797. https://doi.org/10.1074/jbc.M104220200

Li, L., Miles, S., & Breeden, L. L. (2015). A genetic screen for Saccharomyces cerevisiae mutants that fail to enter quiescence. *G3: Genes, Genomes, Genetics*, 5(8), 1783–1795. https://doi.org/10.1534/g3.115.019091

Lin, X., Yu, A. Q., Zhang, C. Y., Pi, L., Bai, X. W., & Xiao, D. G. (2017). Functional analysis of the global repressor Tup1 for maltose metabolism in Saccharomyces cerevisiae: Different roles of the functional domains. *Microbial Cell Factories*, 16(1), 1–12. https://doi.org/10.1186/s12934-017-0806-6

Longtine, M.S.; McKenzie, A.; Demarini, D.J.; Shah, N.G.; Wach, A.; Brachat, A.; Philippsen, P.; Pringle, J.R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast*, 14(10), 953-61. https://doi.org/10.1002/(SICI)1097-0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12), 1–21. https://doi.org/10.1186/s13059-014-0550-8

Malavé, T. M., & Dent, S. Y. R. (2006). Transcriptional repression by Tup1-Ssn6. *Biochemistry and Cell Biology*, 84(4), 437–443. https://doi.org/10.1139/O06-073

Marescal, O., & Cheeseman, I. M. (2020). Cellular Mechanisms and Regulation of Quiescence. *Developmental Cell*, 55(3), 259–271. https://doi.org/10.1016/j.devcel.2020.09.029

McKnight, J. N., Boerma, J. W., Breeden, L. L., & Tsukiyama, T. (2015). Global Promoter Targeting of a Conserved Lysine Deacetylase for Transcriptional Shutoff during Quiescence Entry. *Molecular Cell*, 59(5), 732–743. https://doi.org/10.1016/j.molcel.2015.07.014

McKnight, J. N., Breeden, L. L., & Tsukiyama, T. (2015). A molecular off switch for transcriptional quiescence. *Cell Cycle*, 14(23), 3667–3668. https://doi.org/10.1080/15384101.2015.1112618

McKnight, L. E., Crandall, J. G., Bailey, T. B., Banks, O. G. B., Orlandi, K. N., Truong, V. N., Donovan, D. A., Waddell, G. L., Wiles, E. T., Hansen, S. D., Selker, E. U., & McKnight, J. N. (2021). Rapid and inexpensive preparation of genome-wide nucleosome footprints from model and non-model organisms. *STAR Protocols*, 2(2), 100486. https://doi.org/10.1016/j.xpro.2021.100486

Miles, S., Bradley, G. T., & Breeden, L. L. (2021). The budding yeast transition to quiescence. *Yeast*, 38(1), 30–38. https://doi.org/10.1002/yea.3546

O'Farrell, P. H. (2011). Quiescence: Early evolutionary origins and universality do not imply uniformity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366(1584), 3498–3507. https://doi.org/10.1098/rstb.2011.0079

Ogawa, M., Yaginuma, T., Nakatomi, C., Nakajima, T., Tada-Shigeyama, Y., Addison, W. N., Urata, M., Matsubara, T., Watanabe, K., Matsuo, K., Sato, T., Honda, H., Hikiji, H., Watanabe, S., & Kokabu, S. (2019). Transducin-like enhancer of split 3 regulates proliferation of melanoma cells via histone deacetylase activity. *Oncotarget*, 10(3), 404–414. https://doi.org/10.18632/oncotarget.26552

Papamichos-Chronakis, M., Gligoris, T., & Tzamarias, D. (2004). The Snf1 kinase controls glucose repression in yeast by modulating interactions between the Mig1 repressor and the Cyc8-Tup1 co-repressor. *EMBO Reports*, 5(4), 368–372. https://doi.org/10.1038/sj.embor.7400120

Papamichos-Chronakis, M., Steven, R., Gounalaki, N., Copf, T., Tzamarias, D., Chem, J. B., & Conlan, R. S. (2000). Hrs1 / Med3 Is a Cyc8-Tup1 Corepressor Target in the RNA Polymerase II Holoenzyme. *Journal of Biological Chemistry*, 275(12), 8397–8403. https://doi.org/10.1074/jbc.275.12.8397

Parnell, E. J., & Stillman, D. J. (2011). Shields up: The Tup1-Cyc8 repressor complex blocks coactivator recruitment. *Genes and Development*, 25(23), 2429–2435. https://doi.org/10.1101/gad.181768.111

Peters-Regehr, T., Bode, J. G., Kubitz, R., & Häussinger, D. (1999). Organic osmolyte transport in quiescent and activated rat hepatic stellate cells (Ito cells). *Hepatology*, 29(1), 173–180. https://doi.org/10.1002/hep.510290111

Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S., Dündar, F., & Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Research*, 44(W1), W160–W165. https://doi.org/10.1093/NAR/GKW257

Reimand, J., Aun, A., Vilo, J., Vaquerizas, J. M., Sedman, J., & Luscombe, N. M. (2012). M:Explorer: Multinomial regression models reveal positive and negative regulators of longevity in yeast quiescence. *Genome Biology*, 13(6). https://doi.org/10.1186/gb-2012-13-6-r55

Rittershaus, E. S. C., Baek, S. H., & Sassetti, C. M. (2013). The normalcy of dormancy: Common themes in microbial quiescence. *Cell Host and Microbe*, 13(6), 643–651. https://doi.org/10.1016/j.chom.2013.05.012

Rizzo, J. M., Mieczkowski, P. A., & Buck, M. J. (2011). Tup1 stabilizes promoter nucleosome positioning and occupancy at transcriptionally plastic genes. *Nucleic Acids Research*, 39(20), 8803–8819. https://doi.org/10.1093/nar/gkr557

Rodriguez, J.; McKnight, J.N.; Tsukiyama, T. Genome-wide analysis of nucleosome positions, occupancy, and accessibility in yeast: nucleosome mapping, high-resolution ChIP, and NCAM. *Current Protocols Molecular Biology*, 1(108), 1-16. https://doi.org/10.1002/0471142727.mb2128s108

Sprague, E. R., Redd, M. J., Johnson, A. D., & Wolberger, C. (2000). Structure of the C-terminal domain of Tup1, a corepressor of transcription in yeast. *EMBO Journal*, 19(12), 3016–3027. https://doi.org/10.1093/emboj/19.12.3016

Sun, S., & Gresham, D. (2021). Cellular quiescence in budding yeast. *Yeast*, 38(1), 12–29. https://doi.org/10.1002/yea.3545

Swygert, S. G., Kim, S., Wu, X., Fu, T., Hsieh, T. H., Rando, O. J., Eisenman, R. N., Shendure, J., McKnight, J. N., & Tsukiyama, T. (2019). Condensin-Dependent Chromatin Compaction Represses Transcription Globally during Quiescence. *Molecular Cell*, 73(3), 533-546.e4. https://doi.org/10.1016/j.molcel.2018.11.020

Treitel, M. A., & Carlson, M. (1995). Repression by SSN6-TUP1 is directed by MIG1, a repressor/activator protein. *Proceedings of the National Academy of Sciences*, 92(8), 3132–3136. https://doi.org/10.1073/pnas.92.8.3132

Tümpel, S., & Rudolph, K. L. (2019). Quiescence: Good and Bad of Stem Cell Aging. *Trends in Cell Biology*, 29(8), 672–685. https://doi.org/10.1016/j.tcb.2019.05.002

Uppuluri, P.; Perumal, P.; Chaffin, W.L. (2007). Analysis of RNA species of various sizes from stationary-phase planktonic yeast cells of Candida albicans. *FEMS Yeast Research*, (7), 110-117. https://doi/org/10.1111j.1567-1364.2006.00143.x

Valcourt, J. R., Lemons, J. M. S., Haley, E. M., Kojima, M., Demuren, O. O., & Coller, H. A. (2012). Staying alive: Metabolic adaptations to quiescence. *Cell Cycle*, 11(9), 1680–1696. https://doi.org/10.4161/cc.19879

Vallette, F. M., Olivier, C., Lézot, F., Oliver, L., Cochonneau, D., Lalier, L., Cartron, P. F., & Heymann, D. (2019). Dormant, quiescent, tolerant and persister cells: Four synonyms for the same target in cancer. *Biochemical Pharmacology*, 162(November 2018), 169–176. https://doi.org/10.1016/j.bcp.2018.11.004

van Velthoven, C. T. J., & Rando, T. A. (2019). Stem Cell Quiescence: Dynamism, Restraint, and Cellular Idling. *Cell Stem Cell*, 24(2), 213–225. https://doi.org/10.1016/j.stem.2019.01.001

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D., Stroe, O., Wood, G., Laydon, A., Zídek, A., Green, T., Tunyasuvunakool, K., Petersen, S., Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., … Velankar, S. (2022). AlphaFold Protein Structure Database: Massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Research*, 50(D1), D439–D444. https://doi.org/10.1093/nar/gkab1061

Varanasi, U. S., Klis, M., Mikesell, P. B., & Trumbly, R. J. (1996). The Cyc (Ssn6)-Tup1 corepressor complex is composed of one Cyc8 and four Tup1 subunits. *Molecular and Cellular Biology*, 16(12), 6707–6714. https://doi.org/10.1128/MCB.16.12.6707

Watson, A. D., Edmondson, D. G., Bone, J. R., Mukai, Y., Yu, Y., Du, W., Stillman, D. J., & Roth, S. Y. (2000). Ssn6-Tup1 interacts with class I histone deacetylases required for repression. *Genes and Development*, 14(21), 2737–2744. https://doi.org/10.1101/gad.829100

Williams, F. E., & Trumbly, R. J. (1990). Characterization of TUP1, a mediator of glucose repression in Saccharomyces cerevisiae. *Molecular and Cellular Biology*, 10(12), 6500–6511. https://doi.org/10.1128/mcb.10.12.6500

Williams, F. E., Varanasi, U., & Trumbly, R. J. (1991). The CYC8 and TUP1 proteins involved in glucose repression in Saccharomyces cerevisiae are associated in a protein complex. *Molecular and Cellular Biology*, 11(6), 3307–3316. https://doi.org/10.1128/mcb.11.6.3307

Wong, K. H., & Struhl, K. (2011). The Cyc8-Tup1 complex inhibits transcription primarily by masking the activation domain of the recruiting protein. *Genes and Development*, 25(23), 2525–2539. https://doi.org/10.1101/gad.179275.111

Wu, J., Suka, N., Carlson, M., & Grunstein, M. (2001). TUP1 utilizes histone H3/H2B-specific HDA1 deacetylase to repress gene activity in yeast. *Molecular Cell*, 7(1), 117–126. https://doi.org/10.1016/S1097-2765(01)00160-5

Yanagida, M. (2009). Cellular quiescence: are controlling genes conserved? *Trends in Cell Biology*, 19(12), 705–715. https://doi.org/10.1016/j.tcb.2009.09.006

Yao, G. (2014). Modelling mammalian cellular quiescence. *Interface Focus*, 4(3). https://doi.org/10.1098/rsfs.2013.0074

Yuan, D., Yang, X., Yuan, Z., Zhao, Y., & Guo, J. (2017). TLE1 function and therapeutic potential in cancer. *Oncotarget*, 8(9), 15971–15976. https://doi.org/10.18632/oncotarget.13278

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C., Myers, R. M., Brown, M., Li, W., & Shirley, X. S. (2008). Model-based analysis of ChIP-Seq (MACS). *Genome Biology*, 9(9). https://doi.org/10.1186/gb-2008-9-9-r137

Zhang, Z., & Reese, J. C. (2004a). Redundant mechanisms are used by Ssn6-Tup1 in repressing chromosomal gene transcription in Saccharomyces cerevisiae. *Journal of Biological Chemistry*, 279(38), 39240–39250. https://doi.org/10.1074/jbc.M407159200

Zhang, Z., & Reese, J. C. (2004b). Ssn6-Tup1 requires the ISW2 complex to position nucleosomes in Saccharomyces cerevisiae. *EMBO Journal*, 23(11), 2246–2257. https://doi.org/10.1038/sj.emboj.7600227