***Neisseria gonorrhoeae* diagnostic escape from a *gyrA*-based test for ciprofloxacin sus-**

ceptibility can lead to increased zoliflodacin resistance

Daniel H. F. Rubin[1], Tatum D. Mortimer[1], Yonatan H. Grad[1]

Affiliations

[1] Harvard T.H. Chan School of Public Health, Boston, MA, USA

Corresponding author

Yonatan H. Grad

Department of Immunology and Infectious Diseases

Harvard T.H. Chan School of Public Health

665 Huntington Ave.

Building 1, Room 715

Boston, MA, 02115

Contact information

Email: ygrad@hsph.harvard.edu; tel: 617-432-2275

<u>Abstract</u>

The etiologic agent of gonorrhea, *Neisseria gonorrhoeae*, has become resistant to each of the

first line antibiotics used to treat it. However, most isolates in the US remain susceptible to ciprof-

loxacin, leading to the hypothesis that a diagnostic that accurately reports susceptibility to ciprof-

loxacin may allow its reintroduction into clinical use. One diagnostic approach is to identify codon

91 in *gyrA*, where coding for the wildtype serine is associated with ciprofloxacin susceptibility and

phenylalanine with resistance. Here, we found that escape from this diagnostic method can occur

through *gyrA* F91S reversion and expansion of pre-existing ciprofloxacin non-susceptible *N. gon-*

orrhoeae lineages with *gyrA* 91S. Furthermore, we found that mutations conferring ciprofloxacin

resistance in *gyrA* 91S isolates led to cross-resistance with zoliflodacin, an antibiotic in phase 3

trial for treatment of gonorrhea. These findings define *N. gonorrhoeae* mutations that should be

monitored via surveillance systems, particularly in populations and regions deploying *gyrA* 91-

based diagnostics, and demonstrate how diagnostics that guide antibiotic therapy can have unin-

tended consequences.

Introduction

Neisseria gonorrhoeae, the causative agent of the sexually transmitted disease gonorrhea, has developed resistance to each of the first-line antibiotics used to treat it[1]. This rise in antibiotic resistance has led to fears of untreatable highly drug resistant gonorrhea[2], a concern further heightened by the international circulation of strains resistant to the current first-line antibiotic, ceftriaxone[3-5].

Due to high levels of drug resistance in *N. gonorrhoeae*, there has been substantial interest in the development of new approaches for management of gonorrhea[6]. One promising avenue has been the use of point-of-care diagnostics that test for susceptibility. By understanding alleles associated with antibiotic susceptibility[7], sequence-based diagnostics may enable diagnosis of gonorrhea and inform treatment decisions by using genotype to predict antibiotic susceptibility phenotype.

An early target for this approach has been the fluoroquinolone class antibiotic ciprofloxacin[8,9]. Once first-line treatment of *N. gonorrhoeae* infections, rising levels of resistance and concern for failure of empiric therapy resulted in recommendations against its empiric use[10]. However, surveillance data from the US, the UK, and Europe indicate that ciprofloxacin resistance is present in a minority of cases[11-13], suggesting that a diagnostic that can inform on ciprofloxacin susceptibility would allow for the drug's reintroduction to gonorrhea treatment and thereby aid in reducing the selective pressure on ceftriaxone[14].

Ciprofloxacin resistance in *N. gonorrhoeae* is mainly mediated by mutations in the A subunit of DNA gyrase (encoded by *gyrA*) and the A subunit of topoisomerase IV (encoded by *parC*). Though multiple mutations contribute to ciprofloxacin resistance, including position 95 in GyrA and multiple sites at positions 86-91 in ParC[15], position 91 in GyrA is highly predictive of clinical susceptibility (if a serine) and resistance (if a phenylalanine) to ciprofloxacin[16,17].

The SpeeDx ResistancePlus GC diagnostic test reports the presence of *N. gonorrhoeae* and

ciprofloxacin susceptibility. The ResistancePlus GC test consists of a multiplex qPCR, incorpo-

rating channels for the wild-type (WT) *gyrA* 91S allele, the mutant *gyrA* 91F allele, the *N. gonor-*

rhoeae opa gene, the *N. gonorrhoeae porA* gene, and a PCR positive control. Across multiple

studies, the ResistancePlus assay detects gonorrhea with a sensitivity of 94-100%[8,18,19] and spec-

ificity of 99-100%[8,19]. The test is also highly sensitive (>99%) and specific (>97%) for ciprofloxacin

resistance[18].The ResistancePlus GC assay has been approved in Europe and received a break-

through designation from the FDA[20].

The use of sequence-based diagnostics raises the possibility of diagnostic escape, with a prece-

dent in *N. gonorrhoeae* of mutations that resulted in failure of clinical identification by PCR [21,22].

In this case, the diagnostic applies selective pressure for *N. gonorrhoeae* to develop resistance

through a pathway other than the *gyrA* 91 mutation it targets. False negative tests for antibiotic

susceptibility testing may result in clinical treatment failure for the individual and increased burden

of disease and accelerated spread of resistance in the population[23]. The risk of ciprofloxacin re-

sistance arising through novel pathways is especially pressing due to concern for cross-resistance

to the new *N. gonorrhoeae* antibiotic zoliflodacin, a first-in-class spiropyrimidinetrione antibiotic

that primarily targets the B subunit of DNA gyrase (GyrB)[24] and is in Phase 3 trials for treatment

of gonorrhea[25].

In this study, we investigated the pathways of diagnostic escape from tests that rely on GyrA 91.

We found that some clinical isolates with a 91S allele of GyrA maintain ciprofloxacin resistance

at a level that has been associated with treatment failure, with one such isolate (GCGS0481)

acquiring high-level ciprofloxacin resistance while maintaining the 91S allele. Finally, though

these strains are zoliflodacin naïve, we found that, in the presence of ciprofloxacin, clinical isolates

readily evolve a D429N mutation in GyrB that confers ciprofloxacin resistance and that has been

reported to confer zoliflodacin resistance[26]. Taken together, these results indicate that there are

evolutionary pathways to diagnostic escape from *gyrA* 91-based diagnostics and that one of these

pathways leads to cross-resistance between ciprofloxacin and zoliflodacin. Finally, these results

provide experimental validation for the hypothesis that single antibiotic sequence-based diagnos-

tics may lead to unanticipated consequences, including novel resistance determinants and cross-

resistance.

Methods

Bacterial strains, MIC measurements and Cloning

The strains used in this study are listed in Supplementary Table 1. *N. gonorrhoeae* was grown on

GCB Agar (Difco) supplemented with Kellogg's supplements[27] at 37°C in 5% CO_2. Antibiotic sus-

ceptibility testing was performed on GCB-K agar via agar dilution for zoliflodacin (using the range

0.008 to 4 µg/mL) and Etest (BioMerieux) for ciprofloxacin, ceftriaxone, erythromycin, benzylpen-

icillin, and tetracycline.

Cloning was performed using primers and plasmids listed in Supp. Table 1 and Gibson assembly

[28] into a pUC19[29] backbone using a kanamycin cassette from pDR1[30]. Fragments were amplified

using Phusion (NEB), checked for appropriate size by gel electrophoresis, column purified (Qi-

agen PCR Purification kit), assembled with Gibson Master Mix (NEB), and finally transformed into

chemically competent DH5α *E. coli*. Finally, individual colonies were selected on LB agar with 50

µg/mL kanamycin, picked, and grown overnight. Plasmids were isolated using spin columns (Qi-

agen Spin Miniprep Kit). The resulting plasmids were checked by Sanger sequencing. For inser-

tion of GyrA alleles into *N. gonorrhoeae*, strains of *N. gonorrhoeae* in Supp. Table 2 were grown

overnight on GCB-K. After 16-20 hours, strains were scraped into 0.3M sucrose, electroporated

(V=1.8 kV, exponentially decaying wave) with 200 ng of plasmids DRE77-82, and rescued with

GCP medium (15 g/L proteose peptone 3 (ThermoFisher), 1 g/L soluble starch (ThermoFisher),

4 g/L K_2HPO_4, 1 g/L KH_2PO_4, 5 g/L NaCl)[31] with Kellogg's supplement. After 10 minutes of rescue,

transformants were then plated on nonselective GCB-K agar for 4 hours followed by selection on

GCB-K supplemented with 70 μg/mL kanamycin. Finally, single colonies were re-streaked on

GCB-K agar and checked for GyrA allele by Sanger sequencing. For GyrB cloning, fragments of

GyrB were amplified using primers DR542 and DR543. Electroporation was performed as above,

and single colonies were selected on GCB-K agar supplemented with 0.5 μg/mL ciprofloxacin.

Genomic analysis

The genomic pipeline was conducted with methodology as published.[32] Reads and metadata for

the sequenced *N. gonorrhoeae* isolates were accessed from multiple sources[33-62]. Reads were

then inspected using FastQC (v0.11.7) (https://www.bioinformatics.babraham.ac.uk/pro-

jects/fastqc/) and removed based on divergent GC content (from ~52-54%) or poor base quality.

The remaining reads were mapped to the genome of *N. gonorrhoeae* strain NCCP11945 (RefSeq

accession NC_011035.1) with BWA-MEM (v0.7.17-r1188)[63]. Mapped reads were then dedupli-

cated with Picard (v2.20.1) (https://github.com/ broadinstitute/picard). Samples with <40% cover-

age or <70% reads aligned were discovered with BamQC in Qualimap (v2.2.1)[64] and discarded.

Variants, including those in *gyrA* codons 91 and 95, were called with Pilon (v1.23)[65] with a mini-

mum mapping quality of 20 and a minimum depth of 10. Pseudogenomes were generated from

the resulting Pilon VCFs with inclusion of all PASS sites and alternate alleles with allele frequency

greater than 0.9. All other sites were set to "N", and finally samples with >15% of sites called as

missing were excluded. SPAdes (v3.12.0, with 8 threads, the –careful flag, and paired end reads

where available)[66] was used to create *de novo* assemblies, followed by contig filtering for quality

(coverage >10X, length >500 bp, total genome size ~2.0-2.4 Mbp). These assemblies were an-

notated using Prokka (v1.14.6)[67]. Finally, the phylogenetic tree in this study was assembled in a

recombination-corrected manner using Gubbins (v2.3.4)[68] from all sequenced isolates from Reim-

che *et al.*[55], representing lineages recently transmitting in the US, as well as the five genetically

manipulated strains in Table 1 and all strains that had GyrA^{91S} and GyrA$^{95G/A/N}$. The phylogeny

was visualized in iTOL (v6)[69].

Transformants in this study were sequenced at SeqCenter (seqcenter.com). Sample libraries

were assembled using the Illumina DNA Prep Kit and IDT 10bp UDI indices. The resulting libraries

were sequenced with an Illumina NextSeq 2000. The paired-end 2x151 bp reads were demulti-

plexed, followed by quality control and adapter trimming, using bcl-convert (v3.9.3) (https://sup-

[port.illumina.com/sequencing/sequencing_software/bcl-convert.html](https://support.illumina.com/sequencing/sequencing_software/bcl-convert.html)).   Resulting   reads   were

merged using BBMerge (v38.97)[70]. Reads were mapped to the parental genomic DNA assembly

and variants predicted in Geneious 2021.0 (www.geneious.com) using default settings.

Experimental evolution

Strain GCGS0481_SG (Supp. Table 1) was grown overnight on GCB-K agar. The overnight

streak was scraped, diluted to ~OD 0.1, then plated as 14 replicates overnight onto GCB-K agar

supplemented with 0.125 µg/mL ciprofloxacin. Each replicate was sequentially passaged, follow-

ing the same dilution pattern, onto double the concentration of ciprofloxacin overnight until there

was no visible growth or until 2 µg/mL was reached. Single colonies from the last resulting viable

passage were selected for further analysis.

Results

We selected five clinical *N. gonorrhoeae* isolates (Supp. Table 2) based on three major factors:

high-level ciprofloxacin resistance, genetic and geographic diversity, and varying sets of re-

sistance alleles (Figure 1A). All five of the strains encode GyrA^{S91F} as well as an additional muta-

tion at GyrA position 95. All five strains carried mutations in ParC that are known to cause reduced

susceptibility to ciprofloxacin, including ParC$^{S87R/N}$ and ParC$^{E91G/K}$ [71,72]. Finally, all five strains had

the GyrB^{429D} allele associated with zoliflodacin susceptibility[26].

We introduced six different alleles of GyrA into each strain using kanamycin co-selection, includ-

ing pairwise combinations of codon 91 (91S – susceptible, 91F – resistant) and codon 95 (95D –

susceptible, 95G, 95N – resistant). We were particularly interested in strains with the 91S codon,

as these isolates would be reported as ciprofloxacin susceptible by assays focused on this codon.

In all backgrounds, introduction of the GyrA$^{91S/95D}$ allele, containing the susceptible variant at both

codon 91 and 95, conferred susceptibility to ciprofloxacin. There was substantial MIC variation

even at this baseline, with MICs varying almost 10-fold (Table 1).

Introduction of a mutation at the 95 codon, either GyrA^{95G} or GyrA^{95N}, increased the MIC in all

backgrounds by approximately 10-fold. Some isolates, including NY0842, a 2013 urethral isolate

from New York City, and GCGS0481, a 2006 urethral isolate from U.S. Department of Health and

Human Services Region 10 (Alaska, Idaho, Oregon, and Washington), reached an intermediate

level of ciprofloxacin resistance by the Clinical and Laboratory Standards Institute (CLSI) break-

point (≥0.125 µg/mL)[73]. Notably, MICs for ciprofloxacin ≥0.125 µg/mL are associated with sub-

stantially increased risk of treatment failure[74]. Finally, introduction of GyrA$^{91F/95G/N}$ into all five back-

grounds did not substantially affect MICs compared to the original isolates (Table 1, Supp. Table

2), indicating that the co-selected kanamycin cassette minimally affected the ciprofloxacin re-

sistance profile.

These data suggest that some clinical isolates still retain ciprofloxacin non-susceptibility even

after reversion of the GyrA^{91F} allele to GyrA^{91S}. To evaluate whether examples with this genotype

and phenotype have been observed, we investigated *gyrA* genotypes in 11,355 clinical isolates

with published genomes and ciprofloxacin MICs. We identified 30 isolates bearing GyrA$^{91S/95G/N/A}$,

distributed across the *N. gonorrhoeae* phylogeny as singletons and closely related isolates (Fig-

ure 1A), suggesting that the GyrA$^{91S/95G/N/A}$ alleles can arise in multiple genetic backgrounds and

are sufficiently fit for transmission. The reported MICs for these isolates varied from 0.023 µg/mL

to 0.25 µg/mL, including four with MICs above the clinically defined threshold where treatment

failure is likely to occur (Figure 1B). Thus, diagnostic escape could occur either through reversion

or expansion of specific clades with GyrA^{91S} and reduced susceptibility to ciprofloxacin.

We investigated the possibility that GyrA^{91S} isolates could develop higher level ciprofloxacin re-

sistance through experimental evolution. We focused on the GyrA$^{91S/95G}$ mutant of GCGS0481,

differing from the parental isolate by only the kanamycin cassette and the 91S allele of *gyrA* as

validated by whole-genome sequencing (WGS). We sequentially passaged GCGS0481

GyrA$^{91S/95G}$ on increasing levels of ciprofloxacin, starting at ½ MIC (0.125 µg/mL) and increasing

two-fold up to 2 µg/mL. Of the 14 parallel cultures, 10 (78%) had no viable colonies at the MIC.

Single colonies from the four remaining cultures were then re-streaked and tested for ciprofloxacin

susceptibility. One colony had a high ciprofloxacin MIC (>32 µg/mL), attributable to an S91F mu-

tation; this colony was excluded from further analysis.

All three remaining mutants were isolated at 1 µg/mL ciprofloxacin, suggesting decreased sus-

ceptibility as compared to the GCGS0481 GyrA$^{91S/95G}$ strain (0.25 µg/mL; Table 2) and clinical

resistance to ciprofloxacin. WGS revealed that all three mutants had maintained GyrA^{91S} and

acquired mutations in GyrB. One isolate had a missense mutation encoding a GyrB^{E469D} mutation.

Introduction of this single-nucleotide polymorphism (SNP) in the parental isolate recapitulated the

decreased susceptibility to ciprofloxacin and also conferred a two-fold increase in MIC for zoli-

flodacin (Table 2).

Two of the evolved strains gained GyrB[D429N] mutations, which have been shown to cause de-

creased susceptibility to zoliflodacin[24,75]; we validated by WGS that these GyrB[D429N] strains had

not acquired other genomic mutations. Introduction of the GyrB[D429N] mutation into the parental

isolate recapitulated the phenotype of decreased susceptibility to ciprofloxacin and led to an 8-

fold increase in the MIC for zoliflodacin as compared to the wildtype and parental GCGS0481

isolates (Table 2). These changes in antibiotic susceptibility are specific to ciprofloxacin and zoli-

flodacin (Supp. Table 3), suggesting that the differences in MIC are not due to general mecha-

nisms of reduced drug susceptibility.

Discussion

Our results indicate that GyrB[D429N] is readily acquired by *N. gonorrhoeae* in the presence of ciprof-

loxacin and that GyrB[D429N] increases ciprofloxacin resistance in at least some backgrounds. As

such, this study raises both conceptual and practical implications for the introduction of assays

using *gyrA* codon 91 to infer ciprofloxacin susceptibility. From a conceptual standpoint, this work

provides an example of how escape from a diagnostic used to guide antibiotic therapy can drive

the acquisition and possible spread of novel resistance determinants and how these pathways

may have unintended consequences for resistance profiles to other antibiotics[23]. The presence

of multiple clinical isolates with GyrA[91S], together with resistance-associated mutations at GyrA[95],

strongly suggest that reversion of codon 91 is viable for pathogenesis. Our findings also indicate

that a test that detects both position 91 and position 95 would be less likely to select for reversion

to the susceptible allele, as all of the GyrA[91S/95D] strains in this study remained highly susceptible

to ciprofloxacin. Though the GyrB[E469] mutation identified in this study has not been observed in

clinical isolates of *N. gonorrhoeae*, the corresponding GyrB[E466D] mutation in clinical isolates of

Salmonella enterica serovar Typhimurium confers decreased susceptibility to ciprofloxacin[76]. This

finding provides further evidence consistent with the hypothesis that diagnostic escape from a

gyrA 91S assay may provide evolutionary pressure to drive novel resistance mechanisms.

The finding that GyrB^{D429N} reduces susceptibility to both ciprofloxacin and zoliflodacin was unex-

pected. Earlier work on zoliflodacin showed that, in ciprofloxacin-resistant clinical isolates from

China, introduction of GyrB^{D429N} led to a decrease in ciprofloxacin susceptibility or had no effect

on ciprofloxacin MIC[26]. Other studies have suggested that there is no cross-resistance between

ciprofloxacin and zoliflodacin in recently sampled clinical isolates[77,78], likely due to the finding that

GyrB^{D429N} mutations have not been identified in sequences from ~13,000 clinical isolates[79]. In

contrast, enzymology suggests that ciprofloxacin has modestly decreased potency for the stability

of cleaved complexes with GyrB^{D429N} as compared to GyrB^{429D} (a CC$_{50}$, a measure of the potency

for stability of cleaved complexes, of 3 +/- 0.4 for the mutant vs. 1.9 +/- 0.7 for the WT)[75]. The

cleaved complex is the adduct of enzyme and DNA formed at DNA nicks that are unable to be

repaired in the presence of fluoroquinolones. As these cleaved complexes are thought to be cor-

related with, if not the mechanism of, fluoroquinolone-mediated cell death[80], these data are con-

sistent with zoliflodacin-ciprofloxacin cross-resistance. Further work is required to understand the

genetic background dependence of phenotypic outcomes following introduction of GyrB^{D429N}. One

intriguing possibility is that GyrB^{D429N} is epistatic to GyrA^{S91F} with respect to ciprofloxacin re-

sistance and is therefore less likely to arise in highly ciprofloxacin resistant backgrounds.

The findings in this work have implications for the introduction of zoliflodacin, pending a successful

phase III trial. From the perspective of zoliflodacin resistance, testing for the GyrB^{D429N} allele

should be prioritized in locations where ciprofloxacin is used, as well as in settings where assays

targeting *gyrA* codon 91 inform treatment decisions. From the perspective of ciprofloxacin use,

the emergence of zoliflodacin resistant isolates may also lead to an increase in ciprofloxacin MIC.

GyrB^{429} should therefore be considered amongst the panel of mutations contributing to ciproflox-

acin resistance and informing antibiotic selection for clinical decision-making.

<u>Acknowledgements</u>

We thank the members of the Grad Lab, particularly Sam Palace and the remainder of the GC

subgroup, for their thoughts on and support for this project. We thank SeqCenter (seqcenter.com)

for sequencing isolates.

<u>Financial Support</u>

This work was supported by NIH T32 GM07753 (DHFR), F30 AI160911-01 (DHFR), F32

AI145157 (TDM), R01 AI132606 and R01 AI153521 (YHG), and the Smith Family Foundation

(YHG).

<u>Conflicts of Interest</u>

Y. H. G. receives or has received support from Welcome Trust, Pfizer, and Merck, consulting

fees from GSK, Quidel, and the NBA, and payment for expert testimony from Merck. He serves

on the advisory board for Day Zero Diagnostics. None of these conflicts has bearing on the sub-

mitted work.

References

1. Unemo, M. & Shafer, W. M. Antimicrobial resistance in Neisseria gonorrhoeae in the 21st century: past, evolution, and future. *Clinical microbiology reviews* **27**, 587-613, doi:10.1128/cmr.00010-14 (2014).

2. Bolan, G. A., Sparling, P. F. & Wasserheit, J. N. The emerging threat of untreatable gonococcal infection. *The New England journal of medicine* **366**, 485-487, doi:10.1056/NEJMp1112456 (2012).

3. Unemo, M. & Nicholas, R. A. Emergence of multidrug-resistant, extensively drug-resistant and untreatable gonorrhea. *Future microbiology* **7**, 1401-1422, doi:10.2217/fmb.12.117 (2012).

4. Nakayama, S. *et al.* New Ceftriaxone- and Multidrug-Resistant Neisseria gonorrhoeae Strain with a Novel Mosaic penA Gene Isolated in Japan. *Antimicrobial agents and chemotherapy* **60**, 4339-4341, doi:10.1128/aac.00504-16 (2016).

5. Eyre, D. W. *et al.* Gonorrhoea treatment failure caused by a Neisseria gonorrhoeae strain with combined ceftriaxone and high-level azithromycin resistance, England, February 2018. *Euro Surveill* **23**, doi:10.2807/1560-7917.ES.2018.23.27.1800323 (2018).

6. Rubin, D. H. F., Ross, J. D. C. & Grad, Y. H. The frontiers of addressing antibiotic resistance in Neisseria gonorrhoeae. *Transl Res*, doi:10.1016/j.trsl.2020.02.002 (2020).

7. Mortimer, T. D., Zhang, J. J., Ma, K. C. & Grad, Y. H. Loci for prediction of penicillin and tetracycline susceptibility in Neisseria gonorrhoeae: a genome-wide association study. *Lancet Microbe* **3**, e376-e381, doi:10.1016/s2666-5247(22)00034-9 (2022).

8. Ebeyan, S. *et al.* Evaluation of the ResistancePlus GC (beta) assay: a commercial diagnostic test for the direct detection of ciprofloxacin susceptibility or resistance in Neisseria gonorrhoeae. *The Journal of antimicrobial chemotherapy* **74**, 1820-1824, doi:10.1093/jac/dkz108 (2019).

9. Klausner, J. D. *et al.* Resistance-Guided Treatment of Gonorrhea: A Prospective Clinical Study. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **73**, 298-303, doi:10.1093/cid/ciaa596 (2021).

10. CDC. Update to CDC's sexually transmitted diseases treatment guidelines, 2006: fluoroquinolones no longer recommended for treatment of gonococcal infections. *MMWR. Morbidity and mortality weekly report* **56**, 332-336 (2007).

11. CDC. Sexually Transmitted Disease Surveillance - 2018. (2018).

12. Whittles, L. K., White, P. J., Paul, J. & Didelot, X. Epidemiological Trends of Antibiotic Resistant Gonorrhoea in the United Kingdom. *Antibiotics (Basel, Switzerland)* **7**, doi:10.3390/antibiotics7030060 (2018).

13. ECDC. Gonococcal antimicrobial susceptibility surveillance in Europe - Results summary 2017. *ECDC Surveillance Report* (2019).

14. Allan-Blitz, L. T. *et al.* Implementation of a Rapid Genotypic Assay to Promote Targeted Ciprofloxacin Therapy of Neisseria gonorrhoeae in a Large Health System. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **64**, 1268-1270, doi:10.1093/cid/ciw864 (2017).

15. Mortimer, T. D. & Grad, Y. H. Applications of genomics to slow the spread of multidrug-resistant Neisseria gonorrhoeae. *Annals of the New York Academy of Sciences* **1435**, 93-109, doi:10.1111/nyas.13871 (2019).

16. Allan-Blitz, L. T. & Klausne, J. D. Neisseria gonorrhoeae susceptibility to ciprofloxacin. *MLO Med Lab Obs* **48**, 30 (2016).

17. Hicks, A. L. *et al.* Evaluation of parameters affecting performance and reliability of machine learning-based antibiotic susceptibility testing from whole genome sequencing data. *PLoS computational biology* **15**, e1007349, doi:10.1371/journal.pcbi.1007349 (2019).

18   Cotton, S., McHugh, M. P., Etherson, M., Shepherd, J. & Templeton, K. E. Evaluation of
the molecular detection of ciprofloxacin resistance in Neisseria gonorrhoeae by the
ResistancePlus GC assay (SpeeDx). *Diagnostic microbiology and infectious disease* **99**,
115262, doi:10.1016/j.diagmicrobio.2020.115262 (2021).
19   Lee, D. Y. J. *et al.* Reflex Detection of Ciprofloxacin Resistance in Neisseria gonorrhoeae
by Use of the SpeeDx ResistancePlus GC Assay. *J Clin Microbiol* **59**,
doi:10.1128/jcm.00089-21 (2021).
20   BusinessWire. in *Bloomberg.com* (2019).
21   Whiley, D. M. *et al.* False-negative results using Neisseria gonorrhoeae porA pseudogene
PCR - a clinical gonococcal isolate with an N. meningitidis porA sequence, Australia,
March 2011. *Euro surveillance : bulletin Europeen sur les maladies transmissibles =
European communicable disease bulletin* **16** (2011).
22   Golparian, D., Johansson, E. & Unemo, M. Clinical Neisseria gonorrhoeae isolate with a
N. meningitidis porA gene and no prolyliminopeptidase activity, Sweden, 2011: danger of
false-negative genetic and culture diagnostic results. *Euro surveillance : bulletin Europeen
sur les maladies transmissibles = European communicable disease bulletin* **17** (2012).
23   Tuite, A. R. *et al.* Impact of Rapid Susceptibility Testing and Antibiotic Selection Strategy
on the Emergence and Spread of Antibiotic Resistance in Gonorrhea. *J Infect Dis* **216**,
1141-1149, doi:10.1093/infdis/jix450 (2017).
24   Basarab, G. S. *et al.* Discovery of Novel DNA Gyrase Inhibiting Spiropyrimidinetriones:
Benzisoxazole Fusion with N-Linked Oxazolidinone Substituents Leading to a Clinical
Candidate (ETX0914). *Journal of medicinal chemistry* **58**, 6264-6282,
doi:10.1021/acs.jmedchem.5b00863 (2015).
25   Bradford, P. A., Miller, A. A., O'Donnell, J. & Mueller, J. P. Zoliflodacin: An Oral
Spiropyrimidinetrione Antibiotic for the Treatment of Neisseria gonorrheae, Including
Multi-Drug-Resistant Isolates. *ACS Infect Dis* **6**, 1332-1345,
doi:10.1021/acsinfecdis.0c00021 (2020).
26   Alm, R. A. *et al.* Characterization of the novel DNA gyrase inhibitor AZD0914: low
resistance potential and lack of cross-resistance in Neisseria gonorrhoeae. *Antimicrob
Agents Chemother* **59**, 1478-1486, doi:10.1128/AAC.04456-14 (2015).
27   Kellogg, D. S., Jr., Peacock, W. L., Jr., Deacon, W. E., Brown, L. & Pirkle, D. I. Neisseria
Gonorrhoeae. I. Virulence Genetically Linked to Cloncal Variation. *Journal of bacteriology*
85, 1274-1279, doi:10.1128/jb.85.6.1274-1279.1963 (1963).
28   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred
kilobases. *Nat Methods* **6**, 343-345, doi:10.1038/nmeth.1318 (2009).
29   Norrander, J., Kempe, T. & Messing, J. Construction of improved M13 vectors using
oligodeoxynucleotide-directed mutagenesis. *Gene* **26**, 101-106, doi:10.1016/0378-
1119(83)90040-9 (1983).
30   Rubin, D. H. F. *et al.* Variation in supplemental carbon dioxide requirements defines
lineage-specific antibiotic resistance acquisition in Neisseria gonorrhoeae.
bioRxiv, 2022.2002.2024.481660, doi:10.1101/2022.02.24.481660 (2022).
31   Dillard, J. P. Genetic Manipulation of Neisseria gonorrhoeae. *Current protocols in
microbiology* **0 4**, Unit4A.2, doi:10.1002/9780471729259.mc04a02s23 (2011).
32   Ma, K. C. *et al.* Increased antibiotic susceptibility in Neisseria gonorrhoeae through
adaptation to the cervical environment. *bioRxiv*, 2020.2001.2007.896696,
doi:10.1101/2020.01.07.896696 (2020).
33   Alfsnes, K. *et al.* Genomic epidemiology and population structure of Neisseria
gonorrhoeae in Norway, 2016-2017. *Microb Genom* **6**, doi:10.1099/mgen.0.000359
(2020).

34 Buckley, C. *et al.* Use of whole genome sequencing to investigate an increase in Neisseria
gonorrhoeae infection among women in urban areas of Australia. *Scientific reports* **8**,
1503, doi:10.1038/s41598-018-20015-x (2018).
35 Cehovin, A. *et al.* Identification of Novel Neisseria gonorrhoeae Lineages Harboring
Resistance Plasmids in Coastal Kenya. *J Infect Dis* **218**, 801-808,
doi:10.1093/infdis/jiy240 (2018).
36 Demczuk, W. *et al.* Whole-genome phylogenomic heterogeneity of Neisseria gonorrhoeae
isolates with decreased cephalosporin susceptibility collected in Canada between 1989
and 2013. *J Clin Microbiol* **53**, 191-200, doi:10.1128/jcm.02589-14 (2015).
37 Demczuk, W. *et al.* Genomic Epidemiology and Molecular Resistance Mechanisms of
Azithromycin-Resistant Neisseria gonorrhoeae in Canada from 1997 to 2014. *J Clin
Microbiol* **54**, 1304-1313, doi:10.1128/jcm.03195-15 (2016).
38 De Silva, D. *et al.* Whole-genome sequencing to determine transmission of Neisseria
gonorrhoeae: an observational study. *The Lancet. Infectious diseases* **16**, 1295-1303,
doi:10.1016/s1473-3099(16)30157-8 (2016).
39 Didelot, X. *et al.* Genomic Analysis and Comparison of Two Gonorrhea Outbreaks. *mBio*
7, doi:10.1128/mBio.00525-16 (2016).
40 Eyre, D. W. *et al.* WGS to predict antibiotic MICs for Neisseria gonorrhoeae. *The Journal
of antimicrobial chemotherapy* **72**, 1937-1947, doi:10.1093/jac/dkx067 (2017).
41 Ezewudo, M. N. *et al.* Population structure of Neisseria gonorrhoeae based on whole
genome data and its relationship with antibiotic resistance. *PeerJ* **3**, e806,
doi:10.7717/peerj.806 (2015).
42 Fifer, H. *et al.* Sustained transmission of high-level azithromycin-resistant Neisseria
gonorrhoeae in England: an observational study. *The Lancet. Infectious diseases* **18**, 573-
581, doi:10.1016/s1473-3099(18)30122-1 (2018).
43 Gernert, K. M. *et al.* Azithromycin susceptibility of Neisseria gonorrhoeae in the USA in
2017: a genomic analysis of surveillance data. *Lancet Microbe* **1**, e154-e164,
doi:10.1016/s2666-5247(20)30059-8 (2020).
44 Golparian, D. *et al.* Genomic epidemiology of Neisseria gonorrhoeae elucidating the
gonococcal antimicrobial resistance and lineages/sublineages across Brazil, 2015-16.
The Journal of antimicrobial chemotherapy **75**, 3163-3172, doi:10.1093/jac/dkaa318
(2020).
45 Grad, Y. H. *et al.* Genomic Epidemiology of Gonococcal Resistance to Extended-
Spectrum Cephalosporins, Macrolides, and Fluoroquinolones in the United States, 2000-
2013. *J Infect Dis* **214**, 1579-1587, doi:10.1093/infdis/jiw420 (2016).
46 Harris, S. R. *et al.* Public health surveillance of multidrug-resistant clones of Neisseria
gonorrhoeae in Europe: a genomic survey. *The Lancet. Infectious diseases* **18**, 758-768,
doi:10.1016/s1473-3099(18)30225-1 (2018).
47 de Korne-Elenbaas, J., Bruisten, S. M., de Vries, H. J. C. & Van Dam, A. P. Emergence
of a Neisseria gonorrhoeae clone with reduced cephalosporin susceptibility between 2014
and 2019 in Amsterdam, The Netherlands, revealed by genomic population analysis. *J
Antimicrob Chemother* **76**, 1759-1768, doi:10.1093/jac/dkab082 (2021).
48 Kwong, J. C. *et al.* Whole-genome sequencing reveals transmission of gonococcal
antibiotic resistance among men who have sex with men: an observational study. *Sexually
transmitted infections* **94**, 151-157, doi:10.1136/sextrans-2017-053287 (2018).
49 Lan, P. T. *et al.* Genomic analysis and antimicrobial resistance of Neisseria gonorrhoeae
isolates from Vietnam in 2011 and 2015-16. *The Journal of antimicrobial chemotherapy*
75, 1432-1438, doi:10.1093/jac/dkaa040 (2020).
50 Lee, R. S. *et al.* Genomic epidemiology and antimicrobial resistance of Neisseria
gonorrhoeae in New Zealand. *The Journal of antimicrobial chemotherapy* **73**, 353-364,
doi:10.1093/jac/dkx405 (2018).

51   Mortimer, T. D. *et al.* The Distribution and Spread of Susceptible and Resistant Neisseria
gonorrhoeae Across Demographic Groups in a Major Metropolitan Center. *Clinical*
infectious diseases : an official publication of the Infectious Diseases Society of America
73, e3146-e3155, doi:10.1093/cid/ciaa1229 (2021).

52   Parmar, N. R. *et al.* Genomic Analysis Reveals Antibiotic-Susceptible Clones and
Emerging Resistance in Neisseria gonorrhoeae in Saskatchewan, Canada. *Antimicrobial*
agents and chemotherapy **64**, doi:10.1128/aac.02514-19 (2020).

53   Peng, J. P. *et al.* A Whole-genome Sequencing Analysis of Neisseria gonorrhoeae
Isolates in China: An Observational Study. *EClinicalMedicine* **7**, 47-54,
doi:10.1016/j.eclinm.2019.01.010 (2019).

54   Pinto, M. *et al.* Fifteen years of a nationwide culture collection of Neisseria gonorrhoeae
antimicrobial resistance in Portugal. *Eur J Clin Microbiol Infect Dis* **39**, 1761-1770,
doi:10.1007/s10096-020-03907-7 (2020).

55   Reimche, J. L. *et al.* Genomic Analysis of the Predominant Strains and Antimicrobial
Resistance Determinants Within 1479 Neisseria gonorrhoeae Isolates From the US
Gonococcal Isolate Surveillance Project in 2018. *Sexually transmitted diseases* **48**, S78-
s87, doi:10.1097/olq.0000000000001471 (2021).

56   Ryan, L. *et al.* Antimicrobial resistance and molecular epidemiology using whole-genome
sequencing of Neisseria gonorrhoeae in Ireland, 2014-2016: focus on extended-spectrum
cephalosporins and azithromycin. *Eur J Clin Microbiol Infect Dis* **37**, 1661-1672,
doi:10.1007/s10096-018-3296-5 (2018).

57   Sánchez-Busó, L. *et al.* The impact of antimicrobials on gonococcal evolution. *Nature*
Microbiology **4**, 1941-1950, doi:10.1038/s41564-019-0501-y (2019).

58   Thomas, J. C. *et al.* Phylogenomic analysis reveals persistence of gonococcal strains with
reduced-susceptibility to extended-spectrum cephalosporins and mosaic penA-34. *Nature*
communications **12**, 3801, doi:10.1038/s41467-021-24072-1 (2021).

59   Town, K. *et al.* Phylogenomic analysis of Neisseria gonorrhoeae transmission to assess
sexual mixing and HIV transmission risk in England: a cross-sectional, observational,
whole-genome sequencing study. *The Lancet. Infectious diseases* **20**, 478-486,
doi:10.1016/s1473-3099(19)30610-3 (2020).

60   Jamoralin, M. C., Jr. *et al.* Genomic surveillance of Neisseria gonorrhoeae in the
Philippines, 2013-2014. *Western Pac Surveill Response J* **12**, 17-25,
doi:10.5365/wpsar.2020.11.1.005 (2021).

61   Williamson, D. A. *et al.* Bridging of Neisseria gonorrhoeae lineages across sexual
networks in the HIV pre-exposure prophylaxis era. *Nature communications* **10**, 3988,
doi:10.1038/s41467-019-12053-4 (2019).

62   Yahara, K. *et al.* Genomic surveillance of Neisseria gonorrhoeae to investigate the
distribution and evolution of antimicrobial-resistance determinants and lineages. *Microb*
Genom **4**, doi:10.1099/mgen.0.000205 (2018).

63   Heng, L. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv*, doi:1303.3997 (2013).

64   García-Alcalde, F. *et al.* Qualimap: evaluating next-generation sequencing alignment
data. *Bioinformatics* **28**, 2678-2679, doi:10.1093/bioinformatics/bts503 (2012).

65   Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection
and genome assembly improvement. *PLoS One* **9**, e112963,
doi:10.1371/journal.pone.0112963 (2014).

66   Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J Comput Biol* **19**, 455-477, doi:10.1089/cmb.2012.0021 (2012).

67   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-2069,
doi:10.1093/bioinformatics/btu153 (2014).

68   Croucher, N. J. *et al.* Rapid phylogenetic analysis of large samples of recombinant
bacterial whole genome sequences using Gubbins. *Nucleic acids research* **43**, e15,
doi:10.1093/nar/gku1196 (2015).
69   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. *Nucleic acids research* **49**, W293-w296, doi:10.1093/nar/gkab301
(2021).
70   Bushnell, B., Rood, J. & Singer, E. BBMerge - Accurate paired shotgun read merging via
overlap. *PloS one* **12**, e0185056, doi:10.1371/journal.pone.0185056 (2017).
71   Trees, D. L., Sandul, A. L., Whittington, W. L. & Knapp, J. S. Identification of novel
mutation patterns in the parC gene of ciprofloxacin-resistant isolates of Neisseria
gonorrhoeae. *Antimicrob Agents Chemother* **42**, 2103-2105 (1998).
72   Belland, R. J., Morrison, S. G., Ison, C. & Huang, W. M. Neisseria gonorrhoeae acquires
mutations in analogous regions of gyrA and parC in fluoroquinolone-resistant isolates.
Molecular microbiology **14**, 371-380, doi:10.1111/j.1365-2958.1994.tb01297.x (1994).
73   (CLSI), C. a. L. S. I. *Performance standards for antimicrobial susceptibility testing CLSI*
supplement M100. 30th edn, (Performance standards for antimicrobial susceptibility
testing, 30th ed CLSI supplement M100 Clinical and Laboratory Standards Institute,
2020).
74   Chaudry, A. E. & Klausner, J. D. A Narrative Review of Clinical Treatment Outcomes of
Neisseria gonorrhoeae Infection With Ciprofloxacin by Minimum Inhibitory Concentration
and   Anatomic   Site.   *Sexually   transmitted   diseases*   **48**,   385-392,
doi:10.1097/olq.0000000000001334 (2021).
75   Kern, G. *et al.* Inhibition of Neisseria gonorrhoeae Type II Topoisomerases by the Novel
Spiropyrimidinetrione AZD0914. *The Journal of Biological Chemistry* **290**, 20984-20994,
doi:10.1074/jbc.M115.663534 (2015).
76   Al-Emran, H. M. *et al.* Detection of a Novel gyrB Mutation Associated With
Fluoroquinolone-Nonsusceptible Salmonella enterica serovar Typhimurium Isolated From
a Bloodstream Infection in Ghana. *Clinical infectious diseases : an official publication of*
the Infectious Diseases Society of America **62 Suppl 1**, S47-49, doi:10.1093/cid/civ790
(2016).
77   Unemo, M. *et al.* High susceptibility to zoliflodacin and conserved target (GyrB) for
zoliflodacin among 1209 consecutive clinical Neisseria gonorrhoeae isolates from 25
European countries, 2018. *The Journal of antimicrobial chemotherapy* **76**, 1221-1228,
doi:10.1093/jac/dkab024 (2021).
78   Jacobsson, S. *et al.* High in vitro susceptibility to the first-in-class spiropyrimidinetrione
zoliflodacin among consecutive clinical Neisseria gonorrhoeae isolates from Thailand
(2018) and South Africa (2015-2017). *Antimicrobial agents and chemotherapy* **63**,
doi:10.1128/aac.01479-19 (2019).
79   Adamson, P. C., Lin, E. Y., Ha, S. M. & Klausner, J. D. Using a public database of
Neisseria gonorrhoeae genomes to detect mutations associated with zoliflodacin
resistance.   *The   Journal   of   antimicrobial   chemotherapy*   **76**,   2847-2849,
doi:10.1093/jac/dkab262 (2021).
80   Drlica, K., Malik, M., Kerns, R. J. & Zhao, X. Quinolone-mediated bacterial death.
Antimicrobial agents and chemotherapy **52**, 385-392, doi:10.1128/aac.01617-06 (2008).
532

Figure 1

A

AUNG461
GCGS0481
NY0842
157M
JJJ016

B

Intermediate Resistance

Ciprofloxacin MIC (µg/mL)

Figure 1. (A) Recombination-adjusted phylogenetic tree of 1,510 *N. gonorrhoeae* clinical

isolates. On the innermost ring, the codon at position 91 of GyrA is coded by color, while on the

second ring, the codon at position 95 of GyrA is similarly marked. The ciprofloxacin MIC of each

isolate is shown on the third ring in red, while isolates with GyrA[91S] and GyrA[85N/G/A] are shown on

the outermost ring in black. Isolates from this study are marked by arrows in purple. Tree scale

represents recombination-adjusted number of SNPs. **(B)** Ciprofloxacin MICs of GyrA[91S] and

GyrA[95N/G/A] strains indicated in **(A)**, with the CLSI threshold for intermediate resistance marked

by a red dashed line.

Table 1

Strain	GyrA$^{91S/95D}$	GyrA$^{91S/95G}$	GyrA$^{91S/95N}$	GyrA$^{91F/95D}$	GyrA$^{91F/95G}$	GyrA$^{91F/95N}$
JJJ016	.006	.064	.19	.25	>32	>32
NY0842	.012	.19	.25	.25	>32	>32
AUNG461	.004	.064	.064	.19	12	12
GCGS0481	.023	.25	.25	.50	>32	>32
157M	.004	.047	.094	.25	>32	>32

10   **Table 1.** Ciprofloxacin MICs in µg/mL of five clinical isolates of *N. gonorrhoeae* following

11   introduction of GyrA alleles via kanamycin co-selection.

Table 2

Strain	Ciprofloxacin MIC (µg/mL)	Zoliflodacin MIC (µg/mL)	Ciprofloxacin concentration at isolation
GCGS0481 GyrA$^{91F/95G}$ (parent)	>32	0.25	N/A
GCGS0481 GyrA$^{91S/95G}$	0.25	0.25	N/A
GCGS0481 GyrA$^{91S/95G}$ Cip^{mut} 1	1.5	0.5	1 µg/mL
GCGS0481 GyrA$^{91S/95G}$ GyrB^{E469D}	1.5	0.5	N/A
GCGS0481 GyrA$^{91S/95G}$ Cip^{mut} 2	1.5	2	1 µg/mL
GCGS0481 GyrA$^{91S/95G}$ Cip^{mut} 3	1.5	2	1 µg/mL
GCGS0481 GyrA$^{91S/95G}$ GyrB^{D429N}	1.5	2	N/A

13   **Table 2.** Ciprofloxacin and zoliflodacin MICs of *N. gonorrhoeae* isolates with GyrA[91S/95G]

14   following experimental evolution in the presence of ciprofloxacin.

15