Genetic differences between 129S substrains affect antiretroviral immune responses

Robert Z. Zhang[1,4], Vincent Mele[1,4], Lia Robben[1], Melissa Kane[1,2,3*]

[1]Department of Pediatrics, Division of Infectious Diseases, University of Pittsburgh School of

Medicine, Pittsburgh, PA 15224

[2]RK Mellon Institute for Pediatric Research, UPMC Children's Hospital of Pittsburgh, Pittsburgh,

PA 15224

[3]Center for Microbial Pathogenesis, UPMC Children's Hospital of Pittsburgh, Pittsburgh PA

[4]These authors contributed equally to this work. Author order was determined reverse

alphabetically.

*Address correspondence to Dr. M. Kane, Tel. (412-692-5493); e-mail: kaneme@pitt.edu

Running title: Antiretroviral immune responses in 129S mice

Abstract

Inbred mouse lines vary in their ability to mount protective antiretroviral immune responses, and

even closely related strains can exhibit opposing phenotypes upon retroviral infection. Here, we

found that 129S mice inherit a previously unknown mechanism for the production of anti-murine

leukemia virus (MLV) antibodies and control of infection. The resistant phenotype is controlled

by two dominant loci that are independent from known MLV-resistance genes. We also show

that production of anti-MLV antibodies in 129S7, but not 129S1 mice is independent of

interferon gamma (IFNγ) signaling. Thus, our data indicate that 129S mice inherit an unknown

mechanism for control of MLV infection and demonstrate that there is genetic variability in 129S

substrains that affects their ability to mount antiviral immune responses.

Importance

Understanding the genetic basis for production of protective antiviral immune responses is

crucial for the development of novel vaccines and adjuvants. Additionally, characterizing the

genetic and phenotypic variability in inbred mice has implications for the selection of strains for

targeted mutagenesis, choice of controls, and for broader understanding of the requirements for

protective immunity.

Introduction

Resistance and sensitivity to viral infections depends greatly upon the genetic make-up of the

host. Model organisms such as inbred mice have proved to be invaluable for the investigation of

the pathways underlying protective immune responses due to the inability to perform genetic

manipulations in humans. In this regard, mouse models of murine retroviruses, such as murine

leukemia virus (MLV) and mouse mammary tumor virus (MMTV) have provided essential

insights into the molecular mechanisms underlying anti-viral immune responses. MLV is

transmitted as an exogenous virus through the blood and milk or as an endogenous stably

integrated provirus and primarily infects cells of lymphoid origin (1). Pathogenic MLVs are a

mixture of ecotropic and polytropic [mink cell focus forming (MCF)] viruses, and many contain

an additional replication-defective spleen focus forming virus (SFFV) component that promotes

pathogenesis (1). These viruses cause a range of diseases in susceptible mice, including

lymphomas, leukemias, rhabdomyosarcoma, neurological disorders, and immunosuppression

(2-6). Mice from retrovirus-susceptible strains, such as BALB/c mice, detect retroviral

pathogens, as indicated by the fact that they initiate an antiretroviral response. However, this

response is not long-lasting and is unsuccessful in controlling virus replication (7, 8), which is

most likely due to the numerous mechanisms of immune evasion employed by retroviruses (9-

13). In contrast, pathogen detection in resistant mice translates into a robust, long-lasting, and

virus-neutralizing immune response (7, 9, 14, 15).

37   Classical genetics approaches in mice have identified several genes that play a role in

directly restricting MLV replication, or in the generation of protective antiviral immune responses

[recently reviewed in (16)]. *Fv1* is a capsid-specific restriction factor encoded by an endogenous

retroviral *Gag* gene that governs tropism of MLV. Alleles of *Fv1* are defined by their ability to

block specific subclasses of MLV, with incompatibility resulting in a post-entry block to infection,

(17-20) and in some cases, stimulation of cross-protective neutralizing adaptive immune

responses (21). *Fv2* is a dominant Friend MLV (FV) susceptibility gene that facilitates

splenomegaly and erythroleukemia induction by SFFV (22, 23). *Rfv3* is a dominant FV

resistance gene that is encoded by *Apobec3*. The resistant allele of *Apobec3* inherited by

C57BL/6 and C57BL/10 mice restricts FV replication and promotes neutralizing antibody (Ab)

responses (24). *Vic1* is a recessive MLV and MMTV resistance gene that is encoded by *H2-Ob*,

the beta subunit (Oβ in mice and DOβ in humans) of the nonclassical major histocompatibility

complex class II (MHC-II)-like molecule H2-O (HLA-DO or DO in humans) (25). The resistant,

functionally null allele of *vic1* inherited by I/LnJ mice results in sustained production of virus-

neutralizing Abs. Transfer of the I/LnJ *vic1* locus to multiple retrovirus susceptible backgrounds,

which contain a functional allele of *Ob*, such as BALB/cJ (BALB/c$^{vic1I/LnJ}$ congenic mice) confers

the ability to produce neutralizing Abs against both MMTV and MLV (25, 26). Finally, adaptive

immune responses against retroviruses in mice (and humans) are also affected by classical

MHC class I and II genes, with specific haplotypes directing virus-specific cytotoxic T cell and

Ab responses (16, 22).

Ab responses to soluble protein and carbohydrate antigens in mice are generally

restricted to the IgG1 and IgG3 isotypes (27, 28), while infection of mice with a variety of viral

and parasitic pathogens results in a unique bias in class switching to IgG2a (or IgG2c) Abs (29-

31), including antiretroviral Ab responses in I/LnJ mice (7). In mice, the Th1-type cytokine

interferon gamma (IFNγ) has been well established as the canonical signal for class switching to

the IgG2a/c isotype both in vitro and in the context of infection (32). Specifically, in B6 mice,

IFNγ is required for Ab responses to MLV infection and for long-term viral control (33), and in

I/LnJ mice, IFNγ is essential to produce antibodies of any isotype against retroviral infections (7,

26). However, we recently reported that in BALB/cJ$^{vic1i/i}$ mice, IFNγ is dispensable for the

production of anti-MLV IgG2a Abs (34), indicating that BALB/cJ mice inherit an alternative

mechanism for stimulating antiviral IgG2a Ab production upon viral infection. The ability to

generate IFNγ-independent Ab responses in BALB/cJ mice is controlled by a single recessive

locus on Chromosome (Chr) 9, which we have named *i*nterferon-*g*amma *i*ndependent *I*gG2a

(*igii*) (34).

Reverse genetics approaches in mice are a powerful tool to understand the function of

specific genes and have provided essential research models for numerous diseases. Prior to

the development of CRISPR technology, which allowed for targeted mutagenesis on multiple

genetic backgrounds, most ES cell lines used for targeted mutagenesis came from substrains of

129 mice. The 129 lineage has a complex history, with multiple outcrossing and bottleneck

events, resulting in three major lineages (Parental, Steel, and *Ter*), with substantial substrain

variability ((35, 36) and Figure S1). This complex history has important implications for

investigations using 129 mice, and on analysis of targeted mutations that have been crossed to

other genetic backgrounds. Here, we investigated anti-MLV immune responses in 129S

substrains and determined that 129S1 mice inherit two loci that control antiviral immunity. We

also demonstrate that genetic differences between 129S substrains affect IFNγ-independent

antiviral immune responses.

Results

A dominant genetic mechanism controls anti-MLV immune responses in 129S1 mice.

Previous reports demonstrated that 129P2 mice (representative of the Parental 129 substrains)

inherit the $Fv1^{nr}$ allele, which restricts B-tropic and some N-tropic, but not NB-tropic strains of

MLV (21, 37, 38). 129P2 infected with a B-tropic FV produce robust cross-protective neutralizing

Ab responses due to this $Fv1$-incompatibility. However, although they produce FV-specific Abs

upon infection with an NB-tropic FV, since they inherit susceptible alleles of $Fv2$ and $Rfv3$, they

do not control the infection (21, 39). These reports additionally suggest that the entire 129

lineage is $Fv1^{nr/nr}$ and $Fv2^{s/s}$ (21, 23, 37, 38). Furthermore, genotyping of two Steel substrains

(see Materials and Methods) at the $Rfv3$ locus indicates that the 129 lineage is $Rfv3^{s/s}$. To

investigate antiviral responses in 129 Steel substrains, we infected 129S1/SvImJ (129S1;

developed as a control for steel-derived ES cells (40)) mice with Rauscher-like-MLV (RL-MLV)

and monitored them for antiviral Ab production, development of splenomegaly, and viral titers in

the spleen. RL-MLV is a mixture of a NB-tropic ecotropic virus and a polytropic MCF virus that

causes erythroid leukemias; the mixture does not contain an SFFV component (3), therefore the

Fv2 allele does not affect disease outcome. While susceptible control BALB/cByJ (ByJ) mice do

not produce a robust Ab response and were unable to control infection, 129S1 mice produced a

robust IgG2a Ab response against RL-MLV and the majority (83%) cleared infection and did not

develop splenomegaly (Figure 1). To determine whether the ability to produce protective anti-

MLV Ab responses is inherited in a dominant or recessive fashion, we crossed 129S1 mice to

BALB/cByJ mice (Figure 2A). F_1 mice generated from these crosses were infected with RL-MLV

and screened for antiviral Abs. F_1 mice from crosses of both directions [(129S1xByJ) and

(ByJx129S1) produced antiviral Abs, although IgG2a productions was slightly lower in

(ByJx129S1) F_1 animals (Figure 1A). Additionally, the majority of F_1 mice eliminated the virus

and did not develop splenomegaly (Figure 1B-C), indicating that resistance to RL-MLV is

inherited in a dominant fashion.

Two loci control anti-MLV Ab responses in 129S1 mice. We next investigated how many loci

control anti-MLV Ab production by crossing resistant (129S1xByJ) F_1 mice to susceptible ByJ

mice to generate N_2 mice (Figure 2A). We found that 9 out of 40 (23%) N_2 mice produced

antiviral IgG2a Abs (Figure 2B). We then determined the viral titers in the spleen and spleen

weights of Ab negative and Ab positive N_2 mice and found that the Ab positive (resistant), but

not the Ab negative (susceptible) N_2 mice cleared the infection and had normal spleen weights

(Figure 2B). This three:one ratio of susceptible to resistant N_2 animals indicates that two loci

control the production of protective anti-RL-MLV Ab responses in 129S1 mice. Since the major

histocompatibility complex (MHC) locus is a major factor in controlling antiviral immunity,

including anti-MLV responses (16, 22), and 129S1 mice inherit the protective H2^b haplotype

while ByJ mice inherit the susceptible H2^d haplotype, we next investigated whether the MHC

locus is one of the genetic determinants for anti-MLV Ab responses in 129S1 mice. MHC

haplotypes of N_2 mice were determined by staining peripheral blood lymphocytes with

haplotype-specific Abs for MHC Class I (H2-D^b, H2-D^d) and Class II (I-A^b, I-A^d). We did not

observe significant differences in Ab production, viral titers, and spleen weights in RL-MLV

infected H2$^{b/d}$ and H2$^{d/d}$ N$_2$ mice (Figure 2C and Table 1), indicating that control of infection in N$_2$

mice is independent of the MHC haplotype and that non-MHC loci control antiviral responses in

129S1 mice.

IFNγ-signaling is dispensable for anti-MLV IgG2a-specific Ab production in 129S7 mice.

Following our previous finding that BALB/cJ mice inherit a noncanonical IFNγ-independent

pathway for antiviral IgG2a Ab production (34), we wondered whether this pathway was unique

to this genetic background. Earlier reports indicated that 129S mice also inherit a pathway for

IFNγ-independent production, as IFNγ-receptor 1 (IFNγR)-deficient 129S7 (G129) mice produce

IgG2a Abs against lymphocytic choriomeningitis virus (LCMV) (41) and (lactate dehydrogenase-

elevating virus) LV (42). We therefore investigated whether IFNγR$^{-/-}$ 129S7 mice produce IFNγ-

independent IgG2a Abs against RL-MLV. Control IFNγR$^{-/-}$ ByJ mice did not produce Abs of any

isotype and did not control infection, while IFNγR$^{-/-}$ 129S7 mice produced IgG2a Abs (Figure

3A). Although Ab production was lower than in IFNγR-sufficient 129S1 mice, the majority of

IFNγR$^{-/-}$ 129S7 mice (60%) cleared infection and did not develop splenomegaly (Figure 3 and

Table 1). Therefore, like BALB/cJ mice, 129S7 mice inherit a mechanism for noncanonical,

IFNγ-independent antiretroviral Ab production.

IFNγ-independent anti-MLV IgG2a production in 129S7 mice is a complex trait. We next

sought to investigate whether IFNγ-independent antiretroviral Ab production in 129S7 mice and

BALB/cJ mice is controlled by the same genetic mechanism. Since the resistant allele of *igii-*

inherited by BALB/cJ mice is recessive, we reasoned that IFNγR$^{-/-}$ F$_1$ animals inheriting the

129S7 and BALB/c allele would only produce antiviral IgG2a Abs if the 129S7 allele of *igii* is

resistant. IFNγR-deficient mice are available on the BALB/cByJ background, and we crossed

these mice to IFNγR$^{-/-}$ 129S7 mice. F$_1$ mice generated from these crosses were infected with

RL-MLV and screened for antiviral Abs. These F$_1$ progeny did not display a clear phenotype,

with ~30% of mice producing Abs and controlling infection while the majority did not produce

Abs, developed splenomegaly, and had infectious virus in their spleens (Figure 4 and Table 1).

Therefore, the phenotype is not fully penetrant, and the genetic basis for IFNγ-independent

antiretroviral Ab production in 129S7 mice cannot be determined from these crosses.

Furthermore, although these experiments suggest that 129S7 mice do not inherit a resistant

allele of *igii*, BALB/cByJ and BALB/cJ mice have been separated since 1935, therefore we

cannot exclude the possibility that ByJ and BALB/cJ mice inherit different alleles of *igii*.

Anti-MLV Ab production in 129S1 mice is IFNγ-dependent. Since we were unable to

determine the genetic basis for IFNγ-independent antiretroviral Ab production in 129S7 mice by

crossing to IFNγR$^{-/-}$ ByJ mice, and these strains are deficient for the IFNγ receptor rather than

the cytokine, we decided to generate IFNγ-deficient 129S mice for the investigation of IFNγ-

independent Ab responses. Considering that wild-type mice of the precise genetic background

of 129S7-*Ifngr$^{-/-}$* no longer exist, we selected the 129S1 background for the knock-out of IFNγ,

as they are the control strain recommended by the Jackson Laboratory for lines derived from

multiple 129S substrains. We used a previously validated CRISPR/Cas9 approach that targeted

introns 1 and 3 of the *Ifng* gene, thereby deleting exons 2 and 3, removing four of six alpha

helices ((43) and Figure S2). We infected heterozygous control and IFNγ$^{-/-}$ mice from two

founder lines with RL-MLV and monitored them for antiviral Ab production. Although IFNγ

sufficient controls produced antiviral IgG2a Abs and cleared the infection, IFNγ$^{-/-}$ 129S1 mice

from either founder line failed to produce a robust antiviral Ab response of any isotype (Figure

5). Additionally, less than 20% of IFNγ$^{-/-}$ 129S1 mice cleared the infection, and the majority

developed splenomegaly (Figure 5B-C and Table 1), indicating that unlike 129S7 mice, 129S1

mice do not inherit a pathway for directing IFNγ-independent antiretroviral Ab responses.

Genetic differences, not alternative receptor usage, underlie differential requirements for

IFNγ-signaling in 129S mice. To determine if antiviral IgG2a Ab responses in 129S7 mice are

the result of IFNγ signaling through an alternative receptor, or result from genetic differences

between 129S7 and 129S1 mice, we crossed IFNγR$^{-/-}$ 129S7 mice to IFNγ$^{-/-}$ 129S1 mice and

then intercrossed the resulting F$_1$ generation (Figure 6A). If IFNγ signaling through an

alternative receptor explained virus resistance in IFNγR$^{-/-}$ 129S7 mice, we would expect to

observe antiviral Ab production in IFNγR$^{-/-}$, but not IFNγ$^{-/-}$, or IFNγ/IFNγR$^{-/-}$ F$_2$ mice. On the other

hand, if 129S7 inherit an alternative pathway for IFNγ-independent antiviral Ab production, we

would expect a mixture of resistant and susceptible F$_2$ mice independently of inheritance of

IFNγR- or IFNγ-deficiency. As expected, F$_2$ mice sufficient for both cytokine and receptor

produced anti-MLV IgG2a-specific Abs and cleared the infection (Fig 6B-C). However, antiviral

Ab production was not observed in the majority of IFNγR$^{-/-}$ F$_2$ mice, and only 10% cleared the

infection (Figure 6B-C). Interestingly, more IFNγ$^{-/-}$ and IFNγ/IFNγR$^{-/-}$ F$_2$ mice produced anti-MLV

Abs and cleared infection than IFNγR$^{-/-}$ F$_2$ mice, and 64% of IFNγ$^{-/-}$ and 100% of IFNγ/IFNγR$^{-/-}$ F$_2$

mice cleared the infection (although only four double-deficient mice were infected) (Figure 6B-D

and Table 1). These data strongly suggest that IFNγ does not signal through an alternative

receptor to stimulate antiviral Abs in 129S7 mice and that an unknown genetic factor(s) present

in 129S7, but not 129S1 mice controls IFNγ-independent antiretroviral Ab responses.

Discussion

Here, we demonstrate that 129S1 mice inherit a previously unknown mechanism that controls

the production of protective antiviral Ab production upon MLV infection. This phenotype is

inherited in a dominant fashion and is controlled by two loci (Figures 1-2). Importantly, these loci

are distinct from previously characterized genes that restrict retroviral replication (*Fv1*, *Fv2*,

Rfv3) or promote antiviral immune responses (*vic1*, MHC locus, *Rfv3*), indicating that the

pathway in 129S1 mice potentially encodes for a novel mechanism for the stimulation of

protective immunity. Unlike I/LnJ and C57BL/6 mice, which fully clear RL-MLV infection (15, 26),

MLV resistance in 129S1 mice is not fully penetrant, as ~15% of infected mice fail to produce

antiviral Abs, develop splenomegaly, and retain infectious virus in their spleens (Figures 1 and

5, Table 1). Additionally, although this mechanism does not appear to be sex-linked, there was

a reduction in IgG2a Ab production in (ByJx129S1) F_1 mice, and fewer of them cleared the

infection than 129S1 or (129S1xByJ) F_1 mice (Figure 1 and Table 1), indicating that epigenetic

or environmental factors may play a role in controlling anti-MLV responses in 129S1 mice.

IFNγ is a well-established signal for stimulating CSR to the IgG2a isotype both *in vitro*

and in the context of various viral infections (21, 37, 38), including FV infection in B6 mice (22)

and RL-MLV and MMTV infection in I/LnJ mice (6, 15). We previously demonstrated that

BALB/cJ mice inherit a pathway for the generation of IFNγ-independent IgG2a antiviral Ab

responses that is controlled by a single recessive locus, *igii*, which is mapped to Chr. 9 (34).

Here, we demonstrate that 129S7 mice also inherit a pathway for IFNγ-independent anti-MLV

Ab production. However, the mechanism for IFNγR-independent anti-RL-MLV responses in

129S7 mice is a complex genetic trait, and we were unable to determine whether the pathway is

controlled by *igii* or another locus (or multiple loci). Nevertheless, the ability of both BALB/cJ and

129S7 mice to produce IFNγ-independent Abs against a variety of infections including

retroviruses, herpes simplex virus, LCMV, LV, and parasitic infections (34, 41, 42), indicates

that alternative pathways in addition to the canonical IFNγ-dependent pathway can stimulate

neutralizing Ab responses. These findings additionally highlight the importance of utilizing mice

of multiple genetic backgrounds to investigate the requirements for stimulating protective

antiviral immunity.

Following decades of widespread use of 129 ES cell lines, two reports from 1997

investigated the variation between 129 substrains and found substantial genetic variation

resulting from genetic drift, genetic contamination, and residual heterozygosity from

backcrossing programs (35, 36). This initiated efforts to clarify the nomenclature for the

improvement of gene targeting experiments and the selection and consideration of controls for

comparing phenotypes of genes targeted in ES cells of different 129 substrains (35, 36, 44).

These findings further suggested important features that should be considered for targeted

mutagenesis using 129 substrains 1) chimeras produced using 129 ES cells should be crossed

to a genetically matched substrain for maintenance on the 129 background and 2) selection of a

129 substrain for targeted mutagenesis should involve consideration of different genetic traits

between substrains (35). Like many mice generated prior to these recommendations, chimeric

IFNγR$^{-/-}$ 129S mice were not crossed to a genetically matched substrain, and the appropriate

wild-type control strain is unavailable. The 129S1 substrain is therefore recommended as the

control inbred strain for steel substrain-derived ES cell lines (40). However, we found that while

129S7 mice produce antiviral Ab responses in the absence of IFNγ, this mechanism is not

inherited by 129S1 mice (Figure 5). Some reports have suggested that IFNγ can signal through

an alternative receptor in the absence of *Ifngr1* (45-47), but this does not explain the differences

between 129S1 and 129S7 substrains reported here, as IFNγR$^{-/-}$ (129S1x129S7) F_2 mice did not

produce antiviral Abs (Figure 6). Therefore, for the purposes of investigating the requirements

for IFNγ in antiviral defenses, 129S1 mice are not an appropriate control for 129S7-*Ifngr*$^{-/-}$ mice.

The higher incidence of IFNγ-independent Ab production in IFNγ$^{-/-}$ and IFNγ/IFNγR$^{-/-}$ F_2 mice

than IFNγR$^{-/-}$ F_2 mice (Figure 6) was intriguing. However, while the number of F_2 animals

phenotyped here is sufficient to eliminate alternative receptor usage by IFNγ as the basis for

IFNγR-independent Ab production, retroviral resistance is not a fully penetrant phenotype in

either 129S1 or 129S7-*Ifngr*$^{-/-}$ mice, and it appears that IFNγ-independent antiviral Ab

production in 129S7 mice is a complex trait. Therefore, larger numbers of F_2 mice would be

required to draw any other conclusions regarding the genetic basis for these substrain

differences.

While 129 substrains have the same genotypes at known MLV-resistance loci [(21, 23,

37, 38) and this report], our findings clearly indicate that there are unappreciated differences in

their ability to mount protective antiviral immune responses. As such, it is unknown whether the

129P2/OlaHsd substrain utilized in many investigations on immune responses to FV infection

would exhibit a similar phenotype to 129S1 mice when infected with RL-MLV, or whether they

can produce IFNγ-independent antiviral immune responses. In fact, these substrains have been

separated since the re-establishment of the 129 line in 1948 following the fire at Jackson Labs

and have substantial genetic variation between them ((35, 36, 48) and Figure S1). The different

phenotypes observed here between Steel substrains are not entirely surprising since the 129S1

lineage has been separated from the 129/SvEv lineage, from which the IFNγR$^{-/-}$ 129S7 mice are

derived since 1969. Furthermore, additional crosses occurred in the 129/SvEv lineage prior to

the introduction of the *Ifngr1* mutation in 129S7-derived AB1 ES cells [Figure S1 and (35, 48)],

thereby increasing the genetic divergence of these substrains. Collectively, our findings further

emphasize the need for careful consideration of the genetic background of mutant and control

mice and indicate that genetic differences between 129S substrains can have consequences

well beyond affecting the efficiency of traditional gene targeting methods.

Acknowledgements

We thank Dr. Tatyana Golovkina for support and consultation; S. Gingras and the transgenic

and gene targeting (TGT) core for the generation of *Ifng*$^{-/-}$ mice; and members of the Kane,

Golovkina, and Dr. Alexander Chervonsky laboratories for helpful discussion. This work was

supported by a Pilot Award from the RK Mellon Institute for Pediatric Research (to M.K.). R.Z. is

supported by T32 AI049820. The content is solely the responsibility of the authors.

Author contributions

278 Conception and design: M.K.; Acquisition of data: R.Z.Z., V.M., L.R., and M.K.; Analysis and

279 interpretation of data: R.Z.Z. and M.K.; Drafting the article: M.K.; Revising the article: M.K.,

280 R.Z.Z., and V.M..

Declaration of interests

The authors declare no competing interests

Materials and Methods

Mice

129S1/SvlmJ (129S1 stock #002448) and BALB/cByJ (stock #001026) were purchased from

the Jackson Laboratory. 129-*Ifngr1*^{tm1Agt}/J [(48) also known as G129] mice were purchased from

the Jackson Laboratory (via cryo recovery, stock #002702) and bred and maintained at the

University of Pittsburgh. IFNγR$^{-/-}$ 129 mice were generated using AB1 ES cells derived from

129S7/SvEvBrd-*Hprt*$^{b-m2}$ (129S7) mice, and chimeric founder males were crossed to an

unknown 129/SvEv substrain (although 129S8 mice are commonly referred to as 129/SvEv and

may have been used for these crosses) (48) for simplicity, we refer to these as 129S7-*Ifngr*$^{-/-}$

mice (Figure S1). C.129S7(B6)-*Ifngr1*^{tm1Agt}/J (IFNγR$^{-/-}$ ByJ) were purchased from the Jackson

Laboratory (via cryo recovery, stock #002286) and bred and maintained at the University of

Pittsburgh.

Ifng-deficient 129S1 mice were generated using CRISPR/Cas9 technology. The target

sequences (5′ guide sequence: 5′-GCTGTTTCCCTGCGTAGTTT-3′; 3′ guide sequence: 5′-

TAGAGGCTAACCAGAGCCGA-3′) were previously validated for the generation of conditional

knockout strains on the C57BL/6 background (43). Two male founders with deletions of exons 2

and 3 were selected (Figure S2) and crossed to 129S1 females. Potential off-target sites with

fewer than three mismatches identified by the Cas-OFFinder algorithm (49) were sequenced in

N_1 and N_2 mice, and mice with no mutations at these sites were intercrossed to produce

homozygous *Ifng*-deficient mice.

Mice were genotyped from tail biopsies using real time PCR with specific probes designed for

each gene (Transnetyx, Cordova, TN) or by flow cytometry (see below). Mice of both sexes

were used in equal numbers for each experiment. All animal experiments were performed in the

American Association for the Accreditation of Laboratory Animal Care-accredited, specific-

pathogen-free facility Division of Laboratory Animal Resources, University of Pittsburgh School

of Medicine. Animal protocols were reviewed and approved by the Institutional Animal Care and

Use Committee at The University of Pittsburgh.

Flow cytometry

For MHC genotyping of $[(129S1xByJ)xByJ]N_2$ mice, blood was collected in sodium heparin

collection vials (SAI Infusion Technologies), and peripheral blood lymphocytes were isolated by

overlaying with Ficoll-Paque (Cytivia) and spinning at 11,000xg for 4min at 25^oC. Lymphocytes

were stained with antibodies against Class I [brilliant blue 700 (BB700)-conjugated anti-H-2Db

(BD Biosciences) and phycoerythrin (PE)-conjugated anti-H-2Dd (BD Biosciences)] or Class II

[peridinin chlorophyll (PerCP)-eFluor 710-conjugated anti-H2-Ab1 (I-Ab, Invitrogen) and PE-

conjugated anti-H2-Ad1 (I-Ad, BD Biosciences)] MHC and analyzed on an Attune NxT

cytometer (Life Technologies).

MLV infection

Rauscher-like MuLV (RL-MuLV), a mixture consisting of NB-tropic ecotropic and mink lung cell

focus-forming viruses, was described previously (69) and was provided by T. Golovkina,

University of Chicago. The virus was propagated in SC-1 embryonic mouse fibroblasts (ATCC).

Ecotropic (Eco) viral titers were determined by an infectious center assay (70). Experimental

mice were injected i.p. with 2×10^4 Eco PFUs at 5-8 weeks of age and screened for anti-virus

antibodies and plaque forming units 8-10 weeks later.

ELISA

To detect MuLV Abs in mouse sera, an enzyme-linked immunosorbent assay (ELISA) was

performed as previously described (6, 15). All sera were used at 2×10^{-2} dilutions. Mouse

IgG2a-specific, as well as total IgG-specific secondary antibodies coupled to horseradish

peroxidase (HRP) (Jackson ImmunoResearch) were used to detect anti-virus antibodies.

Backgrounds obtained from incubation with secondary antibodies alone were subtracted from

the values obtained from sera of infected mice.

Statistical analyses

Statistical significance was determined using GraphPad software (one way ANOVA or unpaired

t test).

References

1. Rosenberg N, Jolicoeur P. 1997. Retroviral Pathogenesis, p 475-586. *In* Coffin JM, Hughes SH, Varmus HE (ed), Retroviruses. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

2. Teich N, Wyke J, Bernstein A, Hardy W. 1982. Pathogenesis of retrovirus-induced disease., p 785-988. *In* Weiss R, Teich N, Varmus HE, Coffin JM (ed), RNA Tumor Viruses. Cold Spring Harbor Laboratory.

3. Hook LM, Jude BA, Ter-Grigorov VS, Hartley JW, Morse HC, 3rd, Trainin Z, Toder V, Chervonsky AV, Golovkina TV. 2002. Characterization of a novel murine retrovirus mixture that facilitates hematopoiesis. J Virol 76:12112-22.

4. Davis BR, Brightman BK, Chandy KG, Fan H. 1987. Characterization of a preleukemic state induced by Moloney murine leukemia virus: evidence for two infection events during leukemogenesis. Proc Natl Acad Sci U S A 84:4875-9.

5. Brightman BK, Davis BR, Fan H. 1990. Preleukemic hematopoietic hyperplasia induced by Moloney murine leukemia virus is an indirect consequence of viral infection. J Virol 64:4582-4.

6. Sitbon M, Evans L, Nishio J, Wehrly K, Chesebro B. 1986. Analysis of two strains of Friend murine leukemia viruses differing in ability to induce early splenomegaly: lack of relationship with generation of recombinant mink cell focus-forming viruses. J Virol 57:389-93.

7. Purdy A, Case L, Duvall M, Overstrom-Coleman M, Monnier N, Chervonsky A, Golovkina T. 2003. Unique resistance of I/LnJ mice to a retrovirus is due to sustained interferon gamma-dependent production of virus-neutralizing antibodies. J Exp Med 197:233-43.

8. Chesebro B, Miyazawa M, Britt WJ. 1990. Host genetic control of spontaneous and induced immunity to Friend murine retrovirus infection. Annu Rev Immunol 8:477-99.

376   9.   Dittmer U, He H, Messer RJ, Schimmer S, Olbrich AR, Ohlen C, Greenberg PD,

377   Stromnes IM, Iwashiro M, Sakaguchi S, Evans LH, Peterson KE, Yang G, Hasenkrug

378   KJ. 2004. Functional impairment of CD8(+) T cells by regulatory T cells during persistent

379   retroviral infection. Immunity 20:293-303.

380   10.   Evans DT, Desrosiers RC. 2001. Immune evasion strategies of the primate lentiviruses.

381   Immunol Rev 183:141-58.

382   11.   Jude BA, Pobezinskaya Y, Bishop J, Parke S, Medzhitov RM, Chervonsky AV,

383   Golovkina TV. 2003. Subversion of the innate immune system by a retrovirus. Nat

384   Immunol 4:573-8.

385   12.   Kane M, Case LK, Kopaskie K, Kozlova A, MacDearmid C, Chervonsky AV, Golovkina

386   TV. 2011. Successful transmission of a retrovirus depends on the commensal

387   microbiota. Science 334:245-9.

388   13.   Malim MH, Bieniasz PD. 2012. HIV Restriction Factors and Mechanisms of Evasion.

389   Cold Spring Harb Perspect Med 2:a006940.

390   14.   Kane M, Case LK, Golovkina TV. 2011. Vital role for CD8+ cells in controlling retroviral

391   infections. J Virol 85:3415-23.

392   15.   Kane M, Case LK, Wang C, Yurkovetskiy L, Dikiy S, Golovkina TV. 2011. Innate immune

393   sensing of retroviral infection via Toll-like receptor 7 occurs upon viral entry. Immunity

394   35:135-45.

395   16.   Kane M, Golovkina TV. 2019. Mapping Viral Susceptibility Loci in Mice. Annu Rev Virol

396   6:525-546.

397   17.   Bassin RH, Duran-Troise G, Gerwin BI, Rein A. 1978. Abrogation of Fv-1b restriction

398   with murine leukemia viruses inactivated by heat or by gamma irradiation. J Virol 26:306-

399   15.

400   18.   Best S, Le Tissier P, Towers G, Stoye JP. 1996. Positional cloning of the mouse

401   retrovirus restriction gene Fv1. Nature 382:826-9.

19. Boone LR, Innes CL, Heitman CK. 1990. Abrogation of Fv-1 restriction by genome-

deficient virions produced by a retrovirus packaging cell line. J Virol 64:3376-81.

20. DesGroseillers L, Jolicoeur P. 1983. Physical mapping of the Fv-1 tropism host range

determinant of BALB/c murine leukemia viruses. J Virol 48:685-96.

21. Halemano K, Barrett BS, Li SX, Harper MS, Smith DS, Heilman KJ, Santiago ML. 2013.

Fv1 restriction and retrovirus vaccine immunity in Apobec3-deficient 129P2 mice. PLoS

One 8:e60500.

22. Miyazawa M, Tsuji-Kawahara S, Kanari Y. 2008. Host genetic factors that control

immune responses to retrovirus infections. Vaccine 26:2981-96.

23. Persons DA, Paulson RF, Loyd MR, Herley MT, Bodner SM, Bernstein A, Correll PH,

Ney PA. 1999. Fv2 encodes a truncated form of the Stk receptor tyrosine kinase. Nat

Genet 23:159-65.

24. Takeda E, Tsuji-Kawahara S, Sakamoto M, Langlois MA, Neuberger MS, Rada C,

Miyazawa M. 2008. Mouse APOBEC3 restricts friend leukemia virus infection and

pathogenesis in vivo. J Virol 82:10998-1008.

25. Denzin LK, Khan AA, Virdis F, Wilks J, Kane M, Beilinson HA, Dikiy S, Case LK,

Roopenian D, Witkowski M, Chervonsky AV, Golovkina TV. 2017. Neutralizing Antibody

Responses to Viral Infections Are Linked to the Non-classical MHC Class II Gene H2-

Ob. Immunity 47:310-322 e7.

26. Case LK, Petell L, Yurkovetskiy L, Purdy A, Savage KJ, Golovkina TV. 2008. Replication

of beta- and gammaretroviruses is restricted in I/LnJ mice via the same genetic

mechanism. Journal of virology 82:1438-47.

27. Perlmutter RM, Hansburg D, Briles DE, Nicolotti RA, Davie JM. 1978. Subclass

restriction of murine anti-carbohydrate antibodies. J Immunol 121:566-72.

28. Rosenberg YJ, Chiller JM. 1979. Ability of antigen-specific helper cells to effect a class-

restricted increase in total Ig-secreting cells in spleens after immunization with the

antigen. J Exp Med 150:517-30.

29.   Coutelier JP, van der Logt JT, Heessen FW, Warnier G, Van Snick J. 1987. IgG2a

restriction of murine antibodies elicited by viral infections. J Exp Med 165:64-9.

30.   Nguyen TD, Bigaignon G, Van Broeck J, Vercammen M, Nguyen TN, Delmee M,

Turneer M, Wolf SF, Coutelier JP. 1998. Acute and chronic phases of Toxoplasma

gondii infection in mice modulate the host immune responses. Infect Immun 66:2991-5.

31.   Spinella S, Liegeard P, Hontebeyrie-Joskowicz M. 1992. Trypanosoma cruzi:

predominance of IgG2a in nonspecific humoral response during experimental Chagas'

disease. Exp Parasitol 74:46-56.

32.   Finkelman FD, Katona IM, Mosmann TR, Coffman RL. 1988. IFN-gamma regulates the

isotypes of Ig secreted during in vivo humoral immune responses. J Immunol 140:1022-

7.

33.   Stromnes IM, Dittmer U, Schumacher TN, Schepers K, Messer RJ, Evans LH, Peterson

KE, Race B, Hasenkrug KJ. 2002. Temporal effects of gamma interferon deficiency on

the course of Friend retrovirus infection in mice. J Virol 76:2225-32.

34.   Kane M, Deiss F, Chervonsky A, Golovkina TV. 2018. A Single Locus Controls

Interferon Gamma-Independent Antiretroviral Neutralizing Antibody Responses. J Virol

92.

35.   Simpson EM, Linder CC, Sargent EE, Davisson MT, Mobraaten LE, Sharp JJ. 1997.

Genetic variation among 129 substrains and its importance for targeted mutagenesis in

mice. Nat Genet 16:19-27.

36.   Threadgill DW, Yee D, Matin A, Nadeau JH, Magnuson T. 1997. Genealogy of the 129

inbred strains: 129/SvJ is a contaminated inbred strain. Mamm Genome 8:390-3.

37.   Jung YT, Kozak CA. 2000. A single amino acid change in the murine leukemia virus

capsid gene responsible for the Fv1(nr) phenotype. J Virol 74:5385-7.

38.   Stevens A, Bock M, Ellis S, LeTissier P, Bishop KN, Yap MW, Taylor W, Stoye JP. 2004.

Retroviral capsid determinants of Fv1 NB and NR tropism. J Virol 78:9592-8.

39. Santiago ML, Montano M, Benitez RL, Messer RJ, Yonemoto W, Chesebro B,

Hasenkrug KJ, Greene WC. 2008. Apobec3 Encodes Rfv3, a Gene Influencing

Neutralizing Antibody Control of Retrovirus Infection. Science 321:1343-1346.

40. Anonymous. 129S1/SvlmJ- Development. https://www.jax.org/strain/002448. Accessed

41. van den Broek MF, Müller U, Huang S, Aguet M, Zinkernagel RM. 1995. Antiviral

defense in mice lacking both alpha/beta and gamma interferon receptors. Journal of

virology 69:4792-4796.

42. Markine-Goriaynoff D, van der Logt JT, Truyens C, Nguyen TD, Heessen FW, Bigaignon

G, Carlier Y, Coutelier JP. 2000. IFN-gamma-independent IgG2a production in mice

infected with viruses and parasites. International immunology 12:223-230.

43. Liu C, Chikina M, Deshpande R, Menk AV, Wang T, Tabib T, Brunazzi EA, Vignali KM,

Sun M, Stolz DB, Lafyatis RA, Chen W, Delgoffe GM, Workman CJ, Wendell SG, Vignali

DAA. 2019. Treg Cells Promote the SREBP1-Dependent Metabolic Fitness of Tumor-

Promoting Macrophages via Repression of CD8(+) T Cell-Derived Interferon-gamma.

Immunity 51:381-397 e6.

44. Festing MF, Simpson EM, Davisson MT, Mobraaten LE. 1999. Revised nomenclature for

strain 129 mice. Mamm Genome 10:836.

45. Cantin E, Tanamachi B, Openshaw H, Mann J, Clarke K. 1999. Gamma interferon (IFN-

gamma) receptor null-mutant mice are more susceptible to herpes simplex virus type 1

infection than IFN-gamma ligand null-mutant mice. J Virol 73:5196-200.

46. Espejo C, Penkowa M, Saez-Torres I, Xaus J, Celada A, Montalban X, Martinez-

Caceres EM. 2001. Treatment with anti-interferon-gamma monoclonal antibodies

modifies experimental autoimmune encephalomyelitis in interferon-gamma receptor

knockout mice. Exp Neurol 172:460-8.

47. Lee EY, Schultz KL, Griffin DE. 2013. Mice deficient in interferon-gamma or interferon-

gamma receptor 1 have distinct inflammatory responses to acute viral

encephalomyelitis. PLoS One 8:e76412.

48. Huang S, Hendriks W, Althage A, Hemmi S, Bluethmann H, Kamijo R, Vilcek J,

Zinkernagel RM, Aguet M. 1993. Immune response in mice that lack the interferon-

gamma receptor. Science 259:1742-5.

49. Bae S, Park J, Kim JS. 2014. Cas-OFFinder: a fast and versatile algorithm that searches

for potential off-target sites of Cas9 RNA-guided endonucleases. Bioinformatics

30:1473-5.

Figure 1. Anti MLV-Ab production in 129S1 mice is controlled by a dominant genetic

mechanism.

A) BALB/cByJ, 129S1, or F_1 mice were infected with RL-MLV and monitored for IgG2a-

specific antibodies (left) or total Igs (right) against RL-MLV virion proteins by ELISA 8-10

weeks post infection. ns, not significant; *, $p < 0.05$; ***, $p < 0.001$

B) Spleen cells from RL-MLV infected mice were subjected to an infectious center assay 8-

10 weeks post infection.

C) Spleen weights of RL-MLV infected mice at 8-10 weeks post infection.

Figure 2. Anti MLV-Ab production in 129S1 mice is controlled by two non-MHC loci.

A) Diagram of breeding scheme used to determine Mendelian inheritance of the

mechanism controlling IFN -independent IgG2a Ab production.

B) N_2 mice generated from crossing (129S1xByJ)F_1 and ByJ mice as shown in A) were

infected with RL-MLV. Eight weeks post infection, their sera were tested for IgG2a Abs

against MLV virion proteins by ELISA (left), mice were classified as "resistant" if their

sera reacted with MLV virion proteins with an absorbance greater than 0.1 at OD 450

with an IgG2a-specific secondary. Susceptible and resistant mice were then tested for

presence of infectious virions in their spleens by XC plaque assay (center), and spleen

weights were measured (right).

C) MHC haplotypes of N_2 mice from B) were determined by FACS analysis and then levels

of IgG2a-specific antibodies (left) or total Igs (right) reactive against RL-MLV virion

proteins by ELISA were compared.

D) Presence of infectious virions in spleen (left) and spleen weights (right) of N_2 mice

grouped by MHC haplotype.

Figure 3. IFNγR-independent anti-MLV IgG2a Ab production in 129S7 mice

A) Mice of the indicated genotypes were infected with RL-MLV and monitored for IgG2a-specific antibodies (left) or total Igs (right) against RL-MLV virion proteins by ELISA 8-10 weeks post infection. ns, not significant; *, $p<0.05$; ***, $p<0.001$; ****, $p<0.0001$

B) Spleen cells from RL-MLV infected mice were subjected to an infectious center assay 8-10 weeks post infection.

C) Spleen weights of RL-MLV infected mice at 8-10 weeks post infection.

Figure 4. IFNγR-independent anti-MLV Ab production in 129S7 mice is a complex trait

A) IFNγR[-/-] BALB/cByJ, 129S7, and F_1 mice of the indicated genotypes were infected with RL-MLV and monitored for IgG2a-specific antibodies (left) or total Igs (right) against RL-MLV virion proteins by ELISA eight weeks post infection. ns, not significant

B) Spleen cells from RL-MLV infected mice were subjected to an infectious center assay eight weeks post infection.

C) Spleen weights of RL-MLV infected mice at eight weeks post infection.

Figure 5. Anti-MLV Ab responses in 129S1 mice are IFNγ-dependent

A) IFNγ-deficient and heterozygous control mice from two 129S1 founder lines were

infected with RL-MLV and monitored for IgG2a-specific antibodies (left) or total Igs

(right) against RL-MLV virion proteins by ELISA eight weeks post infection. ns, not

significant; **, p<0.01; ***, p<0.001

B) Spleen cells from RL-MLV infected mice were subjected to an infectious center assay

eight weeks post infection.

C) Spleen weights of RL-MLV infected mice at eight weeks post infection.

Figure 6. Genetic differences between 129S substrains affect IFNγ-independent anti-MLV Ab responses

A) Diagram of breeding scheme used to generate IFNγ, IFNγR, and IFNγ/IFNγR-deficient mice of mixed 129S7 and 129S1 genetic background.

B) F_2 mice of the indicated genotypes generated from intercrossing (129S7-*Ifngr1*[-/-] x 129S1-*Ifng*[-/-])F_1 mice as shown in A) were infected with RL-MLV. 8-10 weeks post infection, their sera were tested for IgG2a-specific antibodies (left) or total Igs (right) against RL-MLV virion proteins by ELISA. +/·, either heterozygous or homozygous wild type. ns, not significant; *, p<0.05; **, p<0.01; ***, p<0.001; ****, p<0.0001

C) Spleen cells from RL-MLV infected F_2 mice were subjected to an infectious center assay eight weeks post infection.

D) Spleen weights of RL-MLV infected F_2 mice at eight weeks post infection.

558 **Table 1.** Summary of mice utilized in this investigation and their phenotype upon MLV infection

Mice	Genotype		MLV infections		
	IFNγ	IFNγR	# Infected	# with ≤ 100PFU/10^6cells	% Cleared
BALB/cByJ	+/+	+/+	5	1	20%
BALB/cByJ	+/+	-/-	12	0	0%
129S1/SvImJ (129S1)	+/+	+/+	24	20	83%
129S1	+/-	+/+	9	8	89%
129S1	-/-	+/+	28	5	18%
(129S1xByJ)F$_1$	+/+	+/+	18	14	78%
(ByJx129S1)F$_1$	+/+	+/+	15	7	47%
([129S1xByJ)xByJ]N$_2$ resistant	+/+	+/+	9	8	89%
([129S1xByJ)xByJ]N$_2$ susceptible	+/+	+/+	31	0	0%
([129S1xByJ)xByJ]N$_2$ H2^d	+/+	+/+	2	19	11%
([129S1xByJ)xByJ]N$_2$ H2$^{d/b}$	+/+	+/+	6	21	29%
129S7	+/+	-/-	20	12	60%
(129S7xByJ)F$_1$	+/+	-/-	30	11	37%
(ByJx129S7)F$_1$	+/+	-/-	38	8	21%
(129S7x129S1)F$_2$	+/·	+/·	34	30	88%
(129S7x129S1)F$_2$	-/-	+/·	22	14	64%
(129S7x129S1)F$_2$	+/·	-/-	21	2	10%
(129S7x129S1)F$_2$	-/-	-/-	4	4	100%

559   +/· - either heterozygous wild-type/knock-out or homozygous wild-type