Cryo-ET of infected cells reveals that a succession of two lattices drives vaccinia virus assembly

Miguel Hernandez-Gonzalez[1], Thomas Calcraft[2], Andrea Nans[3], Peter B Rosenthal[2] and Michael Way[1, 4] *

[1]Cellular signalling and cytoskeletal function laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, UK

[2]Structural Biology of Cells and Viruses Laboratory, The Francis Crick Institute.

[3]Structural Biology Science Technology Platform, The Francis Crick Institute.

[4] Department of Infectious Disease, Imperial College, London W2 1PG, UK

*To whom correspondence should be addressed. Email: michael.way@crick.ac.uk

Running title: Cryo-ET of vaccinia infected cells

Abstract

During replication, vaccinia virus and other poxviruses such as monkeypox first assemble non-infectious spherical immature virions (IV) in the cytoplasm of infected cells. The IV membrane, which is fully coated by a viral D13 lattice, encapsidates both viral proteins and the DNA genome. Subsequently, the D13 lattice is lost and IV transitions into the brick-shaped intracellular mature virus (IMV) with an organised core enclosing the genome. We have an overview of the maturation of IV into IMV but our structural understanding remains limited. Here, we performed cryo-electron tomography of frozen-hydrated vaccinia-infected cells to structurally characterise the maturation process in situ. We found that D13-coated IV have similar diameters and lattice dimensions to spherical D13 assemblies formed in vitro, confirming the size of IV is determined by D13. During IMV formation a new pseudohexagonal lattice forms inside IV to produce the viral core. This lattice is composed of trimeric pillars with outward projections that appears as a palisade in cross-section. Strikingly, the diameter of IV precisely matches the IMV long axis, suggesting the length of the core, and hence the IMV, is determined by the D13 lattice. During maturation the viral membrane becomes corrugated as it adapts to the shape of the core. This suggests that the 50% reduction in volume during the transition of IV to IMV does not require membrane removal. Our observations suggest that the consecutive D13 and palisade lattices structurally define vaccinia assembly and maturation.

Significance statement

Poxviruses such as variola virus (smallpox) and monkeypox induce disease in humans. Other poxviruses, including vaccinia virus and modified vaccinia Ankara (MVA), are used as vaccine vectors. Despite their importance, a structural understanding of how infectious poxvirus virions assemble in the cytoplasm of infected cells is lacking. In this study, we used cryo-electron tomography of frozen-hydrated vaccinia infected cells to structurally characterise the maturation of spherical immature virions into the infectious brick-shaped intracellular mature virus. These high-resolution in situ observations reveal new features concerning the viral membrane and the inner core of mature virions. The model that emerges suggests that two consecutive viral-derived protein lattices drive vaccinia assembly and maturation.

Introduction

Poxviruses are large double stranded DNA viruses that replicate and assemble their virions in cytoplasmic perinuclear viral factories (1). The family includes variola virus, the causative agent of smallpox, and monkeypox as well as vaccinia, the most studied family member which was used as the vaccine to eradicate smallpox (2, 3). During its replication cycle, vaccinia initially assembles into infectious intracellular mature virions (IMV), which are released when infected cells lyse (4). Prior to cell lysis, however, some IMV become enveloped by a Golgi cisterna or endosomal compartment to form intracellular enveloped virions (IEV) (5-7) (Figure 1A). IEV undergo kinesin-1 mediated microtubule transport to the plasma membrane, where they fuse to release extracellular enveloped virions (EEV) that promote the long-range spread of infection (8, 9). Alternatively, after fusion some virions (cell-associated enveloped virus, CEV) remain attached to the cell to stimulate actin-dependent cell-to-cell spread of the virus (10-13).

The first step in the spread of vaccinia infection is the assembly of IMV, which are the precusor to all subsequent forms (IEV, EEV and CEV) of the virus. IMV formation is initiated when endoplasmic reticulum (ER) associated viral membrane assembly proteins (VMAPs) promote the formation of isolated membrane crescents in the cytoplasm of infected cells (14-16). The characteristic shape of the crescents is determined and maintained by the D13 scaffolding viral protein, which forms trimers that assemble into a hexameric lattice coating the outer surface of the membrane (14, 15, 17, 18). In vitro structural analysis of recombinant D13 reveals it has a double-jelly-roll structure composed of 8 antiparallel ß-strands that self assembles into a honeycomb lattice of pseudo-hexagonal trimers (19-22). Ultimately, membrane crescents develop into spherical immature virions (IV) which encapsidate the viral genome and proteins required to produce an IMV (15, 17, 18, 23). In a process that is still not understood but involves the proteolytic cleavage of viral proteins and the formation of disulphide bonds, spherical IV undergo a dramatic reorganization into brick-shaped IMV (24, 25). During this maturation process, the D13 honeycomb lattice is lost, and a biconcave viral core assembles around the viral genome (23). The outer layer of the viral core forms a paracrystalline palisade-like structure that has been reported to be discontinuous (23, 26, 27). In addition, two dense proteinaceous aggregates, known as the lateral bodies, assemble in the cavity between the outer viral membrane and the biconcave viral core (23, 26, 28).

Vaccinia virion assembly has been extensively studied. Nevertheless, the key structural determinants that govern the changes occurring during the IV to IMV transistion as well as

those that define the IMV remain unknown. To obtain detailed ultra-structural insights into

this maturation process, we have now performed cryo-electron tomography (cryo-ET) of

vaccinia infected cells. Our analysis reveals that a succession of two different viral lattices

drives assembly of IMV: D13 shapes the IV membrane, which subsequently coalesces onto

the pseudohexagonal palisade lattice, which is reminiscent of a viral capsid.

Results

Electron cryo-tomography of vaccinia virus in infected cells

Cryo-tomograms of thin cellular regions containing IV and IMV were recorded from the

periphery of HeLa cells infected with vaccinia virus for 8 hours (Figure 1B, C and S1, Movie

1 and 2). IV, fully coated by the viral D13 lattice, are largely spherical and have a diameter

of 351.89 ± 2.88 nm (Table S1). Analysis of tomograms in 3D, however, reveals that a

proportion of IV despite being closed and fully coated by D13 have a single invagination of

the viral membrane (Figure 1B). This suggests that the D13 lattice is flexible and can

accommodate both positive and negative curvatures present on virions with invaginations.

This D13 lattice flexibility is also evident from its ability to coat the exterior of "open

spheres" with different membrane curvature (also known as crescents due to their shape

when seen in a middle view) during IV formation (Figure 2A, B and Movie 3).

The interior of the majority of IV, which contain viral DNA and proteins required for virion

assembly, are homogenous and lack any of the distinctive regions or layers seen in IMV

(Figure 1B). In a few cases, however, we could observe a single dense striated structure

inside IV corresponding to the nucleoid (viral DNA genome) before it decondenses (Figure

2C and S2), as seen in previous ultra-structural analysis of thin EM sections of vaccinia

infected cells (28). We also found examples of related striated structures in the cytoplasm

adjacent to, or in direct association with, assembling IV (Figure 2D, E). Consistent with the

interpretation that the example shown in Figure 2E represents the viral genome being

inserted into assembling IV, structured illumination microscopy together with deconvolution

(125 nm resolution) also reveals the presence of elongated DNA structures associating with

virions, identified by RFP-A3, in the viral factory (Figure 2F). Our 3D tomographic

observations of assembling IV using cryo-ET were consistent with previous ultra-structural

analysis of vaccinia infected cells using electron microscopy (16-18, 29, 30).

In situ structure of the hexagonal D13 lattice

Structural analysis of recombinant D13 in isolation has provided a detailed molecular

understanding of the D13 trimer (19-22). However, the macromolecular arrangements of

the D13 lattice on IV in infected cells have largely been studied by deep-etch EM (17, 18).

Given the clear lattices present on the IV surface in our tomograms (Figure 3A, Movie 4), we

analysed the structure of the D13 lattice in situ by subtomogram averaging. We obtained

maps by averaging surface sectors revealing hexamers of trimer arrangements with a lattice

spacing of 133 Å, in agreement with lattices described for in vitro assembled D13 and deep-

etch EM honeycombs (Figure 3B). Rigid-body docking of the high-resolution structure of a

D13 trimer (PDB: 7VFE) into the map shows the packing of D13 trimers is consistent with

that recently reported (19). While at much lower resolution, precluding a detailed

description of interface residues, the arrangement of D13 on the spherical IV surface does

not appear to include a relative twist of trimeric axes between adjacent trimers as observed

in single particle dimers or tubular assembles.

From the hexagonal packing of D13 in the subtomogram average, we estimate that ~5,300

D13 trimers will cover the surface of a spherical IV. The packing density and radius of the

D13 lattice on IV are compatible with an icosahedrally symmetric architecture with a

triangulation number (T) in the neighborhood of 268. We have not confirmed the presence

of icosahedral symmetry, which would require pentameric arrangements of D13 trimers,

such as those that have been identified in vitro (19). Finally, we observed a gap of ~5.6 nm

between the innermost surface of the D13 lattice and the membrane of the IV. This gap

cannot be explained by any unresolved N- or C-terminal residues of D13 and must therefore

be bridged by another protein such as A17 (21).

Structure of intracellular mature virus in infected cells

The internal organisation, dimensions and shape of IMV, which are "brick-shaped"

membrane-bound virions, contrast dramatically with those of IV (Figure 1). Analysis of the

size and shape of IMV in 3D reveals they can be approximated to a triaxial ellipsoid of 352 x

281 x 198 nm with a calculated volume of 1.02×10^7 nm^3 (Figure 4, Table S1 & Movie 5).

The outer membrane of IMV has a corrugated appearance and the D13 lattice is absent.

There is, however, a new additional outer layer of 6.67 ± 0.11 nm on the viral membrane

that was not present on IV (Figure 4). Underneath the viral membrane there is a palisade-

looking structure consisting of a series of turret-like densities in cross-section (12.5 nm

thick) that is associated with a inner wall that is 3.84 ± 0.09 nm thick (Figure 4A). Depending

on the orientation of the virion, the interior organisation of the IMV appears very different.

In the mid view of its two widest dimensions (352 x 281 nm), the palisade appears in contact

with the viral membrane. However, in the orthogonal lateral view (352 x 198 nm) the

palisade structure, which only contacts the viral membrane at the ends of the virion (Figure

4B), forms a dumbbell shape that results in the formation of two concavities. Each cavity
contains a single dense and amorphous structure corresponding to a lateral body. The space
between the viral membrane and the palisade varies between 9.28 ± 0.18 nm and 46.53 ±
2.72 nm, the latter being where the lateral bodies are accommodated (see Table S1 for
detailed measurements). 3D tomographic analyses reveal that the palisade is a semi-regular
array when viewed from above (Figure 4C, D).
184
The semi-regular organisation of the palisade is most apparent in naked cores, which lack
the viral membrane and are occasionally found in the cytoplasm (Figure 5A, B).
Segmentation of our 3D tomograms reveals the palisade is a continuous structure without
fenestrations that defines the virus core, including the regions that contact the lateral
bodies (Figure 4E, Movie 6). Inside the core, there are no obvious higher order structures.
There are, however, interconnected densities that vary between virions but tend to
accumulate beneath the inner wall of the core (Figure 4A). These densities are especially
apparent in the compressed region underneath the lateral bodies (Figure 4C). Such densities
associated with the inner wall of the core potentially represent the viral genome and its
associated proteins given its high contrast. Another characteristic that is evident in
midplane views of approximately half of the IMV (48.9%, n = 94), is that one corner of the
virion (and in a few cases, two corners) appears as a straight or flattened "cut corner"
(Figure 1C and Figure S3). This characteristic is intrinsic to the core wall as it is also apparent
in naked cores (Figure 5A, Movie 7) and has been previously observed in purified cores (27).
In addition, these naked cores lack the two densities corresponding to lateral bodies but
have associated spaghetti-like structures contacting the palisade surface (Figure 5A). These
flexible polymers, which are 2.6 nm in diameter, form an exclusion zone of ~ 40 nm around
the naked core (Figure 5A). Furthermore, it was noticeable that naked cores have randomly
distributed ring like structures on their surface that were not observed in IMV. These rings
seem to protrude 10-20 nm from the palisade surface and had inner and outer diameters of
4.5 ± 0.1 nm and 9.9 ± 0.1 nm, respectively (Figure 5C).
206
The distinct hexagonal core lattice of infectious vaccinia virions
During viral egress, some IMV become triple-membraned intracellular enveloped virions
(IEV) after envelopment by a Golgi cisterna or endosomal compartment (Figure 1A).
Subsequent fusion of the IEV with the plasma membrane liberates double-membraned
extracellular enveloped virions (EEV) that are known as cell-associated extracellular
enveloped virions (CEV) if they remain attached to the outside of the cell (7).   Our
tomograms reveal that the additional membranes acquired by envelopment are not

corrugated (Figure 6A). In the case of IEV, the outer membrane is also not always in close contact with the underlying membrane (see Table S1 for virion measurements). The dimensions and structure of the IMV are also unaffected by envelopment (Figure 6A). Notably, the palisade fully coats the viral core in all infectious virions and its organisation appears unaltered (Figure 6A-C). To study the architecture of the palisade, we performed subtomogram averaging separately using IMV, IEV and CEV/EEV particles. Maps obtained from the different virion types, all displayed the same organisation and lattice parameters (P6 symmetry with a=b=89 ± 2 Å, θ=120°) (Fig S4). A new combined map obtained by averaging all particle types, reveals that the palisade is composed of trimeric pillars with projecting lobes that interact with neighbouring pillars with local hexagonal symmetry (Figure 6D). These pillars are embedded in an unfeatured inner wall. While further details are required to understand its molecular composition, this arrangement appears to be flexible enough to assemble a continuous biconcave capsid structure.

Maturation of IV to IMV

Radical changes in the organisation, dimensions and shape of IV result in the formation of IMV (Figure 7). Based on our tomograms, IV have an average diameter of 351.89 ± 2.88 nm and a volume of 2.28×10^7 nm^3 for completely spherical particles (Figure 7A). In contrast, "brick-shaped" IMV have a volume of 1.02×10^7 nm^3 based on dimensions of 352 x 281 x 198 nm (Figure 7A). This reduction in volume is accompanied by a dramatic corrugation of the viral membrane, together with the loss of the D13 lattice. To better understand this corrugation process, we measured the middle-plane perimeter of the viral membrane in IV and IMV, following the wrinkles of the viral membrane to obtain its contour length. We found that the membrane contour is virtually identical in IV and IMV, as well as in the equivalent innermost mmembrane of EEV/CEV (~1100 nm, Figure 7B). This suggests the viral membrane folds during maturation, which would explain the reduction in volume without any detectable loss of membrane surface. In addition to reducing their volume by ~50%, IVs also change their shape when they mature into IMV, becoming a triaxial ellipsoid. Strikingly, the longest IMV dimension matches the diameter of IV (Figure 7A, B), suggesting that the major axis of IMV is dictated by the IV diameter. In our tomograms we also found particles that may represent intermediates and/or defective examples of IV maturation. These include examples where the palisade is fully formed but the IV membrane, which either lacks or is partially coated with D13, is not associated with the viral core (Figure 7C). It is also interesting that we did not observe any virions with partially formed cores suggesting that palisade formation is likely to be rapid or occurs en bloc.

Discussion

Previous ultra-structural analyses of vaccinia infected cells by electron microscopy over many decades have provided important insights into the assembly of vaccinia virions. These studies, however, have largely been conducted on fixed sections of infected cells which can have fixation, processing and staining artifacts. Cryo-EM studies, analysing viral entry or purified virions, have demonstrated the potential of using cryo-electron microscopy to study vaccinia structure (26, 27, 31, 32). Here, we apply cryo-ET to image the thin edge of plunge-frozen, vaccinia virus-infected cells, revealing virus architecture in situ. While there is confirmation of past work, important new features emerged from our study. We found that during maturation to IMV, spherical IV lose their D13-coat and reduce their volume by ~50%. During this process, the outer viral membrane becomes corrugated when it associates with the capsid-like palisade layer and the lateral bodies. The palisade, which remains unaltered during subsequent virion morphogenesis, is a continuous regular lattice with pseudohexagonal symmetry that fully covers the viral core. Our observations suggest that these two lattices, D13 and the palisade, drive vaccinia assembly and maturation.

We found that the D13 lattices coating IV have a similar architecture to the D13 lattices formed in vitro (19). Furthermore, Hyun et al. produced D13 spherical IV-like particles with a similar diameter to the D13-coated IV we imaged in situ. This strongly suggests that the D13 lattice, which initially curves open membranes and eventually forms closed particles, determines both the size and shape of the IV. In addition, our images and STA maps show that a 5.6-nm gap separates the D13 lattice from the IV membrane. This gap is most likely occupied by A17, which tethers the D13 lattice to the membrane (21, 33, 34) but whose flexibility would preclude its detection on the IV membrane by cryo-ET. The lack of direct contact of the D13 protein lattice with the viral membrane is reminiscent of clathrin and other vesicle-forming coats in eukaryotic cells where the structural lattice does not directly interact with the membrane to be deformed (35, 36). Instead, an adaptor module mediates the lattice-membrane association. A mechanism by which coat complexes promote membrane curvature is by clustering the adaptor module on the membrane to be deformed (37, 38). Therefore, D13-mediated clustering of A17 could promote membrane deformation during IV formation. In addition, A17 can also contribute to membrane bending independently of D13, since it can deform membranes in vitro and when expressed in non-infected cells (39).

Despite their relatively constant diameter, a proportion of D13-coated IV have an elongated membrane invagination (Figure 1A). This feature has largely gone unnoticed in previous EM

studies imaging thin sections, although IV membrane deformations are apparent in some
studies (28, 40-43). While the cause of this membrane buckling remains to be established,
it may relate to the thinness of the cell periphery or a consequence of forces on the
membrane during IV assembly or DNA insertion into assembling IV (42). As in other studies,
we observe the nucleoid in the IV interior or associated with either open or closed spheres,
as previously suggested for DNA entry events (28, 30). It is tempting to speculate that the
IV invagination may mark the site of DNA entry and IV sealing of the pore.

The palisade, which is largely composed of p4A and A4 (31, 44, 45), consists of a trimeric
assembly in a pseudo-hexagonal lattice arrangement. This new viral lattice, which fully
covers the core and defines its boundary, also dictates the dimensions and shape of IMV.
The viral membrane, no longer covered by D13, wrinkles and adopts the shape and
dimensions of the newly formed palisade. This process may also be responsible for the
compression of the lateral bodies onto the virus core, causing the biconcave deformation of
the latter. The maturation model that emerges from our work provides a simple way by
which a membrane-bound particle adapts to a new internal lattice and changes its shape
and dimensions to become more compact without membrane removal (Figure 7D).
Moreover, our data suggest the palisade length is dictated by the diameter of the IV, which
in turn is established by D13. The palisade would grow inside the D13-coated IV until
reaching its maximum length, that is the IV diameter. In other words, the D13 lattice limits
the longest dimension of the palisade lattice. The size restriction of a growing viral core by
its surrounding viral membrane has also been proposed for the assembly of the HIV core (46,
47). Our observation of a viral particle partly coated by D13 with an assembled palisade
suggests removal of D13 only occurs once the palisade is fully formed (Figure 7C). Moreover,
our observations suggest that palisade assembly and D13 removal are coordinated and rapid.

In the IMV, the lateral bodies define a lateral domain, as opposed to the virion tips, which
are specified by the long axis of the viral core. The lateral bodies prevent direct interactions
between the palisade and the wrinkled viral membrane, which may contribute to the
differences between the lateral and the tip domains. Notably, the virion tips are the site of
polarisation of the set of viral membrane proteins that form the entry fusion complex (EFC),
which is essential for virus infectivity (48-50). How and when EFC polarity is established is
unknown. An appealing possibility is that virion tips are stochastically defined by the growth
of the palisade long axis during assembly and the EFC and other viral factors polarise at the
virion tips after palisade formation by their exclusion from the lateral domains. In such a
scenario, palisade formation would drive virion polarity.

The EFC mediates fusion of the IMV membrane with the plasma membrane or an endocytic
compartment during entry, which releases a naked core into the cytoplasm (51-55).
Subsequently, early proteins are produced by early RNA transcripts released from naked
cores, which are essential to liberate the DNA genome into the cytosol to initiate replication
(1). Our cryo-ET data of naked cores in infected cells indicate the palisade structure does
not require surrounding membranes for stability. Furthermore, the loss of lateral bodies
from naked cores suggests their association with the palisade depends on the IMV
membrane. In addition, we observed flexible 2.6-nm thick polymers fully surrounding naked
cores that might correspond to either RNA or the viral genome which is released from the
naked core. Finally, we detected ring structures protruding from the palisade surface of
naked cores. Analysis of these rings reveals they are compatible in dimension with the
hexameric rings of the viral D5 primase/helicase, which is essential for vaccinia genome
release (56, 57).

A common feature of many viral families is the use of a three-dimensional lattice, or capsid,
which determines the structure and dimensions of the virion. What distinguishes vaccinia
from other viruses is that an initial lattice, D13, is replaced by a second lattice, the palisade,
during virion assembly. This contrasts African Swine Fever virus (ASFV), which retains its
D13-like lattice, in addition to an inner capsid in mature virions (58). Based on the structural
similarities between D13 and the structural components of other viruses with a pseudo-
hexameric structure formed by a trimeric protein containing concatenated beta-barrels,
vaccinia has been included in the PRD1/adenovirus lineage (59). However, the viral proteins
that form the palisade, which is present in the mature particles, may be just as important
as D13 in structural phylogenetic comparisons.

The palisade is a regular capsid-like structure determining the morphology of all forms of
vaccinia virions (IMV, IEV, EEV and CEV). The additional membranes acquired by
envelopment are different to the IMV viral membrane as they are not corrugated or always
in close contact with the IMV surface. Moreover, the acquisition of these membranes is not
associated with major changes in the virion structure or palisade architecture (Figure 5).
This suggests that these additional membranes are used to facilitate viral egress before cell
lysis. In fact, these membranes and associated proteins drive IEV transport on microtubules
to the cell periphery, IEV fusion with the plasma membrane and subsequent actin-based
transport of CEV (7-11, 13). In contrast, the IMV membrane serves to contain the viral core
and lateral bodies and organises the essential components required for entry and, also,
envelopment during viral morphogenesis. Based on the strong protein conservation between

orthopoxviruses, we believe that the virion assembly of monkeypox and variola major are likely to be identical to that of vaccinia virus we have described here.

Methods

Recombinant viruses

To facilitate cryo-electron microscopy (EM) observations, we produced recombinant Western Reserve (WR) vaccinia strains lacking F11 as this prevents infected cells from rounding-up early during infection (60). In addition, the A36-YdF mutation was used to abolish actin tail formation beneath CEV (11, 13, 61). The F11 gene was deleted in A36 YdF (8) or A36 YdF ΔNPF1-3 (62) backgrounds using the same targeting strategy as previously described (63). Fluorescence was used as a selectable marker to isolate recombinant ΔF11-mCherry viruses by successive rounds of plaque purification in BS-C-1 cells. Correct gene replacement and complete isolation of the recombinant viral strains was confirmed by PCR, sequencing of the F11 locus and western blot analysis.

Cell growth, vaccinia infection and vitrification

HeLa cells were maintained in complete MEM (supplemented with 10% Fetal Bovine Serum, 100 ug/ml streptomycin and 100 U/ml penicillin) at 37 °C with 5% CO_2. Cells were washed with PBS, treated with trypsin, seeded on glow discharged (40 sec at 45 mA) Quantifoil R3.5/1 gold grids of 200 mesh and placed in wells of 6-well plates with complete MEM. After overnight growth, the cells were infected with A36 YdF ΔF11 or A36 YdF ΔNPF1-3 ΔF11 vaccinia virus strains in serum-free MEM at a multiplicity of infection of 2. After one hour, the medium was replaced with complete MEM. At 8 hours post infection the grids were washed once with PBS and excess PBS was removed with a Whatman paper before blotting and vitrification using the Vitrobot Mark IV System, which was set to 95% relative humidity at 22 °C. Colloidal gold particles (10-nm diameter) were pipetted onto grids before blotting, which was performed for 14 s with a relative force of -10.

Cryo-electron tomography, image processing and analyses

Grids were first screened on a Talos Arctica TEM (Thermo Fisher) to select edges of cells with an abundance of viral particles that were thin enough for cryo-ET. The precise location of the selected regions on each grid was registered. The grids were subsequently transferred to a Titan Krios (Thermo Fisher) and the mapped grid selections were re-imaged. The Titan Krios was fitted with a K2 Summit direct detector (Gatan) operated in electron counting mode. A Gatan GIF energy filter was used in zero-loss mode with a 20 eV slit width. Dose-symmetric tilt series were collected from -57° to +57° at a 3° increment, a pixel size of

4.31 Å and a defocus of -8 μm using Tomography 5.7 software (Thermo Fisher) (Table S2).

Four movie frames were collected per tilt with a dose of 1.7 e / Å² per tilt, giving a

cumulative dose of 66.3 e / Å² per tilt series. Movie frames were aligned using alignframes

from IMOD (64). Tilt series were aligned using 10-nm Au fiducials or patch tracking in IMOD.

Contrast transfer function (CTF) correction and tomogram reconstruction were also

implemented in IMOD (64, 65). Alternatively, tomograms were CTF corrected and

reconstructed using novaCTF (66). In both cases, a SIRT-like filter equivalent to 5 iterations

was routinely applied. Tomogram segmentations were manually performed using AMIRA

(Thermo Fisher), mainly using tomograms that were further denoised using ISONet with CTF

deconvolution and missing-wedge filling from deep learning (67).

The diameter of IV, the axes of IMV and their perimeters, and the major axes of IEV and EEV

were measured in 3dmod from IMOD using a middle plane of each viral particle. For

perimeters, the viral membranes were manually traced to measure their total contour

length. As reported previously (23), virions with their dumbbell side view perpendicular to

the electron beam were much less frequent than other views. Because of this, for the

calculation of the minor axis of IMV we measured the minor axis of both IMV and IMV inside

EEV. This is also the reason why we did not find enough side views of IEV to directly measure

their minor axis (Table S1). To estimate the IEV minor axis, we measured the thickness that

the additional IEV membranes add, and combined it to the calculated IMV minor axis. For

volumes calculation, we assumed IV are perfectly spherical and IMV, IEV and EEV triaxial

elipsoids.

Subtomogram averaging

Tomograms containing IV or IMV were selected for subtomogram averaging (STA). D13 and

palisade particles in immature and mature virions respectively were picked from the bin4

SIRT-filtered tomograms using oversampled surface models in Dynamo (68), then imported

to Relion 3.1 (69) for subsequent subtomogram averaging. Unless otherwise specified, local

angular searches about the initial orientations generated in Dynamo were used throughout

data processing. For particle extraction, only unbinned weighted back projections

reconstructions from novaCTF were used. Lattice measurements were carried out in real

space using ChimeraX (70). D13 particles were extracted in 96-pixel boxes at the bin2 pixel

size of 4.31Å/pixel. An initial reference was generated using reference-free 3D classification

without applied symmetry, with the resulting map clearly showing the honeycomb lattice

of D13 trimers. Initial classification and refinement was carried out in both C1 and C6

symmetries to verify the symmetry of the D13 hexamer-of-trimers. After initial

classifications and removing of overlapping particles with a distance cutoff of 50Å, particles

were re-extracted without binning (4.31Å/pixel) in 192 pixel boxes. Final refinement in C6

symmetry reached 19Å resolution according to the FSC=0.143 criterion.

Palisade particles were initially extracted at binning level 2 (8.62Å/pixel) in 96-pixel boxes.

Initial references were generated using independent reference-free 3D classifications for

each virion type. This clearly showed the palisade lattice in every case, and the particles

for each virion type were from then on processed independently, but in an identical manner.

For each virion type, alternating rounds of 3D classification and refinement were used to

centre the palisade particles at 2-fold binning. After convergence at 2-fold binning, particles

were re-extracted without binning (4.31Å/pixel) in 128-pixel boxes. Overlapping particles

were removed with a distance cutoff of 40Å before further refinements were carried out

without binning. As all refinements converged on maps with identical lattices, particles from

all virion types except for the naked cores were then combined for further refinements. All

refinements were carried out without application of symmetry up to this point. After the

C1 maps converged and the symmetry of the lattice was apparent, further refinement runs

were carried out with C3 symmetry applied. The final resolutions of the palisade maps

ranged from 20-30Å, according to the FSC=0.143 criterion.

Immunofluorescence imaging

HeLa cells were plated on fibronectin-coated coverslips and infected with an RFP-A3

vaccinia strain (71) for 8 hours and fixed with 4% paraformaldehyde in PBS for 10 min, then

permeabilised with 0.1% Triton X-100 in PBS for 5 min and incubated in blocking buffer (10

mM MES pH 6.1, 150 mM NaCl, 5 mM EGTA, 5 mM $MgCl_2$, and 5 mM glucose) containing 2%

(v/v) fetal calf serum and 1% (w/v) BSA for 30 min prior to addition of 4′,6-diamidino-2-

phenylindole (DAPI) for 5 minutes, before mounting the coverslips using Mowiol. Coverslips

were imaged on an Olympus iX83 Microscope with Olympus 150x/1.45 NA X-Line

Apochromatic Objective Lens, dual Photometrics BSI-Express sCMOS cameras and CoolLED

pE-300 Light Source (Visitech) and was controlled using Micro-Manager 2.0.0. Image stacks

of 10-15 z-slices with 0.1 μm steps were acquired and deconvolved using the express

deconvolution setting on Huygens Software (Scientific Volume Imaging).

Acknowledgements

We thank Alessandro Costa, Neil McDonald and Jeremy Carlton (Francis Crick Institute) as

well as the Way lab for insightful comments on the manuscript as well as LuYan Cao for

providing the initial slide of infected cells used in our SIM analysis (Fig. 2F). We also

acknowledge Philip Walker and Andy Purkiss of the Structural Biology Science Technology Platform and the Scientific Computing Science Technology Platform for computational support. MHG also thanks Pauline McIntosh (Francis Crick Institute) for guidance during the initial stages of the project. The work in the Way and Rosenthal labs was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001209 and FC001143), the UK Medical Research Council (FC001209 and FC001143), and the Wellcome Trust (FC001209 and FC001143). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

Author contributions

All authors designed the research program and strategy. MHG, TC and AN performed research, MHG and TC analyzed data and MHG, TC, PBR and MW wrote the paper.

Competing interests

The authors have no competing financial interests

References

1. B. Moss, "Poxviridae: the viruses and their replication" in Fields Virology, H. P. Knipe DM, Ed. (Lippincott Williams and Wilkins, Philadelphia, 2007), pp. 2905-2946.

2. B. L. Jacobs *et al.*, Vaccinia virus vaccines: past, present and future. *Antiviral Res* **84**, 1-13 (2009).

3. S. R. Walsh, R. Dolin, Vaccinia viruses: vaccines against smallpox and vectors against infectious diseases and tumors. *Expert Rev Vaccines* **10**, 1221-1240 (2011).

4. K. L. Roberts, G. L. Smith, Vaccinia virus morphogenesis and dissemination. *Trends Microbiol* **16**, 472-479 (2008).

5. M. Schmelz *et al.*, Assembly of vaccinia virus: the second wrapping cisterna is derived from the trans Golgi network. *J Virol* **68**, 130-147 (1994).

6. J. Tooze, M. Hollinshead, B. Reis, K. Radsak, H. Kern, Progeny vaccinia and human cytomegalovirus particles utilize early endosomal cisternae for their envelopes. *Eur J Cell Biol* **60**, 163-178 (1993).

7. G. L. Smith, A. Vanderplasschen, M. Law, The formation and function of extracellular enveloped vaccinia virus. *J Gen Virol* **83**, 2915-2931 (2002).

8. J. Rietdorf *et al.*, Kinesin-dependent movement on microtubules precedes actin-based motility of vaccinia virus. *Nat Cell Biol* **3**, 992-1000 (2001).

9.   F. Leite, M. Way, The role of signalling and the cytoskeleton during Vaccinia Virus
egress. *Virus Res* **209**, 87-99 (2015).

10.   S. Cudmore, P. Cossart, G. Griffiths, M. Way, Actin-based motility of vaccinia virus.
Nature **378**, 636-638 (1995).

11.   F. Frischknecht *et al.*, Actin-based motility of vaccinia virus mimics receptor tyrosine
kinase signalling. *Nature* **401**, 926-929 (1999).

12.   M. Hollinshead *et al.*, Vaccinia virus utilizes microtubules for movement to the cell
surface. *J Cell Biol* **154**, 389-402 (2001).

13.   B. M. Ward, B. Moss, Vaccinia virus intracellular movement is associated with
microtubules and independent of actin tails. *J Virol* **75**, 11651-11663 (2001).

14.   A. S. Weisberg *et al.*, Enigmatic origin of the poxvirus membrane from the
endoplasmic reticulum shown by 3D imaging of vaccinia virus assembly mutants. *Proc Natl*
Acad Sci U S A **114**, E11001-E11009 (2017).

15.   B. Moss, Poxvirus membrane biogenesis. *Virology* **479-480**, 619-626 (2015).

16.   P. Chlanda, M. A. Carbajal, M. Cyrklaff, G. Griffiths, J. Krijnse-Locker, Membrane
rupture generates single open membrane sheets during vaccinia virus assembly. *Cell Host*
Microbe **6**, 81-90 (2009).

17.   J. Heuser, Deep-etch EM reveals that the early poxvirus envelope is a single
membrane bilayer stabilized by a geodetic "honeycomb" surface coat. *J Cell Biol* **169**, 269-
283 (2005).

18.   P. Szajner, A. S. Weisberg, J. Lebowitz, J. Heuser, B. Moss, External scaffold of
spherical immature poxvirus particles is made of protein trimers, forming a honeycomb
lattice. *J Cell Biol* **170**, 971-981 (2005).

19.   J. Hyun, H. Matsunami, T. G. Kim, M. Wolf, Assembly mechanism of the pleomorphic
immature poxvirus scaffold. *Nat Commun* **13**, 1704 (2022).

20.   M. W. Bahar, S. C. Graham, D. I. Stuart, J. M. Grimes, Insights into the evolution of
a complex virus from the crystal structure of vaccinia virus D13. *Structure* **19**, 1011-1020
(2011).

21.   D. Garriga *et al.*, Structural basis for the inhibition of poxvirus assembly by the
antibiotic rifampicin. *Proc Natl Acad Sci U S A* **115**, 8424-8429 (2018).

22.   J. K. Hyun *et al.*, Membrane remodeling by the double-barrel scaffolding protein of
poxvirus. *PLoS Pathog* **7**, e1002239 (2011).

23.   R. C. Condit, N. Moussatche, P. Traktman, In a nutshell: structure and assembly of
the vaccinia virion. *Adv Virus Res* **66**, 31-124 (2006).

24.   C. Ansarah-Sobrinho, B. Moss, Role of the I7 protein in proteolytic processing of
vaccinia virus membrane and core components. *J Virol* **78**, 6335-6343 (2004).

25.   T. G. Senkevich, C. L. White, E. V. Koonin, B. Moss, Complete pathway for protein
disulfide bond formation encoded by poxviruses. *Proc Natl Acad Sci U S A* **99**, 6667-6672
(2002).

26.   M. Cyrklaff *et al.*, Cryo-electron tomography of vaccinia virus. *Proc Natl Acad Sci U*
S A **102**, 2772-2777 (2005).

27.   J. Dubochet, M. Adrian, K. Richter, J. Garces, R. Wittek, Structure of intracellular
mature vaccinia virus observed by cryoelectron microscopy. *J Virol* **68**, 1935-1941 (1994).

28.   G. Griffiths, N. Roos, S. Schleich, J. K. Locker, Structure and assembly of
intracellular mature vaccinia virus: thin-section analyses. *J Virol* **75**, 11056-11070 (2001).

29.   A. R. Mohandas, S. Dales, Involvement of spicules in the formation of vaccinia virus
envelopes elucidated by a conditional lethal mutant. *Virology* **214**, 494-502 (1995).

30.   C. Morgan, The insertion of DNA into vaccinia virus. *Science* **193**, 591-592 (1976).

31.   N. Roos *et al.*, A novel immunogold cryoelectron microscopic approach to investigate
the structure of the intracellular and extracellular forms of vaccinia virus. *EMBO J* **15**, 2343-
2355 (1996).

32.   M. Cyrklaff *et al.*, Whole cell cryo-electron tomography reveals distinct disassembly
intermediates of vaccinia virus. *PLoS One* **2**, e420 (2007).

33.   B. Unger, J. Mercer, K. A. Boyle, P. Traktman, Biogenesis of the vaccinia virus
membrane: genetic and ultrastructural analysis of the contributions of the A14 and A17
proteins. *J Virol* **87**, 1083-1097 (2013).

34.   H. Bisht, A. S. Weisberg, P. Szajner, B. Moss, Assembly and disassembly of the capsid-
like external scaffold of immature virions during vaccinia virus morphogenesis. *J Virol* **83**,
9140-9150 (2009).

35.   M. Faini, R. Beck, F. T. Wieland, J. A. Briggs, Vesicle coats: structure, function, and
general principles of assembly. *Trends Cell Biol* **23**, 279-288 (2013).

36.   T. Kirchhausen, Three ways to make a vesicle. *Nat Rev Mol Cell Biol* **1**, 187-198
(2000).

37.   J. C. Stachowiak, F. M. Brodsky, E. A. Miller, A cost-benefit analysis of the physical
mechanisms of membrane curvature. *Nat Cell Biol* **15**, 1019-1027 (2013).

38.   J. C. Stachowiak *et al.*, Membrane bending by protein-protein crowding. *Nat Cell*
Biol **14**, 944-949 (2012).

39.   K. J. Erlandson *et al.*, Poxviruses Encode a Reticulon-Like Protein that Promotes
Membrane Curvature. *Cell Rep* **14**, 2084-2091 (2016).

40.   B. Sodeik *et al.*, Assembly of vaccinia virus: role of the intermediate compartment
between the endoplasmic reticulum and the Golgi stacks. *J Cell Biol* **121**, 521-541 (1993).

41.   J. R. Rodriguez, C. Risco, J. L. Carrascosa, M. Esteban, D. Rodriguez, Vaccinia virus
15-kilodalton (A14L) protein is essential for assembly and attachment of viral crescents to
virosomes. *J Virol* **72**, 1287-1296 (1998).

42.   M. Ericsson *et al.*, Characterization of ts 16, a temperature-sensitive mutant of
vaccinia virus. *J Virol* **69**, 7072-7086 (1995).

43.   W. W. Yeh, B. Moss, E. J. Wolffe, The vaccinia virus A9L gene encodes a membrane
protein required for an early step in virion morphogenesis. *J Virol* **74**, 9701-9711 (2000).

44.   S. Cudmore *et al.*, A vaccinia virus core protein, p39, is membrane associated. *J
Virol* **70**, 6909-6921 (1996).

45.   S. Wilton, A. R. Mohandas, S. Dales, Organization of the vaccinia envelope and
relationship to the structure of intracellular mature virions. *Virology* **214**, 503-511 (1995).

46.   J. A. Briggs *et al.*, The mechanism of HIV-1 core assembly: insights from three-
dimensional reconstructions of authentic virions. *Structure* **14**, 15-20 (2006).

47.   J. A. Briggs, T. Wilk, R. Welker, H. G. Krausslich, S. D. Fuller, Structural organization
of authentic, mature HIV-1 virions and cores. *EMBO J* **22**, 1707-1715 (2003).

48.   R. D. M. Gray *et al.*, Nanoscale polarization of the entry fusion complex of vaccinia
virus drives efficient fusion. *Nat Microbiol* **4**, 1636-1644 (2019).

49.   T. G. Senkevich, B. M. Ward, B. Moss, Vaccinia virus entry into cells is dependent on
a virion surface protein encoded by the A28L gene. *J Virol* **78**, 2357-2366 (2004).

50.   B. Moss, Poxvirus cell entry: how many proteins does it take? *Viruses* **4**, 688-707
(2012).

51.   F. I. Schmidt, C. K. Bleck, A. Helenius, J. Mercer, Vaccinia extracellular virions enter
cells by macropinocytosis and acid-activated membrane rupture. *EMBO J* **30**, 3647-3661
(2011).

52.   J. Mercer, A. Helenius, Vaccinia virus uses macropinocytosis and apoptotic mimicry
to enter host cells. *Science* **320**, 531-535 (2008).

53.   R. A. Janeczko, J. F. Rodriguez, M. Esteban, Studies on the mechanism of entry of
vaccinia virus in animal cells. *Arch Virol* **92**, 135-150 (1987).

54.   G. C. Carter, M. Law, M. Hollinshead, G. L. Smith, Entry of the vaccinia virus
intracellular mature virion and its interactions with glycosaminoglycans. *J Gen Virol* **86**,
1279-1290 (2005).

55.   R. W. Doms, R. Blumenthal, B. Moss, Fusion of intra- and extracellular forms of
vaccinia virus with the cell membrane. *J Virol* **64**, 4884-4892 (1990).

56.   S. Kilcher *et al.*, siRNA screen of early poxvirus genes identifies the AAA+ ATPase D5
as the virus genome-uncoating factor. *Cell Host Microbe* **15**, 103-112 (2014).

57. S. Hutin *et al.*, Domain Organization of Vaccinia Virus Helicase-Primase D5. *J Virol*
90, 4604-4613 (2016).

58. N. Wang *et al.*, Architecture of African swine fever virus and implications for viral
assembly. *Science* **366**, 640-644 (2019).

59. N. G. Abrescia, D. H. Bamford, J. M. Grimes, D. I. Stuart, Structure unifies the viral
universe. *Annu Rev Biochem* **81**, 795-822 (2012).

60. C. H. Durkin *et al.*, RhoD Inhibits RhoC-ROCK-Dependent Cell Contraction via PAK6.
Dev Cell **41**, 315-329 e317 (2017).

61. H. Lynn *et al.*, Loss of cytoskeletal transport during egress critically attenuates
ectromelia virus infection in vivo. *J Virol* **86**, 7427-7443 (2012).

62. X. Snetkov, I. Weisswange, J. Pfanzelter, A. C. Humphries, M. Way, NPF motifs in
the vaccinia virus protein A36 recruit intersectin-1 to promote Cdc42:N-WASP-mediated
viral release from infected cells. *Nat Microbiol* **1**, 16141 (2016).

63. J. V. Cordeiro *et al.*, F11-mediated inhibition of RhoA signalling enhances the spread
of vaccinia virus in vitro and in vivo in an intranasal mouse model of infection. *PLoS One* **4**,
e8506 (2009).

64. J. R. Kremer, D. N. Mastronarde, J. R. McIntosh, Computer visualization of three-
dimensional image data using IMOD. *J Struct Biol* **116**, 71-76 (1996).

65. Q. Xiong, M. K. Morphew, C. L. Schwartz, A. H. Hoenger, D. N. Mastronarde, CTF
determination and correction for low dose tomographic tilt series. *J Struct Biol* **168**, 378-
387 (2009).

66. B. Turonova, F. K. M. Schur, W. Wan, J. A. G. Briggs, Efficient 3D-CTF correction for
cryo-electron tomography using NovaCTF improves subtomogram averaging resolution to
3.4A. *J Struct Biol* **199**, 187-195 (2017).

67. Y.-T. Liu *et al.*, Isotropic Reconstruction of Electron Tomograms with Deep Learning.
bioRxiv 10.1101/2021.07.17.452128, 2021.2007.2017.452128 (2021).

68. D. Castano-Diez, M. Kudryashev, H. Stahlberg, Dynamo Catalogue: Geometrical tools
and data management for particle picking in subtomogram averaging of cryo-electron
tomograms. *J Struct Biol* **197**, 135-144 (2017).

69. T. A. M. Bharat, C. J. Russo, J. Lowe, L. A. Passmore, S. H. W. Scheres, Advances in
Single-Particle Electron Cryomicroscopy Structure Determination applied to Sub-tomogram
Averaging. *Structure* **23**, 1743-1753 (2015).

70. E. F. Pettersen *et al.*, UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82 (2021).

71.   I. Weisswange, T. P. Newsome, S. Schleich, M. Way, The rate of N-WASP exchange limits the extent of ARP2/3-complex-dependent actin-based motility. *Nature* **458**, 87-91 (2009).

Legends

Figure 1. Cryo-ET of vaccinia infected cells.

A. Schematic representation of the different stages in vaccinia assembly and envelopment. D13 coated membrane crescents initially close to form a spherical IV, that mature into brick shaped IMV with dumbbell shaped viral core flanked by two lateral bodies (red). Envelopment of IMV by a Golgi membrane cisterna or endosomal compartment (green) results in the formation of triple membraned IEV, which can fuse with the plasma membrane to liberate extracellular enveloped virions (EEV), which are called CEV if they remain attached to the surface of the cell. **B.** Left: Central section of a tomogram showing fully formed IV. The red arrow points at an IV with a lateral invagination. Invaginations in other IV can be seen in the 3D-segmented model of the membrane of the same tomogram (right) (See Movie 1). **C.** Tomogram middle view (left) and its corresponding segmentation (right) of IMV (See Movie 2). A "cut corner" (magenta arrows) is apparent in some IMV. Scalebars = 100 nm.

Figure 2. The stages of IV assembly

A. Tomogram section showing D13 coated membrane crescents with open ends (indicated magenta arrowheads). **B.** Tomogram sections showing the last stages of IV membrane closure: the middle view (left) reveals membrane discontinuity (indicated magenta arrowheads) while in the higher plane at + 48 nm relative to the middle view (right) the membrane is continuous (see Movie 3). **C.** Closed IV containing a condensed nucleoid (outlined in magenta) with repeated structural features (magenta arrowheads). **D.** Cytoplasmic condensed nucleoid/s next to an IV. **E.** Extended nucleoid/s in association with an open IV. **F.** Structured illumination microscopy image of infected cell showing condensed DNA structures (green) in association with RFP-A3 positive viral particles (magenta) located in a perinuclear viral factory. Tomogram Scalebars = 100 nm.

Figure 3. In situ structure of the D13 lattice.

A. Tomogram section as indicated in the schematic on the left through the middle and top of a representative IV (see Movie 4). The densities corresponding to the D13 lattice are coloured in magenta in panels (right) and show pseudo-hexagonal organisation. Scale bar =

100 nm. **B.** Map based on subtomogram averaging of D13 and the underlying IV membrane

from a top (left) and a side (right) view with fitting of coordinate models of D13 trimers(PDB

7VFE) shown in blue/brown.

Figure 4. Ultrastructure of IMV.

A. Central plane of a tomogram showing the broadest view of a representative IMV together

with a region magnified 2.5-times to highlight the outer layer, viral membrane, palisade

and inner wall of the virion (see Movie 5). **B.** Lateral view of an IMV, showing the lateral

bodies (outlined in red) and the dumbbell shape of the core. The magnified regions outlined

in magenta show the same characteristic IMV layers. **C.** Different sections of the IMV shown

in A at the indicated plane positions in nm. The insert in the +31 nm view shows the surface

of the palisade lattice. In the second row, yellow highlights the palisade structure, while

red indicates the lateral body in the +52 nm view. **D.** The first row shows a schematic

representation of the views displayed in C, while the second row represents the orthogonal

view together with the tomogram position (black line). **E.** Left: Top and side views of a

segmentation model of the IMV in A. The outer layer (blue) and viral membrane (black) are

cut away so the internal palisade (yellow) and the lateral bodies (red) can be seen (see

Movie 6). Right: The dumbbell shaped core is more apparent in a Simulated Digitally

Reconstructed Radiograph (DDR Rendering) of the same segmentation model. Scale bars =

100 nm.

Figure 5. Ultrastructure of the naked viral core.

A. Middle section of a tomogram showing a naked viral core together with a region

magnified 3 times to highlight the palisade and inner wall as well as the polymers that

surround the core. **B.** A higher tomogram section (+ 41 nm) of the same naked viral core

highlighting the pseudohexagonal lattice of the palisade. **C.** At an even higher section (+58

nm) ring structures associated with the palisade are evident. The dimensions of the rings

(n=25) from 5 naked cores are indicated together with the standard error of the mean. Scale

bars = 100 nm.

Figure 6. The palisade is a pseudohexagonal lattice composed of trimeric proteins.

A. Central plane of a tomograms showing that the additional membranes of IEV and CEV

(associated with the plasma membrane, PM) do not alter the ultrastructure of the internal

IMV. Scale bar = 100 nm. **B.** Schematic representation to illustrate the number of membranes

of IMV, IEV and EEV/CEV. **C.** Sections through the palisade layer corresponding to the virions

shown in A. Scale bar = 10 nm. **D.** Map derived from subtomogram averaging showing the

inner wall, palisade layer, viral membrane, and outer layer in surface representation (left) and corresponding sections of the map (right).

Figure 7. Maturation of vaccinia virus: from IV to IMV.

A. Table showing the measured dimensions of middle views of IV (diameter, n = 26; perimeter, n = 15) and IMV (major axis, n = 22; intermediate axis, n = 19; minor axis, n = 10; perimeter, n = 19) tomograms. * indicates that the IMV minor axis was calculated from the side views of 3 IMV and 7 IMV inside EEV (see methods for explanation). For volume calculation IV and IMV were assumed to be spheres and triaxial ellipsoids respectively. **B.** Plots showing the individual values for perimeter and diameter / major axis (indicated by magenta lines in schematics below graph) of the IV, IMV and EEV. IV and IMV measurements were used to calculate the averages shown in A. The averages of the EEV inner most membrane perimeter and major axis are 1074.46 nm ± 7.09 nm and 348.91 ± 1.87 nm, respectively. **C.** Central sections of IV and IMV along with possible intermediates together with magnified regions (2.5-times) highlighting the different layers of each particle. Scale bar = 100 nm. **D.** Vaccinia maturation model: the palisade lattice (yellow) forms inside D13-coated IV, its maximum length being determined by the IV diameter. Following disassembly of the D13 lattice, the viral membrane acquires the shape of the palisade and becomes wrinkled. During this process, the presence of two lateral bodies (red) results in the deformation of the core so it adopts its characteristic biconcave shape.

Supplementary Table 1. Measurements and calculations of virion dimensions.

A. The middle-view perimeters, IV diameter, the major axis of IMV and EEV/CEV are shown, together with the standard error of the mean (SEM) and the number of viral particles measured. The intermediate and minor axes of IMV are also shown, as well as the IEV axes. Finally, the calculated volume of IV, IMV and IEV are provided. See Methods for details of dimension calculations. **B**. For the thickness of the inner wall and the outer layer, 5 different IMV were measured by tracing 5 different lines for each IMV, which correspond to the five measurements shown. To estimate the distance between the palisade and the viral membrane (with no lateral bodies) 5 IMV were used. For the distance between the palisade and the plasma membrane through lateral bodies 1 IMV and 4 EEV were included in the quantification. All values correspond to nm.

Supplementary Table 2. Cryo-ET data collection and STA processing.

Supplementary Figure 1. Low-magnification images of vaccinia infected HeLa cells on grids. A. The pink circle marks the position where the IV in Figure 1B were imaged. The image on the right is a magnification of the circled area and its surroundings. **B.** A cryo-EM image showing IV (magenta arrow) and IMV (green arrow) visualised at low magnification in cells over holes in the carbon support film. Scale bars = 10 (left) and 1 μm (right) (A) and 1 μm (B).

Supplementary Figure 2. Tomographic gallery of IV with nucleoids
Tomographic sections showing IV with an internal condensed nucleoid outlined in magenta in the bottom row. Scale bar = 100 nm.

Supplementary Figure 3. Examples of IMV with cut corners
Gallery of tomographic sections of IMV showing cut or flattened corner/s (black arrowheads). Scale bar = 100 nm.

Supplementary Figure 4. Palisade maps from IMV, IEV and EEV/CEV.
A. Maps derived from subtomogram averaging IMV, IEV or EEV/CEV. A surface (top) and a cut-section view (bottom) are shown. **B.** Fourier Shell Correlation plots for palisade maps as well as for the D13 map. Curves are given for the corrected, unmasked, masked and masked with phase randomisations calculations.

Movie 1. IV overview.
Tomogram of an intracellular region accumulating IV, which corresponds to the section shown in Figure 1B. IV membranes are segmented in red

Movie 2. IMV overview.
Tomogram showing IMV in the cytoplasm, which corresponds to the section shown in Figure 1C. IMV membrane is segmented in blue, microtubules in red and the lumen of an ER-like compartment in yellow.

Movie 3. IV assembly. Tomogram of an IV with a pore on the viral membrane (pore labelled with magenta arrowheads). Sections of this tomograms are shown in Figure 2B.

Movie 4. Ultrastructure of IV.
Tomogram corresponding to the IV shown in Figure 3A. Scale bar = 100 nm.

Movie 5. Ultrastructure of IMV.

Tomogram of the IMV particle shown in Figure 4A. Scale bar = 100 nm.

Movie 6. Segmentation of an IMV. The movie shows the segmentation performed on the

IMV from Figure 4A. Blue is the outer layer and black labels the viral membrane, while

yellow corresponds to the palisade lattice and red to the lateral bodies.

Movie 7. Ultrastructure of the naked viral core.

Tomogram of the naked core shown in Figure 5. The central magenta square marks part of

the palisade, the top magenta rectangle highlights the flexible polymer observed and the

small magenta squares at the end of the movie outline the ring-like structures protruding

from the palisade. Scale bar = 100 nm.

821 **Supplementary Table 1. Measurements and calculations of virion dimensions.**

823 **A.**

Middle view perimeter	Average (nm)	SEM	
IV	1106.39	8.58	n = 15
IMV	1087.75	11.25	n = 19
EEV / CEV	1074.46	7.09	n = 20

Diameter / major axis	Average (nm)	Standard Error of the Mean (SEM)	
IV	351.89	2.88	n = 26
IMV	351.70	3.31	n = 22
EEV / CEV	348.91	1.87	n = 22

IMV	Average (nm)	SEM	
Intermediate axis	280.53	3.66	n = 19
Minor axis*	198.04	4.94	n = 10 (3 IMV, 7 EEV)

IEV	Average (nm)	SEM	
Major axis	437.86	13.88	n = 7
Intermediate axis	383.45112	10.95	n = 7
Minor axis (estimation)	259.11		

	Calculated volume (nm^3)
IV	2.28×10^7
IMV	1.02×10^7
IEV	2.28×10^7

837 **B.**

Inner wall thickness (nm)	IMV1	IMV2	IMV3	IMV4	IMV5			
1	4.89	4.12	2.08	4.43	3.72			
2	4.29	3.88	4.28	3.20	2.71			
3	3.54	4.21	4.75	3.11	4.78			
4	3.25	3.48	3.61	2.38	4.28	Final average	Standard Deviation	SEM
5	3.91	4.02	2.81	6.05	4.20			
average	3.98	3.94	3.51	3.84	3.94	3.84	0.19	0.09

Outer layer thickness (nm)	IMV1	IMV2	IMV3	IMV4	IMV5			
1	7.24	8.15	6.14	6.90	5.61			
2	7.51	7.14	7.25	6.52	7.31			
3	7.48	6.55	5.86	6.37	5.05			
4	6.40	7.12	6.03	6.96	6.63	Final average	Standard Deviation	SEM
5	5.58	5.77	7.71	6.28	7.11			
average	6.84	6.95	6.60	6.61	6.34	6.67	0.11	0.24

Palisade-to-viral membrane distance (nm)	IMV1	IMV3	IMV4	IMV5			
1	9.54	10.12	8.89	9.47			
2	9.57	9.37	7.87	8.08			
3	11.18	9.51	8.32	8.31	Final average	Standard Deviation	SEM
4	8.89	9.57	11.28	10.98			
5	8.66	8.53	7.36	10.04			
average	9.57	9.42	8.74	9.38	9.28	0.18	0.36

Palisade to viral membrane through lateral bodies (nm)	IMV1	EEV1	EEV2	EEV3	EEV4	Average	Standard Deviation	SEM
1	53.20	50.35	43.45	52.01	41.76			
2	31.93	48.19	42.39	62.64	39.35	46.53	8.60	2.72

842 **Supplementary Table S2. Cryo-ET data collection and STA processing.**

	Palisade				D13
Data collection	Combined	EEV/CEV	IMV	IEV	
Voltage (kV)	300				
Cumulative electron dose (e / $Å^2$)	66.3				
Applied defocus (μm)	-8				
Pixel size (Å)	4.31				
Tilt range (°)	±57°, dose symmetric				
STA processing					
Map symmetry	C3	C3	C3	C3	C6
Initial particle number	182938	209703	80698	8256	341376
Final particle number	179822	149662	27590	5686	34092
Map resolution (Å) (FSC=0.143)	21.2	21.2	25.1	30.6	19.2

A

Crescent → IV → IMV → wrapping with Golgi / endosome → IEV → CEV → EEV

cell exit

plasma membrane

non-infectious particles | infectious particles

B

C

Figure 1

A

B +48 nm

C

D

E +31 nm

F Viral particle (RFP-A3)
DNA (DAPI)

5 μm

Figure 2

A

B

D13
lattice

gap

viral
membrane

Figure 3

Figure 4

A

palisade
inner wall

extended
polymers

B   +41 nm

palisade
surface

C   +58 nm

9.9±0.1 nm

4.5±0.1 nm

ring
structures

25 rings
5 naked cores

Figure 5

Figure 6

A

	Immature virus (IV)	Intracellular mature virus (IMV)	
Approximate shape	Sphere	Triaxial ellipsoid	
Dimensions (nm)	Diameter	351.89 ± 2.88	Major axis	351.70 ± 3.31
			Intermediate axis	280.53 ± 3.66
			Minor axis*	198.04 ± 4.94
Perimeter (middle plane, nm)	1106.39 ± 8.58	1087.75 ± 11.25	
Calculated volume (nm³)	2.28 x 10⁷	1.02 x 10⁷	

B

middle plane

perimeter

diameter / major axis

C

Possible intermediates

IV   D13 partial Palisade present   D13 absent Palisade present   IMV

D13 coat
viral membrane

D13 coat
outer layer
viral membrane
palisade
inner wall

outer layer
viral membrane
palisade
inner wall

outer layer
viral membrane
palisade
inner wall

D

D13 lattice   viral membrane   nascent palisade   disassembling D13   palisade   outer layer

90°   lateral body   90°   90°   lateral body

D13

Immature virus   Maturing virus   Palisade   Mature virus

Figure 7

A

B

Supplementary Figure 1

Annotated

Supplementary Figure 2

Cut corner : 48.9%   No cut corner : 51.1%   n = 94 IMV

Supplementary Figure 3

A

B

Supplementary Figure 4